Title: The Role of Vitamin D in *Emiliania huxleyi*: A Microalgal Perspective on UV-B Exposure

Authors: Or Eliason[1], Sergey Malitsky[2], Irina Panizel[2], Ester Feldmesser[2], Martin Sperfeld[1], Einat Segev[1]*

Affiliations:

[1]Department of Plant and Environmental Sciences, Weizmann Institute of Science; Rehovot, 7610001, Israel.

[2]Department of Life Sciences Core Facilities, Weizmann Institute of Science; Rehovot, 7610001, Israel.

*Corresponding author. Email: Einat.Segev@weizmann.ac.il

Abstract

An essential interaction between sunlight and eukaryotes involves the production of vitamin D through exposure to ultraviolet-B (UV-B) radiation. While extensively studied in vertebrates, the role of vitamin D in non-animal eukaryotes like microalgae remains unclear. *Emiliania huxleyi*, a microalga inhabiting shallow ocean depths exposed to UV-B radiation, is well-suited for this research. Our results show that *E. huxleyi* can produce vitamin D_2 and D_3, pointing to their potential role in the algal physiology. We further show that *E. huxleyi* algae respond to vitamin D at the transcriptional level, regulating the expression of protective mechanisms such as the light-harvesting complex stress-related protein (LHCSR) and heme oxygenase, and that vitamin D enhances the algal photosynthetic performance while reducing harmful reactive oxygen species buildup. Understanding the function of vitamin D in *E. huxleyi* has broader implications, shedding light on its role in non-animal eukaryotes and its potential importance in marine ecosystems. This research sets the stage for further investigations into the complex relationship between sunlight, vitamin D, and microalgal physiology, which contributes to our understanding of how eukaryotes adapt to diverse environmental conditions.

Introduction

Life on Earth has a complex relationship with sunlight, relying on its energy for certain processes while simultaneously requiring protection against its potential harmful effects. A molecular process that is tightly linked to sunlight is the formation of vitamin D following exposure to ultraviolet-B (UV-B) radiation emitted from the sun. Vitamin D (calciferol) comprises a group of steroids that result from the photochemical transformation of several sterol precursors by UV-B wavelengths[1]. The most common vitamin D species known to occur naturally are vitamin D_2 and D_3, originating from the conversion of ergosterol and 7-dehydrocholesterol, respectively[2,3].

In mammals and other studied vertebrates, vitamin D functions as a hormone, involved in the regulation of a multitude of intracellular and physiological processes vital for the organism survival and well-being[4]. Vitamin D is pivotal in facilitating the absorption and homeostasis of essential ions such as Ca^{2+} and PO_4^{3-} [5,6], and vitamin D deficiency has been linked to a range of physiological disorders[7,8]. Due to its key role in human health, vitamin D has been the focus of biological and pharmaceutical research endeavors. These efforts have been directed towards understanding the mechanisms through which it operates in humans and human models.

However, mounting evidence suggests that vitamin D has been a constituent of eukaryotes long before the emergence of vertebrates. This is evident not only in the identification of vitamin D in distant eukaryotic lineages like algae[9–11], plants[12], and fungi[13–15] but also in the preservation of vitamin D-related biomarkers, likely from an algal source, in marine sediments dating back over 600 million years[16].

Despite its widespread presence across diverse lineages, our understanding of the physiological role of vitamin D in non-animal eukaryotes remains limited. Non-animal eukaryotes, namely microalgae, have been suggested as potential sources of vitamin D for higher trophic levels in the marine environment[17–19]. But the processes underlying vitamin D production and regulation in microalgae remain largely unexplored[20].

While UV-B radiation is crucial for vertebrate health due to its role in vitamin D formation, it can also be detrimental, causing direct damage to biomolecules like DNA, leading to the generation of reactive oxygen species (ROS)[21–23], and can ultimately result in cell death[24]. Photosynthetic organisms, like algae, are particularly susceptible to UV-B damage, as their energy production

hinges on exposure to solar radiation[25,26]. Although water acts as a UV-B filter[27], significant intensities can still penetrate the upper layers of ocean surfaces[28], potentially impacting organisms such as algae[29].

Microalgae of the species *Emiliania huxleyi*, also named *Gephyrocapsa huxleyi*[30], are widely distributed in modern oceans and play key roles in various biogeochemical cycles[31,32]. These algae are known to flourish in high light environments at shallow depths of about 10 to 20 meters[33], where exposure to UV-B wavelengths is likely. Earlier findings provided intermittent indications that *E. huxleyi* algae might possess the inherent capability to synthesize vitamin D. These reports highlighted the algal capacity to generate vitamin D_2 upon exposure to UV-B irradiation[11], its cholesterol content[34,35], and the presence of a gene analogous to 7-dehydrocholesterol reductase (DHCR7) responsible for converting 7-dehydrocholesterol into cholesterol[36].

In this study we explore the overlooked role of vitamin D in *E. huxleyi*. Specifically, we investigate the relationship between vitamin D formation following exposure to UV-B and the regulation of cellular mechanisms that operate in response to harmful radiation.

Results

E. huxleyi algae produce vitamin D_2 and D_3

To investigate whether vitamin D species are formed by *E. huxleyi* algae upon exposure to UV, we cultivated algal cultures in a chamber with environmentally relevant UV-B radiation levels (see Materials and Methods). Metabolic analyses revealed the presence of both D_2 and D_3 in these algal cultures (Table 1). Our results show that D_2 was significantly enriched in UV-exposed cultures, with levels of approximately ~4 ng/mg dry weight, while it was barely detected in cultures that were not exposed to UV. The D_2 precursor ergosterol was found in both UV-treated and control cultures. Lower amounts of vitamin D_3 (~0.04 ng/mg dry weight) were detected in both UV-treated and control cultures.

Importantly, while D_2 detection was consistent across all analyzed UV-exposed samples, D_3 was identified only in part of the experiments during our research. Interestingly, when D_3 was detected, its precursor 7-dehydrocholesterol, was detected as well. Inconsistent detection of D_3

was previously reported in plants, and was attributed to the sensitivity of the analytical method

used[37]. Our many efforts to resolve the variable measurements of D_3 were not successful (see

detailed description of attempts in Materials and Methods). Collectively, our findings

demonstrate that *E. huxleyi* algae produce D_2 and D_3, suggesting a possible cellular function for

vitamin D.

E. huxleyi algae show a transcriptomic response to UV radiation

UV radiation is necessary for the formation of vitamin D^2. Therefore, we sought to explore the

transcriptomic response of *E. huxleyi* algae to UV exposure, aiming to elucidate cellular

processes that may be related to vitamin D. To achieve this, we analyzed the *E. huxleyi*

transcriptome using cultures that were grown under continuous UV exposure during the light

period of the daily light/dark cycle, in comparison to algal cultures that were protected from the

UV source. Cultures were sampled for RNA-sequencing at three time points representing

different growth phases (days 7, 10, and 13, see Fig. S1).

The transcriptomic analysis revealed differential expression (DE) of 374 genes between UV-

exposed and control cultures (Table S1). Of these genes, 172 were annotated with GO terms

related to a known function or process. The annotated genes that were DE in the transcriptome

under UV exposure were associated with various cellular processes including intracellular

signaling pathways and stress response mechanisms. Notably, genes participating in the inositol

Compound	UV (ng / mg)	Control (ng / mg)
D_2	4.32 ± 1.39 *	0.09 ± 0.01
D_3	0.038 ± 0.001	0.039 ± 0.001
Ergosterol	83.01 ± 28.96	71.76 ± 9.53
7-dehydrocholesterol	0.24 ± 0.04 *	0.16 ± 0.02

Table 1. *E. huxleyi* algae produce vitamin D_2 and D_3. Metabolic analysis of vitamin D species and precursors under UV and control conditions, using algal cultures at day 10 of growth. Values are of ng / mg dry weight. Statistically significant differences ($P < 0.05$) between treatments are marked by *, calculated using two-sample t-test assuming equal variances. ± values indicate standard deviation based on 4 biological replicates.

3-phosphate/calcium (IP_3/Ca^{2+}) and the oxylipin signaling pathways were DE (Table 2), both
playing key roles in stress response mechanisms across different organisms[38–42].

Furthermore, a substantial number of DE genes were involved in various stress responses,
including DNA damage sensing and repair, oxidative stress mitigation, protective pigment
biosynthesis, and maintenance of the photosynthetic machinery. Interestingly, several of the
genes and pathways that were DE in *E. huxleyi* algae, are known to be associated with UV
exposure and vitamin D biosynthesis in vertebrates. For instance, the IP_3/Ca^{2+} and
prostaglandins pathways are involved in UV stress response in mammals[43,44], and vitamin D is
involved in the regulation of these pathways[45–49]. In mammals, vitamin D also plays a role in
oxidative stress mitigation, DNA repair, and the regulation of various enzymes related to stress
response including heme oxygenase, glutathione peroxidase, and tyrosinase[50–54]. Considering
the role of vitamin D in regulating stress response mechanisms in mammals, and the presence
of similar mechanisms regulated by UV in *E. huxleyi*, it seems plausible that vitamin D plays a
role in regulating the algal stress responses.

To further explore the observed transcriptomic response induced by UV using an independent
approach, several genes that were regulated in the algal transcriptome were analyzed by qRT-
PCR. RNA was extracted from cultures that were exposed to UV for a duration of 1 hour at day
10 of growth (Fig. 1, S2). Under these conditions, the investigated genes showed significant
upregulation. The differences observed between the two transcriptomic assays, could be the
result of a different response elicited by a prolonged compared to a brief UV exposure.

Vitamin D upregulates expression of UV-regulated genes

Next, we explored whether vitamin D is indeed involved in regulating stress response
mechanisms in *E. huxleyi* under UV. If vitamin D is synthesized upon UV exposure and plays a
role in regulating the algal stress response to UV, external addition of vitamin D could potentially
induce algae to react as if they were exposed to UV. Therefore, we

Gene ID	Putative protein or domain	Differential expression			Protein function
	Genes related to intracellular signaling	d. 7	d. 10	d. 13	
G10384	Lipoxygenase domain	-0.46	-1.27	2.04	Oxylipin biosynthesis
G14992	Prostaglandin F(2-alpha) synthase	0.38	0.22	1.75	Oxylipin biosynthesis
G12340	Cytosolic phospholipase A2 domain	-0.51	-0.82	1.70	Oxylipin biosynthesis; Intracellular signaling
G14502	Ca-binding domain	-0.57	-1.12	1.67	Shares similarity to *A. thaliana* calmodulin-like protein; Intracellular signaling.
G21784	Phosphoinositide phospholipase C	-0.52	-0.62	1.50	Ip$_3$ signaling initiation; cytosolic calcium regulation
G15496	Phosphoinositide 5-phosphatase	0.54	0.74	1.46	IP$_3$ signaling; cytosolic calcium regulation
G25467	Steroid hydroxylase	1.15	1.49	1.16	Shares similarity to rat *cyp1a2* with a suggested 25-cholesterol hydroxylase activity
G1648	Mannosyl phosphorylinositol ceramide synthase	-0.37	-0.21	0.98	Potentially involved in Calcium signaling
G19702	Phosphatidylinositol 3-kinase	-0.17	0.00	0.95	IP$_3$ signaling
G27192	Calcineurin B-like interacting protein kinase	-0.34	1.25	-0.3	Involved in Ca-mediated signaling; Abiotic stress response
	Genes related to stress response				
G27084	Deoxyribodipyrimidine photolyase	0.93	1.9	4.29	UV-damage DNA repair
G647	3-dehydroquinate synthase domain	-0.31	5.33	4.07	Shikimate pathway; potentially involved in the production of UV-protective compounds
G22973	Sirtuin 2	1.16	0.89	1.49	DNA transcription and repair
G16746	Tyrosinase Cu-binding domain	-0.54	-1.09	1.46	Involved in synthesis of protective pigments and antioxidants
G18590	Light-harvesting protein	0.51	1.22	1.13	Shares similarity to to *C. reinhardtii* LHCSR; Alleviates photo-oxidative stress
G12503	Poly ADP-ribose polymerase Zn-finger domain	0.62	1.13	1.08	DNA damage sensor
G18115	Deoxyribodipyrimidine photolyase	0.24	0.25	0.99	UV-damage DNA repair
G6690	Glutathione peroxidase	0.05	0.32	0.98	Oxidative stress mitigation
G25108	Chalcone synthase 2	-0.36	-0.43	0.82	Potentially involved in UV protection
G2876	DNA-(apurinic or apyrimidinic site) lyase	0.56	0.52	0.81	DNA repair
G26038	ATP-dependent DNA helicase	-0.34	-0.76	0.79	DNA stability and repair
G8907	(6-4)DNA photolyase	0.07	0.43	0.78	UV-damage DNA repair
G2511	Heme oxygenase	0.37	0.96	0.67	Oxidative stress mitigation
G16133	Formamidopyrimidine-DNA glycosylase	0.49	0.89	0.53	DNA repair
G832	Protochlorophyllide reductase	1.23	-0.41	0.2	Chlorophyll biosynthesis
G22197	5'-tyrosyl DNA phosphodiesterase	0.83	0.22	0.10	DNA repair

Table 2. Differential expression (DE) of genes associated with signaling and stress response mechanisms in *E. huxleyi* that were upregulated under UV. DE values calculated according to the transcriptomic analysis are given for days 7, 10 and 13 of growth (designated d. 7, d. 10 and d. 13). Genes are ordered according to DE values at day 13. Full DE and NCBI accession data is presented in Table S1 and Data S1.

supplemented algal cultures with vitamin D and monitored the expression of stress response-

related genes via qRT-PCR.

Algal cultures were treated with D_2, D_3, or with a combination of both, as this combination had a

synergistic impact on algal growth compared to each species alone (Fig. S2). Control cultures

were not treated with vitamin D and were exposed to either normal growth conditions or to UV

radiation for 1 hour. RNA was collected from all cultures 1 hour post treatment.

Our analyses revealed four genes that exhibit upregulated expression both upon vitamin D

treatment and UV radiation (Fig. 1). The upregulation of these genes by UV was observed also

in the transcriptomic analysis (Tables 2, S1). Notably, in vitamin D-treated cultures, the

upregulation was only observed following the addition of both D_2 and D_3. The four upregulated

genes encode for light-harvesting complex stress-related protein (LHCSR, G18590), heme

oxygenase (G2511), steroid hydroxylase (G25467), and a Ca-binding protein (G26534). Both

LHCSR and heme oxygenase are proteins related to the algal stress response.

The LHCSR gene is known to exhibit increased expression in moss and green algae under UV-

B and high-light stress, promoting excess energy dissipation in the light-harvesting complex and

thereby reducing photo-oxidative stress[55–57]. Heme oxygenases are enzymes involved in the

formation of antioxidants in plants and animals, with known upregulated expression in response

to UV-B and other ROS-forming stressors[58–60]. UV-B radiation is a fundamental component of

high-light environments, and vitamin D is produced under UV-B radiation (Table 1) and triggers

the upregulation of oxidative- and photooxidative-stress mitigation pathways. Therefore, it is

likely that vitamin D is part of a cellular signaling cascade that reports on and reacts to harmful

light intensities or radiation.

Figure 1. Combined treatment of vitamin D₂ and D₃ upregulates UV-responsive genes. qRT-PCR analysis of genes following 1 hour of UV exposure or vitamin D treatments. Top title denotes gene products. In brackets: gene identifier in *E. huxleyi* CCMP3266[94] and matching gene transcript in the *E. huxleyi* CCMP1516 reference genome[95]. Statistical significance compared to control was calculated using two-tailed t-test assuming equal variances. One, two or three asterisks indicate $P < 0.05$, $P < 0.01$ and $P < 0.001$, respectively.

Vitamin D treatment improves the algal photosynthetic performance following excess light

To investigate the involvement of vitamin D in the cellular response towards harmful light intensities, we tested the impact of vitamin D treatment on algal physiology, namely photosynthetic performance, under excess light. To this end, algal cultures were exposed to low, non-saturating light levels that are regularly used during incubation[61] (130 μmol photons m^{-2} s^{-1}, or PAR), and to saturating light[62] (1000 PAR) for 2 hours. Algal cultures under each light regime were either treated with both D₂ and D₃, or not treated as control. Algal photosynthetic performance was evaluated by measuring nonphotochemical chlorophyll fluorescence quenching (NPQ) as a proxy for the ability of algae to dissipate excess absorbed light energy into heat[63]. In addition, the maximum PSII quantum yield (F_v/F_m) was measured as an indicator of photosynthetic efficiency[64]. Our results showed that algae under saturating light display decreased NPQ and F_v/F_m, indicative of photoinhibition[61] (Fig. 2a, b). Furthermore, vitamin D-treated cultures exhibited significantly higher NPQ following exposure to excess light (Fig. 2a, S4), in comparison to untreated cultures. The cultures that were treated with vitamin D also demonstrated enhanced F_v/F_m (Fig. 2b). These findings emphasize the potentially central role of vitamin D in algal physiology under harmful light levels. Vitamin D appears to activate cellular

170 processes that enhance excess light energy dissipation
171 and improve photosynthetic quantum yields.

Vitamin D alleviates ROS accumulation under excess light

Under excess light, photosynthetic organisms are likely
to experience oxidative stress[65], and NPQ is a key
process in mitigating oxidative stress induced by
excess light. Our analyses revealed elevated NPQ in
vitamin D-treated cultures following excess light.
Therefore, we assessed whether addition of vitamin D
to algal cultures under excess light, indeed alleviates
oxidative stress resulting in decreased cellular levels of
ROS. To achieve this, we subjected algal cultures to
excess light (1000 PAR) and assessed the intracellular
ROS levels using the cell-permeable fluorescent probe
2,7-Dichlorodihydrofluorescein diacetate (H_2DCF-DA).
The tested algal cultures were either treated with both
D_2 and D_3 or were untreated. Additionally, control
cultures were exposed to regular light intensities (130
PAR) and subjected to the same treatments. Our
findings revealed a substantial decrease in intracellular
ROS levels in algae that were treated with vitamin D
and exposed to excess light, compared with untreated
algae under the same light regime (Fig. 2c). These
findings further support a cellular role of vitamin D in

Figure 2. Vitamin D treatment improves photosynthetic performance and alleviates oxidative stress following exposure to excess light. (a) Non photochemical quenching (NPQ) and (b) F_v/F_m values of vitamin D-treated and control algae, exposed to regular (130 PAR) or excess (1000 PAR) light intensities. (c) Fluorescence values of vitamin D-treated and control algae stained with the intracellular reactive oxygen species (ROS) probe H_2DCF-DA, exposed to regular (130 PAR) or excess light (1000 PAR) intensities. Statistically significant values ($P < 0.05$) compared to control are marked by *, calculated from three biological replicates using two-tailed paired t-test for (a,b) and one-tailed paired t-test for (c).

195 algal cells under excess light conditions, demonstrating that vitamin D is involved in mitigating
196 ROS-induced oxidative stress.

Discussion

The current study reveals the photoprotective role of vitamin D in a globally abundant marine
algal species. Our findings introduce a novel cellular mechanism that utilizes the photochemical
transformation of vitamin D as an indicator of exposure to harmful radiation, consequently
enhancing the algal physiological response to excess-light stress. This enhanced response is
manifested by an increase in photosynthetic efficiency following exposure to conditions that
inhibit photosynthesis, along with an overall reduction in the development of ROS (Fig. 2).
Furthermore, the elevation in NPQ observed after vitamin D treatment, coupled with the
upregulation of LHCSR, suggests that a significant aspect of the response triggered by vitamin
D is aimed at mitigating photo-oxidative stress. Additionally, the fact that the physiological
response in *E. huxleyi* algae to vitamin D is only observed following saturating light conditions
may hint at a light-dependent regulatory mechanism.

Vitamin D can serve as a sensitive light indicator in phytoplankton. For example, phytoplankton
residing in surface mixed layers of oceans often encounter significant fluctuations in PAR
intensity throughout the day. These environments, which can extend depths of over 200
meters[66], may result in cells being transported over considerable vertical distances[67], leading to
rapid change in light intensity of hundreds of PAR within a matter of hours[33,68]. In such a dynamic
scenario, as cells ascend, they experience an increase in UV-B and subsequently may generate
vitamin D. The specificity of the photochemical conversion of vitamin D under UV-B wavelengths,
coupled with its relatively high photochemical quantum yield[69], suggests that it could function as
an independent and sensitive proxy for assessing exposure to UV-B radiation, high light levels,
or fluctuations in light intensity.

Interestingly, the impact of vitamin D on gene expression became evident only under a combined
treatment with D_2 and D_3 (Fig. 1). Similarly, the combined treatment led to a lower cell density
compared to separate D_2 or D_3 treatments (Fig S2). Decreased cell densities might be a result
of cell cycle arrest, which is a phenomenon previously described in *E. huxleyi*[70] and is known to

occur in response to UV radiation across various organisms[71–73]. Whether vitamin D influences

the algal cell cycle, and why a combination of D_2 and D_3 drives a detectable response, merits

further investigation.

When studying algal physiology, it is essential to acknowledge the significance of UV-B as an

influential environmental factor. In experimental setups aimed at studying algal physiology and

ecology, UV-B radiation has traditionally been excluded due to its detrimental effects. However,

this contradicts the natural conditions in which algae thrive, where they regularly encounter low

levels of UV-B. While the omission of UV-B simplifies experimental conditions for precise

variable isolation, our study unveils the profound influence of vitamin D, a product of UV-B

exposure, on the physiological response of *E. huxleyi* to environmental stress. These findings

shed light on the potentially advantageous role of UV-B for algae facing excess-light stress,

encouraging further exploration of the interplay between algae and this often-overlooked

environmental factor.

Our study offers comparative insights on the role of vitamin D in vertebrates and in *E. huxleyi*

algae. The extensive knowledge on vitamin D biology primarily originates from research on

humans and other vertebrates. Transposing this knowledge to *E. huxleyi* presents challenges

due to significant phylogenetic and physiological differences between multicellular organisms

and unicellular entities. Nevertheless, parallels can be drawn. Vitamin D was shown to enhance

cellular defense in human and mice keratinocytes against UV-induced oxidative stress and DNA

damage[53,54]. Vitamin D has also been shown to mitigate oxidative stress in rat liver and

intestine[50,74], partly through the upregulation of heme oxygenase, a response mirrored in vitamin

D-treated algal cells. Furthermore, the upregulation by vitamin D of certain genes in *E. huxleyi,*

potentially involved in intracellular signaling cascades (Fig. 2c, d), suggests the involvement of

vitamin D in initiating or contributing to signaling mechanisms. In mammals, vitamin D plays a

role in the activation of key signaling proteins, such as phospholipase C (PLC), phospholipase

A_2 (PLA$_2$) and phosphatidylinositol-3 kinase (PI3K)[75], which are essential for the rapid generation

of secondary messengers such as Ca^{2+} and IP$_3$. While *E. huxleyi* cells demonstrated the

upregulation of PLC, PLA$_2$, and PI3K following prolonged UV exposure (Table 1), further

investigation is needed to determine if vitamin D also participates in regulating these proteins in

algae. Interestingly, the involvement of Ca^{2+} signaling in photo-oxidative stress mitigation has

been reported in other marine algal species[76]. Another possible similarity between vertebrates and *E. huxleyi* is the activation of vitamin D. In vertebrates, vitamin D requires enzymatic modifications to become hormonally active. These modifications include the hydroxylation of the 1^{st} and 25^{th} carbons, to produce 1,25-$(OH)_2$-vitamin D^4. Similar hydroxylated D_2 and D_3 species were identified in the current research (Fig. S5) based on multiple reaction monitoring (MRM) profiles published previously[77]. Additional work is necessary to validate the detection of these hydroxylated species and the corresponding biosynthetic pathways in algae.

Vitamin D synthesis likely has ancient origins, given its presence across various lineages of eukaryotes[9–15]. Sterols are a defining feature of eukaryotes, and the enzymatic pathways leading to the production of ergosterol and 7-dehydrocholesterol, which are precursor molecules to D_2 and D_3 forms of vitamin D, may have existed in the last eukaryotic common ancestor (LECA)[78,79]. Consequently, it can be hypothesized that the earliest eukaryotes were already synthesizing vitamin D when exposed to solar radiation. While previous studies had postulated the ancient evolutionary origins of vitamin $D^{20,80}$, the understanding of its role beyond the animal kingdom had remained limited. Eukaryotes rely on oxygen (O_2), and their evolution is thought to have necessitated proximity to oxygenic photoautotrophs inhabiting the sunlit ocean surface[81]. This is because during their proposed emergence in the late Archean to early Proterozoic eras[82,83], the deep ocean was likely devoid of O_2 [84–86]. Considering the antioxidant properties exhibited by vitamin D in both animals and *E. huxleyi* algae, coupled with its evolutionary links to oxidative agents such as UV-B and O_2, it prompts the question of whether vitamin D evolved in early eukaryotes as a means of sensing oxidative environments. A similar evolutionary role has been proposed to sterols, and chiefly to cholesterol that has been widely studied in the context of environmental O_2[87,88]. Like sterols, vitamin D possibly played a role in the early stages of life on Earth.

Materials and methods

<u>Algal strain and growth conditions</u>

The axenic algal strain of *E. huxleyi* CCMP3266 was purchased from the National Center for Marine Algae and Microbiota (Bigelow Laboratory for Ocean Sciences, Maine, USA). Algae were grown in artificial sea water according to Goyet and Poisson[89] and supplemented with L1

medium according to Guillard and Hargraves[90], with the exception that Na_2SiO_3 was omitted

following the cultivation recommendations for this strain. Algae were grown in standing cultures

in borosilicate Erlenmeyer flasks with an initial inoculum of 330 cells/ml, placed in a growth

chamber at 18°C under a light/dark cycle of 16/8 hr. PAR intensity during the light period was

130 µmoles photons $m^{-2}\,s^{-1}$. Cultures volume was 50 ml except for cultures used to measure

growth under vitamin D treatment (Fig. S3) which were grown in 20 ml.

Continuous UV irradiation was achieved by placing a UV-emitting light source (Exo Terra Reptile

UVB150, Hagen, Montreal, Canada) inside the algal growth chamber. Algal cultures experienced

UV-A intensity of 0.5 w/m^2, UV-B intensity of 0.07 w/m^2 and UV-C intensity of 0.026 w/m^2,

measured using an ALMENO 2570 device (Ahlborn, Budapest, Hungary) placed within the

Erlenmeyer flask. The UV-B intensity used in this study was selected to replicate the average

UV-B radiation encountered at the ocean surface[29]. The UV light-source was operating daily for

14 hours in parallel to the PAR light period, starting one hour after PAR illumination started, and

ending one hour before PAR illumination ended. This irradiation regime aimed to mimic a

simplified day cycle including dawn and dusk periods.

Algal growth was monitored by a CellStream CS-100496 flow cytometer (Merck, Darmstadt,

Germany) using a 561 nm laser and plotting the chlorophyll fluorescence at 702/87 nm against

side scatter.

<u>Vitamin D treatment</u>

Cultures were treated with 1 µM of vitamin D_2 or D_3 (Sigma-Aldrich, Burlington, Massachusetts,

USA) dissolved in DMSO, as this was the minimal vitamin D concentration in which an effect

was observed on algal growth (Fig. S3b). For combined D_2 and D_3 treatment, 0.5 µM of each

species was added. The final DMSO concentration in cultures was 0.1%. Control cultures were

treated with an equal amount of DMSO.

<u>Vitamin D analysis</u>

Metabolic analysis was conducted following Oberson[77]. Standards for D_2, D_3, ergosterol and 7-

dehydrocholesterol were purchased in dry (Sigma-Aldrich) and dissolved in $CHCl_3$. Standard

solutions of the different metabolites were combined into a single solution and diluted to create

a standard curve. All final standards and samples were spiked with 50 ng of vitamin $D2\text{-}d_3$

(IsoSciences, Ambler, Pennsylvania, USA) serving as an internal standard. Algal samples were centrifuged, lyophilized and stored in -80°C until analysis. Saponification was achieved by resuspending samples in 108 µl 55% KOH, 192 µl ethanol, and 60 µl of 9% NaCl and 7.4% ascorbic acid, followed by homogenization and stirring at room temperature for 18 hours. Samples were then supplemented with 40 µl 10% NaCl and 300 µl of 20% ethyl acetate in heptane, vortexed extensively and centrifuged for 30 minutes. The upper phase was collected, and the process was repeated twice. Samples were evaporated, dissolved in 200 µl of 0.5% isopropanol in hexane and sonicated. Strata SI-silica 55 µm 70 A columns (Phenomenex, Torrance, California, USA) were used for solid phase extraction and were pre-conditioned with 1 ml of 50% $CHCl_3$ in isopropanol, followed by two washes with 1 ml of hexane. Samples were then loaded onto the columns and washed with 0.5 ml of 0.5% isopropanol in hexane which were discarded, and washed again with 2.5 ml of 2.5% isopropanol in hexane which were collected. Samples were evaporated and dissolved in 200 µl of PTAD in acetonitrile, sonicated, stirred at room temperature for 2 hours, centrifuged for 10 minutes and transferred into LC-MS vials. Samples were protected from light as much as possible during the extraction process.

Due to the inconsistency in identification of D_3 in algal samples, several technical adaptations regarding algal growth and sample collection were implemented and evaluated. To examine whether inconsistencies arise due to rapid D_3 enzymatic degradation, algal cultures were immediately placed on ice, centrifuged in a cooled, 4˚C centrifuge, and the supernatant quickly discarded and replaced with 50% methanol in DDW. The samples were then plunged into liquid nitrogen and stored in -80˚C. Later, samples were thawed, evaporated in vacuum to remove the methanol, lyophilized and proceeded to vitamin D extraction. Additional modifications included increasing the intensity of UV-B radiation during algal growth, increasing sample size by combining separate cultures, and using F/2 trace metals instead of L1 trace metals. These attempts did not improve the reproducibility of D_3 detection.

Vitamin D was measured by UPC2-ESI-MS/MS equipped with Acquity UPC2 system (Waters, Milford, Massachusetts, USA). The MS detector (Waters TQ-XS) was equipped with an ESI source. The measurements were performed in the positive ionization mode using MRM. The source and de-solvation temperatures were maintained at 150°C and 500°C, respectively. The capillary voltage was set to 1.5 kV. Nitrogen was used as the de-solvation gas and cone gas at

a flow rate of 700 L h^{-1} and 150 L h^{-1}, respectively. Ionization parameters of ergosterol, 7-

dehydrocholesterol, D_2 and D_3 were adjusted by direct infusion of standards. Ionization

parameters for other compounds were taken from Oberson[77].

UPC2 system: mobile phase A consisted of CO_2, and mobile phase B consisted of 98% MeOH,

2% DDW and 10mM ammonium formate. Make up solvent was 1% formic acid in 90% MeOH

and 10% DDW at a flow rate of 0.4 ml min^{-1}. The column (WATERS Acquity CSH FluoroPhenyl

1.7 μm, 3.0x100 mm, cat. 186006573) was maintained at 45°C, injection volume was 3 μl. At the

first 0.5 min of injection, 99.5% of mobile phase B, and 0.5% of mobile phase A were run at flow

rate of 2.0 ml min^{-1}. Then, mobile phase A was gradually reduced to 92% at 6 min, and further

decreased to 70% at a flow rate of 1.75 ml min^{-1} at 6.5 min. This composition of mobile phase

and flow rate were kept until 7 min, followed by increase in mobile phase A to 99.5% at 7.8 min,

and then increase in flow rate to 2 ml min^{-1} at 8.5 min, and running at those conditions until 9

min.

RNA extraction

Algal cultures were harvested for RNA extraction by centrifugation at 4000 rpm for 5 min at 18°C.

RNA was extracted using the Isolate II RNA mini kit (Meridian Bioscience, London, UK)

according to manufacturer instructions. Cells were ruptured in RLT buffer containing 1% β-

mercapto-ethanol by bead beating for 5 min at 30 mHz. RNA was then treated with 3 μl Turbo

DNAse (ThermoFisher, Waltham, MA, USA) in a 50 μl reaction volume, followed by a cleaning

step using RNA Clean & Concentrator^{TM}-5 kit (Zymo Research, Irvine, CA, USA) according to

manufacturer instructions. RNA was used for generating transcriptomic data and qRT-PCR

analysis.

Transcriptomic analysis

Transcriptomic data was generated using the MARS-seq library preparation protocol[91], and

analyzed with the UTAP pipeline[92]. As part of the pipeline, read counts for each gene were

normalized using the DESeq2's median of ratios method[93]. Differential expression (DE) between

treatments was calculated using the following thresholds: mean number of normalized reads

across all samples ≥ 5, adjusted p-value ≤ 0.05, Log2 fold change ≤ -0.7 or ≥ 0.7. The previously

generated *E. huxleyi* CCMP3226 synthetic genome (sGenome) and annotation file was used as

reference for the UTAP pipeline[94]. Briefly, the *E. huxleyi* CCMP3226 sGenome was generated by *de novo* transcriptome assembly of short-reads and long-reads. The assembled *E. huxleyi* CCMP3226 transcripts were then mapped to the *E. huxleyi* CCMP1516 reference genome[95] to define gene loci. For the current work, functional gene annotations were manually curated by identifying open reading frames in assembled transcripts using the ORF finder tool (www.ncbi.nlm.nih.gov/orffinder; transcript accessions are given in Table S1) and analyzing protein domains in the translated sequences using InterProScan 5[96]. Additionally, transcript sequences were searched against the swissprot database using NCBI blastx[97], and the hit with the highest E-value taken. Specifically, the gene loci analyzed using blastx were G18590, sharing highest similarity to *Chlamydomonas reinhardtii* LHCSR (NCBI accession P93664.1) with E-value of 1e-26 and nucleotide identity of 57%; G25467, sharing highest similarity to rat *cyp1a2* (NCBI accession P04799.2) with E-value of 1e-31 and nucleotide identity of 27%; G14502, sharing highest similarity to *Arabidopsis thaliana* Calmodulin-like protein 12 (NCBI accession P25071.3) with E-value of 6-e5 and nucleotide identity of 22.6%. The putative Ca-binding activity of G26534 was assessed by identifying *bona fide* Ca-binding domains using InterProScan 5. Specifically, we performed blastx[97] and focused on the highest hit that contained an identifiable protein domain using InterProScan 5[96], resulting in the identification of an EF-hand family protein in *Chrysochromulina tobinii* that harbors three EF-hand domain pairs (NCBI accession KOO34173.1, with E-value of 8e-13 and nucleotide identity of 32%).

Quantitative real time PCR (qRT-PCR)

Algal cultures were treated with vitamin D as described earlier. UV-treated cultures were exposed to the same UV intensities as described above and treated with equal amounts of DMSO. All treatments lasted 1 hour. Equal concentrations of RNA taken from 10 days old cultures were utilized for cDNA synthesis using Superscript IV (ThermoFisher), according to manufacturer instructions. qPCR was conducted in 384 well plates using SensiFAST SYBR Lo-ROX Kit (Meridian Bioscience, Cincinnati, OH, USA) in a QuantStudio 5 qPCR cycler (Applied Biosystems, Foster City, CA, USA). The qPCR program ran according to enzyme requirements for 40 cycles. Samples were normalized using three housekeeping genes: *alpha-tubulin*, *beta-tubulin* and *ribosomal protein l13 (rpl13)*. DNA contamination was assessed by applying the same program on RNA samples that were not reverse transcribed (omitting the Superscript IV

enzyme in the reverse transcription reaction mix). Gene expression ratios were analyzed according to Vandesompele[98] by geometric averaging of housekeeping genes. Relative gene expression levels were compared to control samples. Primer efficiencies were determined using the QuantStudio 5 software, by qPCR amplification of serially diluted cDNA. All primers had a measured efficiency between 80-120%. Primer sequences are given in Table S2.

Chlorophyll fluorescence

Algal cultures were divided into four subcultures that were subjected to one of four treatments: 1000 PAR with or without vitamin D (as described earlier) and 130 PAR with or without vitamin D. Chlorophyll A fluorescence parameters were estimated following 2 hours of treatment. Parameters were estimated by pulse amplitude-modulated fluorometry using WATER-PAM II (Heinz Walz GmbH, Effeltrich, Germany). Maximum photosystem II quantum yield (F_v/F_m) was calculated as $F_v/F_m = (F_m - F_0) / F_m$[64], where F_o is the baseline fluorescence under a measuring light of 160 PAR and F_m is the maximum fluorescence measured with a 0.9 s saturating light pulse of 6000 µmol photons $m^{-2} s^{-1}$. Non-photochemical quenching (NPQ) was calculated as $NPQ = (F_m - F'_m) / F'_m$[63], where F_m was the maximum fluorescence yield after dark adaptation and F'_m the maximum fluorescence yield under actinic light of 1150 PAR. Prior to analysis, algal samples were dark-adapted for 5 minutes. No major difference in F_v/F_m was observed between algae that were dark-adapted for 5 or 30 minutes following exposure to regular light (Fig. S6), suggesting that a darkness period of 5 minutes sufficiently relaxed PSII reaction centers.

Intracellular reactive oxygen species (ROS) measurements

Algal cultures were divided into subcultures and treated as described under 'chlorophyll fluorescence'. Vitamin D-treated and control cultures were placed in a growth chamber and exposed to either regular light conditions (130 PAR) or to high-light conditions (1000 PAR) for 5 hours. For intracellular ROS assessment, cultures were stained with 0.5 µM of H_2DCFDA (ThermoFisher) in DMSO. Staining was performed 2 hours after the start of the treatment, and cultures were left in the dark for 20 minutes before they were introduced back into the growth chamber for the remaining 3 hours. Samples were measured using CellStream CS-100496, excited at 488 nm and the signal was collected at 528/46 nm. The algal population was gated by plotting chlorophyll fluorescence (excitation-emission 561-702/87 nm) against side scatter.

Acknowledgments

We appreciate the technical guidance of Dr. Shifra Ben-Dor, Dr. Merav Kedmi and Dr. Hadas Keren-Shaul in RNA-sequencing and are thankful for the help of Dr. Ron Rotkopf with statistical analysis, and of Dr. Alexander Brandis with LC-MS analysis (Life Sciences Core Facilities, Weizmann Institute of Science, Israel). We thank Dr. Shilo Rosenwasser (The Hebrew University of Jerusalem, Israel) for sharing his expertise in pulse amplitude-modulated fluorometry. We thank Prof. Robert Fluhr and Prof. Dan Yakir (Weizmann Institute of Science, Israel) for valuable comments during the study. We are grateful for Dr. Sheera Adar and Yuval Cohen (The Hebrew University of Jerusalem, Israel) for their insights into UV-induced DNA damage, and to Dr. Chana Kranzler (Bar Ilan University, Israel) for her technical and scientific support. Finally, we thank all members of the Segev lab for insightful discussions and input. O.E. received the Sustainability and Energy Research Initiative (SAERI) fellowship. The study was supported by funds received from the Minerva Foundation with funding from the German Federal Ministry for Education and Research, the Israel Science Foundation (ISF 947/18), the European Research Council (ERC StG 101075514) and the de Botton center for marine sciences, granted to E.S.

Authors Contributions

O.E. and E.S. designed the study. O.E., S.M. and I.P. performed and analyzed experiments. O.E., E. F. and M.S. performed computational analyses. O.E. and E.S. wrote the manuscript. All authors discussed the results and contributed to the final manuscript.

Competing interests

The authors declare no competing interests.

References

1. Havinga, E. Vitamin D, example and challenge. *Experientia* **29**, 1181–1193 (1973).

2. Holick, M. F. *et al.* Photosynthesis of Previtamin D3 in Human Skin and the Physiologic Consequences. *Science* **210**, 203–205 (1980).

3. Kalaras, M. D., Beelman, R. B., Holick, M. F. & Elias, R. J. Generation of potentially bioactive ergosterol-derived products following pulsed ultraviolet light exposure of mushrooms (Agaricus bisporus). *Food Chem.* **135**, 396–401 (2012).

4. Hossein-Nezhad, A. & Holick, M. F. Vitamin D for health: A global perspective. *Mayo Clin. Proc.* **88**, 720–755 (2013).

5. Fukumoto, S. Phosphate metabolism and vitamin D. *Bonekey Rep.* **3**, 497 (2014).

6. Veldurthy, V. *et al.* Vitamin D, calcium homeostasis and aging. *Bone Res.* **4**, 1–7 (2016).

7. Lopez Payares, G. M. & Ali, F. A. Vitamin D deficiency. *5-Minute Clin. Consult Stand. 2016 Twenty Fourth Ed.* 266–281 (2015).

8. Schwalfenberg, G. K. A review of the critical role of vitamin D in the functioning of the immune system and the clinical implications of vitamin D deficiency. *Mol. Nutr. Food Res.* **55**, 96–108 (2011).

9. Ljubic, A., Thulesen, E. T., Jacobsen, C. & Jakobsen, J. UVB exposure stimulates production of vitamin D3 in selected microalgae. *Algal Res.* **59**, 102472 (2021).

10. Brown, M. R., Mular, M., Miller, I., Farmer, C. & Trenerry, C. The vitamin content of microalgae used in aquaculture. *J. Appl. Phycol.* **11**, 247–255 (1999).

11. Holick, M. F., Holick, S. A. & Guillard, R. L. Photosynthesis of previtamin D in phytoplankton. *Curr. trends Comp. Endocrinol.* **2**, 1263–1266 (1982).

12. Jäpelt, R. B. & Jakobsen, J. Vitamin D in plants: a review of occurrence, analysis, and biosynthesis. *Front. Plant Sci.* **4**, 1–20 (2013).

13. Mattila, P. H., Piironen, V. I., Uusi-Rauva, E. J. & Koivistoinen, P. E. Vitamin D Contents in Edible Mushrooms. *J. Agric. Food Chem.* **42**, 2449–2453 (1994).

14. Hohman, E. E. *et al.* Bioavailability and Efficacy of Vitamin D2 from UV-Irradiated Yeast in Growing, Vitamin D-Deficient Rats. *J. Agric. Food Chem.* **59**, 2341–2346 (2011).

15. Wang, T., Bengtsson, G., Kärnefelt, I. & Björn, L. O. Provitamins and vitamins D2 and D3 in Cladina spp. over a latitudinal gradient: possible correlation with UV levels. *J. Photochem. Photobiol. B Biol.* **62**, 118–122 (2001).

16. Brocks, J. J. *et al.* The rise of algae in Cryogenian oceans and the emergence of animals. *Nature* **548**, 578–581 (2017).

17. Rao, D. S. & Raghuramulu, N. Food chain as origin of vitamin D in fish. *Comp. Biochem. Physiol. - A Physiol.* **114**, 15–19 (1996).

18. Fraser, D. R. *Evolutionary Biology: Mysteries of Vitamin D in Fish. Vitamin D: Fourth Edition* **1**, (Elsevier Inc., 2018).

19. Atsuko, T., Toshio, O., Makoto, T. & Tadashi, K. Possible origin of extremely high contents of

vitamin D3 in some kinds of fish liver. *Comp. Biochem. Physiol. Part A Physiol.* **100**, 483–487
(1991).

20. Bikle, D. D. Vitamin D: An ancient hormone. *Exp. Dermatol.* **20**, 7–13 (2011).

21. Rastogi, R. P., Richa, Kumar, A., Tyagi, M. B. & Sinha, R. P. Molecular Mechanisms of
Ultraviolet Radiation-Induced DNA Damage and Repair. *J. Nucleic Acids* **2010**, 592980 (2010).

22. de Jager, T. L., Cockrell, A. E. & Du Plessis, S. S. Ultraviolet Light Induced Generation of
Reactive Oxygen Species. in *Ultraviolet Light in Human Health, Diseases and Environment* 15–
23 (2017). doi:10.1007/978-3-319-56017-5_2

23. Czégény, G., Mátai, A. & Hideg, É. UV-B effects on leaves—Oxidative stress and acclimation in
controlled environments. *Plant Sci.* **248**, 57–63 (2016).

24. Tron, V. A. *et al.* p53-regulated apoptosis is differentiation dependent in ultraviolet B- irradiated
mouse keratinocytes. *Am. J. Pathol.* **153**, 579–585 (1998).

25. Kataria, S., Jajoo, A. & Guruprasad, K. N. Impact of increasing Ultraviolet-B (UV-B) radiation on
photosynthetic processes. *J. Photochem. Photobiol. B Biol.* **137**, 55–66 (2014).

26. Szilárd, A., Sass, L., Deák, Z. & Vass, I. The sensitivity of Photosystem II to damage by UV-B
radiation depends on the oxidation state of the water-splitting complex. *Biochim. Biophys. Acta -
Bioenerg.* **1767**, 876–882 (2007).

27. Morris, D. P. *et al.* The attenuation of solar UV radiation in lakes and the role of dissolved
organic carbon. *Limnol. Oceanogr.* **40**, 1381–1391 (1995).

28. Tedetti, M. & Sempéré, R. Penetration of Ultraviolet Radiation in the Marine Environment. A
Review. *Photochem. Photobiol.* **82**, 389 (2006).

29. Häder, D. P. Penetration and effects of solar UV-B on phytoplankton and macroalgae. in *UV-B
and Biosphere* 4–13 (1997). doi:10.1007/978-94-011-5718-6_1

30. Bendif, E. M. *et al.* Repeated species radiations in the recent evolution of the key marine
phytoplankton lineage Gephyrocapsa. *Nat. Commun.* **10**, 4234 (2019).

31. Taylor, A. R., Brownlee, C. & Wheeler, G. Coccolithophore Cell Biology: Chalking Up Progress.
Ann. Rev. Mar. Sci. **9**, 283–310 (2017).

32. Paasche, E. A review of the coccolithophorid emiliania huxleyi (prymnesiophyceae), with
particular reference to growth, coccolith formation, and calcification-photosynthesis interactions.
Phycologia **40**, 503–529 (2001).

33. Nanninga, H. J. & Tyrrell, T. Importance of light for the formation of algal blooms by Emiliania
huxleyi. *Mar. Ecol. Prog. Ser.* **136**, 195–203 (1996).

34. Volkman, J. K., Smith, D. J., Eglinton, G., Forsberg, T. E. V. & Corner, E. D. S. Sterol and fatty
acid composition of four marine haptophycean algae. *J. Mar. Biol. Assoc. United Kingdom* **61**,
509–527 (1981).

35. Eltgroth, M. L., Watwood, R. L. & Wolfe, G. V. Production and cellular localization of neutral long-
chain lipids in the haptophyte algae isochrysis galbana and emiliania huxleyi. *J. Phycol.* **41**,
1000–1009 (2005).

36. Sañé, E. *et al.* The Recent Advanced in Microalgal Phytosterols: Bioactive Ingredients Along
With Human-Health Driven Potential Applications. *Food Rev. Int.* **39**, 1859–1878 (2023).

37.   Jäpelt, R. B., Silvestro, D., Smedsgaard, J., Jensen, P. E. & Jakobsen, J. Quantification of
vitamin D3 and its hydroxylated metabolites in waxy leaf nightshade (Solanum glaucophyllum
Desf.), tomato (Solanum lycopersicum L.) and bell pepper (Capsicum annuum L.). *Food Chem.*
138, 1206–1211 (2013).

38.   Berridge, M. J. The inositol trisphosphate/calcium signaling pathway in health and disease.
Physiol. Rev. **96**, 1261–1296 (2016).

39.   Di Costanzo, F., Di Dato, V., Ianora, A. & Romano, G. Prostaglandins in marine organisms: A
review. *Mar. Drugs* **17**, (2019).

40.   Seltmann, M. A. *et al.* Differential Impact of Lipoxygenase 2 and Jasmonates on Natural and
Stress-Induced Senescence in Arabidopsis. *Plant Physiol.* **152**, 1940–1950 (2010).

41.   Kaye, Y. *et al.* Inositol Polyphosphate 5-Phosphatase7 Regulates the Production of Reactive
Oxygen Species and Salt Tolerance in Arabidopsis . *Plant Physiol.* **157**, 229–241 (2011).

42.   Salmon, J. A. & Higgs, G. A. Prostaglandins and leukotrienes as inflammatory mediators. *Br.*
Med. Bull. **43**, 285–296 (1987).

43.   Wan S., Y., Wang Q., Z., Shao, Y., Voorhees J., J. & Fisher J., G. Ultraviolet irradiation activates
PI 3-kinase/AKT survival pathway via EGF receptors in human skin in vivo. *Int J Oncol* **18**, 461–
466 (2001).

44.   Black, A. K., Greaves, M. W., Hensby, C. N. & Plummer, N. A. Increased prostaglandins E2 and
F2alpha in human skin at 6 and 24 h after ultraviolet B irradiation (290- 320 nm). *Br. J. Clin.*
Pharmacol. **5**, 431–436 (1978).

45.   Moreno, J. *et al.* Regulation of prostaglandin metabolism by calcitriol attenuates growth
stimulation in prostate cancer cells. *Cancer Res.* **65**, 7917–7925 (2005).

46.   Trechsel, U., Carol M. Taylor, Jean-Philippe Bonjour & Fleisch, H. Influence of prostaglandins
and of cyclic nucleotides on the metabolism of 25-hydroxyvitamin D3 in primary chick kidney cell
culture. *Biochem. Biophys. Res. Commun.* **93**, 15–38 (1980).

47.   Tang, W., Ziboh, V. A., Isseroff, R. R. & Martinez, D. Novel regulatory actions of 1α,25-
dihydroxyvitamin D3 on the metabolism of polyphosphoinositides in murine epidermal
keratinocytes. *J. Cell. Physiol.* **132**, 131–136 (1987).

48.   De Boland, A. R., Facchinetti, M. M., Balogh, G., Massheimer, V. & Boland, R. L. Age-associated
decrease in inositol 1,4,5-trisphosphate and diacylglycerol generation by 1,25(OH)2-vitamin D3
in rat intestine. *Cell. Signal.* **8**, 153–157 (1996).

49.   Bourdeau, A., Atmani, F., Grosse, B. & Lieberherr, M. Rapid effects of 1, 25-dihydroxyvitamin d3
and extracellular ca2+ on phospholipid metabolism in dispersed porcine parathyroid cells.
Endocrinology **127**, 2738–2743 (1990).

50.   Wang, P.-F., Yao, D.-H., Hu, Y.-Y. & Li, Y. Vitamin D Improves Intestinal Barrier Function in
Cirrhosis Rats by Upregulating Heme Oxygenase-1 Expression. *Biomol. Ther. (Seoul).* **27**, 222–
230 (2019).

51.   Wu, X. *et al.* Vitamin D–vitamin D receptor alleviates oxidative stress in ischemic acute kidney
injury via upregulating glutathione peroxidase 3. *FASEB J.* **37**, e22738 (2023).

52.   Pavlovitch, J. H., Rizk, M. & Balsan, S. Vitamin D nutrition increases skin tyrosinase response to
exposure to ultraviolet radiation. *Mol. Cell. Endocrinol.* **25**, 295–302 (1982).

53.   Gordon-Thomson, C. *et al.* 1α,25 Dihydroxyvitamin D3 enhances cellular defences against UV-
induced oxidative and other forms of DNA damage in skin. *Photochemical and Photobiological*
Sciences **11**, 1837–1847 (2012).

54.   Wong, C. T. & Oh, D. H. Vitamin D Receptor Promotes Global Nucleotide Excision Repair by
Facilitating XPC Dissociation from Damaged DNA. *J. Invest. Dermatol.* **141**, 1656–1663 (2021).

55.   Pinnola, A. *et al.* Light-Harvesting Complex Stress-Related Proteins Catalyze Excess Energy
Dissipation in Both Photosystems of Physcomitrella patens. *Plant Cell* **27**, 3213–3227 (2015).

56.   Peers, G. *et al.* An ancient light-harvesting protein is critical for the regulation of algal
photosynthesis. *Nature* **462**, 518–521 (2009).

57.   Tilbrook, K. *et al.* UV-B perception and acclimation in chlamydomonas reinhardtii. *Plant Cell* **28**,
966–983 (2016).

58.   Yannarelli, G. G., Noriega, G. O., Batlle, A. & Tomaro, M. L. Heme oxygenase up-regulation in
ultraviolet-B irradiated soybean plants involves reactive oxygen species. *Planta* **224**, 1154–1162
(2006).

59.   Cui, W. *et al.* Haem oxygenase-1 is involved in salicylic acid-induced alleviation of oxidative
stress due to cadmium stress in Medicago sativa. *J. Exp. Bot.* **63**, 5521–5534 (2012).

60.   Applegate, L. A., Luscher, P. & Tyrrell, R. M. Induction of Heme Oxygenase: A General
Response to Oxidant Stress in Cultured Mammalian Cells1. *Cancer Res.* **51**, 974–978 (1991).

61.   Harris, G. N., Scanlan, D. J. & Geider, R. J. Acclimation of Emiliania huxleyi (Prymnesiophyceae)
to photon flux density. *J. Phycol.* **41**, 851–862 (2005).

62.   Xu, J. *et al.* The role of coccoliths in protecting Emiliania huxleyi against stressful light and UV
radiation. 4637–4643 (2016). doi:10.5194/bg-13-4637-2016

63.   Ruban, A. V. Nonphotochemical chlorophyll fluorescence quenching: Mechanism and
effectiveness in protecting plants from photodamage. *Plant Physiol.* **170**, 1903–1916 (2016).

64.   Genty, B., Briantais, J.-M. & Baker, N. R. The relationship between the quantum yield of
photosynthetic electron transport and quenching of chlorophyll fluorescence. *Biochim. Biophys.*
Acta - Gen. Subj. **990**, 87–92 (1989).

65.   Vass, I., Cser, K. & Cheregi, O. Molecular Mechanisms of Light Stress of Photosynthesis. *Ann.*
N. Y. Acad. Sci. **1113**, 114–122 (2007).

66.   Carlson, D. F., Fredj, E. & Gildor, H. The annual cycle of vertical mixing and restratification in the
Northern Gulf of Eilat/Aqaba (Red Sea) based on high temporal and vertical resolution
observations. *Deep Sea Res. Part I Oceanogr. Res. Pap.* **84**, 1–17 (2014).

67.   Chiswell, S., Calil, P. & Boyd, P. Spring blooms and annual cycles of phytoplankton: A unified
perspective. *J. Plankton Res.* **37**, (2015).

68.   Weller, R. A. & Price, J. F. Langmuir circulation within the oceanic mixed layer. *Deep Sea Res.*
Part A. Oceanogr. Res. Pap. **35**, 711–747 (1988).

69.   MacLaughlin, J. A., Anderson, R. R. & Holick, M. F. Spectral Character of Sunlight Modulates
Photosynthesis of Previtamin D3 and Its Photoisomers in Human Skin. *Science* **216**, 1001–1003
(1982).

70.   Pollara, S. B. *et al.* Bacterial Quorum-Sensing Signal Arrests Phytoplankton Cell Division and

Impacts Virus-Induced Mortality. *mSphere* **6**, (2021).

71.   Pavey, S., Russell, T. & Gabrielli, B. G2 phase cell cycle arrest in human skin following UV irradiation. *Oncogene* **20**, 6103–6110 (2001).

72.   Jiang, L., Wang, Y., Björn, L. O. & Li, S. Does cell cycle arrest occur in plant under solar UV-B radiation? *Plant Signal. \& Behav.* **6**, 892–894 (2011).

73.   Rudolph, C. J., Upton, A. L. & Lloyd, R. G. Replication fork stalling and cell cycle arrest in UV-irradiated Escherichia coli. *Genes and Development* **21**, 668–681 (2007).

74.   Özerkan, D., Özsoy, N., Akbulut, K. G., Güney, Ş. & Öztürk, G. The protective effect of vitamin D against carbon tetrachloride damage to the rat liver. *Biotech. Histochem.* **92**, 513–523 (2017).

75.   Hii, C. S. & Ferrante, A. The Non-Genomic Actions of Vitamin D. *Nutrients* **8**, 135 (2016).

76.   Flori, S. *et al.* Diatoms exhibit dynamic chloroplast calcium signals in response to high light and oxidative stress. *bioRxiv* 2008–2023 (2023).

77.   Oberson, J. M., Bénet, S., Redeuil, K. & Campos-Giménez, E. Quantitative analysis of vitamin D and its main metabolites in human milk by supercritical fluid chromatography coupled to tandem mass spectrometry. *Anal. Bioanal. Chem.* **412**, 365–375 (2020).

78.   Brocks, J. J. *et al.* Lost world of complex life and the late rise of the eukaryotic crown. *Nature* **618**, 767–773 (2023).

79.   Desmond, E. & Gribaldo, S. Phylogenomics of Sterol Synthesis: Insights into the Origin, Evolution, and Diversity of a Key Eukaryotic Feature. *Genome Biol. Evol.* **1**, 364–381 (2009).

80.   Carlberg, C. Vitamin D in the Context of Evolution. *Nutrients* **14**, (2022).

81.   Waldbauer, J. R., Newman, D. K. & Summons, R. E. Microaerobic steroid biosynthesis and the molecular fossil record of Archean life. *Proceedings of the National Academy of Sciences of the United States of America* **108**, 13409–13414 (2011).

82.   Knoll, A. H., Javaux, E. J., Hewitt, D. & Cohen, P. Eukaryotic organisms in Proterozoic oceans. *Philos. Trans. R. Soc. B Biol. Sci.* **361**, 1023–1038 (2006).

83.   Han, T. M. & Runnegar, B. Megascopic Eukaryotic Algae from the 2.1-Billion-Year-Old Negaunee Iron-Formation, Michigan. *Science* **257**, 232–235 (1992).

84.   Canfield, D. E. A new model for Proterozoic ocean chemistry. *Nature* **396**, 450–453 (1998).

85.   Lyons, T. W., Reinhard, C. T. & Planavsky, N. J. The rise of oxygen in Earth's early ocean and atmosphere. *Nature* **506**, 307–315 (2014).

86.   Anbar, A. D. & Knoll, A. H. Proterozoic ocean chemistry and evolution: A bioinorganic bridge? *Science* **297**, 1137–1142 (2002).

87.   Smith, L. L. Another cholesterol hypothesis: cholesterol as antioxidant. *Free Radic. Biol. Med.* **11**, 47–61 (1991).

88.   Galea, A. M. & Brown, A. J. Special relationship between sterols and oxygen: were sterols an adaptation to aerobic life? *Free Radic. Biol. Med.* **47**, 880–889 (2009).

89.   Goyet, C. & Poisson, A. New determination of carbonic acid dissociation constants in seawater as a function of temperature and salinity. *Deep Sea Res. Part A. Oceanogr. Res. Pap.* **36**, 1635–1654 (1989).

655   90.   Guillard, R. R. L. & Hargraves, P. E. Stichochrysis immobilis is a diatom, not a chrysophyte.
656   *Phycologia* **32**, 234–236 (1993).

657   91.   Keren-Shaul, H. *et al.* MARS-seq2.0: an experimental and analytical pipeline for indexed sorting
658   combined with single-cell RNA sequencing. *Nat. Protoc.* **14**, 1841–1862 (2019).

659   92.   Kohen, R. *et al.* UTAP: User-friendly Transcriptome Analysis Pipeline. *BMC Bioinformatics* **20**,
660   154 (2019).

661   93.   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for
662   RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

663   94.   Sperfeld, M., Segev, E. & Dayana Yahalomi. Resolving the microalgal gene landscape at the
664   strain level: A novel hybrid transcriptome of Emiliania huxleyi CCMP3266. *Appl. Environ.*
665   *Microbiol.* (2021). doi:10.1128/AEM.01418-21

666   95.   Read, B. A. *et al.* Pan genome of the phytoplankton Emiliania underpins its global distribution.
667   *Nature* **499**, 209–213 (2013).

668   96.   Jones, P. *et al.* InterProScan 5: genome-scale protein function classification. *Bioinformatics* **30**,
669   1236–1240 (2014).

670   97.   Sayers, E. W. *et al.* Database resources of the national center for biotechnology information.
671   *Nucleic Acids Res.* **50**, D20–D26 (2022).

672   98.   Vandesompele, J. *et al.* Accurate normalization of real-time quantitative RT-PCR data by
673   geometric averaging of multiple internal control genes. *Genome Biol.* **3**, research0034.1 (2002).

Figure S1. Continuous UV irradiation during the light period did not affect algal growth. Growth curves of the algal cultures that were grown under control and UV regime and were used for generating the transcriptomic data. Cell density was measured from day 7. Error bars indicate standard deviation based on 3 biological replicates.

Figure S2. UV radiation causes upregulation of various signaling and stress-response mechanisms. qRT-PCR analysis of genes under 1 hour of UV exposure or vitamin D treatments. Top title denotes gene products. In brackets: gene identifier in *E. huxleyi*; matching gene in *E. huxleyi* CCMP1516 reference genome[95]. Statistical significance compared to control was calculated using two-tailed t-test assuming equal variances. One, two or three asterisks indicate $P < 0.05$, $P < 0.01$ and $P < 0.001$, respectively.

Figure S3. The combined treatment of vitamin D₂ and D₃ has a synergistic effect on algal growth, outperforming individual D₂ or D₃ application. (a) Growth dynamics of algal cultures treated with 1μM of vitamin D species. (b) Variation in algal cell densities after 4 days of treatment with different concentrations of vitamin D₂, D₃, and both (indicated by the green shades in the figure legend). Error bars indicate standard deviation from 3 biological replicates.

Figure S4. Temporal dynamics in NPQ and Fm' values of vitamin D-treated algae, exposed to regular or excess light. Temporal dynamics in Fm' and NPQ of both control and vitamin D-treated algae exposed to regular light levels (130 PAR) or excess light levels (1000 PAR) for 2 hours prior to analysis. The timing of saturating pulses is represented by dots. NPQ values were measurable from t=60s onwards. Each line represents a single biological replicate.

Figure S5. *E. huxleyi* algae produce putative hydroxylated vitamin D species. Representative chromatograms of hydroxylated vitamin D species detected in UV-treated and control cultures. The range of the Y axis in chromatograms of the same vitamin D species are identical and therefore comparable. The identification of metabolites is putative according to MRMs taken from Oberson et al.[77]. Further analysis against standards is needed in order to verify the identification of the hydroxylated species.

Figure S6. Dark adaptation of 5 minutes sufficiently relaxes PSII reaction centers. Comparison of maximum PSII quantum yields (F_v/F_m) between algae that were dark-adapted for 5 or 30 minutes, calculated based on three biological replicates.

Gene	Forward	Reverse
G6284	GCCCTACCGGGTGTTTATCC	CTCGACTTGGTTGAGAACTTGC
G18115	CACCGAGCCGACCCAAAAT	TTCATCTCGGTCGTTGACAGG
G18590	CTCCTCGCGATGCAGAACAA	CCCTTTCTGCCAACGTGATCT
G25467	TACGAGAATCGGCTGCTACG	CCCGTCGGACCTTAAGACAG
G12503	TCTCGGTGGAAATGGCGAC	ATTGTCATCGAGGCCGCAAA
G93	GTCACGCCGGCGACAAA	GCGATGTGCGGGTGTATCT
G12340	AACCTGCTTGCCGACATGAT	GGTTGAATCAGCATTGACCCC
G14992	GCGGGCTCTACTGAATCCG	GGGTCCTCGTAGAAGGTGTG
G16746	TCGAAGATCCGGACGACGAT	AGCGCGAGACGAAAATAACG
G2511	TTGTCGCGTCGCTCTACTTT	GCCGAAGTAGTACGCCATGT
G2758	ATGGACCTAGACTCGGACGG	ACAGCCCTCAAGCTCACATC
G8907	CTCTCGTGCTCGTGCTTCTT	TCGTAGATGTACTTTGCCGGG
G26534	CTGGAAGATCGAGGCAACGG	TATGGCGTCGCCGTCAAAG
G26797 *alpha-tubulin*	CGAGAAGGCGTACCACGAG	CTTCGTCTTGATGGTGGCGA
G28192 *beta-tubulin*	CAACATGAAGTGCGCCATCT	CCTCGGTGAACTCCATCTCG
G1895 *rpl13*	ACCAGCACTTCCACAAGACG	TGCCGCAGCTTGTAGTTGTA

Table S2. Primers used in this study. Gene identifiers in *E. huxleyi* CCMP3266[94]. Primers were used to generate qRT-PCR data using *E. huxleyi* CCMP3266 cDNA.