The gene expression signature of electrical stimulation in the human brain

Snehajyoti Chatterjee[1,2], Yann Vanrobaeys[1#], Annie I Gleason[3#], Brian J. Park[4], Shane A. Heiney[1,2,5], Ariane E. Rhone[4], Kirill V. Nourski[4], Lucy Langmack[1,2], Budhaditya Basu[1,2], Utsav Mukherjee[1,2], Christopher K. Kovach[4,6], Zsuzsanna Kocsis[4,7], Yukiko Kikuchi[7], Yaneri A. Ayala[4], Christopher I. Petkov[4,7], Marco M. Hefti[8], Ethan Bahl[3], Jacob J Michaelson[3], Hiroto Kawasaki[4], Hiroyuki Oya[4], Matthew A. Howard III[4], Thomas Nickl-Jockschat[1,2,3,9,10,11], Li-Chun Lin[1,2,12], Ted Abel[1,2,3*]

Affiliations:

[1]Department of Neuroscience and Pharmacology, Iowa Neuroscience Institute, Carver College of Medicine, University of Iowa, Iowa City, IA, USA

[2]Iowa Neuroscience Institute, University of Iowa, Iowa City, IA, USA

[3]Department of Psychiatry, University of Iowa Hospitals and Clinics, Iowa City, IA, USA

[4]Department of Neurosurgery, University of Iowa Hospitals and Clinics, Iowa City, IA, USA

[5]Neural Circuits and Behavior Core, Iowa Neuroscience Institute Carver College of Medicine, University of Iowa, Iowa City, IA, USA

[6]Department of Neurosurgery, University of Nebraska Medical Center, Omaha, NE, USA

[7]Biosciences Institute, Newcastle University Medical School, Newcastle upon Tyne, UK

[8]Department of Pathology, University of Iowa Hospitals and Clinics, Iowa City, IA, USA

[9]Department of Psychiatry and Psychotherapy, Otto-von-Guericke University, Magdeburg, Germany

[10]German Center for Mental Health (DZPG), partner site Halle-Jena-Magdeburg, Germany

[11]Center for Intervention and Research on adaptive and maladaptive brain Circuits underlying mental health (C-I-R-C), Halle-Jena-Magdeburg, Germany

[12]Iowa NeuroBank Core, Iowa Neuroscience Institute, Carver College of Medicine, University of Iowa, Iowa City, IA, USA

[#]Authors contributed equally

*Corresponding author:

ted-abel@uiowa.edu

Keywords: Microglia, anterior temporal neocortex, direct electrical stimulation, single
nuclei multiomics

Abstract

Direct electrical stimulation has been used for decades as a gold standard clinical tool
to map cognitive function in neurosurgery patients[1-8]. However, the molecular impact of
electrical stimulation in the human brain is unknown. Here, using state-of-the-art
transcriptomic and epigenomic sequencing techniques, we define the molecular
changes in bulk tissue and at the single-cell level in the human cerebral cortex following
direct electrical stimulation of the anterior temporal lobe in patients undergoing
neurosurgery. Direct electrical stimulation surprisingly had a robust and consistent
impact on the expression of genes related to microglia-specific cytokine activity, an
effect that was replicated in mice. Using a newly developed deep learning
computational tool, we further demonstrate cell type-specific molecular activation, which
underscores the effects of electrical stimulation on gene expression in microglia. Taken
together, this work challenges the notion that the immediate impact of electrical
stimulation commonly used in the clinic has a primary effect on neuronal gene
expression and reveals that microglia robustly respond to electrical stimulation, thus
enabling these non-neuronal cells to sculpt and shape the activity of neuronal circuits in
the human brain.

Main

Electrical stimulation of the human brain has become an indispensable clinical tool for diagnosis and therapy[1-8]. In neurosurgical patients, cortical stimulation is the gold standard diagnostic tool to identify the location of functionally relevant brain regions that are critical for speech, language, or motor function[9,10] to avoid potential damage to these regions during surgery. Transient changes in gene expression underlie the ability of the brain to adapt to changes in the environment or brain perturbation, such as electrical stimulation. Dynamic transcriptomic patterns are essential for cognition[11-13], affective processing[14], addiction[15], and the initiation of behaviors[16,17]. Thus, electrical stimulation likely exerts neural effects via alterations in gene expression. Although diagnostic and therapeutic brain stimulation is conducted daily in thousands of patients worldwide, the molecular impact of electrical stimulation in the human brain remains unknown. Recent advances in molecular sequencing techniques have revolutionized transcriptomics by enabling mapping of changes in transcription and chromatin accessibility in single cells[18]. However, these experiments have only been conducted in rodents[19,20] and neurons derived from human induced pluripotent stem cells[21]. A recent pioneering study utilized a single nuclei molecular approach to link human brain transcriptomics signatures with oscillatory signatures of memory consolidation in epileptic patients undergoing an episodic memory task, but these gene expression measures were obtained many days after assessment of intracranial recording data during the task[22,23]. Therefore, research is needed to evaluate the more immediate effects on single-cell early gene expression, such as tens of minutes after neural system perturbation with electrical stimulation. Studies investigating gene expression profiles after electrical stimulation are particularly difficult to perform in human brains due to the requirement that tissue is sampled at precise temporal windows before and after stimulation.

Here, in patients undergoing clinical neurosurgical resection of pathological epileptogenic sites in the mesial temporal lobe, samples from tissue in the anterior

temporal lobe (ATL) that required clinical resection to access the deeper epileptogenic

site for treatment were obtained before and after electrical stimulation. By analyzing

gene expression at baseline and minutes after stimulation from the same patient, along

with our analysis of samples taken from patients that did not receive electrical

stimulation, we are able to distinguish genes responsive to electrical stimulation from

the genes altered nonspecifically by surgical or disease-based factors that would be

stable across samples. Our parameters for electrical stimulation were based on those

regularly used in clinical mapping[24,25]; moreover, we referenced the human results to a

similar electrical stimulation paradigm in mice, revealing a similar transcriptional profile.

Finally, using a single-nuclei multi-omics approach, we provide insight into the cell-type-

specific transcriptomic and epigenomic responses to electrical stimulation in the human

brain, highlighting important effects beyond neurons, including microglia. This study

provides fundamental insights into changes in cell-type specific molecular signatures in

the human cortex after electrical stimulation, laying the groundwork for a molecular

understanding of the impact of this fundamental tool in clinical neurosurgery, diagnosis,

and treatment.

Results

We recruited eight adult neurosurgical patients undergoing surgical resection of seizure

foci following clinical monitoring to treat epilepsy. The patients provided informed

consent to take part in this research and were informed that tissue samples would only

be taken from tissue that would require clinically resection for treatment. The patient

participants underwent an anterior temporal lobectomy for access to a medial temporal

lobe epileptogenic site, during which samples were resected from the neocortex and

processed immediately after removal. The participants were evenly distributed between

two experimental paradigms. In the first group with the electrically stimulated paradigm

(4 participants), a baseline tissue sample was resected from the ATL after the

craniotomy and durotomy, exposing the temporal lobe. Then, an adjacent region of the

cortex was stimulated using a stimulation protocol commonly used for bipolar electrical

stimulation (50 Hz) for 2 minutes[24,25] And resected approximately thirty minutes later

(32.5 ± 11.2 min), a sample was taken from the stimulated region **(Fig. 1a;**

Supplementary Fig. 1a, Supplementary data 1). In the second group with the

unstimulated paradigm (4 participants), a sample was taken at baseline, and then a

second sample was taken without stimulation about 30 minutes later (37.3 ± 12.5 mins,

Fig. 1b; Supplementary Fig. 1a). The pre-stimulation sample and the unstimulated

paradigm groups serves as controls for changes in baseline gene expression and

changes in gene expression occurring as a result of the surgical procedure. Among the

8 participants, five were under general anesthesia, whereas three participants were

awake during the surgical procedure. All the resected tissue samples from both the

stimulated and unstimulated paradigms were taken well outside the seizure focus (27.8

± 7mm from the border of seizure focus**, Supplementary Fig. 1b**).

Tissue samples were first subjected to bulk whole-transcriptome RNA sequencing

(RNA-seq) to identify differentially expressed genes following electrical stimulation. Bulk

RNA-seq analysis from the stimulated and corresponding baseline samples revealed

124 differentially expressed genes following electrical stimulation, with 112 up-regulated

and 12 down-regulated genes **(Fig. 1c**; **Supplementary data 2)**. Enrichment network

analysis was used to identify the pathways most represented among the differentially

expressed genes in the stimulated paradigm. The top significant pathways were

enriched with genes involved in cytokine activity, DNA-binding transcription activator

activity (RNA Pol II), cytokine receptor binding, DNA-binding transcription activator

activity, nuclear glucocorticoid receptor binding, chemokine activity, CCR chemokine

receptor binding, RNA pol II specific DNA binding, chemokine receptor binding, and

protein phosphatase activity **(Fig. 1d).** Notably, these genes were not significantly

enriched for previously identified genes induced in regions showing seizure activity in

the human brain **(Supplementary Fig. 2)**[26].

Bulk RNA-seq analysis from the unstimulated paradigm comparing unstimulated

samples with corresponding baseline samples identified differential expression of only

16 genes, with nine up-regulated and seven down-regulated genes **(Fig. 1e,**

Supplementary data 3). Only one gene, *NR4A3*, was found to be differentially

expressed in both the stimulated and unstimulated groups **(Fig. 1g).** The lack of overlap

between the differentially expressed genes observed in our stimulated and unstimulated

groups suggests that the changes in gene expression that we see following electrical

stimulation do not reflect disease state or surgical effects such as craniotomy, brain
temperature, and anesthesia, but reflect changes due to electrical stimulation.

We used a similar electrical stimulation paradigm in the mouse non-primary auditory
cortex to investigate whether these gene expression changes are unique to human
samples and to determine whether they are related to disease state [26,27]. Bulk RNA-seq
was performed from samples collected 30 minutes following electrical stimulation and
unstimulated samples collected at the same time from the contralateral side (**Fig. 2a**).
Bulk RNA-seq identified 44 upregulated and 107 downregulated genes (**Fig. 2b,**
Supplementary data 4). Pathway analysis identified from the upregulated genes were
enriched for cytokine activity, chemokine activity, cytokine receptor binding, and CCR
chemokine receptor binding. (**Fig. 2c**). We also found upregulated pathways linked to
transcription and post-transcription regulatory pathways, such as mRNA 3'-UTR binding,
mRNA3'-UTR AU-rich region binding, and transcription co-repressor activity. Comparing
the stimulation-responsive genes from human bulk RNA-seq and mouse bulk RNA-seq
revealed a significant correlation (R^2=0.0415, p-value <0.00001), especially cytokine-
related genes were commonly upregulated following electrical stimulation in both
humans and mice (**Fig. 2d**). Some of the pathways commonly altered by electrical
stimulation between humans and mice are cytokine activity, chemokine activity,
chemokine receptor binding, cytokine receptor activity, and the CCR chemokine
receptor binding. Some common genes enriched in these chemokine and cytokine-
related pathways are *CCL3*, *CCL4*, *CXCL1*, *IL1A*, and *TNF*. We further validated the
expression of the cytokine activity-related genes *Ccl3* and *Ccl4* using qPCR analysis
from mouse brain following electrical stimulation (**Fig. 2e**). The differentially expressed
genes following electrical stimulation in the human cortex correlated significantly with
learning-induced genes in the mouse cortex (R^2=0.2027, p-value <0.00001). Immediate
early genes such as *Arc*, *Fos*, *Egr1*, *Nr4a1*, and *FosB* were upregulated in both
datasets **(Supplementary Fig. 3)**. However, learning did not induce expression of
cytokine activity related genes in mouse cortex. Thus, our findings of gene expression
signature in response to electrical stimulation in the human and mouse brain reveal a
molecular signature that partially conserved across species.

Next, we investigated the cell types exhibiting differential gene expression following
electrical stimulation in the human brain by utilizing single nuclei multiomics (RNA and
ATAC) on samples from the stimulated paradigm from 3 participants (**Fig 3a,**
Supplementary Fig. 4). Cell clustering analysis identified seven major cell types in our
samples—excitatory neurons, microglia, VIP-Sncg-Lamp5 inhibitory neurons, PValb-Sst
inhibitory neurons, oligodendrocytes, oligodendrocytes precursor cells (OPC), and
astrocytes (**Fig. 3b**). To determine the differentially expressed genes in each cell type
following electrical stimulation, we performed a pseudobulk RNA-seq analysis due to its
superior performance for detecting differential expression in single-cell RNA-sequencing
analyses [28] and the stringent nature of this analysis. The pseudobulk analysis revealed
that the microglia displayed the highest differential gene expression of all cell types with
31 upregulated genes (**Fig. 3c, Supplementary data 5**). Genes related to cytokine
activity (*CCL3*, *CCL4*, *CCL3L1*, and *IL1B*,) were upregulated exclusively within
microglia. Five genes were upregulated in oligodendrocytes (*FOS*, *HSPA1A*, *JUNB*,
GADD45B, and *FOSB*), and only two genes were differentially expressed (one
upregulated: *LHFPL3*, and one downregulated: *ROBO2*) in astrocytes (**Supplementary**
data 5). Surprisingly, no differentially expressed genes were detected within the
neuronal cell types (excitatory and inhibitory neuronal cell types). Comparing the
upregulated genes in microglia with human and mice bulk RNA seq revealed 5 genes
(*Fos*, *Dusp1*, *Ccl3*, *Ccl4*, *Zfp36*) that are commonly upregulated following electrical
stimulation (**Fig. 3d**). *Atf3*, *Cd83*, *Egr3*, *Nr4a1*, *Il1b*, *Mcl1*, *Nedd9, Spp1*, *Nfkbid*, *Rgs1*,
and *Ccl3l1* were found to be human microglia-specific genes induced by electrical
stimulation that do not change in the mice cortex (**Fig 3d**). Pathways enriched among
the upregulated genes in microglia included cytokine-related pathways such as cytokine
activity, chemokine activity, CCR chemokine receptor, and cytokine receptor binding
(**Fig. 3e**). Microglia exhibited activation of transcription regulatory pathways such as
DNA-binding transcription activator activity and RNA polymerase-specific DNA binding
TF binding (**Fig. 3e**). These results reveal that electrical stimulation increases the
expression of genes that are involved in cytokine activity and transcription regulation in
microglia.

To investigate if electrical stimulation has an impact on the epigenome, we assessed

chromatin accessibility using the single nuclear assay for transposase-accessible

chromatin with sequencing (snATAC-seq)[29] on human samples from the stimulated

paradigm. We focused on promoter accessibility (-2 to +2 kb from TSS) to analyze

changes in chromatin accessibility in snATAC-seq data (**Supplementary data 6**). The

differentially accessible regions that were either enriched or depleted following electrical

stimulation in microglia significantly correlated (R^2=0.0262, p-value<0.002) with the

microglia gene expression from snRNA-seq (**Fig. 3f**), suggesting that the genes that

exhibit induced expression following electrical stimulation also exhibit increased

chromatin accessibility. Genes related to transcription regulators, such as NR4A1 and

FOS, showed a positive correlation between transcriptome and chromatin accessibility.

We also found enriched promoter accessibility and increased gene expression for CCL4

in microglia following electrical stimulation (**Fig 3g**). A recent study demonstrated that

ambient RNA from neurons may contaminate non-neuronal cells in single-nuclei

transcriptomic data[30]. Our snRNA seq data revealed the upregulation of several IEGs,

such as *FOS*, *EGR3*, *JUNB,* and *NR4A1* within microglia. These genes are often seen

to be upregulated in neurons following neuronal activation. However, our snATAC-seq

data from the same samples also showed increased chromatin accessibility in microglial

populations within IEG promoters, thus suggesting that the microglia-specific IEG

expression is not due to neuronal ambient RNA contamination. Lastly, we analyzed

genome-wide transcription factor (TF) binding motifs from the upregulated peaks

following electrical stimulation in microglia using chromatin accessibility data.

Genomewide TF motif analysis revealed enrichment of binding motifs for ELK3, ELK1,

YY2, NRF1, HINFP, and ELK4 (**Fig. 3h, Supplementary data 7**). Interestingly, ELK4, a

transcription factor downstream of MAPK signaling, is predicted to positively regulate

the expression of *Ccl3* and *Ccl4*[31]. Thus, our sn-ATAC-seq results reveal a signature of

transcription factor motifs and chromatin accessibility underlying electrical stimulation-

induced gene expression in human microglia.

Our experimental design included internal controls (baseline/unstimulated samples)

from each participant, enabling us to investigate the impact of electrical stimulation

across individual participants. Therefore, we next examined the cell type-specific

response to electrical stimulation from within participants using our newly developed
deep-learning computational tool, NEUROeSTIMator[32], to map the active population of
cells following electrical stimulation in each participant, comparing stimulated samples
with corresponding baseline controls. We estimated cell activity from the single nuclei
RNA data from individual populations of cells with either baseline or stimulated from
each participants (**Fig. 4a**). We used paired linear regression with bootstrap sampling to
assess differences in estimated activity between conditions for each cell type (**Fig. 4b**).
Interestingly, we observed that the microglial cluster showed a significant increase in
activity scores following stimulation, and this effect was consistent across all three
participants [β = 0.100301, p-value < 2e-16] (**Fig. 4c**). This data supports our bulk RNA
seq and snRNAseq data. However, we observed variability in electrical stimulation-
mediated activation of excitatory neurons across participants. We detected significant
differences in activity when using all three participants [β = -0.053920, p-value < 2e-16].
One pair of samples showed a significant increase in activity [β = 0.032528, p-value =
0.000807], while 2 out of 3 pairs showed a significant decrease in activity [β = -
0.022086, -0.083254, p-values = 1.58e-8, 4e-9] (**Fig. 4c**). Variability in activity estimates
was also observed in inhibitory neuronal sub-types across participants (**Fig. 4c**).

To further investigate the epigenomic signature of cellular activity following electrical
stimulation, we estimated activity from snATAC-seq data using a similar approach (**Fig.
5a**). Activity was estimated for cells using gene-level counts in promoter regions and
paired linear regression with bootstrap sampling was used to assess activity between
baseline and electrical stimulation (**Fig. 5b**). Consistent with the transcriptomics data,
microglia showed a significant increase in promoter-informed activity estimates, and this
effect was consistent across all three participants [β = 0.060977, p-value < 2e-16] (**Fig.
5c**). Neuronal clusters showed variability in activation in activity when considering all
three participants [$\beta_{Excitatory}$ = 0.016163, p-value$_{Excitatory}$ = 0.000247; $\beta_{Inhibitory}$ = 0.028302,
p-value$_{Inhibitory}$ = 2.283e-7]. Only one participant showed a significant increase in the
excitatory[β = 0.04712, p-val = 0.000314] and inhibitory neurons (Sst-Pvalb) [β =
0.08863, p-value = 0.00201]. These findings suggest that microglia exhibit robust
transcriptional and epigenomic responses to electrical stimulation and that neuronal cell
types exhibit greater variability between patient participants.

Discussion

Electrical stimulation is a model for studying the human cortex with protocols that are used clinically for mapping the function of specific brain regions. Here, we take a molecular approach, defining the transcriptomic and epigenomic signatures of electrical stimulation at the single cell level in the human cortex in neurosurgical patient participants. Researchers have correlated human brain transcriptomics with prior recorded oscillatory signatures of memory consolidation[22,33], but have not previously examined the changes in gene expression that are rapidly and directly driven by electrical stimulation. Most work thus far has emphasized the role of neurons in responding to external stimuli, but our work reveals the critical role of microglia in sculpting the activity and function of brain circuits. In microglia, genes related to cytokine signaling showed the greatest induction pattern, and this was conserved across species. As previously hypothesized, we observed the induction of activity-dependent genes in our bulk RNA sequencing results, but to our surprise, single nuclei multiomics experiments revealed pronounced microglia-specific transcriptomic activation following electrical stimulation that was supported by analysis using NEUROeSTIMator[32], a deep-learning computational model. Identifying a microglial transcriptomic response following electrical stimulation represents an important conceptual advance in our understanding of the impact of this common form of electrical stimulation used clinically.

Microglia are critical modulators of neuronal function, acting to suppress excessive activity by inhibiting surrounding neurons, including excitatory neurons[34]. Although classified as non-excitable cells, microglia exhibit electrophysiological stimulus-response features, and changes in their membrane potential affect crucial microglial functions including phagocytosis, chemotaxis, and cytokine release[35]. Our analyses identified critical molecular components within microglia that change with electrical stimulation, including chemokine-encoding genes such as *CCL3* and *CCL4*, which act to alter microglial motility, influence neuronal-microglial interactions, and shape neuronal connectivity[34,36-39]. These chemokines are ligands for C-C chemokine receptor type 5 (CCR5), which regulates neuronal excitability and memory allocation in the hippocampus[40]. Previous rodent studies have shown that neuronal activation using

chemogenetic approaches leads to distinct gene expression changes in microglia,
including genes encoding chemokines[34]. Our work also revealed that genes encoding
transcription factors such as FOS and NR4A1 are induced in microglia in response to
electrical stimulation. Further, we observed increased promoter accessibility at these
genes, underscoring the conclusion that these genes are transcriptionally upregulated in
microglia after stimulation and broadening the potential functional impact of these
immediate early genes beyond their more frequently studied role in neurons. Indeed,
NR4A1 acts in neurons as a transcription factor essential for memory consolidation[41,42]
and functions in microglia as a molecular rheostat, contributing to the maintenance of a
threshold that prevents microglial activation[43]. Furthermore, our study of chromatin
accessibility defines a cell-type-specific signature of transcriptional motifs driving these
processes that includes motifs for ELK4 and NRF1, which have been identified in other
studies as transcriptional motifs associated with the differential regulation of molecular
pathways in specific subsets of microglia[44,45].

This study represents a rare opportunity to define the molecular impact of electrical
stimulation in patients undergoing anterior temporal lobectomy due to therapy refractory
seizures. Although presenting a unique opportunity, it is important to note the caveats
that come from this research carried out in a setting focused on clinical efficacy in
patients that have epilepsy. Thus, we can not control all of the parameters as we might
in an experiment carried out in a model organism, although we have worked to address
three potential limitations of our study. First, our experimental design includes a group
that does not receive electrical stimulation, thus controlling for the impact of the surgical
procedure, craniotomy, brain temperature changes, and levels of anesthesia when
present. We also show that electrical stimulation in a mouse model, in which we can
control many additional parameters, gives rise to similar molecular changes.
Importantly, our mouse experiment included the use of an inactive electrode, thus
controlling for mechanically induced changes in gene expression. Second, the
stimulation pattern used is safe and efficacious for brain mapping in human patients but
is not optimized to modulate activity in specific circuits and cell types in the brain.
Indeed, the stimulation protocol used leads to the functional disruption of brain regions[1],
and thus, we may expect to see reductions in activity-dependent gene expression in

excitatory neurons, as seen in two patients in our NEUROeSTIMator analysis. Third, it

goes without saying that these patients have severe therapy refractory epilepsy.

However, the resected samples were taken a considerable distance from the seizure

foci (see Suppl. Fig. 1b), our gene induction signatures do not resemble those seen in

actively spiking tissue[26], and our molecular changes are seen in a mouse model that

does not exhibit seizures.

Despite these caveats, we were able to identify a reliable cell-specific signature of the

impact of electrical stimulation in the living human brain. This study demonstrates for the

first time a unique transcriptomic and epigenomic signature in microglia following direct

electrical stimulation, representing a conceptual advance in our understanding of how

the brain responds to an important clinical tool used to map brain function. These

findings have the potential to inform the clinical practice of diagnostic and therapeutic

brain stimulation. Clinical and translational research has focused mainly on the

immediate effects of stimulation in neuronal populations, while microglia receive little

attention, often being discussed with regard to neuroimmunological aspects[46]. Although,

for clinical mapping, clinicians rely on the immediate neuronal effects on behavior (e.g.,

speech arrest), there is growing evidence that therapeutic effects of electrical

stimulation can accumulate gradually over time[47]. We could not test a range of

stimulation frequencies and intensities or examine a broad range of brain areas

because of clinical limitations, but our results raise the intriguing hypothesis that

microglia, not just neurons, shape plasticity after repetitive stimulation. Our work further

highlights that cytokine and chemokine-based mechanisms enable microglia to respond

to electrical stimulation and sculpt circuit function, making these potential targets to

modify circuit activity with pharmacological approaches. Microglia exist in different

subpopulations depending on transcriptional state and age[48,49], and our epigenetic work

has revealed particular transcriptional motifs that mark microglial subtypes important for

cortical plasticity. It will be especially interesting in future work to extend our studies to

stimulation protocols used therapeutically as well as natural stimuli to probe the critical

role of microglial signaling in mediating the response of the human brain to experience.

Acknowledgments

We thank Rashmi N. Mueller who oversaw anesthesia to manage pain levels before,
during, and after surgical procedures, and Kyle S. Conway for helping with the initial
pathological assessment. We thank the Iowa Institute of Human Genetics (IIHG) core
for RNA seq library preparation and sequencing.

Funding: This work was supported by grants from the National Institute of Health R01
MH 087463 to T.A and The University of Iowa Hawkeye Intellectual and Developmental
Disabilities Research Center (HAWK-IDDRC) P50 HD103556 to T.A., The National
Institute of Health R00 AG068306 to S.C. and the National Institute of Health R01
DC004290 to M.A.H., National Science Foundation BCS-2342847 to C.I.P. T.N.J &
C.I.P. are supported by a Research Program of Excellence of the Carver Trust and the
Iowa Neuroscience Institute. T.A. is supported by the Roy J. Carver Chair of
Neuroscience, JMM is supported byt the Roy J. Carver Professor of Neuroscience and
the Andrew H. Woods Professorship supported T.N.J. The Neural Circuits and Behavior
Core as well as the NeuroBank Core are supported by the Roy J. Carver Charitable
Trust.

Author contributions

S.C., M.A.H., and T.A. conceived the project, designed the experiments, and interpreted
the data. S.C. and T.A. wrote the manuscript with assistance from C.I.P., L.L., and
T.N.J., and inputs from other co-authors. H.K. performed human surgery and electrical
stimulation and resected the tissue. A.E.R and C.K.K. processed with IRB and MRI
imaging registration. H.O, M.A.H, T.N-J, and K.V.L provided advice on surgical
procedures and electrical stimulation. M.M.H. provided a pathological assessment.
L.C.L., K.V.N., and B.J.P. collected the tissue and images during resection. S.C., B.J.P.,
L.C.L., U.M., L.L. performed human or mouse tissue samples experiments. C.I.P. and
Y.A.A. performed distance to the seizure site analysis. U.M. and L.L. performed mouse
brain imaging and data analysis. Y.V., A.I.G., and B.B. performed the bioinformatics
analysis with inputs from E.B. and J.J.M. S.A.H. performed the mouse surgery and
electrical stimulation experiments. K.V.N, C.K.K, Z.K. Y.K, C.I.P, M.H, J.J.M, E.B, U.M,
and T.N.J. provided input on data analysis and interpretation. All authors discussed the
results and commented on the manuscript.

Competing interests

T.A. is a scientific advisor to Aditum Bio and Radius Health and serves on the scientific
advisory board of Embark Neuro. The other authors declare no conflicting interests.

Figure legends

Figure 1. **Electrical stimulation of the human cortex induces changes in genes**
associated with cytokine activity and transcription regulation. a, b. Schematics of
stimulated (**a**) and unstimulated (**b**) paradigms in the human anterior temporal lobe. For
both paradigms sample A was taken at T = 0 minutes then an adjacent sample B was
taken at either 30 min after stimulation (**a**, stimulated paradigm, 4 participants) or 30 min
after sample A was taken (**b**, unstimulated paradigm, 4 participants). **c.** Volcano plot
showing gene expression changes in the stimulated samples. The most significant
genes (FDR < 0.05) are labeled in red (upregulated) or blue (downregulated). **d.** Cnet
plot showing pathway enrichment analysis of the genes significantly (FDR < 0.05)
differentially expressed in the stimulated samples. **e**. Volcano plots showing gene
expression changes in the unstimulated samples. The most significant genes (FDR <
0.05) are labeled in red (upregulated) or blue (downregulated).

Figure 2. Electrical stimulation of mouse cortex induces changes in genes
associated with cytokine activity and transcription regulation. a. The mouse
auditory cortex was stimulated, tissue was collected 30 min later, and the contralateral
auditory cortex was obtained as baseline control. **b.** Volcano plot showing gene
expression changes after stimulation. The most significant genes (FDR < 0.05) are
labeled in red (upregulated) or blue (downregulated). **c.** Cnet plot showing pathway
enrichment analysis of the genes significantly (FDR < 0.05) affected by electrical
stimulation. **d.** Quadrant plot comparing genes induced by electrical stimulation in mice
with genes induced by electrical stimulation in human brain. Genes upregulated in both
mice and human brains after stimulation are labeled. The size, opacity, and color
intensity of each data point denotes the minimum false discovery rate value for a gene
between each transcriptomic datasets. **e.** qPCR analysis of the genes related to
cytokine activity comparing stimulated versus baseline controls in mice cortex.
n=7/group.

Figure 3. Single nuclei multiomics reveal cell type-specific transcriptomic and
epigenomic changes following electrical stimulation. a. Single nucleus multiomic
experimental approach. **b.** UMAP shows the specific cell types from each cluster. **c.**
Volcano plot showing differentially expressed genes in microglia using pseudobulk
analysis comparing stimulated vs baseline samples from human brain (FDR<0.1). **d.**
Upset plot comparing DEGs in mouse, human bulk RNA seq, and human microglia
snRNA-seq after electrical stimulation. **e.** Cnet plot showing pathway enrichment
analysis of the genes significantly (FDR < 0.1) affected by electrical stimulation in
human microglia. **f.** Quadrant plot comparing DEGs from snRNA-seq following
electrical stimulation in human microglia and genes with open chromatin accessibility
from snATAC-seq in microglia following electrical stimulation. Genes upregulated in both
mice and human brains after stimulation are labeled. **g**. Quadrant plot comparing gene
expression changes (log10(FDR)*log2fold-change) between snRNA-seq (x-axis) and
snATAC-seq (y-axis) data in microglia after stimulation. Genes in red in the top-right
corner of the plot are significantly upregulated at the gene expression level and exhibit
increased chromatin accessibility. The dotted lines represent FDR thresholds of 0.01.
The dashed line represents the linear regression applied to this quadrant plot. **g.** DNA
motifs are overrepresented in the set of peaks differentially accessible in microglia after
stimulation. Motifs are ranked based on significance from the most significant left to
right. **h.** The coverage plot shows ATAC peaks at the CCL4 locus. Each track represents
a normalized chromatin accessibility signal from the ATAC assay for each cell type and
condition (baseline or stimulated).

Figure 4. NEUROeSTIMator for snRNA-seq. Cell-type specific differences in
activity estimates. a. UMAP of cells representing the activity estimated from the RNA
gene counts in the baseline and stimulated conditions. Darker blue points are cells with
higher estimated activity. **b.** The bootstrap distribution of the difference between activity
estimates (stimulated – baseline) by cell type. Horizontal bars indicate the 95%
confidence interval of the mean difference in activity estimate. **c.** Activity estimates in
selected cell types in baseline (green) and stimulated (orange) conditions.

Figure 5. NEUROeSTIMator for snATAC-seq. Cell-type specific differences in
activity estimates. a. UMAP of cells representing the activity estimated from the ATAC

counts in the baseline and stimulated conditions. Darker blue points are cells with

higher estimated activity. **b.** The bootstrap distribution of the difference between activity

estimates (stimulated – baseline) by cell type. Horizontal bars indicate the 95%

confidence interval of the mean difference in activity estimate. **c.** Activity estimates in

selected cell types in baseline (green) and stimulated (orange) conditions.

Materials and Methods:

Patient participants: The study participants were 8 adult neurosurgical patients (6

female, 2 male, age 19-63 years old, median age 41 years old) with medically refractory

epilepsy. The patients were undergoing surgical resection of seizure foci following non-

invasive electroencephalography (EEG) or invasive iEEG monitoring. All patients were

diagnosed with intractable epilepsy. All patients underwent ATL resection surgery for

epilepsy of various etiologies **(Extended data fig 1).**

Two of the patient participants, L472 had a cavernoma and L475 had a

dysembryoplastic neuroepithelial tumor. X out of Y participants were required to be

awake during the surgical resection, and the rest were anesthetized. All patient

participants except #6 were non-smokers. Participant's age, sex, surgery, and awake or

sedative information were recorded (**Extended Data Table 1)**. All participants were

native English speakers, 7 were right-handed, 1 was left-handed, and all had left

language dominance as determined by Wada tests. All participants underwent

audiometric evaluation before the study, and none were found to have hearing deficits

or word recognition scores deemed sufficient to affect the findings presented in this

study. The vision was self-reported as normal or corrected to normal with glasses or

contact lenses. As determined by standard neuropsychological assessments, cognitive

function was in the average range in all participants. Research protocols were

approved by the University of Iowa Institutional Review Board (IRB 201910791,

201911084) and the National Institutes of Health. Written informed consent was

obtained from all participants.

Procedure: Surgery was performed awake, under general anesthesia or monitored

anesthetic care. Standard craniotomy was performed by the same senior epilepsy

neurosurgeon in all patients to reach the epilepsy focus for resection, which involved the

anterior and medial temporal lobe in all patients except one who had a temporal
encephalocele and surrounding anterolateral temporal cortical focus. Before
neurosurgical ATL resection, cortical tissue from the anterior temporal lobe was sampled
by the neurosurgeon and handed over to the research team for analysis. The
experimental condition was defined by an electrical "stimulation paradigm" and a control
"no-stimulation paradigm". The location of the sampled tissue is plotted on anatomic
brain reconstructions **(Extended data fig 1).** The 30 minutes in between the control and
experimental samples was clinically required for the clinical EEG team to record inter-
ictal activity with surface recording grids placed gently on the brain surface.

There were 4 participants (1 iEEG and 3 EEG patients) who underwent the "stimulation
paradigm" and 4 (2 iEEG and 2 EEG patients) who underwent the "no-stimulation
paradigm" **(Extended Data Table 5)**.

In the "stimulation paradigm", a baseline sample was obtained from the anterior
temporal cortex that would be resected in the planned surgical resection. The area
directly adjacent to where the baseline sample was collected was stimulated with direct
bipolar electric stimulation (50 Hz frequency, 0.2 ms pulse duration, 2 min stimulation
duration, and 10 V voltage). The stimulated area was then sampled after a period of 30
minutes to allow for gene expression[50,51]. In the "no stimulation paradigm", no direct
electric stimulation was performed and the area directly adjacent to the baseline sample
was collected 30 minutes after initial baseline sampling.

After sampling the tissue samples were immediately placed in a sterile container on dry
ice. The average weight of the baseline sample was 88.0 ± 33.2 mg (mean, standard
deviation) and the adjacent sample was 113.4 ± 62.7 mg. After the collection of all
samples in this fashion, they were weighed and transferred to a freezer at -80° C for
storage until further testing.

Sample Localization to MNI space: All samples were from the same cortical region.
Intraoperative photos of the sample sites were obtained during the time of surgery.
Using patient participant matched preoperative T1 sequence MRI, the sample sites
were mapped onto their anatomic brain reconstructions **(Extended data fig. 1)**. They
were also mapped onto MNI space coordinates.

Human cortical tissue RNA extraction, library preparation, and sequencing: Total
RNA was extracted from sampled human brains using miRNeasy Mini Kit (Qiagen, CA,
USA). The tissue samples were homogenized in QIAzol (Qiagen, CA. USA)
stainless steel beads (Qiagen, CA, USA). Chloroform was then used for phase
separation. RNA containing an aqueous layer was further purified using the
RNeasy MinElute spin column. RNA was finally eluted in RNase-free water. RNA
concentrations were estimated using a Nanodrop (Thermo Fisher Scientific, MA,
USA) and Qubit (Thermo Fisher Scientific, MA, USA). RNA libraries were prepared at
the Iowa Institute of Human Genetics (IIHG), Genomics Division, using the Illumina
Stranded Total RNA Prep, Ligation with Ribo-Zero Plus (Illumina Inc., San Diego, CA).
The KAPA Illumina Library Quantification Kit (KAPA Biosystems, Wilmington, MA) was
used to measure library concentrations. Pooled libraries were sequenced on Illumina
NovaSeq6000 sequencers with 150-bp paired-end chemistry (Illumina) at the Iowa
Institute of Human Genetics (IIHG) core.

Bulk RNA sequencing analysis: RNA-seq data were processed with the bcbio-
nextgen pipeline (https://github.com/bcbio/bcbio-nextgen, version 1.2.9). The pipeline
uses STAR[52] to align reads to the hg38 or mm10 reference genome and quantifies
expression at the gene level with feature Counts [53]. All further analyses were performed
using R. For gene-level count data, the R package EDASeq was used to account for
sequencing depth (upper quartile normalization) [54]. Latent sources of variation in
expression levels were assessed and accounted for using RUVSeq (RUVs mode using
all features) [55]. Appropriate choice of the RUVSeq parameter k was guided through
inspection of principal components analysis (PCA) plots. Specifically, the smallest value
k was chosen where PCA plots demonstrated replicate sample clustering in the first
three principal components[56]. Differential expression analysis was conducted using the
edgeR package[57]. Codes to reproduce the RNA-seq differential gene expression
analysis are available at https://github.com/YannVRB/Human-brain-stimulation.git.

All the transcriptomics data have been deposited in NCBI's Gene Expression Omnibus
and are accessible through GEO Series accession number GSE224952.

Downstream pathway analysis:

The enrichment analysis of differentially expressed genes-associated pathways and
molecular functions from the stimulated and unstimulated RNA-seq was performed with
the Gene Ontology (GO–molecular function) databases using clusterProfiler package in
R. Only the pathways with an adjusted p-value $<$ 0.05 were considered as significant
and displayed. Further, the enrichment data were visualized using 'cnetplot' function of
clusterProfiler.

Single-nuclei multiomics (nuclei isolation, library preparation, sequencing: Nuclei
were isolated from brain tissue using the Chromium Nuclei Isolation Kit (10X
Genomics). Briefly, frozen tissue was dissociated with pestle in lysis buffer, passed
through nuclei isolation column and spun at 16,000 rcf for 20 sec at 4^0C. Flowthrough
was vortexed and spun at 500 rcf for 3 mins at 4^0C. Pellet was resuspended with debris
removal buffer and centrifuged at 700 rcf for 10 mins at 4^0C, nuclei resuspended in
wash buffer and centrifuged again at 500 rcf for 5 mins at 4^0C. Pellet was resuspended
in resuspension buffer and nuclei were counted using a hemocytometer. Nuclei were
directly processed for droplet capture for single cell multiome ATAC + gene expression
using a chromium controller (10X Genomics). Chromium Next GEM Single Cell
Multiome ATAC + Gene v1 chemistry was used to create single nuclei ATAC and RNA
libraries from the same cell. Two baseline and two stimulated samples were used for
independent replicates. Libraries were sequenced on an Illumina Novaseq 6000 with a
150 bp paired end read setup.

Single-nuclei multiomic data processing and analysis: To analyze the RNA part of
the human brain stimulation multiomic data, gene counts were normalized and log
transformed (LogNormalize), and the top 2,000 most variable features between each
nuclei were identified using FindVariableFeatures (selection.method $=$ 'vst'). Features
that are repeatedly variable across nuclei and datasets were selected for integration
(SelectIntegrationFeatures). We then identified anchors (FindIntegrationAnchors), which
took the list of 4 individual Seurat objects for each sample as input and used these
anchors to integrate the four datasets together (IntegrateData). The following analyses
were performed on the integrated Seurat object. Linear dimensionality reduction was

performed by principal component analysis (runPCA, npcs□=□25). A k-nearest-
neighbors graph was constructed based on Euclidean distance in PCA space and
refined (FindNeighbors, npcs□=□30), then nuclei were clustered using the Louvain
algorithm (FindClusters, resolution□=□0.5). Clusters were visualized with UMAP
(runUMAP, dims□=□30). Cell types were annotated by label transfer cell labels from an
existing human primary motor cortex reference dataset from the Allen Institute (doi:
10.1038/s41586-021-03465-8) (FindTransferAnchors and TransferData). Cell types
identification was validated by expression of specific biomarkers. Prior of running
differential gene expression analysis, as recommended by recent publications
(https://doi.org/10.1038/nmeth.4612 and https://doi.org/10.1038/s41467-020-19894-4),
we used an aggregation-based (pseudobulk) workflow. We aggregated all cells within
the same cell type and sample using the AggregateExpression function. This returns a
Seurat object where each 'cell' represents the pseudobulk profile of one cell type in one
individual. After we aggregated cells, we performed celltype-specific differential
expression between stimulated and baseline samples using DESeq2.

To analyze ATAC part of the human brain stimulation multiomic data, prior to integrating
the four Seurat object, the default assay was switched to ATAC, and peak calling was
performed. Since the set of peaks identified by Cellranger often merges distinct peaks
that are close together, creating a problem for motif enrichment analysis and peak-to-
gene linkage, we identified a more accurate set of peaks by calling peaks using MACS2
(CallPeaks) on all cells together. Peaks on nonstandard chromosomes and in genomic
blacklist regions were removed (keepStandardChromosomes and subsetByOverlaps).
Normalization was performed with a frequency-inverse document frequency
normalization which normalizes across cells and peaks (RunTFIDF). Then, a feature
selection was performed using all the peaks as input (FindTopFeatures). The
dimensional reduction was performed on the TF-IDF normalized matrix with the
selected peaks using a singular value decomposition (RunSVD). To mimic the open
chromatin conformation of a gene, a gene activity matrix was calculated using a window
of 1000bp before and after the transcription start site of each protein coding gene
(GeneActivity). Differentially accessible transcription start sites in individual clusters
between baseline and stimulated samples were calculated using a logistic regression

framework (FindMarkers, test.use = 'LR', latent.vars = 'nCount_peaks', Padj < 0.05).

Motif and transcription factor enrichment analysis for Microglia cluster was performed

using FindMotifs on genome-wide all peaks assay of the Seurat object. The top six

enriched motifs in microglia cluster are shown. Genomic locations of typical genes like

CCL4 were presented (CoveragePlot). It also includes co-accessibility between peaks

and transcription start site of genes. Codes to reproduce the multiomic data analysis are

available at https://github.com/YannVRB/Human-brain-stimulation.git.

NEUROeSTIMator analysis. Gene-level counts for each cell were used as input for the

NEUROeSTIMator model[33] to estimate transcriptional signatures of cell activity. Activity

was also estimated for cells using gene-level counts in promoter regions from the ATAC

assay. Estimated activity was assessed for significant differences between the baseline

and stimulated conditions across the donors and cell types using paired linear

regression. P-values were adjusted for multiple testing with the Benjamini-Hochberg

correction. To account for variable sample sizes across cell types and participants,

bootstrap resampling was performed (n = 100 cells per donor, replicates = 1000) for

each cell type in each condition to generate a distribution of the mean difference in

estimated activity between conditions.

To account for variation in sample size across cell types and participants, bootstrap

resampling was performed (n = 100 cells per participant, replicates = 1000) for each

cell type of the mean difference in estimated activity scores between conditions (mean

stimulated – mean baseline). The distributions were analyzed by mean and 95%

confidence interval.

Animals: Adult male C57BL/6J mice were purchased from The Jackson Laboratory

were 3 to 4 months of age during experiments. All mice had free access to food and

water; lights were maintained on a 12-hour light/12-hour dark cycle.

Mouse electrical stimulation: Stimulation experiments were performed in anesthetized

adult male C57BL6/J mice. Anesthesia was induced with 5% isoflurane by inhalation

and maintained at 1.8-2% for the duration of the experiment. The mouse was placed in

a stereotax (Kopf) and a midline incision was made and the skin retracted to expose the

temporal muscle bilaterally. The dorsal insertion of both temporal muscles was

removed, and the muscles retracted. A 2-3 mm craniotomy was made over area AuV

(centered at 2.9 mm posterior, 4.2 mm lateral, 2.8 mm ventral from Bregma based on

Paxinos atlas) bilaterally to expose the cortical surface and a small square of gel foam

soaked in ACSF was applied on top of the dura to prevent it from dehydrating. For each

mouse the side of electrical stimulation and sham stimulation were alternated, with the

sham side serving as the baseline control for gene expression profiling. Electrical

stimulation was delivered through a bipolar ball electrode constructed from two silver

wires in which the uninsulated tips were melted under a butane flame (1 mm tip size, 2

mm tip spacing). For both electrical and sham stimulations, the gel foam was removed

and the electrode was gently lowered to make contact with the cortical surface. For

electrical stimulation, a biphasic pulse train was then delivered for two minutes (8 mA,

50 Hz, 200 μs pulse width). For the sham stimulation no current was delivered but the

electrode was left in place for 2 minutes. Following electrical or sham stimulation the

electrode was slowly retracted, and the exposed dura was covered with gel foam.

Selection for hemisphere was performed randomly, which resulted in the order of

electrical and sham stimulation alternating from mouse to mouse. No more than 5

minutes elapsed between electrical or sham stimulation of both sides. After both sides

were stimulated (electrical or sham) the mouse was left in the stereotax under

anesthesia for 30 minutes before euthanasia and tissue collection. Tissue samples were

immediately stored at -80^{0}C in RNA later solution (Ambion).

RNA extraction, cDNA preparation and qPCRs from mouse auditory cortex: Tissue

samples were homogenized in Qiazol (Qiagen) using stainless steel beads (Qiagen).

Chloroform was added and centrifuged at 12,000*g* at room temperature for 15 min to

separate RNA in the aqueous phase. RNA was precipitated in ethanol and cleared using

RNeasy kit (Qiagen). RNA eluted in nuclease-free water was then treated with DNase

(Qiagen) at room temperature for 25 min to remove genomic DNA. RNA was further

precipitated in ethanol, sodium acetate (pH 5.2) and glycogen overnight at –20°C. RNA

was precipitated by centrifugation at 30,000g for 20 min, precipitate washed with 70%

ethanol and the dried RNA pellet was resuspended in nuclease-free water. RNA

concentration was measured using NanoDrop (Thermo Fisher Scientific). 1 μg of RNA

was used for complementary DNA (cDNA) preparation using the SuperScript IV First-

Strand Synthesis System (Ambion). Real-time reverse transcription polymerase chain

reactions (RT-PCRs) were performed on the QuantStudio 7 Flex Real-Time PCR

System (Applied Biosystems, Life Technologies). Data were normalized to

housekeeping genes (*Tubulin*, *Pgk1*, and *Actin*), and $2^{(-\Delta\Delta Ct)}$ method was used for gene

expression analysis.

Mouse cortex spatial transcriptomics analysis (Yann): Spatial transcriptomics data

were obtained from the Visium Spatial Gene Expression platform, as described

previously[42]. This dataset provides comprehensive gene expression profiles across

various brain regions, including the mouse cortex. To focus on the mouse auditory

cortex, the closest homolog to the human anterior temporal lobe, we manually selected

spatial spots from the Visium data corresponding to this brain region. This selection was

based on anatomical landmarks and known spatial coordinates for the auditory cortex.

For each of the 14 mice in the dataset, we subsetted the data to include only these

selected spots, thereby isolating the gene expression profiles specifically from the

auditory cortex.

The dataset of 14 mice was divided into two experimental conditions: 7 mice underwent

Spatial Object Recognition after 1 hour (SOR-1h), while the other 7 mice were kept in

their homecage environment without specific tasks. Based on the Visium spots data

overlaying the mouse auditory cortex, differential gene expression analysis was

performed to identify genes that were significantly differentially expressed between the

SOR-1h and homecage conditions.

Statistics: For the qPCR analysis, the Wilcoxon matched pairs signed rank test and

one sample Wilcoxon test was performed.

References

1   Penfield, W. & Rasmussen, T. Vocalization and arrest of speech. *Arch Neurol*
Psychiatry **61**, 21-27 (1949).
https://doi.org:10.1001/archneurpsyc.1949.02310070027002

2   Lozano, A. M. *et al.* Deep brain stimulation: current challenges and future
directions. *Nat Rev Neurol* **15**, 148-160 (2019). https://doi.org:10.1038/s41582-
018-0128-2

3   Okun, M. S. Deep-brain stimulation for Parkinson's disease. *N Engl J Med* **367**,
1529-1538 (2012). https://doi.org:10.1056/NEJMct1208070

4   De Risio, L. *et al.* Recovering from depression with repetitive transcranial
magnetic stimulation (rTMS): a systematic review and meta-analysis of
preclinical studies. *Transl Psychiatry* **10**, 393 (2020).
https://doi.org:10.1038/s41398-020-01055-2

5   Kellner, C. H. *et al.* ECT in treatment-resistant depression. *Am J Psychiatry* **169**,
1238-1244 (2012). https://doi.org:10.1176/appi.ajp.2012.12050648

6   Krauss, J. K. *et al.* Technology of deep brain stimulation: current status and
future directions. *Nat Rev Neurol* **17**, 75-87 (2021).
https://doi.org:10.1038/s41582-020-00426-z

7   Sheth, S. A. & Mayberg, H. S. Deep Brain Stimulation for Obsessive-Compulsive
Disorder and Depression. *Annu Rev Neurosci* **46**, 341-358 (2023).
https://doi.org:10.1146/annurev-neuro-110122-110434

8   Goodman, W. K., Storch, E. A. & Sheth, S. A. Harmonizing the Neurobiology and
Treatment of Obsessive-Compulsive Disorder. *Am J Psychiatry* **178**, 17-29
(2021). https://doi.org:10.1176/appi.ajp.2020.20111601

9   George, D. D., Ojemann, S. G., Drees, C. & Thompson, J. A. Stimulation
Mapping Using Stereoelectroencephalography: Current and Future Directions.
Front Neurol **11**, 320 (2020). https://doi.org:10.3389/fneur.2020.00320

10   Grande, K. M., Ihnen, S. K. Z. & Arya, R. Electrical Stimulation Mapping of Brain
Function: A Comparison of Subdural Electrodes and Stereo-EEG. *Front Hum
Neurosci* **14**, 611291 (2020). https://doi.org:10.3389/fnhum.2020.611291

11   Yap, E. L. & Greenberg, M. E. Activity-Regulated Transcription: Bridging the Gap
between Neural Activity and Behavior. *Neuron* **100**, 330-348 (2018).
https://doi.org:10.1016/j.neuron.2018.10.013

12   Sagar, S. M., Sharp, F. R. & Curran, T. Expression of c-fos protein in brain:
metabolic mapping at the cellular level. *Science* **240**, 1328-1331 (1988).
https://doi.org:10.1126/science.3131879

13   Guzowski, J. F., McNaughton, B. L., Barnes, C. A. & Worley, P. F. Environment-
specific expression of the immediate-early gene Arc in hippocampal neuronal
ensembles. *Nat Neurosci* **2**, 1120-1124 (1999). https://doi.org:10.1038/16046

14   Laricchiuta, D. *et al.* Optogenetic Stimulation of Prelimbic Pyramidal Neurons
Maintains Fear Memories and Modulates Amygdala Pyramidal Neuron
Transcriptome. *Int J Mol Sci* **22** (2021). https://doi.org:10.3390/ijms22020810

15   Bali, P. & Kenny, P. J. Transcriptional mechanisms of drug addiction□*Dialogues
Clin Neurosci* **21**, 379-387 (2019).
https://doi.org:10.31887/DCNS.2019.21.4/pkenny

16   Fernandez-Albert, J. *et al.* Immediate and deferred epigenomic signatures of in
vivo neuronal activation in mouse hippocampus. *Nat Neurosci* **22**, 1718-1730
(2019). https://doi.org:10.1038/s41593-019-0476-2

17   Marco, A. *et al.* Mapping the epigenomic and transcriptomic interplay during
memory formation and recall in the hippocampal engram ensemble. *Nat Neurosci*
23, 1606-1617 (2020). https://doi.org:10.1038/s41593-020-00717-0

18   Trevino, A. E. *et al.* Chromatin and gene-regulatory dynamics of the developing
human cerebral cortex at single-cell resolution. *Cell* **184**, 5053-5069 e5023
(2021). https://doi.org:10.1016/j.cell.2021.07.039

19   Reijmers, L. G., Perkins, B. L., Matsuo, N. & Mayford, M. Localization of a stable
neural correlate of associative memory. *Science* **317**, 1230-1233 (2007).
https://doi.org:10.1126/science.1143839

20   Halder, R. *et al.* DNA methylation changes in plasticity genes accompany the
formation and maintenance of memory. *Nat Neurosci* **19**, 102-110 (2016).
https://doi.org:10.1038/nn.4194

21   Boulting, G. L. *et al.* Activity-dependent regulome of human GABAergic neurons
reveals new patterns of gene regulation and neurological disease heritability. *Nat*
Neurosci **24**, 437-448 (2021). https://doi.org:10.1038/s41593-020-00786-1

22   Berto, S. *et al.* Gene-expression correlates of the oscillatory signatures
supporting human episodic memory encoding. *Nat Neurosci* **24**, 554-564 (2021).
https://doi.org:10.1038/s41593-021-00803-x

23   Konopka, G. Cognitive genomics: Linking genes to behavior in the human brain.
Netw Neurosci **1**, 3-13 (2017). https://doi.org:10.1162/NETN_a_00003

24   Berger, M. S. & Ojemann, G. A. Intraoperative brain mapping techniques in
neuro-oncology. *Stereotact Funct Neurosurg* **58**, 153-161 (1992).
https://doi.org:10.1159/000098989

25   Sanai, N., Mirzadeh, Z. & Berger, M. S. Functional outcome after language
mapping for glioma resection. *N Engl J Med* **358**, 18-27 (2008).
https://doi.org:10.1056/NEJMoa067819

26   Brueggeman, L. *et al.* Drug repositioning in epilepsy reveals novel antiseizure
candidates. *Ann Clin Transl Neurol* **6**, 295-309 (2019).
https://doi.org:10.1002/acn3.703

27   Altmann, A. *et al.* A systems-level analysis highlights microglial activation as a
modifying factor in common epilepsies. *Neuropathol Appl Neurobiol* **48**, e12758
(2022). https://doi.org:10.1111/nan.12758

28   Murphy, A. E. & Skene, N. G. A balanced measure shows superior performance
of pseudobulk methods in single-cell RNA-sequencing analysis. *Nat Commun* **13**,
7851 (2022). https://doi.org:10.1038/s41467-022-35519-4

29   Morabito, S. *et al.* Single-nucleus chromatin accessibility and transcriptomic
characterization of Alzheimer's disease. *Nat Genet* **53**, 1143-1155 (2021).
https://doi.org:10.1038/s41588-021-00894-z

30   Caglayan, E., Liu, Y. & Konopka, G. Neuronal ambient RNA contamination
causes misinterpreted and masked cell types in brain single-nuclei datasets.
Neuron (2022). https://doi.org:10.1016/j.neuron.2022.09.010

31   Huang, Y. *et al.* ELK4 exerts opposite roles in cytokine/chemokine production and
degranulation in activated mast cells. *Front Immunol* **14**, 1171380 (2023).
https://doi.org:10.3389/fimmu.2023.1171380

32   Bahl, E. *et al.* Using deep learning to quantify neuronal activation from single-cell
and spatial transcriptomic data. *Nat Commun* **15**, 779 (2024).
https://doi.org:10.1038/s41467-023-44503-5

33   Berto, S., Wang, G. Z., Germi, J., Lega, B. C. & Konopka, G. Human Genomic
Signatures of Brain Oscillations During Memory Encoding. *Cereb Cortex* **28**,
1733-1748 (2018). https://doi.org:10.1093/cercor/bhx083

34   Badimon, A. *et al.* Negative feedback control of neuronal activity by microglia.
Nature **586**, 417-423 (2020). https://doi.org:10.1038/s41586-020-2777-8

35   Laprell, L., Schulze, C., Brehme, M. L. & Oertner, T. G. The role of microglia
membrane potential in chemotaxis. *J Neuroinflammation* **18**, 21 (2021).
https://doi.org:10.1186/s12974-020-02048-0

36   Liu, Y. U. *et al.* Neuronal network activity controls microglial process surveillance
in awake mice via norepinephrine signaling. *Nat Neurosci* **22**, 1771-1781 (2019).
https://doi.org:10.1038/s41593-019-0511-3

37   Stowell, R. D. *et al.* Noradrenergic signaling in the wakeful state inhibits
microglial surveillance and synaptic plasticity in the mouse visual cortex. *Nat*
Neurosci **22**, 1782-1792 (2019). https://doi.org:10.1038/s41593-019-0514-0

38   Cheadle, L. *et al.* Sensory Experience Engages Microglia to Shape Neural
Connectivity through a Non-Phagocytic Mechanism. *Neuron* **108**, 451-468 e459
(2020). https://doi.org:10.1016/j.neuron.2020.08.002

39   Estevao, C. *et al.* CCL4 induces inflammatory signalling and barrier disruption in
the neurovascular endothelium. *Brain Behav Immun Health* **18**, 100370 (2021).
https://doi.org:10.1016/j.bbih.2021.100370

40   Shen, Y. *et al.* CCR5 closes the temporal window for memory linking. *Nature* **606**,
146-152 (2022). https://doi.org:10.1038/s41586-022-04783-1

41   Chatterjee, S. *et al.* Endoplasmic reticulum chaperone genes encode effectors of
long-term memory. *Sci Adv* **8**, eabm6063 (2022).
https://doi.org:10.1126/sciadv.abm6063

42   Vanrobaeys, Y. *et al.* Mapping the spatial transcriptomic signature of the
hippocampus during memory consolidation. *Nat Commun* **14**, 6100 (2023).
https://doi.org:10.1038/s41467-023-41715-7

43   Rothe, T. *et al.* The Nuclear Receptor Nr4a1 Acts as a Microglia Rheostat and
Serves as a Therapeutic Target in Autoimmune-Driven Central Nervous System
Inflammation. *J Immunol* **198**, 3878-3885 (2017).
https://doi.org:10.4049/jimmunol.1600638

44   Chiu, I. M. *et al.* A neurodegeneration-specific gene-expression signature of
acutely isolated microglia from an amyotrophic lateral sclerosis mouse model.
Cell Rep **4**, 385-401 (2013). https://doi.org:10.1016/j.celrep.2013.06.018

45 Olah, M. *et al.* Single cell RNA sequencing of human microglia uncovers a subset
associated with Alzheimer's disease. *Nat Commun* **11**, 6129 (2020).
https://doi.org:10.1038/s41467-020-19737-2

46 Sheng, R., Chen, C., Chen, H. & Yu, P. Repetitive transcranial magnetic
stimulation for stroke rehabilitation: insights into the molecular and cellular
mechanisms of neuroinflammation. *Front Immunol* **14**, 1197422 (2023).
https://doi.org:10.3389/fimmu.2023.1197422

47 Heck, C. N. *et al.* Two-year seizure reduction in adults with medically intractable
partial onset epilepsy treated with responsive neurostimulation: final results of the
RNS System Pivotal trial. *Epilepsia* **55**, 432-441 (2014).
https://doi.org:10.1111/epi.12534

48 Hammond, T. R. *et al.* Single-Cell RNA Sequencing of Microglia throughout the
Mouse Lifespan and in the Injured Brain Reveals Complex Cell-State Changes.
Immunity **50**, 253-271 e256 (2019). https://doi.org:10.1016/j.immuni.2018.11.004

49 Nott, A. *et al.* Brain cell type-specific enhancer-promoter interactome maps and
disease-risk association. *Science* **366**, 1134-1139 (2019).
https://doi.org:10.1126/science.aay0793

50 Hansson, A. C. & Fuxe, K. Time-course of immediate early gene expression in
hippocampal subregions of adrenalectomized rats after acute corticosterone
challenge. *Brain Res* **1215**, 1-10 (2008).
https://doi.org:10.1016/j.brainres.2008.03.080

51 Cullinan, W. E., Herman, J. P., Battaglia, D. F., Akil, H. & Watson, S. J. Pattern
and time course of immediate early gene expression in rat brain following acute
stress. *Neuroscience* **64**, 477-505 (1995). https://doi.org:10.1016/0306-
4522(94)00355-9

52 Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-
21 (2013). https://doi.org:10.1093/bioinformatics/bts635

53 Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose
program for assigning sequence reads to genomic features. *Bioinformatics* **30**,
923-930 (2014). https://doi.org:10.1093/bioinformatics/btt656

54 Risso, D., Schwartz, K., Sherlock, G. & Dudoit, S. GC-content normalization for
RNA-Seq data. *BMC Bioinformatics* **12**, 480 (2011). https://doi.org:10.1186/1471-
2105-12-480

55 Risso, D., Ngai, J., Speed, T. P. & Dudoit, S. Normalization of RNA-seq data
using factor analysis of control genes or samples. *Nat Biotechnol* **32**, 896-902
(2014). https://doi.org:10.1038/nbt.2931

56 Peixoto, L. *et al.* How data analysis affects power, reproducibility and biological
insight of RNA-seq studies in complex datasets. *Nucleic Acids Res* **43**, 7664-
7674 (2015). https://doi.org:10.1093/nar/gkv736

57   Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package
for differential expression analysis of digital gene expression data. *Bioinformatics*
26, 139-140 (2010). https://doi.org:10.1093/bioinformatics/btp616

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5