1

Intestinal microbiota programming of alveolar macrophages influences severity of

respiratory viral infection

Vu L. Ngo[1#], Carolin M. Lieber[1#], Hae-ji Kang[1], Kaori Sakamoto[2], Michal Kuczma[1],

Richard K. Plemper[1&], and Andrew T. Gewirtz[1,3&]

Affiliation:

1. Center for Translational Antiviral Research, Georgia State University Institute for Biomedical

Sciences, Atlanta GA 30303, USA.

2. Department of Pathology, University of Georgia College of Veterinary Science, Athens GA

30602, USA

3. Lead contact

# These authors contributed equally to this work

& Denotes co-senior and co-corresponding authors

Correspondence: rplemper@gsu.edu or agewirtz@gsu.edu

Abstract (146 words)

Susceptibility to respiratory virus infections (RVIs) varies widely across individuals. Because the gut microbiome impacts immune function, we investigated the influence of intestinal microbiota composition on RVI and determined that segmented filamentous bacteria (SFB), naturally acquired or exogenously administered, protected mice against influenza virus (IAV) infection. Such protection, which also applied to respiratory syncytial virus and SARS-CoV-2, was independent of interferon and adaptive immunity but required basally resident alveolar macrophages (AM). In SFB-negative mice, AM were quickly depleted as RVI progressed. In contrast, AM from SFB-colonized mice were intrinsically altered to resist IAV-induced depletion and inflammatory signaling. Yet, AM from SFB-colonized mice were not quiescent. Rather, they directly disabled IAV via enhanced complement production and phagocytosis. Accordingly, transfer of SFB-transformed AM into SFB-free hosts recapitulated SFB-mediated protection against IAV. These findings uncover complex interactions that mechanistically link the intestinal microbiota with AM functionality and RVI severity.

One sentence summary: Intestinal segmented filamentous bacteria reprogram alveolar macrophages promoting nonphlogistic defense against respiratory viruses.

INTRODUCTION

Outcomes following exposure to respiratory viruses, such as influenza A viruses (IAV), respiratory syncytial virus (RSV), and severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), vary widely amongst individuals, ranging from asymptomatic infection to severe lung pathology and/or death. While determinants of such heterogeneity to RVI are likely numerous and complex, we hypothesized that one important factor is gut microbiota composition [1], which is now appreciated to have broad influence over a range of chronic inflammatory diseases and the immune responses with which such diseases are associated [2]. For example, studies comparing mice captured in the wild to those bred for generations in well-controlled vivaria revealed vast differences in their microbiomes that resulted in wild mice having highly and broadly activated immune systems and being relatively resistant to IAV [3,4]. Yet, differences in specific microbial species that can impact RVI outcomes, and how they might do so, have not been well defined. Hence, we studied mice raised in vivaria with discrete defined microbiome differences and, subsequently, mice differing in only the presence or absence of a single common species, namely segmented filamentous bacteria (SFB). We found that gut SFB dictated the phenotype of alveolar macrophages and, consequently, modulated outcomes of RVI.

RESULTS

SFB protects mice from IAV infection

Appreciation of the potential of microbiota to broadly impact health led one commercial rodent supplier to offer "excluded flora (EF)" mice, which were bred in a restricted vivarium that assured absence of a discrete panel of disease-modulating commensal microbes, which might or might not otherwise be present in colonies of "specific pathogen-free (SPF)" mice commonly used in biomedical research [5]. We used such mice as a starting point to probe the influence of microbiota composition on RVI. Specifically, we compared proneness of SPF and EF mice to IAV infection following intranasal administration of 2009 pandemic A/CA/07/2009 (H1N1), herein referred to as CA09, which readily infects mice without need for species adaptation and recapitulates major clinical features of human disease in the animal model [6]. Viral titers in the lung were measured 4 days post-infection (dpi), a time point at which disease symptoms, including mortality, could be assessed although experiments explicitly designed to measure survival were allowed to proceed until 14 dpi, when infectious particles were no longer detectable. Compared to EF mice, SPF mice exhibited significantly reduced lung virus loads. Furthermore, SPF mice did not show hypothermia and weight loss observed in EF mice **(Figure 1A)**. Additionally, some mice of the EF mice required euthanasia. These results suggested that one or more of the microbes specifically excluded from EF mice might confer protection against IAV.

Of the panel of organisms known to be absent in EF mice, segmented filamentous bacteria (SFB) stood out as a potential modulator of IAV proneness in that SFB, despite being a strict anaerobe and, thus, restricted to the intestinal luminal surface, is known to systemically

impact T-lymphocytes [7,8]. Furthermore, SFB is a major contributor to, albeit not the sole

mediator of, spontaneous resistance of mice to rotavirus (RV), an intestinal pathogen, that arose

in some mouse colonies [9]. SFB is very challenging to culture but can be isolated, maintained, and

studied by administering it via fecal microbial transplantation (FMT). Hence, mice lacking SFB

but otherwise carrying a complex SPF microbiota, herein referred to as SFB⁻ mice, were

administered a FMT from other SFB⁻ mice, mice containing SFB as one natural component of

their SPF microbiota, or germ-free mice that had been monoassociated (MA) with SFB. Such

mice were inoculated with CA09 one-week post-FMT and euthanized 4d later to measure lung

viral titers or monitored for 14 days to assess survival. Mice receiving FMT from donors

naturally colonized with SFB exhibited IAV titer reductions of approximately one order of

magnitude compared to animals that lacked SFB and, concomitantly, clinical signs of IAV

infection were ameliorated **(Figure 1B).** FMT from SFB-MA mice fully recapitulated such

protection while administering feces from germ-free mice was without effect **(Figure S1A).**

Thus, administering SFB⁻ mice a FMT from SFB-MA mice, hereafter referred to as

administering SFB, recapitulated the differential proneness of EF and SPF mice to IAV

infection, thus providing a tractable model to investigate how the presence of this specific gut

microbiota constituent, within a complex microbial ecosystem, reduced IAV infection in the

lung.

SFB-mediated protection against IAV infection was evident, albeit seemingly not

maximal, two days post SFB administration and was undiminished three months later (longest

time tested) despite gut SFB levels having declined by over 60-fold from peak to a stable plateau

by this time **(Figure 1C and Figure S1B).** In accord with its relative abundance not being a

critical variable, SFB colonized 24-week-old mice at 10-fold lower levels than 5-week-old mice

but still protected them against IAV infection **(Figure S1C)**. Consistent with previous

observations that SFB only resides in the gut lumen or attached to the apical epithelial gut

surface [10], SFB was readily detected by PCR in the ileum, but was not found in the lung or

spleen **(Figure 1D)**. Histopathologic examination of lung tissue did not detect any impact of SFB

itself on the lung but revealed a striking suppression of IAV-induced lung pathology **(Figure**

1E). Such protection of lung tissue was potentially attributable to the 1-log reduction in viral

titers conferred by SFB colonization but also hinted at the possibility that SFB had reduced IAV-

induced inflammatory signaling.

SFB-mediated protection against IAV is not explained by known candidate mechanisms

113   SFB-mediated lasting protection against IAV contrasted with its transient protection

against RV [9], suggesting that the former might be mediated by adaptive immunity. However,

SFB-mediated protection against IAV was fully maintained in *Rag1*[-/-] mice, which lack most T

and B lymphocytes, arguing against this notion **(Figure 2A)**. Innate antiviral immunity typically

involves interferon and can involve innate lymphoid cells (ILC) and/or IL-22 and IL-17. Yet,

SFB-mediated protection against IAV was also maintained in *Ifnar*[-/-] mice, which lack type I

interferon signaling, *Rag1*[-/-]*-Il2rg*[-/-], which lack type 3 ILC, and upon antibody-mediated

neutralization of IL-17 and IL-22 **(Figure 2B-D)**. SFB-mediated protection against IAV was

furthermore maintained in *Stat1*[-/-] animals, which are broadly compromised in IFN signaling

(Figure 2E). These results argue against a role for any of these candidate host mechanisms in

mediating SFB's protection against IAV infection. We also considered a candidate microbial

mechanism. Specifically, we hypothesized that the protection against IAV conferred by SFB

might reflect that it induced an array of complex changes in other taxa that might then broadly

impact innate immunity. However, SFB's protection was maintained in Altered Schaedler flora

(ASF) mice, which harbor only a minimal 8-species microbial community [11], thus arguing

against this possibility.

SFB-mediated protection against IAV, RSV and SARS-CoV-2 is associated with

preservation of resident lung phagocytes

That candidate mechanisms were not experimentally substantiated led us to utilize less

targeted approaches to investigate how SFB gut colonization protected mice against IAV. First,

we performed transcriptomic profiling of whole lungs harvested from mice that were untreated

(naïve) or administered SFB, CA09, or a combination thereof. The results of this RNA-seq

analysis were presented as a heat map, showing 4-5 individual mice per condition, of all genes

whose expression was significantly altered in response to any of the three treatments **(Figure**

3A). It illustrated that SFB, by itself, had almost no impact on lung gene expression, whereas, as

expected, IAV infection, by itself, markedly remodeled the lung transcriptome. Strikingly, such

IAV-induced changes in gene expression were absent in SFB-colonized mice, indicating that the

presence of SFB in the gut broadly ameliorated IAV's impact on lung tissue. The majority of

genes induced by CA09 were related to immunity and inflammation and thus accorded with the

view that much of the lung pathology in IAV infection is driven by the immune system [12] and,

furthermore, the notion that SFB had dampened such responses.

We next used flow cytometry to examine the impact of SFB and IAV on lung populations

of immune cells, focusing on innate cells given that our results from $Rag1^{-/-}$ mice had suggested

lymphocytes were not required for SFB-mediated protection against IAV. Consistent with other

studies, intranasal administration of CA09, by itself, induced a stark depletion of basally-resident

CD103$^+$ dendritic cells (cDC1) and alveolar macrophages (AM), which was followed by

increased abundance of a panel of inflammatory cells including neutrophils, monocytes, and

natural killer cells [13]. SFB, by itself, had only modest impacts on levels of lung leukocytes,

slightly increasing abundance of AM, but wholly prevented the changes induced by CA09

(Figure 3B and Figure S2). Preventing these CA09-induced changes in lung cellularity and

disease was not SFB strain-specific as similar protective effects resulted from colonization of the

gut by SFB strains isolated in 3 different continents **(Figure S3)**. Nor were SFB impacts specific

to IAV. Rather, SFB colonization also reduced viral loads and prevented changes in lung

cellularity, and reduced pro-inflammatory gene expression, following inoculation of mice with

respiratory syncytial virus (RSV) and mouse-adapted SARS-CoV-2 **(Figure S4)**, which,

irrespective of SFB, did not cause clinical signs of disease.

To investigate the extent to which SFB-mediated prevention of IAV-induced changes in

lung cellularity simply reflected lower IAV loads, we pharmacologically restricted IAV

replication via 4'-fluorouridine (4'-FIU), a broad-spectrum nucleoside analog inhibitor that we

recently reported reduces viral titers and disease symptoms up to 60 hours following

administration of IAV, RSV, and SARS-CoV-2 [14,15]. SFB$^-$ mice received 4'-FIU via oral gavage

at different dose levels in a once daily (q.d.) regimen beginning 24 hours post-inoculation. At

this time, IAV levels and immune cell populations were similar in SFB$^-$ and SFB-administered

mice **(Figure S2C)**, which we hereafter refer to as SFB$^+$ mice. As expected, administration of 4'-

FIU to SFB$^-$ mice dose-dependently suppressed IAV replication to a similar or greater extent

than SFB **(Figure 3C)**. Like SFB, 4'-FIU administration prevented the CA09-induced

appearance of inflammatory cells in the lung. However, even the highest dose of 4'-FIU, which

lowered IAV titers by over three orders of magnitude and markedly reduced inflammatory cell

recruitment, did not mitigate CA09-induced ablation of cDC1 or AM. This result suggested that

the persistence of AM and cDC1 in SFB$^+$ mice was not entirely a consequence of reduced viral

loads and/or inflammation but rather implied that maintenance of these cells may contribute to

suppression of IAV.

Alveolar Macrophages are required for SFB-mediated protection against IAV

Potential roles for cDC1 and AM in SFB-mediated protection against IAV were tested in

mice harboring deficiencies in these cells. We observed that SFB-mediated protection against

IAV was maintained in *Batf3*$^{-/-}$ mice, which wholly lack cDC1 [16], but was greatly reduced in

Csf2$^{-/-}$ mice, which have a 70% reduction in basal AM levels [17] **(Figure 4 A&B)**. The role of

AM was further examined by depleting these cells with clodronate liposomes [18], which we

verified efficiently removed AM but did not alter levels of other lung innate immune cells

(Figure 4Ci). Depletion of AM, by itself, did not significantly impact IAV titers or clinical signs

of disease but eliminated the ability of SFB to lower lung IAV load, ameliorate hypothermia and

weight loss, and prevent death **(Figure 4Cii)**. These results suggested that SFB-mediated impacts

on AM were germane to its protection against RVI. SFB-mediated maintenance of AM levels

could conceivably have reflected increased generation of these cells, which can occur from

monocyte progenitors migrating to the lung or local self-renewal [19], or that SFB colonization had

resulted in AM resisting death upon RVI. SFB's prevention of AM depletion following CA09

inoculation associated with prevention of IAV-induced caspase-3 activation **(Figure 4D)**, which

marks AM commitment to cell death [20], and increased expression of CD71 **(Figure 4E)**, which

associates with enhanced phagocytosis and survival [21,22]. Furthermore, SFB colonization

increased AM expression of proliferation marker Ki-67 **(Figure 4F)**. Collectively, these results

suggested that SFB colonization may have increased AM self-renewal and, furthermore,

"emboldened" AM to resist IAV-induced depletion, resulting in maintenance of a robust AM

population following IAV infection.

SFB's preservation of AM enables their sustained protection of EC from IAV infection

Airway epithelial cells (AEC) are primary replication sites of respiratory viruses,

including IAV, RSV, and SARS-CoV-2. Specifically, in the case of highly pathogenic H1N1

IAV, ciliated AEC are the predominant host for virus replication and generation of progeny

virions [23]. AM are known to protect lung AEC from IAV infection, particularly at early time

points following IAV-infection, i.e. prior to their depletion [24]. Thus, we predicted that SFB's

preservation of AM following CA09 inoculation might result in lower viral loads in AEC in

SFB$^+$ mice. Indeed, the percentage of lung AEC, type I and II, as well as conducting AECs,

displaying detectable levels of a recombinant CA09 that encodes GFP, by flow cytometry, was

significantly reduced in SFB$^+$ mice 4 days post-inoculation **(Figure 4G and Figure S5)**.

Depletion of AM by clodronate liposomes in SFB$^+$ mice increased IAV levels to that of SFB$^-$

mice in all three lung AEC types. These results indicate that SFB-induced AM persistence

following IAV infection enables sustained AM-mediated protection of EC, likely contributing to

lower IAV burden and ameliorated pathology.

Persistence of SFB$^+$ AM upon RVI is cell-intrinsic

Persistence of AM in CA09-infected SFB$^+$ mice was potentially attributable to SFB

altering the lung environment and/or reflect that AM themselves had been intrinsically

transformed to better withstand IAV challenge. To differentiate between these possibilities, we

adoptively transferred FACS-sorted AM from SFB$^+$ and SFB$^-$ mice to SFB$^-$ mice using donor

mice with a CD45.1 subtype, thereby enabling distinction of endogenous and transplanted cells

(Figure 5A). The degree to which AM harvested from SFB$^+$ and SFB$^-$ mice, hereafter referred to

as SFB$^+$ and SFB$^-$ AM, engrafted following transplant was similar, comprising about 15% of the

total AM in their new host. CA09 infection did not significantly alter proportions of endogenous

vs engrafted AM in recipients of SFB$^-$ AM but led to SFB$^+$ AM (i.e. engrafted) becoming their

host's majority AM population. This shift in AM proportions reflected that endogenous AM in

both recipient groups as well as engrafted SFB$^-$ AM were all depleted by CA09 infection,

whereas engrafted SFB$^+$ AM fully resisted CA09-induced depletion and, concomitantly, lacked

Caspse-3 activation, despite having been transplanted into an SFB-free host. Thus, colonization

of the gut with SFB intrinsically altered AM to withstand IAV-induced depletion.

The ability of transplanted SFB$^+$ AM to withstand IAV-induced depletion could

potentially have reflected that, in the donor mice, SFB colonization had phenotypically

transformed resident AM and/or led to these cells being replaced by newly recruited monocytic

precursors, which then became AM. That SFB$^+$ mice did not exhibit a basal or IAV-induced

elevation of monocytes chemo-attractants or monocyte-derived AM, both of which were readily

evident in IAV-infected SFB$^-$ mice, argued against the "AM replacement" mechanism **(Figure**

5B). Furthermore, administering SFB to mice after AM transplant led to similar persistence of

both CD45.2 (endogenous) and CD45.1 (exogenous) cells following IAV infection **(Figure 5C)**,

thus strongly indicating that SFB colonization changed the phenotype of lung resident AM.

SFB$^+$ AM exhibited inflammatory anergy and enhanced C1qa-mediated antiviral function

Differences between SFB⁺ and SFB⁻ AM were characterized *ex vivo* as schematized

(Figure 6A). Profiling of basal gene expression (i.e. without IAV exposure) by RNA-seq found

that only 24 of the 15,510 genes expressed by AM were differentially expressed between SFB⁻

and SFB⁺ AM **(Figure 6B)**, 18 of which were both confirmed by PCR and biologically

replicated in a distinct mouse cohort **(Figure S6A).** Six of the genes upregulated in SFB⁺ AM are

associated with an M2 macrophage phenotype. Yet, quantitation of surface expression markers

previously used to distinguish M1 vs M2 phenotypes [25] did not reveal differences between SFB⁺

and SFB⁻ AM **(Figure S6B)**. The genes most prominently up- and down-regulated, respectively,

in SFB⁺ AM were Tsc22d3 and Notch-4, which, respectively, function in suppressing and

activating pro-inflammatory gene expression [26,27], suggesting such signaling might be dampened

in SFB⁺ AM. Indeed, we found that SFB⁺ AM lacked the robust UV-irradiated IAV-induced

activation of pro-inflammatory gene expression exhibited by SFB⁻ AM **(Figure 6C and Figure**

S7C). Such inflammatory anergy was partially recapitulated by antibody-mediated neutralization

of Notch-4 suggesting that the SFB-induced reduction in AM Notch-4 expression in SFB⁺ AM

contributed to this *ex vivo* phenotype and, moreover, the reduced inflammation observed in IAV-

infected SFB⁺ mice. As noted above, in SFB⁻ mice, IAV infection resulted in depletion of AM,

which are then replaced by newly recruited monocytes, some of which will acquire AM cell

surface markers. Such "replacement AM" lacked any hint of inflammatory anergy. Rather, they

exhibited robust inflammatory responses to IAV, similar to those of the AM they replaced

(Figure S6H).

The small panel of genes upregulated in SFB⁺ AM also included some known to function

in antiviral resistance, including complement C1qa and Ifitm1, thus suggesting that SFB⁺ AM

were not anergic to IAV per se but, rather, might have better virus neutralizing capacity. We

examined this possibility by incubating CA09 stocks with AM for 45 min at which time AM

were removed by centrifugation and supernatant infectious titers and IAV genomes quantified.

SFB⁻ AM lowered the infectious titers of the CA09 stocks to which they were exposed, at 37°C

but not 4°C, but the extent of reduction was 150-fold greater for SFB⁺ AM (**Figure 6Di**). This

greater reduction in titer was accompanied by only 5-fold lower levels of PCR-quantifiable IAV

genomes in supernatants of SFB⁺ AM (**Figure 6Dii**). This led us to hypothesize enhanced

neutralization of IAV by complement and/or other products secreted by SFB⁺ AM. Indeed, we

found that, even when SFB⁺ AM were not exposed to virus, their supernatants from 18h cultures

had greater capacity than those from SFB⁻ AM to lower CA09 infectious titers **(Figure 6Diii)**.

Such enhanced ability of SFB⁺ AM supernatants to lower IAV titers was fully reversed by

antibody-mediated removal of C1qa and, moreover, was not observed in C1qa$^{-/-}$ SFB⁺ AM

(Figure S6C). These results accord with recent observations that C1qa can directly bind IAV

and reduce its infectivity [28]. In contrast to AM themselves, which when incubated with CA09

stocks lowered both levels of infectious virus (i.e. titer) and levels of viral RNA, AM

supernatants did not impact levels of viral RNA (**Figure S6E**) suggesting the enhanced antiviral

capacity of SFB⁺ AM might also involve increased viral absorption and/or uptake, possibly as a

result of viral particles being more decorated with C1qa [29]. In accord with this notion, reductions

in IAV genomes in supernatants of AM/CA09 mixtures were paralleled by increased IAV

genomes in AM lysates **(Figure 6Div)**. The relative increase in IAV genomes in SFB⁺, compared

to SFB⁻, AM lysates was maintained amidst chlorpromazine, but not latrunculin treatment,

neither of which impacted cell viability, suggesting it might reflect enhanced phagocytosis

uptake by SFB⁺ AM (**Figure S6G**). In contrast to our hypothesis, increased uptake of CA09 was

evident in C1qa$^{-/-}$ SFB⁺ AM (Figure S6D) suggesting SFB⁺ AM may be more phagocytic in

general. In accord with this notion, SFB$^+$ AM exhibited modestly greater uptake of IgG-coated

fluorescent beads *ex vivo* and bioparticles *in vivo* **(Figure S6F)**. Exploring the fate of CA09

internalized by AM during the 45 min exposure period found that, 24 h following media

replacement, AM supernatants had few infectious virions while lysates contained few detectable

IAV genomes **(Figure 6Div)** thus suggesting that AM were not permissive for CA09 replication,

but rather constituted a dead end for the virus. Collectively, these results indicate that elevated

C1qa expression and enhanced phagocytosis drove the increased antiviral capacity of SFB$^+$ AM.

Measure of C1qa protein by ELISA confirmed that SFB colonization increased

expression of C1qa in AM isolated from whole lung and bronchoalveolar lavage fluid (BALF)

and, moreover in whole BALF **(Figure 6E)**. Furthermore, experiments in *C1qa$^{-/-}$* mice indicated

that C1qa played a central role in SFB's suppression of IAV in vivo. Specifically, in *C1qa$^{-/-}$*

mice, SFB colonization did not lower viral titers or protect lung epithelial cells from IAV

infection following CA09 inoculation (Figure 6F). Concomitantly, SFB's ability to ameliorate

disease symptoms, including weight loss and hypothermia was markedly reduced in *C1qa$^{-/-}$* mice

relative to *C1qa$^{+/+}$* mice. Nonetheless, colonization of *C1qa$^{-/-}$* mice with SFB still conferred

modest but statistically significant reductions in weight loss and mortality. Moreover, SFB

colonization of *C1qa$^{-/-}$* suppressed CA09-induced pro-inflammatory gene expression, gross lung

pathology, and infiltration of inflammatory cells into the lungs. The reduced but not eliminated

ability of SFB to suppress symptoms of IAV infection in *C1qa$^{-/-}$* mice associated with SFB

colonization partially protecting AM against IAV-induced depletion. Thus, C1qa was germane to

SFB's ability to impede IAV infection and its consequences but nonetheless SFB ameliorated

IAV-induced lung inflammation irrespective of C1qa and viral titer.

SFB-induced reprogramming of AM protects mice against IAV infection

The starkly different transcriptional responses of SFB⁻ and SFB⁺ AM led us to investigate the general metabolic states of these cells via Seahorse analysis. The basal metabolic states of SFB⁻ and SFB⁺ AM differed only modestly with SFB⁺ AM exhibiting a small reduction in glycolysis (Figure S7A). In contrast, inoculating SFB⁻ mice with CA09 led to stark increases in AM ATP production and glycolysis, which were wholly absent in SFB⁺ mice thus paralleling above-described changes in inflammatory parameters in UV-IAV induced changes in AM proinflammatory gene expression observed *ex vivo*. Lastly, we examined the extent to which SFB-induced reprogramming of AM contributed to SFB's protection against IAV infection. Specifically, we performed an additional series of AM transplant experiments designed to measure impacts on IAV levels and clinical-type parameters. Consistent with the notion that AM, prior to their depletion, protect against IAV infection [24], we observed that administration of SFB⁻ AM to a host replete with endogenous AM resulted in a modest reduction in IAV titers, gross lung damage, hypothermia, weight loss, and death following CA09 inoculation **(Figure 7A)**. Administration of SFB⁺ AM provided markedly greater protection against this IAV challenge, approximating the level conferred by administration of SFB itself, as assessed by all parameters measured. Analogous patterns of results were seen following transfer of SFB⁺ and SFB⁻ AM into mice deficient in endogenous AM, achieved via use of *Csf2*⁻ᐟ⁻ mice or chlodronate administration **(Figure 7B and 7C)**. Analogous to results obtained in *C1qa*⁻ᐟ⁻ mice, transplant of C1qa⁻ᐟ⁻ SFB⁺ AM into SFB⁻ mice did not reduce viral titers or protect lung cells from IAV infection following CA09 inoculation but nonetheless reduced indices of inflammation and conferred a modest clinical-type benefit **(Figure S7B)**. *Ex vivo* blockade of Notch-4 on SFB⁻ AM prior to transplanting them also enhanced their ability to withstand IAV-induced depletion,

332   protect EC from IAV infection, and ameliorate CA09-induced disease, but did not impact level

333   of viral genomes in the lung **(Figure S7C)**, suggesting the ability of transplanted SFB$^+$ AM to do

334   so was not mediated solely by dampened inflammatory signaling but also required such AM

335   have enhanced antiviral function. Thus, SFB colonization of the gut stably and broadly altered

336   the AM phenotype enabling these cells to better protect their hosts from lethal RVI.

DISCUSSION

Positioned in the lung lumen to enable their guarding the surfaces of O_2 and CO_2

exchange, AM are amongst the first cells to encounter inhaled threats including respiratory

viruses. We show herein that how AM respond to respiratory viruses is dramatically altered by

the composition of the intestinal microbiota. Specifically, we report that colonization of the

intestine by SFB, naturally acquired, or exogenously administered, markedly attenuates IAV-

induced AM pro-inflammatory gene expression while increasing ability of AM to disable IAV.

This seemingly paradoxical phenotypic differences between SFB^- and SFB^+ AM is reminiscent

of the comparison of blood monocytes and intestinal macrophages wherein the latter are more

phagocytic but yet exhibit inflammatory anergy [30]. An important aspect of SFB-induced

reprogramming of AM is that it resulted in resistance of RVI-induced depletion and thus

sustained protection of lung epithelial cells. Collectively, the consequences of SFB-induced AM

reprogramming resulted in lower viral titers, reduced inflammation, and reduced disease severity.

While the general concept that microbiota composition in total can influence immune

phenotype and, proneness to infection is established [31,32], we were surprised that the presence of

a single common commensal bacterial species, amidst the thousands of different microbial

species that variably inhabit the mouse gut, had such strong impacts in RVI models and,

furthermore, that such impacts were largely attributable to reprogramming of AM. The extent to

which SFB is present in humans is not well defined but it has been reported to be frequently

present in children in some regions [33] and was unequivocally shown to be present in some adult

individuals [34]. Independent of a specific role for SFB, however, our findings speak strongly to

the potential of gut microbiota to influence proneness to RVI, especially via programming of

AM. Both AM pro-inflammatory gene expression and AM depletion are known to correlate with

RVI severity in humans but determinants of inter-individual heterogeneity in AM responses have

not been defined [33]. We consider it highly unlikely that SFB is the only gut microbe capable of

impacting AM phenotype and, consequently, proneness to RVI. Rather, we posit that AM are a

key component of the gut-lung axis wherein an individual's microbiota in its entirety is a

determinant of disease outcome following exposure to an array of respiratory viruses.

The observation that SFB[+] AM maintained an IAV-resistant phenotype following

transplant to SFB[-] mice is consistent with observations that AM have stable cell-type specific

patterns of gene expression and methylation even when cultured extensively *ex vivo* [35].

Moreover, it accords with recently increasing appreciation that AM can be "trained" to display

long-lasting phenotypic alterations that impact their ability to manage challenges. Yet the

consequences and mechanisms of previously observed AM training seems quite distinct from

that reported herein. For example, administration of live, but not heat-inactivated BCG vaccine

resulted in systemic low-grade infection and inflammation that altered gut microbiota

composition and increased gut permeability, potentiating AM pro-inflammatory gene expression

[36]. In addition, compared to BCG, SFB colonization had a somewhat opposite outcome on AM

function in that it suppressed rather than promoted pro-inflammatory gene expression. In further

contrast to BCG, phenotypic impacts of SFB were maintained in gnotobiotic mice, arguing

against broad changes in microbiota composition playing a central role. Another starkly different

example of AM training is the recent observation that IAV infection increases ability of AM to

suppress tumors [37]. Such AM training is mediated by a large-scale depletion of AM, which are

then replaced by cells newly recruited from bone marrow that, even after establishing residency

in the lung, exhibit stark global differences in gene expression profiles. In contrast, SFB

colonization phenotypically transformed resident AM without a detectable influx of monocyte

precursors. Moreover, gene expression profiling of SFB$^-$ and SFB$^+$ AM revealed only modest

differences, likely reflecting that both SFB$^-$ and SFB$^+$ AM had been generated by AM self-

renewal processes that normally maintain AM homeostasis in non-traumatic states.

386   Despite being modest, SFB-induced changes in basal AM gene expression contributed

mightily to the enhanced ability of SFB-colonized mice to manage IAB infection. Most notably,

use of $C1qa^{-/-}$ mice, and AM transferred therefrom, indicated that SFB-induced increases in C1qa

expression were pivotal for lowered viral titers. The comparable genetically-modified mice to

similarly examine the necessity of genes that mediated SFB's suppression of AM inflammatory

signaling are not yet in hand. However, a converse approach, namely use of a Notch-4

neutralizing antibody, argued that the SFB-induced reduction in Notch-4 expression was

germane to SFB-induced AM inflammatory anergy and that this aspect of the SFB$^+$ AM

phenotype also contributed to protection against IAV. In addition to directly disabling IAV, AM

can also protect lung EC from IAV infection via suppressing EC CysLT1 signaling, which might

mediate IAV entry into these cells [24]. Thus, persistence of AM in SFB$^+$ mice, by itself, likely also

contributed to SFB's protection against IAV. That mechanisms driving RVI-induced AM

depletion are not clear, makes determining how SFB colonization prevents such depletion

challenging. Indeed, deciphering the interrelationships between reduced viral titers,

inflammatory anergy, and increased AM survival, is a complex endeavor in which some events

are likely both a cause and consequence of each other. For instance, collectively our results

indicate that AM persistence enabled them to suppress IAV, but yet such suppression of IAV

likely contributed to increased AM survival. Conversely, we envisage that SFB-induced AM

inflammatory anergy promoted AM persistence, which in turn enabled sustained protection of

epithelial cells from IAV infection. Such protection of epithelial cells from IAV infection would

reduce the viral burden, which likely contributes to both inflammation and AM death. Hence,

defining the relative contributions of enhanced antiviral and suppressed inflammation is a

complex undertaking. Nonetheless, our current working hypothesis is that elevated C1qa

expression plays a leading role, and inflammatory anergy plays a key supporting role, in

mediating SFB's protection against RVI.

Less clear is the nature of the signal that is generated by SFB colonization of the luminal

surface of the intestine that drives AM reprogramming. We speculate that SFB produces, or

induces host production of, a metabolite(s) that reach the lung and act upon AM, changing both

their basal gene expression and how they respond to challenge, perhaps by altering DNA

methylation. Additionally/alternatively, it is possible that gut to brain and/or brain to lung neural

signaling is involved. We envisage that deciphering how SFB impacts lung AM may yield

strategies to mitigate disease burden cause by RVI.   Indeed, in addition to mitigating primary

RVI, AM persistence may help prevent secondary consequences of RVI. Many of the severe

consequences of IAV infection in humans result from secondary bacterial infections, to which

the lung is more permissive by loss of IAV-induced AM depletion [38]. Such impaired antibacterial

immunity can persist for prolonged periods perhaps reflecting that the bone-marrow derived

replacement AM are not quickly optimized for mucosal host defense. Thus, microbiota mediated

AM programming may not only influence severity of acute RVI but may also be a long term

post-RVI health determinant.

Acknowledgments:

We thank Tim Denning and Ifor Williams for insightful guidance. We thank RS Baric for SARS-CoV-2

MA10 virus isolate and Georgia State University's High Containment Core and Division of Animal

Resources for support. **Funding**: This work was supported, in part, by Public Health Service grants

AI141222 (to RKP), AI171403 (to RKP), and AI170014 (to ATG) from the NIH/NIAID. The funders had

no role in study design, data collection and interpretation, or the decision to submit the work for

publication.

Author contributions: Conceptualization: RKP, ATG. Investigation: VLN, CML, HK, KS. Resources:

RKP, ATG. Visualization: VLN, CML, KS, RKP, ATG. Validation: RKP, ATG. Funding acquisition:

RKP. Project administration: RKP, ATG. Supervision: RKP, ATG, MK. Writing – original draft: ATG.

Writing – review & editing: VLN, CML, KS, RKP.

Declaration of interest: RKP reports contract testing from Enanta Pharmaceuticals and Atea

Pharmaceuticals, and research support from Gilead Sciences, outside of the described work. All other

authors declare that they have no competing interests.

Data and materials availability: All data are available in the main text or the supplementary materials.

Figure 1. Colonization of the intestine with SFB protected mice from influenza virus infection.

(A) EF and SPF mice were inoculated with A/CA/07/2009 (H1N1) influenza virus (CA09) as schematized. Lung viral titers, survival, bodyweight, and temperature were monitored.

(B) As schematized, conventionally colonized SFB⁻ mice received feces that lacked SFB (SFB⁻), contained SFB (SFB⁺), or contained SFB but no other bacteria (SFB-MA). Seven days post-transplant, mice were inoculated with CA09. Mice were euthanized 4.5-days later or monitored daily for survival, body weight, and temperature for 2 weeks. Statistical analyses were performed with SFB⁻ as a control group.

(C) SFB⁻ mice were administered SFB 3 months or 7 days prior to CA09 inoculation. **(i)** qRT-PCR analysis of SFB colonization level over the course of 14 weeks. **(ii)** Mice were euthanized 4.5-days later or monitored daily for survival, body weight, and temperature for 2 weeks.

(D) qRT-PCR analysis of SFB level in the small intestine, lungs, and spleens after 7 days colonization.

(E) SFB⁻ mice were untreated (naive) or administrated SFB. Seven days later, mice were inoculated (or not) with CA09. Mice were euthanized 4.5 days later. H&E staining lung sections scored for pleuritis, bronchiolitis, alveolitis, vasculitis, interstitial pneumonia (IP), and perivascular cuffing (PVC).

All experiments n = 5 mice per group. Data is representative of two to three independent experiments. yielding an identical pattern of results. Statistical analyses of survival, bodyweight change, and temperature of (B) and (C) were conducted by comparing "SFB-"or "-SFB" group to all other groups. Statistical analysis; Viral titer: One-way ANOVA or student *t* test. survival:

Log-rank Mantel-Cox test. Body weights and temperature: Two-way ANOVA. $*p<0.05$,

$**p<0.01$, $***p<0.001$, ns not significant.

Figure 2. SFB mediated protection against IAV was maintained in an array of knockout

mice and did not require complex gut microbiota.

Lung viral titer, survival rate, body weight change, and temperature of: SFB-negative mice were

colonized with SFB for 7 days then inoculated with CA09.

(A) *rag1$^{-/-}$* mice

(B) C57BL/6 mice received neutralization of IL-17/IL-22 monoclonal antibodies every other day

throughout the experiment

(C) *Ifnar$^{-/-}$* mice

(D) *rag1$^{-/-}$yc$^{-/-}$* mice

(E) *stat1$^{-/-}$* mice

(F) Altered Schaedler flora (ASF) mice.

All experiments n = 5 mice per group. Statistical analysis; Viral titer: student *t* test, Survival:

Log-rank Mantel-Cox test. Body weights and temperature: Two-way ANOVA. $**p<0.01$,

$***p<0.001$.

Figure 3. SFB's protection against IAV infection associated with preservation of resident lung phagocytes.

(A) SFB$^-$ mice were untreated (naïve) or SFB was administered. Seven days later, mice were inoculated (or not) with CA09. Mice were euthanized 4.5-days later. **(i)** Whole lung gene expression assayed by RNA-seq and data are displayed as heatmap of genes that are significantly altered by 2-fold in all treatment groups relative to the naïve group. **(ii)** Quantitation of CA09 level in these extracts by qRT-PCR.

(B) SFB$^-$ mice were untreated (naïve) or SFB was administered. Seven days later, mice were inoculated (or not) with CA09. Mice were euthanized 4.5-days later, and lung assayed by flow cytometry. **(i & ii)** Analysis of dendritic cells and alveolar macrophage; representative flow plot and frequencies and cell numbers **(i)** cDC1. **(ii)** alveolar macrophages. **(iii)** Frequencies and cell numbers of lung neutrophils, inflammatory monocytes, and NK cells.

(C) (i) Experimental schema: SFB$^-$ mice colonized with SFB, inoculated with CA09, and treated with 4'-FlU at indicated doses. On day 4.5 post virus inoculation, mice were euthanized, and lungs were harvested to determine virus titers and FACs analysis. **(ii)** lung viral titer. **(iii)** FACs plot graphical representation, frequencies, and cell numbers of lung cDC1, and AMs. **(iv)** Cell numbers of lung neutrophils, inflammatory monocytes, and NK cells.

All experiment n = 4-5 per group. Data is representative of two to three independent experiments (B and C), yielding an identical pattern of results. Statistical analysis: One-way ANOVA. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$, ns not significant.

Figure 4. SFB-mediated protection against IAV infection and resulting disease required alveolar macrophages.

(A-B) (A) *batf3*⁻/⁻ and **(B)** *csf2*⁻/⁻ mice were administered SFB and then, 7 days later, inoculated with CA09. Lung viral titer, body weight change, temperature, and survival were monitored.

(C) (i-ii) SFB⁻ mice were orally administered with clodronate liposomes one day before mice were euthanized. FACS analysis to determine the AMs depletion efficacy after clodronate treatment. **(i)** FACS graphical representation of lung AMs, frequency of lung AMs and others immune cells after clodronate treatment. **(ii)** SFB⁻ mice were colonized with SFB for 7 days, and clodronate liposomes were orally administered to mice one day before CA09 inoculation. Mice were euthanized on day 4 post-virus inoculation. Lung viral titer, survival rate, body weight, temperature was measured. Statistical analyses of survival, bodyweight change, and temperature were conducted by comparing "SFB+CA09" group to all other groups.

(D) SFB⁻ mice were untreated (naïve) or SFB was administered. Seven days later, mice were inoculated (or not) with CA09. Mice were euthanized 4.5-days later. Level of cleaved caspase-3 of alveolar macrophages in lung were measured with FACS.

(E-F) SFB⁻ mice were untreated or treated as in (D). The level of **(E)** CD71 and **(F)** Ki-67 expression of alveolar macrophages in the lungs were measured with FACS.

(G) SFB⁻ mice treated as described in (C) FACS plot graphical representation, and frequencies CA09 level in T1ACEs, T2AECs, and Conducting AECs were assayed by flow cytometry.

All experiment n = 5 mice per group. Data is representative of two to three independent experiments (B-F), yielding an identical pattern of results. Statistical analysis; Viral titer, relative SFB genomes, MFI, frequencies/cell numbers of immune cell data: One-way ANOVA or student

527   *t* test. Survival data Log-rank Mantel-Cox. Body weights, temperature: Two-way ANOVA.

528   *p<0.05, ***p<0.001, ****p<0.0001, ns not significant.

530 **Figure 5. SFB colonization intrinsically altered AM to withstand IAV-induced depletion.**

531 **(A)** As schematized, SFB⁻ CD45.1 mice were administered with SFB or PBS. Seven days later,

532 AM were isolated by FACs-sorting, and transferred to SFB- CD45.2 mice. One day post-transfer

533 mice were inoculated with CA09. Four days later, lungs were harvested, and AM analyzed by

534 FACs. Representative FACs plots, cell frequencies, cell numbers, and cleaved caspase-3 level

535 are shown.

536 **(B)** SFB⁻ mice were untreated (naïve) or administrated SFB. Seven days later, mice were

537 inoculated (or not) with CA09. Mice were euthanized 4.5-days later. **(i)** qRT-PCR analysis of

538 monocytes chemo-attractants *Ccl2* and *Cxcl10* from lung tissues. **(ii)** Representative FACs plots,

539 cell frequencies, and ratio of tissue residents AM (TR-AM), and monocyte-derived AM (Mo-

540 AM).

541 **(C) (i)** As schematized, AM isolated from CD45.1 mice were transferred to CD45.2 mice. One

542 day post-transferred, mice were administered with SFB or PBS for seven days then mice were

543 inoculated with CA09. For day later, lungs were harvested, and AM analyzed by FACs. **(ii)** Cell

544 numbers are shown.

545 All experiment n = 5 mice per group. Data is representative of two independent experiments (A-

546 B), yielding an identical pattern of results. Statistical test: One-way ANOVA. **p<0.01,

547 ***p<0.001, ****p<0.0001, ns not significant.

548

Figure 6. SFB⁺ AM exhibited inflammatory anergy and enhanced antiviral function *ex vivo*.

(A) Schematic design for *ex vivo* studies of AM.

(B) Comparison of basal SFB⁺ and SFB⁻ AM gene expression by RNA-seq, shown as a volcano

plot.

(C) SFB⁺ and SFB⁻ AM were exposed to UV-CA09 with or without Notch4 neutralizing, or

isotype control antibodies. RNA was harvested 24 hours later, and gene expression assayed by

RT-qPCR shown as a heatmap.

(D) AM and CA09 stocks were incubated at 37°C, or indicated temperature, for 45 minutes and

supernatants were collected. In addition, some cells had new media added and were incubated for

another 24 hours. **(i)** Vius infection titers in supernatant at 45 minutes and 24 hours were

assayed. **(ii)** Supernatants were collected after 45 minutes incubated with CA09 and viral

genomes quantitated by RT-qPCR. **(iii)** Cell-free supernatant of 15 hours AM cultures were

incubated with live CA09 stocks, with or without C1qa-neutralization or isotype antibody for 45

minutes, at which point virus infection titers were assayed. **(iv)** Cell lysates were generated upon

CA09 removal or 24 hours later and viral genomes quantitated by RT-qPCR.

(E) As schematized, mice were administered with SFB. Seven days later, bronchoalveolar lavage

fluid (BALF) and lung were harvested, and AM were isolated via FAC-sorting. C1qa secretions

were assayed by ELISA.

(F) (i) 15 weeks old *C1qa⁺/⁺* and *C1qa⁻/⁻* SFB⁻ mice were untreated (naïve) or administrated SFB

and then, 7 days later, inoculated with CA09. Mice were euthanized and lungs were harvested

4.5-days post CA09 inoculation. **(ii)** Images of lungs. **(iii)** Survival, body weight change,

temperature. **(iv)** Lung viral titer. **(v)** Frequencies CA09 level in T1ACEs, T2AECs, and

571 Conducting AECs were assayed by FACs. **(vi)** Gene expression assayed by RT-qPCR shown as

572 a heatmap. **(vii)** Cell numbers of neutrophils, inflammatory monocytes and AM assayed by

573 FACs. Statistical analyses of survival, body weight change, and temperature were conducted

574 between *C1qa*$^{-/-}$ and *Cq1a*$^{-/-}$+SFB group.

575 All experiment n = 4-5 mice per group. Data is representative of two independent experiments,

576 yielding an identical pattern of results. Statistical analysis: One-way ANOVA. *$p<0.05$,

577 **$p<0.01$, ****$p<0.0001$, ns not significant.

Figure 7. Adoptive transfer of SFB$^+$ AM enabled recipients to better manage IAV infection.

(A) As schematized, SFB$^-$ mice were administered SFB or PBS. Seven days later, AMs were isolated by FACs-sorting and transferred to SFB$^-$ mice. Mice were administered 5×10^5 cells per mouse. One day post adoptive transfer, mice were inoculated with a lethal dose of CA09. Mice were euthanized 4.5 days post-inoculation for viral titer and gross lung assessment or monitored for survival, body weight, and temperature. Statistical analyses of survival, body weight change, and temperature were conducted using the "PBS" group as the control group.

(B) As schematized, AMs from SFB$^-$ *csf2*$^{+/+}$ mice were administered SFB or PBS. Seven days later, AMs were isolated by FACs-sorting and transferred to SFB$^-$ *csf2*$^{-/-}$ mice. One day post-transfer, mice were inoculated with CA09. Mice were monitored for 14 days for clinical symptoms including survival rate, body weight, and temperature.

(C) SFB$^-$ mice were administered clodronate liposomes to remove endogenous AM populations. Seven days later after initiation of depletion, mice were administered 5×10^5 AM cells harvested from the SFB$^-$ or SFB$^+$ mice. One day post transfer, mice were inoculated with CA09. Mice were euthanized on day four to assess lung viral titer and gross lung assessment or monitored for survival, body weight, and temperature.

All experiments have at least n = 5 mice per group. Data is representative of two independent experiments (A), yielding an identical pattern of results. Statistical analyses of survival, bodyweight change, and temperature of (A-C) were conducted by comparing "PBS" group to all other groups. Statistical analysis; One-way ANOVA: viral titers. Survival: Log-rank Mantel-Cox test. Body weight, temperature: Two-way ANOVA. *$p<0.05$, **$p<0.01$, ****$p<0.0001$, ns not significant.

STAR METHODS

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to and will be

fulfilled by the lead contact, Andrew Gewirtz (agewirtz@gsu.edu).

Materials availability

This study did not generate new unique reagents.

Data and code availability

- The raw sequencing data has been deposited at GenBank with access number: PRJEB71689, and PRJEB71449, and all other data used for this study will be shared upon reasonable request from the lead contact.

- This paper does not report original codes.

- Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon reasonable request.

EXPERIMENTAL MODEL DETAILS

Virus strains and growth conditions

All viruses and cells used in this study are listed in the key resources table. Madin-Darby Canine

Kidney (MDCK; ATCC CCL-34), HEp-2 (ATCC CCL-23), and VeroE6 TMPRSS2 (BPS

Bioscience #78081) cells were cultured in Dulbecco's Modified Eagle's Medium (DMEM)

supplemented with 7.5% heat-inactivated fetal bovine serum (FBS) at 37°C and 5% CO_2. All cell

lines used in this study were authenticated and checked for *Mycoplasma* and microbial

contamination. Recombinant Influenza A virus was recovered based on original reports for

627   laboratory adapted IAV [39]. A/CA/07/2009 (H1N1) was grown on MDCK cells using serum-free

628   DMEM supplemented with 0.5% Trypsin. Recombinant RSV strain A2 with line19F reporter

629   gene recRSV-mKate was rescued and amplified as described previously [40], and the SARS-CoV-2

630   MA10 strain [41] was passaged in VeroE6 TMPRSS2 cells. All viruses used were clarified,

631   aliquoted, and stored at -80°C. Virus stock titers were determined through $TCID_{50}$-titration.

Complex microbiota and mono-associated SFB transplantation

634   Fecal samples were collected from donor mice, suspended in 20% glycerol PBS solution at 40

635   mg/mL concentration, passed through a 40 µm filter, and stored at -80C. Frozen fecal

636   suspensions were orally administered to recipient mice at 200 µl per mouse.

Virus infections

All viruses and mice strains used in this study are listed in the key resources table. The

Institutional Animal Care and Use Committee of Georgia State University approved animal

studies. All mice, unless otherwise stated, were purchased from The Jackson Laboratory. The

Jackson Laboratory does not perform testing for segmented filamentous bacteria (SFB) in mice.

Nevertheless, all mice obtained from this vendor were tested and confirmed to be SFB-negative

upon their arrival at GSU. The following mice were purchased from Taconic Biosciences:

C57BL/6 (B6 WT) Excluded Flora (EF) mice and Specific Pathogen-Free (SFP) mice, which we

verified were SFB+ and SFB-, respectively. Unless otherwise stated, mice were 5 weeks of age

at the time of infection. Upon arrival, mice were acclimated for five days and then randomly

assigned to study groups and housed under ABSL-2 or ABSL-3 conditions for infections with

A/CA/07/2009 (H1N1) recombinant, recRSV-mKate, or SARS-CoV-2 MA10 strain,

respectively. Seven days prior to virus infection, mice were subjected to microbiota manipulation

(administration of feces from SPF or SFB-monoassociated mice). Bodyweight was determined

twice daily, and body temperature determined once daily rectally. For infections, animals were

anesthetized with isoflurane [rec A/CA/07/2009 (H1N1) (CA09)] or ketamine/xylazine (recRSV-

mKate and MA-SARS-CoV-2), followed by intranasal inoculation with 2.5×10^2- 2×10^4 $TCID_{50}$

units/animal of GFP-expressing reporter virus version of A/CA/07/2009 (H1N1) (CA09),

5.5×10^5 of recRSV-mKate, or 4.2×10^5 of MA-SARS-CoV-2. In all cases, virus inoculum was

diluted in 50 µl and administered in amounts of 25 µl per nare, followed by treatment with 4'-

FlU at specified dose levels and starting time points through oral gavage. Each individual study

contained animals receiving equal volumes of vehicle through oral gavage. Animals were

euthanized and organs harvested at predefined time points or when animals reached humane

study endpoints. At study end point, lung tissue was harvested for virus titration or

histopathology.

METHODS DETAILS

PCR primers, FACs antibodies, neutralization antibodies and other reagents

All reagents used in these manuscripts can be seen in the key resources table.

Histopathologic analysis

Lungs were perfused using 10% neutral-buffered formalin, dissected, and fixed. Formalin-fixed

lungs were transferred to 70% EtOH, embedded in paraffin, sectioned, stained, and scored by a

board-certified veterinary pathologist (KS), who was blinded to the study groups. For each

parameter, the most severe lesion in the section was scored. Pleuritis, bronchiolitis, and alveolitis

scoring was based on distribution: 1 = focal, 2 = multifocal, 3 = multifocal to coalescing, 4 =

diffuse. For interstitial pneumonia (IP): 1 = alveolar septa expanded by 1 leukocyte thickness, 2

= 2-leukocytes thick, 3 = 3-leukocytes thick, 4 = 4-leukocytes thick. For vasculitis: 1 = vessel

wall infiltrated by leukocytes, 2 = infiltration and separation of myocytes, 3 = same changes as in

2 with fibrinoid change. For perivascular cuffing (PVC): 1 = vessels surrounded by 1 leukocyte

layer, 2 = 2 to 5-leukocytes thick, 3 = 6 to 10-leukocytes thick, 4 = more than 10-leukocytes

thick.

Virus titration

Virus samples were serially diluted (10-fold starting at 1:10 initial dilution) in serum-free

DMEM (recRSV-mKate and MA-SARS-CoV-2) and supplemented with 0.5% Trypsin (Gibco,

recA/California/07/2009). Serial dilutions were added to cells seeded in 96-well plate at

1×10^4 cells per well, 24 hours before infection. Infected plates were incubated for 72 hours at

37°C with 5% CO_2 and scoring of wells based on reporter activity/syncytia.

Virus titration from tissue samples

Organs were weighed and homogenized in 300 µl PBS using a bead, set to 3 cycles of 30

seconds each at 4°C, separated by 30-second rest periods. Homogenates were cleared (10

minutes at 20,000 × g and 4°C), and cleared supernatants were stored at -80°C until virus

titration. Viral titers were expressed as $TCID_{50}$ units per gram input tissue.

Neutralization of IL-17 and IL-22

IL-17 (100 μg) and IL-22 (150 μg) neutralizing monoclonal antibody were administered

intraperitoneally before microbiota transplantation and virus inoculation thereafter every other

day until mice were euthanized.

Lung digestion and flow cytometry analysis/isolation of innate immune cells

Lungs were dissected from mice and finely minced with scissors before digestion (40 minutes, at

37°C) with warm RPMI containing 10% FBS, DNase, and 1mg/mL Collagenase type IV

(Sigma); lungs were further homogenized via swishing with a 3-mL syringe every 10 minutes

during digestion. Following digestion, immune cells were purified with 40%/80% isotonic Percol

gradient centrifugation. For flow cytometry analysis, the prepared cells were blocked with 1

μg/million cells with 2.4G2 (anti-CD16/anti-CD32) in 100 μL PBS for 15 minutes at 4°C,

followed by 2-times wash with PBS to remove residue of 2.4G2 and then incubated in

conjugated mAbs CD45 BV605, MHCII BV650, CD11b APC-Cy7, CD11c BV786, Ly6C

PerCp-Cy5.5, Ly6G AF700, CD64 BV421, CD24 FITC, CD117 GV510, SiglecF PE-CF594,

Fcer1 APC. Intracellular staining for cleaved caspase-3 was done using FoxP3 staining buffer

purchased from eBioscience. Multi-parameter analysis was performed on a CytoFlex (Beckman

Coulter) and analyzed with FlowJo software (Tree Star). Cell sorting was done using Sony

SH800Z cell sorter (Sony). Gating strategy for innate immune cells was based on Yu *et al*., [42]

and gating strategy for airway epithelial cells was based on Cardani *et al*., [24]. Immune cells and

lung airway epithelial subtypes are defined in Table S1.

RNA-Seq

RNA was isolated from lungs or FACs-sorted alveolar macrophages using RNeasy Plus Mini Kit

from Qiagen. The prepared RNA samples were then sent to the Molecular Evolution Core of

Georgia Institute of Technology for library preparation and sequencing on the NextSeq

instrument with a high output 2x75bp run. FASTQC was used for sequencing reads quality

screening. The RNAseq data were then analyzed using the Galaxy server.

Adoptive transfer of alveolar macrophages

Lungs were harvested from CD45.1 mice and digested with Collagenase type IV as described

above. Alveolar macrophages (defined at CD45$^+$MHCII$^+$CD11c$^+$Siglect$^+$) were stained and

FACs-sorted. Small aliquot of isolated cells was used to determined purity. Unless stated

otherwise, $5x10^5$ alveolar macrophages in 30 µl PBS were administered intranasally to recipient

mice. One day post cells transfer, some mice were euthanized to determine transfer efficiency

and survival of transferred cells; and some mice were inoculated with IAV. For survival

experiments, mice were monitored twice daily for 14 days. For lung virus titers, mice were

euthanized on day 4.5 post virus inoculation.

Ex vivo **inhibiting Notch-4 signaling in alveolar macrophages.**

Alveolar macrophages were FACS-sorted as described. After sorting, cells were treated with

either isotype control antibody (Cat# BE0091, BioXCell) or an anti-Notch4 monoclonal antibody

(20µg/mL) (Cat# BE0129, BioXCell), known to neutralize Notch-4 [27] for 30 minutes. Cells

were washed twice and resuspended in PBS and intranasally transferred to mice.

Ex vivo **alveolar macrophages culture, virucidal assay, and anti-body mediated removal of**

C1qa. Alveolar macrophages were FACS-sorted as described. After sorted cells were incubated

for 18h at 37°C. Cells were washed twice with serum-free media and were incubated with live

CA09 (MOI = 1) in serum-free DMEM supplemented with 0.5% Trypsin (flu media) for 45min.

In addition, to assess whether alveolar macrophages are the terminal destination for IAV, some

cells had new flu media added and were incubated for another 24h. Viral titer and genomes were

determined via $TCID_{50}$ and PCR methods. **Antibody-mediated removal of C1qa from *ex vivo***

alveolar macrophages' supernatants. Cell-free supernatants of 18h AM cultures were treated

with C1qa polyclonal antibodies or isotype for 2 hours at 4°C and protein A/G was added and

centrifuged. Supernatant was harvested and incubated with CA09 stocks for 45mins, at which

point IAV infection titers were assayed.

Alveolar macrophages phagocytosis assay. *Ex Vivo* alveolar AM bead uptake assay was

performed with Phagocytosis assay kit (Cat#500290, Cayman chemical). Alveolar macrophages

were FACS-sorted as described. After sorting, 1×10^5 cells were seeded in each well of a 96-well

plate and incubated for 18h at 37°C. Cells were washed twice, and IgG-FITC beads were added

and incubated for 2 hours at 37°C. After incubation period, cells were centrifuged, and

supernatants were removed and resuspended FACS buffer. Level of cell fluorescence, an

indicator of bead uptake, was measured via flow cytometry. For *In vivo* AM bioparticle uptake,

SFB⁻ and SFB⁺ mice were administered pHrodo™ *E.coli* BioParticles™ conjugated (Cat#

P35360, Invitrogen). Three hours later, mice were euthanized and BALF were harvested, and

AM were labeled. Level of cell fluorescence, an indicator of bioparticle uptake, was measured

via flow cytometry.

Quantitative real-time PCR

Total RNA from lungs or from FACs-sorted alveolar macrophages were isolated using the

Qiagen RNeasy Mini Kit with on-column DNase digestion according to manufacturer's protocol.

cDNA was generated using the Superscript First Strand Synthesis kit for RT-PCR and random

hexamer primers (Invitrogen). RT-qPCR was performed with SYBR Green using StepOnePlus

PCR system (Applied Biosystem), and all genes expression was normalized to GAPDH.

C1qa ELISA

BALF was collected from SFB$^+$ and SFB$^-$ mice. Alveolar macrophages (AM) were sorted using

fluorescence-activated cell sorting (FACS) from lung and BALF of SFB$^+$ and SFB$^-$ mice. Sorted

AM were incubated in RPMI + 10% FBS overnight at 37°C in a cell culture incubator. BALF

and supernatants from cultured AM were coated onto 96-well ELISA plates and incubated

overnight at 4°C. Plates were washed with PBST to remove unbound material. Blocking was

performed with 3% bovine serum albumin (BSA) in PBS at room temperature for one hour.

Polyclonal antibody against C1qa was diluted at 1:5000 and added to the plate and incubated for

2h at room temperature. Secondary antibodies (conjugated to horseradish peroxidase, HRP) were

added and incubated at room temperature for 1 hour. Tetramethylbenzidine (TMB) substrate was

added to detect HRP enzyme conjugate activity. Absorbance at 450 nm was measured to quantify

the presence of C1qa. Mouse C1qa recombinant protein was serially diluted to construct a

standard curve for determining the concentration of C1qa in the samples. Absorbance values of

the samples were compared with the standard curve to determine the concentration of C1qa in

the BALF and cultured AM supernatants.

AM metabolic assay.

The real-time metabolism of AM was assessed using the Seahorse XF Cell Mito Stress Test Kit

or the Seahorse XF Glycolytic Stress Test Kit from Agilent Technologies, following the

manufacturer's instructions. AM were FACS-sorted from SFB$^+$ and SFB$^-$ mice. The sorted cells

were then seeded into Seahorse cell culture plates at a density of 1×10^5 cells per well and

cultured overnight in complete RPMI1640 medium. On the following day, AM were incubated

with Seahorse assay medium (Agilent Technologies) for one hour. For the glycolysis stress test,

AM were supplemented with 2mM glutamine. For the mito stress test, AM were supplemented

with 10mM glucose and 1mM pyruvate. All reagents provided in the kits were resuspended in

assay medium, and the final concentrations in the culture wells were as follows: 1.5 μM

oligomycin, 4 μM FCCP, and 0.5 μM rotenone/antimycin A for the mito stress test, and 10 mM

glucose, 1 μM oligomycin, and 50 mM 2-deoxy-D-glucose (2-DG) for the glycolytic stress test.

The oxygen consumption rate (OCR) and extracellular acidification rate (ECAR) were measured

on a Seahorse XF analyzer. Non-glycolytic ECAR is calculated as the average ECAR values

after 2-DG treatment. Basal glycolysis is calculated as the average ECAR value prior to

oligomycin treatment minus non-glycolytic ECAR. Max glycolysis is calculated as the average

ECAR values after oligomycin and before FCCP treatment. Glycolytic reserve is calculated as

max glycolysis minus basal glycolysis. Basal respiration is calculated as the average OCR value

prior to oligomycin treatment minus non-mitochondrial OCR. Max respiration is calculated as

the average OCR value after FCCP and before rotenone/antimycin A addition.

Experimental schematics and graphical abstract.

All experimental schematics and graphical abstract created using BioRender.

https://biorender.com/

Quantification and statistical analysis

Results were expressed as mean ± SEM. All data was plotted in GraphPad Prism version 10.

Statistical significance was assessed by One-way ANOVA, Student's t test, and Two-way

ANOVA. Differences between experimental groups were considered significant if *$p<0.05$,

**$p<0.01$, ***$p<0.001$, ****$p<0.0001$.

References

1. Ngo, V.L., and Gewirtz, A.T. (2021). Microbiota as a potentially-modifiable factor influencing

COVID-19. Curr Opin Virol *49*, 21-26. 10.1016/j.coviro.2021.04.005.

2. Brodin, P. (2022). Immune-microbe interactions early in life: A determinant of health and

disease long term. Science *376*, 945-950. 10.1126/science.abk2189.

3. Beura, L.K., Hamilton, S.E., Bi, K., Schenkel, J.M., Odumade, O.A., Casey, K.A., Thompson, E.A.,

Fraser, K.A., Rosato, P.C., Filali-Mouhim, A., et al. (2016). Normalizing the environment

recapitulates adult human immune traits in laboratory mice. Nature *532*, 512-516.

10.1038/nature17655.

4. Rosshart, S.P., Vassallo, B.G., Angeletti, D., Hutchinson, D.S., Morgan, A.P., Takeda, K., Hickman,

H.D., McCulloch, J.A., Badger, J.H., Ajami, N.J., et al. (2017). Wild Mouse Gut Microbiota

Promotes Host Fitness and Improves Disease Resistance. Cell *171*, 1015-1028 e1013.

10.1016/j.cell.2017.09.016.

5. Phelan, J. (2016). Choosing a Health Standard for Your Model.

https://www.taconic.com/taconic-insights/quality/choosing-rodent-model-health-

standards.html.

6. Korth, M.J., Tchitchek, N., Benecke, A.G., and Katze, M.G. (2013). Systems approaches to

influenza-virus host interactions and the pathogenesis of highly virulent and pandemic viruses.

Semin Immunol *25*, 228-239. 10.1016/j.smim.2012.11.001.

7. Ivanov, II, and Littman, D.R. (2010). Segmented filamentous bacteria take the stage. Mucosal

Immunol *3*, 209-212. 10.1038/mi.2010.3.

8. Teng, F., Klinger, C.N., Felix, K.M., Bradley, C.P., Wu, E., Tran, N.L., Umesaki, Y., and Wu, H.J.

(2016). Gut Microbiota Drive Autoimmune Arthritis by Promoting Differentiation and Migration

of Peyer's Patch T Follicular Helper Cells. Immunity *44*, 875-888. 10.1016/j.immuni.2016.03.013.

9.   Shi, Z., Zou, J., Zhang, Z., Zhao, X., Noriega, J., Zhang, B., Zhao, C., Ingle, H., Bittinger, K., Mattei,

L.M., et al. (2019). Segmented Filamentous Bacteria Prevent and Cure Rotavirus Infection. Cell

179, 644-658 e613. 10.1016/j.cell.2019.09.028.

10.   Zemlin, M., Hummel, M., Anagnostopoulos, I., and Stein, H. (1998). Improved polymerase chain

reaction detection of clonally rearranged T-cell receptor beta chain genes. Diagn Mol Pathol *7*,

138-145. 10.1097/00019606-199806000-00003.

11.   Wymore Brand, M., Wannemuehler, M.J., Phillips, G.J., Proctor, A., Overstreet, A.M., Jergens,

A.E., Orcutt, R.P., and Fox, J.G. (2015). The Altered Schaedler Flora: Continued Applications of a

Defined Murine Microbial Community. ILAR J *56*, 169-178. 10.1093/ilar/ilv012.

12.   Newton, A.H., Cardani, A., and Braciale, T.J. (2016). The host immune response in respiratory

virus infection: balancing virus clearance and immunopathology. Semin Immunopathol *38*, 471-

482. 10.1007/s00281-016-0558-0.

13.   Purnama, C., Ng, S.L., Tetlak, P., Setiagani, Y.A., Kandasamy, M., Baalasubramanian, S.,

Karjalainen, K., and Ruedl, C. (2014). Transient ablation of alveolar macrophages leads to

massive pathology of influenza infection without affecting cellular adaptive immunity. Eur J

Immunol *44*, 2003-2012. 10.1002/eji.201344359.

14.   Lieber, C.M., Aggarwal, M., Yoon, J.J., Cox, R.M., Kang, H.J., Sourimant, J., Toots, M., Johnson,

S.K., Jones, C.A., Sticher, Z.M., et al. (2023). 4'-Fluorouridine mitigates lethal infection with

pandemic human and highly pathogenic avian influenza viruses. PLoS Pathog *19*, e1011342.

10.1371/journal.ppat.1011342.

15.   Sourimant, J., Lieber, C.M., Aggarwal, M., Cox, R.M., Wolf, J.D., Yoon, J.J., Toots, M., Ye, C.,

Sticher, Z., Kolykhalov, A.A., et al. (2022). 4'-Fluorouridine is an oral antiviral that blocks

respiratory syncytial virus and SARS-CoV-2 replication. Science *375*, 161-167.

10.1126/science.abj5508.

16.   Schneider, C., Nobs, S.P., Heer, A.K., Kurrer, M., Klinke, G., van Rooijen, N., Vogel, J., and Kopf,

M. (2014). Alveolar macrophages are essential for protection from respiratory failure and

associated morbidity following influenza virus infection. PLoS Pathog *10*, e1004053.

10.1371/journal.ppat.1004053.

17.   Dranoff, G., Crawford, A.D., Sadelain, M., Ream, B., Rashid, A., Bronson, R.T., Dickersin, G.R.,

Bachurski, C.J., Mark, E.L., Whitsett, J.A., and et al. (1994). Involvement of granulocyte-

macrophage colony-stimulating factor in pulmonary homeostasis. Science *264*, 713-716.

10.1126/science.8171324.

18.   Thepen, T., Van Rooijen, N., and Kraal, G. (1989). Alveolar macrophage elimination in vivo is

associated with an increase in pulmonary immune response in mice. J Exp Med *170*, 499-509.

10.1084/jem.170.2.499.

19.   Rauschmeier, R., Gustafsson, C., Reinhardt, A., N, A.G., Tortola, L., Cansever, D., Subramanian, S.,

Taneja, R., Rossner, M.J., Sieweke, M.H., et al. (2019). Bhlhe40 and Bhlhe41 transcription factors

regulate alveolar macrophage self-renewal and identity. EMBO J *38*, e101233.

10.15252/embj.2018101233.

20.   Behar, S.M., Martin, C.J., Booty, M.G., Nishimura, T., Zhao, X., Gan, H.X., Divangahi, M., and

Remold, H.G. (2011). Apoptosis is an innate defense function of macrophages against

Mycobacterium tuberculosis. Mucosal Immunol *4*, 279-287. 10.1038/mi.2011.3.

21.   Allden, S.J., Ogger, P.P., Ghai, P., McErlean, P., Hewitt, R., Toshner, R., Walker, S.A., Saunders, P.,

Kingston, S., Molyneaux, P.L., et al. (2019). The Transferrin Receptor CD71 Delineates

Functionally Distinct Airway Macrophage Subsets during Idiopathic Pulmonary Fibrosis. Am J

Respir Crit Care Med *200*, 209-219. 10.1164/rccm.201809-1775OC.

22.   Kuczma, M.P., Szurek, E.A., Cebula, A., Chassaing, B., Jung, Y.J., Kang, S.M., Fox, J.G., Stecher, B.,

889   and Ignatowicz, L. (2020). Commensal epitopes drive differentiation of colonic T(regs). Sci Adv *6*,

890   eaaz3186. 10.1126/sciadv.aaz3186.

891   23.   Ruan, T., Sun, J., Liu, W., Prinz, R.A., Peng, D., Liu, X., and Xu, X. (2020). H1N1 Influenza Virus

892   Cross-Activates Gli1 to Disrupt the Intercellular Junctions of Alveolar Epithelial Cells. Cell Rep *31*,

893   107801. 10.1016/j.celrep.2020.107801.

894   24.   Cardani, A., Boulton, A., Kim, T.S., and Braciale, T.J. (2017). Alveolar Macrophages Prevent Lethal

895   Influenza Pneumonia By Inhibiting Infection Of Type-1 Alveolar Epithelial Cells. PLoS Pathog *13*,

896   e1006140. 10.1371/journal.ppat.1006140.

897   25.   Misharin, A.V., Morales-Nebreda, L., Mutlu, G.M., Budinger, G.R., and Perlman, H. (2013). Flow

898   cytometric analysis of macrophages and dendritic cell subsets in the mouse lung. Am J Respir

899   Cell Mol Biol *49*, 503-510. 10.1165/rcmb.2013-0086MA.

900   26.   Berrebi, D., Bruscoli, S., Cohen, N., Foussat, A., Migliorati, G., Bouchet-Delbos, L., Maillot, M.C.,

901   Portier, A., Couderc, J., Galanaud, P., et al. (2003). Synthesis of glucocorticoid-induced leucine

902   zipper (GILZ) by macrophages: an anti-inflammatory and immunosuppressive mechanism shared

903   by glucocorticoids and IL-10. Blood *101*, 729-738. 10.1182/blood-2002-02-0538.

904   27.   Xia, M., Harb, H., Saffari, A., Sioutas, C., and Chatila, T.A. (2018). A Jagged 1-Notch 4 molecular

905   switch mediates airway inflammation induced by ultrafine particles. J Allergy Clin Immunol *142*,

906   1243-1256 e1217. 10.1016/j.jaci.2018.03.009.

907   28.   Varghese, P.M., Kishore, U., and Rajkumari, R. (2022). Human C1q Regulates Influenza A Virus

908   Infection and Inflammatory Response via Its Globular Domain. Int J Mol Sci *23*.

909   10.3390/ijms23063045.

910   29.   Klickstein, L.B., Barbashov, S.F., Liu, T., Jack, R.M., and Nicholson-Weller, A. (1997). Complement

911   receptor type 1 (CR1, CD35) is a receptor for C1q. Immunity *7*, 345-355. 10.1016/s1074-

912   7613(00)80356-8.

913   30.   Smythies, L.E., Sellers, M., Clements, R.H., Mosteller-Barnum, M., Meng, G., Benjamin, W.H.,

914   Orenstein, J.M., and Smith, P.D. (2005). Human intestinal macrophages display profound

915   inflammatory anergy despite avid phagocytic and bacteriocidal activity. J Clin Invest *115*, 66-75.

916   10.1172/JCI19229.

917   31.   Abt, M.C., Osborne, L.C., Monticelli, L.A., Doering, T.A., Alenghat, T., Sonnenberg, G.F., Paley,

918   M.A., Antenus, M., Williams, K.L., Erikson, J., et al. (2012). Commensal bacteria calibrate the

919   activation threshold of innate antiviral immunity. Immunity *37*, 158-170.

920   10.1016/j.immuni.2012.04.011.

921   32.   Ichinohe, T., Pang, I.K., Kumamoto, Y., Peaper, D.R., Ho, J.H., Murray, T.S., and Iwasaki, A. (2011).

922   Microbiota regulates immune defense against respiratory tract influenza A virus infection. Proc

923   Natl Acad Sci U S A *108*, 5354-5359. 10.1073/pnas.1019378108.

924   33.   Chen, H., Wang, L., Wang, X., Wang, X., Liu, H., and Yin, Y. (2020). Distribution and Strain

925   Diversity of Immunoregulating Segmented Filamentous Bacteria in Human Intestinal Lavage

926   Samples. Microb Ecol *79*, 1021-1033. 10.1007/s00248-019-01441-4.

927   34.   Jonsson, H., Hugerth, L.W., Sundh, J., Lundin, E., and Andersson, A.F. (2020). Genome sequence

928   of segmented filamentous bacteria present in the human intestine. Commun Biol *3*, 485.

929   10.1038/s42003-020-01214-7.

930   35.   Subramanian, S., Busch, C.J., Molawi, K., Geirsdottir, L., Maurizio, J., Vargas Aguilar, S., Belahbib,

931   H., Gimenez, G., Yuda, R.A.A., Burkon, M., et al. (2022). Long-term culture-expanded alveolar

932   macrophages restore their full epigenetic identity after transfer in vivo. Nat Immunol *23*, 458-

933   468. 10.1038/s41590-022-01146-w.

934   36.   Jeyanathan, M., Vaseghi-Shanjani, M., Afkhami, S., Grondin, J.A., Kang, A., D'Agostino, M.R., Yao,

935   Y., Jain, S., Zganiacz, A., Kroezen, Z., et al. (2022). Parenteral BCG vaccine induces lung-resident

936   memory macrophages and trained immunity via the gut-lung axis. Nat Immunol *23*, 1687-1702.

937   10.1038/s41590-022-01354-4.

938   37.   Wang, T., Zhang, J., Wang, Y., Li, Y., Wang, L., Yu, Y., and Yao, Y. (2023). Influenza-trained

939   mucosal-resident alveolar macrophages confer long-term antitumor immunity in the lungs. Nat

940   Immunol *24*, 423-438. 10.1038/s41590-023-01428-x.

941   38.   Verma, A.K., Bansal, S., Bauer, C., Muralidharan, A., and Sun, K. (2020). Influenza Infection

942   Induces Alveolar Macrophage Dysfunction and Thereby Enables Noninvasive Streptococcus

943   pneumoniae to Cause Deadly Pneumonia. J Immunol *205*, 1601-1607.

944   10.4049/jimmunol.2000094.

945   39.   Nogales, A., Avila-Perez, G., Rangel-Moreno, J., Chiem, K., DeDiego, M.L., and Martinez-Sobrido,

946   L. (2019). A Novel Fluorescent and Bioluminescent Bireporter Influenza A Virus To Evaluate Viral

947   Infections. J Virol *93*. 10.1128/JVI.00032-19.

948   40.   Hotard, A.L., Shaikh, F.Y., Lee, S., Yan, D., Teng, M.N., Plemper, R.K., Crowe, J.E., Jr., and Moore,

949   M.L. (2012). A stabilized respiratory syncytial virus reverse genetics system amenable to

950   recombination-mediated mutagenesis. Virology *434*, 129-136. 10.1016/j.virol.2012.09.022.

951   41.   Dinnon, K.H., 3rd, Leist, S.R., Schafer, A., Edwards, C.E., Martinez, D.R., Montgomery, S.A., West,

952   A., Yount, B.L., Jr., Hou, Y.J., Adams, L.E., et al. (2020). A mouse-adapted model of SARS-CoV-2 to

953   test COVID-19 countermeasures. Nature *586*, 560-566. 10.1038/s41586-020-2708-8.

954   42.   Yu, Y.R., O'Koren, E.G., Hotten, D.F., Kan, M.J., Kopin, D., Nelson, E.R., Que, L., and Gunn, M.D.

955   (2016). A Protocol for the Comprehensive Flow Cytometric Analysis of Immune Cells in Normal

956   and Inflamed Murine Non-Lymphoid Tissues. PLoS One *11*, e0150606.

957   10.1371/journal.pone.0150606.

Figure 1

Figure 2

Figure 3

Figure 4

A — *Batf3^{-/-}* cDC1 deficient

B — *Csf2^{-/-}*, AMs reduced

C — Clodronate depletion of AMs

i Naive / Clo

ii CA09 / SFB+CA09 / CA09+Clo / SFB+CA09+Clo

D Cleaved Caspase-3

E CD71

F Ki-67

G Naïve / Naïve+Clo / CA09 / SFB+CA09 / SFB+CA09+Clo

T1AECs / T2AECs / Conducting AECs

Figure 5

Figure 6

Figure 7

Figure S1 (related to F1)

Figure S2 (related to F3)

Figure S3 (related to F4)

Figure S4 (related to F4)

Figure S5 (related to F4)

Figure S6 (related to Fig 6)

Figure S7 (related to F6 & F7)