Generation and characterisation of an *Arabidopsis thaliana* *f3h/fls1/ans* triple mutant that accumulates eriodictyol derivatives

Hanna Marie Schilbert [1*] ORCID: 0000-0003-0474-7753

Mareike Busche [1*] ORCID: 0000-0002-2114-7613

Vania Sáez [2] ORCID: 0000-0002-6477-6671

Andrea Angeli [2] ORCID: 0009-0003-8882-5858

Bernd Weisshaar [1] ORCID: 0000-0002-7635-3473

Stefan Martens [2] ORCID: 0000-0001-9913-4882

Ralf Stracke [1#] ORCID: 0000-0002-9261-2279

[1] Bielefeld University, Faculty of Biology, Genetics and Genomics of Plants, 33615 Bielefeld, Germany

[2] Research and Innovation Centre, Fondazione Edmund Mach, San Michele all'Adige (TN), 38098, Italy

[*] These two authors contributed equally to this work.

[#] Corresponding author: ralf.stracke@uni-bielefeld.de

Abstract

Background: Flavonoids are plant specialised metabolites, which derive from phenylalanine and acetate metabolism. They possess a variety of beneficial characteristics for plants and humans. Several modification steps in the synthesis of tricyclic flavonoids cause for the amazing diversity of flavonoids in plants. The 2-oxoglutarate-dependent dioxygenases (2-ODDs) flavanone 3-hydroxylase (F3H, synonym FHT), flavonol synthase (FLS) and anthocyanidin synthase (ANS, synonym leucoanthocyanidin dioxigenase (LDOX)), catalyse oxidative modifications to the central C ring. They are highly similar and have been shown to catalyse, at least in part, each other's reactions. FLS and ANS have been identified as bifunctional enzymes in many species, including *Arabidopsis thaliana*, stressing the capability of plants to bypass missing or mutated reaction steps on the way to flavonoid production. However, little is known about such bypass reactions and the flavonoid composition of plants lacking all three central flavonoid 2-ODDs.

Results: To address this issue, we generated a *f3h/fls1/ans* mutant, as well as the corresponding double mutants and investigated the flavonoid composition of this mutant collection. The *f3h/fls1/ans* mutant was further characterised at the genomic level by analysis of a nanopore DNA sequencing generated genome sequence assembly and at the transcriptomic level by RNA-Seq analysis. The mutant collection established, including the novel double mutants *f3h/fls1* and *f3h/ans*, was used to validate and analyse the multifunctionalities of F3H, FLS1, and ANS *in planta*. Metabolite analyses revealed the accumulation of eriodictyol and additional glycosylated derivatives in mutants carrying the *f3h* mutant allele, resulting from the conversion of naringenin to eriodictyol by flavonoid 3'-hydroxylase (F3'H) activity.

Conclusions: We describe the *in planta* multifunctionality of the three central flavonoid 2-ODDs from *A. thaliana* and identify a bypass in the *f3h/fls1/ans* triple mutant that leads to the formation of eriodictyol derivatives. As (homo-)eriodictyols are known as bitter taste maskers, the annotated eriodictyol (derivatives) and in particular the observations made on

their *in planta* production, could provide valuable insights for creating of novel food supplements.

Keywords: 2-oxoglutarate-dependent dioxygenases, enzyme promiscuity, flavonoid biosynthesis, flavonoids, multifunctionality, triple 2-ODD mutant

Background

Flavonoids are a large group of specialised metabolites with an estimated number of 9,000 flavonoid derivatives and a wide distribution throughout the plant kingdom (1). Flavonoids share a basic structure with two aromatic C6 rings and a heterocyclic ring (2). Reorganisation or modification, such as oxidation or glycosylation, is mainly responsible for the wide variety of flavonoids. They can be classified into different subgroups including flavones, flavanones, flavonols, dihydroflavonols, flavandiols, anthocyanins, and proanthocyanidins (2). These subgroups exhibit manifold functions. Among others, anthocyanins promote pollination and seed dispersal by colouring flowers and fruits (3-5), flavonols play a role in auxin transport, plant fertility, and UV protection (6-8), and proanthocyanins serve as deterrent metabolites for defence against bird predators (9). Moreover, flavonoids are beneficial for human nutrition due to their antioxidative, antigenotoxic and anticarcinogenic properties (10, 11). These health-promoting effects have led to a strong interest in increasing the flavonoid content in plants (12, 13). However, as some flavonoids exhibit antinutritional properties, like a bitter-off taste (14), the engineering of flavonoid composition is of high relevance for the breeding of plants with desirable flavonoid profiles.

Flavonoid biosynthesis (Figure 1) is one of the best characterised pathways in plant specialised metabolism. Due to the visible phenotypes of respective mutants which often display a transparent testa resulting in yellow to pale-brown seed (15) and the large collections of T-DNA- and transposon-tagged lines for *Arabidopsis thaliana* (16-18), the genes involved in flavonoid biosynthesis are well studied. The flavonoid pathway is initiated by its key enzyme chalcone synthase (CHS), which condensates *p*-coumaroyl-CoA and three molecules of malonyl-CoA followed by cyclisation, leading to the formation of naringenin chalcone (19). A heterocyclic ring is introduced spontaneously or through the catalysis of chalcone isomerase (CHI), forming naringenin (20). Flavanone 3-hydroxylase (F3H or FHT) converts naringenin, a flavanone, to dihydrokaempferol, a dihydroflavonol, by hydroxylation at the C-3 position (21). In alternative branches, the flavonoid 3'-hydroxylase (F3'H) converts naringenin to eriodictyol and dihydrokaempferol to dihydroquercetin (22). Naringenin can also be converted to flavones

by flavone synthaseI or II (FNSI or II) (23, 24). However, flavones were thought to be absent in *A. thaliana* until two derivatives (7-2″,3″-diacetylglucoside and pentamethoxydihydroxy flavone) were identified in Columbia-0 (Col-0) leaves (25, 26). DOWNY MILDEW RESISTANT6 (*Ath*DMR6), a 2-oxoglutarate-dependent dioxygenase (2-ODD), has been proposed as the enzyme responsible for their biosynthesis (26). The enzyme flavonol synthase (FLS) uses dihydroflavonols to form flavonols by introducing a double bond between C-2 and C-3 (27). Alternatively, dihydroflavonol 4-reductase (DFR) can reduce dihydroflavonols to the corresponding leucoanthocyanidins, thus competing with FLS for substrates (28, 29). Anthocyanidin synthase (ANS, also known as leucoanthocyanidin dioxygenase (LDOX)) converts leucoanthocyanidins to anthocyanidins (30). Anthocyanidins can be modified by glycosyltransferases (GTs) to form anthocyanins, or they can be converted to proanthocyanidins by anthocyanidin reductase (ANR) (31, 32). Besides anthocyanidins, other flavonoids like flavonols or flavanones can be modified by GTs to form the respective glycosides (33-35).

F3H, FNSI, FLS and ANS belong to the 2-ODD enzyme family (36). 2-ODDs represent the second largest enzyme family in plants (37). They are ubiquitously spread across the plant kingdom but also microorganisms and are involved in a variety of primary and specialised metabolism, including the synthesis of specialised metabolites, hormones, signalling molecules, amino acid modification and DNA-repair (summarised in 38, 39). Plant 2-ODDs can be classified into three classes, DOXA, DOXB and DOXC, based on amino acid sequence similarity, with DOXC harbouring 2-ODDs involved in the synthesis of specialised metabolites, such as flavonoids, including F3H and FNSI (DOXC28) and FLS and ANS (DOXC47) (37).

106
107
Figure 1: Simplified representation of the flavonoid biosynthesis pathway in plants.
Details and abbreviations are given in the text. The 2-ODDs of the flavonoid biosynthesis F3H,
FNSI, FLS, and ANS are shown in bold.

Characteristic hallmarks of 2-ODDs are generally (but not exclusively) the requirement of

2-oxoglutarate as co-reductant and iron as co-factor (36). Elucidation of the crystallographic

structure of *Ath*ANS has identified the conserved amino acid residues involved in substrate-,

co-factor-, and co-substrate binding (30). In addition, residues important for proper folding and

substrate binding have been described (30).

As F3H, FLS and ANS all carry the conserved amino acid residues essential for 2-ODD activity, it has been proposed that they share a common catalytic mechanism, possibly allowing multifunctionality (40, 41). While FNSI and F3H often shows a narrow substrate spectrum (40, 42), FLS and ANS have been characterised as bi- or even multifunctional enzymes in several species (40, 43-46). From an evolutionary perspective, it seems feasible that the 2-ODD enzyme proposed to have evolved first, F3H, may have given rise to the other 2-ODDs involved in flavonoid biosynthesis (summarised in 47) and that *Ath*FLS1 and *Ath*ANS retain *in vitro* F3H (40, 48) and, in the case of *Ath*ANS, additional FLS side activity *in planta* (49). Due to the multifunctional nature of flavonoid 2-ODD enzymes, comprehensive biochemical and functional characterisation is required for unequivocal annotation, especially *in planta*. Only seven amino acid residues distinguish parsley F3H from FNSI, providing an impressive example that amino acid alignments are not sufficient for full functional characterisation (24). The multifunctionalities of flavonoid 2-ODDs can be seen in the "leaky" phenotypes of *f3h*, *fls1* and *ans* mutants of *A. thaliana* (21, 27, 49, 50), attributed to the partial compensation by other 2-ODDs. The *ans* mutant shows a significant reduction in anthocyanin and proanthocyanidin content with almost unchanged flavonol levels compared to the wildtype (49, 51, 52), while the *fls1* mutant shows a significant reduction in flavonol content and increased anthocyanin and proanthocyanidin levels (49, 50, 53). The *f3h* mutant has a reduced flavonoid content, including flavonols, anthocyanins, and proanthocyanidins, due to a generally reduced flux in the flavonoid pathway (21, 54).

To unambiguously annotate the *in planta* multifunctionalities of F3H, FLS and ANS (which does not necessarily reflect the *in vitro* function), multiple mutants are required, like it was done in the case of the *fls1*/*ans* double mutant, which demonstrated the *in planta* FLS side activity of ANS (40, 49).

In this study, we generated novel *A. thaliana f3h*/*fls1* and *f3h*/*ans* double mutants, as well as an *f3h*/*fls1*/*ans* triple mutant to examine the phenotypic changes resulting from the loss of activity of these flavonoid 2-ODDs. The *f3h*/*fls1*/*ans* mutant was characterised in detail, using genomic, transcriptomic and metabolomic methods. Comprehensive phenotyping of the

flavonoid composition of the established flavonoid 2-ODD mutant set (composed of single,

double, and triple mutants) provided new insights into the *in planta* multifunctionalities of

A. thaliana F3H, FLS1 and ANS and the resulting metabolic fluxes. We found a flux into a

family of eriodictyol derivatives, which is most prominent in the *f3h/fls1/ans* triple mutant. The

knowledge gained in this study will be valuable for targeted engineering of flavonoid

composition and content in plants, especially crops.

Results

Genomic characterisation of the *A. thaliana f3h/fls1/ans* mutant

To generate an *A. thaliana f3h/fls1/ans* mutant set, we used previously published insertional

alleles of the respective genes, showing the known loss-of-function phenotypes (*f3h* (*tt6-2*,

GK-292E08, Col-0 background), *fls1* (*fls1-2,* RIKEN_PST16145, Nö-0 background), *ans*

(*tds4-4*, SALK_073183, Col-0 background)). An *fls1/ans* double mutant was crossed with the

f3h single mutant. By screening the F2 progeny, homozygous *f3h/fls1/ans*, *f3h/fls1* and *f3h/ans*

mutants were identified by PCR-based genotyping. A *de novo* long-read assembly of the

f3h/fls1/ans mutant genome was generated to analyse the respective mutated loci at the

genomic level. The 160.57 Mbp assembly comprises 162 contigs with an N50 of 8.17 Mbp

(Supplementary Table S1). The assembly size slightly exceeds the size of the Col-0

Community-Consensus reference genome sequence assembly (133,336,906 bp) likely due to

the increased heterozygosity derived from the Nö-0 background of the *fls1* mutant. In total,

95.5 % complete BUSCOs were identified, indicating a high level of assembly completeness.

The assembly was used to validate the position of the T-DNA insertions at the *f3h* and *ans*

loci, and the position of the immobilised Ds transposon at the *fls1* locus (Figure 2). Two T-DNAs

comprising 11,393 bp sequence are inserted head-to-head into the 5'UTR of *f3h*, precisely

between 19,025,263 - 19,025,278 bp on chromosome three. The 5,417 bp Ds transposon is

inserted between 2,804,241 - 2,804,242 bp on chromosome five into the *fls1* locus in the first

exon. A large T-DNA array comprised of several T-DNAs and the corresponding binary vector

backbone (BVB) sequences was detected between 12,005,371 - 12,005,393 bp on

chromosome four, inserted into the front part of the second exon of *ans*. Due to the high amount

of repetitive sequences derived from this complex T-DNA array, the array is split across two

contigs in the assembly and therefore the exact size (spans >119 kbp) could not be

determined. A T-DNA left border (LB) was located at both ends of this insertion. Two additional

T-DNA insertion events have been identified (Supplementary Figure S1). However, these

insertions do not disrupt any additional annotated genes.

Figure 2: Genomic characterisation of the *f3h*/*fls1*/*ans* mutant. Schematic drawing of the
(A) *F3H* (*AT3G51240*), (B) *FLS1* (*AT5G08640*) and (C) *ANS* (*AT4G22880*) genes illustrating
the mutations in *f3h* (*tt6-2*), *fls1* (*fls1-2*) and *ans* (*tds4-4*), respectively. Black boxes mark the
CDSs and grey boxes indicate UTRs and grey thin lines indicate introns, bent arrows indicate
the transcription start. Dashed lines mark the respective T-DNA or transposon insertion.
Sequences belonging to the T-DNA or the transposon insertion are shown in a blue box in bold
and are flanked by the corresponding genomic sequences. LB, left border.

Transcriptome analysis of the *f3h*/*fls1*/*ans* mutant

To validate the insertion mutant alleles at the transcriptional level and to examine the

transcriptional changes that occur in the *f3h*/*fls1*/*ans* mutant, RNA-Seq data from six-days-old

seedlings were analysed. Col-0 and Nö-0 seedlings were used as controls. First, the

expression profiles of the respective knockout alleles were analysed in the *f3h*/*fls1*/*ans* mutant

(Supplementary Figure S2). For the *ans* and *fls1* mutant alleles, only marginal read coverage

was observed in the upstream regions of the T-DNA and transposon insertion positions,
respectively. These transcripts, which are greatly truncated compared to their normal lengths,
most likely result in non-functional proteins. Surprisingly, the *f3h* mutant allele showed a similar
expression profile as the wildtype allele, indicating that the *F3H* locus is still transcribed in the
f3h/*fls1*/*ans* mutant. However, mapping of the *f3h*/*fls1*/*ans* RNA-Seq reads to the *f3h*/*fls1*/*ans*
genome assembly revealed chimeric RNA-Seq reads containing the T-DNA LB sequence and
the sequence of the 5'UTR of *f3h*, emphasising that part of the T-DNA sequence is also
transcribed (Supplementary Figure S2). Several alternative translation start codons were
identified upstream of the original start site of *f3h* (Supplementary Figure S3), which could be
used for translation from the mutant *f3h* primary transcript and ultimately lead to a frameshift
and/or premature stop codon, resulting in a non-functional protein (refer to the discussion for
details).

In total, 1,643 and 1,203 differentially expressed genes (DEGs) were identified when
comparing *f3h*/*fls1*/*ans* with Col-0 and Nö-0, respectively (Figure 3, Supplementary File S1).
Of the DEGs derived from the comparison between *f3h*/*fls1*/*ans* and Col-0, 768 were
up-regulated and 875 were down-regulated, while 818 genes were up-regulated and 385
genes were down-regulated when *f3h*/*fls1*/*ans* was compared to Nö-0. To investigate the
transcriptional changes occurring in the *f3h*/*fls1*/*ans* mutant, 86 genes which are differentially
expressed between *f3h*/*fls1*/*ans* and Col-0 and between *f3h*/*fls1*/*ans* and Nö-0, but not
additionally between Col-0 and Nö-0, were further analysed (Figure 3). After a manual
screening (see Methods for details), 35 DEGs remained, which are listed in Table 1. Of these,
28 were up-regulated in the *f3h*/*fls1*/*ans* mutant. The *FLS1* gene was the only known gene
related to flavonoid biosynthesis in the list of DEGs.

Figure 3: Venn diagram summarising the number of differentially expressed genes between seedlings of *f3h*/*fls1*/*ans* and the wildtypes Col-0 and Nö-0.

Table 1: List of differentially expressed genes in *f3h*/*fls1*/*ans* vs. Col-0 and Nö-0. Listed are a subset of differentially expressed genes, that were identified as DEGs in both, the *f3h*/*fls1*/*ans* vs. Col-0 and *f3h*/*fls1*/*ans* vs. Nö-0 comparisons, but notably, they were not differentially expressed between Col-0 and Nö-0. AGI, Arabidopsis gene identifier; l2fc, $\log_2$ fold change; n.a., no annotation available.

AGI	l2fc vs. Col-0	l2fc vs. Nö-0	name / annotation
AT2G46765	7,25	5,74	hypothetical protein
AT5G08640	-5,54	-5,71	FLS1, flavonol synthase 1
AT2G20160	-4,71	-5,31	MEO, E3 ubiquitin ligase SCF complex subunit SKP1/ASK1 family protein
AT3G59884	3,35	1,84	MIR827A
AT2G14610	3,31	3,64	PR1 / CAPE9, pathogenesis-related protein 1
AT2G08820	2,3	1,37	n.a.
AT5G43570	2,22	1,38	PR-6 proteinase inhibitor family.
AT4G13493	2,21	1,64	MIR850A
AT4G06075	2,15	2,20	MIR5026 precursor
AT2G08825	2,11	1,40	n.a.
AT5G08855	2,11	1,96	n.a.
AT3G48560	2,08	2,14	AHAS / ALS, acetohydroxy acid synthase
AT3G46900	-1,98	-1,69	COPT2, Copper transporter 2, copper ion transmembrane transporter
AT1G52070	-1,93	-1,67	mannose-binding lectin superfamily protein
AT2G03850	1,93	1,07	LEA, late embryogenesis abundant protein family protein
AT2G11810	1,93	1,36	MGD3 / MGDC monogalactosyl diacylglycerol synthase 3
AT2G20723	1,84	1,79	snoRNA
AT2G20722	1,7	1,65	snoRNA

AT3G54730	1,62	3,16	transcription repressor
AT2G20670	1,52	1,62	DUF506, putative sugar phosphate exchanger
AT3G04210	1,36	1,59	TN13, disease resistance protein
AT2G44470	1,35	1,49	BGLU29, beta glucosidase 29
AT2G02850	1,34	1,16	ARPN, plantacyanin, copper ion binding
AT1G07135	1,29	1,23	glycine-rich protein
AT4G39675	1,28	1,32	hypothetical protein
AT1G60095	-1,27	-1,36	mannose-binding lectin superfamily protein
AT3G61930	1,26	1,17	hypothetical protein
AT2G07698	-1,23	-1,91	ATPase, F1 complex, alpha subunit protein
AT2G29330	1,23	1,54	TRI, tropinone reductase
AT1G36060	1,18	1,03	TG / WIND3, translucent green, Integrase-type DNA-binding superfamily protein
AT4G18980	1,18	1,58	AtS40-3
AT3G44260	1,17	1,13	CAF1A, Polynucleotidyl transferase, ribonuclease H-like superfamily protein
AT4G28420	1,14	1,62	tyrosine transaminase family protein
AT3G19430	-1,12	-1,90	LEA protein-like protein, late embryogenesis abundant protein-related
AT4G24570	1,01	1,10	DIC2, dicarboxylate carrier 2

Phenotypic characterisation of the 2-ODD mutant collection including the novel *f3h*/*fls1*, *f3h*/*ans*, and *f3h*/*fls1*/*ans* mutants

To phenotypically characterise the flavonoid composition of the *f3h*/*fls1*/*ans* mutant compared to the corresponding wildtypes, single and double mutants, the accumulation of flavanones, flavonols, anthocyanins and proanthocyanidins (PA) was analysed (Figure 4, Figure 5). First, the accumulation of flavonols was analysed in norflurazon-bleached seedlings, stained with diphenylboric acid 2-aminoethyl ester (DPBA) (Figure 4A). Both wildtypes, Col-0 and Nö-0, showed an orange colouration indicating the presence of quercetin glycosides. The *chs* mutant, included as a control, is deficient in flavonol glycosides and thus appeared blue due to sinapate derivatives. However, the *ans* mutant displayed an even more intense orange fluorescence than the wildtypes. While the *f3h* mutant showed a red fluorescence, the *fls1* mutant appeared green to light yellow. The *fls1*/*ans* mutant showed a pale yellow colour. The novel *f3h*/*fls1*, *f3h*/*ans*, and *f3h*/*fls1*/*ans* mutants appeared red like the *f3h* mutant.

HPTLC and targeted UPLC-MS/MS analysis (including MRM quantification) of the flavonoid composition in siliques revealed a wide range of flavonol glycosides in both wildtypes (Figure

4C, Supplementary Table S3). Consistent with DPBA staining, the *chs* mutant was deficient in

flavonoids. Accordingly, the level of *p*-coumaric acid was on average 186 µg per g dry weight

in *chs* siliques compared to up to 17 µg in all other lines, showing that this flavonoid precursor

accumulates in the *chs* mutant as the flavonoid biosynthesis pathway is blocked

(Supplementary Table S3). The *f3h*, *ans* and *f3h/ans* mutants showed a similar pattern of

flavonol glycosides as the wildtypes (Figure 4C). Moreover, the *ans* mutant accumulated the

highest amounts of flavonol glycosides with up to 3,549 µg quercetin-3-*O*-rhamnoside (Q-3R)

per g dry silique weight, whereas it was <273 µg in the other mutant lines and <2,656 µg in the

wildtypes (Supplementary Table S3). Accordingly, the *ans* knockout re-routes flavonoid

production to the flavonol biosynthesis branch while the other knockouts cause reduced

flavonol levels. The *fls1* mutant accumulated dihydroflavonols and small amounts of Q-3R,

whereas Q-3R levels decreased further to an average of 31.3 µg per g dry weight in *fls1/ans*

siliques while dihydroflavonol levels increased (Figure 4C and F, Supplementary Table S3).

The *fls1/ans* mutant accumulated the highest dihydroflavonol amounts compared to all lines

analysed, with an average dihydroquercetin level of 1,845 µg per g dry silique weight, showing

that dihydroflavonols accumulate when a further conversion to flavonols or anthocyanins is not

possible. Surprisingly, the *f3h/fls1* and *f3h/fls1/ans* mutants showed marginal and comparable

dihydroflavonols levels of up to 55 µg dihydrokaempferol per g dry weight, indicating the

presence of an (unknown) 2-ODD enzyme with F3H activity (Supplementary Table S3).

Interestingly, all mutant lines carrying the *f3h* allele accumulated a 'red (fluorescent)

metabolite', seen between Q-3R and K-3R-7R in HPTLC, indicating that a disruptive *f3h* allele

is essential for its production (Figure 4C). These results were in line with the red colouration of

in situ DPBA-stained seedlings carrying the *f3h* allele (Figure 4A). In contrast to flavonol

glycosides and dihydroflavonols, the accumulation of the 'red metabolite' was only visible late

after DPBA staining. The accumulation of the 'red metabolite' was further investigated in

DPBA-stained siliques under white and UV light (Supplementary Figure S4). The 'red

metabolite' visibly accumulated in immature seeds (Figure 4B) of all lines carrying the *f3h*

knockout allele and was also observed under white light. The *f3h/ans* and *f3h/fls1/ans* mutants

showed the strongest intensity and the 'red metabolite' appeared to accumulate during seed

maturation (Supplementary Figure S4). In addition, it was detected in young stamens of

mutants harbouring the *f3h* allele (Supplementary Figure S5). Interestingly, the eriodictyol

content of all analysed lines measured by UPLC-MS/MS were similar to the accumulation

pattern of the 'red metabolite' observed by HPTLC (Figure 4C and E). In detail, the average

level of eriodictyol was <1.4 µg per g dry silique weight in lines with a functional *F3H* allele,

whereas the average eriodictyol content of *f3h* knockout lines increased to 170 µg in *f3h*,

300 µg in *f3h/fls1,* 457 µg in *f3h/ans* and 442 µg in *f3h/fls1/ans*, indicating that eriodictyol

accumulates as a consequence of the *f3h* knockout (Figure 4E). To further characterise the

'red metabolite', HPTLC was performed with available eriodictyol glycoside standards together

with methanolic extracts of Col-0 and *f3h/fls1/ans* siliques (Figure 4D). The 'red metabolite'

had the same retention factor (R$_F$) as eriodictyol-7-*O*-glucoside, while the R$_F$ values of the

eriodictyol disaccharides and the eriodictyol aglycone differed from that of the 'red metabolite'.

Taken together, these results suggest that the metabolite underlying the red colouration of

mutants carrying the *f3h* allele is likely to be an eriodictyol hexoside, e.g.,

eriodictyol-7-*O*-glucoside.

Further chromatographic analysis with ion mobility separation in a high resolution mass

spectrometer (UPLC-TWIMS-QToF) on the *f3h/fls1/ans* flavonoid profile in siliques confirmed

the presence of eriodictyol-7-*O*-glucoside compared to a standard, the results show a mass

error below 3 ppm and an average of delta collision cross section (CCS) delta of 0.06 % in

ESI+ and 2.23 % in ESI- mode (Supplementary Table S4). The average content of

eriodictyol-7-*O*-glucoside determined by UPLC-MS/MS (high resolution mass spectrometry) in

the *f3h/fls1/ans* triple mutant is 10.38 µg per g dry weight (Supplementary Table S5).

Interestingly, the profile in *f3h/fls1/ans* is more complex (Supplementary Figures S6-S8). The

annotation of the eriodictyol derivates profile are based on HDMS[E] data, fragmentation

patterns by UPLC-MS/MS and isotopic distribution. The UPLC-TWIMS-QToF HDMS[E] profile

obtained in ESI+ and ESI- mode and subsequently UPLC-MS/MS analysis revealed the

presence of another eriodictyol hexoside, four di-hexosides, two tri-hexosides and two

rutinosides or neohesperidosides, respectively (Supplementary Tables S4 and S6). In ESI-

mode the main loss by UPLC-MS/MS is 162 Da (Supplementary Figure S8), corresponding to

the loss of a hexoside, the fragmentation has the presence of m/z 287, m/z 151 and m/z135

(Supplementary Figure S8), indicating a core with molecular formula $C_{15}H_{12}O_6$. Eriodictyol and

dihydrokaempferol have the same precursor, naringenin (Figure 1), and the same molecular

formula. Therefore, a comparison of fragmentation patterns, under the same methodology, of

eriodictyol and dihydrokaempferol standards by UPLC-MS/MS and UPLC-HDMS[E] low and

high energy is required to improve the metabolite annotation. As is shown in Supplementary

Figures S9-S11, the absence of m/z 177 in all compounds of interest, including the

eriodictyol-7-*O*-glucoside, indicates an eriodictyol core. Two compounds, tentatively annotated

as eriodictyol rutinoside or neohesperidoside have two losses of 162 Da and 308 Da in

ESI- and 308 Da in ESI+ mode (Supplementary Figure S8, Supplementary Tables S4 and S6).

The presence of eriodictyol-7-*O*-rutinoside is excluded by the mismatch in retention time and

CCS values in the triple mutant by HDMS[e] (Supplementary Table S4) and is in agreement with

the results shown in Figure 4D. Moreover, the dihydroquercetin profile of the *f3h*/*fls1*/*ans* triple

mutant compared to Col-0 is interesting, as it contains three different hexosides

(Supplementary Table S4). These results shed light on the accumulation of the eriodictyol and

dihydroquercetin aglycons and their subsequent modifications in the *f3h*/*fls1*/*ans* mutant.

Figure 4: Accumulation of an eriodictyol hexoside in *A. thaliana* 2-ODD mutants carrying the *f3h* allele. (A) Seedlings stained *in situ* with diphenylboric acid 2-aminoethyl ester (DPBA) and visualised under UV light. Quercetin derivatives appear orange, kaempferol derivatives green, and blue corresponds to sinapate acid derivatives. (B) *In situ* DPBA-stained silique of the *f3h/fls1/ans* mutant visualised under UV light. (C) Accumulation of flavonol- and dihydroflavonol glycosides in siliques analysed by HPTLC and visualised under UV light after

treatment with DPBA. Kaempferol derivatives are green, quercetin derivatives are orange and sinapate acid derivatives are blue. G, glucose; K, kaempferol; Q, quercetin; R, rhamnose; DHK-D, dihydrokaempferol derivative; and DHQ-D, dihydroquercetin derivative. (D) Eriodictyol standards analysed by HPTLC and visualised under UV light after treatment with DPBA. E, eriodictyol. (E) Eriodictyol and (F) quercetin-3-*O*-rhamnoside contents in siliques analysed by UPLC-MS/MS MRM. Error bars indicate the standard deviation of four independent measurements.

Besides flavanones and flavonols, other downstream (end-)products of the flavonoid biosynthesis were analysed including anthocyanins and PAs. The accumulation of anthocyanins was analysed in sucrose-induced, norflurazon-bleached seedlings (Figure 5A). While both wildtypes, *f3h*, *fls1*, and *f3h/fls1* mutants show anthocyanin accumulation, *chs*, *ans*, *f3h/ans*, *fls1/ans*, and *f3h/fls1/ans* mutants do not. These results are consistent with the analysis in sucrose-induced, native seedlings and were validated by photometric quantification of anthocyanins in seedlings (Figure 5A and B).

Seed pigmentation is influenced by the levels of PA in the seed coat, with a darker seed colour corresponding to higher PAs levels. Analysis of the native seed colour of the wildtypes revealed characteristic dark brown colour due to the presence of large amounts of PAs, while the *chs* mutant showed a pale yellow seed indicative of the well-known PA deficiency and transparent testa phenotype (Figure 5A). Most of the *2-ODD* mutants showed an intermediate seed colour phenotype. However, the *fls1* mutant showed a similar brown seed colour as the wildtypes. In accordance, UPLC-MS/MS analysis revealed that epicatechin levels of Col-0, Nö-0, and *fls1* averaged between 799 µg and 1,108 µg per g dry silique weight, while all other lines showed average levels <7 µg (Supplementary Table S3). Similar observations were made for procyanidins (Figure 5D, Supplementary Table S3). Thus, PAs and their precursor epicatechin are only produced in considerable amounts when *CHS, F3H*, and *ANS* are intact.

Figure 5: Anthocyanin and proanthocyanidin accumulation in *A. thaliana* 2-ODD mutants. (A) First column: norflurazon (NFZ)-bleached seedlings grown on 4% sucrose for anthocyanin induction. Second column: anthocyanin accumulation in seedlings grown on 0.5 MS plates with 4% sucrose. Third column: native seed colour. (B) Sucrose-induced anthocyanin accumulation in 6-days-old *A. thaliana* seedlings. Three independent biological replicates were measured for each sample. Error bars indicate standard deviation of the mean anthocyanin levels. (C) Procyanidin B2 + B4 content in siliques analysed by UPLC-MS/MS. Error bars indicate the standard deviation of four independent measurements.

Discussion

The three major flavonoid 2-ODD enzymes, F3H, FLS and ANS, are closely related and therefore often have overlapping catalytic activities. We have generated an *A. thaliana* *f3h/fls1/ans* mutant and analysed genomic, transcriptomic and phenotypic changes in order to increase the knowledge about flavonoid biosynthesis related 2-ODDs in *A. thaliana.*

Genome sequence of the *f3h/fls1/ans* triple mutant

To characterise the *f3h/fls1/ans* triple mutant at the genomic level, a *de novo* Oxford Nanopore Technology (ONT) genome assembly was generated. In-depth analysis of the respective *2-ODD* mutant loci revealed the expected insertion positions per mutant allele (49, 51, 55). To our knowledge, these mutant alleles have never been characterised using long-read technology. Thus, in this study, the insertion of repeated T-DNAs and their BVB sequences at the *ans* (*tds4-4*) locus were identified. Moreover, the ONT assembly revealed that two T-DNAs inserted head-to-head in the 5'UTR of *f3h*. Such complex T-DNA insertion events, involving arrays of multiple T-DNAs as well as BVB sequences, have been reported before to be more the rule than the exception for *A. thaliana* T-DNA insertion mutants (56-59).

Transcriptome analysis of the *f3h/fls1/ans* triple mutant

RNA-Seq analysis in seedlings revealed that *FLS1* is the only differentially expressed 2-ODD between the *f3h/fls1/ans* mutant and the wildtypes and not additionally differentially expressed between the wildtypes Col-0 and Nö-0. *ANS* is downregulated in the triple mutant compared to the two wildtypes, but additionally also downregulated in Nö-0 compared to Col-0. *F3H* was not identified as differentially expressed gene (DEGs) due to remaining RNA-Seq read coverage at the *f3h* loci downstream of the T-DNA insertion site (Supplementary Figure S2). For *ANS*, which comprises two exons, read coverage of the most of the first exon was observed upstream of the complex T-DNA insertion array. The resulting transcript is severely truncated and, as previously confirmed by the phenotype of the homozygous mutant (51, 52), results in a non-functional protein.

Interestingly, the expression level of the *f3h* locus was similar to that of the wildtype alleles and

covered the entire *F3H* exons. Comparison of the RNA-Seq data to the *f3h/fls1/ans* genome

assembly identified several chimeric RNA-Seq reads spanning part of the T-DNA and the

5'UTR of *f3h*, emphasising that part of the T-DNA::genome junction is transcribed.

Transcription is likely driven by the 2'promoter of the T-DNA, located 723 bp upstream of the

5'UTR of the *f3h* locus (Supplementary Figure S3). A similar observation was made in a

phytochrome C mutant by *Monte et al.* (60). They analysed a T-DNA insertion line that showed

PHYC transcript levels similar to the wildtype. However, protein analysis revealed, that the

plants did not accumulate the corresponding protein. Similar limitations are conceivable in our

study. The 11,393 bp T-DNA insertion could either lead to an extended and, in consequence,

non-functional protein or one of the start codons located on the T-DNA could be used as an

alternative start codon, ultimately resulting in a frameshift and/or premature stop codon

(Supplementary Figure S3). Taken together, the omics data and the very distinct phenotype of

the *f3h/fls1/ans* mutant strongly suggest a non-functional *f3h* allele, in line with previous results

(17, 55). Furthermore, genomic and transcriptomic data of the *ans* and *fls1* loci showed that

these loci are *null*-alleles.

Apart from the *FLS1* gene, no other genes encoding enzymes, regulators or transporters of

flavonoid biosynthesis were identified as DEGs between the *f3h/fls1/ans* mutant and the

wildtypes. Nevertheless, several identified DEGs are noteworthy. For example, *miR827A* is

upregulated in *f3h/fls1/ans*, compared to the wildtypes. *miR827* expression is known to be

sucrose-induced and it targets ethylene response factors (ERFs) to regulate the expression of

SUCROSE SYNTHASE which has been proposed as a regulator of anthocyanin biosynthetic

genes (61). Accordingly, increased *miR827A* expression could be a means of the *f3h/fls1/ans*

mutant to counteract anthocyanin deficiency. Of course, this requires an unknown feedback

regulation circuit. *TROPINONE REDUCTASE* (*TRI*) transcript abundance is increased in

f3h/fls1/ans seedlings. Moreover, based on publicly available RNA-Seq datasets deposited in

of the European Bioinformatics Institute (EBI) Expression Atlas, *TRI* is expressed in Col-0

siliques (62, 63). TRI has previously been reported to reduce the flavanone backbone (64).

Since the flavonol- and anthocyanin branches of flavonoid biosynthesis are blocked in the

f3h/fls1/ans mutant, flavanones like eriodictyol accumulate in *f3h/fls1/ans* and could be

reduced by TRI activity. *BETA GLUCOSIDASE29* (*BGLU29*) is upregulated in *f3h/fls1/ans*

compared to the wildtypes. BGLU29 belongs to the family of beta-glucosidase enzymes (65).

Although some BGLUs have been shown to be functional glucosyltransferases involved in the

glycosylation of flavonoids (BGLU10 (66) and BGLU6 (67)), it is very unlikely that BGLU29 has

glycosyltransferase activity, as it is most closely related to BGLU28 - BGLU32, all of which

having glycoside hydrolase activity (68). BGLU29 may be involved in the hydrolysis of flavonoid

glycosides. In addition, the expression of *ACETOHYDROXY ACID SYNTHASE* (*AHAS*), which

is involved in the biosynthesis of valine, leucine and isoleucine from pyruvate (69), is increased

in the *f3h/fls1/ans* mutant. Since flavonoids are derived from phenylalanine, increased *AHAS*

expression could hint to an accumulation of phenylalanine in *f3h/fls1/ans* and an effort by the

plant to produce other amino acids instead.

Phenotype

***In planta* multifunctionality of flavonoid 2-ODDs**

Previous studies revealed a "leaky" phenotype of the *2-ODD* single mutants and the *fls1/ans*

double mutant, which was attributed to partial compensation of other 2-ODD dioxygenases,

e.g., the bi- and multifunctionality of FLS and ANS, respectively (21, 27, 49). Owens *et al.* (27)

showed, that FLS1 not only accepts dihydroflavonols as substrate, but can also convert

naringenin to dihydrokaempferol *in planta*, thus exhibiting F3H side activity. ANS has been

shown to produce dihydroquercetin and quercetin from naringenin *in vitro*, indicating its F3H

and FLS side activity (40, 70, 71). Furthermore, the accumulation of flavonol glycosides in the

fls1 mutant and further flavonol reduction in the *fls1/ans* mutant showed that ANS also has an

FLS activity *in planta* (49). This knowledge is consistent with our analyses and the novel

f3h/ans, *f3h/fls1* and *f3h/fls1/ans* mutants allowed an in-depth analysis of the *in planta*

multifunctionality of F3H, FLS, and ANS. The flavonol glycoside patterns of the *f3h*, *ans* and

the novel *f3h/ans* mutants were like the ones of the wildtypes confirming the significant *in*

planta F3H activity of FLS1.

The *fls1/ans* mutant still accumulates flavonol glycosides, suggesting that F3H has additional

in planta FLS activity. Alternatively, FLS3 may have residual *in planta* FLS activity, as FLS *in*

vivo activity of FLS3 has been shown in yeast (72). As all mutant lines with a non-functional

ANS contain no anthocyanins, F3H and FLS do not seem to have any ANS activity. Finally,

the marginal and comparable amounts of dihydroflavonol of up to 55 µg dihydrokaempferol per

g dry silique weight identified in the *f3h/fls1* and *f3h/fls1/ans* mutants indicate that ANS does

not appear to have significant F3H activity *in planta.* However, the identification of residual

dihydroflavonols in the *f3h/fls1/ans* mutant was rather surprising since the two major flavonoid

2-ODD responsible for dihydroflavonol accumulation, F3H and FLS1, were knocked-out in this

mutant.

Although the biological significance of these trace amounts of dihydroflavonols and the

enzyme(s) leading to their production is questionable, it raises the question of which enzyme(s)

could be involved in their biosynthesis. The dihydroflavonols could be produced by an

(unknown) 2-ODD with F3H activity or F3H side activities of one of the FLS paralogs. While

AthFLS2, *AthFLS4*, *AthFLS5* and *AthFLS6* were identified as pseudogenes or encoding

non-functional proteins, minor FLS activity of *Ath*FLS3 was reported under prolonged assay

conditions in yeast (27, 72). Although FLS3 showed no F3H or FLS activity in *Escherichia coli*,

in planta F3H activity was identified for FLS3 homologs from banana and rapeseed, the latter

sharing 67 % amino acid sequence identity with *Ath*FLS3 (42, 46). To investigate a potential

in planta F3H and FLS activity of *Ath*FLS3, an *fls3* knockout allele could be introduced in the

f3h/fls1/ans mutant to generate a *fls3/f3h/fls1/ans* mutant. A reduction of dihydroflavonols and

flavonols in the *fls3/f3h/fls1/ans* mutant compared to the *f3h/fls1/ans* mutant would provide

evidence for an F3H and FLS activity of FLS3 *in planta*.

Alternatively, the production of residual dihydroflavonols in the *f3h/fls1/ans* mutant may involve

FNSI, another flavonoid 2-ODD closely related to F3H. Bifunctional enzymes with FNSI and

F3H activity have been identified in gymnosperms and mosses (73). FNSI enzymes were

thought to be restricted to the Apiaceae family (74) and thus absent from *A. thaliana,* until the

identification of a DOWNY MILDEW RESISTANT6 (DMR6/FNSI) homolog with FNSI activity,

alongside reports of functional FNSIs from liverworts, *Oryza sativa*, and *Zea mays* (26, 73, 75,

76). However, *DMR6*/*FNSI* is primarily expressed in leaves and not in siliques (26), where the

residual amounts of dihydroflavonols were detected. Moreover, the endogenous activity of

DMR6/FNSI is not sufficient to compensate the dihydroflavonol deficient phenotype of the

f3h/*fls1* or *f3h*/*fls1*/*ans* mutants, indicating that DMR6/FNSI does not have substantial F3H

activity *in planta*. Taken together, these findings suggest that if DMR6/FNSI harbours F3H side

activity, it is likely to play a minor role, if any, in the production of the residual dihydroflavonols

detected in *f3h*/*fls1*/*ans* siliques.

In summary, the established 2-ODD mutant collection could be used in the future to analyse

not only potential multifunctionality of DMR6/FNSI, but rather 2-ODDs in general. This is of

particular importance for the FNSI family, as it is no longer confined to the Apiaceae family,

and knowledge on FNSI multifunctionality is rare, especially in angiosperms (73, 77). As

mentioned above, flavonoid 2-ODDs show broad substrate specificities *in vitro,* which do not

necessarily correspond to product accumulation *in vivo* (21). This emphasises the importance

of the established mutant collection for the *in planta* characterisation of flavonoid 2-ODD

multifunctionality, especially for studies characterising their functionalities solely *in vitro* (78,

79). For example, *in planta* complementation experiments using the novel mutant collection

could validate the FLS side activity found *in vitro* and the absence of F3H side activity of

Vitis vinifera's ANS (79).

Flavonoid accumulation in the absence of the three major flavonoid 2-ODDs

The mutant collection was harnessed to investigate changes in the metabolic flux associated

with the knockout of the three major flavonoid 2-ODDs F3H, FLS1, and ANS. Interestingly, all

mutant lines carrying the *f3h* allele accumulate a metabolite in seedlings, immature seeds,

young stamens and siliques, which appears 'red' after DPBA-staining in visible light and under

UV light. This 'red (fluorescent) metabolite' is absent in the wildtypes and is most clearly seen

in an HPTLC of methanolic extracts from siliques (Figure 4B). The characteristic red

fluorescence of the *f3h* seedlings after DPBA staining has previously been observed and

suggested to be derived from an orthodihydroxy- or orthotrihydroxy flavonoid (80). In addition,

3'hydroxylation of the B ring was suggested to be required for the red fluorescence upon

staining, as DPBA-stained *f3h/f3'h* and *chs/f3h* mutants showed a light blue to pale

fluorescence (80). Owens *et al.* (21) suggested that F3'H might convert the accumulating

naringenin to eriodictyol which could be further reduced to a 3-desoxy flavonoid by the FNR

activity of native DFR (21). Wollenweber (81) postulated that the metabolite underlying the red

spot on DPBA-stained TLC plates is an eriodictyol derivative. We show that high levels of

eriodictyol accumulate in siliques of lines carrying an *f3h* knockout allele. This is in accordance

with the high expression of *F3'H* in siliques compared to flowers, leaves, stems, and roots (22).

In addition, we also observed that *F3'H* is expressed in *f3h/fls1/ans* seedlings (Supplementary

Figure S12), which showed the characteristic red colouration after *in situ* DPBA-staining. While

HPTLC-based flavonoid glycoside analysis showed that the 'red metabolite' is most likely an

eriodictyol hexoside, the results by UPLC-MS/MS and UPLC-TWIMS-QToF analyses revealed

a profile with various metabolites annotated as eriodictyol hexosides, including

eriodictyol-7-*O*-glucoside, and additional possible eriodictyols with neohesperidoside or

rutinoside derivatisation.

We therefore postulate that F3'H converts naringenin to eriodictyol, which may then be further

modified by glycosylation via glycosyltransferases. For example, the

flavonoid-7-*O*-glycosyltransferase UGT73B1 has been shown to use eriodictyol as a preferred

substrate for the catalysis to eriodictyol-7-*O*-glycoside (35). Considering the previously

presented DEGs, the tropinone reductase (TRI) may be involved in eriodictyol reduction, as a

previous study showed that TRI can reduce flavanones *in vitro* (82). In the future, knockout

alleles of the identified candidate genes could be introduced in the *f3h/fls1/ans* mutant to

analyse their role in the eriodictyol hexoside biosynthesis and degradation.

Conclusions

We present a detailed characterisation of a novel *A. thaliana f3h/fls1/ans* mutant and the corresponding single and double mutants. We have used genomic, transcriptomic, and metabolite data to extend the knowledge about flavonoid 2-ODDs in *A. thaliana.* The set of *f3h, fls1* and *ans* single, double and triple mutants will enable further work on flavonoid 2-ODDs and their multifunctionalities in *A. thaliana*, as these are often limited to *in vitro* studies or enzyme assays in *E. coli* or yeast (45, 78, 83). The *f3h/fls1/ans* mutant will also facilitate the characterisation of multifunctional 2-ODDs from other species *in planta*. Finally, we have shown that eriodictyol and several eriodictyol derivatives accumulate in the absence of a functional F3H *in planta* with eriodictyol-7-*O*-glucoside being one of the predominant eriodictyol derivatives. (Homo-)eriodictyols are bitter-masking flavanones with no additional strong taste or flavour and are thus promising flavour modifiers for food applications and pharmaceuticals (84).

Methods

Plant material

Columbia-0 (Col-0, NASC ID N1092) and Nössen-0 (Nö-0, NASC ID N3081) accessions were used as wildtype controls. A *chs* mutant (*tt4-11*, SALK_020583, Col-0 background, T-DNA from pROK2 plasmid (18, 85)) was used as a flavonoid deficient control. The *A. thaliana* mutants *f3h* (*tt6-2*, GK-292E08, Col-0 background, T-DNA from pAC161 plasmid (17)), *fls1* (*fls1-2,* RIKEN_PST16145, Nö-0 background, Ds transposon (49,16)), *ans* (*tds4-4*, SALK_073183, Col-0 background (51, 52)) and an *fls1/ans* double mutant (49) were used for crosses and as controls.

Crossing and genotyping of mutants

To generate the novel *f3h/fls1*, *f3h/ans*, and *f3h/fls1/ans* mutants, the *fls1/ans* double mutant was crossed with the *f3h* mutant. Individuals of the F2 progeny were analysed by PCR genotyping using GoTaq2 DNA polymerase (Promega) according to standard protocols.

Sequences of primers used for genotyping are provided in Supplementary Table S2. Plants

homozygous for the respective mutant alleles were used for further analyses.

DNA extraction, library construction, and sequencing

ONT was used to generate long-reads of the *f3h/fls1/ans* mutant. Genomic DNA was isolated

from rosette leaves using the NucleoBond HMW DNA Kit (Macherey-Nagel, Düren, Germany)

according to the manufacturer's instructions. DNA quality was assessed via PicoGreen

(Quant-iT) measurement on a FLUOstar OPTIMA plate reader (BMG labtech), and pulsed-field

gel electrophoresis. The Short Read Eliminator (SRE) Kit (Circulomics, Baltimore, MD, USA)

was used to enrich long DNA fragments and results were validated by PicoGreen

measurements. 1.2 µg of high molecular weight DNA was used for library preparation

according to the SQK-LSK109 protocol (Oxford Nanopore Technologies (ONT), Oxford, UK).

Sequencing was performed on a R9.4.1 flow cell on a GridION. The flow cell was treated with

nuclease flush to load another library twice. Basecalling was performed using Guppy v.4.2.3

(https://community.nanoporetech.com).

***De novo* ONT genome assembly and read mapping**

A *de novo* ONT genome assembly of the *f3h/fls1/ans* mutant was constructed with Canu v2.1.1

(86). No polishing of the assembly was performed, as the assembly was only used to analyse

the respective mutant loci. Contigs <100 kb were removed, followed by the calculation of

assembly statistics as previously described (87). Assembly quality was assessed via BUSCO

v3.0.2 using the embryophyta_odb9 lineage dataset (88). For manual curation of loci of interest

reads were mapped to the *A. thaliana* TAIR10 reference genome sequence (89) using

minimap2 (90). Minimap2 was run in ONT mode (-ax map-ont) and no secondary alignments

were allowed (--secondary=no). BAM files were sorted and indexed using samtools (91).

Manual curation of regions of interest was performed by using Integrative Genomics Viewer

(92) and BLAST (93). The assembly data was screened for the presence of additional T-DNAs

using loreta (57).

RNA-Seq analysis: Ribonucleic acid extraction, library construction, and sequencing

RNA was isolated from 6-days-old seedlings using the NucleoSpin® RNA Plant Kit (Macherey-Nagel, Düren, Germany) according to the manufacturer's instructions. Four biological replicates were used for each genotype (Col-0, Nö-0, and *f3h/fls1/ans*). The RNA quality was validated using a 2100 Bioanalyzer system (Agilent, Santa Clara, CA, USA). Sequencing libraries were constructed from 1 µg of total RNA, following the TrueSeq v2 protocol. Single end sequencing of 84 nt was performed on an Illumina NextSeq500 at the Sequencing Core Facility of the Center for Biotechnology (CeBiTec) at Bielefeld University.

RNA-Seq mapping and count table generation

After assessing read quality using FastQC (94), reads were mapped to the *A. thaliana* Col-0 TAIR10 reference genome sequence (95) using STAR v.2.7.1a (96). STAR was run in Basic mode allowing a maximum of 5 % mismatches per read length and using a minimum of 90 % matches per read length. Mapping statistics were calculated based on STAR.log files using a customised python script (available online via https://github.com/hschilbert/3xODD/blob/main/parse_STAR_log_file_create_mapping_statistic.py). We used featureCounts v2.0.1 (97) with the -O --fraction parameters to generate count tables based on the *A. thaliana* araport11 gff3 file (98) in order to assign a fractional count of 1/y from a reads, where y is the total number of meta-features/features overlapping with the reads.

Analysis of differentially expressed genes

To identify differentially expressed genes (DEGs) between Col-0, Nö-0 and the *f3h/fls1/ans* mutant in young seedlings, DESeq2 (99) was used to identify DEGs with a $\log_2$ fold change >1 and adjusted p-value <0.05 were selected for downstream analysis. Selected DEGs (86 DEGs derived from the comparison of *f3h/fls1/ans* mutant to both wildtypes) were manually curated by in-depth analysis of DEG's RNA-Seq read coverage against the TAIR10 reference genome sequence to exclude DEGs arising from mapping artefacts, e.g., genes with partial read

coverage. The Venn diagram was generated using
https://bioinformatics.psb.ugent.be/webtools/Venn/.

Flavonoid staining in *A. thaliana*

Visualisation of DPBA/Triton X-100-stained flavonoids in norflurazon-bleached *A. thaliana*
seedlings was performed according to Stracke *et al.* (100) using epifluorescence microscopy.
Visualisation of flavonoids in ethanol-bleached siliques of adult *A. thaliana* plants was
performed according to Stracke *et al.* (101).

High performance thin layer chromatography (HPTLC) of flavonols and
dihydroflavonols

Plants for flavonols extraction were grown in a greenhouse with 16 h of light illumination per
day at 22 °C. Extraction and analysis of dihydroflavonol and flavonol glycoside compounds
were performed as described in Stracke *et al.* (49). Siliques were homogenised in 80 %
methanol, incubated at 70 °C for 15 min and centrifuged at 14,000 x g for 10 min. After
vacuum-drying of the resulting supernatants, the dried pellets were redissolved in 80 %
methanol and analysed by HPTLC on a Silica Gel 60 plate (Merck, Darmstadt, Germany). The
mobile phase used was ethyl acetate, formic acid, acetic acid and water (100:26:12:12).
Dihydroflavonols and flavonol glycosides were detected by DPBA staining and UV light.

(Poly)phenol extraction and targeted Ultra Performance Liquid Chromatography
– Multiple Reaction Monitoring Mass Spectrometry (UPLC-MS/MS MRM) analysis

Samples of plant material were taken directly in the laboratory, lyophilized, and stored at room
temperature until used for further analysis. 100 mg of ground tissue were mixed with 1 mL of
80 % methanol for extraction of (poly)phenolic compounds. Samples and solvent were mixed
on an orbital shaker for 1 h at room temperature, sonicated for 30 min and the extraction
continued overnight at 4 °C in the dark. Each extract was filtered through a MILLEX 13-0.22 µm
PTFE membrane into glass vials and analysed as follows.

Sample extracts were injected immediately after filtration. Targeted UPLC-MS/MS with MRM
quantification analysis was performed on a Waters Acquity UPLC system (Milford, MA, USA)

consisting of a binary pump, an online vacuum degasser, an autosampler and a column compartment. Separation of the phenolic compounds was conducted on a Waters Acquity HSS T3 column 1.8 µm, 100 mm × 2.1 mm, kept at 40 °C. The analysis of phenolic compounds was performed according to the method described by Vrhovsek *et al.* (102) using water with 0.1 % formic acid and acetonitrile with 0.1 % formic acid as mobile phases for the gradient.

Mass spectrometry detection was performed on a Waters Xevo TQMS instrument equipped with an electrospray ionisation (ESI) source. The capillary voltage was 3.5 kV in positive mode and −2.5 kV in negative mode; the source was maintained at 150 °C; the desolvation temperature was 500 °C; cone gas flow was 50 L/h; and desolvation gas flow was 800 L/h. Unit resolution was applied to each quadrupole. Prior to UPLC-MS/MS MRM analysis, infusion of each individual standard was performed to optimize the MRM conditions for quantification. Data processing was performed using the Mass Lynx TargetLynx Application Manager (Waters).

Ultra-Performance Liquid Chromatography - Traveling Wave Ion Mobility-Mass Spectrometry - Quadrupole Time of Flight (UPLC-TWIMS-QToF) Mass Spectrometry and tandem mass spectrometry (UPLC-MS/MS) analysis

The metabolite profile of Col-0 and *f3h*/*fls1*/*ans* biological replicate sample extracts were acquired in an UPLC Acquity I system (Waters Corporation) coupled to a high-resolution mass spectrometer HDMS Synapt XS (Waters Corporation). The chromatographic methodology was performed according to Arapitsas *et al.* (103) with some modifications. The chromatographic separation was carried out in an Acquity Premier HSS T3 1.8 µm, 2.1 x 100 mm column (Waters) at 40 °C. The mobile phases A and B consisted of water with 0.1 % of formic acid and methanol with 0.1 % formic acid, respectively, pumped with a flow rate of 0.3 mL/min. High definition MS^e ($HDMS^e$) data acquisition with a scan time of 0.2 s was performed in continuum mode with a mass range of 50-2000 Da in resolution mode (≥ 21500 FWHM at *m/z* 785), for both ionisation modes (ESI+ and ESI-). The source temperature was settled at 120 °C, the desolvation temperature at 450 °C, the nebuliser gas volume flow was 50 L/h and the desolvation gas volume flow rate was 800 L/h. For ESI- acquisition mode, the capillary voltage

was 2.0 kV, and sampling cone was 30 V, whereas for ESI+ mode, the capillary voltage was

3.0 kV, and the sampling cone was 40 V. A collision energy ramp of 20-40 V was applied in

the transfer cell for high energy. For the ion mobility separation, the helium cell flow rate was

180 mL/min and for nitrogen it was 90 mL/min. The wave velocity was set to 650 m/s and the

transfer wave velocity to 141 m/s. The calibration was performed using the Major Mix IMS/ToF

Calibration kit solution (Waters Corporation). A leucine enkephalin solution (200 pg/μL) was

continuously infused as lock mass at flow rate of 10 μL/min during the whole analysis.

MassLynx v.4.2 and UNIFI software (Waters Corporation) were used for data acquisition and

processing. The collision cross sections (CCS) obtained were compared with an *in-house*

standard library.

A target UPLC-MSMS analysis was performed with the same source conditions in resolution

mode in a range of 50-2000 Da, with a scan time of 1 s. The collision energy was applied as a

ramp in the trap from 10-50 V for ESI- and 10-40 V for ESI+.

Determination of anthocyanin accumulation

Anthocyanin accumulation was visualised in 6-days-old seedlings grown on filter paper soaked

with 3 ppm of the bleaching herbicide norflurazon supplemented with 4 % sucrose to induce

anthocyanin accumulation. In addition, anthocyanin content was quantified photometrically in

6-days-old seedlings grown on 0.5 MS plates with 4 % sucrose as described by Mehrtens *et*

al. (104). Three independent biological replicates were measured for each sample.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

All sequence data generated in this study are available under the ENA/NCBI Bioproject ID PRJEB60353. In detail, the RNA-Seq data of *A. thaliana* Col-0, Nö-0, and the *f3h/fls1/ans* mutant can be found with under the ENA/NCBI IDs: ERS14716218 - ERS14716221, ERS14716222 - ERS14716225 and ERS14716226 - ERS14716229, respectively. The genomic ONT data can be accessed using the ENA/NCBI IDs: ERS14716230 (FASTQ) and ERS14716231 (FAST5). The *A. thaliana f3h/fls1/ans* mutant ONT assembly can be accessed at the DOI: 10.4119/unibi/2979807.

Cometing interests

The authors declare that they have no competing interest.

Funding

This research was funded by the BMBF project RaPEQ, grant numbers "FKZ 031B0888A" (RaPEQ-2) and "FKZ 031B1305B" (RaPEQ-3), as well as by core funding of the Chair of Genetics and Genomics of Plants provided by Bielefeld University through Faculty of Biology. We acknowledge support for the publication costs by the Open Access Publication Fund of Bielefeld University and the Deutsche Forschungsgemeinschaft (DFG).

Authors' contributions

MB, HMS, RS and BW conceived and designed the research. MB, HMS, RS and SM carried out experiments and analysed the data. VS performed high resolution mass spectrometry experiments and interpreted the data. AA performed the metabolite quantification methodology. RS and BW supervised the project. MB and HMS drafted the manuscript. RS, SM and BW revised the manuscript.

Acknowledgments

We thank Andrea Voigt, Melanie Kuhlmann and Nicoletta Schmidt for excellent technical assistance and Martin Sagasser for fruitful discussions. Moreover, we thank Nils Kleinbölting

712   for his support regarding *loreta*. We thank the Center for Biotechnology (CeBiTec) at Bielefeld

713   University for providing an environment to perform the computational analyses.

Supplementary Material

716   Supplementary Table S1: *A. thaliana f3h/fls1/ans* mutant *de novo* ONT genome assembly
717   statistics.
718
719   Supplementary Table S2: Oligonucleotide primers used in this work.
720
721   Supplementary Table S3: Polyphenol quantification in siliques of *A. thaliana* wildtypes and
722   2-ODD mutants.
723
724   Supplementary Table S4: Annotation of the eriodictyol and dihydroquercetin profile of siliques
725   of *A. thaliana* wildtypes and *f3h/fls1/ans* mutant.
726
727   Supplementary Table S5: Polyphenol quantification in siliques of *A. thaliana* wildtype and
728   *f3h/fls1/ans* mutant.
729
730   Supplementary Table S6: Dihydroquercetin and eriodictyol profile of siliques of *A. thaliana*
731   *f3h/fls1/ans* mutant.
732
733   Supplementary File S1: Differentially expressed genes between seedlings of different
734   genotypes.
735
736   Supplementary Figure S1: Additional T-DNA insertions in the *f3h/fls1/ans* mutant identified by
737   loreta.
738
739   Supplementary Figure S2: Genomic and transcriptomic characterisation of the mutant loci of
740   the *A. thaliana f3h/fls1/ans* mutant.
741
742   Supplementary Figure S3: The *f3h* locus of the *A. thaliana f3h/fls1/ans* mutant.
743
744   Supplementary Figure S4: *In situ* flavonoid staining of siliques from *A. thaliana* wildtype
745   and 2-ODD mutant plants.
746
747   Supplementary Figure S5: *In situ* flavonoid staining of inflorecences from *A. thaliana* wildtype
748   and 2-ODD mutant plants.
749
750   Supplementary Figure S6: Comparison of the eriodictyol profile between the *A. thaliana*
751   *f3h/fls1/ans* mutant and Col-0 in UPLC-HDMSe ESI+.
752
753   Supplementary Figure S7: Comparison of the eriodictyol profile between the *A. thaliana*
754   *f3h/fls1/ans* mutant and Col-0 in UPLC-HDMSe ESI-.
755
756   Supplementary Figure S8: UPLC-MS/MS profile of one biological replicate from the *A. thaliana*
757   *f3h/fls1/ans* mutant in ESI- mode.
758

759 Supplementary Figure S9: ESI- fragmentation of dihydrokaempferol standard by UPLC-
760 MS/MS and UPLC-TWIM-QToF (HDMSe) high resolution mass spectrometry.
761
762 Supplementary Figure S10: Fragmentation of eriodictyol standard by UPLC-MS/MS and
763 UPLC-TWIM-QToF (HDMSe) high resolution mass spectrometry.
764
765 Supplementary Figure S11: ESI- UPLC-TWIM-QToF (HDMSe) high resolution mass
766 spectrometry of compound 1 and 8 (Supplementary Table S4) in the *f3h/fls1/ans* triple mutant
767 biological replicate.
768
769 Supplementary Figure S12: The *F3'H* locus of the *A. thaliana f3h/fls1/ans* mutant.
770

References

772
1.   Wen W, Alseekh S, Fernie AR. Conservation and diversification of flavonoid
metabolism in the plant kingdom. Current Opinion in Plant Biology. 2020;55:100-8.

2.   Winkel-Shirley B. Flavonoid biosynthesis. A colorful model for genetics, biochemistry,
cell biology, and biotechnology. Plant Physiology. 2001;126:485-93.

3.   Ishikura N, Yoshitama K. Anthocyanin-Flavonol Co-pigmentation in Blue Seed Coats
of Ophiopogon jaburan. Journal of Plant Physiology. 1984;115(2):171-5.

4.   Gronquist M, Bezzerides A, Attygalle A, Meinwald J, Eisner M, Eisner T. Attractive and
defensive functions of the ultraviolet pigments of a flower *(Hypericum calycinum)*. Proceedings
of the National Academie of Sciences of the United States of America. 2001;98(24):13745-50.

5.   Grotewold E. The Science of Flavonoids. Grotewold E, editor. Columbus, Ohio, USA:
Springer New York, NY; 2006. 274 p.

6.   Peer WA, Murphy AS. Flavonoids and auxin transport: modulators or regulators?
Trends in Plant Science. 2007;12(12):556-63.

7.   Mo Y, Nagel C, Taylor LP. Biochemical complementation of chalcone synthase mutants
defines a role for flavonols in functional pollen. Proceedings of the National Academy of
Sciences of the United States of America. 1992;89:7213-7.

8.   Li J, Ou-Lee TM, Raba R, Amundson RG, Last RL. Arabidopsis Flavonoid Mutants Are
Hypersensitive to UV-B Irradiation. The Plant Cell. 1993;5(2):171-9.

9.   Xie P, Shi JY, Tang SY, Chen CX, Khan A, Zhang FX, et al. Control of Bird Feeding
Behavior by Tannin1 through Modulating the Biosynthesis of Polyphenols and Fatty Acid-
Derived Volatiles in Sorghum. Molecular Plant. 2019;12(10):1315-24.

10.   Dauer A, Hensel A, Lhoste E, Knasmüller S, Mersch-Sundermann V. Genotoxic and
antigenotoxic effects of catechin and tannins from the bark of Hamamelis virginiana L. in
metabolically competent, human hepatoma cells (Hep G2) using single cell gel
electrophoresis. Phytochemistry. 2003;63(2):199-207.

11.   Kandil FE, Smith MA, Rogers RB, Pepin MF, Song LL, Pezzuto JM, et al. Composition
of a chemopreventive proanthocyanidin-rich fraction from cranberry fruits responsible for the

inhibition of 12-O-tetradecanoyl phorbol-13-acetate (TPA)-induced ornithine decarboxylase
(ODC) activity. Journal of Agricultural and Food Chemistry. 2002;50(5):1063-9.

12.   Schijlen E, Ric de Vos CH, Jonker H, van den Broeck H, Molthoff J, van Tunen A, et al.
Pathway engineering for healthy phytochemicals leading to the production of novel flavonoids
in tomato fruit. Plant Biotechnology Journal. 2006;4(4):433-44.

13.   Reddy AM, Reddy VS, Scheffler BE, Wienand U, Reddy AR. Novel transgenic rice
overexpressing anthocyanidin synthase accumulates a mixture of flavonoids leading to an
increased antioxidant potential. Metabolic Engineering. 2007;9(1):95-111.

14.   Scharbert S, Hofmann T. Molecular definition of black tea taste by means of quantitative
studies, taste reconstitution, and omission experiments. Journal of Agricultural and Food
Chemistry. 2005;53(13):5377-84.

15.   Koornneef M. Mutations affecting the testa colour in Arabidopsis. Arabidopsis
Information Services. 1990;27:1-4.

16.   Ito T, Motohashi R, Kuromori T, Mizukado S, Sakurai T, Kanahara H, et al. A new
resource of locally transposed Dissociation elements for screening gene-knockout lines in
silico on the Arabidopsis genome. Plant Physiology. 2002;129(4):1695-9.

17.   Rosso MG, Li Y, Strizhov N, Reiss B, Dekker K, Weisshaar B. An *Arabidopsis thaliana*
T-DNA mutagenised population (GABI-Kat) for flanking sequence tag based reverse genetics.
Plant Molecular Biology. 2003;53(1-2):247-59.

18.   Alonso JM, Stepanova AN, Leisse TJ, Kim CJ, Chen H, Shinn P, et al. Genome-wide
insertional mutagenesis of Arabidopsis thaliana. Science. 2003;301(5633):653-7.

19.   Ferrer J-L, Jez JN, Bowman ME, Dixon RA, Noel JP. Structure of chalcone synthase
and the molecular basis of plant polyketide biosynthesis. Nature Structural Biology.
1999;6(8):775-84.

20.   Jez JM, Bowman ME, Dixon RA, Noel JP. Structure and mechanism of the
evolutionarily unique plant enzyme chalcone isomerase. Nature Structural Biology.
2000;7(9):786-91.

21.   Owens DK, Crosby KC, Runac J, Howard BA, Winkel BS. Biochemical and genetic
characterization of Arabidopsis flavanone 3beta-hydroxylase. Plant Physiology and
Biochemistry. 2008;46(10):833-43.

22.   Schoenbohm C, Martens S, Eder C, Forkmann G, Weisshaar B. Identification of the
Arabidopsis thaliana flavonoid 3'-hydroxylase gene and functional expression of the encoded
P450 enzyme. Biological Chemistry. 2000;381:749-53.

23.   Martens S, Forkmann G. Cloning and expression of flavone synthase II from Gerbera
hybrids. The Plant Journal. 1999;20(5):611-8.

24.   Gebhardt YH, Witte S, Steuber H, Matern U, Martens S. Evolution of flavone synthase
I from parsley flavanone 3beta-hydroxylase by site-directed mutagenesis. Plant Physiology.
2007;144(3):1442-54.

25.   Ali MB, McNear DHJ. Induced transcriptional profiling of phenylpropanoid pathway
genes increased flavonoid and lignin content in Arabidopsis leaves in response to microbial
products. BMC Plant Biology. 2014;14(84).

26.   Falcone Ferreyra ML, Emiliani J, Rodriguez EJ, Campos-Bermudez VA, Grotewold E,
Casati P. The Identification of Maize and Arabidopsis Type I FLAVONE SYNTHASEs Links
Flavones with Hormones and Biotic Interactions. Plant Physiology. 2015;169(2):1090-107.

27.   Owens DK, Alerding AB, Crosby KC, Bandara AB, Westwood JH, Winkel BS.
Functional analysis of a predicted flavonol synthase gene family in Arabidopsis. Plant
Physiology. 2008;147(3):1046-61.

28.   Shirley BW, Hanley S, Goodman HM. Effects of ionizing radiation on a plant genome:
analysis of two Arabidopsis transparent testa mutations. The Plant Cell. 1992;4:333-47.

29.   Davies KM, Schwinn KE, Deroles SC, Manson DG, Lewis DH, Bloor SJ, et al.
Enhancing anthocyanin production by altering competition for substrate between flavonol
synthase and dihydroflavonol 4-reductase. Euphytica. 2003;131(3):259-68.

30.   Wilmouth RC, Turnbull JJ, Welford RW, Clifton IJ, Prescott AG, Schofield CJ. Structure
and mechanism of anthocyanidin synthase from *Arabidopsis thaliana*. Structure.
2002;10(1):93-103.

31.   Xie DY, Sharma SB, Paiva NL, Ferreira D, Dixon RA. Role of anthocyanidin reductase,
encoded by BANYULS in plant flavonoid biosynthesis. Science. 2003;299(5605):396-9.

32.   Zhang Y, Butelli E, Martin C. Engineering anthocyanin biosynthesis in plants. Current
Opnion in Plant Biology. 2014;19:81-90.

33.   Tohge T, Nishiyama Y, Hirai MY, Yano M, Nakajima J, Awazuhara M, et al. Functional
genomics by integrated analysis of metabolome and transcriptome of Arabidopsis plants over-
expressing an MYB transcription factor. The Plant Journal. 2005;42(2):218-35.

34.   Jones P, Messner B, Nakajima J, Schaffner AR, Saito K. UGT73C6 and UGT78D1,
glycosyltransferases involved in flavonol glycoside biosynthesis in *Arabidopsis thaliana*.
Journal of Biological Chemistry. 2003;278(45):43910-8.

35.   Kim JH, Kim BG, Park Y, Ko JH, Lim CE, Lim J, et al. Characterization of flavonoid 7-
O-glucosyltransferase from Arabidopsis thaliana. Bioscience, Biotechnology, and
Biochemistry. 2006;70(6):1471-7.

36.   Cheng AX, Han XJ, Wu YF, Lou HX. The function and catalysis of 2-oxoglutarate-
dependent oxygenases involved in plant flavonoid biosynthesis. International Journal of
Molecular Sciences. 2014;15(1):1080-95.

37.   Kawai Y, Ono E, Mizutani M. Evolution and diversity of the 2-oxoglutarate-dependent
dioxygenase superfamily in plants. The Plant Journal. 2014;78(2):328-43.

38.   Prescott AG, Lloyd MD. The iron(II) and 2-oxoacid-dependent dioxygenases and their
role in metabolism. Natural Product Reports. 2000;17(4):367-83.

39.   Trewick SC, Henshaw TF, Hausinger RP, Lindahl T, Sedgwick B. Oxidative
demethylation by Escherichia coli AlkB directly reverts DNA base damage. Nature.
2002;419(6903):174-8.

40.   Turnbull JJ, Nakajima J, Welford RW, Yamazaki M, Saito K, Schofield CJ. Mechanistic
studies on three 2-oxoglutarate-dependent oxygenases of flavonoid biosynthesis:
anthocyanidin synthase, flavonol synthase, and flavanone 3-beta-hydroxylase. Journal of
Biological Chemistry. 2004;279(2):1206-16.

41.   Martens S, Preuß A, Matern U. Multifunctional flavonoid dioxygenases: Flavonol and
anthocyanin biosynthesis in Arabidopsis thaliana L. Phytochemistry. 2010;71(10):1040-9.

42.   Busche M, Acatay C, Martens S, Weisshaar B, Stracke R. Functional Characterisation
of Banana (Musa spp.) 2-Oxoglutarate-Dependent Dioxygenases Involved in Flavonoid
Biosynthesis. Frontiers in Plant Science. 2021;12:701780.

43.   Turnbull JT, Sobey WJ, Aplin RT, Hassan A, Firmin JL, Schofield CJ, et al. Are
anthocyanidins   the   immediate   products   of   anthocyanidin   synthase?   Chemical
Communications. 2000:2473-4.

44.   Yan Y, Chemler J, Huang L, Martens S, Koffas MA. Metabolic engineering of
anthocyanin biosynthesis in Escherichia coli. Applied and Environmental Microbiology.
2005;71(7):3617-23.

45.   Park S, Kim DH, Park BR, Lee JY, Lim SH. Molecular and Functional Characterization
of Oryza sativa Flavonol Synthase (OsFLS), a Bifunctional Dioxygenase. Journal of
Agricultural and Food Chemistry. 2019;67(26):7399-409.

46.   Schilbert HM, Schöne M, Baier T, Busche M, Viehöver P, Weisshaar B, et al.
Characterization of the *Brassica napus* flavonol synthase gene family reveals bifunctional
flavonol synthases. Frontiers in Plant Science. 2021;12:733762.

47.   Yonekura-Sakakibara K, Higashi Y, Nakabayashi R. The Origin and Evolution of Plant
Flavonoid Metabolism. Frontiers in Plant Science. 2019;10:943.

48.   Prescott AG, Stamford NPJ, Wheeler G, Firmin JL. In vitro properties of a recombinant
flavonol synthase from *Arabidopsis thaliana*. Phytochemistry. 2002;60(6):589-93.

49.   Stracke R, De Vos RCH, Bartelniewoehner L, Ishihara H, Sagasser M, Martens S, et
al. Metabolomic and genetic analyses of flavonol synthesis in *Arabidopsis thaliana* support the
in vivo involvement of leucoanthocyanidin dioxygenase. Planta. 2009;229(2):427-45.

50.   Wisman E, Hartmann U, Sagasser M, Baumann E, Palme K, Hahlbrock K, et al. Knock-
out mutants from an En-1 mutagenized *Arabidopsis thaliana* population generate new
phenylpropanoid biosynthesis phenotypes. Proceedings of the National Academy of Sciences
of the United States of America. 1998;95(21):12432-7.

51.   Abrahams S, Lee E, Walker AR, Tanner GJ, Larkin PJ, Ashton AR. The Arabidopsis
TDS4 gene encodes leucoanthocyanidin dioxygenase (LDOX) and is essential for
proanthocyanidin synthesis and vacuole development. The Plant Journal. 2003;35(5):624-36.

52.   Bowerman PA, Ramirez MV, Price MB, Helm RF, Winkel BS. Analysis of T-DNA alleles
of flavonoid biosynthesis genes in Arabidopsis ecotype Columbia. BMC Research Notes.
2012;5.

53.   Routaboul JM, Kerhoas L, Debeaujon I, Pourcel L, Caboche M, Einhorn J, et al.
Flavonoid diversity and biosynthesis in seed of *Arabidopsis thaliana*. Planta. 2006;224(1):96-
107

54.   Pelletier MK, Burbulis IE, Shirley BW. Disruption of specific flavonoid genes enhances
the accumulation of flavonoid enzymes and endproducts in Arabidopsis seedlings. Plant
Molecular Biology. 1999;40(1):45-54.

55.　Appelhagen I, Thiedig K, Nordholt N, Schmidt N, Huep G, Sagasser M, et al. Update
on *transparent testa* mutants from *Arabidopsis thaliana*: characterisation of new alleles from
an isogenic collection. Planta. 2014;240(5):955-70.

56.　Jupe F, Rivkin AC, Michael TP, Zander M, Motley ST, Sandoval JP, et al. The complex
architecture and epigenomic impact of plant T-DNA insertions. PLoS Genetics.
2019;15(1):e1007819.

57.　Pucker B, Kleinbolting N, Weisshaar B. Large scale genomic rearrangements in
selected Arabidopsis thaliana T-DNA lines are caused by T-DNA insertion mutagenesis. BMC
Genomics. 2021;22(1):599.

58.　Rajapriya V, Kannan P, Sridevi G, Veluthambi K. A rare transgenic event of rice with
Agrobacterium binary vector backbone integration at the right T-DNA border junction. J Plant
Biochem Biot. 2021;30(3):421-8.

59.　Wu H, Sparks CA, Jones HD. Characterisation of T-DNA loci and vector backbone
sequences in transgenic wheat produced by Agrobacterium-mediated transformation.
Molecular Breeding. 2006;18(3):195-208.

60.　Monte E, Alonso JM, Ecker JR, Zhang YL, Li X, Young J, et al. Isolation and
characterization of phyC mutants in Arabidopsis reveals complex crosstalk between
phytochrome signaling pathways. The Plant Cell. 2003;15(9):1962-80.

61.　Gao J, Peng H, Chen FB, Luo M, Li WB. Genome-wide analysis of transcription factors
related to anthocyanin biosynthesis in carmine radish (Raphanus sativus L.) fleshy roots.
PeerJ. 2019;7:e8041.

62.　Liu J, Deng S, Wang H, Ye J, Wu HW, Sun HX, et al. CURLY LEAF Regulates Gene
Sets Coordinating Seed Size and Lipid Biosynthesis. Plant Physiology. 2016;171(1):424-36.

63.　Mergner J, Frejno M, List M, Papacek M, Chen X, Chaudhary A, et al. Mass-
spectrometry-based draft of the Arabidopsis proteome. Nature. 2020;579(7799):409-14.

64.　Reinhardt N. Untersuchung der katalytischen Eigenschaften Tropinonreduktase-
ähnlicher kurzkettiger Dehydrogenasen/Reduktasen aus Brassicaceen. Halle (Saale): Martin-
Luther-Universität Halle-Wittenberg; 2014.

65.　Xu Z, Escamilla-Trevino L, Zeng L, Lalgondar M, Bevan D, Winkel B, et al. Functional
genomic analysis of *Arabidopsis thaliana* glycoside hydrolase family 1. Plant Molecular
Biology. 2004;55(3):343-67.

66.　Ishihara H, Tohge T, Viehöver P, Fernie AR, Weisshaar B, Stracke R. Natural variation
in flavonol accumulation in Arabidopsis is determined by the flavonol glucosyltransferase
BGLU6. Journal of Experimental Botany. 2016;67(5):1505-17.

67.　Miyahara T, Sakiyama R, Ozeki Y, Sasaki N. Acyl-glucose-dependent
glucosyltransferase catalyzes the final step of anthocyanin formation in Arabidopsis. Journal
of Plant Physiology. 2013;170(6):619-24.

68.　Sugiyama R, Li R, Kuwahara A, Nakabayashi R, Sotta N, Mori T, et al. Retrograde
sulfur flow from glucosinolates to cysteine in Arabidopsis thaliana. Proceedings of the National
Academy of Sciences of the United States of America. 2021;118(22):e2017890118.

69.　Duggleby RG, McCourt JA, Guddat LW. Structure and mechanism of inhibition of plant
acetohydroxyacid synthase. Plant Physiology and Biochemistry. 2008;46(3):309-24.

70. Turnbull JJ, Nagle MJ, Seibel JF, Welford RW, Grant GH, Schofield CJ. The C-4 stereochemistry of leucocyanidin substrates for anthocyanidin synthase affects product selectivity. Bioorganic & Medicinal Chemistry Letters. 2003;13(21):3853-7.

71. Welford RWD, Turnbull JJ, Claridge TDW, Prescott AG, Schofield CJ. Evidence for oxidation at C-3 of the flavonoid C-ring during anthocyanin biosynthesis. Chemical Communications. 2001(18):1828-9.

72. Preuss A, Stracke R, Weisshaar B, Hillebrecht A, Matern U, Martens S. *Arabidopsis thaliana* expresses a second functional flavonol synthase. FEBS Letters. 2009;583(12):1981–6.

73. Li DD, Ni R, Wang PP, Zhang XS, Wang PY, Zhu TT, et al. Molecular Basis for Chemical Evolution of Flavones to Flavonols and Anthocyanins in Land Plants. Plant Physiology. 2020;184(4):1731-43.

74. Pucker B, Iorizzo M. Apiaceae FNS I originated from F3H through tandem gene duplication. PLoS One. 2023;18(1):e0280155.

75. Han XJ, Wu YF, Gao S, Yu HN, Xu RX, Lou HX, et al. Functional characterization of a *Plagiochasma appendiculatum* flavone synthase I showing flavanone 2-hydroxylase activity. FEBS Letters. 2014;588(14):2307-14.

76. Lee YJ, Kim JH, Kim BG, Lim Y, Ahn JH. Characterization of flavone synthase I from rice. BMB Reports. 2008;41(1):68-71.

77. Wang Y, Shi Y, Li K, Yang D, Liu N, Zhang L, et al. Roles of the 2-Oxoglutarate-Dependent Dioxygenase Superfamily in the Flavonoid Pathway: A Review of the Functional Diversity of F3H, FNS I, FLS, and LDOX/ANS. Molecules. 2021;26(21):6745.

78. Lukacin R, Wellmann F, Britsch L, Martens S, Matern U. Flavonol synthase from *Citrus unshiu* is a bifunctional dioxygenase. Phytochemistry. 2003;62(3):287-92.

79. Zhang JR, Trossat-Magnin C, Bathany K, Negroni L, Delrot S, Chaudiere J. Oxidative Transformation of Dihydroflavonols and Flavan-3-ols by Anthocyanidin Synthase from Vitis vinifera. Molecules. 2022;27(3):1047.

80. Sheahan JJ, Cheong H, Rechnitz GA. The colorless flavonoids of *Arabidopsis thaliana* (Brassicaceae). I. A model system to study the orthodihydroxy structure. American Journal of Botany. 1998;85(4):467-75.

81. E. W. Unusual Flavanones from a Rare American Fern. Zeitschrift für Naturforschung C. 1981;36:604-6.

82. Reinhardt N, Fischer J, Coppi R, Blum E, Brandt W, Drager B. Substrate flexibility and reaction specificity of tropinone reductase-like short-chain dehydrogenases. Bioorganic Chemistry. 2014;53:37-49.

83. Xu F, Li L, Zhang W, Cheng H, Sun N, Cheng S, et al. Isolation, characterization, and function analysis of a flavonol synthase gene from Ginkgo biloba. Molecular Biology Reports. 2012;39(3):2285-96.

84. Ley JP, Krammer G, Reinders G, Gatfield IL, Bertram HJ. Evaluation of bitter masking flavanones from Herba Santa (Eriodictyon californicum (H. and A.) Torr., Hydrophyllaceae). Journal of Agricultural and Food Chemistry. 2005;53(15):6061-6.

85.   Buer CS, Sukumar P, Muday GK. Ethylene modulates flavonoid accumulation and
gravitropic responses in roots of Arabidopsis. Plant Physiology. 2006;140(4):1384-96.

86.   Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. Canu: scalable
and accurate long-read assembly via adaptive k-mer weighting and repeat separation.
Genome Research. 2017;27(5):722-36.

87.   Siadjeu C, Pucker B, Viehöver P, Albach DC, Weisshaar B. High Contiguity De Novo
Genome Sequence Assembly of Trifoliate Yam (Dioscorea dumetorum) Using Long Read
Sequencing. Genes. 2020;11(3):E274.

88.   Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. BUSCO:
assessing genome assembly and annotation completeness with single-copy orthologs.
Bioinformatics. 2015;31(19):3210-2.

89.   Lamesch P, Berardini TZ, Li D, Swarbreck D, Wilks C, Sasidharan R, et al. The
Arabidopsis Information Resource (TAIR): improved gene annotation and new tools. Nucleic
Acids Research. 2012;40(Database issue):D1202-D10.

90.   Li H. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics.
2018;34(18):3094-100.

91.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
Alignment/Map format and SAMtools. Bioinformatics. 2009;25(16):2078-9.

92.   Robinson JT, Thorvaldsdottir H, Winckler W, Guttman M, Lander ES, Getz G, et al.
Integrative genomics viewer. Nature Biotechnology. 2011;29(1):24-6.

93.   Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search
tool. Journal of Molecular Biology. 1990;215(3):403-10.

94.   Andrews S. FastQC, a quality control tool for high throughput sequence data. 2018
[Available from: http://www.bioinformatics.babraham.ac.uk/projects/fastqc/

95.   Lee H, Chawla HS, Obermeier C, Dreyer F, Abbadi A, Snowdon R. Chromosome-Scale
Assembly of Winter Oilseed Rape *Brassica napus*. Frontiers in Plant Science. 2020;11:496.

96.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast
universal RNA-seq aligner. Bioinformatics. 2013;29(1):15-21.

97.   Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. Bioinformatics. 2014;30(7):923-30.

98.   Cheng CY, Krishnakumar V, Chan A, Thibaud-Nissen F, Schobel S, Town CD.
Araport11: a complete reannotation of the Arabidopsis thaliana reference genome. The Plant
Journal. 2017;89(4):789-804.

99.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biology. 2014;15(12):550.

100.   Stracke R, Ishihara H, Huep G, Barsch A, Mehrtens F, Niehaus K, et al. Differential
regulation of closely related R2R3-MYB transcription factors controls flavonol accumulation in
different parts of the *Arabidopsis thaliana* seedling. The Plant Journal. 2007;50(4):660-77.

101.   Stracke R, Jahns O, Keck M, Tohge T, Niehaus K, Fernie AR, et al. Analysis of
PRODUCTION OF FLAVONOL GLYCOSIDES-dependent flavonol glycoside accumulation in

Arabidopsis thaliana plants reveals MYB11-, MYB12- and MYB111-independent flavonol glycoside accumulation. New Phytologist. 2010;188(4):985-1000.

102.   Vrhovsek U, Masuero D, Gasperotti M, Franceschi P, Caputi L, Viola R, et al. A versatile targeted metabolomics method for the rapid quantification of multiple classes of phenolics in fruits and beverages. Journal of Agricultural and Food Chemistry. 2012;60(36):8831-40.

103.   Arapitsas P, Mattivi F. LC-MS Untargeted Protocol for the Analysis of Wine. Methods in Molecular Biology. 2018;1738:225-35.

104.   Mehrtens F, Kranz H, Bednarek P, Weisshaar B. The Arabidopsis transcription factor MYB12 is a flavonol-specific regulator of phenylpropanoid biosynthesis. Plant Physiology. 2005;138(2):1083-96.