1 **Metabotropic signaling within somatostatin interneurons**
2 **controls transient thalamocortical inputs during development**
3
4

5 Deepanjali Dwivedi[1,2†], Dimitri Dumontier[3†], Mia Sherer[1,2], Sherry Lin[1], Andrea MC Mirow[1,2,3],
6 Yanjie Qiu[1,2], Qing Xu[1,2,4], Samuel A Liebman[3], Djeckby Joseph[3], Sandeep R Datta[1], Gord
7 Fishell[1,2,*], Gabrielle Pouchelon[1,2,3*].
8
9 [1] Harvard Medical School, Department of Neurobiology, Boston, Massachusetts, USA.
10 [2] Broad Institute, Stanley Center for Psychiatric Research, Cambridge, Massachusetts, USA.
11 [3] Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, USA
12 [4] Center for Genomics & Systems Biology, New York University Abu Dhabi, Abu Dhabi, UAE
13
14
15 [†] These authors contributed equally to this work
16
17 * Corresponding authors: G.P. (pouchel@cshl.edu); G.F. (gordon_fishell@hms.harvard.edu)

Abstract

During brain development, neural circuits undergo major activity-dependent restructuring. Circuit wiring mainly occurs through synaptic strengthening following the Hebbian "fire together, wire together" precept. However, select connections, essential for circuit development, are transient. They are effectively connected early in development, but strongly diminish during maturation. The mechanisms by which transient connectivity recedes are unknown.

To investigate this process, we characterize transient thalamocortical inputs, which depress onto somatostatin inhibitory interneurons during development, by employing optogenetics, chemogenetics, transcriptomics and CRISPR-based strategies. We demonstrate that in contrast to typical activity-dependent mechanisms, transient thalamocortical connectivity onto somatostatin interneurons is non-canonical and involves metabotropic signaling. Specifically, metabotropic-mediated transcription, of guidance molecules in particular, supports the elimination of this connectivity. Remarkably, we found that this developmental process impacts the development of normal exploratory behaviors of adult mice.

Introduction

Bottom-up afferents from the thalamus provide topographical precise sensory inputs to the neocortex. These sensory inputs target both excitatory and inhibitory cortical neuron populations and undergo dramatic refinement during development. It is well accepted that both genetic and activity-dependent factors are required for appropriate circuit development, but at present it is unclear how these factors interact to recalibrate thalamocortical (TC) circuits for adult function.[1–3]

In the adult somatosensory cortex, feedforward inhibition (FFI) from TC inputs onto parvalbumin (PV) inhibitory cortical interneurons (cINs), which inhibit excitatory neurons, and

feedback inhibition from excitatory neurons to somatostatin (SST) cINs play a crucial role in

regulating adult circuit function.[4–6] The developmental pattern of connectivity expressed by TC

afferents with their neuron targets, the FFI, supports the broad principle that weak connections

are strengthened over time during circuit maturation.[7,8] Specifically, TC afferents only weakly

contact excitatory pyramidal cells and PV cINs during the first postnatal week of development

and are later potentiated.[7,9] This developmental plasticity is thought to follow a Hebbian

mechanism, in which "cells that fire together, wire together".[10–12] The extent to which these are

universal mechanisms or specific to these strengthening processes remains unclear.

Remarkably, during the early postnatal week of development, SST cINs transiently receive TC

inputs. While the mechanisms underlying transient connectivity are unknown, this early

connection is involved in the TC potentiation onto the other cortical neurons.[13,14] As such, SST

cINs, which are the earliest born cINs populations, are thought to orchestrate cortical network

synchrony during development.[15–17] Here we explore the mechanisms by which TC inputs

weaken onto SST cINs and the molecular players that mediate this process.

In contrast to the means by which TC inputs mature onto PV and excitatory neurons in

the somatosensory cortex[12,18], we discovered that transient connectivity to SST cINs involves

non-canonical activity-dependent mechanisms and that this process is modulated by

metabotropic signaling. We identified the developmental postsynaptic expression of the

metabotropic glutamatergic receptor 1 (mGluR1) in SST cINs specifically, as a key player for

the TC input transient refinement through downstream gene transcriptional regulation, such as

semaphorin 3A (Sema3A). These results suggest that postsynaptic SST cINs actively regulate

circuit development by providing feedback signals, which ultimately underlie exploratory

behaviors in adult mice.

Results

Input from VB neurons in the thalamus connect to SST cINs during early postnatal development

TC afferents to SST cINs in layer 5 (L5) of the primary somatosensory cortex (S1) are strong during the first postnatal week of development but substantially weakened as maturation proceeds.[13,14] Active sensory exploration at adulthood occur through distinct thalamocortical neurons. During the first postnatal week, SST cINs primarily receive TC inputs from the VB neuron population in the thalamus, which conveys the primary sensory information[19] and recent work shows that SST cINs contribute to early, fast sensory-driven information transfer prior to active sensory exploration[16]. These results suggest that TC inputs from VB neurons (TC$_{VB}$) transiently contact SST cINs. To explore this hypothesis, we examined TC$_{VB}$ input selectivity onto SST cINs during postnatal development using optogenetic stimulation of TC$_{VB}$ terminals in L5 of S1 and measured their strength onto SST cINs (Fig. 1a). This was accomplished using a mouse driver line specific for VB thalamic neurons (Vipr2-Cre[20,21]) to activate Cre-mediated AAV-driven opsins (Fig. 1a-b). The monosynaptic excitatory postsynaptic currents (EPSCs) in SST cINs, optogenetically evoked by TC$_{VB}$, were observed to decrease over development. During early postnatal stages P6-7, which we define as the *immature* stage, TC$_{VB}$ inputs evoked larger EPSCs onto SST cINs than those observed onto this same population during more *mature* windows, P9-11 and P28-30 (Fig. 1c; Supplementary Fig. 1a), while their strength increases over time onto excitatory neurons as expected (Supplementary Fig. 1b). As, strengthening during development is commonly associated with pruning and synapse elimination, we next examined the synaptic organization underlying transient dynamics. Using pan-TC pre- and postsynaptic markers (VGluT2[22] and Homer1 respectively), we observed that the density of TC synaptic contact follows the respective physiological strength of TC inputs

onto SST and PV neurons at adulthood (Supplementary Fig. 1c-d). We examined TC synaptic

contact density during development and found that the overall density of TC synaptic contacts

onto SST cINs were consistently decreased in L5 (P5, P7, P10 and P30; Fig. 1d;

Supplementary Fig. 1e-f). Together, both anatomical and functional data showed that the

maturation of transient connectivity is associated with weakening and synaptic refinement of

TC_{VB} inputs onto SST cINs.

Postsynaptic activity of SST cINs regulates the maturation of transient TC inputs

The development of TC projections and cortical topographic maps have been shown to

be activity-dependent.[2,23,24] More specifically, Hebbian mechanisms, in which "cells that fire

together, wire together", are thought to underlie the developmental strengthening of TC inputs

onto excitatory neurons.[10–12] In addition, presynaptic activity from TC_{VB} inputs, specifically

instruct the development of their postsynaptic excitatory neuron target.[25] However, the above

experiments indicate that TC_{VB} synapses are distinct on SST cINs from the other cell types,

suggesting a postsynaptic specific mechanism for transient connectivity. Therefore, we next

investigated whether postsynaptic activity of SST cINs controls TC input refinement. We

hypothesized that, similar to Hebbian-based developmental potentiation[9], increasing

postsynaptic activity would maintain strong TC inputs onto SST neurons, while decreasing it

would promote the normal TC input weakening. Accordingly, we used designer receptors

exclusively activated by designer drugs (DREADD)[26,27] and the Clozapine-N-oxide (CNO), to

either activate (Gq) or inhibit (Gi) SST cINs during the first postnatal week (P0-P8; Fig. 2a;

Supplementary Fig. 2a)[28,29] and determine how they affected the strength of TC_{VB} inputs onto

SST cINs (Fig. 2b). Contrary to our expectations, at early mature stage, when the strength of

TC inputs onto SST cINs is normally diminished, DREADD-Gi inhibition within SST cINs failed

to decrease the strength of TC_{VB} inputs. Instead, DREADD-Gq activation promoted the normal weakening. (Fig. 2c-d; Supplementary Fig. 2b). In addition, Gq(+) SST cINs exhibit a decrease in the density of TC synaptic contacts ($VGluT2^+$/$Homer1^+$), compared to Gi(+) and control SST cINs (Fig. 2e; Supplementary Fig. 2c-d). Notably, neither DREADD-manipulations significantly affected the developmental levels of apoptosis of SST cINs (Supplementary Fig. 2e)[28,30] and their intrinsic properties only revealed limited excitability modifications that did not affect their typical electrophysiological identity (Table S1)[31]. After CNO discontinuation (P8), these effects proved temporary, as TC_{VB} inputs were no more reduced onto Gq(+) SST cINs at later mature stage, and instead displayed a mild rebound effect (Supplementary Fig. 2f). We further examined activity-dependent TC_{VB} input maturation onto SST cINs using a well-established genetic tool, the Kir2.1 hyperpolarizing potassium channel. While Kir2.1 inhibition was previously shown to disrupt TC input refinement and cortical map formation[24,32–35] the chronic hyperpolarization of SST cINs during the first postnatal weeks did not significantly affect TC_{VB} transient inputs, similar to the chemogenetic Gi-manipulation (Supplementary Fig. 2g-i). Interestingly, the absence of effect on the excitability of SST cINs upon DREADD-Gq manipulation, in contrast to DREADD-Gi and the hyperpolarizing Kir2.1, suggests that the mechanism involved in the regulation of transient connectivity is non-canonical, through metabotropic signaling rather than postsynaptic neuron excitability.

Altogether, these results suggest that TC transient connectivity onto SST cINs do not involve standard activity-dependent mechanisms during postnatal development, wherein Gq signaling promotes the normal depression of TC inputs.

mGluR1 is highly expressed in SST cINs during development

137   We hypothesize that a metabotropic Gq-signaling, endogenous to SST cINs, comparable to Gq-DREADD, might account for the normal weaking in TC afferents to SST cINs during development. We examined SST cIN-specific gene expression during the formation and maturation of TC transient connectivity. Specifically, we analyzed public single-cell RNA sequencing (scRNAseq) data from developing cINs collected during immature (at P2)[36] and early mature stages (at P10)[37]. We aligned the two datasets using canonical correlation analysis (CCA) followed by clustering using the Seurat pipeline.[38,39] We identified cIN types by curating the known inhibitory cIN markers in these cluster populations[37,40] and compared gene expression specific to SST cIN subtype clusters (Fig. 3a; Supplementary Fig. 3b) compared to PV or all other cIN types (Fig. 3b; Supplementary Fig. 3a; Table S2). Among the three best markers of immature SST cINs was Grm1, a gene coding for the mGluR1, a Gq-coupled postsynaptic metabotropic glutamatergic receptor. mGluR1 expression peaks during the immature time window and persists in this population in mature stages (Fig. 3c-e; Table S3).[41–44] Single molecule Fluorescent *in Situ* Hybridization (smFISH) confirmed the colocalization of Grm1 in SST cINs in L5 of S1 (Fig. 3f). Notably, mGluR1 expression was found to vary within discrete SST cIN subtypes, by P10, an early mature time point, in particular (Supplementary Fig. 3c-d). At both stages, the Myh8[+] SST cIN population, which is found in L5 of the cortex, as well as the Chodl[+] population, exhibit the highest expression of Grm1. Remarkably, mGluR1 immunohistochemistry revealed significant dendritic distribution of the receptor, which persists throughout adulthood, while somatic expression is mainly apparent during the immature time window (P3-P5) (Fig. 3g; Supplementary Fig. 3e). The high mGluR1 level expression at P30 suggests that mGluR1 specificity in SST cINs decreases over time through the acquisition of mGluR1 expression in other cell types, as recently described at adulthood[44,45]. These data

nominate mGluR1 as a candidate for triggering metabotropic postsynaptic signaling in SST

cINs to control transient TC inputs during early postnatal development.

mGluR1 CRISPR-deleted SST cINs retain their strong developmental TC inputs

mGluR1 is a postsynaptic metabotropic receptor primarily coupled with Gq signaling.[46]

Based on the high and specific postnatal expression of mGluR1 in SST cINs during postnatal

development, we hypothesized that mGluR1-Gq signaling promotes the SST cIN-specific TC

input refinement during development. To test this hypothesis, we investigated the function of

mGluR1 in weakening of TC transient connectivity using SST cIN specific loss-of-function. To

avoid non-cell autonomous effects of the mGluR1 deletion, from thalamic neurons in

particular[47], we developed an AAV-based CRISPR strategy to knock down (KD) mGluR1

specifically from SST cINs in S1 (Fig. 4a). Grm1 smFISH coupled with SST immunostaining,

confirmed the Grm1 transcript depletion from SST cINs at P5 and P10 using the CRISPR

strategy (Fig. 4b; Supplementary Fig. 4a-c). mGluR1 KD did not significantly alter cellular

density (Supplementary Fig. 4d) and the physiological properties of SST cINs at the early

mature stage (Table S4). We next examined the maturation of TC_{VB} inputs onto SST cINs in

this model. We measured EPSCs from mGluR1 KD SST cINs in response to optogenetic

stimulation of TC_{VB} inputs (Fig. 4c). At this stage, EPSCs of KD SST cINs were larger than

those recorded from control SST cINs (Fig. 4d; Supplementary Fig. 4e). The persistence of

developmental TC inputs on SST cINs in absence of mGluR1 was corroborated by an increase

of TC synaptic contact density, as labeled with VGluT2$^+$ and Homer1$^+$ onto KD SST cINs (Fig.

4e; Supplementary Fig. 4f). The normal TC input weakening never occurred in absence of

mGluR1, as EPSC evoked in SST cINs by TC_{VB} remained immature through P30, a stage when

S1 is mature (Supplementary Fig. 4g).[48] These results reveal that TC synaptic maturation is

regulated by mGluR1 in SST cIN during development.

mGluR1-dependent transcriptional regulation in SST cINs regulates TC transient

connectivity

mGluRs do not only modulate postsynaptic neuron activity, but have also been shown

to regulate gene transcription[49] and translation[46,50]. We therefore hypothesized that activation

of mGluR1 during the first postnatal week of development modulates transcriptional levels of

genes mediating negative feedback signaling of TC inputs. We select four gene candidates,

the function of which has been shown to be regulated by mGluR1 in the brain, and tested

whether their transcriptional levels were regulated downstream of mGluR1 in postnatal SST

cINs: two glutamate receptors, the subunit 3A of NMDA (Grin3a)[51–55] and GluD1 (Grid1)[56–58],

and two guidance molecules, Semaphorin (Sema) 3A and 7A[59,60]. Using the integrated

scRNAseq dataset, we observed that Grin3A, Grid1 and Sema3A are highly expressed in SST

cINs at immature stage, similarly to the Grm1 expression pattern (Fig. 5a; Supplementary Fig.

5a-d). We examined the transcriptional levels of these genes in the SST cINs in absence of

mGluR1 using smFISH (Fig. 5b-c). In SST cIN deleted for mGluR1, using the same CRISPR

strategy, we observed a downregulation in the RNA expression levels of Grin3A, Sema3A and

Sema7A, while Grid1 was upregulated (Fig. 5c-e; Supplementary Fig. 5e).

Sema3A is secreted and primarily acts as a repulsive guidance cue of axons, notably

through Plexin A1 interactions.[61–63] Remarkably, the examination of the expression of Plexin

A1 in the thalamus using the Allen *in situ* hybridization developing mouse brain atlas[64] revealed

similar dynamics of Plexin A1 expression during postnatal development, as Sema3A

(Supplementary Fig. 5f). Thus, in terms of both Sema3A in SST cINs and its cognate receptor

in TC afferents, the timing of their expression is consistent with these functioning to mediate

mGluR1-mediated downregulation of TC inputs to SST cINs. To further test this hypothesis,

we examined whether Sema3A is necessary to control the maturation of TC transient

connectivity onto SST cINs. To do so, we knocked down Sema3A from SST cINs using a similar

CRISPR strategy as for targeting mGluR1 (Fig. 5f) and measured TC_{VB} input strength onto

Sema3A KD and Ctrl SST cINs (Fig. 5g). At early mature stage, both amplitude and charge

distributions of Sema3A KD SST cINs responses were significantly enlarged compared to

those from control neurons (Fig. 5h), suggesting a role for Sema3A in the maturation of

transient connectivity. Altogether, these results show that mGluR1 regulates expression of

genes that function as feedback regulators of the transient connectivity between TC inputs and

SST cINs.

The persistence of developmental TC inputs alters exploratory behaviors at adulthood

Our TC_{VB} data, as well as a recent study[16], suggest that TC synapses transiently provide

inputs to SST cINs during a critical developmental window when sensory activity and SST cINs

collaborate to regulate cortical circuit maturation and assembly. We therefore hypothesized

that adult circuit functions as reflected by behavior is altered in our model of early postnatal

mGluR1-dependent SST cIN maturation within S1. The complex functions of adult SST cINs,

normally controlling feedback inhibition[4–6], long range motor and association input integration[65]

prompted us to perform unsupervised behavioral analysis, focusing on pseudo-naturalistic

open-field behaviors, using Motion Sequencing (MoSeq) on both male and female mGluR1 KD

and control (Ctrl) animals (Fig. 6a). MoSeq platform identifies and quantifies the use of the brief

action motifs (e.g., rearing, running, sniffing, referred to as syllables; Supplementary Videos)

out of which mouse behavior during exploration is organized (Fig. 6b).[66,67] We observed that,

while gross motor functions (velocity and total distance traveled within a session) were not

affected (Supplementary Fig. 6a-b), the exploration patterns of Ctrl and KD mice were distinct

(Fig. 6c). We further investigated the behavior motifs involved in this difference with syllable

examination and found that no new syllables were used when comparing the mGluR1 KD with

control mice, but that the distribution of syllables usage was significantly altered (Fig. 6d). Most

of these syllables, such as different types of grooming and rearing, are important parts of

mouse exploratory behaviors. Linear discriminant analysis (LDA) clearly separated Ctrl and

mGluR1 KD mice, suggesting clear differences in syllable usage (Fig. 6e). We examined the

syllables that distinguished Ctrl and KD mice, by sorting the absolute model weights in the LDA

and we identify the top 10 syllables, in which rearing was highly enriched (7 rears, 2 grooms

and 1 scrunch, Supplementary videos). While the syllable usages were different between Ctrl

and KD mice, the velocity during the syllables between the two groups were comparable (Table

S5). We repeated LDA on subsets of sessions and further identified two groups of syllables

that robustly distinguished Ctrl and KD mice across iterations by sex. The use of these syllables

alone could predict whether a given mouse was a Ctrl or a KD mouse well above chance (Fig.

6f). These data demonstrate that deficits in mGluR1 function in SST cINs, which we have

shown lead to pervasive developmental phenotypes, are associated with significant

modifications of their natural exploratory behaviors, as identified by unsupervised analysis. We

verified that these behavioral alterations did not originate from cell death or extensive cortical

rewiring, downstream of TC input maturation to SST cINs. While postsynaptic mGluRs have

been shown to prevent cell death[68], the postnatal mGluR1 CRISPR deletion did not induce

significant change in the number of infected SST cINs at P30 (Supplementary Fig. 4d).

Furthermore, anatomical neural pathways connecting to SST cINs using monosynaptic

retrograde rabies tracing were grossly normal, including the adult organization of TC pathways

(Supplementary Fig. 7a-g).[69,70] These results suggest that mGluR1 in SST cINs regulates the

proper maturation of adult cortical functions primarily through synaptic refinement, rather than

gross anatomical developmental events.

Discussion

TC afferents to pyramidal and PV neurons become strengthened as development

proceeds through Hebbian mechanisms. In contrast, here, we show that early TC connectivity

from VB neurons onto SST cINs recruits mGluR1-based transcription, supporting the later

elimination of this connectivity known to regulate the development of cortical networks.

Moreover, the removal of mGluR1 signaling from SST cINs during this critical period also

impacts the development of normal sensory-related exploratory behaviors (Fig. 7).

In electrophysiological development studies, the functional TC inputs to SST cINs during

development are referred as transient[13,14,16,71], as they are thought to be absent at

adulthood[4,5,72,73] However, anatomical studies[19,40,74,75] reveal robust TC connectivity to SST

cINs at adulthood, suggesting that TC synapses are not completely eliminated after maturation

and that remaining synaptic contacts are silent, with respect to fast glutamatergic

neurotransmission. Transient connectivity, therefore, only indicates physiological dynamics of

the network, as measured with electrophysiology and as compared to other cell types.

Remarkably, synaptic pruning or elimination in standard Hebbian developmental plasticity, is

associated with synaptic strengthening. However, we reveal that transient TC connectivity is

associated with a decrease of synaptic contact density. The discrepancy between physiological

and morphological synaptic assembly highlights the essential need to better understand these

types of processes in a given context, transient connectivity in particular, which would warrant

future high-resolution investigations using 3-D electron microscopy.[76]

Guidance molecules are responsible for circuit wiring and are generally associated with

global axon growth during embryonic development.[77] In contrast, postnatal activity-dependent

mechanisms of pruning and strengthening are usually linked to neurotransmitter receptors and

their composition.[23,78,79] Here, our results reveal a tight interplay between these two

mechanisms, wherein metabotropic glutamatergic receptor signaling, rather than cell

excitability controls expression levels of guidance molecules.

While mGluR1 is highly specific to SST cINs, primarily during development and in the

cortex, mGluR1 expression is also found at high levels in the cerebellum and in the thalamus.

Remarkably, previous studies using mGluR1 constitutive KO, demonstrate that mGluR1

promotes synapse elimination, a process leading to synaptic strengthening and synapse

maintenance in the cerebellum and the thalamus.[47,59,80] However, we reveal that mGluR1-

dependent synapse refinement in SST cINs, controls the weakening of presynaptic TC inputs.

To correlate results from cell-type specific CRISPR KD and constitutive KO models, we

measured TC synaptic contact density (VGluT2+/Homer1+) onto SST cINs in a constitutive

mGluR1 KO brain and found that TC synaptic contacts were increased (data not shown), similar

to cell-type specific CRISPR mGluR1 KD. This suggests that mGluR1-dependent synaptic

elimination observed in the constitutive KO includes cell-autonomous effects and that the

deletion of mGluR1 in SST cINs within the KO occludes the contrasting maturation effects

found in the presynaptic thalamic neurons. Overall, the comparison of the models, indicates

that mGluR1 functions are highly contextualized based on both regional and temporal contexts.

Moreover, the CRISPR KD approach enabled us to study behavioral consequences of cell-

specific mGluR1 functions, as the constitutive KO mice display strong ataxia and important

motor deficits, precludes any behavioral study.

Linking neural activity with genetic programs of synaptic assembly development is essential for a better understanding of neurodevelopmental disorder etiologies. Indeed, a lack of general synaptic refinement has been associated with autism spectrum disorders (ASD).[81,82] In addition, disruption of transient brain structures and defects in inhibitory and excitatory balance have been found in autism spectrum disorders (ASD) in particular.[83] However, the role of TC transient connectivity to SST cINs in cortical dysfunctions found in neurodevelopmental disorders remains unexplored.

SST cINs in L5 of S1 are among the earliest born populations within this region. Previous works have demonstrated that they play a critical developmental role in regulating the PV cIN maturation and the formation of the feedforward inhibition to pyramidal neurons. These works, in combination with our present findings, illustrate that SST cINs play a key role in regulating the maturation of S1 cortex that is distinct from those involving Hebbian plastic mechanisms utilized to shape the TC afferents to other cortical neurons. Instead, here, we demonstrate that the dynamic connectivity of TC inputs is regulated through SST cINs using mGluR1-dependent transcriptional regulation, of Sema3A in particular, to directly weaken TC inputs during postnatal development, when S1 cortex begins to assume its mature function. The discovery of this mechanism illustrates how genetically-programmed signals within the SST cIN population allows the thalamic afferents to activate a critical developmental signal that reconfigures S1 cortical circuits. During this period, SST cINs hand over the bottom-up FFI from the thalamus to PV cINs and in turn assume their adult role in regulating local feedback signaling to L5 pyramidal neurons. Hence, in contrast to the typical activity-based mechanism by which network dynamics drive cortical maturation, the L5 SST cINs provides a critical signaling mechanism that initiates the transition from developmental bottom-up signaling to the critical period rebalancing of bottom-up and top-down cortical functions in S1 cortex.

The source of glutamate triggering mGluR1 signaling feedback to TC inputs remains to

be investigated. However, we have previously observed using retrograde labeling, that within

glutamatergic inputs received by SST cINs, VB thalamocortical inputs is one of the earliest to

connect during development, in contrast to cortico-cortical inputs that increase throughout

maturation[19], suggesting mGluR1 activation by TC inputs during early development. Since

adult mGluR1 function appear to be distinct than the one during the first postnatal week in S1[44],

the timing of glutamatergic input integration could be critical for triggering feedback signaling

from SST cINs. While retrograde labeling does not reveal major disruptions from other inputs

onto mGluR1 KD SST cINs, it would be of interest to examine compensatory effects on SST

cINs at the level of synaptic physiology.

The feedback mechanism of sensory TC transient connectivity onto SST cINs occurs

during critical periods of plasticity, when somatosensory topographical maps are highly

sensitive to sensory TC inputs.[78,84] In addition, SST cINs have been directly involved in inducing

these critical periods of plasticity in the visual cortex.[85,86] Therefore, our findings provide a

potential genetically-encoded mechanism, by which SST cINs controls TC input-driven cortical

plasticity and ultimately somatosensory cortical functions. Notably, this mGluR1-based

feedback mechanism is specific to development, as mGluR1 at adulthood has been shown to

trigger SST cIN potentiation in adult prefrontal cortex.[44] The opposite effects of mGluR1 at

different stages, suggests a distinct SST cIN-specific molecular context, such as Sema3A

expression, controlling age-dependent functions. Moreover, this highlights the essential role of

these developmental programs, rather than adult plasticity mechanisms for the formation of

proper adult cortical circuits.

It remains to be determined whether these developmental genetic programs in SST cINs

are sufficient to instruct feedback transient connectivity and input/output circuit reorganization.

Interestingly, our postnatal scRNAseq analysis revealed that mGluR1 is also expressed

developmentally in VIP cINs (Sncg[+] and Vip[+] cIN clusters; Fig. 3a,d), which have been shown

to receive transient TC inputs during development.[87] Therefore, mGluR1 is a strong candidate

for intrinsic encoding of developmental transient TC connectivity and bottom-up/top-down

circuit wiring. The model in which cell type genetic identity is sufficient to determine feedback

TC input maturation provides for a powerful framework to generates circuit diversity from

common TC afferents, such as the formation of feedforward versus feedback inhibition formed

by PV and SST cINs respectively.

Methods

Mice

All experiments were approved by and in accordance with Harvard Medical School IACUC

protocol number IS00001269 and by Cold Spring Harbor Laboratory IACUC protocol number

22-4. Animals were group housed and maintained under standard, temperature-controlled

laboratory conditions. Mice were kept on a 12:12 light/dark cycle and received water and food

ad libitum. C57Bl/6 mice were used for breeding with transgenic mice. Transgenic mice, SST-

Cre (stock number: 013044)[88], SST-FlpO (stock number: 031629)[89], Vipr2-Cre-neo (stock

number: 031332; RRID:IMSR_JAX:031332)[90], Calb2-Cre (stock number: 010774;

RRID:IMSR_JAX:010774)[88], LSL-Cas9-eGFP (stock number: 026175)[91], RCE:LoxP (stock

number: 032037)[92], PV-Cre (stock number: 017320; RRID:IMSR_JAX:017320)[93] are available

at Jackson Laboratories. Mice were injected at P0 and experiments conducted between ages

P3-P7 for developmental stages and between ages P9-11 and P28-37 for adult/mature time

points. Both female and male animals were used for all experiments.

Histology

Immunohistochemistry (IHC) - Mice were deeply anesthetized with sodium pentobarbital by intraperitoneal injection and transcardially perfused with PBS 1X followed by paraformaldehyde (PFA) dilutes at 4% in PBS. Brains were postfixed for 4 hrs in 4% PFA at 4 °C, except for monosynaptic rabies tracing, for which brains were postfixed overnight. 50 μm vibratome (Leica) sections were incubated 1-2 hrs at room temperature in a blocking solution containing 3% Normal Donkey serum and 0.3% Triton X-100 in PBS and incubated 48 hrs at 4°C with primary antibodies: rat anti-RFP (1:1,000; Chromotek #5f8), chicken anti-GFP (1:1,000; Aves Labs #1020), rabbit anti-Homer1b/c (1:500, Synaptic Systems #160023), guinea-pig anti-VGlut2 (1:2000, Millipore #AB2251), guinea-pig anti-VGlut2 (1:1000, Synaptic Systems #135404), rabbit anti-somatostatin (1:3,000; Peninsula Laboratories International T-4103.0050) and rabbit anti-mGluR1a (1:1000; Af811 Frontier Institute), which exhibits the best labeling compared to multiple anti-mGluR1 we tested (negative controls in constitutive mGluR1 KO did not show any signal, data not shown). Sections were rinsed three times 15 min with 0.1% Triton X-100 in PBS and incubated for 60–90 min at room temperature or overnight at 4°C with Alexa Fluor 488- , 594- or 647-conjugated donkey secondary antibodies (1:500; Thermo Fisher Science or Jackson ImmunoResearch). Sections were mounted in Fluoromount-G (Southern Biotechnology, #100241-874) before imaging.

Dual single molecule Fluorescent in Situ Hybridization (smFISH) and Immunohistochemistry (IHC) - For combined smFISH and IHC, mice were deeply anesthetized with sodium pentobarbital by intraperitoneal injection and transcardially perfused with PBS followed by 4% PFA. Brains were fixed 1hr at room temperature and 2hrs in 4% PFA at 4 °C. The brains were cryopreserved in 30% sucrose before sectioned at 20 μm using a sliding microtome (Leica). Sections were stored in a cryo-solution containing 28% (w/v) sucrose, 30% (v/v) ethylene glycol in 0.1M sodium phosphate buffer, pH 7.4, before performing smFISH experiments.

RNAscope® fluorescent *in situ* hybridization and immunohistochemistry (Dual FISH-IHC)

assay was performed using RNAscope® Multiplex Fluorescent Detection Kit v2 (ACDBio

323110) purchased from Advanced Cell Diagnostics (ACDBio). Manufacturer's protocol for

fixed frozen tissue was followed. Probes used in this study include Mm-Grm1 (ACD 449781),

Mm-Sema7A (ACD 437261-C3), Mm-Sema3A (ACD 412961). After smFISH, IHC was

performed. Sections were washed in Tris-buffered saline, pH 7.4, (TBS) with 10% Tween®20

(TBST) and blocked in 10% Normal donkey serum in TBS-0.1% Bovine serum albumin (BSA)

for 1 hour at room temperature. Primary antibodies, rat anti-RFP (1:1,000; Chromotek #5f8) to

reveal AAV-sgRNA-DIO-dTomato reporters, and rabbit anti-somatostatin (1:3,000; Peninsula

Laboratories International T-4103.0050) diluted in TBS-0.1% BSA, was applied to the sections

for 2 hours at room temperature. Sections were washed in TBST and incubated in

corresponding Alexa-conjugated secondary antibodies (1:500; Thermo Fisher A-10040), in

TBS-0.1% BSA, for 1 hour at room temperature. Sections were washed in TBST and DAPI (5

uM; Sigma D9542) was applied to each section for 30 seconds. Prolong Gold antifade mounting

medium (Thermo Fisher P36930) was used to adhere glass coverslip.

Mouse neonate stereotactic injections

For postnatal time points stereotaxic injections were possible using a neonate adaptor (Harvard

apparatus). Mouse pups were anesthetized by hypothermia 5 min and stereotaxically micro-

injected with AAV (volume 80-100 nl) at P0-P1 with a micro-injector (Nanoject III). Primary

somatosensory cortex (S1) was targeted with the following coordinates: AP+1.40, ML-1.85,

DV-0.20 from Lambda. For slice physiology experiments, double injection of in S1 cortex and

the somatosensory thalamus of Vipr2-Cre or Cab2-Cre mice were performed at P0. Thalamus

coordinates for VB and PO were as follows: AP+0.9, ML-1.0, DV-2.4 from Lambda. For more

consistent targeting, cortical injections were performed over 5 min and thalamic injections over

10 min, following a 20 nl, 4-5 time with 10s delay program. After surgery, mice were given

Meloxicam (Metacam) subcutaneously at 5 mg/kg of body weight and, upon recovery on a

heating pad set at 37 °C, were placed back in the home cage with the mother.

Chronic DREADD activation

For DREADDs experiments, Clozapine-N-Oxide (CNO, Tocris) or CNO dihydrochloride (Hello

Bio) was dissolved in 5% dimethyl sulfoxide (Sigma) or 0.9% saline respectively, at 1 mg/ml

(10x solution) and stored at 4 °C for the duration of the chronic activation (7 days). Every day,

fresh 1x CNO solution was made by fresh dilution with 0.9% saline to 0.1 mg/ml. Pups were

injected with CNO (10 ul/g equivalent to 1 mg/Kg) subcutaneously for 7 days (from P1-P8),

twice daily.

Adult stereotactic injections

For monosynaptic rabies retrograde labeling, stereotactic injection of EnVA-CVS-N2c-dG-H2B-

tdTomato (70 nL; dilution 1/10, Addgene plasmid: #175441)[19] was performed at P30 in animals

previously injected with AAV-3xgRNA-DIO-TVA-N2cG at P0 (80 nl), using a Nanoject III at 1

nl/second according to stereotaxic coordinates (from Bregma AP+1.00, ML-3.00, DV-0.89).

Animals were perfused 7 days later and processed for immunohistochemistry.

Viruses

The enhancers, reporters, and effectors were cloned using the Gibson Cloning Assembly Kit

(New England BioLabs, catalog no. NEB-E5510S). After cloning and sequencing, the growth

time of the transformed DH10B Competent Cells was kept below 12 hours on plates and 10

hours in flask at 37 C. DNA from several clones was recovered with an endotoxin free midi-kit

(Zymo D4202). This allows for consistently obtaining low recombination rates detected by PCR.

Primers were designed for amplification of plasmids at the junction of each possible

recombined form and compared to recombination-free controls. Only clones with a

recombined/not recombined ratio of 1.0E+05 or greater were considered for further AAV

production. The rAAVs were produced using standard production methods. Polyethylenimine

was used for transfection and OptiPrep gradient (Sigma) was used for viral particle purification.

Titer was estimated by quantitative PCR with primers for the WPRE sequence that is common

to all constructs.

AAV2/1-hSyn-fDIO-DREADD-Gi-mCherry and hSyn-fDIO-DREADD-Gq-mCherry were

produced from gifts from U. Gether (http://n2t.net/addgene:154867/addgene #154867;

http://n2t.net/addgene:154868/#154868). Titer: 1.80E+12 and 1.72E+12 vg/ml.

AAV2/1-hSyn-DIO-DREADD-Gi-P2A-tagBFP and DREADD-Gq-P2A-tagBFP were generated

from VTKS2 backbone[94] and produced for this manuscript. Titer: 3.6E+12 and 9.1E+11

vg/ml.

AAV8-hSyn-DIO-HA-hM3D(Gq)-IRES-mCitrine and -HA-hM4D(Gq)-IRES-mCitrine were gifts

from B. Roth (http://n2t.net/addgene:50454/addgene #50454-AAV8;

http://n2t.net/addgene:50455/addgene #50455-AAV8). Titer: 1.9E+13 and vg/ml.

AAV2/1-3x Grm1 gRNA-DIO-dTomato was generated from the VTKS2 backbone and

produced for this manuscript. Titer: 2.42E+12 vg/ml.

AAV2/1-3x Grm1 gRNA-DIO-N2cG-TVA was generated and produced for this manuscript.

Titer: 7.8E+12 vg/ml.

AAV2/1-3x Grm1 gRNA-fDIO-dTomato was generated from the VTKS3 backbone and

produced for this mansucript. Titer: 2.0E+12 vg/ml.

AAV2/1-3x Sema3a gRNA-DIO-mCherry was generated from the VTKS2 backbone and

produced for this manuscript. Titer: 1.07E+12 vg/ml.

AAV2/9-Ef1a-DIO-ChRmine-mScarlet was produced by the Children's Hospital Viral Core

from a plasmid, which was a gift from K. Deisseroth[95]

(http://n2t.net/addgene:130998/addgene#130998) produced by. Titer 6.26E+14 vg/ml.

AAV2/1-DIO-mCherry was generated from the VTKS2 backbone and produced for this

manuscript. Titer: 1.4E+12 vg/ml.

AAV PHP.eB-EF1a-DIO-hChR2(H134R)-eYFP and AAV9-EF1a-fDIO-eYFP were gifts from

K. Deisseroth (http://n2t.net/addgene:20298/addgene #20298-AAV-PHPeB[96];

http://n2t.net/addgene:55641/addgene #55641-AAV9)[97]. Titer: 1.00E+13 and 2.00E+13

vg/ml.

AAV2/1-hSyn-fDIO-HA-Kir2.1 was generated from the VTKS3 backbone and produced for

this manuscript. Titer: 2.60E+12 vg/ml.

AAV9-EF1a-fDIO-Cre was a gift from E. Engel & A. Nectow[98]

(http://n2t.net/addgene:121675/addgene #121675-AAV9). Titer: 2.50E+13 vg/ml.

AAV-PHP.eB-S5E2-dTomato was a generous gift from J. Dimidschstein. Titer: 8.3E+09

vg/ml.[99]

EnvA-pseudotyped CVS-N2c(deltaG-H2B:tdTomato) Rabies construct that we previously

generated (Addgene #175441)[19] was utilized for monosynaptic retrograde labeling. Titer:

3.7E+09U/ml. The nuclear labeling from H2B:tdTomato improves the automated cell detection

and the overall accuracy. Rabies were generously shared by K. Ritola. Titer: 1.4E+08U/ml.

CRISPR strategy

For proper deletion of Sema3A and Grm1 genes, three guide RNAs (gRNA) were packaged

into one single AAV. Each gRNA is under human U6 promoters and with a scaffold gRNA and

a polyA sequence at their 3' end. The whole triple gRNA (3x gRNAs) construct was synthesized

by GeneScript with an ApaI site at each end, to be directly inserted into a VTKS2 or VTKS3

AAV backbone[94], upstream of the human Synapsin promoter, which controlled a DIO- or a

fDIO-reporter (dTomato or N2cG rabies helpers). Each 3x gRNA was packaged into single

AAV1. Grm1 gRNA with spCas9 PAM sequences were designed by cross-validating best ON-

and OFF-target scores from Benchling CRISPR gRNA Design Tool, ChopChop[100,101] and

CRISPick (previously known as Genetic Perturbation Platform (GPP)). gRNA suggestions were

picked in Exons 1 to 3, to increase the probability of full gene deletion. Sequence of selected

Grm1 gRNAs are: 5'-CGATGCTTGATATCGTCAAG-3'; 5'-CGACCGCGTCTTCGCCACAA-3'

and 5'-GTCGCTCAGGTCTATGCTCG-3'. Sequence selected for Sema3A gRNAs are: 5'-

TACTCCGTTCTTCATCCAGA-3'; 5'-TGTGGCCAGTATCTTACACA-3'; 5'-

GAGACGTTAGTGTTGCCATG-3'.

Of note, since each guide have their own off-target effects, it has been increasingly accepted

that scrambled gRNAs are not proper controls for CRISPR, in contrast to shRNA or siRNA. In

this manuscript, Cas9(+) or animals without gRNA infection were considered as controls.

Alternatively, gRNA-DIO-reporter(+) cells were used as controls. In electrophysiology

experiments, Vipr2-Cre mouse line was utilized for opsin expression in the thalamus. Vipr2-

Cre recombination is specific to the first-order thalamus and practically absent from the entire

cortex (Allen Brain Atlas Transgenic Characterization)[64], preventing Cas9 recombination in

non-SST cINs. Moreover, SST-Cre alone was not sufficient to trigger Grm1 deletion by P5 and

P10, when injected at P0 (Data not shown). Therefore, we used a combination of Vipr2-Cre

and SST-FlpO mouse lines, together with an AAV-fDIO-Cre and the AAV-sgRNAs to trigger

CRISPR-deletion in SST cINs early in postnatal development.

10x single cell RNAseq

Publicly available scRNAseq datasets for P2 and P10 cINs were used[36,37] (Gene Expression

Omnibus (GEO) at accession number GSE165233 and GSE104158, respectively). P2

scRNAseq was obtained from Dlx6a-Cre; Ai14(+) sorted cINs and prepared with 10x v3.1

Genomics Chromium platform, with a total of 5,384 cells sequenced. P10 scRNAseq was

obtained from Dlx6a-Cre; RCELoxP sorted cINs and prepared with 10x Genomics Chromium

system, with a total of 6,346 cells sequenced. While public data are already processed, we

aligned the same and higher quality criteria to both datasets, by removing cells with a number

of UMI of 500 minimum and 10,000 maximum and cells with higher than 10% mitochondria.

Clustering of both datasets was performed using Seurat pipeline in "R"[38]. The number of

principal components used for clustering analysis were determined with the ElbowPlot function

(20 dimensions). Cells within P10 and P2 were assigned to cIN type via Seurat canonical

correlation analysis (CCA) and clusters assigned based on gene marker expression.

Cells within non-cIN clusters (excitatory neurons and glial cells) were defined based on the

expression of their developmental markers[102] and removed from the UMAP representation

(data not shown; Slc17a7, Neurod6, Mki67, Mbp, Gfap).

Marker expression of cIN types were defined as described in[37] for the main cIN types (such

SST and PV cINs) and supported with [40] to assign SST cIN subpopulations (such as Myh8+

and Chodl+ SST cINs). Marker genes for early postnatal PV cINs are: Mef2c, Erbb4, Plcxd3,

Grp149; marker genes for early postnatal SST cINs are: Sst, Tspan7, Satb1. For

subpopulations, we verified the expression of adult marker genes in the integrated P2-P10

datasets. SST cIN subtype marker genes expressed at P2/P10 are: Cbln4 (T-shaped Martinotti

cINs), Hpse (L4-projecting cINs), Myh8 (T-shaped Martinotti cINs), nNos1/Chodl (long-range

projecting cINs).

Using Seurat, we performed differential gene expression (DGE) between identified SST cIN

and PV cIN clusters, between SST cINs and all other cIN clusters, and between SST-Myh8

subtype with all other cIN subtypes, based on the non-parametric Wilcoxon rank sum test. Log2

(fold change(FC)) and adjusted p-values of the top candidates are represented in Table S3 and

S4. Adjusted p-values smaller than 2.22E−16 cannot be computed by "R". For visible volcano

plot representations, a constant corresponding to one-digit value lower than the last positive

decimal of the lowest computer p-value, as added to p-values = 0. Comparisons were done at

each age, P2 and P10, separately

Whole-cell slice electrophysiology

For P6-7 time point, injection of DIO-Chrmine at P0 generated stronger TC input stimulation

and improved optogenetic variability from the restricted AAV-expression in such short time

windows. From P10 throughout adulthood, DIO-ChR2 was sufficient to evoke more consistent

responses. P6-7 mice were anesthetized by hypothermia, followed by decapitation. P9-11 and

P28-32 mice were anesthetized using isoflurane, followed by decapitation. Brains were

dissected out in ice-cold oxygenated (95% O_2 and 5% CO_2) sucrose cutting solution containing

(in mM): 87 NaCl, 2.5 KCl, 2 MgCl2, 1 CaCl2, 1.25 NaH2PO4, 26 NaHCO3, 10 glucose and

75 sucrose (pH = 7.4). 300-µm thick coronal slices were cut using a Leica VT1000S vibratome

through the primary somatosensory cortex. Slices recovered in a holding chamber with artificial

cerebrospinal fluid (ACSF) containing (in mM) 125 NaCl, 20 Glucose, 2.5 KCl, 1.25 NaH2PO4,

26 NaHCO3, 2 CaCl2, 1MgCl2 (pH = 7.4) at 32°C for 30 minutes and room temperature for at

least 45 minutes prior to recordings. For recordings, slices were transferred to an upright

microscope (Scientifica) with oblique illumination Olympus optics. Cells were visualized using

a 60x or a 20x water immersion objective. Slices were perfused with oxygenated ACSF in a

recording chamber at room temperature at 1-2 ml/min. Recording electrodes (3–6 MΩ) were

pulled from borosilicate glass (1.5 mm OD, Harvard Apparatus) with a horizontal P-1000

Flaming Micropipette Puller (Sutter Instrument). For current-clamp recordings, electrodes were

filled with an internal solution containing (in mM): 130 K-Gluconate, 10 KCl, HEPES, 0.2 EGTA,

4 MgATP, 0.3 NaGTP, 5 Phosphocreatine and 0.4% biocytin, equilibrated with KOH at pH =

7.4. For voltage-clamp recordings, electrodes were filled with an internal solution containing (in

mM): 125 Cs-gluconate, 2 CsCl, 10 HEPES, 1 EGTA, 4 MgATP,0.3Na-GTP,8

Phosphocreatine-Tris, 1 QX-314-Cl, equilibrated with CsOH at pH = 7.4. Recordings were

performed using a Multiclamp 700B amplifier (Molecular Devices) and digitized using a

Digidata 1550A and the Clampex 10 program suite (Molecular Devices) or, using an EPC 10

amplifier (HEKA) and the Patchmaster software (HEKA). Voltage-clamp signals were filtered

at 3 kHz and recorded with a sampling rate of 10 kHz. Recordings were performed at a holding

potential of -70 mV. Cells were only accepted for analysis if the initial series resistance was

less than 40 MΩ and did not change by more than 20% during the recording period. The series

resistance was compensated at least ~50% in voltage-clamp mode, and no corrections were

made for the liquid junction potential. Whole-cell patch clamp recordings were done from

reporter-expressing SST cINs and nearby non-fluorescent pyramidal neurons in L5. fDIO-

mCherry was used for developmental analysis and all control conditions throughout the study.

To stimulate thalamic afferents expressing ChR2, blue light (5.46-6.94 mW from the objective)

was transmitted from a collimated LED (Mightex) to the epifluorescence port of the microscope.

We used a 5 ms pulse of light for the stimulation of P7 and P10 slices and a 1 ms pulse of light

for P30 slices. Pulses were delivered once every 5 s, for a total of 15 trials. To isolate

monosynaptic EPSCs from thalamus to cortex, recordings were performed in the presence of

1 μM TTX and 1 mM 4-AP (Tocris). Data analysis was performed using MATLAB and Prism 8

(Graphpad). The peak amplitude of the evoked EPSCs were averaged across the 15 trials per

cell.

For intrinsic properties, SST cINs were recorded in current-clamp mode and the data were

acquired at 20 KHz. In supplementary Table 1, Resting Membrane Potential (RMP) of the cell

is the potential when the net injected current (holding current + IV current-clamp step) is the

closest to zero. Input Resistance (IR) was computed using Ohm's law (V = I/R) by finding the

slope of the IV curve obtained using the net current from the IV current-clamp protocol sweep

with no action potential (AP) and the potential from the corresponding sweeps. Rheobase was

defined as the minimum net injected current to evoke a sweep with more than one AP. AP

threshold was defined as the max of the second derivative of the first AP from the first sweep

with more than one AP. In supplementary Table 2, Resting Membrane Potential (RMP) of the

cell was recorded as mean membrane potential from a 1 min long recording in I = 0 mode or

by taking the mean potential. Input Resistance (IR) was computed using Ohm's law (V = I/R)

by finding the slope of the IV curve obtained using current injection from -10 to 10 pA in steps

of 5 pA. Rheobase was defined as the minimum input current to evoke firing from a cell. Spike

threshold was defined as the membrane potential where dv/dt > 5 mV/ms before spike initiation.

mGluR1 KD intrinsic properties were compared to publicly available Control data of SST cINs

in somatosensory cortex at P10, recorded the same way.[19] Analyses were done using clampfit,

Easy electrophysiology and Python. Statistical analyses were done using Python (scipy)[103] and

in GraphPad prism 9.

Motion Sequencing (MoSeq)

SST-Cre:Cas9fl mice were injected with AAVs expressing mGluR1-gRNA, fDIO-Cre bilaterally

in the somatosensory cortex at P0-P1. Control animals were SST-Cre:Cas9fl injected with

fDIO-Cre-AAV or gRNA-AAV only, or SST-Cre animals injected with both fDIO-Cre- and gRNA-

AAVs. All mice were perfused after recording and analysis to confirm the bilateral targeting of

the somatosensory cortices, as in Figure 5B. Mice were recorded around 2.5 months of age.

MoSeq recording were carried out as previously described.[66,67] 3D depth video is recorded in

a circular 17'' diameter open-filed arena (OFA) using a Microsoft Kinect v2 depth camera.

Briefly, mice were freely moving in the arena for 45 mins in the dark, and put back to their home

cage. The arena is wiped down with 10% bleach, followed by 70% ethanol between each

session. After removing corrupted sessions due to a significant amount of noise in the

recording, in total 44 female mice (24 Ctrl and 20 KD) and 32 male mice (20 Ctrl and 12 KD)

were used in the analysis.

MoSeq is an unsupervised machine learning algorithm that segments animal behavior in a

novel, but neutral environment (bucket) into repeated modulated motifs called syllables. The

set of syllables represent the grammar expressed in the exploration (body language) of each

animal and each group. Each syllable is uniquely identified with a number and a global behavior

label, such as "rearing" or "grooming", which are identified with a color label in the figures and

grouped into global classes (corresponding colors).

MoSeq pipeline (http:www.moseq4all.org) takes 3D depth videos as input and produces frame

by frame syllable labels for each frame in the videos. Depth videos first go through image

processing in MoSeq package suite to crop out and align the mouse such that the center of the

mouse is at the center of the frame, and the nose of the mouse is pointing right. To

accommodate noise and morphological variations, the processed aligned video goes through

a deep neural network for denoising. The denoised data goes through dimensionality reduction

and modeling steps to generate the syllable labels. Syllables are labeled by sorted usage

across all sessions such that syllable 0 is the most used syllable, and the syllables that make

up 99% of the total frames are included in the analysis. Syllable usages by group are

aggregated across all sessions in the group (i.e. male Ctrl, male KD, female Ctrl, female KD)

and normalized such that syllable usages within one session add up to 1. Chi-square tests are

done on the unnormalized frame counts aggregated by group. MoSeq pipeline outputs frame-

by-frame kinematic values such as velocity and the coordinates for the mouse centroid in

addition to the syllable labels. The position occupancy heatmap for each session is computed

by binning the mouse centroid with 100 bins for each axis to compute 2D histogram, and the

histogram is smoothed by a median filter with kernel size 11 pixels. The group position density

is the average of all the session position density within the group. The heatmap differences are

computed by subtracting the control histogram from the KD histogram form each sex. The

average velocity in a session is the average of the frame-by-frame velocity. The total distance

traveled within a session is the sum of between frame mouse centroid movement.

Linear Discriminant Analysis (LDA) - To find the syllables that distinguished Ctrl from KD

animals, we used LDA on syllable usages across all sessions to find the linear projection that

maximizes group separability between Ctrl and KD mice, and sessions from both male and

female data in each genotype are combined.

To find a set of syllables that robustly distinguish Ctrl and KD mice by sex, we applied leave-

one-out strategy and trained multiple LDA embedding on a subset of the sessions within the

same sex, leaving one session out. The top 10 high absolute loading syllables are recorded at

each iteration and the most frequent 10 syllables among all the iterations by sex are used in

the linear classifier. There was a total of 27 unique syllables found in the iterations for each

sex. The 19 syllables that were present in male and female were used in the classifier.

Linear Classifier on syllable usages- We applied a linear classifier to the usages of the 19

syllables identified in the leave-one-out LDA described above. Since the number of Ctrl and

KD mice included in both males and females are unbalanced, the data is randomly subsampled

such that both Ctrl and KD have the same number of mice. The subsample process is repeated

10 times. After subsampling, the input data goes through a standard scaler for data

preprocessing, and logistic regression is used to predict genotype. Leave one out cross

validation is used to compute model accuracy.

Quantification and statistical analysis

All statistical details can be found both in the results and the legends of the corresponding figure. In the manuscript, "N" represents the number of animals, while "n" represents the number of cells. All statistical analyses were performed using GraphPad Prism, Python (scipy) and R software. Unless otherwise tested, statistical significance was tested with one-way ANOVA, followed by post-hoc Tukey's multiple comparison test or Student t-test for parametric data and with one-way Kruskal-Wallis followed by post-hoc Dunn's test or Mann-Whitney test for non-parametric data. P values < 0.05 were considered statistically significant. For all figures, $*p < 0.05$; $**p < 0.01$; $***p < 0.001$; $****p < 0.0001$.

The normal distribution of the synaptic density was formally tested with a Kolmogorov-Smirnov test. The data distribution of the synaptic density during development, in CRISPR KD, in Kir2.1 and in DREADD experiments did not show a normal distribution and were tested with non-parametric tests (Kruskal-Wallis and Mann-Whitney).

The data distribution of cell density and monosynaptic retrograde labeling was assumed to be normal and analyzed with parametric tests, based on previous studies, but was not formally tested.

RNA particle distribution (from smFISH) appears to be marker-specific, as previous studies use parametric and non-parametric tests. We formally tested the data distribution of Grm1 RNA particles, found they are primarily normal (with D'Agostino Pearson test. The exception was P10 data, when tested with a Kolmogorov-Smirnov test only, p-value = 0.0132) and used parametric tests. In contrast, Sema3A smFISH data distribution was not normal (Kolmogorov-Smirnov test p-value <0.0001) and non-parametric tests were performed.

The data distribution of optogenetics responses was assumed to be non-gaussian[4,104,105] and analyzed with non-parametric tests, based on previous studies, but was not formally tested.

Image acquisition and analysis

For all analyses, images were taken in layer 5 of the somatosensory cortex, except for cell number quantification, which was performed in all layers. For analysis of synapse density and smFISH, tissue samples were imaged on an upright ZEISS LSM 800 confocal using a 40X oil immersion objective, 1.4 NA, 2.5 digital zoom, 2,048 x 2,048 pixels, 16 bits. For cell density and monosynaptic retrograde labeling, tissue samples were imaged on a ZEISS Axio Imager using a 10X dry objective (with tiling mode).

Synaptic density - For TC synapse analysis onto PV and SST cINs, single planes were analyzed using a custom script in Fiji (ImageJ) software, as previously described before[106]. In brief, noise reduction and smoothing were applied to all channels and imaged converted to RGB. A color threshold was automatically set to identify the cell body of SST cINs and PV cINs labeled with the RCE:LoxP mouse line. In contrast, the reporter distribution of DREADD-Gi/Gq and as well as the somatostatin labeling were not homogeneous. In that case, manual delimitation of cell bodies was performed. Cell body perimeter was automatically measured and a masked binary image with the cell body only was created. For bouton segmentation, a watershed-based method is used such that boutons were separated based on the local minima of the pixel gray values. For the presynaptic boutons (VGluT2) and postsynaptic clusters (Homer1), a color threshold was selected to segment boutons as isolated puncta. The comparison between the original images and the masks was used to guide the choice of the threshold value, which was determined to detect all putative presynaptic boutons or postsynaptic clusters in each condition. The "Analyze Particles" (VGluT2 minimum size = 0.20 mm; Homer1 minimum size= 0.10 mm) and "Watershed" tools were applied and a mask was generated. A merged image from all masks was created, converted to an 8-bit image and the

707 overlap between presynaptic TC puncta, postsynaptic clusters and the cell body was

708 automatically detected.

709 *smFISH analysis* - single RNA molecules were quantified within SST cINs, as labeled with

710 somatostatin antibody. Control cINs were imaged in the non-injected contralateral side, while

711 KD cINs were imaged in the injected ipsilateral side and identified by the AAV-reporter

712 colocalization. Single plane images were analyzed using a similar adapted custom script in Fiji

713 (ImageJ) software. Images were converted in RGB, cell bodies first identified as control or KD

714 and then manually delimited based on the somatostatin labeling. Grm1 RNA, in particular,

715 formed dense clusters within the cells and the watershed-based method was used to separate

716 particles. A color threshold was applied to all images of the same condition to segment RNA

717 particles. The ''Analyze Particles'' (minimum size for Grm1 was determined as ~ 0.30 mm) and

718 ''Watershed'' tools were applied and a mask was generated. A merged image of the 2 masks

719 was created, converted to an 8-bit image and the overlap between SST cINs and RNA dots

720 automatically detected.

721 *Cell density* - For SST cIN density in DREADD experiments at P10, SST cINs were labeled

722 with the AAV-DIO-reporter, mCitrine for DREADD-Gi/Gq. SST cIN infected with AAV-DIO-

723 DREADD-Gi-mCitrine, from mice injected with saline instead of CNO, were the control

724 condition. For SST cIN density in mGluR1 KD CRISPR experiments at P30, SST INs were

725 identified from the LSL-Cas9eGFP labeling. Control cINs were infected by fDIO-Cre only

726 without AAV-sgRNA. Using Fiji (ImageJ) software, areas covering ~ 3-4 topographical columns

727 of the somatosensory cortex were drawn from layer 1 to the white matter. Area (mm^2) was

728 measured to normalized cell number and SST cINs were manually counted using the "cell

729 counting" plugin. Quantification of 1-to-3 sections per DREADD(+) and 4 sections per CRISPR

KD and Ctrl brains were averaged to represent 1 biological N each. The identification of each

condition was blind to the experimenter.

Monosynaptic retrograde labeling quantification - Every 4th section of a whole brain (from

orbital cortex to brainstem) was imaged and uploaded into NeuroInfo® software (MBF

Bioscience). All sections were manually reordered from rostral to caudal direction of the brain.

The software's section detection parameters were adjusted to properly recognize the borders

of each brain section. Sections were aligned, first using the software's "Most Accurate"

alignment option, and then re-adjusted manually if necessary. The distance between each

section (200 µm) was specified for the sections to be registered to a standardized 3-D mouse

brain atlas, using the "Section Registration" function of the software. For P30 mouse brains,

the last version of the Mouse Allen Brain atlas was selected. Non-linear registration was run

on each section to account for the slight distortions, such as imperfections from

sectioning/mounting and/or asymmetry from the sectioning angle. In the "Cell Detection"

function, parameters for cell size and distance from background were adjusted, although the

use of H2B:tdTomato rabies, which are nuclear, greatly improves the detection specificity

compared to other labeling. Neural Network with "pyramidal-p64-c1-v15.pbx" preset was used

to automatically detect rabies infected nuclei in the red channel. Detection results were

reviewed manually to correct for any detection mistakes (false positives or negatives). Final

results were exported as a CSV file. Retrogradely labeled neurons were grouped into functional

regions and normalized by the total of retrogradely labeled neurons per brain. Depending on

the analysis, functional groups or subgroups of regions were defined as followed: local = S1

cortex respectively; contralateral = any neurons found in the non-injected contralateral side;

other areas = all ipsilateral cortical areas except S1; primary = primary sensory areas: M1, S1,

V1, A1; association = "other areas" - "primary areas"; First order thalamic nuclei: dLG, VB, LD,

AV, VAL, MGB; Higher order = LP, PO, VM, MD, AM; Limbic = Pf, AD, CM; CL; PCN; Rhe/RH.

Hierarchical functional organization of the thalamus, as first, higher order and Limbic

populations reflect the genetic and functional identity of thalamic nuclei[106–108]. This organization

of rabies retrogradely labeled thalamic neurons was previously used to identify thalamocortical

pathway changes after sensory deprivation[19] and help us to distinguish whole thalamus versus

subpopulation changes in the function of thalamocortical pathways.

Data availability

All quantified data are available in main, supplementary figures and supplementary tables. All

original physiological, imaging, histological data are archived at Harvard Medical School Fishell

lab server and Cold Spring Harbor Laboratory, Pouchelon lab server. Any additional

information required to reanalyze the data reported in this work paper is available upon request.

Acknowledgements

We would like to thank Elaine Sevier for the discussions and none-included preliminary

analysis, Marian Fernandez-Otero for the extensive support with mouse colony/genotyping at

Harvard Medical School. This work was supported with an EMBO Long-Term fellowship, an

early- and an advanced-Swiss Foundation postdoctoral fellowships, a Hearst foundation grant

and the Pershing Square Innovation Fund (PSIF) (to G.P.), and grants from the National

Institutes of Health (NIH), MH071679, NS08297, NS074972, MH095147, as well as support

from the Simons Foundation (SFARI) (to GF).

Author contributions

G.P. and G.F. conceived the project, developed the methodology and wrote the manuscript.

G.P., D.Dwivedi and D.Dumontier analyzed and interpreted the results. D.Dumontier and G.P.

processed experimental data and designed the figures. G.P. designed the CRISPR and viral

strategies/constructs. D.Dwivedi performed electrophysiological experiments and analysis.

D.Dumontier recorded intrinsic properties from DREADD experiments and performed the

corresponding analysis. G.P. analyzed scRNAseq datasets. A.MC.M and M.S. established and

performed smFISH, respectively. S.Liebman. and D.Dumontier performed DREADD-

dependent cell death experiments. D.J. quantified cell density and was blind to the

experiments. G.P., D.Dwivedi, S.Liebman., M.S. and D.Dumontier performed stereotactic

injections. G.P. performed the synaptic density experiments and analysis. Q.X. cloned and

produced AAVs. Y.Q. completed the automated monosynaptic retrograde labeling analysis.

M.S, G.P. and Y.Q. collected MoSeq recordings. S. Lin. and B.D. processed, analyzed and

interpret the MoSeq behavioral experiments. B.D. helped writing the manuscript.

Competing interests

Gord Fishell is a founder of Regel Therapeutics, which has no competing interests with the

present manuscript.

References

1. Wamsley, B. & Fishell, G. Genetic and activity-dependent mechanisms underlying interneuron diversity. *Nat Rev Neurosci* **18**, 299–309 (2017).
2. Faust, T. E., Gunner, G. & Schafer, D. P. Mechanisms governing activity-dependent synaptic pruning in the developing mammalian CNS. *Nat Rev Neurosci* **22**, 657–673 (2021).
3. Cossart, R. & Garel, S. Step by step: cells with multiple functions in cortical circuit assembly. *Nat Rev Neurosci* 1–16 (2022) doi:10.1038/s41583-022-00585-6.
4. Porter, J. T., Johnson, C. K. & Agmon, A. Diverse Types of Interneurons Generate Thalamus-Evoked Feedforward Inhibition in the Mouse Barrel Cortex. *J. Neurosci.* **21**, 2699–2710 (2001).
5. Cruikshank, S. J., Urabe, H., Nurmikko, A. V. & Connors, B. W. Pathway-Specific Feedforward Circuits between Thalamus and Neocortex Revealed by Selective Optical Stimulation of Axons. *Neuron* **65**, 230–245 (2010).
6. Yu, J., Hu, H., Agmon, A. & Svoboda, K. Recruitment of GABAergic Interneurons in the Barrel Cortex during Active Tactile Behavior. *Neuron* **104**, 412-427.e4 (2019).
7. Isaac, J. T. R., Crair, M. C., Nicoll, R. A. & Malenka, R. C. Silent Synapses during Development of Thalamocortical Inputs. *Neuron* **18**, 269–280 (1997).
8. Vardalaki, D., Chung, K. & Harnett, M. T. Filopodia are a structural substrate for silent synapses in adult neocortex. *Nature* **612**, 323–327 (2022).
9. Daw, M. I., Ashby, M. C. & Isaac, J. T. R. Coordinated developmental recruitment of latent fast spiking interneurons in layer IV barrel cortex. *Nat Neurosci* **10**, 453–461 (2007).
10. Katz, L. C. & Shatz, C. J. Synaptic Activity and the Construction of Cortical Circuits. *Science* **274**, 1133–1138 (1996).
11. Morris, R. G. M. D.O. Hebb: The Organization of Behavior, Wiley: New York; 1949. *Brain Research Bulletin* **50**, 437 (1999).
12. Feldman, D. E., Nicoll, R. A. & Malenka, R. C. Synaptic plasticity at thalamocortical synapses in developing rat somatosensory cortex: LTP, LTD, and silent synapses. *Journal of Neurobiology* **41**, 92–101 (1999).
13. Tuncdemir, S. N. *et al.* Early Somatostatin Interneuron Connectivity Mediates the Maturation of Deep Layer Cortical Circuits. *Neuron* **89**, 521–535 (2016).
14. Marques-Smith, A. *et al.* A Transient Translaminar GABAergic Interneuron Circuit Connects Thalamocortical Recipient Layers in Neonatal Somatosensory Cortex. *Neuron* **89**, 536–549 (2016).
15. Picardo, M. A. *et al.* Pioneer GABA cells comprise a subpopulation of hub neurons in the developing hippocampus. *Neuron* **71**, 695–709 (2011).
16. Ghezzi, F. *et al.* GABAergic circuits reflect different requirements for emergent perception in postnatal mouse neocortex. *bioRxiv* 2023.11.21.568139 (2023) doi:10.1101/2023.11.21.568139.
17. Mòdol, L., Moissidis, M. & Marín, O. Somatostatin interneurons control the timing of developmental desynchronization in cortical networks. *bioRxiv* 2023.12.21.572767 (2023) doi:10.1101/2023.12.21.572767.
18. Foeller, E. & Feldman, D. E. Synaptic basis for developmental plasticity in somatosensory cortex. *Current Opinion in Neurobiology* **14**, 89–95 (2004).
19. Pouchelon, G. *et al.* The organization and development of cortical interneuron presynaptic circuits are area specific. *Cell Reports* **37**, (2021).

20. Zhuang, J. *et al.* The spatial structure of feedforward information in mouse primary visual
cortex. *bioRxiv* 2019.12.24.888156 (2020) doi:10.1101/2019.12.24.888156.
21. Ibrahim, L. A. *et al.* Bottom-up inputs are required for establishment of top-down
connectivity onto cortical layer 1 neurogliaform cells. *Neuron* **109**, 3473-3485.e5 (2021).
22. Graziano, A., Liu, X.-B., Murray, K. D. & Jones, E. G. Vesicular glutamate transporters
define two sets of glutamatergic afferents to the somatosensory thalamus and two
thalamocortical projections in the mouse. *Journal of Comparative Neurology* **507**, 1258–
1276 (2008).
23. Narboux-Nême, N. *et al.* Neurotransmitter Release at the Thalamocortical Synapse
Instructs Barrel Formation But Not Axon Patterning in the Somatosensory Cortex. *J.*
Neurosci. **32**, 6183–6196 (2012).
24. Antón-Bolaños, N. *et al.* Prenatal activity from thalamic neurons governs the emergence
of functional cortical maps in mice. *Science* **364**, 987–990 (2019).
25. Pouchelon, G. *et al.* Modality-specific thalamocortical inputs instruct the identity of
postsynaptic L4 neurons. *Nature* **511**, 471–474 (2014).
26. Armbruster, B. N., Li, X., Pausch, M. H., Herlitze, S. & Roth, B. L. Evolving the lock to fit
the key to create a family of G protein-coupled receptors potently activated by an inert
ligand. *Proceedings of the National Academy of Sciences* **104**, 5163–5168 (2007).
27. Urban, D. J. & Roth, B. L. DREADDs (Designer Receptors Exclusively Activated by
Designer Drugs): Chemogenetic Tools with Therapeutic Utility. *Annual Review of*
Pharmacology and Toxicology **55**, 399–417 (2015).
28. Wong, F. K. *et al.* Pyramidal cell regulation of interneuron survival sculpts cortical
networks. *Nature* **557**, 668–673 (2018).
29. Devienne, G. *et al.* Regulation of Perineuronal Nets in the Adult Cortex by the Activity of
the Cortical Network. *J Neurosci* **41**, 5779–5790 (2021).
30. Priya, R. *et al.* Activity Regulates Cell Death within Cortical Interneurons through a
Calcineurin-Dependent Mechanism. *Cell Reports* **22**, 1695–1709 (2018).
31. Karayannis, T., De Marco Garcia, N. & Fishell, G. Functional adaptation of cortical
interneurons to attenuated activity is subtype-specific. *Frontiers in Neural Circuits* **6**,
(2012).
32. De Marco García, N. V., Karayannis, T. & Fishell, G. Neuronal activity is required for the
development of specific cortical interneuron subtypes. *Nature* **472**, 351–355 (2011).
33. Vitali, I. *et al.* Progenitor Hyperpolarization Regulates the Sequential Generation of
Neuronal Subtypes in the Developing Neocortex. *Cell* **174**, 1264-1276.e15 (2018).
34. Martini, F. J., Guillamón-Vivancos, T., Moreno-Juan, V., Valdeolmillos, M. & López-
Bendito, G. Spontaneous activity in developing thalamic and cortical sensory networks.
Neuron **109**, 2519–2534 (2021).
35. Ibrahim, L. A. *et al.* Nova proteins direct synaptic integration of somatostatin interneurons
through activity-dependent alternative splicing. *eLife* **12**, e86842 (2023).
36. Allaway, K. C. *et al.* Genetic and epigenetic coordination of cortical interneuron
development. *Nature* **597**, 693–697 (2021).
37. Mayer, C. *et al.* Developmental diversification of cortical inhibitory interneurons. *Nature*
555, 457–462 (2018).
38. Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902.e21
(2019).
39. Satija, R., Farrell, J. A., Gennert, D., Schier, A. F. & Regev, A. Spatial reconstruction of
single-cell gene expression data. *Nat Biotechnol* **33**, 495–502 (2015).

40. Wu, S. J. *et al.* Cortical somatostatin interneuron subtypes form cell-type-specific circuits.
Neuron (2023) doi:10.1016/j.neuron.2023.05.032.
41. Baude, A. *et al.* The metabotropic glutamate receptor (mGluRIα) is concentrated at
perisynaptic membrane of neuronal subpopulations as detected by immunogold
reaction. *Neuron* **11**, 771–787 (1993).
42. McBain, C. J., DiChiara, T. J. & Kauer, J. A. Activation of metabotropic glutamate
receptors differentially affects two classes of hippocampal interneurons and potentiates
excitatory synaptic transmission. *J. Neurosci.* **14**, 4433–4445 (1994).
43. Lapointe, V. *et al.* Synapse-specific mGluR1-dependent long-term potentiation in
interneurones regulates mouse hippocampal inhibition. *The Journal of Physiology* **555**,
125–135 (2004).
44. Maksymetz, J. *et al.* mGlu1 potentiation enhances prelimbic somatostatin interneuron
activity to rescue schizophrenia-like physiological and cognitive deficits. *Cell Reports* **37**,
109950 (2021).
45. Carly B. Fabian *et al.* Parvalbumin interneuron mGlu$_5$ receptors govern sex differences in
prefrontal cortex physiology and binge drinking. *bioRxiv* 2023.11.20.567903 (2023)
doi:10.1101/2023.11.20.567903.
46. Lüscher, C. & Huber, K. M. Group 1 mGluR-dependent synaptic long-term depression
(mGluR-LTD): mechanisms and implications for circuitry & disease. *Neuron* **65**, 445–459
(2010).
47. Narushima, M. *et al.* The Metabotropic Glutamate Receptor Subtype 1 Mediates
Experience-Dependent Maintenance of Mature Synaptic Connectivity in the Visual
Thalamus. *Neuron* **91**, 1097–1109 (2016).
48. Micheva, K. D. & Beaulieu, C. Quantitative aspects of synaptogenesis in the rat barrel
field cortex with special reference to GABA circuitry. *Journal of Comparative Neurology*
373, 340–354 (1996).
49. Wang, H. & Zhuo, M. Group I metabotropic glutamate receptor-mediated gene
transcription and implications for synaptic plasticity and diseases. *Front Pharmacol* **3**,
189 (2012).
50. Bear, M. F., Huber, K. M. & Warren, S. T. The mGluR theory of fragile X mental
retardation. *Trends in Neurosciences* **27**, 370–377 (2004).
51. Murillo, A. *et al.* Temporal Dynamics and Neuronal Specificity of Grin3a Expression in the
Mouse Forebrain. *Cerebral Cortex* **31**, 1914–1926 (2021).
52. Bossi, S. *et al.* GluN3A excitatory glycine receptors control adult cortical and amygdalar
circuits. *Neuron* **110**, 2438-2454.e8 (2022).
53. Yuan, T. *et al.* Expression of Cocaine-Evoked Synaptic Plasticity by GluN3A-Containing
NMDA Receptors. *Neuron* **80**, 1025–1038 (2013).
54. Bossi, S., Pizzamiglio, L. & Paoletti, P. Excitatory GluN1/GluN3A glycine receptors
(eGlyRs) in brain signaling. *Trends in Neurosciences* **46**, 667–681 (2023).
55. Roberts, A. C. *et al.* Downregulation of NR3A-Containing NMDARs Is Required for
Synapse Maturation and Memory Consolidation. *Neuron* **63**, 342–356 (2009).
56. Benamer, N. *et al.* GluD1, linked to schizophrenia, controls the burst firing of dopamine
neurons. *Mol Psychiatry* **23**, 691–700 (2018).
57. Andrews, P. C. & Dravid, S. M. An emerging map of glutamate delta 1 receptors in the
forebrain. *Neuropharmacology* **192**, 108587 (2021).
58. Fossati, M. *et al.* Trans-Synaptic Signaling through the Glutamate Receptor Delta-1
Mediates Inhibitory Synapse Formation in Cortical Pyramidal Neurons. *NEURON* **104**,
1081-1094.e7 (2019).

59. Uesaka, N. *et al.* Retrograde semaphorin signaling regulates synapse elimination in the
developing mouse brain. *Science* **344**, 1020–1023 (2014).
60. Kreibich, T. A., Chalasani, S. H. & Raper, J. A. The Neurotransmitter Glutamate Reduces
Axonal Responsiveness to Multiple Repellents through the Activation of Metabotropic
Glutamate Receptor 1. *J. Neurosci.* **24**, 7085–7095 (2004).
61. Giger, R. J., Pasterkamp, R. J., Holtmaat, A. J. G. D. & Verhaagen, J. Chapter 12
Semaphorin III: Role in neuronal development and structural plasticity. in *Progress in*
Brain Research (ed. Callow, J. A.) vol. 117 133–149 (Elsevier, 1998).
62. Shelly, M. *et al.* Semaphorin3A Regulates Neuronal Polarization by Suppressing Axon
Formation and Promoting Dendrite Growth. *Neuron* **71**, 433–446 (2011).
63. Wang, N., Dhumale, P., Chiang, J. & Püschel, A. W. The Sema3A receptor Plexin-A1
suppresses supernumerary axons through Rap1 GTPases. *Sci Rep* **8**, 15647 (2018).
64. Lein, E. S. *et al.* Genome-wide atlas of gene expression in the adult mouse brain. *Nature*
445, 168–176 (2007).
65. Letzkus, J. J., Wolff, S. B. E. & Lüthi, A. Disinhibition, a Circuit Mechanism for Associative
Learning and Memory. *Neuron* **88**, 264–276 (2015).
66. Markowitz, J. E. *et al.* The Striatum Organizes 3D Behavior via Moment-to-Moment Action
Selection. *Cell* **174**, 44-58.e17 (2018).
67. Wiltschko, A. B. *et al.* Mapping Sub-Second Structure in Mouse Behavior. *Neuron* **88**,
1121–1135 (2015).
68. Rong, R. *et al.* PI3 kinase enhancer–Homer complex couples mGluRI to PI3 kinase,
preventing neuronal apoptosis. *Nat Neurosci* **6**, 1153–1161 (2003).
69. Yu, C., Derdikman, D., Haidarliu, S. & Ahissar, E. Parallel Thalamic Pathways for
Whisking and Touch Signals in the Rat. *PLOS Biology* **4**, e124 (2006).
70. Sherman, S. M. Tonic and burst firing: dual modes of thalamocortical relay. *Trends in*
Neurosciences **24**, 122–126 (2001).
71. Molnár, Z., Luhmann, H. J. & Kanold, P. O. Transient cortical circuits match spontaneous
and sensory-driven activity during development. *Science* **370**, eabb2153 (2020).
72. Sermet, B. S. *et al.* Pathway-, layer- and cell-type-specific thalamic input to mouse barrel
cortex. *eLife* **8**, e52665 (2019).
73. Audette, N. J., Urban-Ciecko, J., Matsushita, M. & Barth, A. L. POm Thalamocortical Input
Drives Layer-Specific Microcircuits in Somatosensory Cortex. *Cerebral Cortex* **28**, 1312–
1328 (2018).
74. Ährlund-Richter, S. *et al.* A whole-brain atlas of monosynaptic input targeting four different
cell types in the medial prefrontal cortex of the mouse. *Nat Neurosci* **22**, 657–668 (2019).
75. Wall, N. R. *et al.* Brain-Wide Maps of Synaptic Input to Cortical Interneurons. *J. Neurosci.*
36, 4000–4009 (2016).
76. Wildenberg, G., Li, H., Sampathkumar, V., Sorokina, A. & Kasthuri, N. Isochronic
development of cortical synapses in primates and mice. *Nat Commun* **14**, 8018 (2023).
77. Dorskind, J. M. & Kolodkin, A. L. Revisiting and refining roles of neural guidance cues in
circuit assembly. *Current Opinion in Neurobiology* **66**, 10–21 (2021).
78. Erzurumlu, R. S. & Gaspar, P. How the Barrel Cortex Became a Working Model for
Developmental Plasticity: A Historical Perspective. *J Neurosci* **40**, 6460–6473 (2020).
79. Sidorov, M. S., Kaplan, E. S., Osterweil, E. K., Lindemann, L. & Bear, M. F. Metabotropic
glutamate receptor signaling is required for NMDA receptor-dependent ocular
dominance plasticity and LTD in visual cortex. *Proceedings of the National Academy of*
Sciences **112**, 12852–12857 (2015).

80. Narushima, M., Yagasaki, Y., Takeuchi, Y., Aiba, A. & Miyata, M. The metabotropic
glutamate receptor subtype 1 regulates development and maintenance of lemniscal
synaptic connectivity in the somatosensory thalamus. *PLOS ONE* **14**, e0226820 (2019).
81. Penzes, P., Cahill, M. E., Jones, K. A., VanLeeuwen, J.-E. & Woolfrey, K. M. Dendritic
spine pathology in neuropsychiatric disorders. *Nat Neurosci* **14**, 285–293 (2011).
82. Nelson, S. B. & Valakh, V. Excitatory/Inhibitory Balance and Circuit Homeostasis in
Autism Spectrum Disorders. *Neuron* **87**, 684–698 (2015).
83. Hadders-Algra, M. Emerging signs of autism spectrum disorder in infancy: Putative neural
substrate. *Developmental Medicine & Child Neurology* **64**, 1344–1350 (2022).
84. Hensch, T. K. Critical period plasticity in local cortical circuits. *Nat Rev Neurosci* **6**, 877–
888 (2005).
85. Morishita, H., Miwa, J. M., Heintz, N. & Hensch, T. K. Lynx1, a Cholinergic Brake, Limits
Plasticity in Adult Visual Cortex. *Science* **330**, 1238–1240 (2010).
86. Sadahiro, M. *et al.* Activation of Somatostatin Interneurons by Nicotinic Modulator Lypd6
Enhances Plasticity and Functional Recovery in the Adult Mouse Visual Cortex. *J*
Neurosci **40**, 5214–5227 (2020).
87. Kastli, R. *et al.* Developmental divergence of sensory stimulus representation in cortical
interneurons. *Nat Commun* **11**, 5729 (2020).
88. Taniguchi, H. *et al.* A Resource of Cre Driver Lines for Genetic Targeting of GABAergic
Neurons in Cerebral Cortex. *Neuron* **71**, 995–1013 (2011).
89. He, M. *et al.* Strategies and Tools for Combinatorial Targeting of GABAergic Neurons in
Mouse Cerebral Cortex. *Neuron* **91**, 1228–1243 (2016).
90. Daigle, T. L. *et al.* A Suite of Transgenic Driver and Reporter Mouse Lines with Enhanced
Brain-Cell-Type Targeting and Functionality. *Cell* **174**, 465-480.e22 (2018).
91. Platt, R. J. *et al.* CRISPR-Cas9 Knockin Mice for Genome Editing and Cancer Modeling.
Cell **159**, 440–455 (2014).
92. Sousa, V. H., Miyoshi, G., Hjerling-Leffler, J., Karayannis, T. & Fishell, G.
Characterization of Nkx6-2-Derived Neocortical Interneuron Lineages. *Cerebral Cortex*
19, i1–i10 (2009).
93. Hippenmeyer, S. *et al.* A Developmental Switch in the Response of DRG Neurons to ETS
Transcription Factor Signaling. *PLOS Biology* **3**, e159 (2005).
94. Pouchelon, G. *et al.* A versatile viral toolkit for functional discovery in the nervous system.
Cell Reports Methods **2**, 100225 (2022).
95. Marshel, J. H. *et al.* Cortical layer–specific critical dynamics triggering perception. *Science*
365, eaaw5202 (2019).
96. Chan, K. Y. *et al.* Engineered AAVs for efficient noninvasive gene delivery to the central
and peripheral nervous systems. *Nat Neurosci* **20**, 1172–1179 (2017).
97. Fenno, L. E. *et al.* Targeting cells with single vectors using multiple-feature Boolean logic.
Nat Methods **11**, 763–772 (2014).
98. Schneeberger, M. *et al.* Regulation of Energy Expenditure by Brainstem GABA Neurons.
Cell **178**, 672-685.e12 (2019).
99. Vormstein-Schneider, D. *et al.* Viral manipulation of functionally distinct interneurons in
mice, non-human primates and humans. *Nat Neurosci* **23**, 1629–1636 (2020).
100. Labun, K., Montague, T. G., Gagnon, J. A., Thyme, S. B. & Valen, E. CHOPCHOP v2: a
web tool for the next generation of CRISPR genome engineering. *Nucleic Acids*
Research **44**, W272–W276 (2016).
101. Labun, K. *et al.* CHOPCHOP v3: expanding the CRISPR web toolbox beyond genome
editing. *Nucleic Acids Research* **47**, W171–W174 (2019).

102. Favuzzi, E. *et al.* Activity-Dependent Gating of Parvalbumin Interneuron Function by the
Perineuronal Net Protein Brevican. *Neuron* **95**, 639-655.e10 (2017).
103. Virtanen, P. *et al.* SciPy 1.0: fundamental algorithms for scientific computing in Python.
Nat Methods **17**, 261–272 (2020).
104. Collins, D. P., Anastasiades, P. G., Marlin, J. J. & Carter, A. G. Reciprocal Circuits
Linking the Prefrontal Cortex with Dorsal and Ventral Thalamic Nuclei. *Neuron* **98**, 366-
379.e4 (2018).
105. Hay, Y. A., Naudé, J., Faure, P. & Lambolez, B. Target Interneuron Preference in
Thalamocortical Pathways Determines the Temporal Structure of Cortical Responses.
Cerebral Cortex **29**, 2815–2831 (2019).
106. Phillips, J. W. *et al.* A repeated molecular architecture across thalamic pathways. *Nat*
Neurosci **22**, 1925–1935 (2019).
107. Clascá, F., Rubio-Garrido, P. & Jabaudon, D. Unveiling the diversity of thalamocortical
neuron subtypes. *European Journal of Neuroscience* **35**, 1524–1532 (2012).
108. Frangeul, L. *et al.* A cross-modal genetic framework for the development and plasticity of
sensory pathways. *Nature* **538**, 96–98 (2016).
1044

Fig. 1: TC input from VB transiently project onto SST cINs during postnatal development

Fig. 1: TC inputs from VB transiently project onto SST cINs during postnatal development

a, AAV Cre-dependent DIO-ChRmine or -ChR2 controlled under Vipr2-Cre line, targeting VB thalamic neurons, and Flp-dependent fluorescence (fDIO) reporters under the control of SST-FlpO allow for specific recording of SST cIN responses to TC_{VB} inputs. AAVs were injected at P0 and responses were recorded at multiple time point defining 3-time windows: immature stage: P6-P7; early mature stage: P9-11; later mature stage: P28-32. **b**, TC_{VB} projections pattern at P30. Scale bar 200µm. **c**, Average traces of unitary examples of SST cIN responses at P6 and P30 upon light stimulation of TC_{VB} projections. The peak amplitude of EPSCs (excitatory postsynaptic currents) in L5, decreases over development. Kruskal-Wallis test (p-value=0.0029), followed by Dunn's multiple comparison tests (p-value for P6-P10=0.056; P6-P30=0.003; P6: 74±12 n=15 N=4, P10: 35.08±4.67 n=13 N=3; P30 26.92±6.01 n=12 N= 3). EPSC charges in L5 also decrease over development. Kruskal-Wallis test (p-value=0.0004) followed by Dunn's multiple comparison tests (p-values for P6-P30=0.0238; P6: 4.34±1.18, P10: 1.24±0.21, P30: 0.89±0.28). **d**, Staining of VGluT2 (presynaptic) and Homer1 (postsynaptic) TC synaptic contacts onto SST cINs, labeled with SST-cre mouse line and fluorescent reporters. Masks of puncta colocalizations on the soma at P5 and P30 in L5. Scale bar 5 µm. Quantification of TC synaptic contact density onto SST cINs in L5. During development, the number of contacts onto SST soma is decreased. Kruskal-Wallis (p-value <0.0001), followed by Dunn's multiple comparison test (p-values for P5-P7/P5-P30/P5/P10<0.0001, P7-P30=0.290; P7-P10=0.151; P10-P30>0.999; P5: 0.282±0.01 n=89 N=3, P7: 0.141±0.009 n=87 N=3, P10: 0.115±0.011 n=86 N=2; P30: 0.105±0.009 n=54 N=3).

Fig. 2: Postsynaptic activity of SST cINs regulates the maturation of transient TC inputs.

Fig. 2: Postsynaptic activity of SST cINs regulates the maturation of transient TC inputs

a, Designer Receptors Exclusively Activated by Designer Drugs (DREADD)-based chemogenetic tools used to test activity-dependent maturation of TC inputs onto SST cINs. DREADD coupled with Gi inhibits cINs, while DREADD coupled with Gq activates neurons upon delivery of the designer drug Clozapine-N-Oxide (CNO). 12-24hrs following local AAV injections in the cortex and in the thalamus at postnatal day 0 (P0), DREADD-Gi or -Gq are chronically activated during TC connection to SST cIN maturation, with intraperitoneal (i.p.) injection of CNO twice a day until P8. Animals are sacrificed for recording or synaptic analysis after maturation occurs during early and later mature windows (P9-11 and P28-32). **b**, AAV-driven Cre-dependent ChR2 expression in VB is controlled by Vipr2-Cre mouse line at P10. **c**, Example response traces from control SST cINs and SST cINs chronically-inhibited with Gi or -activated with Gq. **d**, During the early mature window (P9-11), EPSC peak amplitudes and charges are significantly decreased after Gq expression compared to Control and Gi-expressing SST cINs. Amplitudes: Kruskal-Wallis test (p-value=0.0066) followed by Dunn's multiple comparison test (p-values for Ctrl-Gq=0.0038, Gi-Gq=0.0222; Ctrl: 35.08±4.67 n=13 N=3 includes Ctrl from Fig.1, Gi: 31.22±5.17 n=10 N=3, Gq: 12.68±2.021 n=8 N=3). Charges: Kruskal-Wallis test (p-value=0.0002) followed by Dunn's multiple comparison test (Ctrl 1.24±0.21, Gi 1.68±0.69, Gq 0.08±0.02). **e**, VGluT2 (presynaptic) and Homer1 (postsynaptic) staining of TC synaptic contacts onto SST cINs and masks of colocalization measured on the soma of Gi- and Gq-expressing SST cINs in L5. Scale bar: 10 µm. Quantification of TC synaptic contact density onto SST cINs in L5 shows a decrease of the number of synaptic contacts onto SST cINs expressing Gq compared to SST cINs expressing Gi and to P10 Ctrl cells from Fig.1d. Kruskal-Wallis test (p-value <0.0001) followed by Dunn's multiple comparison tests (p-values for Ctrl-Gq=0.0078, Ctrl-Gi=0.4542, Gi-Gq<0.0001; Ctrl: 0.116±0.0.011 n=86 N=2, Gq: 0.124±0.015 n=56 N=2, Gi: 0.123±0.009 n=105 N=3, Gq 0.074±0.007 n=94 N=3).

Fig. 3: mGluR1 is highly expressed in SST cINs during development

Fig. 3: mGluR1 is highly expressed in SST cINs during development

a, UMAP representation of the integration of P2 and P10 cortical GABAergic neurons scRNA-seq public databases[36,37] (representing gene expression during the immature and early mature windows), allowing for the identification of SST cINs at both stages. Left UMAP is color-coded by time-point, right UMAP is color-coded by major cIN classes based on the expression markers as previously described.[37,40] **b**, Differential gene expression between PV and SST cIN clusters using Seurat non-parametric Wilcoxon rank sum test. Volcano plot representing the average of the Fold Change (FC) and adjusted p-value of genes specifically expressed in PV and SST cIN clusters. During immature stage (P2), Grm1 coding for mGluR1 is in one of the 3 highest significant genes in SST cINs and the top 3rd differentially expressed gene (FC=4.53, adj. p-value=9E-264). While still significant during the early mature stage (P10) (adj. p-value=7.43E-20), the FC is only x1.64. **c,d**, UMAP representation of Sst and Grm1 expression levels onto scRNAseq immature (P2) and mature (P10) datasets, showing the reverse temporal dynamics in expression levels. **e**, Violin plots representing the level of Grm1 expression in cells from SST and PV cIN clusters at both immature and mature stages (P2/P10). **f**, smFISH of Grm1 combined with Sst immunostaining at P5 and P10 shows the high colocalization of Grm1 RNA molecules in SST cINs. Scale bar: 10 μm. **g**, Immunofluorescence of mGluR1 in S1 cortex. Expression is high throughout development including both immature an mature time windows, and is associated with a soma-to-neurites redistribution of the receptor. Left scale bars: 100 μm; Right scale bar: 20 μm

Fig. 4: mGluR1 CRISPR-deleted SST cINs retain their strong developmental TC inputs

Fig. 4: mGluR1 CRISPR-deleted SST cINs retain their strong developmental TC inputs

a, mGluR1 is a postsynaptic metabotropic receptor coupled to Gq protein activated by the presynaptic release of glutamate, similarly to DREADD-Gq. Strategy for early deletion of mGluR1 (KD) in SST cINs specifically, using CRISPR/Cas9 system: AAV-driven Flp-dependent Cre expression is driven by SST-FlpO mouse line and controls LSL-Cas9eGFP mouse line. 3x CRISPR single guide RNAs targeting mGluR1 are driven by an AAV expressing a Cre-dependent reporter. AAVs were injected in S1 at P0. **b**, Top: example of Grm1 expression in control (Ctrl) I and KD SST cINs using smFISH of Grm1 combined with Sst immunostaining at P10. Scale bar: 10 µm. Bottom: Quantification of Grm1 RNA molecules per SST cINs using smFISH, showing major downregulation of Grm1 expression in the mGluR1 CRISPR KD at P10. Student t-test at P10 (p-value<0.0001, Ctrl: 30.89±2.17 n=27 N=2, KD: 10.35±2.35 n=23 N=2). **c**, AAV-driven Cre-dependent ChR2 expression in VB is controlled by Vipr2-Cre mouse line injected at P0. SST cINs were recorded in L5 during early and late mature windows (P9-11 and P29-32). **d**, Averaged traces from unitary example of labeled Ctrl and KD SST cINs in response to VB fiber stimulation at early mature stage. EPSCs peak amplitudes of SST cINs are significantly increased in SST cINs deleted for mGluR1 compared to Ctrl cINs at early mature stage. Mann-Whitney test (p-value=0.0112; Ctrl from Fig. 1: 35.08±4.67 n=13 N=3; KD: 185.6±66.4 n=15 N=4). The variance (Var.) of the responses is also significantly increased for KD SST cIN compared to Ctrl cells. Bartlett's test (p-value=4.43E-12). EPSC charges of KD SST cINs, as well as their variance, are significantly increased compared to Ctrl SST cINs. Mann-Whitney test (p-value=0.0004; Ctrl: 1.24±0.21, KD: 9.20±2.84). Bartlett's test (p-value=7.76E-12). **e**, TC synapse density is increased onto mGluR1 KD SST cINs in L5 compared to Ctrl cells. Example staining of VGluT2 (presynaptic) and Homer1 (postsynaptic) labeling of TC synaptic contacts onto SST cINs at P30. Masks of puncta colocalization. Scale bar: 5 µm. Quantification of TC synapses. Mann-Whitney test (p-value=0.0014; Ctrl 0.077±0.004 n=152 N=4; mGluR1 KD 0.101±0.005 n=140 N=4).

Fig. 5: mGluR1-dependent transcriptional regulation in SST cINs regulates transient TC connectivity

Fig. 5: mGluR1-dependent transcriptional regulation in SST cINs regulates transient TC connectivity

a, Early postnatal expression levels of the candidate genes Grin3a (coding for NR3A), Grid1 (coding for GluD1), Sema3A and Sema7A in SST cINs using P2/P10 integrated scRNAseq. Violin plots represent the expression of these genes in SST and PV cIN clusters (inset) at P2 and P10, compared to Sst and Grm1. **b**, Strategy for investigation of candidate gene expression in SST cINs in absence of mGluR1. CRISPR/Cas9 strategy for mGluR1 deletion in SST cINs with AAV injection at P0. smFISH analysis at P10. **c**, Example of smFISH in Ctrl and mGluR1 KD SST cINs. Colocalization of Sema3A RNA using smFISH in SST cINs as labeled with Sst immunostaining at P10 in Control and mGluR1 KD. Scale bar: 10 µm. **d**, Quantification of RNA molecule number per SST cIN of the select candidate genes using smFISH. The number of Grin3a, Grid1, Sema3A and Sema7A transcripts is decreased in SST cINs depleted for mGluR1 compared to Ctrl SST cINs. Student t-test for Grin3a: p-value=0.0092 (Ctrl: 31.13±1.65 n=86 N=2, KD: 25.46±1.37 n=84 N=2); Mann-Whitney test for Grid1: p-value=0.0389 (Ctrl: 7.30±0.77 n=44 N=2, KD: 9.26±0.74 n=54 N=2), for Sema3A: p-value=0.0041 (Ctrl: 16.64±1.21 n=148, N=4; KD 12.04±1.05 n=140, N=4), for Sema7A: p-value=0.0119 (Ctrl 12.04±1.17 n=138 N=4, KD 8.14±0.76 n=145 N=4). **e**, Fold change (mean of molecule number per cell in each animal (N)). **f**, Strategy for early deletion of Sema3A in SST cINs specifically, using CRISPR/cas9 system: AAV-driven Flp-dependent Cre expression is driven by SST-FlpO mouse line and controls LSL-Cas9eGFP mouse line. 3x CRISPR single guide RNAs targeting Sema3A are driven by an AAV expressing a Cre-dependent reporter. **g**, Recordings of labeled Ctrl and Sema3A CRISPR KD SST cINs upon VB projection stimulation in L5 during early mature window (P9-11). **h**, Averaged traces of example response from Ctrl and Sema3A KD SST cINs labeled in response to VB fiber stimulation at P10. While EPSC peak amplitudes and charges of SST cINs show only a mild increase in Sema3A KD SST cINs compared to Ctrl at early maturation stage, their variance is significantly increase, similar to mGluR1 KD SST cINs. Mann-Whitney test (Amplitude p-value=0.077; Ctrl from Fig. 1: 35.08±4.67 n=13 N=3; Sema3A KD: 66.75±12.78 n=12 N=3; Charge p-value=0.068; Ctrl: 1.24±0.21; Sema3A KD: 3.37±0.87); Bartlett's test (Amplitude: p-value=0.0023; Charge: p-value=3.15E-05).

Fig. 6: The persistence of developmental TC inputs alters exploratory behavior at adulthood

Fig. 6: The persistence of developmental TC inputs disrupts exploratory behavior at adulthood

a, Strategy for early deletion of mGluR1 in SST cINs specifically, using CRISPR/Cas9 system: AAV-driven Flp-dependent Cre expression is driven by SST-FlpO mouse line and controls LSL-Cas9eGFP mouse line. 3x CRISPR single guide RNAs targeting mGluR1 are driven by an AAV expressing a Cre-dependent reporter. AAVs were injected bilaterally in S1 at P0. Animals were recorded around P60 for MoSeq. **b**, Motion Sequencing pipeline. Examples of behavioral motifs, so-called syllables identified with MoSeq. **c**, Position occupancy heatmap by sex. The top two rows are the min-max scaling of the occupancy percentages, where 1 is the maximum value and 0 is the minimum value. The bottom row is the difference between occupancy percentages for mGluR1 KD mice and the Control mice (∆ index). **d.i,** Syllables for Ctrl and mGluR1 KD male mice, sorted by syllable usage differences between mGluR1 KD and Ctrl. The syllable usage distributions within the two groups are tested with a Chi-square test on the unnormalized syllable usages (p-value<0.01) and the two distributions are significantly different. The prominent syllables (see methods) are labeled with red stars (27 syllables for male, and 27 syllables for female). The syllable behavior types are color coded (i.e., light gray for dart, green for groom, red for rear, etc.). The inset subpanel above is aggregated usages by syllable classes and the differences between two groups are also tested with a Chi-square test on the unnormalized syllable usages (p-value<0.01). **d.ii**, Same as d but for female mice. **e**, Linear Discriminant Analysis (LDA) plot shows that Ctrl and mGluR1 KD mice are separated by the syllable ensemble (males and females combined). The top 10 syllables with the highest absolute model weights consist of 7 rears, 2 grooms and 1 scrunch. **f**, Confusion matrices for classification accuracy of a linear classifier trained on the 19 prominent syllables identified from 76 LDA iterations by sex (see methods). The use of these syllables alone could predict Ctrl or KD mice. The classifier yields to 0.69 and 0.84 accuracy across cross-validation for females and males, respectively (top/bottom). Labels per-mouse are shuffled for control (right).

Fig. 7: Model for transient TC connectivity development

Fig. 7: Model for transient TC connectivity development
In contrast to other neuron types, to which TC connectivity develops according to Hebbian mechanisms, SST cINs receive transient TC inputs from the VB thalamus. These TC inputs regress onto SST cINs following a non-Hebbian mechanism and involves postsynaptic metabotropic-dependent transcriptional regulation. This process is required for the maturation of exploratory behaviors in adult mice.