New observations of fluorescent organisms in the Banda Sea and in the Red Sea

Lars Henrik Poding [1] *#,

Budiono Senen [2], Gino Valentino Limmon [3][4],

Stefan Herlitze [1] +, Mareike Huhn [1]

[1] Department of General Zoology and Neurobiology, Institute of Biology and Biotechnology,

Ruhr-University Bochum, 44801 Bochum, Germany

[2] Universitas Banda Neira, Maluku Tengah, Indonesia

[3] Department of Aquatic Resources Management, Faculty of Fisheries and Marine Science,

Pattimura University Ambon, Indonesia

[4] Maritime and Marine Science Center for Excellence of Pattimura University Ambon,

Indonesia

* first author

corresponding author: lars.poding@rub.de

+ corresponding author: sxh106@gmail.com

Abstract

Fluorescence is a widespread phenomenon found in animals, bacteria, fungi, and plants. In

marine environments fluorescence has been proposed to play a role in physiological and

behavioral responses. Fluorescence has been described in jellyfish, corals and fish, and many

fluorescent proteins and molecules have been identified. Here we describe new fluorescent

species, which we found during night dives in the Banda Sea, Indonesia, and in the Red Sea,

Egypt. Among various phyla we found fluorescence in sponges, molluscs, tunicates, and fish

and, with this, extend the knowledge on the number of fluorescent species in marine

environments.

Keywords: Fluorescence, marine organsims, coral reefs, Banda Islands, Red Sea

Introduction

Fluorescence is the property of a molecule to absorb light of a certain wavelength followed by the emission of light with longer wavelength. In bacteria, plants, fungi and animals, various fluorescent molecules have been described such as chitin, minerals, carotenoids, flavonoids, porphyrins, chlorophyll, phycobiliproteins or green fluorescent protein (GFP)-like fluorescent proteins. These molecules can be excited by either sun- or moonlight [1,2].

In marine environments many fluorescent species have been described and for various of these species an ecological and/or physiological function has been suggested. Only a few studies addressed the behavioral and physiological role of fluorescence [2]. In jellyfish, for example, where the first fluorescent protein (avGFP) was characterized, GFP is expressed as an acceptor molecule for bioluminescent light in its tentacles [3]. The green fluorescence is used to attract prey [4]. So far fluorescent proteins have been found in Cnidaria, Vertebrata, Cephalochordata (lancelets) and Arthropoda, but have not been described in Porifera (sponges), Ctenophora (comb jellies), Tunicata (tunicates), Hemichordata (hemichordates), Echinodermata (echinoderms), Mollusca (molluscs), Annelida (segmented worms) and Nematoda (roundworms) [5–7]. Most fluorescent proteins with blue to red emission spectra have been identified in Anthozoa (corals). Here the fluorescence has been shown to be involved in attraction of photosynthetic symbionts, in photoprotection of the symbionts and in converting blue light to longer wavelength light in mesophotic environments (photoenhancement) [2]. Note that the green to red components of the sunlight are selectively removed with water depth resulting in a blue light environment [8]. In fish red fluorescent body coloration and red fluorescent eye rings have been shown to be involved in intra- and interspecific communication [9–13]. In addition, differences in the fluorescence pattern between males and females have been suggested to play a role in sexual communication [10,14].

In contrast to jellyfish, corals, sea anemones, deep-sea benthic copepods, lancelets, eels and

moray eels, where fluorescence is based on the expression of fluorescent proteins, non-FP

based fluorescence has been observed in Porifera (sponges), Heterobranchia (slugs and

snails), Annelida (segmented worms), Crustacea (copepods), Osteichthyes (bony fish) and

Chondrichthyes (cartilaginous fishes) [8,15–19]. For most species, excluding jellyfish, corals

and fish, the physiological function of fluorescence and the fluorescent molecules have not

been identified.

Here we describe several new fluorescent species, which we found during night dives in the

Banda Sea, Banda Islands, Indonesia and the Red Sea, Dahab, Egypt. These new fluorescent

species include nudibranchs, tunicates, sponges, various other invertebrates, and fish. In

addition to the new species, we show detailed images of known fluorescent species to give

precise information about the distribution of body fluorescence.

Methods

Photo acquisition in the field

Images of new fluorescent species were acquired during SCUBA diving in the Red Sea in Dahab, Egypt (28°29'20.0"N 34°30'57.2"E) and in the Banda Sea at the Banda Islands, Maluku, Indonesia (4°30'55.62"S 129°53'35.92"E). All dives were started after sunset and limited to a maximum depth of 15 meters, which was monitored by a dive computer. Field observations were carried out in May 2019 (Red Sea) and in February-March 2019 and September 2022 (Indo Pacific).

Fluorescence was excited with "Sola" light source (NightSea, USA; λ max = 449 nm), Flashlight (D2000; INON; Japan), Ex-Inon excitation filter (NightSea, USA; λ max = 454 nm) or "Blue Star" light source (NightSea, USA; λ max = 470 nm). The transmission filter (BF 2, NightSea, USA) in front of the underwater housing showed a low cut at λ < 500 nm. Fluorescence photography was performed with Powershot G15 (Canon, Japan) placed in WP DC 48 (Canon, Japan) underwater housing or Sony α 6000 (Sony, Japan) placed in SF-A6500 (Seafrogs, Hongkong) underwater housing. All images were processed with Lightroom (Version 2015.1.1, Adobe, USA) and Corel Draw (Version 20.0.0.633, Corel Corporation, Canada), by correcting the brightness, contrast, removing haze and cropping the images.

Species were identified with standard literature [20–29]. The species were identified by comparison of digital underwater pictures with the description and pictures of the literature. Since we did not take DNA or tissue samples, the species information must be interpreted with caution for the less distinct taxa.

Photo acquisition by Leica Thunder

Detailed images of *Anampses meleagrides* [Randall, 1972]*, Cirrhilabrus aquamarines* [Tea, 2021]*, Cirrhilabrus rubrisquamis* [Randall & Emery, 1983]*, Doryrhamphus excisus* [Kaup, 1856]*, Eviota atriventris* [Greenfield & Suzuki 2012], *Eviota nigriventris* [Giltay 1933]*,

Lybia tessellata [Latreille in Milbert, 1812]*, Nembrotha kubaryana* [Bergh, 1877]*,*

Odontodactylus scyllarus [Linnaeus, 1758]*,* one species of Polyplacophora*,*

Synchiropus sycorax [Tea & Gill, 2016], were acquired with the fluorescence stereoscope

Leica M205 FCA equipped with a DFC90000 GT Camera and a GFP (λ_{Ex}: 395–455 nm, λ_{Em}:

480 LP), mCherry (λ_{Ex}: 540-580 nm, λ_{Em}: 593-667 nm) and a CY5 (λ_{Ex}: 590–650 nm, λ_{Em}:

663–737 nm) filter cube. The fish and invertebrates were purchased from the wholesale trader

DeJong Marinelife (Netherlands) in 2021/2022 or from Korallenfarm Joe & Co (Germany). All

images were processed with LASX (Version 3.7.6.25997, Leica, Germay), Lightroom (Version

2015.1.1, Adobe, USA) and Corel Draw (Version 20.0.0.633, Corel Corporation, Canada) by

correcting and/or cropping the images.

Results

During night dives at the Banda Islands, Indonesia and Dahab, Egypt new fluorescent species

belonging to the phyla Porifera, Mollusca, Arthropoda, Annelida and Chordata could be

identified.

Porifera - We found green, yellow and orange fluorescence in different unidentified sponges in

Egypt (Fig. 1 C) and in Indonesia (Fig. 1 D). *Gelliodes fibulata* [Ridley, 1884] revealed green

fluorescent spots on its surface (Fig. 1 A, B).

Corals – We found a variety of shallow-water stony corals that exhibited yellow fluorescence

(Fig. 2), namely *Physogyra lichensteini* [Milne Edwards & Haime, 1851], three different species

of *Montipora* and two *Fungia* species.

Molluscs - In molluscs we found fluorescence in the families Octopodidae, Strombidae,

Cardiidae, Facelinidae (Fig. 3), Polyceridae (Fig. 4) and unidentified families of Polyplacophora

(Fig. 5). The octopus *Abdopus aculeatus* [Férussac & d'Orbigny], 1834] revealed orange

fluorescent areas on its body. Bright fluorescent patches were located below the yellow

fluorescent eyes and on every arm (Fig. 3 A & B). Other species (unspecified) of the same

family showed no fluorescence, but had yellow fluorescent eyes, most likely representing

yellow intraocular filters (Fig. 3 B, D, E). We found green and red fluorescence in several snails.

The bubble conch *Euprotomus bulla* [Röding, 1798] had a bright green fluorescent operculum

and weak green fluorescent eyes. The giant clam *Tridacna crocea* [Lamarck, 1819] showed a

very bright red fluorescence that could be related to photosynthetic symbiotic algae (Fig. 3 G).

In addition, the nudibranch *Facelina rhodopos* [Yonow, 2000] (Fig. 3 H) showed bright orange

fluorescence at the tip of the cerata, which most likely was related to ingested metabolites of

the prey-hydrozoan *Millepora sp.*. Another nudibranch *Nembrotha kubaryana* that we found in

the Banda Sea in its natural habitat (Fig. 4 A), and had purchased from Dejong Marinelife

(Netherlands) revealed strong red fluorescence (Fig. 4 B). We were able to localize the

fluorescence in more detail under laboratory settings and found fluorescent patches around

the mouth (Fig. 4 D), the rhinophores (Fig. 4 E - G), gills (Fig. 4 B) and the dorsal rim (Fig. 4 B

& H). We also found a variety of different Polyplacophora that exhibited bright green, yellow

and red fluorescence (Fig. 5 A - D).

Echinodermata – We found bright green, yellow and red fluorescent crinoid species of different

genera at the Banda Islands and in the Red Sea (Figure 6).

Arthropoda - Several fluorescent decapoda of the family Inachidae, Scyllaridae and

Palaemonidae were found. The decorator crab *Camposcia retusa* [Latreille, 1829] (Fig. 7 A

Banda Sea) showed a bright green fluorescent carapace. In the Red Sea we discovered a

bright green fluorescing slipper lobster *Scyllarides sp*. (Fig. 7 B). In addition, we found bright

green fluorescence in the carapace of Coleman`s Shrimp*, Periclimenes colemani* [Bruce,

1975] (Fig. 7 C), living in symbiosis with fire urchins that do not fluoresce. Green and red

fluorescence of the carapace was discovered in peacock mantis shrimps

Odontodactylus scyllarus. The images revealed fluorescence in filamentous structures on

pleopods (Fig. 8 C), uropods (Fig. 8 E), paraeopods (Fig. 8 D), and on maxillipeds (Fig. 8 A,

B, F). Furthermore, we identified fluorescent body parts of the Mosaic Boxer Crab

(*Lybia tessellata*) (Fig. 8 G)

Annelida - We also found fluorescence in two types of bristle worms (polychaetes) from the

family Eunicidae and Sabellidae. In the subclass Eunicida only the bristles fluoresced bright

green (Fig. 7 D, F). In addition, we found bright yellow fluorescence in a feather duster worm

(Sabellidae) (Fig. 7 E).

Chordata – Two fluorescent tunicate species and three new species of fluorescent bony fish

were photographed in the Banda Sea, Indonesia. Both tunicates, the ascidians

Clavelina coerulea [Oka, 1934] (Fig. 9 A & B) and *Clavelina robusta* [Kott, 1990] (Fig.9 C & D)

revealed a bright green rim-like fluorescence around their siphons. In both species the

fluorescent rims appeared yellow under white light conditions. We found two green and orange

fluorescent scarlet frogfish *Antennatus coccineus* [Lesson, 1831] (Fig. 6 A & B). The green

fluorescence was distributed over the entire body, while the orange fluorescence was found in

patches and in the lures. In addition, we found a bright fluorescing green leopard flounder

Bothus pantherinus [Rüppell, 1830] (Fig. 10 C). Green fluorescence was more prominent in

the white banded parts of the skin. We also found individual differences in fluorescence in the

banded pipefish *Corythoichthys intestinalis* [Ramsay, 1881]. Individuals revealed either green

or orange fluorescent skin patterns (Fig. 10 D-F). However, all individuals revealed a bright

yellow/orange fluorescence around the eyes and the caudal fin (Fig. 10 D-F). Moreover, we

observed two individuals of *Pleurosicya mossambica* [Smith, 1959] in the Banda Sea with

bright orange-red fluorescence around the eyes and the chorda (Fig. 11 A & B) and two

individuals of *Scorpaenopsis possi* [Randall & Eschmeyer, 2002] with orange and red

fluorescent patches in their tissue (Fig. 11 C & D). In addition, we found two different color

morphs of *Soleichthys heterorhinos* [Bleeker, 1856] from the Banda Sea (E) and Red Sea (F)

and a green fluorescent undefined species of Lutjanidae (G), and *Brachysomophis henshawi*

[Jordan & Snyder, 1904] with a red fluorescent head (H). Detailed microscope images of green

and red fluorescent fish are provided in Figure 12 and 13.

Discussion

During night dives in the Banda Sea, Indonesia and the Red Sea, Egypt we photographed marine organisms, in which, to our knowledge, fluorescence had not been described before. These include sponges, crustaceans, polychaetes, slugs, snails, octopus, ascidians, and fish.

Fluorescence has been described in various marine organisms and many behavioural and physiological functions have been suggested. Only a few studies were conducted, which show a correlation between fluorescence and visual guided behaviour or physiological responses [2]. We will, therefore, relate these functions to the new fluorescent species we found:

Fluorescence in sponges has been described in a few species, and it was suggested that fluorescence might be used for photoenhancement, i.e. the conversion of blue light into longer wavelength light. Photoenhancement has been discussed in mesophotic environments. Since yellow green fluorescent sponges can be found at depths of up to 40 m [16], and fluorescence is expressed in archaeocytes [15], which ingest and digest food, the fluorescence might be used to attract prey. Wether the green fluorescent spots we observed for *Gelliodes fibulata* are intrinsic to the sponge or are secondary to some symbionts or parasites needs to be investigated.

To date, fluorescence has been described and characterized mainly in corals and many FPs have been identified. Yellow to red fluorescent corals are usually found in the mesophotic coral ecosystem, where fluorescence most likely plays a role in photoenhancement [30,31]. It was therefore surprising that we found various yellow fluorescent corals in the shallow reefs (<10m) in Egypt and Indonesia.

Several crustaceans have been described to be fluorescent [32–35], where fluorescence was shown to be produced by autofluorescence. This could also be the case for the green fluorescing cuticula of the slipper lobster (Fig. 4 B). Symbionts (fluorescent microalgae), or

fluorescent molecules such as chitin, lipofuscin, or tryptophan associated molecules could

cause the fluorescence here [2]. For isopods that live in symbiosis with red fluorescent

microbes, it has been suggested that the red fluorescence might serve as a warning signal for

predators [19]. Pontellidae, a family of copepods show sexual dimorphism in eye design with

different fluorescent patterns [33,35,36]. It has also been suggested that the different patterns

might be involved in intraspecific recognition. In addition, countershading has been suggested

for blue and green fluorescence [35]. This could also be the case for the decorator crab (Fig.

4 A) and the Coleman`s Shrimp (Fig. 4 C). Whether fluorescence is also used for

communication in the newly discovered fluorescent species, similar to the mantis shrimp [37],

or it is an unfunctional epiphenomenon [38], still needs to be investigated.

Fluorescence has also been described in various segmented worms. The intertidal worm

Eulalia sp. (Polychaeta) secretes a blue-green fluorescent mucus, which has been suggested

to protect the worm against high-energy radiation. The gossamer worm (*Tomopteri*s *spp., a*

pelagic annelid) uses fluorescence to enhance bioluminescent light [39]. In the tubeworm

Chaetopterus variopedatus the mucus of the worm contains blue and green fluorescence

mediated by riboflavin and related derivatives [2,40,41]. In addition, blue fluorescence and

bioluminescence has also been observed in the fireworm *Odontosyllis phosphorea* [42,43].

While the fluorescent molecules are most likely used to enhance light spectra within the

luminescent worms, a physiological/behavioral function has not been demonstrated [2]. We

also found fluorescence in fire or bristle worms from the family Eunicidae and Sabellidae with

so far unknown function (Fig. 4 D-F).

Fluorescence is known from molluscs [17] . The giant clams *Tridacna* (Fig. 2 G) and *Hippopus*

live in symbiosis with red fluorescent photosynthetic algae, which are hosted in the mantle

tissue. The fluorescence seen for *Tridacna* is therefore most likely associated to the chlorophyll

photosystem II of the algae [44–46]. In snails fluorescence is often found in the cerata, which

originate either from ingested food (green fluorescence) or symbiotic algae (red) [17]. It could

be possible, that the Nudibranch *Facelina rhodopos* deposits ingested fluorescent food in their

cerata since this snail feeds on the hydrozoan *Millepora,* that has a weak fluorescence.

Behavioural experiments with *Goniobranchus splendidus* revealed that predators learn to

avoid feeding on the unpalatable snail. The avoidance behaviour of the fish is triggered by the

yellow rim of the snail [47]. These experiments suggest that the red fluorescent rim of

Nebrotha kubaryana may be an aposematic signal.

In Cephalopoda fluorescence had not been described so far. It is therefore surprising to find

green and red fluorescence in the octopus *Abdopus aculeatus*. Based on the distribution and

pattern of the fluorescence one could speculate that fluorescence is involved in intraspecific

communication.

Another surprising result was the occurrence of fluorescence in two species of ascidians. In

both species bright green fluorescence was seen in a rim-like structure around the oral and

atrial siphon (Fig. 5). Since food is ingested via the oral siphon, fluorescence might be involved

in attraction of plankton which was demonstrated for the green fluorescence of the jellyfish

Olindias formosa [4]. A similar function of fluorescence might also play a role in the green to

red fluorescence of echinoderms.

Green and red fluorescence is common in fish. In crypto-benthic fish, such as the Scarlet

frogfish and the Leopard flounder (Figure 10) fluorescence may facilitate background matching

(camouflage) [8,48]. Red fluorescent body coloration in the red-eyed wrasse

Cirrhilabrus solorensis has been shown to increase aggressive behavior by increased

recognition of fluorescent possible rivals and might also be involved in sexual communication

based on the difference in the fluorescence pattern between males and females [9,10,14]. Red

fluorescent eye rings in the black-faced blenny *Tripterygion delaisi* are most likely used for

intraspecific communication since the fish can perceive and respond to red fluorescence [11–

13]. Thus, the red fluorescent eye rings we observed in the pipefish may play a similar role for

short distance communication.

This study contributes twenty one new species to the database of known fluorescent marine

organisms. Three of these species belonged to the phylum Porifera, seven to the phylum

Mollusca, three to the subphylum Crustacea, three to the phylum of Annelida, two to the class

Ascidiacea and three to the subphylum Vertebrata. We describe for the first-time fluorescence

in Octopoda and Ascidiacea, and also show the first case of fluorescence in Nudibranchia,

where the fluorescence is not linked to ingested food. This study, therefore, extends the palette

of fluorescent marine species.

Acknowledgements

We thank all RUB students who contributed to this article by taking underwater pictures. Especially Dr. Peter Jägers for supporting in field and proofreading of the manuscript, Lukas Menzen, Andre Marcel Haase, Leonard Klaus, Denise Wagner and Jan-Niklas Schnalke for taking underwater pictures in Banda and in the Red Sea. Thanks to the dive instructors and dive guides who showed us fluorescence in coral reefs during night dives.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by DFG He2471/18-2 Priority Program (SPP1926) and funds of the Ruhr-University Bochum

Availability of data and material

All data generated or analysed during this study are included in this published article [and its supplementary information files].

Authors' contributions

Conceptualization LHP, MH, SH

Investigation LHP, BS, GL

Writing – Original LHP, BS, GL, MH, SH

Writing – Review & Editing LHP, MH, SH

Figures

Figure 1: Fluorescence in Porifera in the Banda Sea: The thorny stem sponge *Gelliodes fibulata* ((A & B) white light in A, fluorescence in B) show green fluorescent spots over the whole tissue. Two unidentified sponges (C, D) fluoresce yellow and green.

Figure 2: Fluorescence in corals: Yellow fluorescent corals in the Banda Sea and in the Red Sea. *Physogyra lichensteinii* (A), *Montipora sp.* (B - D) and *Cycloseris* sp./*Fungia sp.* (E & F). A, B, D - F: Banda Sea, C: Red Sea.

Figure 3: Fluorescence in Mollusca: Three different types of fluorescent Octopods; *Abdopus aculeatus* (A & B) with orange fluorescent parts of skin, two unidentified species of Octopodidae (C – E) with fluorescent eyes, green fluorescent operculum of *Euprotomus bulla* (F), red fluorescent mantle of *Tridacna crocea* (G), orange fluorescent *Facelina rhodopus* (H) on *Millepora sp.*. A – G: Banda Sea, H: Red Sea

Figure 4: Fluorescence in the nudibranch *Nebrotha kubaryana:* (A) natural habitat (Banda Sea), white light & Fluorescence (B), whole body (B, C), oral tentacles (D), rhinophores (E-G), gill (H) fluorescence. Pictures C – H created with a Thunder microscope (Leica, Germany) with CY5 filter.

Figure 5: Polyplacophora: Red, yellow and green fluorescent Polyplacophora in the Red Sea (A-C) and in the Banda Sea (D). Bright green and red fluorescent Polyplacophora in a reef tank in E-F.

Figure 6: Fluorescence in Echinoderms: New observations of red, yellow and green fluorescence in crinoids from the Banda Sea (A-E) and in the Red Sea (F-H).

Figure 7: Fluorescence in Arthropoda and Annelida: Green fluorescent *Camposcia retusa* (A)*,*

Scyllarides sp. (B), *Periclimenes colemani* (C), Eunicida (D,F)*,* Yellow fluorescent chaeta of an

unidentified species of Sabellidae (E). A, C, D, F: Banda Sea, B & E: Red Sea

Figure 8: Fluorescence of *Odontodactylus scyllarus* and *Lybia tessellata:* Green (GFP filter)

and red fluorescing (Cy5 filter) structures of maxillipeds (A, B, F), pleopods (C), uropods (E)

and paraeopods (D). Green and red fluorescing *Lybia tessellata* (G; GFP & CY5 filter). Pictures

were taken on a Thunder microscope (Leica, Germany).

Figure 9: Fluorescence in Tunicata in the Banda Sea: White light and fluorescent picture of

Clavelina coerulea (A, B)*, Clavelina robusta* (C,D). The green fluorescence concentrates on

the siphons of the ascidians.

Figure 10: Fluorescence in fish in the Banda Sea: Green and orange fluorescing

Antennatus coccineus (A,B), green fluorescing *Bothus pantherinus* (C) and different

fluorescing individuals of *Corythoichthys intestinalis* (D-F).

Figure 11 Fluorescence in fish: Red fluorescing *Pleurosicya mossambica* (A & B)*,*

Scorpaenopsis possi (C & D), different colormorphs of orange fluorescing

Soleichthys heterorhinos (E & F), undefined species of Lutjanidae (G) and *Brachysomophis*

henshawi (H) with a red fluorescing head. A, B, D, F-H: Banda Sea, C & E: Red Sea.

Figure 12: Fluorescence in fish: Detailed fluorescent pictures of *Anampses meleagrides* (A,

B; both overlay of GFP and CY5 filter), *Doryrhamphus excisus* (C; overlay of mCherry and

CY5 filter), *Eviota atriventris* (D, E; both overlay of GFP & CY5 filter), *Eviota nigriventris* (F-I;

F: overlay of GFP, CY5, mCherry filter; G: mCherry and CY5 filter; H: brightfield & CY5 filter;

I: mCherry filter), *Cirrhilabrus aquamarinus* (J; mCherry filter), Pictures were taken on a

Thunder microscope (Leica, Germany).

Figure 13: Fluorescence in fish: Detailed fluorescent pictures of *Cirrhilabrus aquamarinus* (A;

brightfield & Cy5 filter), Pair of *Synchiropus sycorax* (B; ♀ left and ♂ right; overlay of

brightfield and Cy5 filter), first dorsal fin of *Synchiropus sycorax* ♂ (C; brightfield and Cy5

filter), detailed fluorophores of *Synchiropus sycorax* (D-F; Cy5 filter), fluorescent patch

around the eye of *Synchiropus sycorax* ♂ (G; Cy5 filter). Pictures were taken on a Thunder

microscope (Leica, Germany).

References

1.   Johnsen S. The Optics of Life. A Biologist's Guide to Light in Nature. Princeton:

Princeton University Press; 2011.

2.   Poding L, Jägers P, Huhn M, Herlitze S. Diversity and function of fluorescent molecules

in marine organisms. EvoEcorxiv; 2023. preprint doi: 10.32942/X2CW2R

3.   Shimomura O. Discovery of Green Fluorescent Protein. In: Chalfie M, editor. Green

fluorescent protein. Properties, applications and protocols. 2nd ed. Hoboken, NJ: Wiley-

Interscience; 2006. pp. 1–13.

4.   Haddock SHD, Dunn CW. Fluorescent proteins function as a prey attractant:

experimental evidence from the hydromedusa Olindias formosus and other marine

organisms. Biol Open. 2015; 4:1094–104. Epub 2015/07/31. doi: 10.1242/bio.012138

PMID: 26231627.

5.   Macel M-L, Ristoratore F, Locascio A, Spagnuolo A, Sordino P, D'Aniello S. Sea as a

color palette: the ecology and evolution of fluorescence. Zoological Lett. 2020; 6:9.

Epub 2020/06/10. doi: 10.1186/s40851-020-00161-9 PMID: 32537244.

6.   Alieva NO, Konzen KA, Field SF, Meleshkevitch EA, Hunt ME, Beltran-Ramirez V, et al.

Diversity and evolution of coral fluorescent proteins. PLOS ONE. 2008; 3:e2680. Epub

2008/07/16. doi: 10.1371/journal.pone.0002680 PMID: 18648549.

7. Chudakov DM, Matz MV, Lukyanov S, Lukyanov KA. Fluorescent proteins and their

applications in imaging living cells and tissues. Physiological Reviews. 2010; 90:1103–

63. doi: 10.1152/physrev.00038.2009 PMID: 20664080.

8. Sparks JS, Schelly RC, Smith WL, Davis MP, Tchernov D, Pieribone VA, et al. The

covert world of fish biofluorescence: a phylogenetically widespread and phenotypically

variable phenomenon. PLoS One. 2014; 9:e83259. Epub 2014/01/08.

doi: 10.1371/journal.pone.0083259 PMID: 24421880.

9. Gerlach T, Sprenger D, Michiels NK. Fairy wrasses perceive and respond to their deep

red fluorescent coloration. Proc Biol Sci. 2014; 281. doi: 10.1098/rspb.2014.0787 PMID:

24870049.

10. Gerlach T, Theobald J, Hart NS, Collin SP, Michiels NK. Fluorescence characterisation

and visual ecology of pseudocheilinid wrasses. Front Zool. 2016; 13:13. Epub

2016/03/15. doi: 10.1186/s12983-016-0145-1 PMID: 26981144.

11. Wucherer MF, Michiels NK. Regulation of red fluorescent light emission in a cryptic

marine fish. Front Zool. 2014; 11:1. Epub 2014/01/08. doi: 10.1186/1742-9994-11-1

PMID: 24401080.

12. Kalb N, Schneider RF, Sprenger D, Michiels NK. The Red-Fluorescing Marine Fish

Tripterygion delaisi can Perceive its Own Red Fluorescent Colour. Ethology. 2015;

121:566–76. doi: 10.1111/eth.12367.

13. Bitton P-P, Harant UK, Fritsch R, Champ CM, Temple SE, Michiels NK. Red

fluorescence of the triplefin Tripterygion delaisi is increasingly visible against

background light with increasing depth. R Soc Open Sci. 2017; 4:161009. Epub

2017/03/22. doi: 10.1098/rsos.161009 PMID: 28405391.

14. Anthes N, Theobald J, Gerlach T, Meadows MG, Michiels NK. Diversity and Ecological

Correlates of Red Fluorescence in Marine Fishes. Front Ecol Evol. 2016; 4.

doi: 10.3389/fevo.2016.00126.

15. Liaci L. Natura e localizzazione di un pigmento fluorescents in Aaptos aaptos O. S.

(Demospongiae). Bolletino di zoologia. 1962; 29:425–31.

doi: 10.1080/11250006209440630.

16. Read KR, Davidson JM, Twarog BM. Fluorescence of sponges and coelenterates in

blue light. Comparative Biochemistry and Physiology. 1968; 25:873–82.

doi: 10.1016/0010-406X(68)90575-6.

17. Betti F, Bavestrello G, Cattaneo-Vietti R. Preliminary evidence of fluorescence in

Mediterranean heterobranchs. Journal of Molluscan Studies. 2021; 87.

doi: 10.1093/mollus/eyaa040.

18. Rodrigo AP, Lopes A, Pereira R, Anjo SI, Manadas B, Grosso AR, et al. Endogenous

Fluorescent Proteins in the Mucus of an Intertidal Polychaeta: Clues for Biotechnology.

Mar Drugs. 2022; 20. Epub 2022/03/25. doi: 10.3390/md20040224 PMID: 35447897.

19. Lindquist N, Barber PH, Weisz JB. Episymbiotic microbes as food and defence for

marine isopods: unique symbioses in a hostile environment. Proc Biol Sci. 2005;

272:1209–16. doi: 10.1098/rspb.2005.3082 PMID: 16024384.

20. Burgess WE, Axelrod HR, Hunziker RE. Dr. Burgess's Atlas of marine aquarium fishes.

Neptune City, NJ: T.F.H. Publ; 1990.

21. Debelius H. Fischführer Indischer Ozean. Rotes Meer bis Thailand ; über 900 Farbfotos

aus dem natürlichen Lebensraum der Meeresfische. 2nd ed. Melle: Tetra; 1996.

22. Patzner RA, Baensch HA. Perciformes (Barschartige). Ausgenommen Falter- und

Kaiserfische. 1st ed. Melle: Mergus Verl. für Natur- und Heimtierkunde Baensch; 1998.

23. Patzner RA, Mossleitner H. Non-Perciformes (Nicht-Barschartige). Sowie Falter- und

Kaiserfische. 1st ed. ; 1999.

24. Erhardt H. Wirbellose. 1st ed. Melle: Mergus Verlag für Natur- und Heimtierkunde

Baensch; 1998.

25. Erhardt H. Wirbellose. 1st ed. Melle: Mergus Verlag für Natur- und Heimtierkunde

Baensch; 2000.

26. Debelius H. Fisch-Führer Indopazifik. Malediven bis Philippinen ; 700 Fischarten in

ihrem natürlichen Lebensraum fotografiert. 1st ed. Münster: NTV; 2002.

27. Debelius H, Kuiter RH. Nacktschnecken der Weltmeere. 1200 Arten weltweit. 1st ed.

Stuttgart: Kosmos; 2007.

28. Kuiter RH, Debelius H. Atlas der Meeresfische. Die Fische an den Küsten der

Weltmeere. 3rd ed. Stuttgart: Kosmos; 2009.

29. Kuiter RH, Debelius H. Atlas der wirbellosen Meerestiere. Weichtiere, Würmer,

Stachelhäuter, Krebstiere. Stuttgart: Kosmos; 2009.

30. Bollati E, Lyndby NH, D'Angelo C, Kühl M, Wiedenmann J, Wangpraseurt D. Green

fluorescent protein-like pigments optimise the internal light environment in symbiotic

reef-building corals. Elife. 2022; 11. Epub 2022/07/08. doi: 10.7554/eLife.73521 PMID:

35801683.

31. Eyal G, Wiedenmann J, Grinblat M, D'Angelo C, Kramarsky-Winter E, Treibitz T, et al.

Spectral Diversity and Regulation of Coral Fluorescence in a Mesophotic Reef Habitat

in the Red Sea. PLOS ONE. 2015; 10:e0128697. Epub 2015/06/24.

doi: 10.1371/journal.pone.0128697 PMID: 26107282.

32. Masuda H, Takenaka Y, Yamaguchi A, Nishikawa S, Mizuno H. A novel yellowish-green

fluorescent protein from the marine copepod, Chiridius poppei, and its use as a reporter

protein in HeLa cells. Gene. 2006; 372:18–25. Epub 2006/02/14.

doi: 10.1016/j.gene.2005.11.031 PMID: 16481130.

33. Evdokimov AG, Pokross ME, Egorov NS, Zaraisky AG, Yampolsky IV, Merzlyak EM, et

al. Structural basis for the fast maturation of Arthropoda green fluorescent protein.

EMBO Rep. 2006; 7:1006–12. Epub 2006/08/25. doi: 10.1038/sj.embor.7400787 PMID:

16936637.

34. Hunt ME, Scherrer MP, Ferrari FD, Matz MV. Very bright green fluorescent proteins

from the Pontellid copepod Pontella mimocerami. PLOS ONE. 2010; 5:e11517. Epub

2010/07/14. doi: 10.1371/journal.pone.0011517 PMID: 20644720.

35.   Shagin DA, Barsova EV, Yanushevich YG, Fradkov AF, Lukyanov KA, Labas YA, et al.

GFP-like proteins as ubiquitous metazoan superfamily: evolution of functional features

and structural complexity. Mol Biol Evol. 2004; 21:841–50. Epub 2004/02/12.

doi: 10.1093/molbev/msh079 PMID: 14963095.

36.   Ohtsuka S, Huys R. Sexual dimorphism in calanoid copepods: morphology and

function. Hydrobiologia. 2001; 453/454:441–66. doi: 10.1023/A:1013162605809.

37.   Mazel CH, Cronin TW, Caldwell RL, Marshall NJ. Fluorescent enhancement of signaling

in a mantis shrimp. Science. 2004; 303:51. Epub 2003/11/13.

doi: 10.1126/science.1089803 PMID: 14615546.

38.   Mazel C. Method for Determining the Contribution of Fluorescence to an Optical

Signature, with Implications for Postulating a Visual Function. Front Mar Sci. 2017; 4.

doi: 10.3389/fmars.2017.00266.

39.   Gouveneaux A, Flood PR, Erichsen ES, Olsson C, Lindström J, Mallefet J. Morphology

and fluorescence of the parapodial light glands in Tomopteris helgolandica and allies

(Phyllodocida: Tomopteridae). Zoologischer Anzeiger. 2017; 268:112–25.

doi: 10.1016/j.jcz.2016.08.002.

40.   Deheyn DD, Enzor LA, Dubowitz A, Urbach JS, Blair D. Optical and physicochemical

characterization of the luminous mucous secreted by the marine worm Chaetopterus

sp. Physiol Biochem Zool. 2013; 86:702–5. Epub 2013/10/16. doi: 10.1086/673869

PMID: 24241067.

41.   Branchini BR, Behney CE, Southworth TL, Rawat R, Deheyn DD. Chemical Analysis of

the Luminous Slime Secreted by the Marine Worm Chaetopterus (Annelida,

Polychaeta). Photochem Photobiol. 2014; 90:247–51. Epub 2013/10/17.

doi: 10.1111/php.12169 PMID: 24004150.

42.   Deheyn DD, Latz MI. Internal and secreted bioluminescence of the marine polychaete

Odontosyllis phosphorea (Syllidae). Invertebrate Biology. 2009; 128:31–45.

doi: 10.1111/j.1744-7410.2008.00149.x.

43. Meulenaere E de, Puzzanghera C, Deheyn DD. Self-powered bioluminescence in a

marine tube worm. The FASEB Journal. 2020; 34:1.

doi: 10.1096/fasebj.2020.34.s1.04618.

44. Carlos AA, Baillie BK, Maruyama T. Diversity of dinoflagellate symbionts

(zooxanthellae) in a host individual. Mar Ecol Prog Ser. 2000; 195:93–100.

doi: 10.3354/meps195093.

45. Maxwell K, Johnson GN. Chlorophyll fluorescence—a practical guide. Journal of

Experimental Botany. 2000; 51:659–68. doi: 10.1093/jexbot/51.345.659.

46. Yau AJ-Y, Fan T-Y. Size-dependent photosynthetic performance in the giant clam

Tridacna maxima, a mixotrophic marine bivalve. Mar Biol. 2012; 159:65–75.

doi: 10.1007/s00227-011-1790-8.

47. Winters AE, Green NF, Wilson NG, How MJ, Garson MJ, Marshall NJ, et al. Stabilizing

selection on individual pattern elements of aposematic signals. Proc Biol Sci. 2017; 284.

doi: 10.1098/rspb.2017.0926 PMID: 28835556.

48. Brauwer M de, Hobbs J-PA, Ambo-Rappe R, Jompa J, Harvey ES, McIlwain JL.

Biofluorescence as a survey tool for cryptic marine species. Conserv Biol. 2018;

32:706–15. Epub 2018/02/19. doi: 10.1111/cobi.13033 PMID: 28984998.

A

B

C

D
863,3 µm

E
932,7 µm

F
256,5 µm

G
256,5µm

H
216,8 µm

5,7 mm

A
2,1 mm

B
5,8 mm

C
5,8 mm

D
5,8 mm

E
5,8 mm

F
5,7 mm

G
5,8 mm