Sediment source and dose influence the larval performance of the threatened coral

Orbicella faveolata

Xaymara M. Serrano[1,2*,#a], Stephanie M. Rosales[1,2], Margaret W. Miller[3], Ana M. Palacio-

Castro[1,2], Olivia M. Williamson[4], Andrea Gomez[5], and Andrew C. Baker[4*]

[1]Cooperative Institute of Marine and Atmospheric Studies, Rosenstiel School of Marine,

Atmospheric, and Earth Science, University of Miami, Miami, Florida, USA

[2]National Oceanic and Atmospheric Administration, Atlantic and Oceanographic Meteorological

Laboratory, Miami, Florida, USA

[3]SECORE International, Hilliard, Ohio, USA

[4]Department of Marine Biology and Ecology, Rosenstiel School of Marine, Atmospheric, and

Earth Science, University of Miami, Miami, Florida, USA

[5]National Oceanic and Atmospheric Administration, Greater Atlantic Regional Fisheries Office,

Gloucester, Massachusetts, USA

[#a]Current address: National Marine Fisheries Service, Habitat Conservation Division, West Palm

Beach, Florida, USA

[*]Corresponding authors

Emails: xaymara.serrano@noaa.gov (XS), abaker@earth.miami.edu (AB)

Abstract

The effects of turbidity and sedimentation stress on early life stages of corals are poorly understood, particularly in Atlantic species. Dredging operations, beach nourishment, and other coastal construction activities can increase sedimentation and turbidity in nearby coral reef habitats and have the potential to negatively affect coral larval development and metamorphosis, reducing sexual reproduction success. In this study, we investigated the performance of larvae of the threatened Caribbean coral species *Orbicella faveolata* exposed to sediments collected from a reef site in southeast Florida recently impacted by dredging (Port Miami), and compared it to the performance of larvae exposed to sediments collected from the offshore, natal reef of the parent colonies. In a laboratory experiment, we tested whether low and high doses of each of these sediment types affected the survival, settlement, and respiration of coral larvae compared to a no-sediment control treatment. In addition, we analyzed the sediments used in the experiments with 16S rRNA gene amplicon sequencing to assess differences in the microbial communities present in the Port versus Reef sediments, and their potential impact on coral performance. Overall, *O. faveolata* larvae exposed to high doses of either sediment type (Port or Reef) exhibited reduced survival and settlement rates, but only the Port sediments resulted in adverse effects in the low-dose treatment. Sediments collected near the Port also contained different microbiomes than Reef sediments, and higher relative abundances of the bacteria Desulfobacterales, which has been associated with coral disease. We hypothesize that differences in microbiomes between the two sediments may be a contributing factor in explaining the observed differences in larval performance. Together, these results suggest that the settlement success and survival of *O. faveolata* larvae are more readily compromised by encountering port inlet sediments compared to

45   reef sediments, with potentially important consequences for the recruitment success of this species

46   in affected areas.

Introduction

Dredging and port construction activities can induce chronic sedimentation and turbidity stress in surrounding coral reef habitats [1-6]. Sedimentation caused by dredging is often more harmful to corals than sedimentation caused by other activities, due to its acute onset and often long duration, typically lasting many months to years [1, 7]. Sedimentation effects vary depending on grain size and composition (e.g., [7-12]), and dredging activities can release large quantities of sediments in the water column that are typically of finer grain size compared to naturally occurring sediments (<63 μm, [7]). These finer sediments tend to attenuate more light and are more prone to being resuspended, thus causing a greater net reduction in light essential for photosynthesis [10]. Finer sediments also have an adhesive, clay-like texture that is more resistant to bioturbation when deposited [13], can lead to a change in the bacterial community [12], and is more likely to become anoxic [8, 14]. Finally, suspended sediment plumes, especially those with finer grain size, can also travel several km away from the dredging site (e.g., [9, 15, 16]), and can also release contaminants [17-19] and pathogens [11, 20, 21] which may potentially transmit coral diseases such as Stony Coral Tissue Loss Disease [22].

Sediments can negatively affect corals throughout their life cycle, suppressing coral health, condition, and survival via multiple mechanisms (reviewed in [7, 12, 13, 23]). Reported effects on early life stages of corals include reductions in the fecundity of parent colonies [24], fertilization [25, 26], larval settlement [27-31], and recruit survival [5, 27, 32, 33]. Tolerance to sedimentation exposure is estimated to be an order of magnitude lower for coral recruits compared to adults [13, 23, 34], although most of this work has been conducted on Pacific coral species (reviewed in [12, 13, 23]). Much less is known regarding the effects of sediments on coral larvae and recruits for Atlantic coral species, and this information is critical for developing appropriate conservation

plans for these species. Poor understanding of coral responses to sediment disturbances during

their most susceptible stages can result in inappropriate management of dredging projects, lead to

preventable coral losses, and unnecessarily high costs to adaptively manage operations (corrective

actions) and provide compensatory mitigation to offset impacts.

Sediments may interfere with the ability of coral larvae to find a suitable substrate on which

to settle, which is key to successful recruitment and survival [31, 35, 36]. Even a thin layer of

sediments not harmful to adult corals can be detrimental to coral larvae [23, 28, 31, 37-39], both

by physically obstructing settlement surfaces [27, 30], and impairing biotic settlement cues such

as those from crustose coralline algae (CCA) [30]. However, the mechanisms and cues by which

sediments can affect settlement and recruitment success, or benthic habitat quality (e.g., [31]) are

poorly understood. Recent work suggests that coral larvae show a preference for distinct microbial

and/or chemical signature types in certain types of CCA [40-43], suggesting that specific bacterial

communities can serve as environmental cues for coral settlement [40, 42]. How and whether

microbial communities present in sediments can negatively affect the ability of coral larvae to

identify these cues remains an open question.

The aim of the present study was to investigate the performance of coral larvae of an

important Caribbean reef-building species (*Orbicella faveolata*) following a short-term (pulse)

sediment exposure. *O. faveolata* (Ellis and Solander, 1786) is listed as threatened under the U.S.

Endangered Species Act [44] but is present in southeast Florida in the vicinity of two large ports

with expansions (deepening/widening) planned in the near future (Port Everglades and Port

Miami). The most recent large-scale deepening project at Port Miami (which occurred from 2013-

2015) resulted in severe sedimentation impacts to local coral reef habitats [1, 3, 15, 45], although

the severity of those impacts has been disputed [46]. In addition, recent work exposed adult *O.*

faveolata corals to Port Miami sediments during a 96-hr period and showed adverse effects on

tissue regeneration capacity compared to no sediment controls [47]. In this study, we exposed *O.*

faveolata larvae to sediments collected near the Port Miami channel and compared their

performance with larvae exposed to reef sediments collected from the natal reef of the parent

colonies in the Florida Keys. We compared the effects of the two different sediment types on the

survival, settlement, and respiration of coral larvae, and also characterized the bacterial

communities of these sediments using 16S rRNA gene amplicon sequencing to assess the potential

role of microbial communities in affecting coral performance. To our knowledge, this is the first

study to investigate the effects of suspended sediments on larvae of western Atlantic (Caribbean)

coral species, and the first to use sediments that have not been dried or sterilized prior to exposure,

which may have masked some of the biological or chemical effects of sedimentation in previous

studies.

Methods

Coral larvae collection

O. faveolata larvae were collected and cultured as described in [48] and [49]. Gamete

bundles were collected seven nights after the full moon (night of 14th August 2017) from colonies

that spawned at two reefs in the upper Florida Keys (Horseshoe and Sand Island reefs). Once the

bundles broke apart, equal volumes of gametes from 3-4 parental genets were combined for

fertilization. Resulting embryos were maintained in water collected from their natal reef at a land-

based facility in the Keys (either in static bins with regular water changes or in a recirculating

system with mesh-floored bins and a constant drip input) and cultured for ~2 days so that

unfertilized eggs would disintegrate, leaving behind only developing larvae. Embryos were kept

in a shaded area under ambient light conditions and a mean temperature of 29^{o}C. Larvae were then

transferred to the University of Miami for experiments.

Sediment collections and maintenance

Immediately prior to starting the larval experiments (24 - 48 h), two batches of sediments

were collected for experiments, one from a "Reef" origin and another from a "Port" origin. Reef

sediments were collected at Horseshoe Reef in Key Largo, from the same site where the coral

parent colonies occurred (25°8'19"N, 80°17'41"W, Fig 1). Port sediments were collected from a

site located ~200 m north of the recently-dredged channel of Port Miami (25°45'50"N, 80°5'56"W,

Fig 1), reported in [3] as having detrimental effects from dredging-induced sedimentation on adult

corals. This site was shown by [3] as the most severely impacted location surveyed, with the

highest sediment cover (43%) and highest prevalence of sediment "halos" (indicating partial

mortality) compared to the reference stations.

Fig 1. (A) Map of locations on the coast of southeastern Florida where Port and Reef

sediments were collected from (near Port Miami and Horseshoe reef in Key Largo). (B) Map

of coral reef habitats around the Port of Miami federal channel. (C) Port sediments were collected

from a site within the Inner Reef located ~200 m north of the recently dredged channel of Port

Miami (circled in red), which was the most severely impacted location surveyed due to

sedimentation compared to the reference stations by [3]. (D) Representative photo of the sediments

collected near the Port of Miami. (E) Representative photo of the sediments collected at Horseshoe

reef.

All sediments were taken from the upper ~3 cm of the sediment layer (surficial layer) by

sliding a metal tool below the sediment-water interface and then scooping them into a certified-

clean HPDE sample container. Three to four containers were collected per site and combined to

produce a single "batch" for experiments. Sediments were then brought to the laboratory,

homogenized in a beaker of seawater and kept in continuous movement and aeration until used in

experiments. Prior to use, sediments were allowed to settle and water was siphoned out with care

to avoid disturbing the fine particles on the surface. Sediments were then wet-weighed (adjusting

for water content) to avoid killing any microorganisms present while retaining as many fine

particles as possible. The treatments were conducted using freshly collected sediments to be able

to assess any physical (e.g., abrasion) and biological (e.g., pathogenic) stress responses.

A subset of the sediments was analyzed for grain size by sieving and pipetting after drying

the sediments thoroughly in an oven. Grain size limits were defined according to the Udden-

Wentworth US standard classification [50], while statistical parameters were determined using

graphical measures from the cumulative size distributions as shown in S1 Fig. A summary of these

parameters are provided in S1 Table. Reef sediments can be described as a well-sorted medium

sand with virtually no silt or gravel. Conversely, Port sediments can be described as a poorly-

sorted, medium sand with small amounts of gravel (approx. 1.97% >2000 μm) and silt and clay-

sized particles (also referred as "mud", approx. 1.25% <63 μm). No chemical or organic matter

analysis of the sediments were conducted.

Experimental setup

Fifty 8-day-old larvae were added to individual glass vials containing 45 mL filtered

seawater (5 μm). Vials were allocated randomly to one of five different treatments: (1) a control

treatment (no sediment), (2) a low "Reef" suspended sediment treatment (20 mg/L of homogenized

sediment from Horseshoe Reef added), (3) a low "Port" sediment treatment (20 mg/L sediment

from the Port site added), (4) a high "Reef" sediment treatment (100 mg/L sediment from

Horseshoe reef added), and (5) a high "Port" sediment treatment (100 mg/L sediment from the

Port site added). Nine to ten replicates were used per treatment. Vials were attached to a titer plate

shaker set to 300 rpm to continuously maintain sediments in suspension for 24 h. Initial trials were

conducted to assess the best exposure times that caused minimal mortality in control larvae and

maintained both fine and coarse sediment particles in continuous suspension.

Treatments were chosen to characterize sub-lethal and lethal responses to suspended

sediment concentrations for "tolerant" versus "sensitive" species as described by [12]. In addition,

our goal was to test a low-concentration of suspended sediment (20 mg/L) representing typical

conditions near dredging operations. Jones et al. [13] analyzed suspended sediment concentrations

over a 30-day running mean period during the Barrow Island dredging project in Australia and

found that the 95th percentile was 21 mg/L for 7 sites located at distances <1 km from the dredging.

Similarly, although our 100 mg/L (high-dose) suspended sediment treatments were approximately

5 times higher than those typically observed during dredging operations, such conditions do occur

over short periods (see [51]).

24 hour treatment exposure: larval survival assays

After 24 h of exposure, swimming larvae remaining in each vial were pipetted out, counted

and examined with a microscope to determine the proportion that had survived. The vial was also

carefully examined for potential settlers. A subset of larvae (N=6 per vial) was used to assess

changes in respiration rates. Remaining larvae were allowed to recover for one week in filtered

seawater (without sediments) in the same vials before being transported to the Florida Keys to

conduct the settlement assays.

1-week post-exposure: larval settlement assays

Settlement assays were conducted by pooling all the larvae remaining in each treatment

and then redistributing N=15 larvae to individual vials containing filtered seawater and a piece of

rubble collected from Horseshoe reef to act as settlement cue. A total of 10 replicate glass vials

per treatment were used to quantify larval settlement per treatment. After 24 h, the total number of

larvae still swimming, or that had settled on the rubble, was manually scored using fluorescence

microscopy. Larvae were classified as "settlers" only if they displayed visible signs of settlement

(attachment to the substrate) and metamorphosis (i.e., transition from pear-shaped to flat/disc

shape). To determine the cumulative percentage of larval settlement, the total number of settled

corals (spat) was divided by the initial number of larvae added to each container.

24 hour and 1-week assessment of coral larval respiration

Respiration was measured using a microplate reader developed by Loligo Systems

(Denmark), which allows measuring real-time respiration rates in each one of 24 wells using an

optical fluorescence oxygen sensing technology (SDR SensorDish Reader®, PreSens, Germany)

as described in [52-54]. This system comes pre-calibrated specifically for the reader used, and

oxygen solubility is continuously calculated with oxygen sensors (optrodes) attached to the bottom

of each well. Larval respiration rates were assessed at two time points, immediately after the 24 h

treatment exposure and 1-week post-exposure. At each time point, larvae (N=6) were taken at

random from each vial and placed in each of the wells of a sealed glass 125-µL microplate (without

any sediments). A total of 8-10 replicate wells were used per treatment. Four additional control

wells were used in each plate without any larvae to measure any background respiration in the

treatment water.

Larvae were visually inspected for normal swimming behavior prior to each run. After each

plate was prepared, wells were inspected to ensure there were no air bubbles, and an oxygen-

impermeable seal was created using a silicone membrane covered with parafilm and a compression

block. To maintain constant temperature, the microplate was placed inside a temperature-

controlled flow-through water bath (included as part of the system), in a light-controlled room.

The plate was then placed in a titer shaker at a low speed to maintain constant water motion.

Finally, measurements of oxygen concentrations were taken every 15 seconds throughout the

duration of the run and recorded using the SDR v 4.0.0 software (PreSens, Germany). Oxygen

concentration was plotted as a function of time for individual wells and the first 10-20% (pO_2 in

% air saturation) linear decreases in oxygen were used to calculate the oxygen consumption rate

(an example plot of oxygen consumption over time for 6 *O. faveolata* larvae placed in a single

well of the microplate is shown in S2 Fig). Any portion of the slope that dipped below 70% air

saturation was not used for analysis. Respiration rates were then corrected for background

respiration and calculated as nanomoles of oxygen consumed per larva per minute.

16S sequencing of sediment samples

Immediately upon collection, ~1 gram from each of the Port and Reef sediment samples

collected per site was preserved in RNAlater for 16S sequencing. The DNA was extracted and

PCR amplified with 16S rRNA gene V4 primers [55] with the Platinum Hot Start PCR Master Mix

(2X) (ThermoFisher Scientific, Waltham, MA). The master mix was a 48-μl reaction and 2 μl of

DNA template was used. The DNA was then amplified as follows: 94 °C for 3 minutes (1X), 94 °C

for 45 seconds (35X), 50 °C for 60 seconds (35X), 72 °C for 90 seconds (35X), and 72 °C for

10 minutes (1X). Amplified products were cleaned using AMPure XP beads (Beckman Coulter,

Brea, CA), normalized to 4 nM, and 5 µl of each normalized sample was pooled. The samples were

sequenced on a MiSeq with PE-300v3 kits at the Halmos College of Natural Sciences and

Oceanography at Nova Southeastern University.

Statistical analysis

Larval performance

All analyses of larval performance data were done in R v.4.1.2. The effects of the sediment

treatments on the proportion of larval survivorship and settlement were analyzed using binomial

generalized linear models (GLM) with the lme4 package [56]. The models included sediment

treatment as a fixed effect and replicate (vial) as a random effect. Model predictions (survivorship

and settlement probabilities as well as the odds ratio between the control and the treatments) were

plotted using the plot_model function from the sjPlot package [57] including 95% confidence

intervals. Larval respiration rates after 24 h of treatment exposure and 1-week post-exposure were

analyzed using a linear mixed-effects model (LMEM) with the lme4 package. The model included

sediment treatment and time point (exposure and recovery) as fixed interactive effects, and vial,

plate, and well as random effects. All data and code for larva survivorship, settlement and oxygen

consumption can be found at Zenodo [58].

Microbiome analysis of sediment samples

DNA sequences were demultiplexed and further processed with Qiime2-2022.11 [59].

Since the reverse reads were of poor quality only the forward reads were analyzed. Using the

program Cutadapt the primers were removed [60], and Amplicon Sequence Variants (ASVs) were

generated with the program DADA2 [61] trimmed at the 10 and 205 bp positions. The ASVs were

taxonomically classified with the classify-consensus-vsearch function and the silva 138 99

database [62]. If the sequences were annotated as chloroplast or mitochondria they were removed

from the analysis. The data were then uploaded to R and converted into a Phyloseq v1.26.1 data

object for analysis and graphical display of phylogenetic sequencing data.

Alpha-diversity was evaluated by rarefying to a minimum sequence depth of 60,000. Three

alpha-diversity metrics, richness, Shannon-Wiener, and the Inverse Simpson index, were

generated with the phyloseq estimate_richness function. The respective values were then tested

for normality using qqnorm, qqline, and the Shapiro test. Upon passing normality, significance

between sites was tested with an Analysis of Variance (ANOVA).

For beta-diversity, ASVs were filtered if they summed to <10 in 30% of the data. The

filtered data were then transformed to centered log ratios (CLR) using the package Microbiome

v0.9.99. The Vegan v2.5.4 package was used for beta-diversity testing for both within and between

group differences. Dispersion of the samples (within group) was tested by using the function

Vegdist (method = "Euclidean", Permutations = 999) and Betadisper and was tested for

significance with a permutation test. A permutational multivariate analysis of variance

(PERMANOVA, function Adonis, method = "Euclidean", Permutations = 999) was used to

evaluate significance between groups.

To identify differentially abundant microbial ASVs between sites, the program Analysis of

Compositions of Microbiomes with Bias Correction (ANCOM_BC) was used [63]. ANCOM_BC

was used with an alpha <0.001 and a W statistic was >90. The ASVs were further evaluated if they

had a log-fold change < -3.5 or > 3.5.

Results

Larval performance

Sediment treatments significantly impacted the performance of *O. faveolata* larvae (Fig 2, S2 Table). In general, sediment exposure negatively impacted larval survival and settlement for both Port and Reef sediment types. However, the effect of sediment concentration was important depending on the sediment type used in the treatment. Whereas both high and low Port sediment concentrations negatively impacted larval survival and settlement, only the high-dose Reef sediment treatment resulted in adverse effects (Figs 2 and S3). Specifically, survival proportions after 24 h in the treatments were above 93% for Control and Low Reef, but exposure to Low Port, High Reef, and High Port reduced survival probabilities to 90%, 86%, and 81%, respectively ($p < 0.05$, df = 4, S2 and S3 Tables), resulting in significantly lower odds ratios of survivorship for these treatments (<0.6) compared to the Control (S3 Fig). Larval settlement probabilities 1-week after treatments were 21-22% in the Control and Low Reef treatments, but were reduced to less than 12% in the Low Port, High Reef, and High Port treatments ($p < 0.05$, df = 4, S2 and S3 Tables). This resulted in lower (<0.5) odds ratios for settlement in the Low Port, High Reef, and High Port treatments compared to the Control (S3 Fig).

Fig 2. Effects of sediments on *Orbicella faveolata* larvae (model estimated values ± 95% CI). (A) Proportion of larval survival after a 24 h exposure to experimental treatments. (B) Proportion of larval settlement after one week of recovery from experimental treatments. Asterisks denote significant differences in a sediment treatment compared to controls with no sediment added (*: $p < 0.05$; **: $p < 0.01$; ***: $p < 0.001$). Colors match Figs 3 and 4 and denote either Port or Reef sediment treatments.

None of the sediment treatments affected the larval respiration rates at either of the time points (S4 Table, S4 Fig). Average respiration rates per larva were $9.33 \times 10^{-4} \pm 2.53 \times 10^{-4}$ nmol O_2 min^{-1} (mean $\pm$ SD) across treatments after 24 h of exposure and declined to $6.01 \times 10^{-4} \pm 1.97 \times 10^{-4}$ nmol O_2 min^{-1} after a week of recovery, but this was only marginally significant (p<0.06, df = 1, F = 15.83, S4 Table).

Microbiome analysis

A total of seven sediment samples (Port N=3 and Reef N=4) resulted in sequence counts between 93,626 and 419,403 (median=171,281). After processing and filtering, 1,852 ASVs were used for the analysis. The median alpha-diversity was slightly higher in the Reef sediments (richness: Reef=3604 vs Port=3001, Shannon-Wiener: Reef=7.26 vs Port=7.20, and Inverse Simpson: Reef=654.24 vs Port=584.8), but this difference was not significant between sites (Fig 3A). In contrast, within-group (permutest, p=0.023) and between-group beta-diversity were significantly different (PERMANOVA, Fmodel=6.127, R^2=0.55 p=0.03, Fig 3B), with the microbial communities in Reef sediments having a more dispersed beta-diversity compared to Port sediments (average distance to centroid: Reef=55.76 vs Port=41.99). Of the seven most relatively abundant bacteria orders, the Reef and Port sediments shared four taxa (57%) with NB1-J and Cyanobacteriales being unique to Reef sediments, and Desulfobacterales being unique to the Port sediments (Fig 3C).

Fig 3. Microbial communities found in Port and Reef sediments. (A) Alpha diversity differences between Reef and Port samples using three diversity metrics: (1) Species richness (2) Shannon diversity Index, and (3) Inverse Simpson Index. (B) Beta diversity of microbial

composition between Port and Reef sediment samples. (C) The cumulative relative abundances of

the most abundant (0.05>) microbial order per sample grouped by the site of sediment collections.

Differential abundance analysis was conducted to identify the differences between sites,

resulting in 293 significant ASVs (Fig 4). Of these, there were a total of 74 orders, 82 families,

and 96 genera. The most prevalent differentially abundant orders were Steroidobacterales (N=29),

Desulfobacterales (N=19), and Pirellulales (N=15). The Port sediments had more enriched ASVs

(N=223) compared to Reef sediments (N=70). More specifically, the ASVs most enriched in Port

sediments were from the orders Ectothiorhodospirales (family *Ectothiorhodospiraceae*, genus

Thiogranum), an uncultured genus from the Class Gammaproteobacteria, Desulfobacterales

(*Desulfosarcinaceae*, Sva0081 sediment group), and Steroidobacterales (*Woeseiaceae*, *Woeseia*).

In the Reef sediments, the most enriched orders were Alteromonadales (*Moritellaceae* from an

uncultured genus), Steroidobacterales *(Woeseiaceae,   Woeseia*), and Nitrosococcales

(*Nitrosococcaceae*, *AqS1*).

Fig 4. Microbial Amplicon Sequence Variants (ASVs) associated with Reef or Port

sediments. Differential abundances between Reef and Port, where the y-axis shows significant

ASVs (padj<0.001, W statistic >90) grouped by order. Only ASVs with a log-fold change <-3.5

and >3.5 were visualized.

Discussion

In this study, we investigated differences in the survival and settlement of larvae from the

threatened Caribbean coral *Orbicella faveolata* after short-term exposure to sediments collected

near the recently dredged Port Miami channel (considered anthropogenically modified) versus

sediments collected from the natal reef of the parent colonies in the Florida Keys (considered

natural or unmodified). Neither of the sediment types were dried or sterilized to avoid changing

their biological or chemical properties. Overall, we found that although *O. faveolata* larvae

exposed to high doses of both sediment types ("Port" and "Reef") exhibited adverse survival and

settlement rates, only the Port sediments resulted in adverse effects when exposed to low-dose

treatments, characteristic of conditions near actual dredging projects (e.g., [13]). The low-dose

Reef sediment treatment was not detrimental to coral larvae, and showed similar outcome odds of

survival and settlement compared to the no sediment (control) treatment (see S3 Fig).

Port sediments had different microbiome profiles and significantly higher abundances of

potentially pathogenic bacteria compared to Reef sediments, which may explain the reduced larval

survival and settlement observed even in the low-dose treatment. We speculate that Port sediments

could have been more detrimental to larvae compared to reef sediments due to differences in their

physical and chemical composition (e.g., grain size and sediment composition), and other

biological differences (e.g., different microbial communities). Together, these findings suggest

that, although suspended sediments may be generally detrimental to the survival and recruitment

potential of *O. faveolata* larvae, anthropogenic sediments, such as those occurring adjacent to

recent dredging projects, may incur greater harm compared to natural reef sediments.

Effects on larval survival

Both of our sediment sources significantly impacted the survival of *O. faveolata* larvae,

however, exposure to both the low- and high-dose Port sediment treatments reduced survival

proportions to 86% and 81%, respectively (compared to the control treatment), while the high-

dose Reef sediment treatment only reduced the survival proportion to 90% (Fig 2A). Suspended

sediments may reduce larval survival through physical abrasion [38] and reduced light over

prolonged periods can also lead to decreased photosynthetic efficiency of larval symbionts and

subsequent larval mortality from starvation [23]. However, in our study, decreased light was

unlikely the driver of our results, given the short exposure to sediments (24 h) and the use of

aposymbiotic (symbiont-free) larvae. Instead, we suspect that reduced larval survival in both of

our high-dose sediment treatments (Fig 2A) could have been caused by continuous physical

abrasion from suspended sediments during exposure. Given the pelagic nature of coral larvae, this

stage is likely to be more prone to physical abrasion from moving sediment particles compared to

coral recruits, on which sediment can accumulate on the growing coral tissue.

Increased mortality of *O. faveolata* larvae in both of our Port sediment treatments

(compared to Reef sediments) could have also been due to chemical or microbial differences in

the sediment composition and/or grain size. A recent study [32] assessed the effects of

sedimentation on recruits from the brooding coral *Porites astreoides* using either reef sediments

collected from the site of adult colonies, or sediments collected near Port Everglades (Fort

Lauderdale, Florida) to resemble the grain size composition often found in dredging areas. Similar

to our study, the authors found that reef sediments did not negatively affect coral survival, but Port

sediments did. However, since there were only trace amounts of mud-sized particles in our Port

sediments (approximately 1.25%), we hypothesize that Port sediments may be more likely to have

affected coral larvae through microbial processes leading to decreased survival.

Effects on larval settlement

Except for the low-dose Reef sediment treatment, we showed significant effects of both sediment types (Port and Reef) after one week of sediment exposure, resulting in approximately 50-60% reduction in settlement in *O. faveolata* larvae compared to the control treatment (Figs 2B and S1). Reduced larval settlement has been documented for a few Pacific coral species in the presence of suspended sediments (in concentrations less than 10 mg/L, [23]), presumably due to changes in light quality and quantity (as a result of increased light attenuation). Sediment-covered surfaces have also been shown to decrease larval settlement in various coral species such as *Pocillopora damicornis* [64], *Acropora digitifera* [38], *Acropora millepora* [38], and *O. faveolata* [31], likely due to limited space to settle (preventing larvae from accessing and attaching to suitable substrate), higher risk of abrasion, smothering/burial by deposited sediments, and masking of appropriate chemical cues for settlement [23].

An important distinction of our study compared to previous work on the effects of sedimentation on coral larvae is that we tested potential latent effects of sediment exposure on eventual settlement one week after acute exposure. Thus, the settlement environment (i.e., light and condition or exposure of settlement substrates) were the same across treatments, and consequently neither sediment deposition on the coral surfaces or settlement substrates, nor changes in light, were drivers of the reduced settlement observed in the exposed larvae. Furthermore, since our low-dose Reef sediment treatment was not detrimental to coral settlement (but the low-dose Port sediment treatment was), we speculate that disruption of potential settlement cues in larvae exposed to Port sediments may have resulted from biological/chemical damage (potential interactions with any contaminants or bacterial pathogens present in the Port sediments). Sediment-stressed larvae could also have had lower energy reserves available for

allocation to settlement and subsequent metamorphosis, although no significant respiration rate

differences were observed among treatments (see S4 Fig). One reason why our study may have

found latent effects is that previous studies conducted experiments using sediments that had been

dried prior to exposure. Oven drying can change the chemical composition of the sediments and

kill bacteria [5], masking these potentially important biological or chemical effects. The inclusion

of wet sediments in our study may therefore have helped preserve these properties and allowed us

to explore how bacterial assemblages can affect larval survival and settlement.

Microbiome analysis of sediment samples and potential implications for coral survival and settlement

Coral settlement is the result of complex behavioral responses that are modulated by

distinct microbial signatures in CCA [40-43]. In this study, we hypothesized that microbial

differences among the two sediment types may have resulted in decreases in survivorship and

potential impairment of the ability of larvae to detect cues for settlement and metamorphosis. The

microbiome analysis of the sediment samples collected near the Port revealed differences in

microbial beta-diversity and differentially abundant bacteria taxa compared to Reef sediments

(Figs 3BC and 4). Relative abundance analysis of the most abundant taxa showed that

Desulfobacterales were unique to the Port sediments (Fig 3C). These taxa have been found to be

enriched in microbial communities from other inlets in southeast Florida [65] and have also been

associated with black band disease [66], suggesting that this bacterial taxa may play a role in coral

health and its enrichment in Port sediments may have contributed to the reduced survival and

settlement in larvae exposed to both low- and high-doses of these sediments (Fig 2A). Moreover,

Desulfobacterales are anaerobic sulfate-reducing bacteria [67] that reduce sulfate (SO_4^{2-}) to

hydrogen sulfide (H_2S). Desulfobacterales are abundant in organic-rich and high nutrient

environments linked to eutrophication [68], and their waste product, H_2S, is correlated with coral

tissue degradation [11], and algal mortality [69]. Port sediments enriched in Desulfobacterales (Fig

4) may therefore reflect elevated H_2S and stressful hypoxic conditions that may have resulted in

lower survival and settlement rates.

Sediments may also act as vectors for the transmission of diseases such as Stony Coral

Tissue Loss Disease [22], which is documented to have appeared for the first time near Port Miami

in 2014 [70, 71]. Also, bacteria genera from two common human pathogens, *Staphylococcus* and

Streptococcus, have been found across multiple time points in surface water and sediment samples

near Port Everglades [65, 72], another southeast Florida inlet subject to dredging impacts. Our

findings add to these studies and also suggest that port inlets may carry potentially pathogenic

bacteria which may be transported to nearby reefs through various sediment transport processes

(natural or anthropogenic), resulting in potential exposure to coral larvae and other downstream

health impacts.

440 Effects on coral larval respiration

Despite significant effects of sedimentation on *O. faveolata* larval survival and settlement

(Figs 2 and S3), none of the sediment treatments had significant effects on coral larval respiration

(S4 Fig and S4 Table). Sedimentation has been shown to decrease photosynthesis to respiration

(P/R) ratios in adult corals, via decreased photosynthesis of the algal symbionts and increased

respiration of the host (e.g., [73]). However, in this study, we used aposymbiotic larvae, which

may explain the lack of significant effects on respiration. Alternatively, changes in respiration may

have not been detected because it is not possible to maintain the sediment treatments in the

microplate reader while respiration measurements are taken. Lower respiration rates across all

treatments at the 1-week recovery period (compared to the 24-hr sediment exposure, see S3 Fig)

could have resulted from ontogenetic and/or behavioral differences during those two time points

(active swimming versus exploration of suitable substrate for settlement), e.g. [74]).

Management implications for the conservation of corals

Our study is the first to examine the effects of suspended sediments on larvae from an

Atlantic coral species. With the two major commercial shipping ports in southeast Florida planning

major expansions which require multi-year dredge projects, this information is important for

understanding the impacts of sedimentation on susceptible life stages of corals, and assisting with

the development of protective management strategies. *O. faveolata* is listed under the US

Endangered Species Act (ESA) and already shows poor recruitment [49, 75, 76]. Recent work

confirms the ESA Critical Habitat (legal) requirements that larvae require substrates free of

sediment for settlement [31]. Importantly, our study showed that while sediments have detrimental

effects on the survival and settlement of *O. faveolata* larvae at high doses, even brief exposures to

low doses of anthropogenic sediments found near a recently dredged Port during the pelagic phase

could reduce larval survival and disrupt settlement, potentially compromising the recruitment

success and recovery potential of this species. Furthermore, we showed substantial direct mortality

from short-term sediment exposure, but also an additional significant latent effect on the surviving

larvae's capacity to successfully settle one week after sediment exposure. These results indicate

that sediments which remain adjacent to a dredging project even after ~2 years may still have

significant effects on early life stages of corals during relatively short-term resuspension events

(e.g., storms). This highlights the potential for long-term impacts of dredging (e.g., critical habitat

modification) as a result of resuspension of anthropogenic sediments. However, it is unclear how

long such sediments can persist in the area, or how often resuspension events might happen.

Our findings also highlight the importance of setting temporary moratoriums as a

management tool during dredging operations. Since 1993 dredging projects in Western Australia

conducted near coral reefs have been required to implement a coral environmental ("stand-down")

window spanning five days before spawning to seven days after spawning [13, 77]. The *O.*

faveolata larvae used in our experiments were eight days old and were adversely impacted by both

sediment types with just 24 h of exposure, with detrimental latent effects still detectable after one

week. Consequently, a one-week "stand down" period may be insufficient for broadcast spawner

coral species such as *O. faveolata* given the length of time it takes for larvae to become competent

and find suitable substrate for settlement. Based on our results, we recommend setting

environmental windows which encompass the months of peak spawning (typically August and

September for *O. faveolata* [78, 79]), plus an additional period of at least two to three weeks after

spawning to prevent negative effects on settlement. These months are also typically the peak period

for heat-stressed induced coral bleaching, and avoiding dredging during these months would

therefore have multiple benefits.

Finally, we showed that sediments collected near the Port contained higher abundances of

potentially pathogenic bacteria compared to Reef sediments, aligning with previous work

conducted in the vicinity of the Port Everglades inlet [65, 72]. These studies highlight the

importance of monitoring for potential contaminants in sediments and changes in the microbiome

which may result from dredging. In addition, although we only looked at the effects of suspended

sediments on coral larval performance, suspended sediments can act in concert with other stressors

typically associated with dredging. For example, [51] showed that elevated suspended sediment

concentrations, when simultaneously combined with reduced light levels, resulted in partial

mortality for multiple Pacific coral species. These findings emphasize the importance of

monitoring both suspended sediments and benthic light availability (PAR) during large-scale

dredging projects, and highlight the need for similar studies for Atlantic coral species.

Acknowledgements

The authors thank C. Sinigalliano, J. Hendee, B. Jensen, J. Destache, C. Pasparakis, and B. Young for help in the laboratory. Field assistance was provided by scientists from the NOAA Southeast Fisheries Science Center (D. Williams, A. Bright and A. Peterson). J. Wolfe provided GIS assistance with Fig 1. *O. faveolata* gametes were collected under permit FKNMS-2016-047-A1 to M.W. Miller. Sediment collections were determined to be exempt from requirements for an Environmental Resource Permit (ERP) by the Florida's Department of Environmental Protection (DeMinimus exemption). This research was carried out in part under the auspices of the Cooperative Institute for Marine and Atmospheric Studies (CIMAS), a cooperative institute of the University of Miami and the National Oceanic and Atmospheric Administration (NOAA), Cooperative Agreement NA 20OAR4320472. The scientific results and conclusions, as well as any opinions expressed herein, are those of the author(s) and do not necessarily reflect the views of NOAA or the Department of Commerce.

References

1.   Cunning R, Silverstein RN, Barnes BB, Baker AC. Extensive coral mortality and critical habitat loss following dredging and their association with remotely-sensed sediment plumes. Mar Pollut Bull. 2019;145:185-99. Epub 2019/10/09. doi: 10.1016/j.marpolbul.2019.05.027. PubMed PMID: 31590775.

2.   Foster T, Corcoran E, Erftemeijer P, Fletcher C, Peirs K, Dolmans C, et al. Dredging and port construction around coral reefs. PIANC Environmental Commission. Report, 2010.

3.   Miller MW, Karazsia J, Groves CE, Griffin S, Moore T, Wilber P, et al. Detecting sedimentation impacts to coral reefs resulting from dredging the Port of Miami, Florida USA. PeerJ. 2016;4:e2711. Epub 2016/11/30. doi: 10.7717/peerj.2711. PubMed PMID: 27896036; PubMed Central PMCID: PMCPMC5119242.

4.   Rosales SM, Sinigalliano C, Gidley M, Jones PR, Gramer LJ. Oceanographic habitat and the coral microbiomes of urban-impacted reefs. PeerJ. 2019;7:e7552. Epub 2019/10/01. doi: 10.7717/peerj.7552. PubMed PMID: 31565557; PubMed Central PMCID: PMCPMC6743471.

5.   Rushmore ME, Ross C, Fogarty ND. Physiological responses to short-term sediment exposure in
adults of the Caribbean coral Montastraea cavernosa and adults and recruits of Porites astreoides. Coral
Reefs. 2021;40(5):1579-91. doi: 10.1007/s00338-021-02156-0.
6.   Walker BK, Gilliam DS, Dodge RE, Walczak J. Dredging and shipping impacts on southeast Florida
coral reefs. Proceedings of the 12th International Coral Reef Symposium, Cairns, Australia, 9-13 July
2012. 2012.
7.   Jones R, Bessell-Browne P, Fisher R, Klonowski W, Slivkoff M. Assessing the impacts of sediments
from dredging on corals. Mar Pollut Bull. 2016;102(1):9-29. Epub 2015/12/15. doi:
10.1016/j.marpolbul.2015.10.049. PubMed PMID: 26654296.
8.   Weber M, Lott C, Fabricius KE. Sedimentation stress in a scleractinian coral exposed to
terrestrial and marine sediments with contrasting physical, organic and geochemical properties. Journal
of Experimental Marine Biology and Ecology. 2006;336(1):18-32. doi: 10.1016/j.jembe.2006.04.007.
9.   Jones R, Fisher R, Bessell-Browne P. Sediment deposition and coral smothering. PLoS One.
2019;14(6):e0216248. Epub 2019/06/20. doi: 10.1371/journal.pone.0216248. PubMed PMID: 31216275;
PubMed Central PMCID: PMCPMC6584000
10.   Storlazzi CD, Norris BK, Rosenberger KJ. The influence of grain size, grain color, and suspended-
sediment concentration on light attenuation: Why fine-grained terrestrial sediment is bad for coral reef
ecosystems. Coral Reefs. 2015;34(3):967-75. doi: 10.1007/s00338-015-1268-0.
11.   Weber M, de Beer D, Lott C, Polerecky L, Kohls K, Abed RM, et al. Mechanisms of damage to
corals exposed to sedimentation. Proc Natl Acad Sci U S A. 2012;109(24):E1558-67. Epub 2012/05/23.
doi: 10.1073/pnas.1100715109. PubMed PMID: 22615403; PubMed Central PMCID: PMCPMC3386076.
12.   Erftemeijer PL, Riegl B, Hoeksema BW, Todd PA. Environmental impacts of dredging and other
sediment disturbances on corals: a review. Mar Pollut Bull. 2012;64(9):1737-65. Epub 2012/06/12. doi:
10.1016/j.marpolbul.2012.05.008. PubMed PMID: 22682583.
13.   Jones R, Ricardo GF, Negri AP. Effects of sediments on the reproductive cycle of corals. Mar
Pollut Bull. 2015;100(1):13-33. Epub 2015/09/20. doi: 10.1016/j.marpolbul.2015.08.021. PubMed PMID:
26384866.
14.   Piniak GA. Effects of two sediment types on the fluorescence yield of two Hawaiian scleractinian
corals. Mar Environ Res. 2007;64(4):456-68. Epub 2007/06/15. doi: 10.1016/j.marenvres.2007.04.001.
PubMed PMID: 17568664.
15.   Barnes BB, Hu C, Kovach C, Silverstein RN. Sediment plumes induced by the Port of Miami
dredging: Analysis and interpretation using Landsat and MODIS data. Remote Sensing of Environment.
2015;170:328-39. doi: 10.1016/j.rse.2015.09.023.
16.   Fisher R, Stark C, Ridd P, Jones R. Spatial Patterns in Water Quality Changes during Dredging in
Tropical Environments. PLoS One. 2015;10(12):e0143309. Epub 2015/12/03. doi:
10.1371/journal.pone.0143309. PubMed PMID: 26630575; PubMed Central PMCID: PMCPMC4667927
17.   Eggleton J, Thomas KV. A review of factors affecting the release and bioavailability of
contaminants during sediment disturbance events. Environ Int. 2004;30(7):973-80. Epub 2004/06/16.
doi: 10.1016/j.envint.2004.03.001. PubMed PMID: 15196845.
18.   Jones RJ. Spatial patterns of chemical contamination (metals, PAHs, PCBs, PCDDs/PCDFS) in
sediments of a non-industrialized but densely populated coral atoll/small island state (Bermuda). Mar
Pollut Bull. 2011;62(6):1362-76. Epub 2011/05/10. doi: 10.1016/j.marpolbul.2011.01.021. PubMed
PMID: 21549399.
19.   Su SH, Pearlman LC, Rothrock JA, Iannuzzi TJ, Finley BL. Potential long-term ecological impacts
caused by disturbance of contaminated sediments:a case study. Environ Manage. 2002;29(2):234-49.
Epub 2002/01/30. doi: 10.1007/s00267-001-0005-3. PubMed PMID: 11815826.
20.   Hodgson G. Tetracycline reduces sedimentation damage to corals. Marine Biology.
1990;104:493-6.

21.   Voss JD, Richardson LL. Coral diseases near Lee Stocking Island, Bahamas: patterns and potential
drivers. Diseases of Aquatic Organisms. 2006;69(1):33-40.
22.   Studivan MS, Rossin AM, Rubin E, Soderberg N, Holstein DM, Enochs IC. Reef Sediments Can Act
As a Stony Coral Tissue Loss Disease Vector. Frontiers in Marine Science. 2022;8. doi:
10.3389/fmars.2021.815698.
23.   Tuttle LJ, Donahue MJ. Effects of sediment exposure on corals: a systematic review of
experimental studies. Environ Evid. 2022;11(1):4. Epub 2022/02/15. doi: 10.1186/s13750-022-00256-0.
PubMed PMID: 35154667; PubMed Central PMCID: PMCPMC8818373.
24.   Ricardo GF, Jones RJ, Negri AP, Stocker R. That sinking feeling: Suspended sediments can
prevent the ascent of coral egg bundles. Sci Rep. 2016;6:21567. Epub 2016/02/24. doi:
10.1038/srep21567. PubMed PMID: 26898352; PubMed Central PMCID: PMCPMC4761919.
25.   Ricardo GF, Jones RJ, Clode PL, Humanes A, Negri AP. Suspended sediments limit coral sperm
availability. Sci Rep. 2015;5:18084. Epub 2015/12/15. doi: 10.1038/srep18084. PubMed PMID:
26659008; PubMed Central PMCID: PMCPMC4677285.
26.   Ricardo GF, Jones RJ, Clode PL, Humanes A, Giofre N, Negri AP. Sediment characteristics
influence the fertilisation success of the corals Acropora tenuis and Acropora millepora. Mar Pollut Bull.
2018;135:941-53. Epub 2018/10/12. doi: 10.1016/j.marpolbul.2018.08.001. PubMed PMID: 30301119.
27.   Babcock R, Davies P. Effects of sedimentation on settlement of Acropora millepora. Coral reefs.
1991;9:205-8.
28.   Babcock R, Smith L, editors. Effects of sedimentation on coral settlement and survivorship.
Proceedings of the Ninth International Coral Reef Symposium, Bali, 23-27 October 2000; 2000: Citeseer.
29.   Perez K, 3rd, Rodgers KS, Jokiel PL, Lager CV, Lager DJ. Effects of terrigenous sediment on
settlement and survival of the reef coral Pocillopora damicornis. PeerJ. 2014;2:e387. Epub 2014/06/03.
doi: 10.7717/peerj.387. PubMed PMID: 24883248; PubMed Central PMCID: PMCPMC4034646.
30.   Ricardo GF, Jones RJ, Nordborg M, Negri AP. Settlement patterns of the coral Acropora
millepora on sediment-laden surfaces. Sci Total Environ. 2017;609:277-88. Epub 2017/07/28. doi:
10.1016/j.scitotenv.2017.07.153. PubMed PMID: 28750231.
31.   Speare KE, Duran A, Miller MW, Burkepile DE. Sediment associated with algal turfs inhibits the
settlement of two endangered coral species. Mar Pollut Bull. 2019;144:189-95. Epub 2019/06/11. doi:
10.1016/j.marpolbul.2019.04.066. PubMed PMID: 31179987.
32.   Fourney F, Figueiredo J. Additive negative effects of anthropogenic sedimentation and warming
on the survival of coral recruits. Sci Rep. 2017;7(1):12380. Epub 2017/09/30. doi: 10.1038/s41598-017-
12607-w. PubMed PMID: 28959051; PubMed Central PMCID: PMCPMC5620051.
33.   Moeller M, Nietzer S, Schils T, Schupp PJ. Low sediment loads affect survival of coral recruits: the
first weeks are crucial. Coral Reefs. 2016;36(1):39-49. doi: 10.1007/s00338-016-1513-1.
34.   Fabricius KE. Effects of terrestrial runoff on the ecology of corals and coral reefs: review and
synthesis. Mar Pollut Bull. 2005;50(2):125-46. Epub 2005/03/02. doi: 10.1016/j.marpolbul.2004.11.028.
PubMed PMID: 15737355.
35.   Price N. Habitat selection, facilitation, and biotic settlement cues affect distribution and
performance of coral recruits in French Polynesia. Oecologia. 2010;163(3):747-58. Epub 2010/02/20.
doi: 10.1007/s00442-010-1578-4. PubMed PMID: 20169452; PubMed Central PMCID: PMCPMC2886133.
36.   Ritson-Williams R, Paul VJ, Arnold SN, Steneck RS. Larval settlement preferences and post-
settlement survival of the threatened Caribbean corals Acropora palmata and A. cervicornis. Coral Reefs.
2009;29(1):71-81. doi: 10.1007/s00338-009-0555-z.
37.   Birrell CL, McCook LJ, Willis BL. Effects of algal turfs and sediment on coral settlement. Mar
Pollut Bull. 2005;51(1-4):408-14. Epub 2005/03/11. doi: 10.1016/j.marpolbul.2004.10.022. PubMed
PMID: 15757739.

38.   Gilmour J. Experimental investigation into the effects of suspended sediment on fertilisation,
larval survival and settlement in a scleractinian coral. Marine Biology. 1999;135:451-62.
39.   Goh B, Lee C, editors. A study of the effect of sediment accumulation on the settlement of coral
larvae using conditioned tiles. Proceedings Eleventh International Coral Reef Symposium, Ft Lauderdale,
Florida; 2008.
40.   Jorissen H, Galand PE, Bonnard I, Meiling S, Raviglione D, Meistertzheim AL, et al. Coral larval
settlement preferences linked to crustose coralline algae with distinct chemical and microbial
signatures. Sci Rep. 2021;11(1):14610. Epub 2021/07/18. doi: 10.1038/s41598-021-94096-6. PubMed
PMID: 34272460; PubMed Central PMCID: PMCPMC8285400.
41.   Quinlan ZA, Ritson-Williams R, Carroll BJ, Carlson CA, Nelson CE. Species-Specific Differences in
the Microbiomes and Organic Exudates of Crustose Coralline Algae Influence Bacterioplankton
Communities. Front Microbiol. 2019;10:2397. Epub 2019/11/30. doi: 10.3389/fmicb.2019.02397.
PubMed PMID: 31781048; PubMed Central PMCID: PMCPMC6857149.
42.   Siboni N, Abrego D, Puill-Stephan E, King WL, Bourne DG, Raina J-B, et al. Crustose coralline
algae that promote coral larval settlement harbor distinct surface bacterial communities. Coral Reefs.
2020;39(6):1703-13. doi: 10.1007/s00338-020-01997-5.
43.   Sneed JM, Ritson-Williams R, Paul VJ. Crustose coralline algal species host distinct bacterial
assemblages on their surfaces. ISME J. 2015;9(11):2527-36. Epub 2015/04/29. doi:
10.1038/ismej.2015.67. PubMed PMID: 25918832; PubMed Central PMCID: PMCPMC4611515.
44.   NOAA CD. Endangered and threatened wildlife and plants: Final listing determinations on
proposal to list 66 reef-building coral species and to reclassify elkhorn and staghorn corals. Federal
Register. 2014:53851-4123.
45.   Vazquez C. Explosive report finds PortMiami dredging caused extensive coral reef damage. Local
10 News [Internet]. 2023 Sept 06. Available from:
https://www.local10.com/news/local/2023/09/06/explosive-report-finds-portmiami-dredging-caused-
extensive-coral-reef-
damage/#:~:text=MIAMI%20%E2%80%93%20A%20new%20federal%20report,along%20the%20port's%2
0entrance%20channel.
46.   Gintert BE, Precht WF, Fura R, Rogers K, Rice M, Precht LL, et al. Regional coral disease outbreak
overwhelms impacts from a local dredge project. Environ Monit Assess. 2019;191(10):630. Epub
2019/09/15. doi: 10.1007/s10661-019-7767-7. PubMed PMID: 31520148.
47.   May LA, Miller CV, Moffitt ZJ, Balthis L, Karazsia J, Wilber P, et al. Acute turbidity exposures with
Port of Miami sediments impact Orbicella faveolata tissue regeneration. Marine Pollution Bulletin.
2023;193:115217.
48.   Szmant AM, Miller MW, editors. Settlement preferences and post-settlement mortality of
laboratory cultured and settled larvae of the Caribbean hermatypic corals Montastraea faveolata and
Acropora palmata in the Florida Keys, USA. Proc 10th Int Coral Reef Symp; 2006.
49.   Miller MW. Post-settlement survivorship in two Caribbean broadcasting corals. Coral Reefs.
2014;33(4):1041-6. doi: 10.1007/s00338-014-1177-7.
50.   Wentworth CK. A scale of grade and class terms for clastic sediments. The journal of geology.
1922;30(5):377-92.
51.   Bessell-Browne P, Negri AP, Fisher R, Clode PL, Duckworth A, Jones R. Impacts of turbidity on
corals: The relative importance of light limitation and suspended sediments. Mar Pollut Bull.
2017;117(1-2):161-70. Epub 2017/02/07. doi: 10.1016/j.marpolbul.2017.01.050. PubMed PMID:
28162249.
52.   Pasparakis C, Mager EM, Stieglitz JD, Benetti D, Grosell M. Effects of Deepwater Horizon crude
oil exposure, temperature and developmental stage on oxygen consumption of embryonic and larval

mahi-mahi (Coryphaena hippurus). Aquat Toxicol. 2016;181:113-23. Epub 2016/11/10. doi:
10.1016/j.aquatox.2016.10.022. PubMed PMID: 27829195.
53.   Serrano XM, Miller MW, Hendee JC, Jensen BA, Gapayao JZ, Pasparakis C, et al. Effects of
thermal stress and nitrate enrichment on the larval performance of two Caribbean reef corals. Coral
Reefs. 2017;37(1):173-82. doi: 10.1007/s00338-017-1645-y.
54.   Yashchenko V, Fossen EI, Kielland ON, Einum S. Negative relationships between population
density and metabolic rates are not general. J Anim Ecol. 2016;85(4):1070-7. Epub 2016/03/13. doi:
10.1111/1365-2656.12515. PubMed PMID: 26970102.
55.   Apprill A, McNally S, Parsons R, Weber L. Minor revision to V4 region SSU rRNA 806R gene
primer greatly increases detection of SAR11 bacterioplankton. Aquatic Microbial Ecology.
2015;75(2):129-37. doi: 10.3354/ame01753.
56.   Kuznetsova A, Brockhoff PB, Christensen RHB. lmerTest Package: Tests in Linear Mixed Effects
Models. Journal of Statistical Software. 2017;82(13). doi: 10.18637/jss.v082.i13.
57.   Ludecke MD. Package sjPlot: Data Visualization for Statistics in Social Science. 2022.
58.   Palacio-Castro AM, Serrano XM. Data and code for "Sediment source and concentration
influence larval performance of the threatened coral Orbicella faveolata" submitted to PlosOne (Version
v1) [Data set]. Zenodo. 2023. doi: https://doi.org/10.5281/zenodo.8022069.
59.   Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, et al. Reproducible,
interactive, scalable and extensible microbiome data science using QIIME 2. Nat Biotechnol.
2019;37(8):852-7. Epub 2019/07/26. doi: 10.1038/s41587-019-0209-9. PubMed PMID: 31341288;
PubMed Central PMCID: PMCPMC7015180.
60.   Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet journal. 2011;17(1):10-2.
61.   Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJ, Holmes SP. DADA2: High-resolution
sample inference from Illumina amplicon data. Nat Methods. 2016;13(7):581-3. Epub 2016/05/24. doi:
10.1038/nmeth.3869. PubMed PMID: 27214047; PubMed Central PMCID: PMCPMC4927377.
62.   Bokulich NA, Kaehler BD, Rideout JR, Dillon M, Bolyen E, Knight R, et al. Optimizing taxonomic
classification of marker-gene amplicon sequences with QIIME 2's q2-feature-classifier plugin.
Microbiome. 2018;6(1):90. Epub 2018/05/19. doi: 10.1186/s40168-018-0470-z. PubMed PMID:
29773078; PubMed Central PMCID: PMCPMC5956843 The authors declare that they have no competing
interests. PUBLISHER'S NOTE: Springer Nature remains neutral with regard to jurisdictional claims in
published maps and institutional affiliations.
63.   Lin H, Peddada SD. Analysis of compositions of microbiomes with bias correction. Nat Commun.
2020;11(1):3514. Epub 2020/07/16. doi: 10.1038/s41467-020-17041-7. PubMed PMID: 32665548;
PubMed Central PMCID: PMCPMC7360769.
64.   Hodgson G. Sediment and the settlement of larvae of the reef coral Pocillopora damicornis.
Coral Reefs. 1990;9:41-3.
65.   Krausfeldt LE, Lopez JV, Bilodeau CM, Won Lee H, Casali SL. Change and stasis of distinct
sediment microbiomes across Port Everglades Inlet (PEI) and the adjacent coral reefs. PeerJ.
2023;11:e14288. Epub 2023/01/20. doi: 10.7717/peerj.14288. PubMed PMID: 36655050; PubMed
Central PMCID: PMCPMC9841897.
66.   Klaus JS, Janse I, Fouke BW. Coral black band disease microbial communities and genotypic
variability of the dominant cyanobacteria (CD1C11). Bulletin of Marine Science. 2011;87(4):795.
67.   Waite DW, Chuvochina M, Pelikan C, Parks DH, Yilmaz P, Wagner M, et al. Proposal to reclassify
the proteobacterial classes Deltaproteobacteria and Oligoflexia, and the phylum Thermodesulfobacteria
into four phyla reflecting major functional capabilities. Int J Syst Evol Microbiol. 2020;70(11):5972-6016.
Epub 2020/11/06. doi: 10.1099/ijsem.0.004213. PubMed PMID: 33151140.

68.   Nie S, Zhang Z, Mo S, Li J, He S, Kashif M, et al. Desulfobacterales stimulates nitrate reduction in
the mangrove ecosystem of a subtropical gulf. Sci Total Environ. 2021;769:144562. Epub 2021/01/19.
doi: 10.1016/j.scitotenv.2020.144562. PubMed PMID: 33460836.
69.   Clausing RJ, Annunziata C, Baker G, Lee C, Bittick SJ, Fong P. Effects of sediment depth on algal
turf height are mediated by interactions with fish herbivory on a fringing reef. Marine Ecology Progress
Series. 2014;517:121-9. doi: 10.3354/meps11029.
70.   Dobbelaere T, Muller EM, Gramer LJ, Holstein DM, Hanert E. Coupled Epidemio-Hydrodynamic
Modeling to Understand the Spread of a Deadly Coral Disease in Florida. Frontiers in Marine Science.
2020;7. doi: 10.3389/fmars.2020.591881.
71.   Precht WF, Gintert BE, Robbart ML, Fura R, van Woesik R. Unprecedented Disease-Related Coral
Mortality in Southeastern Florida. Sci Rep. 2016;6:31374. Epub 2016/08/11. doi: 10.1038/srep31374.
PubMed PMID: 27506875; PubMed Central PMCID: PMCPMC4979204.
72.   O'Connell L, Gao S, McCorquodale D, Fleisher J, Lopez JV. Fine grained compositional analysis of
Port Everglades Inlet microbiome using high throughput DNA sequencing. PeerJ. 2018;6:e4671. Epub
2018/05/16. doi: 10.7717/peerj.4671. PubMed PMID: 29761039; PubMed Central PMCID:
PMCPMC5947159.
73.   Abdel-Salam HA, Porter JW, Hatcher B, editors. Physiological effects of sediment rejection on
photosynthesis and respiration in three Caribbean reef corals. Proc 6th int coral Reef Symp; 1988.
74.   Okubo N, Yamamoto HH, Nakaya F, Okaji K. Oxygen consumption of a single embryo/planula in
the reef-building coral Acropora intermedia. Marine Ecology Progress Series. 2008;366:305-9. doi:
10.3354/meps07562.
75.   Brainard R, Birkeland C, Eakin CM, McElhany P, Miller MW, Patterson M, et al. Status review
report of 82 candidate coral species petitioned under the US Endangered Species Act: Citeseer; 2011.
76.   Hughes TP, Tanner JE. Recruitment failure, life histories, and long⁠⁠‐term decline of Caribbean
corals. Ecology. 2000;81(8):2250-63.
77.   Protection AE. Technical Guidance: Environmental Impact Assessment of Marine Dredging
Proposals. Environmental Protection Authority, Perth, Western Australia. 2016.
78.   Szmant AM. Reproductive ecology of Caribbean reef corals. Coral reefs. 1986;5:43-53.
79.   Szmant AM. Sexual reproduction by the Caribbean reef corals Montastrea annularis and M.
cavernosa. Mar Ecol Prog Ser. 1991;7(4):13-25.

Supporting information

S1 Fig. Composite display of grain size analysis of sediment samples. Two replicates per

sediment type (Port versus Reef) were used.

S2 Fig. Example plot of oxygen consumption over time for 6 *Orbicella faveolata* larvae

placed in a single well of the microplate. The R^2 value is given for the linear trend. The first

10–20% (pO2 in % air saturation) linear decreases in oxygen were used to calculate respiration

rates per larva. Any portion of the slope that dipped below 70% air saturation was not used for

data analysis.

S3 Fig. Odds ratios (OR) for the association between *Orbicella faveolata* larvae exposure to

sediment treatments. (A) larval survival after 24 h of treatments, and (B) larval settlement after

one week of recovery from experimental treatments. OR=1 Exposure does not affect odds of

outcome, OR>1 Exposure associated with higher odds of outcome, and OR<1 Exposure

associated with lower odds of outcome.

S4 Fig. Respiration rates (mean ± 95 CI) in *Orbicella faveolata* larvae after a 24 h exposure to

experimental treatments (left panel) and after one week of recovery from experimental

treatments (right panel). White dots denote individual replicate wells.

S1 Table. Size fraction analysis of sediment samples. Values correspond to graphic in S1 Fig.

S2 Table. Generalized linear mixed-effect models outputs for *Orbicella faveolata* larvae

performance. DF: Degrees of freedom

S3 Table. Results of the final generalized linear mixed-effect models for *Orbicella faveolata*

performance. A) larval survival after 24 h of treatments, and B) larval settlement after one week

of recovery from experimental treatments (shown in Fig 1). The coefficient estimates describe

the change in the log odds for each treatment level compared to the base level (control).

Asterisks denote significant differences in a sediment treatment compared to controls with no

sediment added (*: $p < 0.05$; **: $p < 0.01$; ***: $p < 0.001$).

S4 Table. Linear mixed-effect model outputs for *Orbicella faveolata* larvae respiration.

Figure 1

Figure 2

Figure 3

Figure 4