Prrx1-driven LINC complex disruption _in vivo_ does not significantly alter bone properties

in 8-week male mice nor after 6-weeks voluntary wheel running

Scott Birks[1], Sean Howard[2], Christian S. Wright[3], Caroline O'Rourke[4], Anthony Lau[4], William R.

Thompson[3], Gunes Uzer[2] [†]

[1]Boise State University, Micron School of Materials Science and Engineering

[2]Boise State University, Mechanical and Biomedical Engineering

[3]Indiana University, Department of Physical Therapy, School of Health and Human Sciences

[4]The College of New Jersey, Biomedical Engineering

Word count: 3277

[†]**Corresponding author:**

Gunes Uzer PhD

Boise State University

Department of Mechanical & Biomedical Engineering

1910 University Drive, MS-2085

Boise, ID 83725-2085

Ph. (208) 426-4461

Email: gunesuzer@boisestate.edu

<u>Abstract</u>

The Linker of Nucleoskeleton and Cytoskeleton (LINC) complex serves to connect the nuclear

envelope and the cytoskeleton, influencing cellular processes such as nuclear arrangement,

architecture, and mechanotransduction. The role LINC plays in mechanotransduction pathways

in bone progenitor cells has been well studied; however, the mechanisms by which LINC

complexes govern *in vivo* bone formation remain less clear. To bridge this knowledge gap, we

established a murine model disrupting LINC using transgenic Prx-Cre mice and floxed Tg(CAG-

LacZ/EGFP-KASH2) mice. Prx-Cre mice express the Cre recombinase enzyme controlled by

the paired-related homeobox gene-1 promoter, a pivotal regulator of skeletal development.

Tg(CAG-LacZ/EGFP-KASH2) mice carry a lox-stop-lox flanked LacZ gene allowing for the

overexpression of an EGFP-KASH2 fusion protein via cre recombinase mediated deletion of the

LacZ cassette. This disrupts endogenous Nesprin-Sun binding in a dominant negative manner

disconnecting nesprin from the nuclear envelope. By combining these lines, we generated a

Prrx1(+) cell-specific LINC disruption model to study its impact on the developing skeleton and

subsequently exercise-induced bone accrual. The findings presented here indicate Prx-driven

LINC disruption (PDLD) cells exhibit no change in osteogenic and adipogenic potential

compared to controls *in vitro* nor are there bone quality changes when compared to in sedentary

animals at 8 weeks. Although PDLD animals displayed increased voluntary running activity, a 6-

week exercise intervention did not significantly alter bone microarchitecture or mechanical

properties.

Key Words: Paired-related homeobox gene-1 (Prrx1), Mechanical Signals, Mesenchymal Stem

Cells, Mechanobiology, LINC, Bone

Introduction

Bone is a highly dynamic tissue influenced by both biochemical and mechanical stimuli to retain proper structure and function [1]. While the anabolic effects of mechanical challenge (i.e. weight-bearing exercise) on bone quality is well accepted, how Linker of Nucleoskeleton and Cytoskeleton (LINC) complexes may affect bone progenitor cell response to these signals remains underexplored.

LINC complexes serve as a physical link between the cytoskeleton and the nucleoskeleton, facilitating the transmission of mechanical forces and signals between the extracellular matrix and the nucleus [2]. LINC complexes are comprised of cytoskeletal nesprins (nuclear envelope spectrin repeats) and perinuclear Sun (Sad1/Unc84) proteins with the N-terminal calponin homology domains of nesprins binding to cytoskeletal F-actin microfilaments and the evolutionarily conserved C-terminal KASH (Klarsicht/Anc-1, Syne Homology) domains binding to Sun domains within the perinuclear space. Sun proteins directly engage with the intranuclear lamina, completing the connection from the cytoskeleton to the nucleus [3-5]. Beyond their structural role, LINC complexes have emerged as critical components in transducing mechanical signals within cells [6].

Earlier *in vitro* studies have effectively illustrated the dynamic involvement of LINC complexes within mechanotransduction pathways [6, 7]. For instance, disabling LINC complexes through RNA interference against SUN1 and SUN2 proteins, or transfecting cells with a dominant negative Nesprin KASH domain, leads to the suppression of FAK and Akt phosphorylation. This, in turn, circumvents the mechanical repression of adipogenesis when mesenchymal stromal cells are exposed to low-magnitude, high-frequency signals [6]. While the role of LINC complexes in mechanotransduction has been demonstrated *in vitro*, their role in orchestrating

the mechanical regulation of bone in an *in vivo* remains largely uncharted. The current study

aims to implement an *in vivo* strategy to investigate the ramifications of LINC complex disruption

during mechanical loading, specifically within bone progenitor cells. This was achieved by

creating a cre/lox Prrx1-driven LINC disruption (PDLD) murine model.

The paired-related homeobox gene-1 (*Prrx1*) is expressed in early limb bud mesenchyme,

playing a crucial role in regulating skeletal development and is also found in adult

undifferentiated mesenchyme and periosteum [7, 8]. Thus, using transgenic Prrx1-cre mice we

are able to target skeletal tissues throughout development and into adulthood. LINC disruption

in Prx(+) cells was then achieved by crossing these mice with floxed mice created by Razafsky

and Hodzic that overexpress an EGFP-KASH2 fusion protein upon Cre-recombination [9, 10].

This fusion protein binds to perinuclear Sun proteins in a dominant negative manner, causing

displacement of cytoplasmic nesprins into the endoplasmic reticulum and thus, disrupting LINC

connectivity between the nucleus and cytoskeleton.

While there are several modalities for mechanical stimulation *in vivo*, exercise is considered a

potent modality that can trigger adaptive responses in bone tissue, enhancing bone density and

strength [11]. A variety of exercise models including forced/trained treadmill running,

voluntary/forced wheel running, swimming, and resistance training (loaded/unloaded ladder

climbing) have been utilized in rodent studies, all of which have been shown to improve bone

mass, bone strength, and bone mineral density [12] [13]. We chose voluntary wheel running for

the present study to reduce the amount of stress on the animals.

To interrogate the effect of disabling LINC complex function in Prx positive bone cell

progenitors, we created a LINC disruption model under the Prx-Cre driver and measured

primary cell differentiation potential *in vitro*, bone microarchitecture via micro-computed

tomography (micro-CT) and mechanical properties via 3-point bending at an 8-week baseline.

We then subjected a cohort of animals to either a running or non-running intervention starting at

8-weeks of age and lasting for 6 weeks with a 1-week acclimation period. Micro-CT and

biomechanical properties were also measured post-exercise to assess any effects of LINC

disruption on exercise induced bone accrual.

<u>Materials and Methods</u>

Animals

Mouse strains included B6.Cg-Tg(Prrx1-cre))1Cjt/J aka *Prx-Cre* (The Jackson Laboratory, Stock

Number: 005584)[7], Tg(CAG-LacZ/EGFP-KASH2) aka *KASH2* (Hodzic, Wash U)[9], and

B6.Cg-Gt(ROSA)26Sor$^{tm14(CAG-tdTomato)Hze}$/J aka *Ai14* (generously provided by Dr. Richard

Beard)[14].

Hemizygous *Prx-Cre* mice were crossed with floxed *KASH2* mice to generate Prrx1-driven LINC

disrupted (PDLD) murine model. LINC disruption mechanism described by Razafsky and Hodzic

[9, 10]. Briefly, upon cre recombination, a LacZ containing lox-stop-lox cassette upstream of an

eGFP-KASH2 ORF is excised allowing for the overexpression an eGFP-KASH2 fusion protein

which saturates available Sun/KASH binding in a dominant-negative manner. Genotyping was

performed on tissue biopsies via real-time PCR probing for LacZ and Cre (Transnetyx, Cordova,

TN) to determine experimental animals and controls. Cre(+)/LacZ(+) animals were considered

experimental and all other genotypes were considered controls. Animals were housed in

individually ventilated cages under controlled conditions until placement in conventional cages

for the duration of the exercise intervention. *Ad libitum* access to food and water allowed.

Additionally, *Prx-Cre* mice were crossed with Ai14 RFP reporter mice to assess Prx-Cre activity

in skeletal tissue and bone marrow cells. All procedures approved by Boise State University

Institutional Animal Care and Use Committee.

Bone Marrow Cell Isolation, Expansion, and Differentiation

Bone marrow cells were isolated from tibiae and femora via centrifugation as previously

described [15]. Briefly, tibia and femur were dissected, cleared of all soft tissue, and centrifuged

at >12,000g to remove the bone marrow. This bone marrow pellet was then resuspended in

media and plated in supplemented growth media (alpha-MEM +20% FBS, +100U/mL penicillin,

+100µg/mL streptomycin). Cells were washed with PBS after 48 hours to remove non-adherent

cells and allowed to proliferate to 50% confluency. At this confluency, cells were lifted, split onto

two separate plates to produce enough cells for assays, and allowed to proliferate to ~80%

confluency. Cells were then lifted and plated for differentiation assays.

For osteogenic assays (n=2/group), cells were seeded at 200K per well on 6-well plates in

normal growth media for 24 hours prior to switching to osteogenic media (alpha-MEM with 20%

FBS, 100U/mL penicillin, 100µg/mL streptomycin, 10mM beta-glycerophosphate, 50µg/µL

ascorbic acid) for 14 days.

For adipogenic assays (n=2/group), cells were seeded at 150K per well on 6-well plates in

normal growth media for 24 hours prior to switching to adipogenic media (alpha-MEM with 10%

FBS, 100U/mL penicillin, 100µg/mL streptomycin, 5µg/mL insulin, 0.1µM dexamethasone, 50µM

indomethacin) for 7 days.

Immunofluorescence staining and microscopy

Nesprin staining was performed on bone marrow cultures from 12-week Prx-Cre(+)/eGFP-

KASH2(+) and aged-matched control mice (n=2/group). After isolation and culture as described

above, cells were rinsed with PBS, fixed in 4% paraformaldehyde, permeabilized with 0.3%

Triton buffer and blocked with 5% goat serum (Jackson ImmunoResearch 005-000-001) prior to

being tagged with a rabbit anti-nesprin2 antibody (1:300; ImmuQuest IQ565). After primary

antibody tagging, cells were incubated with an anti-rabbit secondary antibody (1:300 AlexaFluor

594nm goat anti-rabbit, Invitrogen A11037), nuclei were stained with Hoechst 33342

(Invitrogen™ R37605) and visualized via fluorescent microscopy (Zeiss LSM 900). No

additional staining was needed to visualize endogenous eGFP.

Oil Red O and Alizarin Red Staining

After 7 days in adipogenic media, cultures were stained against Oil Red O as described

previously [16]. Briefly, cells were rinsed with PBS, fixed in 10% neutral buffered formalin for 30

mins at room temperature, and washed with deionized water. Cultures were then incubated in

100% propylene glycol, stained with 0.5% Oil Red O in propylene glycol, and finally incubated in

85% propylene glycol. Unbound dye was removed by rinses with tap water until clear and

images were taken. Oil Red O stain was then eluted with 100% propan-2-ol and optical

absorbance values were read on a microplate spectrophotometer (ThermoFisher Multiskan GO)

at 510nm.

After 21 days in osteogenic media, cells were rinsed in PBS, fixed with 70% ethanol, and

stained with 40mM Alizarin Red solution for visualization of calcium depsosits. The stain was

then eluted using 10% w/v cetylpyridium chloride (CPC) and diluted 4-fold to avoid absorbance

saturation. Absorbance values were read on a microplate spectrophotometer (ThermoFisher

Multiskan GO) at 562nm.

Exercise Regimen

Eight-week old male PDLD or control mice were randomly assigned to a voluntary wheel

running exercise intervention or a non-running control group without access to a running wheel

(n≥10/group). Animals were allotted a one-week acclimation period in these new cages followed

by a six-week tracked exercise intervention. All mice were individually housed and running

metrics including elapsed time, total distance, average speed, and maximum speed were

measured with a cyclocomputer (CatEye Velo 7) and recorded daily. Ad libitum access to food

and water was allowed. Body mass and food intake was also tracked weekly.

Micro-computed tomography

Proximal tibiae stored frozen in PBS soaked gauze prior to scanning and were scanned

immersed in PBS using an x-ray microtomograph (SkyScan 1172F). Settings as follows: power

of 55kV/181mA, a voxel size of $10.33\mu m^3$, and 230ms integration time. The volume of interest

for trabecular quantitative analysis started 10 slices distal to the proximal physis extending 1

mm. The cortical volume of interest started 2.15mm distal to the physis, extending another

0.5mm. Image reconstruction was performed with NRecon software (Bruker). After any

necessary image reorientation, quantitative analysis was performed with CTAn (Bruker) and 3D

image rendering was performed with CTVol. Reported variables and nomenclature used

according to ASBMR guidelines [17].

Three-point Bend Mechanical Testing

3-point bending mechanical testing was performed using an Instron Universal Testing System

model 5967. Force and deflection data was collected and ran through a MATLAB code in order

to determine the mechanical properties of the bone (i.e. linear flexural stiffness, maximum yield

force, and maximum force).

Statistical Analysis

Results are expressed as mean ± standard deviation. Significance was determined by 2-tailed

unpaired student's t-test or 2-way ANOVA with Tukey post-hoc test (Prism).

Results

Expression of EGFP-KASH2 fusion protein physically disrupts LINC complex in Prrx1 positive

cells

We crossed Prx-Cre(+) mice with the Ai14 RFP reporter line to visualize Cre activity in skeletal

tissues and bone marrow cells. P7 pups were placed in an *in vivo* fluorescent scanner (IVIS®

Spectrum) and notable red fluorescent in the appendicular skeleton and calvaria was visible

(Figure 1A), consistent with previous reports [18]. This confirms Cre recombinase activity is

present in Prrx1(+) tissues. We then isolated bone marrow from these animals at 12 weeks and

successfully visualized RFP containing cells within the bone marrow (Figure 1B) providing

evidence for the presence of Prrx1 in bone marrow.

A Prx-driven LINC disruption (PDLD) model was then generated by crossing hemizygous Prx-

Cre(+) mice with floxed Tg(CAG-LacZ/EGFP-KASH2) mice. The mechanism for LINC disruption

in this model relies on Cre recombinase mediated excision of the lox-stop-lox LacZ cassette in

the *KASH2* mice allowing for the overexpression of an eGFP-KASH2 fusion protein. This fusion

protein then binds to perinuclear Sun proteins, displacing nesprins from the nuclear envelope.

To confirm this model disrupts LINC function via nesprin displacement, we flushed bone marrow

from 12-week old PDLD and control animals (N=2/group) to check for cells expressing

endogenous eGFP localized to the nuclear envelope, indicating the overexpression of the

eGFP-KASH2 protein. We additionally immunostained these cells with an anti-nesprin2 antibody

and secondary fluorophore to confirm nesprin detachment from the nuclear envelope. As seen

in Figure 1C, there is a clear nuclear halo of eGFP fluorescence in PDLD mice compared to no

eGFP visualized in controls. Nesprin-2 staining also has a punctate and scattered appearance

in PDLD mice compared to the clear nuclear positioning seen in controls indicating successful

displacement of nesprin from the nuclear envelope where eGFP is present. These data provide

evidence of successful LINC disruption in cells expressing eGFP.

Prrx1-driven LINC disruption does not change adipogenic and osteogenic differentiation

potential from LINC-intact controls

To determine the differentiation potential of LINC disrupted cells, bone marrow was flushed from

tibia and femur of 12-week male PDLD and negative control mice (2 animals per group, 3

technical replicates per animal), and plated for osteogenic/adipogenic differentiation assays

(Figure 2). After 7 days, adipogenic cultures were stained with Oil Red O and eluted stain was

quantified via spectrophotometry at 510nm. After 14 days, osteogenic cultures were stained with

Alizarin Red and eluted stain was quantified via spectrophotometry at 562nm.

Relative absorbance values of Oil Red O dye from PDLD adipogenic cultures was 24.8% lower

than adipogenic LINC-intact controls (p=0.15) and 120.4% greater than undifferentiated controls

(p<0.05). Relative absorbance values of Alizarin Red stain from PDLD osteogenic cultures was

5.0% less than that of LINC intact osteogenic controls (p=0.8), and 531.4% greater than

undifferentiated LINC-intact controls (p<0.01). This suggests a trend towards diminished

adipogenic differentiated potential in PDLD cells compared to LINC-intact controls, but no

appreciable difference in osteogenic potential.

Bone microarchitecture unchanged in PDLD mice compared to LINC-intact controls at 8-week

baseline

Prior to beginning the exercise intervention, we established baseline bone microarchitecture

measurements via micro-computed tomography (micro-CT) using an 8-week cohort of mice

covering all four possible genotypes (n=3/grp): Prx-Cre(+)/EGFP-KASH2(+), Prx-Cre(+)/EGFP-

KASH2(-), Prx-Cre(-)/EGFP-KASH2(+), Prx-Cre(-)/EGFP-KASH2(-). There was no significant

difference found between genotypes (data not shown), so Prx-Cre(+)/EGFP-KASH2(+) mice

remained our experimental model while all other genotypes were indiscriminately used as

controls.

Each mouse from this cohort was weighed and both tibiae and femora were harvested from Prx-

Cre(+)/EGFP-KASH2(+) and control (n=10/group). Each tibia was cleared of soft tissue for

micro-CT analysis. We noted no chance in body weight in PDLD mice vs. control recorded at

21.59 ± 1.7 g and 21.20 ± 0.77 g (p=0.68) respectively. A trend towards decreased bone

microarchitecture with a 5.6% decrease in BV/TV (p=0.30), 5.29% decrease in trabecular

thickness (p=0.19), and 6.10% decrease in cortical bone area (p=0.12) in PDLD mice was

observed between control and PDLD mice (Figure 3).

Prrx1-driven LINC disrupted mice run longer, farther, and faster than controls

A second cohort of 8-week male mice were given voluntary access to a running wheel (or no

wheel and allowed a one-week acclimation period before beginning the six-week tracked

exercise intervention. Tracked running metrics including elapsed time, average distance,

average velocity, and max speed were recorded daily. Our data indicate that PDLD mice ran

significantly longer, farther, and faster than their LINC intact counterparts with an average run

time of 169.2 ± 6.8 mins compared to 146.6 ± 9.4 mins (p<0.001), an average distance of 2.35 ±

0.16 km compared to 1.82 ± 0.1 km (p<0.0001), and an average velocity was 0.53 ± 0.05 km/hr

for PDLD mice and 0.48 ± 0.02 km/hr for controls (p<0.05).

LINC disruption does not significantly affect bone microarchitecture or mechanical properties in

PDLD mice subjected to six-week running protocol

Bone microarchitecture was measured via micro-CT to assess any effects of LINC disruption on

exercise-induced bone accrual in the proximal tibial metaphysis and diaphysis for trabecular and

cortical parameters, respectively. There was no notable difference in trabecular bone fraction

(BV/TV), trabecular thickness (Tb.Th.), trabecular separation (Tb.Sp.), trabecular number

(Tb.N), total cortical tissue area (Tt.Ar.), or cortical thickness (Ct.Th.) between any of the

groups. There was, however, a 10.5% decrease (p<0.05) in cortical bone area (Ct.Ar.) in

PDLD+Ex compared to sedentary LINC-intact controls and a 6.15% decrease (p<0.05) in

cortical area fraction (Ct.Ar./Tt.Ar.) in PDLD+Ex compared to PDLD-Ex. No significant changes

in measured bone mechanical properties were noted (Figure 5).

<u>Discussion</u>

While the *in vitro* role of LINC complex in mechanotransduction pathways in bone progenitor cells has been well studied, its function *in vivo* remains underexplored. Here we sought to elucidate if LINC complex disruption in *Prx(+)* cells influences bone marrow stromal cell phenotype and compromises bone tissue quality throughout development and during bone response to voluntary wheel running. After successfully generating a LINC disruption mouse model under the Prrx1-Cre driver, we found no change in osteogenic and adipogenic differentiation potential in PDLD cells, nor did we find any bone architecture changes in PDLD mice at an 8-week baseline. There were also no remarkable differences in bone microarchitecture or mechanical properties between PDLD mice subjected to a six-week exercise intervention compared to LINC intact animals or sedentary controls.

Studies involving LINC complex disruption via KASH overexpression *in vitro* show decreases osteogenic activity in human bone marrow mesenchymal stem cells by reducing Runx2 via increased histone acetylation [19]. We recently reported that LINC disruption also decreases adipogenic potential *in vitro* [20]. Despite this evidence, our results indicate no change in adipogenic and osteogenic potential of these cells when extracted from bone marrow. To exclude the possibility of these cells not surviving the marrow extraction we first we confirmed that GFP-KASH, expressing, LINC-deficient cells were present in our cultures after marrow extraction. Shortly, we harvested cells from female Prx-Cre(+)/Ai14(+)/eGFP-KASH2(+) animals that express eGFP and tdTomato in Prx(+) cells. Cells were visualized for green and red fluorescence. Average cell counts from 24 representative images with 683 total cells yielded 98.07 ± 7.2% red cells and 86.3 ± 14.52 % green cells indicating that eGFP- KASH2 expressing cells were present in our cultures (Fig. S1 A/B). Next to confirm participation of LINC-deficient cells in osteogenic nodules we have differentiated these cultures into osteogenic lineage and stained with calcein blue. Our results in Fig.S1 C confirms that Prx-Cre(+)/eGFP-KASH2(+) cells

were present within calcein blue stained nodules (Fig. S1 C). Finally, it is worth noting that the

duration of *in vitro* studies involving KASH overexpression is relatively short (1-2 days to 1

week), while the animals in this study express the dominant negative KASH proteins chronically

from conception. To test the possible effect of acute versus chronic KASH protein expression,

we utilized transient transfection with a previously published dominant negative form of the

nesprin KASH domain (DNK)[21, 22]. RNA-seq results from transient DNK transfection show

decreased osteogenic gene transcripts *in vitro* when compared to empty mCherry plasmid

transfected controls *(*Fig.S2*)*. It is not clear why we did not see any effect on cell phenotypes

from animals, it is possible bone cells in our animals may have adapted to this chronic

expression of eGFP-KASH fusion protein.

Prrx1 is a well-established marker for early limb bud development [7, 8], and this study provides

valuable insights into the impact of LINC disruption across various developmental stages, from

embryogenesis through adolescence into adulthood. Interestingly, our findings suggest that this

disruption has no significant influence on the differential potential of bone marrow cells or whole

bone structural or mechanical properties. One possible explanation for this observation is that

cells with LINC disruption might be selectively excluded from the population of bone-forming

cells, possibly directed towards apoptotic pathways. However, it's worth noting that we do detect

Prx1-Cre activity in adult bone marrow cell aspirates, indicating that these cells are still present

in the adult skeletal system. Yet, their precise function and role in bone biology remain unclear.

Some studies propose that cells originating from the Prx1 lineage may not be actively involved

in the homeostatic mechanisms or bone remodelling [18]. This raises questions about the

specific contributions of these cells within the intricate landscape of bone development and

maintenance, prompting further investigation into their functions and potential implications for

bone health.

Similar voluntary wheel running studies also with a six-week intervention but using older (16

week) female mice found increased trabecular quality and quantity post-exercise in animals on

a regular diet [23] [24]. While bones of male mice also shown to respond to voluntary wheel

running [25] [26] [27], other studies using male mice reported minimal effects of exercise on

bone after 10 or 21 weeks of voluntary wheel running [28, 29]. Compared to reported daily

running metrics in these studies (ranging from 4-10 km/day), our mice ran less on average (2.1

km/day). The reason for this is unknown, but may have affected our outcomes. Furthermore, our

mice were 8-weeks old when placed in the exercise intervention, which is before the 12-week

reported age of skeletal maturity [28, 30]. Therefore, it is possible that mechanical challenge

provided by voluntary wheel running did not affect the ongoing bone growth in these mice.

Further studies using different modalities of LINC disruption (i.e. SUN & Nesprin depletion

models) or the use of different mesenchymal/osteochondroprogenitor cre drivers (i.e. Dermo1 or

Sox-9) would shed further light on the impact of impaired mechanotransduction in skeletal

stem/progenitor cells.

Acknowledgements

This study was supported by AG059923, 1S10OD032354, AR075803, AR074473,

P20GM109095, NSF1929188, 2025505 and TCNJ Mentored Undergraduate Summer

Experience. Authors declare no competing interests financial or otherwise.

Author Contributions: All authors have read and approved the final submitted manuscript.

Scott Birks: concept/design, data analysis/interpretation, manuscript writing.

Sean Howard: data analysis/interpretation, manuscript writing.

Caroline O'Rourke: data analysis/interpretation, manuscript writing.

Matthew Goelzer: data analysis/interpretation, manuscript writing.

William R Thompson: data analysis/interpretation, financial support, manuscript writing.

Anthony Lau: data analysis/interpretation, financial support, manuscript writing.

Gunes Uzer: concept/design, data analysis/interpretation, financial support, manuscript writing.

References

1.   Gusmão, C.V. and W.D. Belangero, *HOW DO BONE CELLS SENSE MECHANICAL*

LOADING? Rev Bras Ortop, 2009. **44**(4): p. 299-305.

2.   Bouzid, T., et al., *The LINC complex, mechanotransduction, and mesenchymal stem cell*

function and fate. Journal of Biological Engineering, 2019. **13**(1): p. 68.

3.   Crisp, M., et al., *Coupling of the nucleus and cytoplasm: role of the LINC complex.* J Cell

Biol, 2006. **172**(1): p. 41-53.

4.   Padmakumar, V.C., et al., *The inner nuclear membrane protein Sun1 mediates the*

anchorage of Nesprin-2 to the nuclear envelope. J Cell Sci, 2005. **118**(Pt 15): p. 3419-

30.

5.   Haque, F., et al., *SUN1 interacts with nuclear lamin A and cytoplasmic nesprins to*

provide a physical connection between the nuclear lamina and the cytoskeleton. Mol Cell

Biol, 2006. **26**(10): p. 3738-51.

6.   Birks, S. and G. Uzer, *At the nuclear envelope of bone mechanobiology.* Bone, 2021.

151: p. 116023.

7.   Logan, M., et al., *Expression of Cre Recombinase in the developing mouse limb bud*

driven by a Prxl enhancer. Genesis, 2002. **33**(2): p. 77-80.

8.   Martin, J.F. and E.N. Olson, *Identification of a prx1 limb enhancer.* Genesis, 2000. **26**(4):

p. 225-9.

9.   Razafsky, D. and D. Hodzic, *Temporal and tissue-specific disruption of LINC complexes*

in vivo. Genesis, 2014. **52**(4): p. 359-65.

10.   Razafsky, D., C. Potter, and D. Hodzic, *Validation of a Mouse Model to Disrupt LINC*
413   *Complexes in a Cell-specific Manner.* J Vis Exp, 2015(106): p. e53318.

414   11.   Benedetti, M.G., et al., *The Effectiveness of Physical Exercise on Bone Density in*
415   *Osteoporotic Patients.* Biomed Res Int, 2018. **2018**: p. 4840531.

416   12.   Hastings, M.H., et al., *Animal Models of Exercise From Rodents to Pythons.* Circulation
417   Research, 2022. **130**(12): p. 1994-2014.

418   13.   Guo, S., et al., *Impacts of exercise interventions on different diseases and organ*
419   *functions in mice.* J Sport Health Sci, 2020. **9**(1): p. 53-73.

420   14.   Madisen, L., et al., *A robust and high-throughput Cre reporting and characterization*
421   *system for the whole mouse brain.* Nat Neurosci, 2010. **13**(1): p. 133-40.

422   15.   Amend, S.R., K.C. Valkenburg, and K.J. Pienta, *Murine Hind Limb Long Bone Dissection*
423   *and Bone Marrow Isolation.* J Vis Exp, 2016(110).

424   16.   Kraus, N.A., et al., *Quantitative assessment of adipocyte differentiation in cell culture.*
425   Adipocyte, 2016. **5**(4): p. 351-358.

426   17.   Bouxsein, M.L., et al., *Guidelines for assessment of bone microstructure in rodents using*
427   *micro-computed tomography.* Journal of Bone and Mineral Research, 2010. **25**(7): p.
428   1468-1486.

429   18.   Bragdon, B.C., et al., *Post natal expression of Prx1 labels appendicular restricted*
430   *progenitor cell populations of multiple tissues.* Journal of Cellular Physiology, 2022.
431   **237**(5): p. 2550-2560.

432   19.   Killaars, A.R., C.J. Walker, and K.S. Anseth, *Nuclear mechanosensing controls MSC*
433   *osteogenic potential through HDAC epigenetic remodeling.* Proc Natl Acad Sci U S A,
434   2020. **117**(35): p. 21258-21266.

435   20.   Goelzer, M., et al., *Depletion of Sun1/2 Induces Heterochromatin Accrual in*
436   *Mesenchymal Stem Cells during Adipogenesis.* bioRxiv, 2022: p. 2022.02.15.480528.

437   21.   Newberg, J., et al., *Isolated Nuclei Stiffen in Response to Low Intensity Vibration.*

438   Journal of Biomechanics, 2020: p. 110012.

439   22.   Uzer, G., et al., *Cell Mechanosensitivity to Extremely Low-Magnitude Signals Is Enabled*

440   *by a LINCed Nucleus.* STEM CELLS, 2015. **33**(6): p. 2063-2076.

441   23.   Styner, M., et al., *Exercise Decreases Marrow Adipose Tissue Through ss-Oxidation in*

442   *Obese Running Mice.* J Bone Miner Res, 2017. **32**(8): p. 1692-1702.

443   24.   McGrath, C., et al., *Exercise Degrades Bone in Caloric Restriction, Despite Suppression*

444   *of Marrow Adipose Tissue (MAT).* J Bone Miner Res, 2020. **35**(1): p. 106-115.

445   25.   McCabe, L.R., et al., *Exercise prevents high fat diet-induced bone loss, marrow*

446   *adiposity and dysbiosis in male mice.* Bone, 2019. **118**: p. 20-31.

447   26.   Isaksson, H., et al., *Physical exercise improves properties of bone and its collagen*

448   *network in growing and maturing mice.* Calcif Tissue Int, 2009. **85**(3): p. 247-56.

449   27.   Isaksson, H., et al., *Long-term voluntary exercise of male mice induces more beneficial*

450   *effects on cancellous and cortical bone than on the collagenous matrix.* Exp Gerontol,

451   2009. **44**(11): p. 708-17.

452   28.   Dirkes, R.K., et al., *Voluntary Wheel Running Partially Compensates for the Effects of*

453   *Global Estrogen Receptor-α Knockout on Cortical Bone in Young Male Mice.* Int J Mol

454   Sci, 2021. **22**(4).

455   29.   Ma, H., et al., *Effects of Diet-Induced Obesity and Voluntary Wheel Running on Bone*

456   *Properties in Young Male C57BL/6J Mice.* Calcified Tissue International, 2010. **86**(5): p.

457   411-419.

458   30.   Somerville, J.M., et al., *Growth of C57BL/6 mice and the material and mechanical*

459   *properties of cortical bone from the tibia.* Calcif Tissue Int, 2004. **74**(5): p. 469-75.

461   **Figure Legends**

Figure 1. Prrx1-driven LINC disruption (PDLD) model characterization. A) IVIS® Spectrum *in vivo* fluorescent scan of P7 pups from Prrx1-Cre/Ai14 cross indicating skeletal Cre activity in Prx-Cre(+) mice (red) vs. no Cre activity in Prx-Cre(-) littermates (blue). **B)** Representative image of bone marrow aspirates from 8-week male Prx-Cre(+)/Ai14 mice indicating the presence of Prx-Cre(+) cells within the bone marrow space. **C)** Bone marrow aspirates from 12-week male Prrx1-driven LINC disrupted (PDLD) and control littermates. Confirmation of eGFP expression localized to the nuclear envelope with nesprin displacement compared to no eGFP visualization and intact nesprin localization to the nuclear envelope in controls.

Figure 2. Bone marrow aspirates from PDLD mice do not display differences in osteogenic and adipogenic differentiation potential compared to LINC intact controls. A) Schematic for differentiation assay timeline as described in Methods **B)** Oil Red O staining after 7 days in adipogenic or normal growth media for 12 week male ODLD mice vs LINC intact controls (n=2/grp). **C)** Absorbance values for eluted Oil Red O stain read at 510nm. **D)** Alizarin Red staining after 21 days in osteogenic or normal growth media. **E)** Absorbance values for eluted Alizarin Red stain read at 562nm. Results presented as mean +/- SD. Significance determined via ordinary one-way ANOVA. *$p<0.05$, **$p<0.01$, ***$p<0.001$

Figure 3. Bone microarchitecture and body mass not significantly different between PDLD and control mice at 8-week baseline. Trabecular measurements include bone volume fraction (BV/TV), trabecular thickness (Tb.Th.), trabecular spacing (Tb.Sp.), and trabecular number (Tb.N). Cortical measurements include total cross-sectional area inside the periosteal envelope (Tt.Ar.), cortical bone area (Ct.Ar.), cortical area fraction (Ct.Ar./Tt.Ar.), and average cortical thickness (Ct.Th.) Significance determined via non-parametric Student's t-test. *$p<0.05$, **$p<0.01$, ***$p<0.001$

Figure 4. PDLD mice run significantly longer, farther, and faster than controls. A) Body

mass steadily increased throughout exercise intervention with no significant difference between

groups. **B)** No significant difference in food intake between groups. **C)** PDLD mice ran longer,

farther, and faster than LINC intact controls. Running metrics tracked by cyclometer measuring

elapsed time, average distance, average velocity, and max velocity (data not shown). Significance

determined via 2-way ANOVA for body weight and food intake and unpaired Student's t-test for

elapsed time, average distance, and average velocity. *$p<0.05$, **$p<0.01$, ***$p<0.001$

Figure 5. Trabecular and cortical microarchitecture of proximal tibiae unaffected by LINC

disruption after 6-week voluntary wheel running. Trabecular measurements include bone

volume fraction (BV/TV), trabecular thickness (Tb.Th.), trabecular spacing (Tb.Sp.), and

trabecular number (Tb.N). Cortical measurements include total cross-sectional area inside the

periosteal envelope (Tt.Ar.), cortical bone area (Ct.Ar.), cortical area fraction (Ct.Ar./Tt.Ar.), and

average cortical thickness (Ct.Th.) *$p<0.05$, **$p<0.01$, ***$p<0.001$

Figure S1 eGFP-KASH expressing cells in primary cell culture. A) Representative images

of bone marrow aspirated from 12-week female Prx-Cre(+)/Ai14(+)/KASH(+) Number of cells

per field expressing the GFP-KASH construct was determined by fluorescent microscopy in the

control cells of Prx lineage were determined by TD tomato expression in both the control All

counts were done utilizing FIJI. **B)** Graph of each cell percentages to total, error bars represent

standard deviation. **C) KASH-GFP expressing cells contribute to the formation of**

osteogenic nodules. Images were taken at 10X using a) FITC filter showing eGFP expression

b), DAPI filter showing calcein blue fluorescence c) an overlay to qualitatively show that the

green KASH-GFP expressing cells were located in the blue Calcein Blue stained nodule region

Figure S2. RNA-seq Analysis of DNK and MC Expressing MSCs During Osteogenesis: (**A**)

Heatmap of FPKM values for mCherry tagged dominant negative KASH (DNK) and empty

mCherry (MC) plasmids expressing MSCs during 7 days of osteogenesis all shown values were

$p<0.05$. (**B**) FPKM values of osteogenesis genes of Ibsp, Bglap, Alpl, and Phospho1 in MSCs

expressing DNK or MC plasmid during 7-day osteogenesis, both undifferentiated cintrols (UD)

and osteogenic media (OS) was shown.

Figure 1. Prrx1-driven LINC disruption (PDLD) model characterization. A) IVIS®
Spectrum *in vivo* fluorescent scan of P7 pups from Prrx1-Cre/Ai14 cross indicating skeletal
Cre activity in Prx-Cre(+) mice (red) vs. no Cre activity in Prx-Cre(-) littermates (blue). **B)**
Representative image of bone marrow aspirates from 8-week male Prx-Cre(+)/Ai14 mice
indicating the presence of Prx-Cre(+) cells within the bone marrow space. **C)** Bone marrow
aspirates from 12-week male Prrx1-driven LINC disrupted (PDLD) and control littermates.
Confirmation of eGFP expression localized to the nuclear envelope with nesprin
displacement compared to no eGFP visualization and intact nesprin localization to the
nuclear envelope in controls.

Figure 2. Bone marrow aspirates from PDLD mice do not display differences in osteogenic and adipogenic differentiation potential compared to LINC intact controls. A) Schematic for differentiation assay timeline as described in Methods **B)** Oil Red O staining after 7 days in adipogenic or normal growth media for 12 week male ODLD mice vs LINC intact controls (n=2/grp). **C)** Absorbance values for eluted Oil Red O stain read at 510nm. **D)** Alizarin Red staining after 21 days in osteogenic or normal growth media. **E)** Absorbance values for eluted Alizarin Red stain read at 562nm. Results presented as mean +/- SD. Significance determined via ordinary one-way ANOVA. *p<0.05, **p<0.01, ***p<0.001

Figure 3. Bone microarchitecture and body mass not significantly different between PDLD and control mice at 8-week baseline. Trabecular measurements include bone volume fraction (BV/TV), trabecular thickness (Tb.Th.), trabecular spacing (Tb.Sp.), and trabecular number (Tb.N). Cortical measurements include total cross-sectional area inside the periosteal envelope (Tt.Ar.), cortical bone area (Ct.Ar.), cortical area fraction (Ct.Ar./Tt.Ar.), and average cortical thickness (Ct.Th.) Significance determined via non-parametric Student's t-test. $*p<0.05$, $**p<0.01$, $***p<0.001$

Figure 4. PDLD mice run significantly longer, farther, and faster than controls. A) Body mass steadily increased throughout exercise intervention with no significant difference between groups. **B)** No significant difference in food intake between groups. **C)** PDLD mice ran longer, farther, and faster than LINC intact controls. Running metrics tracked by cyclometer measuring elapsed time, average distance, average velocity, and max velocity (data not shown). Significance determined via 2-way ANOVA for body weight and food intake and unpaired Student's t-test for elapsed time, average distance, and average velocity. *p<0.05, **p<0.01, ***p<0.001

Figure 5. Trabecular and cortical microarchitecture of proximal tibiae unaffected by LINC disruption after 6-week voluntary wheel running. Trabecular measurements include bone volume fraction (BV/TV), trabecular thickness (Tb.Th.), trabecular spacing (Tb.Sp.), and trabecular number (Tb.N). Cortical measurements include total cross-sectional area inside the periosteal envelope (Tt.Ar.), cortical bone area (Ct.Ar.), cortical area fraction (Ct.Ar./Tt.Ar.), and average cortical thickness (Ct.Th.) *p<0.05, **p<0.01, ***p<0.001

Supplementary Figures:

Prrx1-driven LINC complex disruption *in vivo* does not significantly alter bone properties in 8-week male mice nor after 6-weeks voluntary wheel running

Scott Birks[1], Sean Howard[2], Christian S. Wright[3], Caroline O'Rourke[4], Anthony Lau[4], William R. Thompson[3], Gunes Uzer[2] †

[1]Boise State University, Micron School of Materials Science and Engineering
[2]Boise State University, Mechanical and Biomedical Engineering
[3]Indiana University, Department of Physical Therapy, School of Health and Human Sciences
[4]The College of New Jersey, Biomedical Engineering

Figure S1 eGFP-KASH expressing cells in primary cell culture. A) Representative images of bone marrow aspirated from 12-week female Prx-Cre(+)/Ai14(+)/KASH(+) Number of cells per field expressing the GFP-KASH construct was determined by fluorescent microscopy in the control cells of Prx lineage were determined by TD tomato expression in both the control All counts were done utilizing FIJI. **B)** Graph of each cell percentages to total, error bars represent standard deviation. **C) KASH-GFP expressing cells contribute to the formation of osteogenic nodules.** Images were taken at 10X using a) FITC filter showing eGFP expression b), DAPI filter showing calcein blue fluorescence c) an overlay to qualitatively show that the green KASH-GFP expressing cells were located in the blue Calcein Blue stained nodule region

Figure S2. RNA-seq Analysis of DNK and MC Expressing MSCs During Osteogenesis: (**A**) Heatmap of FPKM values for mCherry tagged dominant negative KASH (DNK) and empty mCherry (MC) plasmids expressing MSCs during 7 days of osteogenesis all shown values were p<0.05. (**B**) FPKM values of osteogenesis genes of Ibsp, Bglap, Alpl, and Phospho1 in MSCs expressing DNK or MC plasmid during 7-day osteogenesis, both undifferentiated cintrols (UD) and osteogenic media (OS) was shown.