Antifungal efficacy of natural antiseptic products against *Candida auris*

Wing-Gi WU[a#], Kristine Shik LUK[a,b,c,d], Mei-Fan HUNG[a], Wing-Yi TSANG[a], Kin-Ping LEE[a,c], Bosco Hoi-Shiu LAM[a], Ka-Lam CHENG[c], Wing-Sze CHEUNG[b], Hau-Ling TANG[d], Wing-Kin TO[a,b]

[a]Department of Pathology, Princess Margaret Hospital, Hong Kong Special Administrative Region, China

[b]Infection Control Team, Princess Margaret Hospital, Hong Kong Special Administrative Region, China

[c]Infection Control Team, Caritas Medical Centre, Hong Kong Special Administrative Region, China

[d]Infection Control Team, Yan Chai Hospital, Hong Kong Special Administrative Region, China

#Correspondence to: Dr Wing-Gi WU, Department of Pathology, Princess Margaret Hospital, 2-10 Princess Margaret Hospital Road, Lai Chi Kok, Kowloon, Hong Kong. Email: wwg396@ha.org.hk

Keywords: Candida auris, natural, decolonization, tea tree, manuka

Authors' Contact Details

WU WG – wwg396@ha.org.hk

LUK KS – sluk@ha.org.hk

HUNG MF – hungmf@ha.org.hk

TSANG WY – twy029@ha.org.hk

LEE KP – may.lee@ha.org.hk

LAM BHS - bhslam@gmail.com

CHENG KL - BCKLZ09@ha.org.hk

CHEUNG WS - heungws@ha.org.hk

TANG HL - gigi_tang3@yahoo.com.hk

TO WK – towk@ha.org.hk

ABSTRACT

Candida auris is an emerging fungal pathogen responsible for healthcare associated infections and outbreaks with high mortality around the world. It readily colonizes the skin, nares, respiratory and urinary tract of hospitalized patients, and such colonization may lead to invasive *Candida* infection in susceptible patients. However, there is currently a lack of decolonization protocol for *C. auris*. The aim of this study is to evaluate the susceptibility of *C. auris* to commonly used synthetic and natural antiseptic products. Synthetic antiseptics including chlorhexidine, povidone-iodine, and nystatin were shown to be fungicidal against *C. auris*. Among the natural antiseptics tested, tea tree oil and manuka oil were fungicidal at concentrations less than 1.25% and 0.63% (v/v), respectively. Manuka honey inhibited *C. auris* at 25% (v/v) concentrations. Among the commercial products tested, manuka body wash and mouthwash were fungicidal against *C. auris* at concentrations less than 0.10% (w/v) and 6.25% (v/v), respectively, while tea tree body wash and Medihoney™ wound gel demonstrated fungistatic properties. In conclusion, this study demonstrated good in vitro antifungal efficacy of tea tree oil, manuka oil, manuka honey, and commercially available antiseptic products containing these active ingredients. Future studies are warranted to evaluate the effectiveness of these antiseptic products in clinical settings.

INTRODUCTION

The fungal pathogen *Candida auris* was first described in 2009 after isolation from the ear of a patient in Japan [1]. It has since been reported in more than 40 countries across six continents, causing a wide range of healthcare-associated outbreaks and infections [2], with the mortality rate varying from 30 to 60%, due to the high virulence and resistance to several antifungals [3]. The recent inaugural World Health Organization fungal priority pathogen list highlights the urgent threat of *C. auris* to global public health [4]. With its biofilm-forming ability and tolerance to high temperature and salt concentrations, *C. auris* can persist and survive in the human skin and environmental surfaces for several months following exposures [5-7]. This provides additional opportunities for transmission and colonization within the healthcare settings. Furthermore, the minimum time to acquire *C. auris* from a patient or their immediate environment is 4 hours or less, reinforcing the importance of vigilant infection control measures [8]. This fungal superbug can also resist certain disinfectants and is well adapted to healthcare environments [9].

In 2019, a cluster of *C. auris* colonization affecting 15 patients was reported in Princess Margaret Hospital (PMH), Kowloon West (KW) cluster, Hong Kong. Whole genome sequencing identified the strains as Clade 1/South Asia, associated with ERG11 mutation conferring azole resistance [10]. The clade is also known to cause invasive infections and outbreaks frequently [6]. Four regional hospitals (a total of 3,878 beds) serve an estimated population of 1.4 million in the KW cluster. In spite of enhanced surveillance and infection control measures, *C. auris* cases and transmission have continued to increase due to pandemic-related strain on the healthcare system. As of April 2023, there were 364 cases of *C. auris* reported in KW Cluster, Hong Kong.

C. auris readily colonizes the axillae, groin, nares, respiratory, and urinary tract of hospitalized patients, and less frequently the intestinal tract than other *Candida* species [11].

Colonization predisposes to invasive infection, with mechanical ventilation and placement of invasive devices identified as major risk factors [6]. A descriptive analysis of *C. auris* outbreaks in New York State reported that 4% of *C. auris* colonized patients developed bloodstream infection, with a rate of 0.3 per 1,000 patient-days [12]. These studies provided evidence that an important strategy for preventing severe candidiasis is the elimination of *C. auris* carriage to preclude subsequent risk of endogenous infection. Nevertheless, recommendations on decolonization protocol for *C. auris* are currently lacking. Successful decolonization by a five-day protocol with twice-daily chlorhexidine body wash, mouthwash containing 0.2% chlorhexidine, and oral nystatin has been reported in a healthcare worker having positive nasal swabs [8]. Both chlorhexidine, povidone-iodine, and nystatin have been used as cutaneous antiseptics prior to surgical procedures, wound irrigation, and the insertion of intravascular devices [13-16]. In search of an effective decolonization protocol for *C. auris*, different topical antiseptics, such as chlorhexidine, povidone-iodine, isopropyl alcohol, and nystatin have been studied, with their yeasticidal activities against *C. auris* demonstrated in vitro [17-19]. Chlorhexidine is one of the most common disinfectants used in healthcare settings. Despite laboratory evidence showing excellent in vitro activity of chlorhexidine against *C. auris*, clinical experience from several outbreaks suggested that colonization persisted even with daily chlorhexidine washes [8, 9]. A murine model study demonstrated that *C. auris* could enter the dermis without causing overt histopathologic signs of inflammation, resulting in deeper skin layer residence [20]. One factor contributing to *C. auris* persistent colonization may be the limited permeation into deeper skin layers by chlorhexidine, which was illustrated by a study based on a model of ex vivo human skin [21]. Poor effectiveness of chlorhexidine in decolonization, as well as subsequent recolonization of *C. auris* in patient who had repeated interval negative swabs, is anticipated.

The development of novel and effective decolonization regimen against *C. auris* using

alternative antiseptics has become crucial. Natural substances with antimicrobial properties are potential good candidates as they would not mount additional drug-selecting pressure and they are well tolerated as skin washes in general. *Fernades et al.* have shown that tea tree (*Melalenca alternifolia*) oil had good in vitro antifungal activity against *C. auris* in both planktonic cells and biofilms [22]. *Johnson et al.* have also shown that the active components of tea tree oil could augment the activity of chlorhexidine for *C. auris* decolonization in the porcine skin model [23]. Another potential effective natural antiseptic is manuka (*Leptospermum scoparium*), which is also known as the "New Zealand tea tree". It has been a part of the traditional Māori medicine, including topical treatment for wounds, cuts, and sores, with fungicidal properties towards *C. albicans* and *C. glabrata* [24]. Notably, manuka honey was found to be effective in inhibiting various *Candida* species [25]. There are multiple antimicrobial constituents of the natural extracts, namely monoterpenes and sesquiterpenes (tea tree and manuka oil), triketones (manuka oil), methylglyoxal, polyphenolic compounds and bee defensin-1 (manuka honey) [26-28]. Therefore, unlike other synthetic antimicrobials, there have been no reports of clinical isolates with acquired resistance to the natural antiseptics [29, 30]. Clinical trials have demonstrated the efficacy of manuka honey and tea tree oil in the treatment of oral candidiasis, vulvovaginal candidiasis and successful decolonization for methicillin-resistant Staphylococcus aureus (MRSA) carriers [31-36], making them attractive alternatives against *C. auris*. These natural components are also widely used as active ingredients for their antiseptic effects in commercial hygiene products such as body wash, mouthwash and wound gel. To pave the way for evaluating the clinical application of the natural antiseptics against *C. auris*, this study aimed to determine the in-vitro antifungal susceptibility of *C. auris* to the natural antiseptics and their commercial products; and to compare the antifungal susceptibility with that of synthetic antiseptics.

MATERIALS AND METHODS

Characterization of *Candida auris* isolates: A total of 40 *C. auris* isolates belonging to clade I/South Asia (unpublished data) were obtained in the KW Cluster of Hong Kong from 2019 to 2022. All tested isolates were identified by matrix-assisted laser desorption ionization-time of flight mass spectrometry (Bruker Daltonics; Database version: IVD-9468 MSP) [37]. Reference strains *C. auris* JCM 15448, *C. albicans* ATCC 90028, *C. glabrata* ATCC MYA-2950, *C. krusei* ATCC 6258, and *C. parapsilosis* ATCC 22019 were also included. The study protocol was approved by the Research Ethics Committee of the Kowloon West Cluster, Hospital Authority, Hong Kong (KW/EX-21-098(160-14)).

Antiseptics and preparations: Our experiment studied three groups of topical antiseptics. The first group included synthetic antiseptics products commonly used in healthcare settings: chlorhexidine gluconate (Microshield® 4%, Australia), povidone-iodine (Betadine® 10%, Turkey) and nystatin (100,000 units/mL, Pharmascience, Canada). The second group comprised natural antiseptics: tea tree oil (15%, The Body Shop, UK; 100% Kiwi Manuka®, New Zealand), manuka oil (100%, Manuka Lab®, New Zealand; 100%, Melora®, New Zealand), manuka honey (UMF 15+ and 20+, Comvita®, New Zealand; UMF 15+ and 20+, Melora®, New Zealand), manuka hydrosol (Bioactives®, New Zealand). The third group consisted of commercial antiseptic products containing active components from tea tree oil and/or manuka derivatives: manuka body wash (Manuka Lab®, New Zealand), tea tree clearing body wash (The Body Shop®, UK), manuka mouthwash (Manuka Lab®, New Zealand), and Medihoney™ antibacterial wound gel (Comvita®, New Zealand). Dimethyl sulfoxide (DMSO, Sigma-Aldrich, UK) was added to water-insoluble products, including tea tree oil, manuka oil, antibacterial wound gel, and nystatin to enhance solubility in RPMI 1640 medium (Sigma, Paisley, UK).

In vitro antifungal susceptibility of antiseptics: In vitro antifungal susceptibility of antiseptics was performed by broth microdilution method according to CLSI M27-A4 [38].

Briefly, 100 µL of RPMI 1640 medium with different concentrations of the antiseptics were added to 96-well microtiter plates (Saining Biotechnology, China). *Candida* isolates were suspended in 2 mL sterile physiological saline to obtain 0.5 McFarland standard suspension using the DensiCHEK Plus instrument (BioMerieux, France), corresponding to 1-5 x 10^6 cells per mL, then further diluted with RPMI 1640 medium to obtain a 1:1000 final inoculum containing 1-5 x 10^3 cells per mL. The yeast suspensions (100 µL) were added to the 96-well plate with the product suspensions (100 µL) and incubated aerobically at 35°C for 24 h. The minimum inhibitory concentration (MIC) endpoint of a product was defined as the lowest drug concentration that caused a prominent decrease in growth (50%) in relation to the growth control using a reading mirror. The minimum fungicidal concentration (MFC) endpoint was defined as the lowest drug concentration resulting in the death of 99.9% or more of the initial inoculum. To determine MFC values, 100 µL of broth was taken from the wells without microbial growth and inoculated onto Sabouraud dextrose agar (BD DIFCO^{TM}, USA). Following incubation at 35°C for 48 h, the colony-forming unit (CFU) was counted to assess viability. Each isolate was tested in duplicate separately and the assay was repeated if the MIC or MFC differed by more than one dilution. Antifungal drug susceptibility of *C. krusei* ATCC 6258 and *C. parapsilosis* ATCC 22019 was performed in each run of broth microdilution as controls [39].

For antiseptics demonstrating fungistatic effects, log growth reduction in CFU after 24-hour incubation was measured, and the test was repeated in triplicate. Mann-Whitney U test was used to analyze the significance of difference in CFU count compared to the growth control, and one-way ANOVA, Kruskal-Wallis, or Mann-Whitney U tests were used to compare the log growth reduction of the antiseptics at different concentrations. Pairwise comparisons were performed using Bonferroni or Dunn's test when suitable. P value <0.05 was considered statistically significant. All statistical analyses were performed using STATA18.0 (StataCorp.

2023. Stata Statistical Software: Release 18. College Station, TX: StataCorp LLC.).

In vitro antifungal susceptibility testing of antifungal drugs: To confirm the local

resistance pattern of our isolates to conventional antifungal drugs, in vitro susceptibility was

determined using Sensititre^{TM} YeastONE^{TM} YO10 AST Plate (ThermoFisher Scientific, UK).

The antifungal drugs included anidulafungin, micafungin, caspofungin, 5-flucytosine,

posaconazole, voriconazole, itraconazole, fluconazole and amphotericin B. The test was

performed according to the procedures described by the manufacturer [40]. Centres for Disease

Control and Prevention (CDC)'s tentative breakpoints for *C. auris* were used as a reference

[41].

RESULTS

Table 1 summarizes the antifungal activities of topical antiseptics against *C. auris*. For the synthetic antiseptics, chlorhexidine, povidone-iodine and nystatin were fungicidal against *C. auris* at concentrations less than 21.99 mg/L (0.004%), 312.50 mg/L (0.313%), and 9.77 µg/mL (0.049%), respectively. The MICs and MFCs of chlorhexidine against *C. auris* strains, including the JCM 15448, were relatively higher when compared with that against *C. krusei* ATCC 6258, *C. parapsilosis* ATCC 22019, *C. albicans* ATCC 90028, and *C. glabrata* ATCC MYA-2950.

For the natural antiseptics, manuka honey demonstrated fungistatic properties against clinical *C. auris* isolates at 25% (v/v). Pure tea tree and manuka oil were fungicidal against *C. auris* at concentrations less than 1.25% and 0.63% (v/v), respectively. Commercial products containing active ingredients of manuka honey, manuka oil and/or tea tree oil were shown to be active in vitro against *C. auris* and other *Candida* species. Manuka body wash and mouthwash were fungicidal against *C. auris* at concentrations less than 0.10% (w/v) and 6.25% (v/v), respectively. Manuka body wash, however, exhibited weaker antifungal activity against *C. parapsilosis* (MFC 3.13% (w/v)). Interestingly, the MICs of natural antiseptics against *C. auris* JCM 15448 were 4- to 128-fold lower than that against the clinical counterparts; tea tree body wash and Medihoney™ wound gel even became fungicidal against *C. auris* JCM 15488 at concentrations of 25% (w/v) and 12.5% (w/v), respectively. Manuka hydrosol was not effective against all tested *Candida* species.

Manuka honey, Medihoney™ wound gel and tea tree body wash demonstrated fungistatic properties against clinical *C. auris* isolates in vitro. At a concentration of 37.5%, the natural antiseptics exhibited an average 4 log reduction in CFU after 24 h of incubation, which is statistically significant in comparison with the growth control ($p < 0.01$), as shown in Figure 1. The inhibitory effects of manuka honey and tea tree body wash against *C. auris* were

218   concentration-dependent (p <0.05). Medihoney^{TM} wound gel at 25% and 37.5% resulted in a

219   similar reduction (6-log) in CFU.

220   The antifungal susceptibility of common systemic antifungals against *C. auris* was shown

221   in table 2. All tested isolates (N=40) were resistant to fluconazole and 90% (N=36) were also

222   resistant to Amphotericin B. Susceptibility to other azoles and echinocandins was variable.

DISCUSSION

The emergence of multidrug-resistant *C. auris* to conventional antifungal drugs and its high propensity to cause nosocomial transmission make it urgent to develop novel and effective decolonization strategy. This in vitro study is a pilot study to test antifungal susceptibility of several natural antiseptics against *C. auris*. Herein, we demonstrated excellent in vitro antifungal activity of natural antiseptics, including manuka honey, manuka oil and tea tree oil, and their commercial products, against *C. auris*. This suggests the current decolonization strategy could be much improved by combining these well-tolerated, natural antiseptics. A clinical trial on *C. auris* decolonization using manuka body wash, manuka mouthwash, tea tree oil, povidone-iodine, and nystatin oral suspension in 41 patients showed about 60% success rate and no adverse effects were reported (Luk, 2023, unpublished data).

In the present study, the local *C. auris* isolates were highly resistant to both fluconazole (MIC_{50} of 256 µg/mL) and amphotericin B (MIC_{50} of 2 µg/mL). Genomic and proteomic analyses revealed mutation of the fluconazole target enzyme (cytochrome 450 lanosterol 14α-demethylase) [10] and the role of extracellular vesicles in the development of amphotericin B resistance [42].

Our results confirmed the high susceptibility of *C. auris* to chlorhexidine (MFC 0.004%), povidone iodine (MFC 0.312%) and nystatin (MFC 9.77 µg/mL) at concentrations used for clinical indications [43, 44]. Nonetheless, the MICs an MFCs of chlorhexidine against *C. auris* strains were higher than that against other *Candida* species. Similar findings were reported by *Abdolrasouli et al* [44], showing that their *C. auris* outbreak strains were less susceptible to chlorhexidine than other *Candida* species though the testing methods were different and the mechanism is currently unknown. Likewise, *C. auris* was highly susceptible to tea tree oil (MFC 1.25%) and manuka oil (MFC 0.63%); the fungicidal effects of tea tree and manuka oil were comparable to synthetic antiseptics, including chlorhexidine, povidone-iodine and

nystatin. Our findings are consistent with a previous report of high antifungal activity of tea

tree oil against *C. auris* isolates (MFC 1.56% (w/v)) [22]. Tea tree oil and its main component,

terpinene-4-ol, have been shown to cause damage to the fungal cell wall and cytoplasmic

membrane thereby causing cellular dysfunctions and death though the exact mechanism is

unknown [45, 46]. Interestingly, The Body Shop® formulation contains a diluted content of

15% tea tree oil but a lower MIC_{90} of 0.004% (v/v) was observed compared to 100% tea tree

oil. This enhanced inhibition may be attributed to the synergistic or additive interactions of the

supplementary ingredients, including denatured alcohol, *Calophyllum inophyllum* seed oil, and

Leptospermum petersonii oil, which were shown to have antimicrobial properties [47, 48].

Moore et al. also showed the addition of isopropyl alcohol to disinfectant formulation clearly

enhanced the killing of *C. auris* [17]. On the other hand, the antifungal properties of manuka

oil were less frequently evaluated for *C. auris* specifically. *Parker et al.* demonstrated

significant inhibitory effects of manuka oil on *C. auris* AR0381 (clade II) and AR0385 (clade

IV), with MIC values less than 1.0% [49], which agree well with our findings. More importantly,

the low fungicidal concentrations of tea tree oil and manuka oil against *C. auris* are considered

safe and tolerable for dermal use. An in vitro skin model proved manuka oil is non-irritant, with

cell viability unaffected after treatment with an undiluted sample [50]. Another clinical trial

also demonstrated that 5% tea tree oil shampoo was well tolerated in the treatment for dandruff

[51]. The additional property of enhanced skin permeation of essential oils [52] further supports

the role of essential oils in eliminating *C. auris* colonization.

Regarding manuka honey, *C. auris* were inhibited at a concentration of 25%. The growth

inhibition was presented in a dose-dependent manner. This is in line with several studies

showing fungistatic effect of manuka honey against *Candida* species. [25, 53]. Manuka honey

belongs to the "non-peroxide-based" honey group, and its main antimicrobial action is

contributed by methylglyoxal (MGO) [54], which is converted non-enzymatically from the

dihydroxyacetone present in the nectar of the flowers of *Leptospermum scoparium*. The

characteristically high level of MGO in manuka honey makes it a unique marker for

authentication. In this study, manuka honey with greater Unique Manuka Factor (UMF), which

is correlated with the MGO and total phenol components [55], resulted in greater CFU reduction

of *C. auris*. Increasing the concentration of honey from 25% to 50% led to an average 3-log

reduction in the CFU of *C. auris* ($p < 0.05$). Surprisingly, the *C. auris* JCM 15448 (clade II)

had a much lower MIC (0.2% v/v) towards manuka honey. Similar phenomenon was also

observed for other natural antiseptics. Comparative genomic analysis revealed that multiple

genes involved in cell wall proteins and stress response-related functions were absent in the

Japanese *C. auris* strains, including JCM 15448, while a more recent phenotypic analysis also

supported clade-specific phenotypic profiles of *C. auris* with lowest tolerance to osmotic stress

and chemicals observed in clade II [56, 57]. These genomic structural variations and metabolic

profiles may highlight the differences in antifungal resistance, environmental survival, and

virulence among *C. auris* isolates. Nevertheless, previous study did not suggest difference in

antifungal activity of medical grade honey formulation among *C. auris* clades [58]. Studies

including larger numbers of *C. auris* of different clades should determine whether susceptibility

to natural antiseptics is clade specific.

As expected, commercial products having the active ingredients of natural antiseptics

showed promising antifungal effect in vitro against *C. auris*. Manuka body wash and

mouthwash were fungicidal against *C. auris* at low concentrations. Product labels of manuka

body wash and mouthwash disclose the ingredients of 525+ MGO (mg/kg), which is equivalent

to UMF 15+ manuka honey. Besides, the aforementioned products contain tea tree and manuka

oil with antifungal effects demonstrated in our study. Alternatively, Medihoney^{TM} antibacterial

wound gel containing 80% medical-grade manuka honey and The Body Shop$^{®}$ tea tree body

wash, also caused a 6-log reduction in *C. auris* CFU. Together with the anti-inflammatory and

anti-oxidant properties of manuka honey [30], the commercial hygiene products serve as

important options in the development of an efficacious and tolerable decolonization regimen.

There are some limitations in this study. First, extraction and evaluation of the bioactive

chemical markers in the natural antiseptics were not performed. The specific quantities of the

active ingredients were often not disclosed in the product labels. Besides, all our *C. auris*

isolates belonged to clade I and the results of this study may not be applicable to isolates of

other clades. We had included *C. auris* JCM 15448, a clade II strain, in our testing and found

significantly lower MIC and MFC values towards the natural antiseptics. Moreover, the effect

of antifungal drug resistance on the antiseptics could not be studied due to the low variability

of the resistance pattern of our *C. auris* isolates. Nonetheless, these antiseptic products had an

overall high in vitro antifungal efficacy, which supported their application as topical or

decolonization therapies for *C. auris.*

CONCLUSION

313   *C. auris* are highly susceptible to natural antiseptics including tea tree oil, manuka oil and

314   manuka honey, and their commercial hygiene products in the forms of body wash, mouthwash,

315   and wound gel. The promising in vitro results provide insight into the development of novel

316   and effective *C. auris* decolonization regimen. Future studies should demonstrate whether these

317   natural antiseptics are effective in the clinical settings.

ACKNOWLEDGEMENT

The funding of this work was supported by the Clinical Research Centre Mini Research Grant 2023 (2023-CRC-MRG-02), Kowloon West Cluster, Hospital Authority, Hong Kong. We wish to thank the hospital infection control teams in the KW cluster for their efforts in *C. auris* control and the microbiology laboratory staff of Princess Margaret Hospital for the antifungal testing.

CONFLICT OF INTEREST

We declare no conflict of interest.

DATA AVAILABILITY

All data are incorporated into the article and its online supplementary material.

REFERENCE

1. Satoh K, Makimura K, Hasumi Y, Nishiyama Y, Uchida K, Yamaguchi H. *Candida auris* sp. nov., a novel ascomycetous yeast isolated from the external ear canal of an inpatient in a Japanese hospital. *Microbiology and Immunology*. 2009;53(1):41-44. doi:10.1111/j.1348-0421.2008.00083.x

2. de Cássia Orlandi Sardi J, Silva DR, Soares Mendes-Giannini MJ, Rosalen PL. Candida auris: Epidemiology, risk factors, virulence, resistance, and therapeutic options. *Microb Pathog*. 2018;125:116-121. doi:10.1016/j.micpath.2018.09.014

3. Al-Rashdi A, Al-Maani A, Al-Wahaibi A, Alqayoudhi A, Al-Jardani A, Al-Abri S. Characteristics, Risk Factors, and Survival Analysis of Candida auris Cases: Results of One-Year National Surveillance Data from Oman. *JoF*. 2021;7(1):31. doi:10.3390/jof7010031

4.   Fisher MC, Denning DW. The WHO fungal priority pathogens list as a game-changer.

Nat Rev Microbiol. 2023;21(4):211-212. doi:10.1038/s41579-023-00861-x

5.   Piedrahita CT, Cadnum JL, Jencson AL, Shaikh AA, Ghannoum MA, Donskey CJ.

Environmental Surfaces in Healthcare Facilities are a Potential Source for Transmission

of *Candida auris* and Other *Candida* Species. *Infect Control Hosp Epidemiol*.

2017;38(9):1107-1109. doi:10.1017/ice.2017.127

6.   Ahmad S, Alfouzan W. Candida auris: Epidemiology, Diagnosis, Pathogenesis,

Antifungal Susceptibility, and Infection Control Measures to Combat the Spread of

Infections   in   Healthcare   Facilities.   *Microorganisms*.   2021;9(4):807.

doi:10.3390/microorganisms9040807

7.   Berrio I, Caceres DH, Coronell R W, et al. Bloodstream Infections With *Candida auris*

Among Children in Colombia: Clinical Characteristics and Outcomes of 34 Cases.

Journal of the Pediatric Infectious Diseases Society. 2021;10(2):151-154.

doi:10.1093/jpids/piaa038

8.   Schelenz S, Hagen F, Rhodes JL, et al. First hospital outbreak of the globally emerging

Candida auris in a European hospital. *Antimicrob Resist Infect Control*. 2016;5(1):35.

doi:10.1186/s13756-016-0132-5

9.   Biswal M, Rudramurthy SM, Jain N, et al. Controlling a possible outbreak of Candida

auris infection: lessons learnt from multiple interventions. *Journal of Hospital Infection*.

2017;97(4):363-370. doi:10.1016/j.jhin.2017.09.009

10. Tse H, Tsang AKL, Chu YW, Tsang DNC. Draft Genome Sequences of 19 Clinical

Isolates of Candida auris from Hong Kong. *Microbiol Resour Announc*.

2021;10(1):e00308-20. doi:10.1128/MRA.00308-20

11. Piatti G, Sartini M, Cusato C, Schito AM. Colonization by Candida auris in critically ill

patients: role of cutaneous and rectal localization during an outbreak. *Journal of*

Hospital Infection. 2022;120:85-89. doi:10.1016/j.jhin.2021.11.004

12. Southwick K, Adams EH, Greenko J, et al. 2039. New York State 2016–2018:

Progression from *Candida auris* Colonization to Bloodstream Infection. *Open Forum*

Infectious Diseases. 2018;5(suppl_1):S594-S595. doi:10.1093/ofid/ofy210.1695

13. Phillips M, Rosenberg A, Shopsin B, et al. Preventing Surgical Site Infections: A

Randomized, Open-Label Trial of Nasal Mupirocin Ointment and Nasal Povidone-

Iodine Solution. *Infect Control Hosp Epidemiol*. 2014;35(7):826-832.

doi:10.1086/676872

14. Bigliardi PL, Alsagoff SAL, El-Kafrawi HY, Pyon JK, Wa CTC, Villa MA. Povidone

iodine in wound healing: A review of current concepts and practices. *International*

Journal of Surgery. 2017;44:260-268. doi:10.1016/j.ijsu.2017.06.073

15. Bodin MB, Godoy G, Philips JB. Topical Nystatin for the Prevention of Catheter-

Associated Candidiasis in ELBW Infants. *Advances in Neonatal Care*. 2015;15(3):220-

224. doi:10.1097/ANC.0000000000000170

16. Jalalzadeh H, Groenen H, Buis DR, et al. Efficacy of different preoperative skin

antiseptics on the incidence of surgical site infections: a systematic review, GRADE

assessment, and network meta-analysis. *The Lancet Microbe*. 2022;3(10):e762-e771.

doi:10.1016/S2666-5247(22)00187-2

17. Moore G, Schelenz S, Borman AM, Johnson EM, Brown CS. Yeasticidal activity of

chemical disinfectants and antiseptics against Candida auris. *J Hosp Infect*.

2017;97(4):371-375. doi:10.1016/j.jhin.2017.08.019

18. Ku TSN, Walraven CJ, Lee SA. Candida auris: Disinfectants and Implications for

Infection Control. *Front Microbiol*. 2018;9:726. doi:10.3389/fmicb.2018.00726

19. Voorn MG, Kelley AM, Chaggar GK, Li X, Teska PJ, Oliver HF. Contact time and

disinfectant formulation significantly impact the efficacies of disinfectant towelettes

against Candida auris on hard, non-porous surfaces. *Sci Rep*. 2023;13(1):5849.

doi:10.1038/s41598-023-32876-y

20. Huang X, Hurabielle C, Drummond RA, et al. Murine model of colonization with fungal

pathogen Candida auris to explore skin tropism, host risk factors and therapeutic

strategies. *Cell Host & Microbe*. 2021;29(2):210-221.e6.

doi:10.1016/j.chom.2020.12.002

21. Karpanen TJ, Worthington T, Conway BR, Hilton AC, Elliott TSJ, Lambert PA.

Penetration of Chlorhexidine into Human Skin. *Antimicrob Agents Chemother*.

2008;52(10):3633-3636. doi:10.1128/AAC.00637-08

22. Fernandes L, Ribeiro R, Costa R, Henriques M, Rodrigues ME. Essential Oils as a Good

Weapon against Drug-Resistant Candida auris. *Antibiotics*. 2022;11(7):977.

doi:10.3390/antibiotics11070977

23. Johnson CJ, Eix EF, Lam BC, et al. Augmenting the Activity of Chlorhexidine for

Decolonization of Candida auris from Porcine skin. *JoF*. 2021;7(10):804.

doi:10.3390/jof7100804

24. Lee YS, Kim J, Lee SG, Oh E, Shin SC, Park IK. Effects of plant essential oils and

components from Oriental sweetgum (Liquidambar orientalis) on growth and

morphogenesis of three phytopathogenic fungi. *Pesticide Biochemistry and Physiology*.

2009;93(3):138-143. doi:10.1016/j.pestbp.2009.02.002

25. Anand S, Deighton M, Livanos G, Pang ECK, Mantri N. Agastache honey has superior

antifungal activity in comparison with important commercial honeys. *Sci Rep*.

2019;9(1):18197. doi:10.1038/s41598-019-54679-w

26. Borotová P, Galovičová L, Vukovic NL, Vukic M, Tvrdá E, Kačániová M. Chemical

and Biological Characterization of Melaleuca alternifolia Essential Oil. *Plants*.

2022;11(4):558. doi:10.3390/plants11040558

27. Mathew C, Tesfaye W, Rasmussen P, et al. Mānuka Oil—A Review of Antimicrobial

and Other Medicinal Properties. *Pharmaceuticals*. 2020;13(11):343.

doi:10.3390/ph13110343

28. Pleeging CCF, Wagener FADTG, De Rooster H, Cremers NAJ. Revolutionizing non-

conventional wound healing using honey by simultaneously targeting multiple

molecular mechanisms. *Drug Resistance Updates*. 2022;62:100834.

doi:10.1016/j.drup.2022.100834

29. Blair SE, Cokcetin NN, Harry EJ, Carter DA. The unusual antibacterial activity of

medical-grade Leptospermum honey: antibacterial spectrum, resistance and

transcriptome analysis. *Eur J Clin Microbiol Infect Dis*. 2009;28(10):1199-1208.

doi:10.1007/s10096-009-0763-z

30. Cooper RA, Jenkins L, Henriques AFM, Duggan RS, Burton NF. Absence of bacterial

resistance to medical-grade manuka honey. *Eur J Clin Microbiol Infect Dis*.

2010;29(10):1237-1241. doi:10.1007/s10096-010-0992-1

31. Jandourek A, Vaishampayan JK, Vazquez JA. Efficacy of melaleuca oral solution for

the treatment of fluconazole refractory oral candidiasis in AIDS patients. *AIDS*.

1998;12(9):1033-1037.

32. Caelli M, Porteous J, Carson CF, Heller R, Riley TV. Tea tree oil as an alternative

topical decolonization agent for methicillin-resistant Staphylococcus aureus. *J Hosp

Infect*. 2000;46(3):236-237. doi:10.1053/jhin.2000.0830

33. Seifinadergoli Z, Nahidi F, Safaiyan A, Javadzadeh Y, Eteraf-Oskouei T. Comparison

of the efficacy of honey gel and clotrimazole cream in the treatment of vaginal

candidiasis symptoms: a randomized clinical trial. *Electron Physician*. 2018;10(6):6904-6911. doi:10.19082/6904

34. Yang C, Gong G, Jin E, et al. Topical application of honey in the management of chemo/radiotherapy-induced oral mucositis: A systematic review and network meta-analysis. *Int J Nurs Stud*. 2019;89:80-87. doi:10.1016/j.ijnurstu.2018.08.007

35. McLoone P, Tabys D, Fyfe L. Honey Combination Therapies for Skin and Wound Infections: A Systematic Review of the Literature. *Clin Cosmet Investig Dermatol*. 2020;13:875-888. doi:10.2147/CCID.S282143

36. Poovelikunnel TT, Gethin G, Solanki D, McFadden E, Codd M, Humphreys H. Randomized controlled trial of honey versus mupirocin to decolonize patients with nasal colonization of meticillin-resistant Staphylococcus aureus. *J Hosp Infect*. 2018;98(2):141-148. doi:10.1016/j.jhin.2017.10.016

37. Kathuria S, Singh PK, Sharma C, et al. Multidrug-Resistant Candida auris Misidentified as Candida haemulonii: Characterization by Matrix-Assisted Laser Desorption Ionization-Time of Flight Mass Spectrometry and DNA Sequencing and Its Antifungal Susceptibility Profile Variability by Vitek 2, CLSI Broth Microdilution, and Etest Method. *J Clin Microbiol*. 2015;53(6):1823-1830. doi:10.1128/JCM.00367-15

38. CLSI. Reference Method for Broth Dilution Antifungal Susceptibility Testing of Yeasts; Approved Standard. 4[th] ed. M27. Wayne (PA): *Clinical and Laboratory Standards Institute*.

39. CLSI. Performance Standards for Antifungal Susceptibility Testing of Yeasts. 2nd Ed. M60. Wayne (PA): *Clinical and Laboratory Standards Institute*.

40. Thermo Fisher ScientificTM SENSITITRE® YEASTONE® For in Vitro Diagnostic Use. MCS Diagnostics. 2012. [(accessed on 14 April 2023)]; Available online: https://docplayer.net/21149741-Sensititre-yeastone-for-in-vitro-diagnostic-use.html

41. Centers for Disease Control and Prevention, Antifungal Susceptibility Testing and

Interpretation. [(accessed on 14 April 2023]; Available online:

https://www.cdc.gov/fungal/candida-auris/c-auris-antifungal.html

42. Chan W, Chow FWN, Tsang CC, et al. Induction of amphotericin B resistance in

susceptible *Candida auris* by extracellular vesicles. *Emerging Microbes & Infections*.

2022;11(1):1900-1909. doi:10.1080/22221751.2022.2098058

43. Alserehi H, Filippell M, Emerick M, et al. Chlorhexidine gluconate bathing practices

and skin concentrations in intensive care unit patients. *American Journal of Infection*

Control. 2018;46(2):226-228. doi:10.1016/j.ajic.2017.08.022

44. Abdolrasouli A, Armstrong-James D, Ryan L, Schelenz S. In vitro efficacy of

disinfectants utilised for skin decolonisation and environmental decontamination during

a hospital outbreak with *Candida auris*. *Mycoses*. 2017;60(11):758-763.

doi:10.1111/myc.12699

45. Yu D, Wang J, Shao X, Xu F, Wang H. Antifungal modes of action of tea tree oil and

its two characteristic components against Botrytis cinerea. *J Appl Microbiol*.

2015;119(5):1253-1262. doi:10.1111/jam.12939

46. Li WR, Li HL, Shi QS, et al. The dynamics and mechanism of the antimicrobial activity

of tea tree oil against bacteria and fungi. *Appl Microbiol Biotechnol*.

2016;100(20):8865-8875. doi:10.1007/s00253-016-7692-4

47. Sahu B, R. A, Mohammed Abu Javid. Application of Calophyllum inophyllum oil as

antifungal fat-liquor for leather industry. *Industrial Crops and Products*. 2017;105:104-

112. doi:10.1016/j.indcrop.2017.04.064

48. Kim E, Park IK. Fumigant antifungal activity of Myrtaceae essential oils and

constituents from Leptospermum petersonii against three Aspergillus species.

Molecules. 2012;17(9):10459-10469. doi:10.3390/molecules170910459

49. Parker RA, Gabriel KT, Graham KD, Butts BK, Cornelison CT. Antifungal Activity of

Select Essential Oils against Candida auris and Their Interactions with Antifungal

Drugs. *Pathogens*. 2022;11(8):821. doi:10.3390/pathogens11080821

50. Campo Research PTE Ltd. 2015. Manuka Oil Extract Leptospermum Scoparium.

Campo Research PTE Ltd.; Singapore.

51. Satchell AC, Saurajen A, Bell C, Barnetson RS. Treatment of dandruff with 5% tea tree

oil shampoo. *J Am Acad Dermatol*. 2002;47(6):852-855. doi:10.1067/mjd.2002.122734

52. Herman A, Herman AP. Essential oils and their constituents as skin penetration

enhancer for transdermal drug delivery: a review. *Journal of Pharmacy and*

Pharmacology. 2015;67(4):473-485. doi:10.1111/jphp.12334

53. Fernandes L, Oliveira A, Henriques M, Rodrigues ME. Honey as a Strategy to Fight

Candida tropicalis in Mixed-Biofilms with Pseudomonas aeruginosa. *Antibiotics*

(Basel). 2020;9(2):43. doi:10.3390/antibiotics9020043

54. Mandal MD, Mandal S. Honey: its medicinal property and antibacterial activity. *Asian*

Pacific Journal of Tropical Biomedicine. 2011;1(2):154-160. doi:10.1016/S2221-

1691(11)60016-6

55. Scepankova H, Combarros-Fuertes P, Fresno JM, et al. Role of Honey in Advanced

Wound Care. *Molecules*. 2021;26(16):4784. doi:10.3390/molecules26164784

56. Sekizuka T, Iguchi S, Umeyama T, et al. Clade II Candida auris possess genomic

structural variations related to an ancestral strain. Chowdhary A, ed. *PLoS ONE*.

2019;14(10):e0223433. doi:10.1371/journal.pone.0223433

57. Brandt P, Mirhakkak MH, Wagner L, et al. High-Throughput Profiling of Candida auris

Isolates Reveals Clade-Specific Metabolic Differences. *Microbiol Spectr*.

2023;11(3):e0049823. doi: 10.1128/spectrum.00498-23

58. de Groot T, Janssen T, Faro D, Cremers NAJ, Chowdhary A, Meis JF. Antifungal

Activity of a Medical-Grade Honey Formulation against Candida auris. *J Fungi (Basel)*.

2021;7(1):50. doi:10.3390/jof7010050

LEGENDS TO TABLE AND FIGURE

521   **Table 1:** In vitro susceptibility of *Candida* isolates to synthetic antiseptics, natural antiseptics and their commercial products

	Clinical *Candida auris* isolates (N=40)				*Candida auris* JCM 15448		*Candida krusei* ATCC 6258		*Candida parapsilosis* ATCC 22019		*Candida albicans* ATCC 90028		*Candida glabrata* ATCC MYA-2950	
	MIC_{50}	MIC_{90}	MIC range	MFC range	MIC	MFC	MIC	MFC	MIC	MFC	MIC	MFC	MIC	MFC
Chlorhexidine (%)	0.001	0.001	0.0005-0.001	0.002-0.004	0.001	0.002	0.0002	0.0005	0.00006	0.0002	0.0005	0.001	0.0005	0.0005
Povidone-iodine (%)	0.078	0.078	0.039-0.078	0.078-0.313	0.039	0.16	0.039	0.31	0.078	0.625	0.16	0.31	0.16	0.31
Nystatin (µg/mL)	2.44	2.44	1.22-2.44	2.44-9.77	1.22	1.22	1.22	2.44	0.61	2.44	1.22	4.88	1.22	2.44
Manuka honey 15+ (Comvita) (v/v) (%)	25	25	6.25-25	>50	0.20	>50	25	>50	25	>50	25	>50	1.56	>50
Manuka honey 20+ (Comvita) (v/v) (%)	25	25	3.13-25	>50	0.20	>50	12.5	>50	25	>50	25	>50	1.56	>50
Manuka honey 15+ (Kiwi Manuka) (v/v) (%)	25	25	3.13-25	>50	0.20	>50	12.5	>50	25	>50	25	>50	1.56	>50
Manuka honey 20+ (Kiwi Manuka) (v/v) (%)	25	25	6.25-25	>50	0.20	>50	12.5	>50	25	>50	25	>50	1.56	>50
Tea tree oil 15% (Body Shop) (v/v) (%)	0.004	0.004	0.004	2.08-4.17	0.004	1.04	0.002	2.08	0.004	4.17	0.039	0.63	0.16	0.63
Tea tree oil 100% (Kiwi Manuka) (v/v) (%)	0.31	0.31	0.31	0.63-1.25	0.31	0.63	0.63	1.25	0.63	1.25	0.63	1.25	0.63	1.25
Manuka oil 100% (Manuka Lab) (v/v) (%)	0.31	0.31	0.16-0.31	0.63	0.08	0.63	0.31	1.25	0.31	1.25	0.63	2.5	0.078	0.63
Manuka oil 100% (Melora) (v/v) (%)	0.31	0.31	0.31	0.63	0.31	0.31	0.31	1.25	0.31	1.25	1.25	5.0	0.078	0.63
Manuka hydrosol (v/v) (%)	>50	>50	>50	>50	>50	>50	>50	>50	>50	>50	>50	>50	>50	>50
Manuka body wash (w/v) (%)	0.01	0.01	0.006-0.01	0.05-0.10	0.006	0.02	0.006	0.05	0.006	3.13	0.024	0.098	0.049	1.56
Tea tree body wash (w/v) (%)	0.012	0.012	0.006-0.39	≥50	0.003	25	0.002	25	0.003	50	0.049	25	0.20	>50
MEDIHONEY wound gel (w/v) (%)	3.13	3.13	1.56-6.25	≥25	0.20	12.5	0.39	25	0.20	>37.5	6.25	>37.5	0.098	37.5
Manuka mouthwash (v/v) (%)	0.39	0.39	0.20-0.39	3.13-6.25	0.39	6.25	0.10	6.25	0.20	6.25	1.56	6.25	1.56	6.25

524 **Figure 1:** The fungistatic effect of natural antiseptics on *Candida auris*. Three *Candida auris* isolates were incubated with 4 types of manuka

525 honey, tea tree body wash – 25%, 37.5% and 50%, and antibacterial wound gel – 25% and 37.5% (50% wound gel is not available for testing due

526 to its insolubility at high concentrations). Log reductions in CFU were shown. Significant differences ($p < 0.05$) are indicated with an asterisk.

528   **Table 2:** Antifungal drug susceptibility of *Candida auris* isolates

	Clinical *C. auris* isolates (N=40)				*Candida auris* JCM 15448
	MIC$_{50}$ (mg/L)	MIC$_{90}$ (mg/L)	MIC range (mg/L)	Resistance, n (%)	MIC (mg/L)
Amphotericin B	2	4	1-4	36 (90)	1
Fluconazole	256	>256	64 - >256	40 (100)	8
Anidulafungin	0.25	0.25	0.12 - 8	1 (2.5)	0.12
Micafungin	0.25	0.25	0.06 - 1	0 (0)	0.06
Caspofungin	0.25	0.5	0.06 - >8	3 (7.5)	0.03
Itraconazole	0.12	0.25	0.06 - >16	NA	0.06
Voriconazole	0.5	1	0.12 - >8	NA	0.06
Posaconazole	0.03	0.06	0.015 - >8	NA	0.03
5-flucytosine	0.12	0.12	<0.06 - >64	NA	0.25

530   CDC's tentative nonsusceptible breakpoints: ≥2 mg/L for amphotericin B, ≥32 mg/L for fluconazole, ≥4 mg/L for echinocandins anidulafungin

531   and micafungin and ≥2 mg/L for caspofungin. For the triazoles itraconazole, voriconazole and posaconazole, and 5-flucytosine, no breakpoints are

532   available. NA: Not available.