Antifungal efficacy of natural antiseptic products against *Candida auris*

Wing-Gi WU[a#], Kristine Shik LUK[a,b,c,d], Mei-Fan HUNG[a], Wing-Yi TSANG[a], Kin-Ping LEE[a,c], Bosco Hoi-Shiu LAM[a], Ka-Lam CHENG[c], Wing-Sze CHEUNG[b], Hau-Ling TANG[d], Wing-Kin TO[a,b]

[a]Department of Pathology, Princess Margaret Hospital, Hong Kong Special Administrative Region, China
[b]Infection Control Team, Princess Margaret Hospital, Hong Kong Special Administrative Region, China
[c]Infection Control Team, Caritas Medical Centre, Hong Kong Special Administrative Region, China
[d]Infection Control Team, Yan Chai Hospital, Hong Kong Special Administrative Region, China

#Correspondence to: Dr Wing-Gi WU, Department of Pathology, Princess Margaret Hospital, 2-10 Princess Margaret Hospital Road, Lai Chi Kok, Kowloon, Hong Kong. Email: wwg396@ha.org.hk

Keywords: Candida auris, natural, decolonization, tea tree, manuka

ABSTRACT

Candida auris is an emerging fungal pathogen responsible for healthcare associated infections and outbreaks with high mortality around the world. It readily colonizes the skin, nares, respiratory and urinary tract of hospitalized patients, and such colonization may lead to invasive *Candida* infection in susceptible patients. However, there is no recommended decolonization protocol for *C. auris* by international health authorities. The aim of this study is to evaluate the susceptibility of *C. auris* to commonly used synthetic and natural antiseptic products using an in vitro, broth microdilution assay. Synthetic antiseptics including chlorhexidine, povidone-iodine, and nystatin were shown to be fungicidal against *C. auris*. Among the natural antiseptics tested, tea tree oil and manuka oil were both fungicidal against *C. auris* at concentrations less than or equal to 1.25% (v/v). Manuka honey inhibited *C. auris* at 25% (v/v) concentrations. Among the commercial products tested, manuka body wash and mouthwash were fungicidal against *C. auris* at concentrations less than or equal to 0.39% (w/v) and 6.25% (v/v) of products as supplied for use, respectively, while tea tree body wash and Medihoney™ wound gel demonstrated fungistatic properties. In conclusion, this study demonstrated good in vitro antifungal efficacy of tea tree oil, manuka oil, manuka honey, and commercially available antiseptic products containing these active ingredients. Future studies are warranted to evaluate the effectiveness of these antiseptic products in clinical settings.

INTRODUCTION

The fungal pathogen *Candida auris* was first described in 2009 after isolation from the ear of a patient in Japan [1]. It has since been reported in more than 40 countries across six continents, causing a wide range of healthcare-associated outbreaks and infections [2], with the mortality rate varying from 30 to 60%, due to the high virulence and resistance to several antifungals [3]. The recent inaugural World Health Organization fungal priority pathogen list highlights the urgent threat of *C. auris* to global public health [4]. With its biofilm-forming ability and tolerance to high temperature and salt concentrations, *C. auris* can persist and survive in the human skin and environmental surfaces for several months following contact with colonized patients [5-7]. This provides additional opportunities for transmission and colonization within the healthcare settings. Furthermore, the minimum time to acquire *C. auris* from a patient or their immediate environment is 4 hours or less, reinforcing the importance of vigilant infection control measures [8]. This fungal superbug can also resist certain disinfectants and is well adapted to healthcare environments [9].

In 2019, a cluster of *C. auris* colonization affecting 15 patients was reported in Princess Margaret Hospital (PMH), Kowloon West (KW) cluster, Hong Kong. Whole genome sequencing identified the strains as Clade I/South Asia, associated with ERG11 mutation conferring azole resistance [10]. The clade is also known to cause invasive infections and outbreaks frequently [6]. Four regional hospitals (a total of 3,878 beds) serve an estimated population of 1.4 million in the KW cluster. In spite of enhanced surveillance and infection control measures, *C. auris* cases and transmission have continued to increase due to pandemic-related strain on the healthcare system. As of April 2023, there were 364 cases of *C. auris* reported in KW Cluster, Hong Kong.

C. auris readily colonizes the axillae, groin, nares, respiratory, and urinary tract of hospitalized patients [6]. Compared to other *Candida* species, gastrointestinal tract colonization

is less frequent for *C. auris*. [11]. Colonization predisposes to invasive infection, with mechanical ventilation and placement of invasive devices identified as major risk factors [6]. A descriptive analysis of *C. auris* outbreaks in New York State reported that 4% of *C. auris* colonized patients developed bloodstream infection, with a rate of 0.3 per 1,000 patient-days [12]. These studies provided evidence that an important strategy for preventing severe candidiasis is the elimination of *C. auris* carriage to preclude subsequent risk of endogenous infection. Nevertheless, there are no well-established guidelines for *C. auris* decolonization. For this reason, more evidence from in vitro antifungal susceptibility testing of topical antiseptics against *C. auris* is needed to provide the basis of decolonization regimen before clinical trials. Synthetic antiseptics including chlorhexidine, povidone-iodine, isopropyl alcohol, and nystatin have been most studied for their antifungal activities against *C. auris* in vitro [13-15]. While being one of the commonly used disinfectants in healthcare settings, chlorhexidine has been shown to have variable effectiveness for *C. auris* depending on the formulation and in vitro testing methods. *Sherry et al.* and *Abdolrasouli et al.* showed chlorhexidine was effective for *C. auris* using broth dilution antifungal susceptibility testing [16-17]. *Moore et al.* reported that 2% chlorhexidine solution with isopropyl alcohol formation was effective in killing *C. auris* with a 2-min contact time, while chlorhexidine alone without isopropyl alcohol failed to kill *C. auris* [13].

By far, most clinical experience of *C. auris* decolonization was reported in outbreak settings as part of the infection control measures. *Schelenz et al.* first reported a decolonization regimen using a five-day protocol with twice-daily 2% chlorhexidine body washes, 0.2% chlorhexidine mouthwash, and oral nystatin during hospital outbreak management [8]. Based on this early experience, decolonization on our *C. auris* colonized cohorts using a protocol of twice-daily chlorhexidine bath, chlorhexidine mouthwash and povidone iodine applied to axilla, groin and nares was attempted as an outbreak control measure. Frequent chlorhexidine baths

were poorly tolerated. More importantly, none of our patients were successfully decolonized (0/14 patients; unpublished). In addition, subsequent studies have reported poor clinical efficacy of chlorhexidine-based regimens for decolonizing *C. auris,* with high rates of persistent colonization [9]. A murine model study demonstrated that *C. auris* could enter the dermis without causing overt histopathologic signs of inflammation, resulting in deeper skin layer residence [18]. Another study based on ex vivo human skin model has illustrated chlorhexidine had limited permeation into deeper skin layers [19], and this might contribute to persistent *C. auris* colonization despite its in vitro efficacy.

The limited success of traditional decolonization using chlorhexidine has highlighted the need to explore alternative regimens and approaches. Natural substances with antimicrobial properties are potentially good candidates as their antimicrobial effects are based on multiple compounds and would not mount drug-selecting pressure. They are also well tolerated as skin washes in general. *Fernades et al.* have shown that tea tree (*Melalenca alternifolia*) oil had good in vitro antifungal activity against *C. auris* in both planktonic cells and biofilms [20]. *Johnson et al.* have also shown that the active components of tea tree oil could augment the activity of chlorhexidine for *C. auris* decolonization in the porcine skin model [21]. Another potential natural antiseptic is manuka (*Leptospermum scoparium*), which is also known as the "New Zealand tea tree". It has been a part of the traditional Māori medicine with fungicidal properties towards *C. albicans* and *C. glabrata* [22]. Notably, manuka honey was found to be effective in inhibiting various *Candida* species [23]. There are multiple antimicrobial constituents of the natural extracts, namely monoterpenes and sesquiterpenes (tea tree and manuka oil), triketones (manuka oil), methylglyoxal, polyphenolic compounds and bee defensin-1 (manuka honey) [24-26]. Therefore, unlike other synthetic antimicrobials, there have been no reports of clinical isolates with acquired resistance to the natural antiseptics [27-28]. Clinical trials have demonstrated the efficacy of manuka honey and tea tree oil in the

treatment of oral candidiasis, vulvovaginal candidiasis and successful decolonization for methicillin-resistant *Staphylococcus aureus* (MRSA) carriers [29-34], making them attractive alternatives against *C. auris*. These natural components are also widely used as active ingredients for their antiseptic effects in commercial hygiene products such as body wash, mouthwash and wound gel. To pave the way for evaluating the clinical application of the natural antiseptics against *C. auris*, this study aimed to determine the in vitro antifungal susceptibility of *C. auris* to the natural antiseptics and their commercial products.

MATERIALS AND METHODS

Characterization of *Candida auris* isolates: A total of 40 *C. auris* isolates belonging to clade I/South Asia (unpublished data) were obtained in the KW Cluster of Hong Kong from 2019 to 2022. All tested isolates were identified by matrix-assisted laser desorption ionization-time of flight mass spectrometry (Bruker Daltonics; Database version: IVD-9468 MSP) [35]. Reference strains *C. auris* JCM 15448 (clade II/East Asia), *C. auris* ATCC MYA-5002 (clade III/South Africa), *C. auris* ATCC MYA-5003 (clade IV/South America), *C. albicans* ATCC 90028, *C. glabrata* ATCC MYA-2950, *C. krusei* ATCC 6258, and *C. parapsilosis* ATCC 22019 were included. The study protocol was approved by the Research Ethics Committee of the Kowloon West Cluster, Hospital Authority, Hong Kong (KW/EX-21-098(160-14)).

Antiseptics and preparations: Our experiment studied three groups of topical antiseptics. The first group included synthetic antiseptics products commonly used in healthcare settings: chlorhexidine gluconate (Microshield® 4%, Australia), povidone-iodine (Betadine® 10%, Turkey) and nystatin (100,000 units/mL, Pharmascience, Canada). The second group comprised natural antiseptics: tea tree oil (15%, The Body Shop, UK; 100% Kiwi Manuka®, New Zealand), manuka oil (100%, Manuka Lab®, New Zealand; 100%, Melora®, New Zealand), manuka honey (UMF 15+ and 20+, Comvita®, New Zealand; UMF 15+ and 20+, Melora®, New Zealand), manuka hydrosol (Bioactives®, New Zealand). The third group consisted of commercial antiseptic products containing active components from tea tree oil and/or manuka derivatives: manuka body wash (Manuka Lab®, New Zealand), tea tree clearing body wash (The Body Shop®, UK), manuka mouthwash (Manuka Lab®, New Zealand), and Medihoney™ antibacterial wound gel (Comvita®, New Zealand). Dimethyl sulfoxide (DMSO, Sigma-Aldrich, UK) at 2% was added to water-insoluble products, including tea tree oil, manuka oil, antibacterial wound gel, and nystatin to enhance solubility in RPMI 1640 medium (Sigma, Paisley, UK). The effects of 2% DMSO in growth control was also studied to confirm no

antifungal activity was added by the solvent.

In vitro antifungal susceptibility of antiseptics: In vitro antifungal susceptibility of antiseptics was performed by broth microdilution method according to CLSI M27-A4 [36]. Briefly, 100 µL of RPMI 1640 medium with different concentrations of the antiseptics were added to 96-well microtiter plates (Saining Biotechnology, China). *Candida* isolates were suspended in 2 mL sterile physiological saline to obtain 0.5 McFarland standard suspension using the DensiCHEK Plus instrument (BioMerieux, France), corresponding to 1-5 x 10^6 cells per mL, then further diluted with RPMI 1640 medium to obtain a 1:1000 final inoculum containing 1-5 x 10^3 cells per mL. The yeast suspensions (100 µL) were added to the 96-well plate with the product suspensions (100 µL) and incubated aerobically at 35°C for 24 h. The minimum inhibitory concentration (MIC) endpoint was determined visually and defined as the lowest concentration of a product that caused a 50% decrease in growth (prominent decrease in turbidity) relative to the growth in the control well. To assess the minimal fungicidal concentration (MFC) endpoint, the broth was first assessed for any visible growth. For those without visible growth, 100 µL of broth was taken from the wells and inoculated onto Sabouraud dextrose agar (BD DIFCOTM, USA). Following incubation at 35°C for 48 h, the colony-forming unit (CFU) was counted to assess viability. The antiseptic was defined as fungicidal if there was no growth on inoculated plates, whereas it was defined as fungistatic if it inhibited growth but not fungicidal. Antifungal was defined as a product with either fungistatic or fungicidal effects. Each isolate was tested in duplicate separately and the assay was repeated if the MIC or MFC differed by more than one dilution. Antifungal drug susceptibility of *C. krusei* ATCC 6258 and *C. parapsilosis* ATCC 22019 was performed in each run of broth microdilution as controls [37].

For antiseptics demonstrating fungistatic effects on clinical *C. auris* isolates, log growth reduction in CFU compared to the growth control (yeast suspension in RPMI) after 24-hour

incubation was measured, and the test was repeated in triplicate. The yeast suspension preparation and incubation procedures were the same as the aforementioned section. Mann-Whitney U test was used to analyze the significance of difference in CFU count compared to the growth control, and one-way ANOVA, Kruskal-Wallis, or Mann-Whitney U tests were used to compare the log growth reduction in CFU by the antiseptics at different concentrations. Pairwise comparisons were performed using Bonferroni or Dunn's test when suitable. P value <0.05 was considered statistically significant. All statistical analyses were performed using STATA18.0 (StataCorp. 2023. Stata Statistical Software: Release 18. College Station, TX: StataCorp LLC.).

In vitro antifungal susceptibility testing of antifungal drugs: To confirm the local resistance pattern of our isolates to conventional antifungal drugs, in vitro susceptibility was determined using Sensititre^{TM} YeastONE^{TM} YO10 AST Plate (ThermoFisher Scientific, UK). The antifungal drugs included anidulafungin, micafungin, caspofungin, 5-flucytosine, posaconazole, voriconazole, itraconazole, fluconazole and amphotericin B. The test was performed according to the procedures described by the manufacturer [38]. Centres for Disease Control and Prevention (CDC)'s tentative breakpoints for *C. auris* were used as a reference [39].

RESULTS

Table 1 summarizes the antifungal activities of topical antiseptics against *Candida auris*. For the synthetic antiseptics, chlorhexidine, povidone-iodine and nystatin were fungicidal against *C. auris* at concentrations less than or equal to 21.99 mg/L (0.004%), 312.50 mg/L (0.313%), and 9.77 μg/mL (0.049%), respectively. The MICs and MFCs of chlorhexidine against *C. auris* strains, including the JCM 15448, were relatively higher when compared with that against *C. krusei* ATCC 6258, *C. parapsilosis* ATCC 22019, *C. albicans* ATCC 90028, and *C. glabrata* ATCC MYA-2950.

For the natural antiseptics, whilst manuka honey exhibited fungistatic properties against clinical *C. auris* isolates at 25% (v/v), it was not fungicidal. Pure (equates to 100% in neat formula) tea tree and manuka oil were fungicidal against clinical *C. auris* isolates at concentrations less than or equal to 1.25% and 0.63% (v/v), respectively. Commercial products containing active ingredients of manuka honey, manuka oil and/or tea tree oil were shown to be active in vitro against *C. auris* and other *Candida* species. Manuka body wash and mouthwash were fungicidal against *C. auris* at concentrations less than or equal to 0.39% (w/v) and 6.25% (v/v), respectively. Manuka body wash, however, exhibited weaker antifungal activity against *C. parapsilosis* [MFC 3.13% (w/v)]. Interestingly, the MICs of manuka honey against *C. auris* JCM 15448 (clade II) were 128-fold lower than those against the clinical counterparts (clade I) and the two ATCC strains of clades III and IV. Tea tree body wash and Medihoney™ wound gel even became fungicidal against *C. auris* JCM 15488 at concentrations of 25% (w/v) and 12.5% (w/v), respectively.

Manuka honey [25-50% (v/v)], tea tree body wash [25-50% (w/v)] and Medihoney™ wound gel [25-37.5% (w/v)] demonstrated fungistatic properties against clinical *C. auris* isolates in vitro, but they were not fungicidal. They resulted in a statistically significant log reduction in CFU compared to the growth control ($p < 0.01$). The growth control yielded around

1.5-5 x 10^7 CFU/mL after 24 h incubation. At a concentration of 37.5%, tea tree body wash and the Medihoney™ wound gel achieved >4 log reduction in CFU while manuka honey achieved 2 to 3 log reduction in CFU after 24 h of incubation, as shown in Figure 1. The inhibitory effects of manuka honey and tea tree body wash against *C. auris* were concentration-dependent ($p < 0.05$). Medihoney™ wound gel at 25% and 37.5% resulted in a similar reduction (6-log) in CFU.

The antifungal susceptibility of common systemic antifungals against 40 clinical *C. auris* strains (clade I) was shown in table 2. All tested isolates (N=40) were resistant to fluconazole and 90% (N=36) were also resistant to Amphotericin B. Susceptibility to other azoles and echinocandins was variable.

DISCUSSION

The emergence of multidrug-resistant *Candida auris* having a high propensity to cause nosocomial transmission render the urgency to develop a novel and effective decolonization strategy. Herein, we demonstrated excellent in vitro fungicidal and/or fungistatic activity of natural antiseptics, including manuka honey, manuka oil and tea tree oil, and their commercial products, against *C. auris*. This suggests the current decolonization strategy can be much improved by combining these well-tolerated, natural antiseptics. A clinical trial on *C. auris* decolonization using manuka body wash, manuka mouthwash, tea tree oil, povidone-iodine, and nystatin oral suspension in 41 patients achieved about 60% success rate and no adverse effects were reported (*Luk*, 2023, unpublished data).

In the present study, the local *C. auris* isolates were highly resistant to both fluconazole (MIC$_{50}$ of 256 µg/mL) and amphotericin B (MIC$_{50}$ of 2 µg/mL). Genomic and proteomic analyses revealed mutation of the fluconazole target enzyme (cytochrome 450 lanosterol 14α-demethylase) [10] and the role of extracellular vesicles in the development of amphotericin B resistance [40].

Our results confirmed the high susceptibility of *C. auris* to 4% chlorhexidine (MFC 0.004%), 10% povidone iodine (MFC 0.312%) and 33.3 µg/mL (100,000 units/mL) nystatin (MFC 9.77 µg/mL) at concentrations used for clinical indications [17, 41]. Nonetheless, the MICs an MFCs of chlorhexidine against *C. auris* strains were higher than those against other *Candida* species. Similar findings were reported by *Abdolrasouli et al* [17], showing that their *C. auris* outbreak strains were less susceptible to chlorhexidine than other *Candida* species though the testing methods were different and the mechanism is currently unknown. Likewise, our clinical *C. auris* strains were highly susceptible to tea tree oil [MFC 1.25% (v/v)] and manuka oil [MFC 0.63% (v/v)]; their fungicidal effects were comparable to synthetic antiseptics, including chlorhexidine, povidone-iodine and nystatin. Our findings are consistent

with a previous report of potent antifungal activity of tea tree oil against *C. auris* isolates (MFC 1.56% (w/v)) [20]. Tea tree oil and its main component, terpinene-4-ol, have been shown to cause damage to the fungal cell wall and cytoplasmic membrane thereby causing cellular dysfunctions and death though the exact mechanism is unknown [42- 43]. Interestingly, The Body Shop® formulation contains a diluted content of 15% tea tree oil but a lower MIC_{90} of 0.004% (v/v) was observed compared to that of 100% tea tree oil [0.31% (v/v)]. This enhanced inhibition may be attributed to the synergistic or additive interactions of the supplementary ingredients, including denatured alcohol, *Calophyllum inophyllum* seed oil, and *Leptospermum petersonii* oil, which were shown to have antimicrobial properties [44-45]. *Moore et al.* also showed the addition of isopropyl alcohol to disinfectant formulation enhanced the killing of *C. auris* [13]. On the other hand, the antifungal properties of manuka oil were less frequently studied for *C. auris* specifically. *Parker et al.* demonstrated significant inhibitory effects of manuka oil on *C. auris* AR0381 (clade II) and AR0385 (clade IV), with MIC values less than 1.0% [46], of which the findings agree well with ours. More importantly, the low fungicidal concentrations of tea tree oil and manuka oil against *C. auris* are considered safe and tolerable for dermal use. An in vitro skin model proved manuka oil is non-irritative, with cell viability unaffected after treatment with an undiluted sample [47]. Another clinical trial also demonstrated that daily use of 5% tea tree oil shampoo was well tolerated in the treatment for dandruff [48]. The additional property of enhanced skin permeation of essential oils [49] further supports the role of essential oils in eliminating *C. auris* colonization.

Regarding manuka honey, *C. auris* isolates were modestly inhibited (1.5 log reduction in CFU) at a concentration of 25%. The growth inhibition was presented in a dose-dependent manner. This accords with several studies showing fungistatic effect of manuka honey against *Candida* species. [23, 50]. Manuka honey belongs to the "non-peroxide-based" honey group, and its main antimicrobial action is contributed by methylglyoxal (MGO) [51], which is

converted non-enzymatically from the dihydroxyacetone present in the nectar of the flowers of *Leptospermum scoparium*. The characteristically high level of MGO in manuka honey makes it a unique marker for authentication. In this study, manuka honey with greater Unique Manuka Factor (UMF), which is correlated with the MGO and total phenol components [52], resulted in greater CFU reduction of *C. auris*. Increasing the concentration of honey from 25% to 50% led to an average 3-log reduction in the CFU of *C. auris* ($p < 0.05$). Although the inhibitory effect of manuka honey against *C. auris* is modest, honey is seldom applied topically in a clinical setting. Alternatively, honey can act as an important ingredient in the manufacturing of antiseptic products. Of note, the *C. auris* JCM 15448 (clade II) had a much lower MIC (0.2% v/v) towards manuka honey. Similar phenomenon was also observed for manuka oil, tea tree body wash and Medihoney™ wound gel. Comparative genomic analysis revealed that multiple genes involving in cell wall proteins and stress response-related functions were absent in the Japanese *C. auris* strains, including JCM 15448, while a more recent phenotypic analysis also supported clade-specific phenotypic profiles of *C. auris* with clade II having the lowest tolerance to osmotic stress and chemicals [53-54]. These genomic structural variations and metabolic profiles may explain the differences in antifungal resistance, environmental survival, and virulence among different *C. auris* clades. Nevertheless, *de* Groot T *et al.* did not report any difference in antifungal activity of medical grade honey among *C. auris* clades [55]. Studies including larger numbers of *C. auris* isolates of different clades should determine whether susceptibility to natural antiseptics is clade specific.

As expected, commercial products having the active ingredients of natural antiseptics showed promising antifungal effect in vitro against *C. auris*. Manuka body wash and mouthwash were fungicidal against *C. auris* at low concentrations. Product labels of manuka body wash and mouthwash disclose one of the ingredients being 525+ MGO, which is equivalent to UMF 15+ manuka honey. Besides, the aforementioned products contain tea tree

and manuka oil with antifungal effects demonstrated in our study. In contrast, Medihoney™ antibacterial wound gel containing 80% medical grade manuka honey and The Body Shop® tea tree body wash were not fungicidal against *C. auris* at concentrations tested though they caused a 6-log reduction in *C. auris* CFU. In addition, the anti-inflammatory and anti-oxidant properties of manuka honey [28] confer the commercial hygiene products as prominent options in the development of an efficacious and tolerable decolonization regimen.

In vitro antifungal susceptibility testing was performed by broth microdilution assay in this study, which is traditionally the gold standard. The MIC and MFC endpoints obtained from broth dilution allows comparison with conventional antifungal agents, for which their clinical correlation has been established; and comparison among different antiseptics or *Candida* species when MIC or MFC data are available. This is particularly useful for evaluating novel antifungal agents for an emerging pathogen. In particular, only limited data of in vitro antifungal efficacy of manuka honey and essential oils are currently available for *Candida* species, including *C. auris*. Our MIC and MFC findings of natural antiseptics were consistent with previous studies using broth dilution assays [20, 23, 46, 50]. Nevertheless, this testing method has some limitations. Contact time exposure is not taken into account during in vitro testing. The exposure time of disinfectants or topical antiseptics in use is often much shorter. Inhibitory and fungicidal endpoints of antiseptics using broth microdilution at 24 hours might therefore overestimate the clinical effectiveness. Alternatively, a time-dependent quantitative assay seems to reflect more about the real-life situation. *Moore et al.* has comprehensively described how to translate the contact time and dilution considerations of antiseptics from clinical use to in vitro testing for *C. auris* [13]. The antimicrobial wash, chlorhexidine 4%, was diluted with water to produce a 2% solution in order to simulate the bath process. When tested with contact time ≤ 2 min, 2% chlorhexidine without isopropyl alcohol failed to eliminate *C. auris*, which was in contrast to the susceptibility demonstrated by broth microdilution after overnight

incubation. This highlights the importance of time exposure in the evaluation of antiseptics with regard to the actual use in clinical settings. Moreover, the contact time should be well controlled by neutralizing the antimicrobial activity in vitro after the exposure. *Sutton et al.* has proposed the criteria for identifying the appropriate neutralizing substances, which are specific to the antimicrobial agents and the organisms challenged [56]. Interestingly, a previous study used glyoxalase to neutralize MGO for its main antimicrobial effects in manuka honey. MGO-neutralized manuka honey retained activity against various organisms, indicating the presence of other antimicrobial constituents besides MGO [57]. In addition, most neutralization tests were performed on disinfectants against bacterial organisms and how these results could be applied to fungal organisms was not known. Similarly, there is no universally-agreed neutralizing agent for most natural antiseptics. Alternative approach using membrane filtration or dilution is also unproven on natural antiseptics. Future studies on the active constituents of natural antiseptics will facilitate the validation of appropriate neutralization tests and development of time-dependent assays.

There are some limitations in this study. First, extraction and evaluation of the bioactive chemical markers in the natural antiseptics were not performed. The specific quantities of the active ingredients were often not disclosed in the product labels. The antifungal effects could be contributed by ingredients other than the natural antiseptics we tested individually. We could only tell the overall in vitro antifungal efficacy of the antiseptic products instead of each constituent. Besides, all our *C. auris* isolates belonged to clade I and the results of this study may not be applicable to isolates of other clades. We had included *C. auris* JCM 15448 (clade II/East Asia), ATCC MYA-5002 (clade III/South Africa), and ATCC MYA-5003 (clade IV/South America) in our testing, and found JCM 15448 had significantly lower MIC and/or MFC values towards several natural antiseptics and their commercial products. Moreover, the effect of antifungal drug resistance on the antiseptics could not be studied due to the low

variability of the resistance pattern of our *C. auris* isolates. Nonetheless, these antiseptic products had an overall high in vitro antifungal efficacy. Future development of an appropriate time-dependent assay can provide more evidence to support their application as topical or decolonization therapies for *C. auris*.

CONCLUSION

C. auris are highly susceptible to natural antiseptics including tea tree oil, manuka oil and manuka honey, and their commercial hygiene products in the forms of body wash, mouthwash, and wound gel. The promising in vitro results provide insight into the development of novel and effective *C. auris* decolonization regimen. Future studies should demonstrate whether these natural antiseptics are effective in clinical settings.

ACKNOWLEDGEMENT

The funding of this work was supported by the Clinical Research Centre Mini Research Grant 2023 (2023-CRC-MRG-02), Kowloon West Cluster, Hospital Authority, Hong Kong. We wish to thank the hospital infection control teams in the KW cluster for their efforts in *C. auris* control and the microbiology laboratory staff of Princess Margaret Hospital for the antifungal testing.

CONFLICT OF INTEREST

We declare no conflict of interest.

DATA AVAILABILITY

All data are incorporated into the article and its online supplementary material.

REFERENCE

1. Satoh K, Makimura K, Hasumi Y, Nishiyama Y, Uchida K, Yamaguchi H. *Candida auris* sp. nov., a novel ascomycetous yeast isolated from the external ear canal of an inpatient in a Japanese hospital. *Microbiology and Immunology*. 2009;53(1):41-44. doi:10.1111/j.1348-0421.2008.00083.x

2. de Cássia Orlandi Sardi J, Silva DR, Soares Mendes-Giannini MJ, Rosalen PL. Candida auris: Epidemiology, risk factors, virulence, resistance, and therapeutic options. *Microb Pathog*. 2018;125:116-121. doi:10.1016/j.micpath.2018.09.014

3. Al-Rashdi A, Al-Maani A, Al-Wahaibi A, Alqayoudhi A, Al-Jardani A, Al-Abri S. Characteristics, Risk Factors, and Survival Analysis of Candida auris Cases: Results of One-Year National Surveillance Data from Oman. *JoF*. 2021;7(1):31. doi:10.3390/jof7010031

4. Fisher MC, Denning DW. The WHO fungal priority pathogens list as a game-changer. *Nat Rev Microbiol*. 2023;21(4):211-212. doi:10.1038/s41579-023-00861-x

5. Piedrahita CT, Cadnum JL, Jencson AL, Shaikh AA, Ghannoum MA, Donskey CJ. Environmental Surfaces in Healthcare Facilities are a Potential Source for Transmission of *Candida auris* and Other *Candida* Species. *Infect Control Hosp Epidemiol*. 2017;38(9):1107-1109. doi:10.1017/ice.2017.127

6. Ahmad S, Alfouzan W. Candida auris: Epidemiology, Diagnosis, Pathogenesis, Antifungal Susceptibility, and Infection Control Measures to Combat the Spread of Infections in Healthcare Facilities. *Microorganisms*. 2021;9(4):807. doi:10.3390/microorganisms9040807

7. Berrio I, Caceres DH, Coronell R W, et al. Bloodstream Infections With *Candida auris* Among Children in Colombia: Clinical Characteristics and Outcomes of 34 Cases. *Journal of the Pediatric Infectious Diseases Society*. 2021;10(2):151-154. doi:10.1093/jpids/piaa038

8. Schelenz S, Hagen F, Rhodes JL, et al. First hospital outbreak of the globally emerging Candida auris in a European hospital. *Antimicrob Resist Infect Control*. 2016;5(1):35. doi:10.1186/s13756-016-0132-5

9. Biswal M, Rudramurthy SM, Jain N, et al. Controlling a possible outbreak of Candida auris infection: lessons learnt from multiple interventions. *Journal of Hospital Infection*. 2017;97(4):363-370. doi:10.1016/j.jhin.2017.09.009

10. Tse H, Tsang AKL, Chu YW, Tsang DNC. Draft Genome Sequences of 19 Clinical Isolates of Candida auris from Hong Kong. *Microbiol Resour Announc*. 2021;10(1):e00308-20. doi:10.1128/MRA.00308-20

11. Piatti G, Sartini M, Cusato C, Schito AM. Colonization by Candida auris in critically ill patients: role of cutaneous and rectal localization during an outbreak. *Journal of Hospital Infection*. 2022;120:85-89. doi:10.1016/j.jhin.2021.11.004

12. Southwick K, Adams EH, Greenko J, et al. 2039. New York State 2016–2018: Progression from *Candida auris* Colonization to Bloodstream Infection. *Open Forum Infectious Diseases*. 2018;5(suppl_1):S594-S595. doi:10.1093/ofid/ofy210.1695

13. Moore G, Schelenz S, Borman AM, Johnson EM, Brown CS. Yeasticidal activity of chemical disinfectants and antiseptics against Candida auris. *J Hosp Infect*. 2017;97(4):371-375. doi:10.1016/j.jhin.2017.08.019

14. Ku TSN, Walraven CJ, Lee SA. Candida auris: Disinfectants and Implications for Infection Control. *Front Microbiol*. 2018;9:726. doi:10.3389/fmicb.2018.00726

15. Voorn MG, Kelley AM, Chaggar GK, Li X, Teska PJ, Oliver HF. Contact time and disinfectant formulation significantly impact the efficacies of disinfectant towelettes against Candida auris on hard, non-porous surfaces. *Sci Rep*. 2023;13(1):5849. doi:10.1038/s41598-023-32876-y

16. Sherry L, Ramage G, Kean R, et al. Biofilm-Forming Capability of Highly Virulent, Multidrug-Resistant Candida auris. *Emerging Infectious Diseases*. 2017;23(2):328-331. doi: 10.3201/eid2302.161320

17. Abdolrasouli A, Armstrong-James D, Ryan L, Schelenz S. In vitro efficacy of disinfectants utilised for skin decolonisation and environmental decontamination during a hospital outbreak with *Candida auris*. *Mycoses*. 2017;60(11):758-763. doi:10.1111/myc.12699

18. Huang X, Hurabielle C, Drummond RA, et al. Murine model of colonization with fungal pathogen Candida auris to explore skin tropism, host risk factors and therapeutic

strategies. *Cell Host & Microbe*. 2021;29(2):210-221.e6. doi:10.1016/j.chom.2020.12.002

19. Karpanen TJ, Worthington T, Conway BR, Hilton AC, Elliott TSJ, Lambert PA. Penetration of Chlorhexidine into Human Skin. *Antimicrob Agents Chemother*. 2008;52(10):3633-3636. doi:10.1128/AAC.00637-08

20. Fernandes L, Ribeiro R, Costa R, Henriques M, Rodrigues ME. Essential Oils as a Good Weapon against Drug-Resistant Candida auris. *Antibiotics*. 2022;11(7):977. doi:10.3390/antibiotics11070977

21. Johnson CJ, Eix EF, Lam BC, et al. Augmenting the Activity of Chlorhexidine for Decolonization of Candida auris from Porcine skin. *JoF*. 2021;7(10):804. doi:10.3390/jof7100804

22. Lee YS, Kim J, Lee SG, Oh E, Shin SC, Park IK. Effects of plant essential oils and components from Oriental sweetgum (Liquidambar orientalis) on growth and morphogenesis of three phytopathogenic fungi. *Pesticide Biochemistry and Physiology*. 2009;93(3):138-143. doi:10.1016/j.pestbp.2009.02.002

23. Anand S, Deighton M, Livanos G, Pang ECK, Mantri N. Agastache honey has superior antifungal activity in comparison with important commercial honeys. *Sci Rep*. 2019;9(1):18197. doi:10.1038/s41598-019-54679-w

24. Borotová P, Galovičová L, Vukovic NL, Vukic M, Tvrdá E, Kačániová M. Chemical and Biological Characterization of Melaleuca alternifolia Essential Oil. *Plants*. 2022;11(4):558. doi:10.3390/plants11040558

25. Mathew C, Tesfaye W, Rasmussen P, et al. Mānuka Oil—A Review of Antimicrobial and Other Medicinal Properties. *Pharmaceuticals*. 2020;13(11):343. doi:10.3390/ph13110343

26. Pleeging CCF, Wagener FADTG, De Rooster H, Cremers NAJ. Revolutionizing non-conventional wound healing using honey by simultaneously targeting multiple molecular mechanisms. *Drug Resistance Updates*. 2022;62:100834. doi:10.1016/j.drup.2022.100834

27. Blair SE, Cokcetin NN, Harry EJ, Carter DA. The unusual antibacterial activity of medical-grade Leptospermum honey: antibacterial spectrum, resistance and transcriptome analysis. *Eur J Clin Microbiol Infect Dis*. 2009;28(10):1199-1208. doi:10.1007/s10096-009-0763-z

28. Cooper RA, Jenkins L, Henriques AFM, Duggan RS, Burton NF. Absence of bacterial resistance to medical-grade manuka honey. *Eur J Clin Microbiol Infect Dis*. 2010;29(10):1237-1241. doi:10.1007/s10096-010-0992-1

29. Jandourek A, Vaishampayan JK, Vazquez JA. Efficacy of melaleuca oral solution for the treatment of fluconazole refractory oral candidiasis in AIDS patients. *AIDS*. 1998;12(9):1033-1037.

30. Caelli M, Porteous J, Carson CF, Heller R, Riley TV. Tea tree oil as an alternative topical decolonization agent for methicillin-resistant Staphylococcus aureus. *J Hosp Infect*. 2000;46(3):236-237. doi:10.1053/jhin.2000.0830

31. Seifinadergoli Z, Nahidi F, Safaiyan A, Javadzadeh Y, Eteraf-Oskouei T. Comparison of the efficacy of honey gel and clotrimazole cream in the treatment of vaginal candidiasis symptoms: a randomized clinical trial. *Electron Physician*. 2018;10(6):6904-6911. doi:10.19082/6904

32. Yang C, Gong G, Jin E, et al. Topical application of honey in the management of chemo/radiotherapy-induced oral mucositis: A systematic review and network meta-analysis. *Int J Nurs Stud*. 2019;89:80-87. doi:10.1016/j.ijnurstu.2018.08.007

33. McLoone P, Tabys D, Fyfe L. Honey Combination Therapies for Skin and Wound Infections: A Systematic Review of the Literature. *Clin Cosmet Investig Dermatol*. 2020;13:875-888. doi:10.2147/CCID.S282143

34. Poovelikunnel TT, Gethin G, Solanki D, McFadden E, Codd M, Humphreys H. Randomized controlled trial of honey versus mupirocin to decolonize patients with nasal colonization of meticillin-resistant Staphylococcus aureus. *J Hosp Infect*. 2018;98(2):141-148. doi:10.1016/j.jhin.2017.10.016

35. Kathuria S, Singh PK, Sharma C, et al. Multidrug-Resistant Candida auris Misidentified as Candida haemulonii: Characterization by Matrix-Assisted Laser Desorption Ionization-Time of Flight Mass Spectrometry and DNA Sequencing and Its Antifungal Susceptibility Profile Variability by Vitek 2, CLSI Broth Microdilution, and Etest Method. *J Clin Microbiol*. 2015;53(6):1823-1830. doi:10.1128/JCM.00367-15

36. CLSI. Reference Method for Broth Dilution Antifungal Susceptibility Testing of Yeasts; Approved Standard. 4th ed. M27. Wayne (PA): *Clinical and Laboratory Standards Institute*.

37. CLSI. Performance Standards for Antifungal Susceptibility Testing of Yeasts. 2nd Ed. M60. Wayne (PA): *Clinical and Laboratory Standards Institute*.

38. Thermo Fisher ScientificTM SENSITITRE® YEASTONE® For in Vitro Diagnostic Use. MCS Diagnostics. 2012. [(accessed on 14 April 2023)]; Available online: https://docplayer.net/21149741-Sensititre-yeastone-for-in-vitro-diagnostic-use.html

39. Centers for Disease Control and Prevention, Antifungal Susceptibility Testing and Interpretation. [(accessed on 14 April 2023]; Available online: https://www.cdc.gov/fungal/candida-auris/c-auris-antifungal.html

40. Chan W, Chow FWN, Tsang CC, et al. Induction of amphotericin B resistance in susceptible *Candida auris* by extracellular vesicles. *Emerging Microbes & Infections*. 2022;11(1):1900-1909. doi:10.1080/22221751.2022.2098058

41. Alserehi H, Filippell M, Emerick M, et al. Chlorhexidine gluconate bathing practices and skin concentrations in intensive care unit patients. *American Journal of Infection Control*. 2018;46(2):226-228. doi:10.1016/j.ajic.2017.08.022

42. Yu D, Wang J, Shao X, Xu F, Wang H. Antifungal modes of action of tea tree oil and its two characteristic components against Botrytis cinerea. *J Appl Microbiol*. 2015;119(5):1253-1262. doi:10.1111/jam.12939

43. Li WR, Li HL, Shi QS, et al. The dynamics and mechanism of the antimicrobial activity of tea tree oil against bacteria and fungi. *Appl Microbiol Biotechnol*. 2016;100(20):8865-8875. doi:10.1007/s00253-016-7692-4

44. Sahu B, R. A, Mohammed Abu Javid. Application of Calophyllum inophyllum oil as antifungal fat-liquor for leather industry. *Industrial Crops and Products*. 2017;105:104-112. doi:10.1016/j.indcrop.2017.04.064

45. Kim E, Park IK. Fumigant antifungal activity of Myrtaceae essential oils and constituents from Leptospermum petersonii against three Aspergillus species. *Molecules*. 2012;17(9):10459-10469. doi:10.3390/molecules170910459

46. Parker RA, Gabriel KT, Graham KD, Butts BK, Cornelison CT. Antifungal Activity of Select Essential Oils against Candida auris and Their Interactions with Antifungal Drugs. *Pathogens*. 2022;11(8):821. doi:10.3390/pathogens11080821

47. Campo Research PTE Ltd. 2015. Manuka Oil Extract Leptospermum Scoparium. Campo Research PTE Ltd.; Singapore.

48. Satchell AC, Saurajen A, Bell C, Barnetson RS. Treatment of dandruff with 5% tea tree oil shampoo. *J Am Acad Dermatol*. 2002;47(6):852-855. doi:10.1067/mjd.2002.122734

49. Herman A, Herman AP. Essential oils and their constituents as skin penetration enhancer for transdermal drug delivery: a review. *Journal of Pharmacy and Pharmacology*. 2015;67(4):473-485. doi:10.1111/jphp.12334

50. Fernandes L, Oliveira A, Henriques M, Rodrigues ME. Honey as a Strategy to Fight Candida tropicalis in Mixed-Biofilms with Pseudomonas aeruginosa. *Antibiotics (Basel)*. 2020;9(2):43. doi:10.3390/antibiotics9020043

51. Mandal MD, Mandal S. Honey: its medicinal property and antibacterial activity. *Asian Pacific Journal of Tropical Biomedicine*. 2011;1(2):154-160. doi:10.1016/S2221-1691(11)60016-6

52. Scepankova H, Combarros-Fuertes P, Fresno JM, et al. Role of Honey in Advanced Wound Care. *Molecules*. 2021;26(16):4784. doi:10.3390/molecules26164784

53. Sekizuka T, Iguchi S, Umeyama T, et al. Clade II Candida auris possess genomic structural variations related to an ancestral strain. Chowdhary A, ed. *PLoS ONE*. 2019;14(10):e0223433. doi:10.1371/journal.pone.0223433

54. Brandt P, Mirhakkak MH, Wagner L, et al. High-Throughput Profiling of Candida auris Isolates Reveals Clade-Specific Metabolic Differences. *Microbiol Spectr*. 2023;11(3):e0049823. doi: 10.1128/spectrum.00498-23

55. de Groot T, Janssen T, Faro D, Cremers NAJ, Chowdhary A, Meis JF. Antifungal Activity of a Medical-Grade Honey Formulation against Candida auris. *J Fungi (Basel)*. 2021;7(1):50. doi:10.3390/jof7010050

56. Sutton SV, Proud DW, Rachui S, Brannan DK. Validation of microbial recovery from disinfectants. PDA J Pharm Sci Technol. 2002 Sep-Oct;56(5):255-66.

57. Kwakman PHS, Te Velde AA, De Boer L, Vandenbroucke-Grauls CMJE, Zaat SAJ. Two Major Medicinal Honeys Have Different Mechanisms of Bactericidal Activity. Cardona PJ, ed. *PLoS ONE*. 2011;6(3):e17709. doi:10.1371/journal.pone.0017709

Table 1: In vitro susceptibility of *Candida* isolates to synthetic antiseptics, natural antiseptics and their commercial products

	Clinical *Candida auris* isolates (N=40) (clade I/South Asia)				*Candida auris* JCM 15448 (clade II/East Asia)		*Candida auris* ATCC MYA 5002 (clade III/South Africa)		*Candida auris* ATCC MYA 5003 (clade IV/South America)		*Candida krusei* ATCC 6258		*Candida parapsilosis* ATCC 22019		*Candida albicans* ATCC 90028		*Candida glabrata* ATCC MYA-2950	
	MIC50	MIC90	MIC range	MFC range	MIC	MFC	MIC	MFC	MIC	MFC	MIC	MFC	MIC	MFC	MIC	MFC	MIC	MFC
Synthetic antiseptics																		
Chlorhexidine (%)	0.001	0.001	0.0005-0.001	0.002-0.004	0.001	0.002	0.001	0.002	0.001	0.002	0.0002	0.0005	0.00006	0.0002	0.0005	0.001	0.0005	0.0005
Povidone-iodine (%)	0.078	0.078	0.039-0.078	0.078-0.313	0.039	0.16	0.078	0.156	0.078	0.156	0.039	0.31	0.078	0.625	0.16	0.31	0.16	0.31
Nystatin (µg/mL)	2.44	2.44	1.22-2.44	2.44-9.77	1.22	1.22	1.22	2.44	1.22	2.44	1.22	2.44	0.61	2.44	1.22	4.88	1.22	2.44
Natural antiseptics																		
Manuka honey 15+ (Comvita) (v/v) (%)	25	25	6.25-25	>50	0.20	>50	25	>50	25	>50	25	>50	25	>50	25	>50	1.56	>50
Manuka honey 20+ (Comvita) (v/v) (%)	25	25	3.13-25	>50	0.20	>50	25	>50	25	>50	12.5	>50	25	>50	25	>50	1.56	>50
Manuka honey 15+ (Kiwi Manuka) (v/v) (%)	25	25	3.13-25	>50	0.20	>50	25	>50	25	>50	12.5	>50	25	>50	25	>50	1.56	>50
Manuka honey 20+ (Kiwi Manuka) (v/v) (%)	25	25	6.25-25	>50	0.20	>50	25	>50	25	>50	12.5	>50	25	>50	25	>50	1.56	>50
Tea tree oil 15% (Body Shop) (v/v) (%)	0.004	0.004	0.004	2.08-4.17	0.004	1.04	0.001	0.31	0.001	0.31	0.002	2.08	0.004	4.17	0.039	0.63	0.16	0.63
Tea tree oil 100% (Kiwi Manuka) (v/v) (%)	0.31	0.31	0.31	0.63-1.25	0.31	0.63	0.31	1.25	0.31	1.25	0.63	1.25	0.63	1.25	0.63	1.25	0.63	1.25
Manuka oil 100% (Manuka Lab) (v/v) (%)	0.31	0.31	0.16-0.31	0.63	0.08	0.63	0.31	0.63	0.31	0.63	0.31	1.25	0.31	1.25	0.63	2.5	0.08	0.63
Manuka oil 100% (Melora) (v/v) (%)	0.31	0.31	0.31	0.63	0.04	0.31	0.31	1.25	0.31	1.25	0.31	1.25	0.31	1.25	1.25	5.0	0.08	0.63
Manuka hydrosol (v/v) (%)	>50	>50	>50	>50	>50	>50	>50	>50	>50	>50	>50	>50	>50	>50	>50	>50	>50	>50
Commercial products																		
Manuka body wash (w/v) (%)	0.012	0.012	0.006-0.012	0.05-0.10	0.006	0.024	0.024	0.39	0.012	0.098	0.006	0.05	0.006	3.13	0.024	0.098	0.049	1.56
Tea tree body wash (w/v) (%)	0.012	0.012	0.006-0.39	≥50	0.003	25	0.006	50	0.006	50	0.002	25	0.003	50	0.049	25	0.20	>50
MEDIHONEY wound gel (w/v) (%)	3.13	3.13	1.56-6.25	≥25	0.20	12.5	3.13	>37.5	6.25	>37.5	0.39	25	0.20	>37.5	6.25	>37.5	0.098	37.5
Manuka mouthwash (v/v) (%)	0.39	0.39	0.20-0.39	3.13-6.25	0.39	6.25	0.78	3.13	1.56	3.13	0.10	6.25	0.20	6.25	1.56	6.25	1.56	6.25

Figure 1: Concentration-dependent antifungal effect of natural antiseptics on *Candida auris*. *Candida auris* isolates (n = 3) were selected with MICs representative for MIC_{90} of the 40 isolates. The test products were diluted using RPMI to a test concentration of 25%, 37.5%, and 50% for manuka honey (v/v), tea tree body wash (w/v), and Medihoney[TM] wound gel (w/v) (50% wound gel is not available for testing due to its insolubility at high concentrations). Log reductions in CFU compared with growth control (yeast suspension in RPMI) are shown. Significant differences ($p < 0.05$) between concentrations of antiseptics are indicated with an asterisk.

Table 2: Antifungal drug susceptibility of clinical *Candida auris* isolates

	Clinical *C. auris* isolates (N=40)			
	MIC_{50} (mg/L)	MIC_{90} (mg/L)	MIC range (mg/L)	Resistance, n (%)
Amphotericin B	2	4	1-4	36 (90)
Fluconazole	256	>256	64 - >256	40 (100)
Anidulafungin	0.25	0.25	0.12 - 8	1 (2.5)
Micafungin	0.25	0.25	0.06 - 1	0 (0)
Caspofungin	0.25	0.5	0.06 - >8	3 (7.5)
Itraconazole	0.12	0.25	0.06 - >16	NA
Voriconazole	0.5	1	0.12 - >8	NA
Posaconazole	0.03	0.06	0.015 - >8	NA
5-flucytosine	0.12	0.12	<0.06 - >64	NA

CDC's tentative nonsusceptible breakpoints: ≥ 2 mg/L for amphotericin B, ≥ 32 mg/L for fluconazole, ≥ 4 mg/L for echinocandins anidulafungin and micafungin and ≥ 2 mg/L for caspofungin. For the triazoles itraconazole, voriconazole and posaconazole, and 5-flucytosine, no breakpoints are available. NA: Not available.