Caveolin-1 Regulates Neurogenesis and Learning and Memory by Modulation of Mitochondrial Dynamics

Terilyn K. L. Stephen[1*], Deepika Patel[1*], Luis Aponte Cofresi[1], Elvis Quiroz[1], Kofi Owusu-Ansah[1], Yomna Ibrahim[1], Ellis Qualls[1], Jeffery Marshall[1], Wenping Li[2], Pavan Kumar[1], Aashutosh Shetti[1], Jacqueline A Bonds [3], Richard D. Minshall[4,5], Stephanie M. Cologna[2], Orly Lazarov[1+]

[1]Department of Anatomy and Cell Biology, University of Illinois at Chicago, Chicago, IL, USA

[2]Department of Chemistry, University of Illinois at Chicago, IL, USA

[3]Departmet of Anesthesiology, University of California San Diego, CA, USA

[4]Deparment of Pharmacology and Regenerative Medicine, University of Illinois at Chicago, IL, USA

[5]Department of Anesthesiology, University of Illinois at Chicago, IL USA

[*]Co-first authors

[+]Correspondence: olazarov@uic.edu

Keywords

Abstract

Hippocampal neurogenesis plays instrumental roles in learning and memory. However, the mechanisms underlying neurogenesis are not fully understood. Here we show that the expression of Caveolin-1 (Cav-1), the principal component of caveolae, peaks in neural progenitor cells (NPCs) during neurogenesis. Using NestinCreER^{T2};Cav-1$^{fl/fl}$ male mice, and CRISPR-sham (Cav-1 Ctrl) and CRISPR/Cas9-edited (Cav-1 KO) human induced-pluripotent stem cells, we observed that Cav-1 deletion led to reduced stem cell proliferation and enhanced differentiation into neurons. This was manifested by increased neuronal dendritic tree surface area and enhanced mouse performance in contextual discrimination. Proteomic analysis revealed that Cav-1 plays a role in mitochondrial pathways in NPCs. Cav-1 localized to the mitochondria in NPCs and co-immunoprecipitated with mitofusion 2. Mitochondrial morphology was elongated in Cav-1 KO NPCs and the expression of mitofusion 2 was increased in mitochondrial fractions. Restoration of Cav-1 levels rescued elongated mitochondrial morphology and altered neuronal differentiation in Cav-1 KO NPCs. Together, this study identifies Cav-1 as a novel regulator of neurogenesis and -dependent learning and memory.

Short running title: Caveolin-1, Neurogenesis and Mitochondrial dynamics

Significance Statement

The hippocampal dentate gyrus (DG) orchestrates adult hippocampal neurogenesis (AHN). The precise mechanisms governing AHN remain elusive. Caveolin-1 regulates neurogenesis through mitochondrial fission-fusion process, suggesting Caveolin-1 as a novel regulator of AHN and underscoring the impact of AHN on cognition.

Introduction

Adult hippocampus neurogenesis (AHN) is the postnatal continuous formation of new neurons and their functional integration in the granular cell layer of the dentate gyrus (DG) of the hippocampus. Radial glia-like neural stem cells (NSC) and neural progenitor cells (NPCs, together referred to NSPC) that reside in the subgranular zone (SGZ) of the DG, undergo proliferation and differentiation to produce new granule neurons (Toda et al., 2019). New neurons contribute to spatial and contextual learning and memory (Sahay et al., 2011; Niibori et al., 2012; Miller and Sahay, 2019; Masachs et al., 2021). However, AHN declines with age. In mice, marked reductions in NSPC proliferation and neurogenesis are observed as early as 6 months of age (Encinas et al., 2011; Demars et al., 2013; Britton et al., 2022). Impaired AHN occurs in various brain disorders, including Alzheimer's Disease (AD), both in animal models and humans (Demars et al., 2010; Moreno-Jimenez et al., 2019; Tobin et al., 2019; Terreros-Roncal et al., 2021). In human cohorts the number of neuroblasts is inversely correlated with cognitive diagnosis (Tobin et al., 2019). Interesting, strategies that enhance neurogenesis, such as genetic and pharmacologic interventions restore memory deficits in AD mouse models (Hu et al., 2010; Choi et al., 2018; Mishra et al., 2022). While many intrinsic and extrinsic factors modulate AHN (Demars et al., 2011; Gadadhar et al., 2011; Artegiani et al., 2017; Vicidomini et al., 2020; Zhou et al., 2022), the mechanisms underlying the maintenance of NSPC and neurogenesis-dependent memory function are still not fully understood.

Caveolin-1 (Cav-1) is a 21-24 kDa scaffolding protein of the caveolin gene family that is crucial for the formation of caveolae (Williams and Lisanti, 2004). Cav-1 governs numerous signaling pathways (Parton, 2018; Parton et al., 2020). Cav-1 is enriched in endothelial cells (Ikezu et al., 1998; Virgintino et al., 2002) and is essential for blood brain barrier integrity and neurovascular coupling (Knowland et al., 2014; Blochet et al., 2020; Chow et al., 2020; Filchenko et al., 2020).

Previously, we showed that Cav-1 expression is depleted in the hippocampus of a Type II

diabetes mouse model, and restoration of Cav-1 improved hippocampal memory. Cav-1

knockout mice show diverse neurological deficits, including impaired cholinergic function and

hippocampal plasticity (Trushina et al., 2006; Gioiosa et al., 2008; Head et al., 2010).

Overexpressing Cav-1 in hippocampal neurons enhances dendritic growth and arborization,

likely via increased lipid raft formation and enhanced trafficking of synaptic receptors (Head et

al., 2011; Egawa et al., 2018). Few studies exist examining Cav-1 in neurogenic cell types. In

the developing cortex, Cav-1 expression is essential for internalization of cell adhesion proteins

that regulate proper migration and dendritic pruning in immature neurons (Shikanai et al., 2018).

In early stages of neuronal differentiation from NPCs derived from human inducible pluripotent

stem cells (iPSCs), Cav-1 phosphorylation at its tyrosine 14 site is needed for axonal growth

(Wang et al., 2019). However, a role for Cav-1 in AHN has never been described. Here, we

show that Cav-1 is expressed in both hippocampal NSPC isolated from mice and human NPCs

and neurons derived from iPSCs. Cell-specific deletion of Cav-1 in AHN in mice, resulted in

significant reductions in NSC proliferation, increased neuronal differentiation and enhanced

performance of mice in the AHN-dependent contextual discrimination. Similarly, Cav-1 deletion

from iPSCs induced NPCs resulted in stalled proliferation and enhanced mature neuron

formation. Proteomic analysis revealed that Cav-1 regulates AHN via mitochondria-related

protein pathways. Prior reports linked mitochondrial fusion to neuronal differentiation (Khacho

and Slack, 2018; Kochan et al., 2024). However, the molecular signals that regulate this

process is largely unknown. Importantly, we observed that Cav-1 localizes to the mitochondria

in iPSCs-derived NPCs and regulates fission fusion via interaction with mitofusion-2. Reinstating

Cav-1 in Cav-1 KO iPSCs-derived NPCs restored structural alterations in mitochondria. This

study determines that Cav-1 is a novel regulator of AHN in the mouse and human brain and

governs neuronal differentiation.

Results

Cav-1 deletion in NSPCs shift neurogenesis towards neuronal differentiation in mice

Nestin expression is high in actively proliferating NSPC and downregulated as cells exit the cell cycle during differentiation (Frederiksen and McKay, 1988; Wilhelmsson et al., 2019). To investigate the role of Cav-1 in AHN, we generated a NSPC specific inducible Cav-1 knockout model, where we bred a Cav-1 floxed mice (Cav-1$^{fl/fl}$) (Cao et al., 2003; Oliveira et al., 2017) with a tamoxifen inducible Cre recombinase driven by a Nestin promoter (NestinCre$^{ERT2/+}$) (Mishra et al., 2022). NestinCre$^{ERT2/+}$;Cav-1$^{fl/fl}$ mice were injected at 4-5 weeks of age with either tamoxifen to conditionally delete Cav-1 from NSPCs (iNSC Cav-1 KO) or with corn oil to generate control mice (iNSC Cav-1 WT; **Figure S1 A)**. We confirmed Cav-1 recombination in hippocampal NSPC isolated from iNSC Cav-1 KO and iNSC Cav-1 WT mice by quantitative real-time PCR (qPCR), immunoblotting and immunostaining **(Figure S1 B-D).** Ultrastructural transmission electron microscopy was used to determine if Cav-1 expression in NSPC correlated with the presence of caveolae. Flask-shaped invaginations and vesicles resembling caveolae were identified in the iNSC Cav-1 WT NSPC, whereas larger (>150 nm) electron dense vesicles resembling clathrin-coated vesicles were predominantly found in the iNSC Cav-1 KO NSPC **(Figure S1 E).** Together, this indicates that Cav-1 is expressed in adult hippocampal NSPCs, associated with the presence of caveolae, and that tamoxifen-induced recombination leads to a complete deletion of Cav-1 in iNSC Cav-1 KO mice.

To assess the impact of Cav-1 deletion on neurogenesis, we analyzed NSC proliferation in the DG of iNSC Cav-1 WT and KO mice. At both 3 and 6 months of age, there were no significant differences in the total NSC population (GFAP$^+$Nestin$^+$) or in the quiescent NSC subpopulation (GFAP$^+$Nestin$^+$MCM2-) between genotypes **(Figure 1 A-C and 1 E-G)**. However, the number of proliferating NSCs (GFAP$^+$Nestin$^+$MCM2$^+$) was significantly reduced in Cav-1 KO mice at both ages **(Figure 1 D and 1 H)**. In line with this, expression of ASCL-1 and TBR-2-proxies of proliferating NSCs and intermediate progenitor cells (IPCs), respectively-was also

significantly reduced in Cav-1 KO mice, indicating impaired expansion of these cell populations

at both ages (**Figure S1 F-M).** Supporting these in vivo findings, neurosphere assays revealed

that Cav-1 KO NSPCs had reduced clone formation, smaller clone size, fewer total cells **(Figure**

1 I-L) and reduced incorporation of EDU compared to WT controls **(Figure S1 N-O).**

We next utilized a 12 day 5-bromo-2'-deoxyuridine (BrdU) pulse labeling paradigm

followed by analysis 24 hr after the final BrdU injection to assess NSPC proliferation and lineage

progression (**Figure 1 M**). A trending reduction (p = 0.05) in the total level of BrdU positive cells

in the DG was observed in the iNSC Cav-1 KO mice compared to the iNSC Cav-1 WT mice

(Figure 1 N-O). Consistent with our proliferation analysis, BrdU$^+$ GFAP$^+$ Sox2$^+$ NSCs were

significantly reduced in iNSC Cav-1 KO mice **(Figure 1 P-Q),** and a trend toward fewer

BrdU$^+$GFAP$^-$Sox2$^+$ NPCs was also detected **(p = 0.07; Figure 1 R)**. More notably, there was a

significant shift in the lineage of BrdU$^+$ cells away from NSPCs pool in the iNSC Cav-1 KO mice,

suggesting that Cav-1 loss alters early neurogenic fate decisions **(Figure 1 S)**.

Given that stalled proliferation of NSC can influence early stages of differentiation, we

quantified triple-labeled BrdU$^+$DCX$^+$NeuN$^-$ neuroblasts and BrdU$^+$DCX$^+$NeuN$^+$ immature

neurons **(Figure 2 A-D)**. While no significant differences were observed in BrdU$^+$ neuroblasts

(Figure 2 B), the number of BrdU$^+$ immature neurons were significantly elevated in Cav-1 KO

mice compared with WT **(Figure 2 C-D).** Consistent with these findings, at 6 months of age the

iNSC Cav-1 KO mice exhibited a significant increase in total number of DCX$^+$ NPCs,

neuroblasts and immature neurons (DCX$^+$NeuN$^+$) compared to iNSC Cav-1 WT mice (**Figure 2**

E-H, **Figure S1 P-S).** These findings support the hypothesis that Cav-1 loss promotes neuronal

lineage commitment.

The effect of Cav-1 deletion on differentiation was also examined in hippocampal

neurosphere cultures isolated from iNSC Cav-1 WT and iNSC Cav-1 KO mice (**Figure 2 I**)

(Bonds et al., 2020; Gadadhar et al., 2011). During in vitro differentiation over 5–7 days, Cav-1

transcript and protein levels were significantly downregulated in WT NSPCs **(Figure 2 J, M-N).**

Differentiation progression was confirmed by decreased expression of nestin and Sox2 and

concurrent upregulation of the neuronal markers Map2 and βIII-tubulin **(Figure 2 K-L, O-P).**

Consistent with in vivo findings, Cav-1 KO NSPCs exhibited significantly higher expression of

Map2 and βIII-tubulin compared with WT controls during differentiation **(Figure 2 L, P).**

Together these findings suggest that Cav1 negatively regulates NSPC differentiation. Its loss

facilitates the transition from proliferation to neuronal commitment, enhancing the maturation of

new neurons.

Cav-1 deletion in NSPC increases newborn neuron dendritic tree volume and improves

context discrimination learning and memory

Dendritic arborization of immature neurons is a critical component of neuronal

maturation, influencing the ability of these adult-born neurons to integrate into existing circuits

and participate in hippocampal-dependent functions such as learning and memory (Toni et al.,

2007; Zhao et al., 2008). To investigate whether Cav-1 deletion alters dendritic structure in

newly generated neurons, we infected dentate gyrus neurons in iNSC Cav-1 WT and iNSC Cav-

1 KO mice with retrovirus-green fluorescence protein (RV-GFP) and sacrificed them 30 days

later to allow the visualization of the dendritic structure of newly post-mitotic neurons. Dendrite

morphology was quantified by Sholl analysis **(Figure□3A).** We observed no significant

differences in the number of dendritic branch intersections and total dendritic length **(Figure 3**

B-D). Yet, both surface area and more significantly, dendritic volume were increased in the RV-

GFP labeled newly born neurons in the iNSC Cav-1 KO mice compared to iNSC Cav-1 WT

(Figure 3 E-F). These findings suggest that Cav-1-deficient newborn neurons, despite

unchanged dendritic branching patterns, develop thicker dendrites with potentially greater

membrane and cytoplasmic volume. These features are often associated with enhanced

structural stability and increased potential for synaptic integration (van Elburg and van Ooyen,

2010).

Given that newborn neurons (4-6 weeks post-mitotic) become selectively activated

during memory formation and are essential for contextual discrimination, the ability to

discriminate between similar contexts (Clelland et al., 2009; Sahay et al., 2011; Nakashiba et

al., 2012), we next asked whether the increase in immature neurons in the iNSC Cav-1 KO mice

results in improved neurogenesis-dependent learning and memory. Thus, 6 month old iNSC

Cav-1 KO and iNSC Cav-1 WT mice underwent a contextual discrimination test (Clemenson et

al., 2014) **(Figure 3 G).** Both iNSC Cav-1 KO and iNSC Cav-1 WT mice displayed higher

freezing levels in Context A (Cxt A) compared to Context B (Cxt B) at 30 mins post-shock, albeit

with no difference in the discrimination index **(Figure 3 H-I).** Interestingly 24 hours post-shock

on Day 3, the iNSC Cav-1 KO mice had a significantly higher level of freezing in Cxt A

compared to Cxt B whereas the iNSC Cav-1 WT mice exhibited the equal amount of freezing in

both Cxt A and Cxt B **(Figure 3 J)**. A trending increase ($p = 0.06$) in the discrimination index

was observed in the iNSC Cav-1 KO mice (**Figure 3 K**), suggesting enhanced neurogenesis-

dependent memory.

We then examined whether the iNSC Cav-1 KO mice displayed enhanced context

generalization compared to the iNSC Cav-1 WT mice 24 hours post-shock. Studies show that

significant alterations in context geometry are sufficient to induce generalization rather than

discrimination behavior (Gonzalez et al., 2003; Huckleberry et al., 2016) and that newborn

neurons maintain generalization behavior (Guo et al., 2018). To evaluate context generalization,

a similar paradigm was used with a Context C (Cxt C), which is modified in geometry and tactile

cues **(Figure S2 A)**. Notably on Day 2 (30 mins post shock), iNSC Cav-1 WT mice were unable

to discriminate Cxt A from Cxt C whereas iNSC Cav-1 KO mice had a significantly higher level

of freezing in Cxt A compared to Cxt C with a trending significant discrimination index **(Figure**

S2 B-C). Notably, 24 hours post shock, both groups of mice were able to equally discriminate

Cxt A from Cxt C 24 hr post-shock **(Figure S2 D-E).** No differences in level of anxiety-like

behavior **(Figure S2 F-I)** nor performance in the spatial novel object location (NOL) task (**Figure**

SS J-L) were observed between the iNSC Cav-1 KO and iNSC Cav-1 WT mice. Together these

findings indicate that Cav-1 deletion in NSPCs promotes the generation of immature neurons

with enhanced dendritic volume, leading to improved neurogenesis-dependent context

discrimination without affecting general anxiety or spatial memory.

Proteomic analysis reveals that Cav-1 regulates mitochondrial signaling pathways and

morphology in hippocampal NSPCs

Building on our findings that Cav-1 deletion in NSPCs enhances hippocampal

neurogenesis, we hypothesized that Cav-1 regulates pathways involved cell cycle cessation and

differentiation. Using quantitative proteomics, we assessed the molecular mechanisms by which

Cav-1 regulates adult hippocampal neurogenesis in mice **(Figure 4 A).** For that, we isolated

hippocampal NSPCs from the dentate gyrus of iNSC Cav-1 KO and iNSC Cav-1 WT mice and

examined their protein profile by mass spectrometry. We identified 4,730 proteins, with 326

differentially expressed (DEPs; ANOVA, p < 0.05, **Figure 4 B Table S1**) — 228 downregulated

and 98 upregulated in Cav-1 KO NSPCs. Ingenuity Pathway Analysis (IPA) and Gene Ontology

(GO) revealed that the top altered pathways included Mitochondrial Dysfunction, TCA Cycle II,

Cholesterol Biosynthesis, and Oxidative Phosphorylation (**Figure 4 C-D, Table S2**). GO

analysis also pointed to changes in ATP biosynthesis, the electron transport chain (ETC),

mitochondria and ribosome subunits, and lipid oxidation.

To explore Cav-1's role in mitochondrial regulation in NSPCs, we compared the DEPs to

the MitoCarta3.0 gene dataset (Rath et al., 2021), which catalogs 1,140 mouse genes localized

to mitochondria. Of the 326 DEPs identified in iNSC Cav-1 KO vs. WT NSPCs, 84 (24%)

matched MitoCarta3.0 genes (**Figure 4 E**). Downregulated proteins in Cav-1 KO NSPCs were

enriched in pathways related to mtDNA maintenance, mtRNA processing and translation,

protein import, sorting, and mitochondrial homeostasis. Several metabolic pathways including

glycolysis, TCA, OXPHOS, lipid, nucleotide, amino acid, vitamin, and metal metabolism were

also downregulated (**Figure 4 E**). Noticeably, we found that mitochondrial dynamics and

surveillance associated proteins including Metaxin-2 (MTX2) and Voltage-dependent Anion-

selective Channel (Vdac1) were downregulated, whereas Mitochondrial Rho GTPase 2 (Rhot2,

also called Miro2) was significantly enriched in the iNSC Cav-1 KO NSPCs compared to iNSC

Cav-1 WT NSPCs (**Figure 4 E)**. Miro2 interacts with mitofusion-2 (Mfn-2) to coordinate

fission/fusion events (Saotome et al., 2008) and in neurons, Miro2 is required for retrograde

trafficking of mitochondria (Misko et al., 2010; Ruggiero et al., 2017) suggesting that part of

Cav-1's function may involve regulating mitochondrial positioning and dynamics in NPCs.

In hippocampal neurogenesis, changes in mitochondrial morphology and dynamics have

been found to govern self-renewal and cell fate (Khacho et al., 2016). We first examined the co-

localization of Cav-1 to mitochondria. Caveolae and caveolin-1 were previously shown to

localize to the outer membrane of mitochondria (Patel and Insel, 2009; Fridolfsson et al., 2012).

Cav-1 co-localizes with Mitochondrial import receptor subunit TOM20 homolog (Tom-20), a

translocase located on the mitochondrial outer membrane mitochondria, in WT hippocampal

NSPCs by utilizing 3D confocal microcopy (**Figure 4 F**). Next, we examined whether Cav-1

deletion in hippocampal NSPCs alters mitochondrial morphology by Ultrasound EM and live cell

imaging analysis. Ultrastructural EM analysis revealed that Cav-1 KO NSPCs had fewer

mitochondria per cell, yet significantly increased mitochondrial area and perimeter, with reduced

circularity indicating a shift toward elongated morphology **(Figure 4 G-K).** Live cell imaging

using TMRM confirmed these findings: Cav-1 KO mitochondria were larger (**Figure S3 A-C**),

more elliptical (**Figure S3 D**), and more branched (**Figure S3 E**), with elevated TMRM

fluorescence (**Figure S3 F**) indicating a hyperpolarized membrane potential. Time-lapse

imaging of mitochondria was analyzed using the Trackmate plugin in ImageJ (Ershov et al.,

2022) (**Figure S3 G**). Mitochondria in the iNSC Cav-1 KO NSPC had a significant increase in

velocity and total distance traveled compared to iNSC Cav-1 WT NSPCs **(Figure S3 H-I)**.

Together, these data suggest that Cav-1 regulates mitochondrial structure and dynamics in

hippocampal NSPCs. Its deletion promotes elongation, branching, and mobility, features of

fusion and fission dynamics, potentially driving neuronal differentiation (Khacho and Slack,

2018).

Cav-1 deletion enhances differentiation of human iPSCs-derived NSCs

To explore whether Cav-1 plays similar role in human neurogenesis, we next utilized

iPSCs model. Using this approach, we examined the effects of Cav-1 deletion on neuronal

lineage progression and maturation across defined stages of neural development via generation

of a CRISPR/Cas9-edited Cav-1 knockout iPSCs (Cav-1 KO). Cav-1 Ctrl and Cav-1 KO iPSCs

were differentiated into forebrain neurons using neural induction method **(Figure 5 A).** This

method defines 3 neuronal differentiation stages, with neural progenitors (NPCs) having the

most stemness, precursors cells resembling early stages of neuron lineage (similar to

neuroblasts) and lastly neurons.

Cav-1 displayed a cell-type-specific, biphasic expression pattern during neuronal

differentiation, with peak levels in NPCs and mature neurons (**Figure 5 B–C**). Cav-1 expression

was low in iPSCs, increased in NPCs, and significantly decreased in precursor cells. Following

differentiation, Cav-1 expression rose ~6-fold in 10-week-old neurons compared to precursors.

Cav-1 levels were also significantly higher in mature neurons than in iPSCs, NPCs, or

precursors, suggesting a cell-autonomous role throughout differentiation. As expected, Cav-1

was expressed at all stages in control cells but was absent in Cav-1 KO cells, validating the

model (**Figure 5 D–E**). Neuronal differentiation progression was further assessed by quantifying

βIII-Tubulin expression (**Figure 5 D, F**).

Next, we asked whether Cav-1 similarly regulates neurogenesis in human NPCs, as

observed in primary hippocampal NSPCs from Cav-1 WT and KO mice. DCX expression,

assessed by immunocytochemistry in human NPCs, precursor cells, and 5-week-old neurons,

was significantly higher in Cav-1 KO precursor cells than in WT controls (**Figure 5 G, I**). Cav-1

KO precursor cells and neurons showed significantly elevated β-III-tubulin expression,

confirming enhanced neuronal differentiation relative to controls (**Figure 5 H, J**). Supporting our

mouse data, Cav-1 KO iPSC-derived NPCs had significantly fewer EdU$^+$ cells compared to WT,

indicating reduced proliferation (**Figure 5 K-L**). Taken together, findings from iPSCs derived

human NPCs and in vivo quantification of NSPC in mice confirmed that cell autonomous loss of

Cav-1 halts cell proliferation while enhancing neuron differentiation in AHN.

Rescue of Cav-1 expression in iPSCs-derived NPCs lacking Cav-1 restores mitochondrial

morphology.

To further assess the mechanism of Cav-1 as a direct regulator of mitochondria

morphology, we used a lentivirus expressing Cav-1 and GFP (LV-Cav-1-p2A-GFP; LV Cav-1) or

a GFP control (LV-GFP) to restore Cav-1 protein expression in our iPSCs model (**Figure 6 A**).

Consistent with our proteomic and mouse NPCs model, we observed Cav-1 co-localization with

the mitochondrial outer membrane marker TOM20 in iPSCs derived NPCs Cav-1 Ctrl and

restoration of Cav-1 expression in Cav-1 KO NPCs using confocal microscopy (**Figure 6 B**).

Next we performed live imaging using TMRM to examine rescue of mitochondria morphology

(**Figure 6 C**). Consistent with our mouse NPC model, iPSCs-derived Cav-1 KO NPCs displayed

significantly increased mitochondrial area, perimeter, and branch length, along with reduced

circularity indicating a more elongated morphology compared to WT NPCs (**Figure 6 D–G**).

Importantly, restoration of Cav-1 expression in the Cav-1 KO NPCs significantly reduced

mitochondrial area, perimeter, and branch length, while increasing circularity compared to LV-

GFP controls, confirming that Cav-1 regulates mitochondrial morphology in NPCs.

To further investigate the mechanism by which Cav-1 regulates mitochondrial

morphology, we assessed its interaction with key mitochondrial dynamics regulators: mitofusin 2

(Mfn2) and dynamin-related protein 1 (Drp1). Mfn2 promotes mitochondrial fusion, enabling the

elongation and interconnection of mitochondrial networks, while Drp1 mediates mitochondrial

fission, leading to mitochondrial fragmentation and distribution. Co-immunoprecipitation

confirmed that both Mfn2 and Drp1 directly interact with Cav-1 (**Figure 6 H**). Next, alterations in

the mitochondrial localization of Drp1 and Mfn2 in the absence of Cav-1 were evaluated through

subcellular fractionation of Cav-1 WT and Cav-1 KO NPCs (**Figure 6 I-M**). Western blot analysis

revealed a significant increase in Mfn2 levels in the mitochondrial fraction with no change in

Drp-1 levels as result of Cav-1 loss (**Figure 6 K).** Interestingly, a significant increase in

membrane Drp-1 levels were found in the Cav-1 KO NPCs compared to control (**Figure 6 L).**

No change in total expression levels or cytosol expression levels of Mfn2 or Drp1 were found in

Cav-1 KO NPCs compared to controls (**Figure 6 J, M**). Together, our findings suggest that Cav-

1 regulates mitochondrial morphology by modulating the subcellular localization of fusion-

fission-linked proteins and therefore, Cav-1 loss promotes neuronal lineage commitment by

enhancing mitochondrial fusion in NPCs.

Restoring Cav-1 levels reverses neuronal differentiation phenotype.

To examine whether reinstating Cav-1 in Cav-1KO NPCs would reverse enhanced

neurogenesis, Cav-1 in Cav-1 KO NPCs were infected with LV-GFP (control) or LV-Cav-1.

Examination of Cav-1 expression in protein lysate of these cells across neural differentiation

stages validated successful transduction procedure **(Figure 7A, B)**. Notably DCX protein levels

is significantly reduced in precursors with Cav-1 overexpression, indicating normalization of

enhanced neuronal lineage commitment at this stage **(Figure 7C)**. Strong trends toward

decreased β-III-Tubulin, were observed in Cav-1-overexpressing lines, though these differences

do not reach statistical significance at precursor and neurons stage **(Figure 7D)**. Together with

Figure 6 data, these results suggest that Cav-1 regulates neurogenesis by modulating

mitochondrial remodeling **(Figure 7E).**

Discussion

This study revealed several key insights. First, Cav-1 deletion in mouse NSPCs reduced

NSPC proliferation and increased the number of immature neurons in the DG, suggesting that

Cav-1 acts as a negative regulator of neuronal differentiation. This shift in cell fate was

associated with enhanced context discrimination memory and greater dendritic volume in

newborn neurons, indicating functional relevance. Next, to extend these findings to human

neurogenesis, we used iPSCs-derived NPCs and found that, similarly to the mouse, Cav-1

deletion impaired proliferation and promoted neuronal differentiation, supporting a conserved

role for Cav-1 in regulating neurogenesis. Finally, we identified Cav-1 as a novel regulator of

mitochondrial dynamics in neurogenesis. Cav-1 loss in human NPCs led to elongated

mitochondria, consistent with increased fusion, which was reversed by Cav-1 restoration.

Mechanistically, Cav-1 directly bound Mfn2 and Drp1, key mediators of fusion and fission,

respectively. In Cav-1 KO NPCs, mfn2 was enriched in mitochondrial fractions, while Drp1

remained in membrane fraction, indicating that Cav-1 supports balanced mitochondrial

dynamics by facilitating proper protein localization. Together, these findings establish Cav-1 as

a critical coordinator of neurogenesis and mitochondrial function.

The balance between quiescence, proliferation, and differentiation of neural stem cells

(NSCs) is essential for maintaining hippocampal neurogenesis. While NSCs integrate both

extrinsic and intrinsic signals to determine their fate (Urban et al., 2019), our findings identify

Cav-1 as a key intrinsic regulator that suppresses neuronal differentiation. Although Cav-1 is

enriched in endothelial cells and may exert non-cell autonomous effects, our data show that its

deletion in NSCs leads to a reduction in NSPC proliferation and a marked increase in immature

neurons, suggesting a shift toward neuronal fate commitment. This was accompanied by

enhanced dendritic volume in newborn neurons, a hallmark of functional maturation and

integration into hippocampal circuits.

Dendritic growth is critical for synaptic integration, input specificity, and neuronal

survival. In the dentate gyrus, adult-born granule cells must extend complex dendritic trees to

reach the molecular layer and receive afferent input from the entorhinal cortex. Increased

dendritic volume and complexity have been correlated with enhanced synaptic input, improved

synaptic plasticity, and stronger contributions to hippocampal-dependent learning tasks (Zhao et

al., 2006; Toni et al., 2007). Thus, the observed dendritic phenotype in iNSC Cav-1 KO mice

likely enhances the functional relevance of increased neuronal differentiation by potentially

facilitating synaptic incorporation and circuit maturation.

Recent studies suggest that mitochondrial morphology is tightly linked to stem cell

identity and cell fate, with mitochondrial fusion promoting neuronal differentiation (Khacho et al.,

2016; Khacho and Slack, 2018; Coelho et al., 2022). NSC in the adult DG exhibit fragmented

and globular mitochondria that become more elongated as cells progress into committed NPCs

and immature neurons (Beckervordersandforth et al., 2017). *In vivo* and *in vitro* studies show

that when NSPC differentiate into neurons a metabolic switch occurs from glycolysis to

OXPHOS (Zheng et al., 2016; Beckervordersandforth et al., 2017). Moreover, mitochondrial

fission-fusion dynamics, orchestrated by Drp1 and Mfn2, are increasingly recognized as

essential for newborn neuron plasticity and maturation (Kochan et al., 2024).

Our findings add to the growing body of evidence implicating mitochondrial dynamics in

neural stem cell fate by identifying Cav-1 as a novel regulator of mitochondrial morphology in

neurogenesis. We show that Cav-1 physically interacts with both Drp1 and Mfn2, suggesting it

may serve as a scaffold to coordinate their subcellular localization. Deletion of Cav-1 in both

mouse NSPCs and human NPCs led to elongated mitochondrial morphology, consistent with

enhanced mitochondrial fusion, and increased mitochondrial velocity and displacement in

mouse NSPCs. In human Cav-1 KO NPCs, we observed a significant redistribution of Mfn2 to

the mitochondria and Drp1 to the plasma membrane, without changes in total protein levels,

indicating a spatial rather than transcriptional regulation. Notably, prior studies have shown that

loss of Cav-1 or mutation at its phosphorylation site disrupts its interaction with Mfn2, thereby

permitting Mfn2 translocation to the mitochondrial outer membrane and thus, alter cell migration

patterns (Jiang et al., 2022). Importantly, re-expression of Cav-1 in Cav-1 KO NPCs restored

mitochondrial morphology, further supporting its regulatory role in mitochondrial dynamics. This

shift toward a fused mitochondrial state aligns with our observed increase in neuronal

differentiation in Cav-1-deficient models, suggesting that Cav-1 normally acts to restrain lineage

commitment by maintaining mitochondrial dynamics.

Our previous studies further underscore the relevance of Cav-1 function in disease

states associated with impaired neurogenesis (Bonds et al., 2019; Bonds et al., 2020). We

found that Cav-1 levels were significantly reduced in the hippocampi of patients with type 2

diabetes mellitus (T2DM) and inversely correlated with β-amyloid (Aβ) levels. Similarly, in the

db/db diabetic mouse model, Cav-1 expression was diminished alongside elevated levels of

amyloid precursor protein (APP) and hyperphosphorylated tau—hallmarks of Alzheimer's

disease (AD) pathology (Bonds et al., 2019). Notably, these mice also exhibited impaired

hippocampal neurogenesis (Bonds et al., 2020), consistent with our current findings that link

Cav-1 deficiency to altered proliferation of NSC in the DG. Furthermore, we previously

demonstrated that reduced endothelial Cav-1 expression contributes to insulin signaling deficits

observed in T2DM (Shetti et al., 2023). Given the link between mitochondrial dysfunction and

both T2DM and AD (Carvalho and Moreira, 2023), these findings suggest that Cav-1 may be a

contributing factor in the neurogenesis and cognitive impairments associated with these

disorders.

In conclusion, this study identifies Cav-1 as a novel intrinsic regulator of adult

hippocampal neurogenesis that exerts its effects, in part, by governing mitochondrial dynamics.

Cav-1 directly interacts with Mfn-2 and Drp-1, supporting the recruitment of these proteins to

mitochondria and maintaining balanced fusion and fission. Loss of Cav-1 favors mitochondrial

fusion, accelerates NSPC differentiation, and enhances neurogenesis-dependent cognitive

function. These findings broaden our understanding of the molecular interplay between

membrane scaffolding proteins and mitochondrial dynamics in the regulation of adult

neurogenesis.

Materials and Methods

Mouse Models

All mouse experiments were approved by the University of Illinois at Chicago Institutional

Animal Care and Use Committee. Mice were housed on a 12-hour light/dark cycle and provided

food and water ad libitum. C57Bl/6J wildtype (WT) and global Cav-1 knockout (gCav-1 KO)

used in experiments were from Jackson Laboratories (Strain #:000664 and Strain #: 007083,

respectively). Inducible NSPC specific-Cav-1 knockout mice (NestinCre$^{ERT2/+}$;Cav-1$^{fl/fl}$) were

generated by crossing NestinCre$^{ERT2/+}$ mice (Sahay et al., 2011; Mishra et al., 2022) with Cav-

1$^{fl/fl}$ mice (Cao et al., 2003; Oliveira et al., 2017). The NestinCre$^{ERT2/+}$ transgene was maintained

as hemizygous and the Cav-1$^{fl/fl}$ homozygous. Male mice were used in experiments unless

otherwise stated in text.

Tamoxifen Injections

Tamoxifen (TAM, Sigma-Aldrich) was dissolved in corn oil (Sigma-Aldrich) at 20 mg/mL

at 37°C and then stored at 4°C for 5 days. To induce recombination, 4–5-week-old mice were

intraperitoneally injected at a dose of 130 mg/kg TAM or equal volume of corn oil once a day for

5 consecutive days.

Viral Vectors

The Retrovirus/Moloney Murine Leukemia Virus (MMLV) construct carrying GFP (RV-

GFP) was generously provided by Dr. Jenny Hsieh (University of Texas at San Antonio, TX).

The viral particles were packaged as previously described (Lybrand et al., 2021). The packaging

process was carried out at the Vector Core Facility of the University of Illinois Chicago (UIC)

under the guidance of the Hsieh lab. The viral titer was 1.67 × 10□ TU/ml for this study.

Mouse NSPC Isolation and Culture

Primary hippocampal NSPCs were isolated from 6–8-week-old mice similarly as

described (Bonds et al., 2020; Ahmed et al., 2021). Hippocampi were dissected in ice-cold

HBSS and pooled from 4-6 mice. Hippocampi were transferred to a tissue culture hood, minced

using a sterile scalpel until no visible pieces remained (3-5 mins) and transferred to 3-5 mL of

warm culture media (DMEM/F-12 with 20 mM KCl, 2 µg/mL heparin, 1% penicillin–streptomycin,

20% B27 supplement, 10% N2 supplement). Tissue was spun at 200g for 2 mins and

dissociated with 0.1% Trypsin-EDTA (diluted in DMEM/F12) at 37°C for 10 mins. After

incubation, tissue was triturated 5 times with a P1000 pipette. Next, 3 mL of trypsin-inhibitor

(0.139 mg/mL and 1U/mL DNase I in HBSS-/-) was added, triturated with a P1000 pipette 5

times and centrifuged at 300g for 5 mins. The cell pellet was singly resuspended in 1 mL of

culture media by pipetting an additional 25 times with a P1000 pipette followed by filtration

through a 40 µm cell strainer and filter washed with 15-20 mL of culture media. Cells were

centrifuged 300g for 5 mins and resuspended in 1 mL of culture media plus growth factors (20

ng/mL EGF and 10 ng/mL bFGF) by pipetting 10 times with a P1000 pipette. Cells were plated

in a 24-well plate (1 well per mouse) containing 2 mL of culture media plus growth factors. Half

media changes were preformed every 48 hrs for 7-10 days with growth factors added to culture

media immediately prior to use.

After the first 7-10 days, cells were collected and centrifuged 300g for 5 mins. The cell

pellet was singly dissociated by pipetting 20 times with a P1000 pipette in 1 mL of culture media

containing growth factors. The cell suspension was filtered through a 40 µm cell strainer and

filter washed with 15-20 mL of culture media. Cells were centrifuged 300g for 5 mins,

resuspended in 1 mL of culture media plus growth factors and plated in 2-3 wells of 6-well plate

containing 3 mL of culture media plus growth factors per well. Half media changes were

preformed every 48 hrs until ~100 µm in diameter neurospheres are formed (7-10 days). All

experiments were performed using NSPCs between passage numbers 3 to 8, where

neurospheres were dissociated using Accutase (StemCell Technologies) at 37°C for 7 mins

followed by the addition of 5 mL culture media and centrifuged 300g for 5 mins. Cells were

counted and plated at 10,000 cells/cm^2 for floating neurosphere cultures. For experiments

requiring monolayers, singly dissociated NSPCs were grown on poly-L-orthinine/laminin

(PLO/laminin) coated plastic dishes or acid washed coverslips. Briefly, plastic dishes and

coverslips were incubated at 37°C with 15 µg/mL PLO diluted in 1X PBS for 24-48 hr, washed

with 1X PBS and incubated at 37°C with 10 µg/mL laminin in 1X PBS for 24 hr. PLO/laminin

coated dishes or coverslips were washed 1 time with 1X PBS prior to plating of cells.

Human iPSCs Culture, NPC differentiation and Neuron Maturation

CRISPR-sham- and CRISPR/Cas9-edited induced pluripotent stem cell lines (Cav-1WT

and Cav-1KO iPSCs, n = 4) were generated by Synthego. These iPSCs lines were derived from

adults with no known trauma or genetic dispositions. One vial from the master stocks was

thawed at 37°C and were plated on matrigel-coated plates (MG, Corning) with mTeSR plus

media and Y-27632 (ROCK inhibitor) used at a final concentration of 10µM. The ROCK inhibitor

was added during the initial 24 h medium which was replaced daily. Individual colonies were

mechanically dissected and were passaged weekly. All reagents were purchased from

STEMCELL Technologies unless otherwise specified.

The neural differentiation protocol was followed from STEMCELL Technologies and the

NPCs, precursor cells, 10 weeks old neurons were validated by ICC staining. Refer figure 5A.

for the differentiation timeline. STEMdiff Neural Induction Medium + SMADi was used to

generate CNS-type NPCs from hiPSCs using embryoid body (EB) protocol. STEMdiff Neural

Progenitor Medium was used to culture and passage neural progenitor cells. STEMdiff™

Forebrain Neuron Differentiation Kit was used to generate neuronal precursors from NPCs. The

neuronal precursors generated were matured using STEMdiff™ Forebrain Neuron Maturation

Kit to produce forebrain-type neurons.

Confluent iPSCs were dissociated using Gentle Cell Dissociation Reagent and plated at

3×10^6 cells/well in neural induction medium (NIM) with SMADi of the AggreWell™800 plate,

treated with Anti-Adherence Rinsing Solution. This resulted in 10,000 cells/microwell. Y-27632

was added to STEMdiff™ Neural Induction Medium + SMADi to obtain a final concentration of

10μM for the first 24h only. NIM was partially replaced daily for 5 days, after which EBs were

harvested from a single well of an AggreWell™800 plate and plated onto a tissue culture-treated

6-well plate coated with PLO/laminin for the formation of neural rosettes. After a week, neural

rosettes were selected using STEMdiff™ Neural Rosette Selection Reagent and replated on

PLO/Laminin coated 6-well plate for the generation for NPCs differentiation. Cells were either

expanded for banking or kept for neuronal differentiation. One fifth of cells were retained at each

passage for marker analysis. Cells were expanded up to 4 passages.

NPCs were cultured for a week in neural progenitor medium. After seven days, cells

were dissociated using accutase, and re-plated in Forebrain Neuron Differentiation media for

neural precursor cells differentiation for a week. The neuronal precursors generated were

matured using STEMdiff™ Forebrain Neuron Maturation Kit to produce a mixed population of

excitatory and inhibitory forebrain-type neurons (≥ 90% class III β-tubulin-positive neurons). Cell

culture at all stages was carried out on tissue culture plastic well plates and for imaging, 24 well

glass bottom plates were used.

Cav-1 Lentivirus Transduction

A custom-designed lentiviral vector was purchased from VectorBuilder. The construct

was designed to express Cav-1 and GFP separated by a self-cleaving P2A linker to allow

equimolar co-expression from a single transcript under the control of the Nestin promoter

(Vector ID: VB200519-1185asg). The control lentiviral construct was designed to express GFP

under the control of the Nestin promotor (VB200803-1257vht). The viral titer was >2x10^9 TU/ml

for this study.

NPCs were cultured as described above. Approximately 50,000 NPCs were seeded per

well in 24-well glass bottom plates. Approximately 80% confluent NPCs were transduced using

produced virus particles at multiplicity of infection (MOI) of 10. The medium was changed after 2

h and the cells were analyzed 5 days post-transduction, after TMRM incubation for 30 min,

using a confocal microscope.

Similarly, for the neuronal differentiation rescue experiment, approximately 1.2 x 10^6

NPCs were seeded per well in a 6-well plate. After 24 hours, cells were transduced with

lentivirus (LV) carrying either GFP or Cav-1 for 2 hours, followed by a medium change.

Neuronal differentiation was then carried out as described previously. NPCs were maintained in

neural progenitor medium for one week, differentiated into precursor cells using Forebrain

Neuron Differentiation Medium for another week, and subsequently matured using the

STEMdiff™ Forebrain Neuron Maturation Kit for up to 10 weeks.

BrdU and EdU Labeling

For the 12-day *in vivo* experiments, 5-bromo-2'-deoxyuridine (BrdU, Sigma) was

prepared fresh daily by dissolving BrdU at 20 mg/mL in 1X PBS at 37°C and then sterile filtered

via 0.22µm syringe filter. Mice were intraperitoneally injected at a dose of 100 mg/kg once daily

for 12 consecutive days and tissue collected 24 hr after the last injection. Mice were sacrificed

and tissue collected 4 weeks after the last injection. For *in vitro* BrdU and EdU experiments,

NSPCs were seeded on PLO/laminin coated coverslips at a density of 50,000 cells/cm^2 and

incubated overnight in culture media containing growth factors. A 5 mM EdU stock was

prepared in DMSO and stored at -20°C until use. EdU stock solutions were diluted to 5 µM in

culture media, sterile filtered via 0. 22µm syringe filter and growth factors added immediately

prior to the start of the experiment.

Clonogenic Proliferation Assay

Hippocampal NSPCs were singly dissociated and plated as floating cultures in 96-well

plates at 1000 cells per well. Cells were grown in culture media containing growth factors (20

ng/mL EGF and 10 ng/mL bFGF) and growth factors were added to media every 48 hr. On day

6, cultures were imaged using a Keyence BZ-X800 followed by dissociation with Accutase to

count the number of cells per well. The number and diameter of neurospheres (clones) per well

was quantified using ImageJ.

Differentiation Assay

Hippocampal NSPCs were seeded on PLO/laminin coated plastic dishes or coverslips at

a density of 50,000 cells/cm^2. Cells were grown in culture media containing growth factors (20

ng/mL EGF and 10 ng/mL bFGF) overnight (8-12 hours). Media was then changed to

differentiation media consisting of culture media (DMEM/F-12 with 20 mM KCl, 2 µg/mL heparin,

1% penicillin–streptomycin, 20% B27 supplement, 10% N2 supplement) with 1 µM retinoic acid

(Sigma) and 5 µM forskolin (Sigma). Half of the media was changed every 48-72 hours. Cells

were fixed or collected after 3 or 7 days of differentiation.

Immunohistochemistry and Immunocytochemistry

Mice were transcardially perfused with ice cold PBS followed by 4% PFA in 1X PBS.

Brains were post-fixed for 24 hr in 4% PFA followed by immersion in 10% sucrose in 1X PBS for

24 hr, 20% sucrose in 1X PBS for 24 hr and 30% sucrose 1X PBS for 24 hr. Brains were

sectioned at 40 µm using a sliding microtome and floating coronal sections were stored in

cryoprotectant consisting of glycerol (20% v/v) and ethylene glycol (24% v/v) in 1X PBS at -

20°C. Sections were washed with 1X PBS, incubated with 1% sodium borohydride in 1X PBS

for 10 mins at RT and washed 3 times with 1X PBS for 10 min per wash. For antibodies

requiring heat induced antigen retrieval, sections were incubated in 10 mM sodium citrate buffer

containing 0.05% Tween 20, pH 6.0 at 99°C for 15 mins in a vegetable steamer. For BrdU

immunodetection, sections were washed with 1X PBS, treated in pre-warmed 1N HCl for 40 min

at 37°C followed by incubation with 0.1 M sodium borate buffer pH 8.5 for 10 min at RT.

Following, antigen retrieval or HCl pre-treatment, sections were washed 3 times with 1X PBS for

10 min per wash. Sections were blocked in 1X PBS containing 0.3 M Glycine, 0.2% Triton X-

100 and 5% normal donkey serum (NDS) for 1 hr at RT. After blocking, sections were incubated

with primary antibodies diluted in block solution for 48 hr at 4°C. Primary antibodies used were,

Rat anti-BrdU (ab6326), Rabbit anti-Cav-1 (3267), Rabbit anti-DCX (4604), Mouse anti-DCX

(sc-271390), Guinea pig anti-DCX (AB2263), Rabbit anti-GFAP (Z0334), Rat anti-GFAP (13-

0300), Rabbit anti-MCM2 (4007), Chicken anti-Nestin (AvesLab), Rabbit anti-NeuN (ab104225),

Mouse anti-NeuN (ab104224), Rabbit anti-Sox2 (ab97959), Goat anti-Sox2 (AF2018), Rat anti-

Tbr2 (14-4875-82). After primary incubation, sections were washed 3 times in PBS containing

0.1% Tween-20 (PBST) for 15 mins per wash. Sections were incubated with secondary

antibodies diluted in block solution for 1.5-2 hr at RT. Secondary donkey antibodies used were

ordered from Jackson ImmunoResearch Labs. Sections were washed 3 times in PBST for 15

mins per wash, counterstained with DAPI, and mounted on Superfrost Plus slides

(ThermoFisher) with Prolong Gold Antifade Mountant (Invitrogen).

For immunocytochemistry (ICC), culture media was removed, coverslips containing cells

were washed two times with 1X PBS and fixed in 4% PFA in 1X PBS for 20 min at RT. Cells

were washed two times with 1X PBS and blocked in 1X PBS containing 0.3 M Glycine, 0.2%

Triton X-100 and 5% NDS for 30 minutes at RT. For EdU detection, cells were processed

according to the Click-iT EdU Cell Proliferation Kit (Invitrogen) instructions prior to blocking step.

For BrdU immunodetection, cells were treated in pre-warmed 1N HCl for 40 min at 37°C

followed by incubation with 0.1 M sodium borate buffer pH 8.5 for 10 min at RT prior to blocking

step. Primary antibodies were diluted in blocking solution and incubated 24 hr at 4°C. Primary

antibodies used were, Rabbit anti-DCX (ab18723) and Mouse anti-βIII-tubulin (MA1-118). After

primary incubation, cells were washed 3 times in PBST for 5 mins per wash and incubated with

secondary antibodies diluted in blocking solution for 45 min at RT. Cells were then washed 3

times in PBST for 5 mins per wash, counterstained with DAPI and mounted with Prolong Gold

Antifade Mountant (Invitrogen). Images were acquired at 40x or 63x magnification using

confocal microscopy (Zeiss LSM 710).

GFP Retrovirus Hippocampal Stereotaxic Injection

The Retrovirus/Moloney Murine Leukemia Virus (MMLV) construct carrying GFP (RV-

GFP) was generously provided by Dr. Jenny Hsieh (University of Texas at San Antonio, TX).

The viral particles were packaged as previously described (Lybrand et al., 2021). The packaging

process was carried out at the Vector Core Facility of the University of Illinois Chicago (UIC)

under the guidance of the Hsieh lab. The viral titer was $1.67 \times 10\square$ TU/ml for this study.

Stereotaxic surgeries were conducted on mice aged 4 to 4.5 months (16-18 weeks).

Anesthesia was induced using a 2.5% isoflurane and 100% oxygen mixture, and buprenorphine

(0.01 mg/kg) was administered preoperatively for analgesia. A total of 1 µl of retrovirus was

bilaterally injected into the dentate gyrus (DG) at three stereotaxic coordinates: AP = −1.50 mm,

ML = ±0.60 mm, and DV = −2.25 mm; AP = −2.00 mm, ML = ±1.30 mm, and DV = −2.15 mm;

and AP = −2.55 mm, ML = ±1.90 mm, and DV = −2.20 mm. Injections were delivered at a rate of

70 nl/min using a 1 µl Hamilton microsyringe. The needle was positioned at the target site for 5

minutes before injection and remained in place for an additional 10 minute post-injection to

prevent viral backflow. Mice were sacrificed 28 days after the procedure, and injection sites

were confirmed through histological and immunohistochemical analysis.

Neuronal Reconstruction, Sholl and Dendritic Analysis

Imaging of new neurons (labeled with RV-GFP) in the dentate gyrus of CORN and TAM

injected NestinCre$^{ERT2/+}$; CAV-1$^{fl/fl}$ was accomplished with Neurolucida 360 (MBF Bioscience,

Inc.) and a Zeiss LSM 710 confocal microscope. Neurons were imaged under 20X magnification

and centralized to include the soma and all neurites. Once image was acquired, Neurolucida

would be used to create a 3D tracing of the cell. Briefly, soma tracing would be used to create

and automated tracing of the cell body. Once the soma was finished, dendritic tree tracing was

used, using the user guide mode (a semi-automated setting where Neurolucida automatically

detects the thickness of the processes while their path is being guided by the user) to create the

final tracing of the cell.

For the Sholl analysis, the Neurolucida explorer software (MBF Bioscience, Inc.) was

used to report the number of intersections per pre-determined radius for each neuronal trace.

Starting at a radius of 10 μm from the center of the soma, a continuous shell of concentric

circles with 10μm between them was used to determine the number of intersections per radius.

For determining the total surface area (μm^2) and volume (μm^3) of the dendritic tree, Neurolucida

explorer software was used. Using the neuron summary function in the branched structure

analysis of the software, a total surface area and volume for all the dendritic branches of the cell

were analyzed. For the Sholl analysis, the number of intersections per radius was analyzed

using a mixed effects model with the Geisser-Greenhouse correction followed by Bonferroni

post-hoc test for multiple comparisons.

Transmission Electron Microscopy

Cells were washed 3 times with 1X PBS and centrifuged at 300g for 5 min to form

pellets. Cell pellets were fixed in 1.6% glutaraldehyde in 100 mM sodium phosphate, pH 7.4 for

1 hr at room temperature as similarly described(Sandhu JK, 2021). Samples were post-fixed

with 1% osmium tetroxide for 1 hr and dehydrated using an ascending series of

ethanol (through 100% absolute). Sample were then embedded in LX112 epoxy resin and

polymerized at 60°C for 3 days. Ultrathin sections (~75 nm) were collected onto copper grids

and stained with uranyl acetate and lead citrate, respectively. Specimens were examined using

a JEOL JEM-1400F transmission electron microscope at 80 kV. Micrographs were acquired

using an AMT Side-Mount Nano Sprint Model 1200S-B and Biosprint 12M-B cameras, loaded

with AMT Imaging software V.7.0.1.

RNA Isolation

To isolate RNA, cells were washed 2 times with 1X PBS and RNA was isolated using a

RNAeasy Plus Mini Kit (Qiagen).

Quantitative Real-Time PCR

Quantitative real-time PCR (qRT-PCR) was used to measure RNA with Luna Universal

One-Step RT-qPCR Kit (New England Biolabs) via CFX Connect Real-Time PCR Detection

System (Bio-Rad). Target gene expression was normalized to gene expression of GAPDH or β-

Actin.

Western blotting

Cells were washed 3 times with 1X PBS and lysed on ice for 10 mins in RIPA buffer

containing protease and phosphatase inhibitor cocktails (ThermoFisher). Lysed samples were

sonicated on ice at 20% power 3 times for 15 s with 5 s rest between sonication. Samples were

centrifuges at 10,000g for 15 mins to remove insoluble material and cellular debris. The

supernatant was collected and protein concentration determined by BCA Protein Assay Kit

(ThermoFisher). Samples were prepared in sample buffer consisting of NuPAGE LDS Sample

Buffer and NuPAGE reducing agent followed by boiling for 5 mins at 95°C. Protein were

separated on Bolt Bis-Tris Plus SDS-PAGE gels (Invitrogen) with MES SDS running buffer

(Invitorgen) and transferred to 0.2 μm nitrocellulose membranes via the iBlot 2 Dry Blotting

System (Invitrogen) Membranes were blocked for 1 hr at RT in 5% non-fat dry milk (milk) diluted

in TBS containing 0.1% Tween-20 (TBST). Primary antibodies were diluted in 5% milk in TBST

and membranes incubated overnight at 4°C. Primary antibodies used were, Rabbit-anti-

caveolin-1 (3267), Mouse anti-βIII-tubulin (MA1-118), Rabbit-anti-TOM20 (42406S), Rabbit-anti-

DRP1 (8570), and Mouse-anti-MFN2 (67487-1-Ig). Membranes were washed 3 times for 15

mins per wash with TBST followed by incubation with HRP conjugated secondary antibodies

[Goat anti-mouse IgG-HRP (W402B), and Goat anti-rabbit IgG-HRP (W4011)] diluted in 5% milk

in TBST for 2 hr at RT. Membranes were washed 3 times for 15 mins per wash and developed

with ECL Super Signal Kit via Kodak RP X-OMAT developer or Azure Biosystems 300q Image

Western Blot Imaging System. Band intensities were quantified in Fiji (NIH) from scanned

images and total protein expression was normalized to GAPDH or β-Actin protein expression.

Cell Quantification

For *in vivo* experiments, every sixth section of brain tissue was quantified using

unbiased stereology (StereoInvestigator, MBF Biosciences). Under the optical fractionator

workflow, contours of the DG were traced under 10x magnification and cells counted under 63x

magnification. A 120 μm x 120 μm counting frame with a 2 μm guard zone on both sides of the

section and a counting grid size determined by sampling 35% or 50% of the contour was used.

The volume of DG (μm^3) was determined by multiplying the area of contour by the measured

mounted thickness of the section. Total cell counts were normalized to volume of DG (μm^3)

counted per mouse. Alternatively, for DCX and NeuN cell quantification, 30 µm z-stacks of the

DG were acquired at 25x magnification using confocal microscopy (Zeiss LSM 710) and positive

cells counted from maximum projection images in Fiji (NIH). For in vitro experiments, 10 µm z-

stacks were acquired at 40x or 63x magnification using confocal microscopy (Zeiss LSM 710)

and positive cells counted from maximum projection images in Fiji (NIH).

Behavior Tests

All mice were handled 3-5 days for 2 min per mouse per day prior to the start of behavior

testing.

Elevated Plus Maze

The elevated plus maze (EPM) test was used to examine anxiety-like behavior. The

EPM apparatus consisted of two open arms without walls and two closed arms with opaque

walls. Mice were placed at the center of the apparatus facing an open arm and allowed to freely

explore for 5 mins. The EPM apparatus was cleaned with 70% ethanol between mice. Video

recordings were analyzed by Ethovision XT v16 software (Noldus) and the time spent in the

open and closed arms as well as frequency of entries into the arms was calculated.

Novel Object Location

The novel object location (NOL) test was used to examine spatial learning and memory

behavior as similarly described in X. A 38 cm x 51 cm x 30 cm opaque white plastic chamber

box with one short end containing a black circle wall print and the other short end containing a

black vertical line wall print was utilized. On day 1 of the test, mice were habituated in the empty

box for 10 mins followed by placement of 2 identical objects equal distance a part on the short

end of the box containing the black circle wall print for 10 mins. Objects were removed for 5 min

and then replaced in same locations for another 10 min session. Mice were placed back in

home cages for 24 hrs. On day 2, mice were habituated in the empty box for 10 mins and tested

by placing one of the objects in a novel location diagonally to the other one along the short end

of the box with black vertical line wall print. Mice were allowed to explore the objects for 5 mins.

Video was captured for every trial and exploration time with each object was manually scored

blinded. Chambers were cleaned with 70% ethanol between mice. The discrimination index (DI)

was calculated as $DI = (T_N - T_o)/(T_n + T_o)$, where T_N is the exploration time with the object in the

new location and T_o the exploration time with the object in the old location.

Contextual Fear Discrimination

This contextual fear discrimination test was conducted similarly as described(Clemenson

et al., 2014). All contexts consisted of 17.8 cm x 17.8 cm x 30.5 cm chamber housed in a sound

isolation cubicle (Coulbourn Instruments). Context A consisted of two translucent plexiglass

walls, two walls comprised of alternating black and white metal tiles and a stainless-steel rod

floor. Context A also had a 28V exhaust fan. Context B had no fan and consisted of two walls

comprised of alternating black and white metal tiles, two walls of black vertical line print and a

stainless-steel rod floor. To test fear generalization, Context B was exchanged with Context C

that had no fan, a circular wall insert made of up of a black and white plastic sheet and a black

plastic floor covering. On day 1 of the test, mice were exposed to Context A for 10 mins. On day

2, mice were placed in Context A, received a 2s foot shock (0.7mA) immediately upon entry and

remained in Context A for an additional 28s (a total of 30s in Context A). Thirty mins later, mice

were placed back in Context A for a 3 min exposure trial followed by placement in Context B for

3 min exposure trial. On day 3 (24 hr post-shock), mice were placed in Context A for 3 mins or

Context B for 3 mins in a counter-balanced order. For a less difficult discrimination test, mice

followed the same sequence as stated but were placed in Context C instead of Context B on

day 2 and 3. FreezeFrame software (Actimetrics) was used for video recording and analysis of

freezing behavior in each context. Chamber walls and floor were cleaned with 70% ethanol

between mice. The discrimination index (DI) was calculated as $DI = (F_A - F_x)/(F_A + F_x)$, where F_A is

the percentage of freezing in Context A and F_x is the percentage of freezing in either Context B

or Context C.

Nano-LC-MS/MS analysis

Cells were isolated as described in Section 2.2.4 and washed 3 times in 1X PBS prior to

freezing in liquid nitrogen. Cell pellets were lysed in 10% sodium dodecyl sulfate (SDS)

containing 100 mM triethylammonium bicarbonate (TEAB) with Pierce protease inhibitor

(ThermoFisher Scientific) supplemented, sonicated, and centrifuged at 14,000 rpm for 10

minutes. Protein supernatant was collected and quantified using a BCA Protein Assay Kit

(ThermoFisher). 100 ug of protein per sample was processed, trypsin digested in S-Trap

microcolumns (Protifi), and peptides labeled according to the TMT10plex Isobaric Label

Reagent Set (ThermoFisher) manufacture instructions. 100 fmol green fluorescent protein

(GFP) per μg of protein was spiked in each sample before digestion. Each individual isobaric

labeled sample was combined, lyophilized, and resuspended in 10 mM ammonium hydroxide

prior to fractionation by high pH reversed-phase liquid chromatography. 60 fractions were

collected and further concatenated into 20 fractions by combining 3 fractions for every 20

fractions apart(Yang et al., 2012). All fractions were dried down completely then resuspended in

0.1% formic acid. Peptide chromatographic separations and mass detection occurred with an

Agilent 1260 nano/capillary HPLC system (Agilent Technologies) coupled to an Q-Exactive

Orbitrap mass spectrometer (MS, ThermoFisher Scientific). Peptides were loaded onto an

Acclaim PepMap 100 trap column (75 μm × 2 cm nanoViper, C18, 3 μm 100 Å, ThermoFisher

Scientific) at flow rate of 2 μL/min. Peptides were further separated using a Zorbax 300SB-C18

column (0.075×150 mm, 3.5 μm 300 Å) (Agilent Technologies) at a flow rate of 0.25 μL/min, and

eluted using a 5-30% mobile phase B gradient consisting of 0.1% formic acid in acetonitrile over

90 minutes. Data was collected in the data-dependent acquisition (DDA) mode at a mass

resolution of 70,000 and scan range 375-2000 m/z. Automatic gain control (AGC) target was set

at 1×10^6 for a maximum injection time (IT) of 100 ms. The top 10 most abundant precursors

(charge state between 2 and 5) were selected for MS/MS analysis. MS/MS spectra were

acquired at a resolution of 35,000, AGC target at 1×10^6, and maximum IT of 50 ms. All raw

mass spectrometry data is publicly available on MassIVE with project ID MSV000090289

(ftp://massive.ucsd.edu/MSV000090289/).

The raw MS/MS data for each sample was searched against the curated SwissProt *Mus*

musculus database in Proteome Discoverer software (v2.3.0.523, ThermoFisher Scientific),

where trypsin was selected as the protease with 2 or less missed cleavage sites, precursor and

fragment mass error tolerances set to 10 ppm and ± 0.02 Da, and only peptide precursors of +2,

+3, +4 were analyzed. Peptide variable modifications allowed during the search were: oxidation

((+15.995 Da; M), TMT6 (+229.163 Da; S, T), and acetylation (+42.011 Da; N-terminus),

whereas carbamidomethyl (+57.021 Da; C) and TMT (+229.163 Da; any N-terminus) were set

as static modifications. Samples were grouped as iNSC Cav-1 KO (n=3) and iNSC Cav-1 WT

(n=3). Protein identifications were accepted if they contained at least 2 unique peptides and

abundances normalized to total peptide abundance. Differentially expressed proteins (DEP) for

the iNSC Cav-1 KO NSPCs relative to the iNSC Cav-1 WT NSPCs were determined by applying

ANOVA with a p-value of < 0.05.

Protein pathway and mitochondria protein analysis

Protein pathway analysis was conducted on DEP using Ingenuity Pathway Analysis

(IPA, Qiagen) and Gene Ontology (GO) enrichment analysis (Ashburner et al., 2000; Sandhu

JK, 2021) using g:Profiler (Raudvere et al., 2019) (ve107_eg54_p17_bf42210). Functional

enrichment maps were constructed in Cytoscape (Shannon et al., 2003) with EnrichmentMap

pipeline collection applications (Reimand et al., 2019). Mitochondria pathways and functional

groups were determined by manual curation using the MitoCarta 3.0 dataset (Rath et al., 2021).

Mitochondria and membrane isolation

Cells were washed 3 times with 1X PBS and mitochondria were isolated according to the manufacturer instructions using the Mitochondria Isolation Kit (ThermoFisher). Briefly, cells were pelleted by centrifugation at 300g for 5 mins at 4^{o}C. Reagent A containing protease and phosphatase inhibitor cocktails (ThermoFisher) was added to the cell pellet and placed on ice for 2 mins. Cells were then lysed by 100 strokes using a Dounce tissue homogenizer. Following the addition of Reagent C, the lysate was centrifuged at 700g for 10 mins at 4^{o}C. The supernatant was then collected and centrifuged at 3000g for 15 min at 4^{o}C to pellet mitochondria from the cytosol. The cytosol containing supernatant was collected and the mitochondria enriched pellet was washed with Reagent C and then centrifuged for 12000g for 5 mins at 4^{o}C. The supernatant was centrifuged at 5.5×10^{5} rpm for 1 h to collect the membrane fraction. The pellets corresponding to mitochondria and plasma membrane were dissolved with 2% CHAPS in TBS and processed for BCA protein estimation and western blotting as described.

TMRM Live Cell Imaging and mitochondria morphology analysis

Tetramethylrhodamine, methyl ester (TMRM) live cell imaging was conducted similarly as described(Jiang et al., 2022). Singly dissociated NPCs were grown on PLO/laminin coated glass bottom 24 well at a density of 50,000 cells/well. 5 days after plating, media was replaced with proliferation media containing 50 nM TMRM. Cells were incubated with TMRM for 30 mins at 37^{o}C. Immediately prior to imaging, cells were washed 2 times with PBS followed by replacement of proliferation media lacking phenol red. 2D images were captured with a 63X objective using a Zeiss LSM880 META confocal microscope with heated stage and a DPSS 561-10 laser set at 0.1% laser power with GaAsP detection.

For quantification of mitochondria morphology, images were analyzed with ImageJ as described (Chaudhry et al., 2020). For optimal results, all images were pre- and postprocessed

to reduce noise. The workflow and procedures for image processing and thresholding followed:

1) 2D images, operating on each slice in the stack, were preprocessed using "subtract

background" (radius = 1 µm) to remove background noise; *2)* "sigma filter plus"

(radius = 0.1 µm, 2.0 sigma) to reduce noise and smooth object signal while preserving

edges; *3)* "enhance local contrast" (block size = 64, slope = 2.0 for 2D) to enhance dim

areas while minimizing noise amplification; and *4)* "gamma correction" (value = 0.80 for 2D) to

correct any remaining dim areas. To identify mitochondria in the images, the "adaptive

threshold" method was used. In the adaptive threshold plugin, block size was set to an

equivalent of 1.25 µm. The thresholded images were then postprocessed using "despeckle" and

then "remove outliers" (radius = 0.15 µm) to remove residual noise. Further to analyze

mitochondrial 2D morphology, the image was first processed and thresholded (steps above),

and the resulting binary image was used as the input for the "analyze particles" command

(size = 0.06 μm^2-infinity, circularity = 0.00–1.00), measuring for "area," "perimeter," and

"shape descriptors." For branch length analysis, the "skeletonize 2D/3D" command was applied

to the thresholded image to produce a skeleton map, and the "analyze skeleton" command was

used to calculate branch lengths in the skeletonized network.

For 2D time lapse videos, images were captured with 3x digital zoom for 120 s at frame

rate of 3 Hz. Mitochondria velocity and total distance traveled were measured using the

Trackmate (Ershov et al., 2022) plugin in ImageJ.

Statistics

In all graphs, data is shown as mean ± SEM. Prism (Graphpad) was used for statistical

analysis with tests indicated in figure legends. The following was used for p values: ns > 0.05,

*p < 0.05, **p < 0.01, ***p < 0.001, and ****p < 0.0001.

Acknowledgments

We thank Dr. Trongha Phan for his expertise and guidance on behavioral experiments

and scientific discussion. We thank Dr. Pavan Kumar, Dr. Muskan Gupta, Abhi Ramakrishnan,

Stephanie Dunning for help with animal husbandry, genotyping, and IP injections. We thank

Karen Rakowiecki for assistance with cell culture experiments. We thank Dr. Christian Peters

and Shana Netherton for use of Elevated Plus Maze equipment. We thank Dr. Ying Jiang for

expertise and guidance on the mitochondrial assays. We also thank Dr. Peter Toth and Dr. Ke

Ma at the UIC Fluorescence Imaging Core for guidance on microscopy imaging, Figen Seiler at

the UIC Electron Microscopy Core for TEM imaging, and Dr. Balaji Ganesh at the UIC Flow

Cytometry Core for assistance with the Mitochondria Seahorse Assay. Graphical abstract,

Figure 2A, 3A, and 4E were created Biorender.com. This work was financially supported by

National Institutes of Health, National Institution on Aging (NIA) F30AG071144 (TKLS),

T32AG067468, AG033570, AG076940, AG062251, AG060238 (OL),

NINDS/NIA:R01NS114413 (SMC) and Together Strong NPC Foundation (SMC).

890 Author Contributions

TKLS and DP contributed equally to the design and execution of experiments, analysis

of the data, writing and revisions of the manuscript. LA, EQ, YI, KO, EQ, PK, JM and WL

assisted with data collection and analysis. AS and JB assisted with the design of the studies

and generation of the mouse models used. RDM and SMC assisted with the data analysis,

interpretation, and revision of the manuscript. OL supervised the project, assisted with the data

analysis, interpretation and revised the manuscript. All authors contributed to the review of the

manuscript.

899 Declarations of Interests

The authors declare no conflict of interests.

References

Ahmed A, Isaksen TJ, Yamashita T (2021) Protocol for mouse adult neural stem cell isolation

and culture. STAR Protoc 2:100522.

Artegiani B, Lyubimova A, Muraro M, van Es JH, van Oudenaarden A, Clevers H (2017) A

Single-Cell RNA Sequencing Study Reveals Cellular and Molecular Dynamics of the

Hippocampal Neurogenic Niche. Cell Rep 21:3271-3284.

Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight

SS, Eppig JT, Harris MA, Hill DP, Issel-Tarver L, Kasarskis A, Lewis S, Matese JC,

Richardson JE, Ringwald M, Rubin GM, Sherlock G (2000) Gene ontology: tool for the

unification of biology. The Gene Ontology Consortium. Nat Genet 25:25-29.

Beckervordersandforth R et al. (2017) Role of Mitochondrial Metabolism in the Control of Early

Lineage Progression and Aging Phenotypes in Adult Hippocampal Neurogenesis.

Neuron 93:560-573 e566.

Blochet C, Buscemi L, Clement T, Gehri S, Badaut J, Hirt L (2020) Involvement of caveolin-1 in

neurovascular unit remodeling after stroke: Effects on neovascularization and

astrogliosis. J Cereb Blood Flow Metab 40:163-176.

Bonds JA, Shetti A, Stephen TKL, Bonini MG, Minshall RD, Lazarov O (2020) Deficits in

hippocampal neurogenesis in obesity-dependent and -independent type-2 diabetes

mellitus mouse models. Sci Rep 10:16368.

Bonds JA, Shetti A, Bheri A, Chen Z, Disoukey A, Tai L, Mao M, Head BP, Bonini MG, Haus

JM, Minshall RD, Lazarov O (2019) Depletion of Caveolin-1 in Type-2 Diabetes Model

Induces Alzheimer's disease Pathology Precursors. J Neurosci.

Britton R, Liu AT, Rege SV, Adams JM, Akrapongpisak L, Le D, Alcantara-Lee R, Estrada RA,

Ray R, Ahadi S, Gallager I, Yang CF, Minami SS, Braithwaite SP, Czirr E, Campbell MK

(2022) Molecular and histological correlates of cognitive decline across age in male

C57BL/6J mice. Brain Behav 12:e2736.

Cao G, Yang G, Timme TL, Saika T, Truong LD, Satoh T, Goltsov A, Park SH, Men T, Kusaka

N, Tian W, Ren C, Wang H, Kadmon D, Cai WW, Chinault AC, Boone TB, Bradley A,

Thompson TC (2003) Disruption of the caveolin-1 gene impairs renal calcium

reabsorption and leads to hypercalciuria and urolithiasis. Am J Pathol 162:1241-1248.

Carvalho C, Moreira PI (2023) Metabolic defects shared by Alzheimer's disease and diabetes: A

focus on mitochondria. Curr Opin Neurobiol 79:102694.

Chaudhry A, Shi R, Luciani DS (2020) A pipeline for multidimensional confocal analysis of

mitochondrial morphology, function, and dynamics in pancreatic beta-cells. Am J Physiol

Endocrinol Metab 318:E87-E101.

Choi SH, Bylykbashi E, Chatila ZK, Lee SW, Pulli B, Clemenson GD, Kim E, Rompala A, Oram

MK, Asselin C, Aronson J, Zhang C, Miller SJ, Lesinski A, Chen JW, Kim DY, van Praag

H, Spiegelman BM, Gage FH, Tanzi RE (2018) Combined adult neurogenesis and

BDNF mimic exercise effects on cognition in an Alzheimer's mouse model. Science 361.

Chow BW, Nunez V, Kaplan L, Granger AJ, Bistrong K, Zucker HL, Kumar P, Sabatini BL, Gu C

(2020) Caveolae in CNS arterioles mediate neurovascular coupling. Nature 579:106-

110.

Clelland CD, Choi M, Romberg C, Clemenson GD, Jr., Fragniere A, Tyers P, Jessberger S,

Saksida LM, Barker RA, Gage FH, Bussey TJ (2009) A functional role for adult

hippocampal neurogenesis in spatial pattern separation. Science 325:210-213.

Clemenson GD, Lee SW, Deng W, Barrera VR, Iwamoto KS, Fanselow MS, Gage FH (2014)

Enrichment rescues contextual discrimination deficit associated with immediate shock.

Hippocampus 25:385-392.

Coelho P, Fao L, Mota S, Rego AC (2022) Mitochondrial function and dynamics in neural stem

cells and neurogenesis: Implications for neurodegenerative diseases. Ageing Res Rev

80:101667.

Demars M, Hu YS, Gadadhar A, Lazarov O (2010) Impaired neurogenesis is an early event in

the etiology of familial Alzheimer's disease in transgenic mice. J Neurosci Res 88:2103-

2117.

Demars MP, Bartholomew A, Strakova Z, Lazarov O (2011) Soluble amyloid precursor protein:

a novel proliferation factor of adult progenitor cells of ectodermal and mesodermal origin.

Stem Cell Res Ther 2:36.

Demars MP, Hollands C, Zhao Kda T, Lazarov O (2013) Soluble amyloid precursor protein-

alpha rescues age-linked decline in neural progenitor cell proliferation. Neurobiol Aging

34:2431-2440.

Egawa J, Zemljic-Harpf A, Mandyam CD, Niesman IR, Lysenko LV, Kleschevnikov AM, Roth

DM, Patel HH, Patel PM, Head BP (2018) Neuron-Targeted Caveolin-1 Promotes

Ultrastructural and Functional Hippocampal Synaptic Plasticity. Cereb Cortex 28:3255-

3266.

Encinas JM, Michurina TV, Peunova N, Park JH, Tordo J, Peterson DA, Fishell G, Koulakov A,

Enikolopov G (2011) Division-coupled astrocytic differentiation and age-related depletion

of neural stem cells in the adult hippocampus. Cell Stem Cell 8:566-579.

Ershov D, Phan MS, Pylvanainen JW, Rigaud SU, Le Blanc L, Charles-Orszag A, Conway

JRW, Laine RF, Roy NH, Bonazzi D, Dumenil G, Jacquemet G, Tinevez JY (2022)

TrackMate 7: integrating state-of-the-art segmentation algorithms into tracking pipelines.

Nat Methods 19:829-832.

Filchenko I, Blochet C, Buscemi L, Price M, Badaut J, Hirt L (2020) Caveolin-1 Regulates

Perivascular Aquaporin-4 Expression After Cerebral Ischemia. Front Cell Dev Biol 8:371.

Frederiksen K, McKay RD (1988) Proliferation and differentiation of rat neuroepithelial precursor

cells in vivo. J Neurosci 8:1144-1151.

Fridolfsson HN et al. (2012) Mitochondria-localized caveolin in adaptation to cellular stress and

injury. FASEB J 26:4637-4649.

979 Gadadhar A, Marr R, Lazarov O (2011) Presenilin-1 regulates neural progenitor cell

980   differentiation in the adult brain. J Neurosci 31:2615-2623.

981 Gioiosa L, Raggi C, Ricceri L, Jasmin JF, Frank PG, Capozza F, Lisanti MP, Alleva E,

982   Sargiacomo M, Laviola G (2008) Altered emotionality, spatial memory and cholinergic

983   function in caveolin-1 knock-out mice. Behav Brain Res 188:255-262.

984 Gonzalez F, Quinn JJ, Fanselow MS (2003) Differential effects of adding and removing

985   components of a context on the generalization of conditional freezing. J Exp Psychol

986   Anim Behav Process 29:78-83.

987 Guo N, Soden ME, Herber C, Kim MT, Besnard A, Lin P, Ma X, Cepko CL, Zweifel LS, Sahay A

988   (2018) Dentate granule cell recruitment of feedforward inhibition governs engram

989   maintenance and remote memory generalization. Nat Med 24:438-449.

990 Head BP, Hu Y, Finley JC, Saldana MD, Bonds JA, Miyanohara A, Niesman IR, Ali SS, Murray

991   F, Insel PA, Roth DM, Patel HH, Patel PM (2011) Neuron-targeted caveolin-1 protein

992   enhances signaling and promotes arborization of primary neurons. J Biol Chem

993   286:33310-33321.

994 Head BP, Peart JN, Panneerselvam M, Yokoyama T, Pearn ML, Niesman IR, Bonds JA,

995   Schilling JM, Miyanohara A, Headrick J, Ali SS, Roth DM, Patel PM, Patel HH (2010)

996   Loss of caveolin-1 accelerates neurodegeneration and aging. PLoS One 5:e15697.

997 Hu YS, Xu P, Pigino G, Brady ST, Larson J, Lazarov O (2010) Complex environment

998   experience rescues impaired neurogenesis, enhances synaptic plasticity, and attenuates

999   neuropathology in familial Alzheimer's disease-linked APPswe/PS1DeltaE9 mice.

1000   FASEB J 24:1667-1681.

1001 Huckleberry KA, Ferguson LB, Drew MR (2016) Behavioral mechanisms of context fear

1002   generalization in mice. Learn Mem 23:703-709.

1003 Ikezu T, Ueda H, Trapp BD, Nishiyama K, Sha JF, Volonte D, Galbiati F, Byrd AL, Bassell G,

1004   Serizawa H, Lane WS, Lisanti MP, Okamoto T (1998) Affinity-purification and

characterization of caveolins from the brain: differential expression of caveolin-1, -2, and

-3 in brain endothelial and astroglial cell types. Brain Res 804:177-192.

Jiang Y, Krantz S, Qin X, Li S, Gunasekara H, Kim YM, Zimnicka A, Bae M, Ma K, Toth PT, Hu

Y, Shajahan-Haq AN, Patel HH, Gentile S, Bonini MG, Rehman J, Liu Y, Minshall RD

(2022) Caveolin-1 controls mitochondrial damage and ROS production by regulating

fission - fusion dynamics and mitophagy. Redox Biol 52:102304.

Khacho M, Slack RS (2018) Mitochondrial dynamics in the regulation of neurogenesis: From

development to the adult brain. Dev Dyn 247:47-53.

Khacho M, Clark A, Svoboda DS, Azzi J, MacLaurin JG, Meghaizel C, Sesaki H, Lagace DC,

Germain M, Harper ME, Park DS, Slack RS (2016) Mitochondrial Dynamics Impacts

Stem Cell Identity and Fate Decisions by Regulating a Nuclear Transcriptional Program.

Cell Stem Cell 19:232-247.

Knowland D, Arac A, Sekiguchi KJ, Hsu M, Lutz SE, Perrino J, Steinberg GK, Barres BA,

Nimmerjahn A, Agalliu D (2014) Stepwise recruitment of transcellular and paracellular

pathways underlies blood-brain barrier breakdown in stroke. Neuron 82:603-617.

Kochan SMV, Malo MC, Jevtic M, Jahn-Kelleter HM, Wani GA, Ndoci K, Perez-Revuelta L,

Gaedke F, Schaffner I, Lie DC, Schauss A, Bergami M (2024) Enhanced mitochondrial

fusion during a critical period of synaptic plasticity in adult-born neurons. Neuron

112:1997-2014 e1996.

Lybrand ZR, Goswami S, Zhu J, Jarzabek V, Merlock N, Aktar M, Smith C, Zhang L, Varma P,

Cho KO, Ge S, Hsieh J (2021) A critical period of neuronal activity results in aberrant

neurogenesis rewiring hippocampal circuitry in a mouse model of epilepsy. Nat Commun

12:1423.

Masachs N, Charrier V, Farrugia F, Lemaire V, Blin N, Mazier W, Tronel S, Montaron MF, Ge S,

Marsicano G, Cota D, Deroche-Gamonet V, Herry C, Abrous DN (2021) The temporal

origin of dentate granule neurons dictates their role in spatial memory. Mol Psychiatry

26:7130-7140.

Miller SM, Sahay A (2019) Functions of adult-born neurons in hippocampal memory interference

and indexing. Nat Neurosci 22:1565-1575.

Mishra R, Phan T, Kumar P, Morrissey Z, Gupta M, Hollands C, Shetti A, Lopez KL,

Maienschein-Cline M, Suh H, Hen R, Lazarov O (2022) Augmenting neurogenesis

rescues memory impairments in Alzheimer's disease by restoring the memory-storing

neurons. J Exp Med 219.

Misko A, Jiang S, Wegorzewska I, Milbrandt J, Baloh RH (2010) Mitofusin 2 is necessary for

transport of axonal mitochondria and interacts with the Miro/Milton complex. J Neurosci

30:4232-4240.

Moreno-Jimenez EP, Flor-Garcia M, Terreros-Roncal J, Rabano A, Cafini F, Pallas-Bazarra N,

Avila J, Llorens-Martin M (2019) Adult hippocampal neurogenesis is abundant in

neurologically healthy subjects and drops sharply in patients with Alzheimer's disease.

Nat Med 25:554-560.

Nakashiba T, Cushman JD, Pelkey KA, Renaudineau S, Buhl DL, McHugh TJ, Rodriguez

Barrera V, Chittajallu R, Iwamoto KS, McBain CJ, Fanselow MS, Tonegawa S (2012)

Young dentate granule cells mediate pattern separation, whereas old granule cells

facilitate pattern completion. Cell 149:188-201.

Niibori Y, Yu TS, Epp JR, Akers KG, Josselyn SA, Frankland PW (2012) Suppression of adult

neurogenesis impairs population coding of similar contexts in hippocampal CA3 region.

Nat Commun 3:1253.

Oliveira SDS, Castellon M, Chen J, Bonini MG, Gu X, Elliott MH, Machado RF, Minshall RD

(2017) Inflammation-induced caveolin-1 and BMPRII depletion promotes endothelial

dysfunction and TGF-beta-driven pulmonary vascular remodeling. Am J Physiol Lung

Cell Mol Physiol 312:L760-L771.

Palmer TD, Takahashi J, Gage FH (1997) The adult rat hippocampus contains primordial neural

stem cells. Mol Cell Neurosci 8:389-404.

Parton RG (2018) Caveolae: Structure, Function, and Relationship to Disease. Annu Rev Cell

Dev Biol 34:111-136.

Parton RG, McMahon KA, Wu Y (2020) Caveolae: Formation, dynamics, and function. Curr

Opin Cell Biol 65:8-16.

Patel HH, Insel PA (2009) Lipid rafts and caveolae and their role in compartmentation of redox

signaling. Antioxid Redox Signal 11:1357-1372.

Rath S et al. (2021) MitoCarta3.0: an updated mitochondrial proteome now with sub-organelle

localization and pathway annotations. Nucleic Acids Res 49:D1541-D1547.

Raudvere U, Kolberg L, Kuzmin I, Arak T, Adler P, Peterson H, Vilo J (2019) g:Profiler: a web

server for functional enrichment analysis and conversions of gene lists (2019 update).

Nucleic Acids Res 47:W191-W198.

Reimand J, Isserlin R, Voisin V, Kucera M, Tannus-Lopes C, Rostamianfar A, Wadi L, Meyer M,

Wong J, Xu C, Merico D, Bader GD (2019) Pathway enrichment analysis and

visualization of omics data using g:Profiler, GSEA, Cytoscape and EnrichmentMap. Nat

Protoc 14:482-517.

Ruggiero A, Aloni E, Korkotian E, Zaltsman Y, Oni-Biton E, Kuperman Y, Tsoory M, Shachnai L,

Levin-Zaidman S, Brenner O, Segal M, Gross A (2017) Loss of forebrain MTCH2

decreases mitochondria motility and calcium handling and impairs hippocampal-

dependent cognitive functions. Sci Rep 7:44401.

Sahay A, Scobie KN, Hill AS, O'Carroll CM, Kheirbek MA, Burghardt NS, Fenton AA, Dranovsky

A, Hen R (2011) Increasing adult hippocampal neurogenesis is sufficient to improve

pattern separation. Nature 472:466-470.

Sandhu JK R-LM, Abulrob A. (2021) Molecular and Functional Characterization of Caveolae in

Mixed Cultures of Human NT-2 Neurons and Astrocytes. Neuroglia 2:68-88.

Saotome M, Safiulina D, Szabadkai G, Das S, Fransson A, Aspenstrom P, Rizzuto R,

Hajnoczky G (2008) Bidirectional Ca2+-dependent control of mitochondrial dynamics by

the Miro GTPase. Proc Natl Acad Sci U S A 105:20728-20733.

Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B,

Ideker T (2003) Cytoscape: a software environment for integrated models of

biomolecular interaction networks. Genome Res 13:2498-2504.

Shetti AU, Ramakrishnan A, Romanova L, Li W, Vo K, Volety I, Ratnayake I, Stephen T,

Minshall RD, Cologna SM, Lazarov O (2023) Reduced endothelial caveolin-1 underlies

deficits in brain insulin signalling in type 2 diabetes. Brain 146:3014-3028.

Shikanai M, Nishimura YV, Sakurai M, Nabeshima YI, Yuzaki M, Kawauchi T (2018) Caveolin-1

Promotes Early Neuronal Maturation via Caveolae-Independent Trafficking of N-

Cadherin and L1. iScience 7:53-67.

Terreros-Roncal J, Moreno-Jimenez EP, Flor-Garcia M, Rodriguez-Moreno CB, Trinchero MF,

Cafini F, Rabano A, Llorens-Martin M (2021) Impact of neurodegenerative diseases on

human adult hippocampal neurogenesis. Science 374:1106-1113.

Tobin MK, Musaraca K, Disouky A, Shetti A, Bheri A, Honer WG, Kim N, Dawe RJ, Bennett DA,

Arfanakis K, Lazarov O (2019) Human Hippocampal Neurogenesis Persists in Aged

Adults and Alzheimer's Disease Patients. Cell Stem Cell 24:974-982 e973.

Toda T, Parylak SL, Linker SB, Gage FH (2019) The role of adult hippocampal neurogenesis in

brain health and disease. Mol Psychiatry 24:67-87.

Toni N, Teng EM, Bushong EA, Aimone JB, Zhao C, Consiglio A, van Praag H, Martone ME,

Ellisman MH, Gage FH (2007) Synapse formation on neurons born in the adult

hippocampus. Nat Neurosci 10:727-734.

Trushina E, Du Charme J, Parisi J, McMurray CT (2006) Neurological abnormalities in caveolin-

1 knock out mice. Behav Brain Res 172:24-32.

Urban N, Blomfield IM, Guillemot F (2019) Quiescence of Adult Mammalian Neural Stem Cells:

A Highly Regulated Rest. Neuron 104:834-848.

van Elburg RA, van Ooyen A (2010) Impact of dendritic size and dendritic topology on burst

firing in pyramidal cells. PLoS Comput Biol 6:e1000781.

Vicidomini C, Guo N, Sahay A (2020) Communication, Cross Talk, and Signal Integration in the

Adult Hippocampal Neurogenic Niche. Neuron 105:220-235.

Virgintino D, Robertson D, Errede M, Benagiano V, Tauer U, Roncali L, Bertossi M (2002)

Expression of caveolin-1 in human brain microvessels. Neuroscience 115:145-152.

Wang S, Zhang Z, Almenar-Queralt A, Leem J, DerMardirossian C, Roth DM, Patel PM, Patel

HH, Head BP (2019) Caveolin-1 Phosphorylation Is Essential for Axonal Growth of

Human Neurons Derived From iPSCs. Front Cell Neurosci 13:324.

Wilhelmsson U, Lebkuechner I, Leke R, Marasek P, Yang X, Antfolk D, Chen M, Mohseni P,

Lasic E, Bobnar ST, Stenovec M, Zorec R, Nagy A, Sahlgren C, Pekna M, Pekny M

(2019) Nestin Regulates Neurogenesis in Mice Through Notch Signaling From

Astrocytes to Neural Stem Cells. Cereb Cortex 29:4050-4066.

Williams TM, Lisanti MP (2004) The caveolin proteins. Genome Biol 5:214.

Yang F, Shen Y, Camp DG, 2nd, Smith RD (2012) High-pH reversed-phase chromatography

with fraction concatenation for 2D proteomic analysis. Expert Rev Proteomics 9:129-134.

Zhao C, Deng W, Gage FH (2008) Mechanisms and functional implications of adult

neurogenesis. Cell 132:645-660.

Zhao C, Teng EM, Summers RG, Jr., Ming GL, Gage FH (2006) Distinct morphological stages

of dentate granule neuron maturation in the adult mouse hippocampus. J Neurosci 26:3-

11.

Zheng X, Boyer L, Jin M, Mertens J, Kim Y, Ma L, Ma L, Hamm M, Gage FH, Hunter T (2016)

Metabolic reprogramming during neuronal differentiation from aerobic glycolysis to

neuronal oxidative phosphorylation. Elife 5.

Zhou Y et al. (2022) Molecular landscapes of human hippocampal immature neurons across

lifespan. Nature 607:527-533.

1135

Figure Legends

Figure 1. iNSC Cav-1 KO mice have reduced levels of proliferating NSPC in the DG.

(A) Representative confocal images of GFAP (white), Nestin (red), MCM2 (green) and DAPI

(blue) markers in the DG of iNSC Cav-1 WT and iNSC Cav-1 KO mice at 3 months of age.

Yellow arrowheads indicate GFAP$^+$Nestin$^+$MCM2$^+$ cells. Scale bar, 25 μm.

(B-D) Quantification of total NSC (GFAP$^+$Nestin$^+$), quiescent NSC (GFAP$^+$Nestin$^+$MCM2$^-$) and

proliferating NSC (GFAP$^+$Nestin$^+$MCM2$^+$) in the DG of iNSC Cav-1 WT and iNSC Cav-1 KO

mice at 3 months of age. n=5 mice per group.

(E) Representative confocal images of GFAP (white), Nestin (red), MCM2 (green) and DAPI

(blue) positive cells in the DG of iNSC Cav-1 WT and iNSC Cav-1 KO mice at 6 months of age.

Yellow arrowheads indicate GFAP$^+$Nestin$^+$MCM2$^+$ cells. Scale bar, 25 μm.

(F-H) Quantification of total NSC (GFAP$^+$Nestin$^+$), quiescent NSC (GFAP$^+$Nestin$^+$MCM2$^-$) and

proliferating NSC (GFAP$^+$Nestin$^+$MCM2$^+$) in the DG of iNSC Cav-1 WT and iNSC Cav-1 KO

mice at 6 months of age. n=5 mice per group.

(I-L) Clonogenic proliferation assay in NSPC isolated neurospheres from iNSC Cav-1 WT and

iNSC Cav-1 KO mice. The number of clones (neurospheres), average clone diameter and

number of cells after dissociation of clones on day 5 of the assay were quantified per well. Scale

bar, 100 μm.

(M) 5-bromo-2'-deoxyuridine (BrdU) pulse strategy to quantify changes in NSC fate and

differentiation in DG of iNSC Cav-1 WT and iNSC Cav-1 KO mice at 6 months of age. Mice

were injected daily for 12 consecutive days and sacrificed 24 hr after the last injection.

(N-O) Confocal images of BrdU immunostaining and quantification total number of BrdU$^+$ cells in

the DG of 6-month-old iNSC Cav-1 KO and iNSC Cav-1 WT mice. Scale bar, 25 µm. n=4 mice

per group.

(P) Representative confocal images of BrdU (red), Sox2 (white), GFAP (green) and DAPI (blue)

markers in the DG of iNSC Cav-1 WT and iNSC Cav-1 KO mice at 6 months of age. Solid

yellow arrowheads indicate BrdU$^+$GFAP$^+$Sox2$^+$ cells. Outlined yellow arrowheads indicate

BrdU$^+$GFAP$^-$Sox2$^-$ cells. Scale bar, 25 µm.

(Q-S) Quantification of total BrdU$^+$ NSCs (GFAP$^+$Sox2$^+$), BrdU$^+$ NPCs (GFAP$^+$Sox2$^-$) and

percentage of BrdU$^+$ cell phenotypes normalized to total number of BrdU $^+$ cells in the DG of

iNSC Cav-1 WT and iNSC Cav-1 KO mice. n=4 mice per group.

Data represented as mean ± SEM. Data analyzed by unpaired two-tailed Student's t-test except

(S) which was analyzed by two-way ANOVA with Tukey multiple comparisons correction. ns p>

0.05, *p< 0.05, **p < 0.01, ***p < 0.001.

1170

Figure 2. Cav-1 regulates *in vivo* and *in vitro* differentiation of hippocampal NSPCs

(A) Representative confocal images of BrdU (white), DCX (Green), NeuN (red) and DAPI (blue)

markers in the DG of iNSC Cav-1 WT and iNSC Cav-1 KO mice at 6 months of age. Outlined

yellow arrowheads indicate BrdU$^+$DCX$^+$NeuN$^+$ cells. Scale bar, 25 µm.

(B-D) Quantification of BrdU$^+$ DCX expressing NPCs and neuroblasts (DCX$^+$NeuN$^-$), immature

neurons (DCX$^+$NeuN$^+$), and percentage of BrdU$^+$ immature neurons (DCX$^+$NeuN$^+$) normalized to

total number of BrdU$^+$ cells in the DG of iNSC Cav-1 WT and iNSC Cav-1 KO mice. n=4 mice

per group.

(E) Confocal images of DCX and NeuN immunostaining in the DG of iNSC Cav-1 WT and iNSC

Cav-1 KO mice at 6 months of age. Scale bar, 25 µm.

(F-H) Quantification of NPCs and neuroblasts (DCX$^+$NeuN$^-$) and immature neurons

(DCX$^+$NeuN$^+$) and total DCX expressing cells in the DG of iNSC Cav-1 WT and iNSC Cav-1 KO

mice. n=4 mice per group.

(I) Schematic representation of the protocol used for neural differentiation of primary

hippocampal NSPCs.

(J-L) Quantification of Cav-1, Nestin, and MAP2 transcript expression by RT-qPCR between

iNSC Cav-1 WT and iNSC Cav-1 KO NSPCs undergoing differentiation for 5 days. Expression

level normalized to d0 of iNSC Cav-1 WT NSPC (n=3 replicates).

(M) Immunoblot of Cav-1, Sox2 and β-III-Tubulin in NSPC isolated from iNSC Cav-1 KO and

iNSC Cav-1 WT mice undergoing differentiation for 7 days.

(N-P) Quantification of Cav-1, Sox2 and β-III-Tubulin normalized to GAPDH expression and

then normalized to d0 iNSC Cav-1 WT NSPC (n=3 replicates).

Data represented as mean ± SEM. Data analyzed by unpaired two-tailed Student's t-test (B-D

and F-H) and two-way ANOVA with Tukey multiple comparisons correction (J-L and N-P). ns p

> 0.05, *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, and ****$p < 0.0001$.

1196

Figure 3. iNSC Cav-1 KO mice display enhanced dendritic volume of newborn neurons

and improved contextual discrimination learning and memory.

(A) GFP retrovirus (RV-GFP) labeling strategy of newborn neurons. Mice were stereotaxic

injected with RV-GFP in the DG and then, sacrificed 30 days post injection for analysis.

(B) Confocal images of GFP$^+$ newborn neurons with zoomed dendritic tree arborization in the

DG of 5.5-month-old iNSC Cav-1 KO and iNSC Cav-1 WT mice. Scale bar, 25 μm and 10 μm.

(C) Sholl analysis data represented as mean number of intersections for each successive

30 μm segment as a function of the radial distance of the corresponding segment from the

soma.

(D-F) Quantification of total dendritic tree length, surface area and volume respectively. n=4

mice per group, 28-30 cells analyzed total per group.

(G) Schematic of contextual fear discrimination paradigm between Context A and B. See

Materials and methods for details.

(H-I) Quantification of percent freeze percent freeze (30 min post-shock) and discrimination

index in context A and B on Day 2 between the iNSC Cav-1 WT and iNSC Cav-1 KO mice.

n=12 per genotype.

(J-K) Quantification of percent freeze (24 hr post-shock) and discrimination index in context A

and B on Day 3 between the iNSC Cav-1 WT and iNSC Cav-1 KO mice. n=12 per genotype.

Data represented as mean $\pm$ SEM. Data analyzed in (C) by mixed model a mixed effects model

with the Geisser-Greenhouse correction followed by Bonferroni post-hoc test for multiple

comparisons; in D-F, I and K analyzed by unpaired two-tailed Student's t-test; and in H and J by

a two-way ANOVA with Tukey's multiple comparisons correction. ns $p > 0.05$, *$p < 0.05$, **$p <$

0.01.

1220

Figure 4. Loss of Cav-1 in mouse hippocampal NSPCs alters mitochondrial protein

pathway expression and promotes mitochondrial elongation.

(A) Schematic of proteomic workflow for hippocampal NSPCs isolated from iNSC Cav-1 KO

and iNSC Cav-1 WT mice. See Materials and methods for details.

(B) Volcano plot of proteins identified in iNSC Cav-1 KO vs iNSC Cav-1 WT cells. Significantly

upregulated proteins in red and significantly downregulated proteins in blue. Data represented

as Log_2FoldChange (FC) of mean abundance of proteins normalized to iNSC Cav-1 WT (n=3

iNSC Cav-1 KO and n=3 iNSC Cav-1 WT). DEP determined by a one-way ANOVA with a p-

value of < 0.05.

(C) Bar chart showing the top 10 Ingenuity Pathway Analysis (IPA) altered pathways in the

iNSC Cav-1 KO compared to iNSC Cav-1 WT hippocampal NSPCs (ANOVA, $p < 0.05$).

(D) Cytoscape cluster mapping of altered GO pathways in hippocampal NSPCs isolated from

iNSC Cav-1 KO and iNSC Cav-1 WT mice. Analysis based on Log_2FC ratio of DEPs with a

significant cut off value of $p < 0.05$.

(E) Volcano plot of Mitocarta 3.0 (Rath et al., 2021) proteins identified in iNSC Cav-1 KO vs

iNSC Cav-1 WT cells. Metabolism related protein in orange; Protein Import, Sorting &

Homeostasis in green; Signaling and Molecule Transport in blue, Mitochondria Central Dogma

in purple; and Mitochondria Dynamics and Surveillance in pink.

(F) Representative confocal image of Caveolin-1 (Cav-1, Green), Mitochondrial import receptor

subunit TOM20 homolog (Tom-20, green) and DAPI (blue) immunostaining in WT hippocampal

NSPCs. Scale bar 10 µm.

(G) Representative transmission electron microscopy (TEM) images of NSPCs isolated from

iNSC Cav-1 KO and iNSC Cav-1 WT mice. Mitochondria are shaded in green; scale bar = 1 µm

with zoomed image scale bar = 500 nm.

(H-K) Quantification of the number of mitochondria (mito) per cell imaged, area per

mitochondria, perimeter per mitochondria and circularity per mitochondria using ultrastructural

electron microscopy. n=20 cells per genotype with n=460 mitochondria for iNSC Cav-1 WT and

n=370 mitochondria for iNSC Cav-1 KO analyzed.

Data represented as mean $\pm$ SEM. Data analyzed by unpaired two-tailed Student's t-test. **p

< 0.01, ***$p < 0.001$, and ****$p < 0.0001$.

1251

Figure 5. Cav-1 regulates proliferation and differentiation in human iPSCs-derived NPCs.

(A) Schematic representation of the protocol used for neural induction of human induced-

pluripotent stem cells (iPSCs) to neural progenitor cells (NPC), neural precursor cells

(Precursor) and forebrain neurons (Neurons).

(B) Immunoblot of Cav-1 isolated from Cav-1 WT iPSCs, NPC, Precursor and 10 weeks old

Neurons.

(C) Quantification of Cav-1 normalized to GAPDH expression and then normalized to Cav-1 WT

iPSCs (n=3 replicates).

(D) Immunoblots of Cav-1, β-III-Tubulin and GAPDH from iPSCs derived Cav-1 KO and Cav-1

WT NPCs, Precursor cells and 10 weeks old neurons.

(E-F) Quantification of Cav-1 and β-III-Tubulin normalized to GAPDH expression in iPSCs

derived Cav-1 KO and Cav-1 WT (n=3 replicates).

(G) Representative confocal images of DCX (red) and DAPI (blue) positive cells in the iPSCs

derived Cav-1 KO and Cav-1 WT NPCs, precursor cells and 5 weeks old neurons. Scale bar,

100 μm.

(H) Representative confocal images of β-III-Tubulin (red) and DAPI (blue) positive cells in the

iPSCs derived Cav-1 KO and Cav-1 WT NPCs, precursor cells and 5 weeks old neurons. Scale

bar, 100 μm.

(I) Quantification of DCX^+DAPI^+ cells iPSCs derived Cav-1 KO and Cav-1 WT NPCs, precursor

cells and 5 weeks old neurons. n=9 images per group.

(J) Quantification of $β-III-Tubulin^+DAPI^+$ cells iPSCs derived Cav-1 KO and Cav-1 WT NPCs,

precursor cells and 5 weeks old neurons. n=9 images per group.

(K) EdU cell proliferation assay in iPSCs derived Cav-1 WT and Cav-1 KO neural progenitor

cells. Yellow arrowheads indicate EdU^+DAPI^+ cells. Scale bar, 50 μm.

(L) Quantification of the percentage of EdU^+ cells to the total DAPI comparing iPSCs derived

Cav-1 WT and Cav-1 KO NPCs.

Data represented as mean ± SEM. Data analyzed by unpaired two-tailed Student's t-test (E, F,

L), one-way ANOVA with Tukey multiple comparisons correction (C), two-way ANOVA with

Sidak's multiple comparisons correction (E, F, I, J). ns $p > 0.05$, *$p < 0.05$, **$p < 0.01$, ***$p <$

0.001, and ****$p < 0.0001$.

1282

Figure 6. Cav-1 rescue restores mitochondrial morphology and dynamics via interaction

with Mfn2 and Drp1 in iPSCs derived NPCs.

(A) Live imaging experimental schematic representation with (Cav-1 rescue) and without

lentivirus (LV) infection to NPCs.

(B) Representative confocal image of Caveolin-1 (Cav-1, Green), Mitochondrial import receptor

subunit TOM20 homolog (Tom-20, green) and DAPI (blue) immunostaining in Ctrl and KO

NPCs. Below left image is of the Cav-1 KO NPCs infected with LV-GFP (Ctrl; green) and below

right is with LV-Cav-1 (green). Scale bar 50 µm.

(C) Representative live-cell images of TMRM staining from Cav-1 Ctrl, Cav-1 KO, Cav-1 KO

NPCs with LV-GFP, and LV-Cav-1. NPCs were incubated with 50 nM TMRM for 30 min

followed by confocal microscopy visualization. Mitochondria were skeletonized in ImageJ. Scale

bar, 50 µm.

(D-F) Quantification of mitochondria area per cell, mitochondria perimeter per cell, and

circularity per cell in Cav-1 Ctrl, Cav-1 KO, Cav-1 KO LV- GFP, Cav-1 KO LV- Cav-1 NPCs.

N=25 cells per group.

(G) Quantification of skeletonized mitochondria branch length per mitochondria in Cav-1 Ctrl,

Cav-1 KO, Cav-1 KO LV- GFP, Cav-1 KO LV-Cav-1 NPCs. N=25 cells per group.

(H) Association of Cav-1 with MFN2 and DRP1. Cav-1 Ctrl NPCs lysates were

immunoprecipitated with a Cav-1 polyclonal antibody and subjected to Western blot analysis

with Cav-1, MFN2 and DRP1 Abs to assess co-IP.

(I) Immunoblot of Cav-1, Drp-1, mitofusion-2 (Mfn-2) and Tom-20 in total cell lysate (Whole cell),

mitochondrial (Mito) enriched, membrane, and cytosol fraction of Cav-1 Ctrl and Cav-1 KO

NPCs.

(J-M) Normalization of Cav-1 KO Drp-1 and Mfn-2 protein expression in whole cell, mito,

membrane, and cytosol enriched fractions to Cav-1 Ctrl expression.

Data representative of n=3, 6-well plate per genotype. Data represented as mean ± SEM. Data

analysed by unpaired two-tailed Student's t-test except in (D-G) was analyzed by two-way

ANOVA with Tukey's multiple comparisons correction. $*p < 0.05$, $**p < 0.01$, $***p < 0.001$ and

$****p < 0.0001$.

1312

Figure 7. Restoring Cav-1 levels normalizes neuronal differentiation process.

(A) Representative Western blots showing expression levels of Caveolin-1 (Cav-1), DCX, β-III-

Tubulin, and GAPDH across neural progenitor cells (NPCs), precursors (PRE), and 10-week-old

neurons, transduced with either LentiVirus (LV)-GFP (control) or LV-Cav-1.

(B-D) Quantifications of Cav-1, DCX, and β-III-Tubulin normalized to GAPDH expression in

iPSCs derived Cav-1 KO NPCs, transduced with either LV-GFP or LV-Cav-1, NPCs-induced

differentiation into precursor and neurons. Data representative of n=3, 6-well plate per

treatment. Data represented as mean ± SEM. Data analyzed by unpaired two-tailed Student's

t-test (neurons), and one-way ANOVA with Tukey multiple comparisons correction (NPCs and

PRE). $*p < 0.05$, $***p < 0.001$.

(E) Schematic illustrating Cav-1's role in facilitating mitochondrial fusion-fission dynamics in

NPCs, thereby regulating neuronal differentiation.

1325

Figure S1. Generation of inducible Nestin specific Cav-1 knockout (iNSC Cav-1 KO)

mouse model. iNSC Cav-1 KO mice displays altered levels of NSC proliferation yet no

changes immature neuron formation at 3 months of age.

(A) Schematic of NestinCre$^{ERT2/+}$;Cav-1$^{fl/fl}$ mice, where mice were injected at 4-5 weeks of age

with either corn oil (Corn) or tamoxifen (TAM) for 5 consecutive days to generate control mice

(iNSC Cav-1 WT) and Cav-1 knockout mice (iNSC Cav-1 KO), respectively.

(B) RT-qPCR quantification of Cav-1 transcript expression from NSPC isolated from the iNSC

Cav-1 KO and iNSC Cav-1 WT mice.

(C) Cav-1 immunoblot, and quantification normalized to β-actin of protein lysate from NSPC

isolated from the iNSC Cav-1 KO and iNSC Cav-1 WT mice.

(D) Immunocytochemistry of Cav-1 and confocal imaging of Cav-1 (green) and DAPI (blue) in

hippocampal NSPC isolated from iNSC Cav-1 WT and iNSC Cav-1 KO mice shown. Scale bar,

20 μm.

(E) Ultrastructural electron micrographs of caveolae and clathrin coated vesicles in hippocampal

NSPC isolated from iNSC Cav-1 WT and iNSC Cav-1 KO mice. Solid green arrowhead

indicates caveolae. Open green arrowhead indicates clathrin coated vesicle. Scale bar, 200 nm.

(F) Representative images of TBR2 (red), ASCL1 (green), and DAPI (blue) markers in the DG of

iNSC cav-1 WT and iNSC Cav-1 KO mice at 3 months of age. Yellow arrowheads indicate

TBR2$^+$ASCL1$^+$ cells. Scale bar, 50 μm.

(G-I) Quantification of total intermediate progenitor cells (TBR2$^+$ ASCL1$^+$) in the DG of iNSC

Cav-1 WT and iNSC Cav-1 KO mice at 3 months of age. n=4 mice per group.

(J) Representative images of TBR2 (red), ASCL1 (green), and DAPI (blue) positive cells in the

DG of iNSC cav-1 WT and iNSC Cav-1 KO mice at 6 months of age. Yellow arrowheads

indicate TBR2$^+$ASCL1$^+$ cells. Scale bar, 50 μm.

(K-M) Quantification of total intermediate progenitor cells (TBR2$^+$ ASCL1$^+$) in the DG of iNSC

Cav-1 WT and iNSC Cav-1 KO mice at 6 months of age. n=4 mice per group.

(N-O) EdU uptake assay in NSPC isolated from iNSC Cav-1 WT and iNSC Cav-1 KO mice.

Quantification of the percentage of EdU+ cells to the total DAPI between NSPC isolated from

iNSC Cav-1 WT and iNSC Cav-1 KO mice. Scale bar, 20 μm.

(P) Representative confocal images of NeuN (red), DCX (green) and DAPI (blue) in the DG of

iNSC Cav-1 WT and iNSC Cav-1 KO mice at 3 months of age. Scale bar, 25 μm.

(Q-S) Quantification of total DCX expressing cells, NPC and Neuroblasts (DCX$^+$NeuN$^-$) and

immature neurons (DCX$^+$NeuN$^+$) in the DG of iNSC Cav-1 WT and iNSC Cav-1 KO mice at 3

months of age. n=5-6 mice per group.

Data represented as mean $\pm$ SEM. Data analyzed by unpaired two-tailed Student's t-test. ns p>

0.05, *p< 0.05, **p < 0.01, ***p < 0.001 ****p < 0.0001.

1362

Figure S2. iNSC Cav-1 KO mice display no anxiety-like behavior or altered performance

in a context generalization or spatial recognition learning and memory task compared to

iNSC Cav-1 WT mice at 6 months of age.

(A) Schematic of contextual fear generalization paradigm between Context A and C. See

Materials and methods for details.

(B-C) Quantification of percent freeze (30 min post-shock) and discrimination index in context A

and C on Day 2 between the NSC Cav-1 WT and iNSC Cav-1 KO mice. n=12 iNSC Cav-1 WT

and n=8 iNSC Cav-1 KO.

(D-E) Quantification of percent freeze (24 hr post-shock) and discrimination index in context A

and C on Day 3 between the NSC Cav-1 WT and iNSC Cav-1 KO mice. n=12 iNSC Cav-1 WT

and n=8 iNSC Cav-1 KO.

(F) Schematic of elevated plus maze (EPM). Mice were placed in the center with head towards

the open arm and allowed to explore for 5 mins. See Materials and methods for details.

(G-I) Quantification of time spent in the open arm, number of entries in the open arm and total

distance traveled in the EPM of iNSC Cav-1 WT and iNSC Cav-1 KO mice. n=7 per genotype.

(J) Schematic of novel object location (NOL). Mice were trained during two 10 min trials on day

1 followed by a 5 min testing trial on day 2 with one object moved to a new location. See

Materials and methods for details.

(K-L) Quantification of the percentage of time spent with objects in each location and the

discrimination index determined between new and old locations on Day 2 of task between the

iNSC Cav-1 KO and iNSC Cav-1 WT mice. n=7 per iNSC Cav-1 WT and n=5 iNSC Cav-1 KO.

Data represented as mean $\pm$ SEM. Data analyzed by unpaired two-tailed Student's t-test. ns p

> 0.05, **p < 0.01, ***p < 0.001.

1386

Figure S3. Loss of Cav-1 in mouse hippocampal NSPS alters trafficking dynamics.

(A) Representative live-cell images of TMRM staining in NSPCs isolated from iNSC Cav-1 KO

and iNSC Cav-1 WT mice. NSPCs were incubated with 50 nM TMRM for 30 min followed by

confocal microcopy visualization. Mitochondria were skeletonized in ImageJ. Scale bar, 10 µm.

(B-D) Quantification of mitochondria area per cell, mitochondria perimeter per cell, and

circularity per cell in iNSC Cav-1 WT and iNSC Cav-1 KO NSCPs. N=25 cells per group.

(E) Quantification of skeletonized mitochondria branch length per mitochondria in iNSC Cav-1

WT and iNSC Cav-1 KO NSCPs. N=25 cells per group.

(F) Quantification of TMRM fluorescence in iNSC Cav-1 WT and iNSC Cav-1 KO NSPC. N=25

cells per group.

(G) Representative images of mitochondria trafficking per minute using Trackmate (Ershov et

al., 2022) in iNSC Cav-1 WT and iNSC Cav-1 KO NSPCs. Scale bar 10 µm.

(H) Quantification of mitochondria velocity. n=10 cells per genotype with n=948 particles

(mitochondria) for iNSC Cav-1 WT and n=1259 particles (mitochondria) for iNSC Cav-1 KO

analyzed.

(I) Quantification of mitochondria total distanced traveled. n=10 cells per genotype with n=948

particles (mitochondria) for iNSC Cav-1 WT and n=1259 particles (mitochondria) for iNSC Cav-1

KO analyzed.

Data represented as mean $\pm$ SEM. Data analyzed in by unpaired two-tailed Student's t-test. ns

p > 0.05, *p < 0.05, **p < 0.01, ***p < 0.001 and ****p < 0.0001.

1407

1408

1409 **Table S1.** Differentially Expressed Proteins in iNSC Cav-1 KO vs iNSC Cav-1 WT hippocampal

1410 NSPCs (ANOVA p<0.05).

Accession	Gene Name	Abundance Ratio: iNSC Cav-1 KO / iNSC Cav-1 WT	Log2 (Abundance Ratio): iNSC Cav-1 KO / iNSC Cav-1 WT	Abundance Ratio p-Value: iNSC Cav-1 KO / iNSC Cav-1 WT	-log (p-value): iNSC Cav-1 KO / iNSC Cav-1 WT
P14824	Anxa6	1.554	0.640	4.81E-07	6.3181
P55012	Slc12a2	1.56	0.640	5.49E-07	6.2601
Q9WV92	Epb41l3	2.767	1.470	9.40E-07	6.0270
Q9WUR2	Eci2	0.845	-0.240	4.15E-06	5.3818
P13020-2	Gsn	1.352	0.430	6.37E-06	5.1959
Q3V3R1	Mthfd1l	0.867	-0.210	7.93E-06	5.1008
P19324	Serpinh1	0.849	-0.240	3.81E-05	4.4188
O70318	Epb41l2	1.107	0.150	4.90E-05	4.3099
P68404-2	Prkcb	0.768	-0.380	4.94E-05	4.3066
P97447	Fhl1	0.697	-0.520	5.20E-05	4.2841
P60710	Actb	1.091	0.130	6.97E-05	4.1567
Q9D394	Rufy3	0.871	-0.200	1.16E-04	3.9351
O88342	Wdr1	1.106	0.140	1.23E-04	3.9118
Q8R0Y6	Aldh1l1	0.764	-0.390	1.25E-04	3.9029
P16015	Ca3	1.531	0.610	1.46E-04	3.8367
Q9CZN7	Shmt2	0.874	-0.190	1.75E-04	3.7576
Q9DBV4	Mxra8	0.843	-0.250	4.09E-04	3.3885
Q3TVI8	Pbxip1	0.887	-0.170	4.38E-04	3.3584
Q80YX1	Tnc	0.905	-0.140	4.62E-04	3.3350
Q9Z127	Slc7a5	0.878	-0.190	4.66E-04	3.3315
Q9DCZ1	Gmpr	1.398	0.480	4.67E-04	3.3306
O88958	Gnpda1	0.826	-0.280	4.68E-04	3.3296
Q920E5	Fdps	1.143	0.190	5.19E-04	3.2850
P25911	Lyn	1.218	0.290	5.36E-04	3.2705
Q80X90	Flnb	1.108	0.150	5.61E-04	3.2509
Q91WG8	Gne	0.889	-0.170	6.18E-04	3.2090
Q99LC5	Etfa	0.932	-0.100	6.85E-04	3.1644
Q640R3	Hepacam	0.897	-0.160	6.89E-04	3.1618
Q9CQB5	Cisd2	0.939	-0.090	7.56E-04	3.1213
Q05186	Rcn1	0.918	-0.120	7.71E-04	3.1127

Q8BGT5	Gpt2	0.9	-0.150	8.46E-04	3.0725
P26041	Msn	1.197	0.260	9.34E-04	3.0295
Q8K2B3	Sdha	0.915	-0.130	0.0010	2.9933
O55137	Acot1	0.886	-0.180	0.0010	2.9812
Q9CZ13	Uqcrc1	0.909	-0.140	0.0011	2.9709
Q8BNU0	Armc6	0.861	-0.220	0.0012	2.9217
Q8BH04	Pck2	0.755	-0.410	0.0012	2.9147
Q99N94	Mrpl9	0.885	-0.180	0.0012	2.9111
Q9R0P9	Uchl1	1.153	0.210	0.0013	2.8723
Q8BTM8	Flna	1.164	0.220	0.0013	2.8711
Q9Z110	Aldh18a1	0.848	-0.240	0.0014	2.8472
P80313	Cct7	1.052	0.070	0.0016	2.8061
Q06185	Atp5me	0.903	-0.150	0.0016	2.7835
P25322	Ccnd1	0.944	-0.080	0.0017	2.7775
Q9WTR5	Cdh13	0.766	-0.380	0.0018	2.7441
P16045	Lgals1	1.136	0.180	0.0020	2.7095
P28652	Camk2b	0.933	-0.100	0.0020	2.7042
P48962	Slc25a4	0.876	-0.190	0.0021	2.6828
P09242	Alpl	0.762	-0.390	0.0021	2.6786
Q9CS84	Nrxn1	0.715	-0.480	0.0021	2.6771
Q9CQI7	Snrpb2	1.076	0.110	0.0021	2.6770
B2RRD7	Brpf1	1.212	0.280	0.0022	2.6649
P39447	Tjp1	0.871	-0.200	0.0022	2.6558
P97821	Ctsc	0.872	-0.200	0.0022	2.6495
Q61823	Pdcd4	0.813	-0.300	0.0023	2.6372
Q03265	Atp5f1a	0.927	-0.110	0.0023	2.6351
Q3UHD6	Snx27	0.915	-0.130	0.0026	2.5907
Q922D8	Mthfd1	0.931	-0.100	0.0027	2.5646
Q9JI59	Jam2	0.885	-0.180	0.0028	2.5511
Q8K2Y7	Mrpl47	0.898	-0.160	0.0029	2.5382
P48722	Hspa4l	1.087	0.120	0.0029	2.5353
Q9D9E0	Slc22a17	0.793	-0.340	0.0029	2.5324
P06745	Gpi	0.925	-0.110	0.0030	2.5211
Q64727	Vcl	1.218	0.280	0.0031	2.5152
Q9CYH2	Prxl2a	0.915	-0.130	0.0031	2.5037
Q925I1	Atad3	0.915	-0.130	0.0032	2.4956
Q91YJ3	Thyn1	0.648	-0.630	0.0033	2.4874
Q8BWT1	Acaa2	0.904	-0.150	0.0033	2.4849
Q6P3B9	Rbfa	0.8	-0.320	0.0036	2.4491
P09925	Surf1	0.935	-0.100	0.0036	2.4407

Q8JZN7	Rhot2	1.212	0.280	0.0037	2.4278
Q8BMS1	Hadha	0.911	-0.130	0.0039	2.4123
Q8BH44	Coro2b	0.838	-0.250	0.0039	2.4119
Q9JLJ2	Aldh9a1	1.149	0.200	0.0039	2.4104
Q61881	Mcm7	1.078	0.110	0.0039	2.4080
Q99LI2	Clcc1	0.901	-0.150	0.0040	2.3983
Q9R059	Fhl3	0.861	-0.220	0.0040	2.3982
Q91YT7	Ythdf2	1.045	0.060	0.0044	2.3596
Q61024	Asns	0.91	-0.140	0.0044	2.3596
Q3UGR5	Hdhd2	0.892	-0.170	0.0044	2.3570
Q9CQF0	Mrpl11	0.903	-0.150	0.0045	2.3488
Q00915	Rbp1	2.203	1.140	0.0045	2.3478
P70404	Idh3g	0.921	-0.120	0.0045	2.3434
Q8BH95	Echs1	0.936	-0.100	0.0046	2.3417
Q9CPQ8	Atp5mg	0.902	-0.150	0.0046	2.3332
Q8K009	Aldh1l2	0.858	-0.220	0.0047	2.3275
P59328	Wdhd1	1.146	0.200	0.0048	2.3219
P17182	Eno1	1.114	0.160	0.0050	2.3035
Q60932	Vdac1	0.91	-0.140	0.0050	2.2972
P05480	Src	1.092	0.130	0.0051	2.2946
Q8C050	Rps6ka5	0.883	-0.180	0.0051	2.2939
Q9CZD3	Gars	0.96	-0.060	0.0053	2.2753
P10852	Slc3a2	0.87	-0.200	0.0053	2.2721
Q9DBE0	Csad	0.892	-0.160	0.0053	2.2720
P97820	Map4k4	0.926	-0.110	0.0054	2.2686
Q812A2	Srgap3	0.903	-0.150	0.0054	2.2651
P18242	Ctsd	0.943	-0.080	0.0055	2.2632
Q61739	Itga6	0.918	-0.120	0.0057	2.2451
Q8R422	Cd109	1.129	0.170	0.0060	2.2231
Q99PU5	Acsbg1	0.916	-0.130	0.0060	2.2189
Q07235	Serpine2	0.858	-0.220	0.0062	2.2102
Q8CGY8	Ogt	0.94	-0.090	0.0062	2.2100
Q91ZX7	Lrp1	0.919	-0.120	0.0063	2.2015
Q80UM7	Mogs	0.913	-0.130	0.0064	2.1945
Q921N7	Tmem70	0.931	-0.100	0.0067	2.1771
Q9WVA4	Tagln2	1.321	0.400	0.0068	2.1670
Q99JY0	Hadhb	0.929	-0.110	0.0069	2.1613
Q8BPU7	Elmo1	0.873	-0.200	0.0069	2.1609
Q8JZZ7	Adgrl2	0.846	-0.240	0.0070	2.1578
O35604	Npc1	0.879	-0.190	0.0070	2.1560

Q9D0F9	Pgm1	0.884	-0.180	0.0072	2.1426
Q8BL66	Eea1	1.066	0.090	0.0073	2.1366
P56564	Slc1a3	0.914	-0.130	0.0074	2.1330
Q80XL6	Acad11	0.871	-0.200	0.0074	2.1313
Q8BXV2	Bri3bp	0.912	-0.130	0.0074	2.1283
Q6W8Q3	Pcp4l1	1.694	0.760	0.0076	2.1169
Q8JZK9	Hmgcs1	1.165	0.220	0.0077	2.1124
P38647	Hspa9	0.87	-0.200	0.0078	2.1105
Q62193	Rpa2	1.068	0.090	0.0078	2.1102
Q9DBT5	Ampd2	0.91	-0.140	0.0078	2.1102
Q64737	Gart	1.07	0.100	0.0078	2.1070
Q6P9J9	Ano6	1.026	0.040	0.0085	2.0683
Q9QY06	Myo9b	0.914	-0.130	0.0086	2.0660
Q8BH86	Dglucy	0.864	-0.210	0.0088	2.0534
Q05920	Pc	1.066	0.090	0.0089	2.0521
Q80SY5	Prpf38b	0.89	-0.170	0.0090	2.0457
Q91UZ5	Impa2	1.233	0.300	0.0094	2.0262
P70158	Smpdl3a	0.838	-0.250	0.0097	2.0134
Q8VHY0	Cspg4	1.117	0.160	0.0097	2.0112
O70475	Ugdh	1.084	0.120	0.0098	2.0105
Q3UTH8	Arhgef9	1.243	0.310	0.0099	2.0049
Q61070	Ei24	0.879	-0.190	0.0099	2.0025
Q9D2G2	Dlst	0.931	-0.100	0.0103	1.9869
Q61474	Msi1	0.905	-0.140	0.0103	1.9859
P56480	Atp5f1b	0.919	-0.120	0.0105	1.9798
P97868	Rbbp6	0.906	-0.140	0.0106	1.9765
P51881	Slc25a5	0.913	-0.130	0.0106	1.9736
Q9CXZ1	Ndufs4	1.08	0.110	0.0108	1.9680
Q9CWG8	Ndufaf7	0.834	-0.260	0.0108	1.9675
Q923W1	Tgs1	0.882	-0.180	0.0108	1.9648
P37913	Lig1	1.083	0.120	0.0109	1.9640
P51174	Acadl	0.957	-0.060	0.0110	1.9596
Q99MK8	Grk2	0.913	-0.130	0.0110	1.9583
P13864	Dnmt1	1.115	0.160	0.0112	1.9520
Q8CAF4	Nhsl1	0.89	-0.170	0.0113	1.9481
Q6PIC6	Atp1a3	0.445	-1.170	0.0113	1.9459
P42227	Stat3	1.236	0.310	0.0115	1.9401
Q6P5G6	Ubxn7	1.16	0.210	0.0117	1.9334
Q8CCJ3	Ufl1	0.891	-0.170	0.0118	1.9271
Q9D1E6	Tbcb	1.087	0.120	0.0119	1.9261

Q80SW1	Ahcyl1	0.94	-0.090	0.0119	1.9237
Q61879	Myh10	0.907	-0.140	0.0119	1.9235
Q9D2R0	Aacs	1.114	0.160	0.0120	1.9195
Q3U9G9	Lbr	1.061	0.090	0.0122	1.9152
P17751	Tpi1	0.96	-0.060	0.0122	1.9135
Q5SSZ5	Tns3	1.085	0.120	0.0124	1.9075
O35129	Phb2	0.9	-0.150	0.0131	1.8829
Q9DC50	Crot	0.916	-0.130	0.0133	1.8774
P97311	Mcm6	1.079	0.110	0.0134	1.8735
P24472	Gsta4	0.955	-0.070	0.0136	1.8652
Q9Z1Q5	Clic1	1.121	0.160	0.0137	1.8618
Q9DCJ9	Npl	0.822	-0.280	0.0139	1.8583
P54276	Msh6	1.032	0.050	0.0140	1.8554
Q9CY66	Gar1	0.945	-0.080	0.0141	1.8499
Q9CXW2	Mrps22	0.934	-0.100	0.0142	1.8472
Q9CQA3	Sdhb	0.916	-0.130	0.0142	1.8469
P97742	Cpt1a	0.849	-0.240	0.0143	1.8460
P61028	Rab8b	1.111	0.150	0.0143	1.8448
Q8CG47	Smc4	1.095	0.130	0.0144	1.8425
Q9ERG0	Lima1	1.215	0.280	0.0144	1.8411
Q9CX56	Psmd8	1.083	0.110	0.0149	1.8257
Q640M1	Utp14a	0.904	-0.140	0.0150	1.8241
Q9QYJ0	Dnaja2	0.934	-0.100	0.0152	1.8192
Q9ESP1	Sdf2l1	1.198	0.260	0.0154	1.8136
Q8BFU3	Rnf214	0.866	-0.210	0.0155	1.8111
Q8BFR5	Tufm	0.915	-0.130	0.0164	1.7857
Q6DIB4	Nol4l	1.342	0.420	0.0164	1.7857
Q80UG5	Sept9	0.97	-0.040	0.0164	1.7840
Q8VCW8	Acsf2	1.139	0.190	0.0166	1.7805
Q9DCW4	Etfb	0.91	-0.140	0.0166	1.7796
O88441	Mtx2	0.927	-0.110	0.0169	1.7727
Q3TDD9	Ppp1r21	0.918	-0.120	0.0170	1.7708
Q6P9S7	Galnt10	0.824	-0.280	0.0171	1.7668
Q93092	Taldo1	1.078	0.110	0.0172	1.7645
Q91VC9	Ghitm	0.85	-0.230	0.0176	1.7545
Q9DB77	Uqcrc2	0.91	-0.140	0.0176	1.7542
Q9D8N0	Eef1g	1.039	0.050	0.0178	1.7506
Q9CXL3		1.125	0.170	0.0187	1.7280
Q8C156	Ncaph	1.125	0.170	0.0189	1.7241
Q505B7	Zbtb8	0.88	-0.180	0.0191	1.7200

P00405	Mtco2	0.927	-0.110	0.0191	1.7191
Q9CQJ8	Ndufb9	1.111	0.150	0.0193	1.7147
O08749	Dld	0.903	-0.150	0.0195	1.7099
Q3USB7	Plcl1	0.695	-0.530	0.0196	1.7083
Q9CXI0	Coq5	0.841	-0.250	0.0199	1.7016
P13595-2	Ncam1	0.893	-0.160	0.0200	1.6998
O35316	Slc6a6	0.878	-0.190	0.0201	1.6974
P58242	Smpdl3b	0.802	-0.320	0.0202	1.6943
Q8QZT1	Acat1	0.926	-0.110	0.0202	1.6936
Q9JKF7	Mrpl39	0.921	-0.120	0.0206	1.6859
O09117	Sypl1	1.188	0.250	0.0208	1.6810
P60469	Ppfia3	0.847	-0.240	0.0209	1.6805
Q9CZX7	Pip4p2	0.928	-0.110	0.0212	1.6728
Q60931	Vdac3	0.916	-0.130	0.0213	1.6706
Q8R404	Mic 13	0.914	-0.130	0.0216	1.6653
Q76MZ3	Ppp2r1a	1.017	0.020	0.0216	1.6652
Q8JZX4	Rbm17	0.937	-0.090	0.0223	1.6516
Q9D8S3	Arfgap3	0.844	-0.240	0.0225	1.6486
Q9D328	Tmem35a	0.775	-0.370	0.0227	1.6447
Q04887	Sox9	1.132	0.180	0.0227	1.6439
Q8BMI3	Gga3	0.833	-0.260	0.0230	1.6389
Q8K4Z3	Naxe	0.928	-0.110	0.0232	1.6352
Q8BFZ9	Erlin2	0.91	-0.140	0.0234	1.6311
Q9Z0H4	Celf2	0.906	-0.140	0.0238	1.6243
Q9JME2	Chst11	0.913	-0.130	0.0239	1.6225
P53702	Hccs	0.901	-0.150	0.0246	1.6092
P05213	Tuba1b	1.024	0.030	0.0247	1.6074
P57780	Actn4	1.107	0.150	0.0249	1.6042
Q8R550	Sh3kbp1	1.179	0.240	0.0249	1.6031
Q64133	Maoa	0.915	-0.130	0.0253	1.5976
Q8BGD5	Cpt1c	0.762	-0.390	0.0255	1.5936
P47199	Cryz	1.166	0.220	0.0262	1.5812
Q501J6	Ddx17	0.977	-0.030	0.0264	1.5781
Q9JMH9-1	Myo18a	0.883	-0.180	0.0265	1.5764
Q62420	Sh3gl2	0.891	-0.170	0.0266	1.5749
Q78PY7	Snd1	0.965	-0.050	0.0268	1.5723
P80318	Cct3	1.023	0.030	0.0269	1.5696
P35951	Ldlr	1.061	0.090	0.0273	1.5644
Q6A028	Swap70	0.937	-0.090	0.0278	1.5560
Q922W5	Pycr1	0.738	-0.440	0.0279	1.5539

Q8C8R3-4	Ank2	0.961	-0.060	0.0280	1.5529
Q8VD75	Hip1	1.025	0.040	0.0281	1.5520
P11214	Plat	0.869	-0.200	0.0287	1.5429
A6H611	Mipep	0.916	-0.130	0.0288	1.5408
P46935	Nedd4	1.111	0.150	0.0289	1.5397
Q6ZPE2	Sbf1	0.864	-0.210	0.0290	1.5377
Q3UPF5	Zc3hav1	1.192	0.250	0.0291	1.5356
P12960	Cntn1	1.275	0.350	0.0297	1.5268
Q9WUM5	Suclg1	0.923	-0.120	0.0299	1.5237
Q7TNP2	Ppp2r1b	1.177	0.240	0.0300	1.5227
Q6PDL0	Dync1li2	0.937	-0.090	0.0302	1.5205
P70670	Naca	1.054	0.080	0.0303	1.5192
Q922K7	Nop2	0.946	-0.080	0.0307	1.5128
Q8BP56	Pgghg	0.742	-0.430	0.0307	1.5124
Q8R4N0	Clybl	0.871	-0.200	0.0308	1.5114
P70704-2	Atp8a1	0.814	-0.300	0.0310	1.5088
P26040	Ezr	0.856	-0.220	0.0310	1.5087
Q9CQN1	Trap1	0.939	-0.090	0.0310	1.5086
P67778	Phb	0.906	-0.140	0.0313	1.5049
Q9Z0X1	Aifm1	0.938	-0.090	0.0315	1.5012
Q9CQN7	Mrpl41	0.873	-0.200	0.0317	1.4992
Q99N87	Mrps5	0.957	-0.060	0.0319	1.4958
Q8R0F8	Fahd1	0.902	-0.150	0.0319	1.4955
O35621	Pmm1	0.898	-0.160	0.0323	1.4906
Q9WV55	Vapa	0.948	-0.080	0.0330	1.4816
O35691	Pnn	0.925	-0.110	0.0337	1.4727
Q9CRA4	Msmo1	1.114	0.160	0.0338	1.4713
P49615	Cdk5	0.927	-0.110	0.0343	1.4653
Q8CJ67	Stau2	0.757	-0.400	0.0348	1.4579
Q921L3	Tmco1	0.882	-0.180	0.0350	1.4559
Q8K354	Cbr3	0.966	-0.050	0.0351	1.4553
Q99N85	Mrps18a	0.888	-0.170	0.0352	1.4538
P97287	Mcl1	0.837	-0.260	0.0355	1.4497
Q61792	Lasp1	0.832	-0.270	0.0356	1.4485
Q8C0M9	Asrgl1	1.085	0.120	0.0357	1.4473
P18872-2	Gnao1	0.844	-0.250	0.0358	1.4459
Q99KI0	Aco2	0.883	-0.180	0.0362	1.4412
Q00PI9	Hnrnpul2	0.952	-0.070	0.0362	1.4407
Q69ZS7	Hbs1l	0.91	-0.140	0.0362	1.4407
Q6ZPT1	Klhl9	0.853	-0.230	0.0365	1.4374

Q810J8	Zfyve1	0.833	-0.260	0.0365	1.4374
Q07646	Mest	0.793	-0.330	0.0367	1.4348
Q9JM14	Nt5c	0.94	-0.090	0.0370	1.4315
Q91VD9	Ndufs1	0.971	-0.040	0.0371	1.4304
Q9CQV1	Pam16	0.905	-0.140	0.0376	1.4252
Q91WQ3	Yars	0.933	-0.100	0.0378	1.4228
Q9CQW2	Arl8b	0.958	-0.060	0.0380	1.4207
P62313	Lsm6	1.105	0.140	0.0382	1.4177
Q9D379	Ephx1	0.884	-0.180	0.0383	1.4173
Q3UQ44	Iqgap2	0.933	-0.100	0.0385	1.4140
Q9D8B3	Chmp4b	0.939	-0.090	0.0387	1.4125
Q9ESW8	Pgpep1	0.868	-0.200	0.0387	1.4121
Q9R0Y5	Ak1	0.816	-0.290	0.0389	1.4102
Q9Z2Z9	Gfpt2	1.289	0.370	0.0395	1.4034
Q8R2S9	Actr8	1.08	0.110	0.0396	1.4022
Q61127	Nab2	0.854	-0.230	0.0400	1.3975
P06151	Ldha	1.033	0.050	0.0408	1.3894
Q9QZQ8	H2afy	0.904	-0.150	0.0409	1.3881
P35922	Fmr1	1.094	0.130	0.0409	1.3881
Q9WUU9	Mcm3ap	0.805	-0.310	0.0412	1.3851
Q9D6R2	Idh3a	0.888	-0.170	0.0415	1.3821
P56382	Atp5f1e	0.908	-0.140	0.0416	1.3807
Q9CQE3	Mrps17	0.91	-0.140	0.0422	1.3749
Q9CZT8	Rab3b	1.307	0.390	0.0426	1.3708
Q9CQL4	Mrpl20	0.924	-0.110	0.0427	1.3691
Q8R4C2	Rufy2	0.895	-0.160	0.0429	1.3679
Q91YJ5	Mtif2	0.825	-0.280	0.0431	1.3659
Q2TPA8	Hsdl2	0.836	-0.260	0.0431	1.3658
Q99J62	Rfc4	1.069	0.100	0.0431	1.3656
Q9Z315	Sart1	0.923	-0.120	0.0434	1.3625
P63038	Hspd1	0.962	-0.060	0.0438	1.3587
Q9WUA3	Pfkp	1.189	0.250	0.0450	1.3469
Q9CQS8	Sec61b	0.936	-0.100	0.0450	1.3469
Q8VDC0	Lars2	0.862	-0.210	0.0452	1.3447
Q60875	Arhgef2	0.959	-0.060	0.0453	1.3438
Q06335	Aplp2	0.902	-0.150	0.0454	1.3432
Q4VBD2	Tapt1	0.89	-0.170	0.0454	1.3426
Q8CG48	Smc2	1.063	0.090	0.0456	1.3411
Q9JKN5	Olig1	0.864	-0.210	0.0458	1.3392
Q8BM55	Tmem214	1.064	0.090	0.0464	1.3334

G5E829	Atp2b1	1.142	0.190	0.0466	1.3318
Q8BP48	Metap1	1.124	0.170	0.0468	1.3298
P21956	Mfge8	0.889	-0.170	0.0468	1.3298
Q922B2	Dars	0.942	-0.090	0.0472	1.3259
Q9CR88	Mrps14	0.864	-0.210	0.0475	1.3232
Q9R0H0-2	Acox1	0.889	-0.170	0.0477	1.3214
P35550	Fbl	0.942	-0.090	0.0480	1.3191
Q62426	Cstb	1.178	0.240	0.0480	1.3186
Q01279	Egfr	0.921	-0.120	0.0481	1.3175
Q8BLV3	Slc9a7	0.915	-0.130	0.0483	1.3163
Q8BGW1	Fto	1.064	0.090	0.0484	1.3155
Q9JK38	Gnpnat1	0.887	-0.170	0.0488	1.3118
Q8BIJ6	Iars2	0.913	-0.130	0.0489	1.3111
P09055	Itgb1	1.077	0.110	0.0489	1.3104
O55028	Bckdk	0.837	-0.260	0.0494	1.3062
Q9CRB2	Nhp2	0.9	-0.150	0.0495	1.3053

1411

1412 **Table S2.** Ingenuity Pathway Analysis (IPA) of differentially expressed proteins (DEP) in iNSC

1413 Cav-1 KO vs Cav-1 WT hippocampal NSPCs.

Ingenuity Canonical Pathways	-log(p-value)	% DEP in pathway	Molecules
Mitochondrial Dysfunction	12.6	13.1	ACO2,AIFM1,Atp5e,ATP5F1A,ATP5F1B,ATP5MG,CPT1A,CPT1C,MAOA,MT-CO2,NDUFB9,NDUFS1,NDUFS4,RHOT2,SDHA,SDHB,SURF1,UQCRC1,UQCRC2,VDAC1,VDAC3
TCA Cycle II (Eukaryotic)	8.87	36.4	ACO2,DLD,DLST,IDH3A,IDH3G,SDHA,SDHB,SUCLG1
Superpathway of Cholesterol Biosynthesis	7.92	28.6	ACAA2,ACAT1,FDPS,HADHA,HADHB,HMGCS1,LBR,MSMO1
Oxidative Phosphorylation	7.87	12.7	Atp5e,ATP5F1A,ATP5F1B,ATP5MG,MT-CO2,NDUFB9,NDUFS1,NDUFS4,SDHA,SDHB,SURF1,UQCRC1,UQCRC2
Sirtuin Signaling Pathway	7.61	7.4	ACADL,ATP5F1A,ATP5F1B,CPT1A,CPT1C,LDHA,NDUFB9,NDUFS1,NDUFS4,PAM16,PCK2,SDHA,SDHB,SLC25A4,SLC25A5,STAT3,TUBA1B,UQCRC2,VDAC1,VDAC3
Superpathway of Geranylgeranyldiphosphate Biosynthesis I (via Mevalonate)	6.68	35.3	ACAA2,ACAT1,FDPS,HADHA,HADHB,HMGCS1
Isoleucine Degradation I	6.51	33.3	ACAA2,ACAT1,DLD,ECHS1,HADHA,HADHB
Ketogenesis	6.29	45.5	ACAA2,ACAT1,HADHA,HADHB,HMGCS1
Mevalonate Pathway I	5.86	38.5	ACAA2,ACAT1,HADHA,HADHB,HMGCS1
Fatty Acid β-oxidation I	5.84	20.0	ACAA2,ACSBG1,ACSF2,ECHS1,ECI2,HADHA,HADHB

ILK Signaling	5.42	7.3	ACTB,ACTN4,CCND1,FLNA,FLNB,ITGB1,MYH10,MYO18A,NACA,PPP2R1A,PPP2R1B,RHOT2,RPS6KA5,VCL
Ketolysis	5.06	44.4	ACAA2,ACAT1,HADHA,HADHB
Neutrophil Extracellular Trap Signaling Pathway	4.93	5.2	ATP5F1A,ATP5F1B,ITGB1,LYN,MCL1,NDUFB9,NDUFS1,NDUFS4,PAM16,PLCL1,PRKCB,SDHA,SDHB,SLC25A4,SLC25A5,SRC,UQCRC2,VDAC1,VDAC3
Folate Polyglutamylation	4.36	60.0	MTHFD1,MTHFD1L,SHMT2
Tetrahydrofolate Salvage from 5,10-methenyltetrahydrofolate	4.36	60.0	GART,MTHFD1,MTHFD1L
RHOGDI Signaling	4.22	6.1	ACTB,ARHGEF2,ARHGEF9,CDH13,EZR,GNAO1,ITGA6,ITGB1,MSN,MYH10,MYO18A,RHOT2,SRC
Germ Cell-Sertoli Cell Junction Signaling	4.08	6.8	ACTB,ACTN4,GSN,ITGA6,ITGB1,RAB8B,RHOT2,SRC,TJP1,TUBA1B,VCL
Regulation of Cellular Mechanics by Calpain Protease	4.02	9.2	ACTN4,CCND1,EGFR,EZR,ITGA6,ITGB1,SRC,VCL
Caveolar-mediated Endocytosis Signaling	3.69	9.6	ACTB,EGFR,FLNA,FLNB,ITGA6,ITGB1,SRC
Glutaryl-CoA Degradation	3.63	21.1	ACAA2,ACAT1,HADHA,HADHB
Mitochondrial L-carnitine Shuttle Pathway	3.63	21.1	ACSBG1,ACSF2,CPT1A,CPT1C
Folate Transformations I	3.46	33.3	MTHFD1,MTHFD1L,SHMT2
Valine Degradation I	3.45	19.0	DLD,ECHS1,HADHA,HADHB
tRNA Charging	3.42	13.2	DARS1,GARS1,IARS2,LARS2,YARS1
LPS/IL-1 Mediated Inhibition of RXR Function	3.31	5.2	ACOX1,ACSBG1,ACSF2,ALDH18A1,ALDH1L1,ALDH1L2,ALDH9A1,CHST11,CPT1A,CPT1C,HMGCS1,MAOA

Xenobiotic Metabolism CAR Signaling Pathway	3.28	5.9	ALDH18A1,ALDH1L1,ALDH1L2,ALDH9A1,CHST11,EGFR,PPP2R1A,PPP2R1B,PRKCB,SRC
Actin Cytoskeleton Signaling	3.23	5.1	ACTB,ACTN4,EZR,FLNA,GSN,IQGAP2,ITGA6,ITGB1,MSN,MYH10,MYO18A,VCL
Tight Junction Signaling	3.2	5.8	ACTB,ARHGEF2,JAM2,MYH10,MYO18A,PPP2R1A,PPP2R1B,TJP1,VAPA,VCL
Glycolysis I	3.15	16.0	ENO1,GPI,PFKP,TPI1
Tryptophan Degradation III (Eukaryotic)	3.08	15.4	ACAA2,ACAT1,HADHA,HADHB
UDP-N-acetyl-D-galactosamine Biosynthesis II	3.05	25.0	GNPDA1,GNPNAT1,GPI
Colanic Acid Building Blocks Biosynthesis	2.85	21.4	GPI,PMM1,UGDH
2-ketoglutarate Dehydrogenase Complex	2.8	50.0	DLD,DLST
Proline Biosynthesis I	2.8	50.0	ALDH18A1,PYCR1
BEX2 Signaling Pathway	2.76	7.8	CCND1,LGALS1,PPP2R1A,PPP2R1B,STAT3,SWAP70
Cell Cycle Control of Chromosomal Replication	2.75	9.4	CDK5,LIG1,MCM6,MCM7,RPA2
Sertoli Cell-Sertoli Cell Junction Signaling	2.73	5.0	ACTB,ACTN4,ITGA6,ITGB1,JAM2,RAB8B,SRC,TJP1,TUBA1B,VCL
Xenobiotic Metabolism PXR Signaling Pathway	2.65	5.3	ALDH18A1,ALDH1L1,ALDH1L2,ALDH9A1,CAMK2B,CDK5,CHST11,MAOA,PRKCB
Mismatch Repair in Eukaryotes	2.52	16.7	LIG1,MSH6,RFC4
Aryl Hydrocarbon Receptor Signaling	2.49	5.4	ALDH18A1,ALDH1L1,ALDH1L2,ALDH9A1,CCND1,CTSD,MCM7,SRC
Mechanisms of Viral Exit from Host Cells	2.42	10.3	ACTB,CHMP4B,PRKCB,SH3GL2

GDP-mannose Biosynthesis	2.41	33.3	GPI,PMM1
UDP-N-acetyl-D-glucosamine Biosynthesis II	2.41	33.3	GFPT2,GNPNAT1
Zymosterol Biosynthesis	2.41	33.3	LBR,MSMO1
Leukocyte Extravasation Signaling	2.4	4.8	ACTB,ACTN4,EZR,ITGB1,JAM2,MSN,PRKCB,SRC,VCL
Remodeling of Epithelial Adherens Junctions	2.39	7.8	ACTB,ACTN4,SRC,TUBA1B,VCL
Signaling by Rho Family GTPases	2.38	4.2	ACTB,ARHGEF2,ARHGEF9,CDH13,EZR,GNAO1,ITGA6,ITGB1,MSN,RHOT2,SEPTIN9
Huntington's Disease Signaling	2.25	4.1	ATP5F1A,ATP5F1B,CDK5,CTSD,EGFR,HIP1,HSPA9,PRKCB,PSMD8,SDHA,SDHB
GM-CSF Signaling	2.22	7.1	CAMK2B,CCND1,LYN,PRKCB,STAT3
AMPK Signaling	2.2	4.2	ACTB,AK1,CCND1,CPT1A,CPT1C,GNAO1,PCK2,PFKP,PPP2R1A,PPP2R1B
IL-8 Signaling	2.19	4.5	CCND1,CSTB,EGFR,GNAO1,LASP1,MAP4K4,PRKCB,RHOT2,SRC
Histidine Degradation III	2.15	25.0	MTHFD1,MTHFD1L
Estrogen Receptor Signaling	2.15	3.5	ATP5F1A,ATP5F1B,CCND1,EGFR,GNAO1,NDUFB9,NDUFS1,NDUFS4,PLCL1,PRKCB,SDHA,SDHB,SRC,UQCRC2
Reelin Signaling in Neurons	2.14	5.2	ARHGEF2,ARHGEF9,CAMK2B,CDK5,ITGB1,LYN,SRC
MSP-RON Signaling In Cancer Cells Pathway	2.12	5.2	CCND1,EGFR,FLNA,ITGA6,ITGB1,SRC,STAT3
Paxillin Signaling	2.1	5.7	ACTB,ACTN4,ITGA6,ITGB1,SRC,VCL
Role of Tissue Factor in Cancer	1.97	5.4	EGFR,ITGA6,ITGB1,LYN,RPS6KA5,SRC

Thrombin Signaling	1.97	4.1	ARHGEF2,ARHGEF9,CAMK2B,EGFR,GNAO1,PLCL1,PRKCB,RHOT2,SRC
3-phosphoinositide Degradation	1.96	4.4	ALPL,CA3,NT5C,PIP4P2,PPFIA3,PPP2R1A,PPP2R1B,TNS3
HER-2 Signaling in Breast Cancer	1.96	4.1	CCND1,EGFR,ITGB1,LYN,MT-CO2,PRKCB,SRC,STAT3,SURF1
Virus Entry via Endocytic Pathways	1.95	5.3	ACTB,FLNA,FLNB,ITGB1,PRKCB,SRC
D-myo-inositol-5-phosphate Metabolism	1.91	4.3	ALPL,CA3,NT5C,PIP4P2,PPFIA3,PPP2R1A,PPP2R1B,TNS3
Neuregulin Signaling	1.9	5.2	CDK5,EGFR,ITGA6,ITGB1,PRKCB,SRC
Insulin Secretion Signaling Pathway	1.9	3.8	CAMK2B,DLD,LYN,PC,PLCL1,PRKCB,RPS6KA5,SEC61B,SRC,STAT3
Glucocorticoid Receptor Signaling	1.86	3.1	ACTB,ATP5F1A,ATP5F1B,EGFR,HSPA9,NDUFB9,NDUFS1,NDUFS4,PC,PCK2,RPS6KA5,SDHA,SDHB,SRC,STAT3,UQCRC2
Xenobiotic Metabolism Signaling	1.85	3.7	ALDH18A1,ALDH1L1,ALDH1L2,ALDH9A1,CAMK2B,CHST11,MAOA,PPP2R1A,PPP2R1B,PRKCB
Gap Junction Signaling	1.82	4.2	ACTB,EGFR,NT5C,PLCL1,PRKCB,SRC,TJP1,TUBA1B
Macrophage Classical Activation Signaling Pathway	1.8	4.5	ACO2,DLD,NDUFS1,SDHA,SDHB,STAT3,UQCRC2
Asparagine Biosynthesis I	1.78	100.0	ASNS
Fcγ Receptor-mediated Phagocytosis in Macrophages and Monocytes	1.74	5.4	ACTB,EZR,LYN,PRKCB,SRC
Cholesterol Biosynthesis I	1.72	15.4	LBR,MSMO1
Cholesterol Biosynthesis II (via 24,25-dihydrolanosterol)	1.72	15.4	LBR,MSMO1

Cholesterol Biosynthesis III (via Desmosterol)	1.72	15.4	LBR,MSMO1
Integrin Signaling	1.69	3.9	ACTB,ACTN4,GSN,ITGA6,ITGB1,RHOT2,SRC,VCL
VEGF Signaling	1.67	5.2	ACTB,ACTN4,PRKCB,SRC,VCL
Cell Cycle Regulation by BTG Family Proteins	1.67	8.3	CCND1,PPP2R1A,PPP2R1B
Aldosterone Signaling in Epithelial Cells	1.66	4.2	HSPA4L,HSPA9,HSPD1,PLCL1,PRKCB,SLC12A2,TRAP1
WNT/β-catenin Signaling	1.65	4.1	CCND1,GNAO1,LRP1,PPP2R1A,PPP2R1B,SOX9,SRC
ERK/MAPK Signaling	1.64	3.9	ITGA6,ITGB1,PPP2R1A,PPP2R1B,PRKCB,RPS6KA5,SRC,STAT3
D-myo-inositol (1,4,5,6)-Tetrakisphosphate Biosynthesis	1.63	4.1	ALPL,CA3,NT5C,PPFIA3,PPP2R1A,PPP2R1B,TNS3
D-myo-inositol (3,4,5,6)-tetrakisphosphate Biosynthesis	1.63	4.1	ALPL,CA3,NT5C,PPFIA3,PPP2R1A,PPP2R1B,TNS3
Fatty Acid Activation	1.61	13.3	ACSBG1,ACSF2
Agrin Interactions at Neuromuscular Junction	1.6	6.0	ACTB,EGFR,ITGB1,SRC
Melatonin Signaling	1.56	5.8	CAMK2B,GNAO1,PLCL1,PRKCB
NAD Signaling Pathway	1.51	4.2	ACADL,AIFM1,ETFA,HSPD1,LDHA,SLC7A5
Necroptosis Signaling Pathway	1.51	4.2	CAMK2B,PAM16,SLC25A4,SLC25A5,VDAC1,VDAC3
Colorectal Cancer Metastasis Signaling	1.49	3.4	CCND1,EGFR,GNAO1,GRK2,LRP1,MSH6,RHOT2,SRC,STAT3
Superpathway of Inositol Phosphate Compounds	1.48	3.6	ALPL,CA3,NT5C,PIP4P2,PPFIA3,PPP2R1A,PPP2R1B,TNS3
Alanine Biosynthesis II	1.48	50.0	GPT2
Alanine Degradation III	1.48	50.0	GPT2

Asparagine Degradation I	1.48	50.0	ASRGL1
Glycine Biosynthesis I	1.48	50.0	SHMT2
Taurine Biosynthesis	1.48	50.0	CSAD
Thyroid Hormone Biosynthesis	1.48	50.0	CTSD
UDP-D-xylose and UDP-D-glucuronate Biosynthesis	1.48	50.0	UGDH
Regulation of Actin-based Motility by Rho	1.47	4.6	ACTB,GSN,ITGA6,ITGB1,RHOT2
γ-linolenate Biosynthesis II (Animals)	1.46	11.1	ACSBG1,ACSF2
Macropinocytosis Signaling	1.45	5.3	ACTN4,ITGB1,PRKCB,SRC
CDK5 Signaling	1.42	4.5	CDK5,ITGA6,ITGB1,PPP2R1A,PPP2R1B
Synaptic Long Term Depression	1.4	3.7	GNAO1,LYN,NT5C,PLCL1,PPP2R1A,PPP2R1B,PRKCB
Epithelial Adherens Junction Signaling	1.38	3.9	EGFR,MYH10,PPP2R1A,PPP2R1B,SRC,VCL
Xenobiotic Metabolism AHR Signaling Pathway	1.38	5.1	ALDH18A1,ALDH1L1,ALDH1L2,ALDH9A1
Putrescine Degradation III	1.37	10.0	ALDH9A1,MAOA
Clathrin-mediated Endocytosis Signaling	1.36	3.6	ACTB,HIP1,ITGB1,LDLR,SH3GL2,SH3KBP1,SRC
TR/RXR Activation	1.36	5.0	ENO1,LDLR,PFKP,RAB3B
3-phosphoinositide Biosynthesis	1.35	3.6	ALPL,CA3,NT5C,PPFIA3,PPP2R1A,PPP2R1B,TNS3
5-aminoimidazole Ribonucleotide Biosynthesis I	1.31	33.3	GART
L-carnitine Biosynthesis	1.31	33.3	ALDH9A1
N-acetylglucosamine Degradation I	1.31	33.3	GNPDA1

NADH Repair	1.31	33.3	NAXE
Retinoate Biosynthesis II	1.31	33.3	RBP1

1414