Long-read genome assembly of the Japanese parasitic wasp *Copidosoma*

***floridanum* (Hymenoptera: Encyrtidae)**

Kouhei Toga [1,2], Takuma Sakamoto [3], Miyuki Kanda [1,4], Keita Tamura [1], Keisuke Okuhara [2,4], Hiroko

Tabunoki [3,5], Hidemasa Bono [1,2]*

[1]Laboratory of BioDX, PtBio Co-Creation Research Center, Genome Editing Innovation Center,

Hiroshima University, 3-10-23 Kagamiyama, Higashi-Hiroshima city, Hiroshima, 739-0046, Japan

[2]Laboratory of Genome Informatics, Graduate School of Integrated Sciences for Life, Hiroshima

University, 3-10-23 Kagamiyama, Higashi-Hiroshima city, Hiroshima, 739-0046, Japan

[3]Department of Science of Biological Production, Graduate School of Agriculture, Tokyo University

of Agriculture and Technology, 3-5-8 Saiwai-cho, Fuchu, Tokyo, 183-8509, Japan

[4]Research and Development Department, PtBio Inc., 3-10-23 Kagamiyama, Higashi-Hiroshima city,

Hiroshima, 739-0046, Japan

[5]Cooperative Major in Advanced Health Science, Graduate School of Bio-Applications and System

Engineering, Tokyo University of Agriculture and Technology, 3-5-8 Saiwai-cho, Fuchu, Tokyo, 183-

8509, Japan

* Correspondence:

Hidemasa Bono

Email: bonohu@hiroshima-u.ac.jp

Tel.: +81 82 424 4013

Keywords: *Copidosoma floridanum*, Pacific Biosciences high-fidelity reads, genome assembly

Abstract

Copidosoma floridanum is a cosmopolitan species and an egg-larval parasitoid of the Plusiine moth. *C. floridanum* has a unique development mode called polyembryony, in which thousands of genetically identical embryos are produced from a single egg. Some embryos develop into sterile soldier larvae, and their developmental patterns differ between the US and Japanese *C. floridanum* strains. Genome sequencing can accelerate our understanding of the molecular bases underlying polyembryony, including the production of soldier castes. However, only the genome sequence of the US strain has been reported. In the present study, we determined the genome sequence of the Japanese strain using Pacific Biosciences high-fidelity reads and generating a highly contiguous assembly (552.7 Mb, N50: 17.9 Mb). Gene prediction and annotation identified 13,886 transcripts derived from 10,786 gene models. We searched the genomic differences between US and Japanese strains. Among gene models predicted in this study, 100 gene loci in the Japanese strain had extremely different gene structure from those in the US strain. This was accomplished through the functional annotation (GGSEARCH) and long-read sequencing. Genomic differences between strains were also reflected to amino acid sequences of *vasa* that plays a central role in caste determination in this species. The genome assemblies constructed in this study will facilitate the genomic comparisons between Japanese and US strains, leading to our understanding of detail genomic regions responsible for the ecological and physiological characters of *C. floridanum.*

Introduction

Copidosoma floridanum (Hymenoptera: Encyrtidae) is a worldwide distributed egg-larval parasitoid of the Plusiinae moth (Lepidoptera, Noctuidae, Plusiinae) (Guerrieri and Noyes 2005). This species reproduces by polyembryony, producing >2,000 genetically identical individuals from a single egg (Ode and Strand 1995). Polyembryony has evolved independently in the four Hymenoptera families (Encyrtidae, Platygastridae, Braconidae, and Dryinidae); however, encyrtid wasps have the most extreme form regarding clone size (Segoli et al. 2010). Embryos develop into larvae that feed on the host tissues until pupation. In some encyrtid wasps, the embryos develop into two types of larvae: reproductive larvae and the sterile soldier larvae. Sterile soldier larvae possesses large mandibles, and contribute to the defense of their clonal siblings by attacking against a competitor in the host (Cruz 1981; Strand 2003; Iwabuchi 2019). The phenomenon of solitary individuals forming cooperative groups with a reproductive division of labor can be considered as one of the major evolutionary events that brought complexity to life on Earth (Szathmáry and Smith 1995; Whyte 2021).

The process and molecular basis of soldier development in *C. floridanum* have been described and investigated mainly in the US and Japanese strains (Ode and Strand 1995; Smith et al. 2017; Sakamoto et al. 2020). Like other hymenopteran insects (Cook 1993), *C. floridanum* is haplodiploid species in which the females develop from the fertilized eggs and the males develop from the unfertilized egg (Iwabuchi 2019). Soldier larvae develop from both male and female embryo, and the development and behavioral patterns of male soldiers differed between intraspecific strains. In the US strain, male soldiers appear in the penultimate larval stage of the host (Grbic et al. 1997), whereas in the Japanese strain, they appear in the first- or second-instar larvae of the host (Yamamoto et al. 2007). Male soldiers were functional in the Japanese strain but are not as aggressive as the female soldiers (Uka et al. 2013). Male soldiers in the US strain are nonaggressive (Giron et al. 2007). *vasa*,

encoding a DEAD-box RNA helicase protein (Hay et al. 1988; Lasko and Ashburner 1988), has been

shown to play a central role in caste determination of *C. floridanum* (Donnell et al. 2004; Vladimir

Zhurov et al. 2004). The *vasa* gene was originally identified in *Drosophila melanogaster* as a

maternal-effect factor for germ-cell specification and development of abdominal segments

(Schupbach and Wieschaus 1986), and they have been found in invertebrates and vertebrates (Raz

2000). *vasa* has been compared between the Japanese and the US strains of *C. floridanum*, and a 10

amino acid sequence difference in the N-terminal region was observed (Ohno et al. 2019).

Each strain should optimize its behavior and developmental patterns for its habitat. Genome

or transcriptome analyses should accelerate our understanding of the molecular basis underlying the

intraspecific differences of parasitic ability including polyembryony and caste system (Sakamoto et

al. 2020). The genome of the US strain has been sequenced using short-read sequencing (i5K

Consortium 2013), while that of the Japanese strain has not. Long-read sequencing offers several

advantages over short-read sequencing (Pollard et al. 2018). High-quality (high contiguity of the

assembly and completeness of gene sets) insect genomes have recently been achieved using long-read

sequencing (Hotaling et al. 2021). Single-molecule real-time (SMRT) sequencing developed by

Pacific Biosciences (PacBio) employ the DNA polymerase reaction from a single circular DNA

template (Eid et al. 2009), generating highly accurate long high-fidelity reads (HiFi reads, >10 kb,

>99%) (Rhoads and Au 2015; Ardui et al. 2018; Wenger et al. 2019). HiFi reads are more suitable

than other long- or short-read methods for identifying genes containing repetitive sequences

(Kawahara et al. 2022; Hotaling et al. 2023).

Comparisons among intraspecific genomes should be useful in understanding of the various

parasitic ability in *C. floridanum*. In this study, we aimed to construct a highly contiguous genome

assembly of the Japanese *C. floridanum* using PacBio long-read sequencing. We performed the quality

check, the gene prediction and functional annotation for the *de novo* assembly. Gene annotation and

amino acid sequences of *vasa* were compared to clarify the genomic differences between Japanese

and U.S. strains.

101   Materials and Methods

102   DNA extraction and genome sequencing

Genomic DNA was extracted from adult *C. floridanum* males using the Blood & Cell Culture DNA

Maxi Kit (Qiagen Co. Ltd., Valencia, CA, USA). The genetic system of *C. floridanum* and other

hymenopteran insects is haplodiploid; females develop from fertilized eggs (diploid), and males

develop from unfertilized eggs (haploid). In this study, the male adults were used. The male adults are

clones born from a single egg and are genetically nearly identical. The sequence library was prepared

using the SMRTbell Express Template Preparation Kit 2.0 or SMRTbell Prep Kit 3.0 (Pacific

Biosciences, Menlo Park, CA, USA). Libraries were sequenced using the Sequel II or IIe (Pacific

Biosciences).

112   *De novo* assembly and quality assessment

Subreads and HiFi reads generated from the PacBio sequencers were assembled using Flye v 2.9

(Kolmogorov et al. 2019) and Hifiasm v 0.16.1 using the '-primary' option (Cheng et al. 2021),

respectively. In the PacBio long-read system, polymerase reads are yielded by sequencing circularized

DNA. Then, the adaptors are removed from the polymerase reads, offering the subreads (refer to

https://www.pacb.com/technology/hifi-sequencing/). HiFi reads are defined as the consensus

sequence called from the subreads. Hifiasm v 0.16.1 can produce the primary contig and alternate

contig. The primary contig indicates the haplotype, and the alternate contig is composed of short

haplotype-specific contigs (haplotigs). Quast v 5.2.0 was used to assess the genome assembly statistics

(Gurevich et al. 2013). Gene set completeness was assessed using the Benchmark of Universal Single-

Copy Orthologs (BUSCOs) v 5.2.2, with '-l insecta_odb10 (2020-09-10)' (Manni et al. 2021). k-mer-

based validation was performed using Merqury v1.3 and Meryl db (k=21) (Rhie et al. 2020) for the

assembly constructed with HiFi reads. Contigs were assigned to taxonomic classifications using

BlobTools v 1.1(Laetsch and Blaxter 2017) with the BLAST (BLASTN) program in the National

Center for Biotechnology Information BLAST software package (v 2.14.0) against the NCBI non-

redundant nucleotide sequence database (nt), prepared using ncbi-blast-dbs v 10.5.0

(https://github.com/yeban/ncbi-blast-dbs, accessed on May 18, 2023). minimap2 v 2.26-r1175(Li

2018) was used to obtain mapping coverage.

Dot plots were visualized using the dotPlotly program with '-s -t -m 10000 -q 25000 -l'

(https://github.com/tpoorten/dotPlotly, accessed on July 17, 2023) to compare the genomic differences

between the US and Japanese strains. The 1 Mb or longer contigs were extracted using SeqKit v 2.0.0,

aligned by a nucmer with default settings, and delta files were filtered using a delta-filter with '-q -r'

options in MUMmer4 v4.0.0rc (Marçais et al. 2018).

Gene prediction and functional annotation

Gene prediction was performed using the BRAKER2 and BRAKER3 databases (Brůna et al. 2021;

Gabriel et al.). Public RNA sequencing (RNA-Seq) data (morula: DRR138914, DRR138915,

DRR138916, and DRR138917; male adult: SRR1864697) and OrthoDB protein sets of Arthropoda

(Kuznetsov et al. 2023) (https://bioinf.uni-greifswald.de/bioinf/partitioned_odb11/, accessed on June

17, 2023) were used as extrinsic evidence data to perform the BRAKER pipeline. The RNA-Seq reads

were processed using fastp v 0.23.2 with default settings (Chen 2023) and mapped to the *de novo*

assembled genome using HISAT2 v 2.2.1(Kim et al. 2019). The generated SAM files were converted

to BAM files using SAMtools v 1.12 (Danecek et al. 2021) and the BAM files were imported together

when the BRAKER pipeline was run. Repetitive assembly sequences were soft-masked using

RepeatModeler v 2.0.3 and RepeatMasker v 4.0.6 in Extensive de novo TE Annotator v 2.1.3 (Ou et

al. 2019). The soft-masked genome assembly was inputted into the BRAKER pipeline. GFFRead

v0.12.1 (Pertea and Pertea 2020) was used to convert the transcript or protein sequences from the GTF

file created by BRAKER. Protein sequences were assessed using BUSCO (protein mode) with '-l

insecta_odb10' (2020-09-10).

The functional annotation workflow of Fanflow4Insects (Bono et al. 2022a) was modified

and used for protein sequences, as described below. The protein sequences were aligned against

reference datasets of *Homo sapiens*, *Mus musculus*, *Caenorhabditis elegans*, *Drosophila*

melanogaster, *C. floridanum* (US strain), and UniProtKB/Swiss-Prot using GGSEARCH v 36.3.8g in

the FASTA package (https://fasta.bioch.virginia.edu/, accessed on July 19, 2023). Protein sequences

were also searched using HMMSCAN in HMMER v 3.3.2. Pfam v 35.0 was used as the protein

domain database for HMMSCAN (Mistry et al. 2021). Enrichment analysis was performed using

Metascape software (https://metascape.org/, accessed on July 19, 2023) with default settings (Zhou et

al. 2019).

Results and Discussion

***De novo* genome assembly of Japanese *C. floridanum* and the quality assessment**

This study generated HiFi reads and subreads using Japanese strains (Cflo-J). HiFi reads were

assembled, and the primary contig comprised 149 contigs with an N50 length of 17.9 Mb and a total

assembly size of 552,713,363 bp (50.8-fold coverage; Table 1 and Table S1). Hifiasm also produced

alternate contigs (1,162 contigs, Table S1). The genome assembly generated by subreads comprised

996 contigs with an N50 length of 4.6 Mb and a total assembly size of 534,638,157 bp (244.7-fold

coverage, Table 1 and Table S1). Cflo_2.0 (NCBI RefSeq assembly GCF_000648655.2) was used as

the genome assembly of US strain (Cflo-US). The assembly quality between the Japanese strain (Cflo-

J) and the US strain (Cflo-US) was evaluated. The N50 value of Cflo-J assemblies was much higher

than that of Cflo-US (1.2 M). However, the GC content of the assemblies was similar. The BUSCO

completeness of Cflo-J was also comparable to that of Cflo-US; however, the fragmentation score was

lower for the Cflo-J assemblies (Table 1). The N50 values and number of contigs were the best for the

HiFi assembly (Table 1 and Figure 1A). We confirmed the differences in genomic structures among

all the assemblies using dot-plot visualization (Figures 1B and C). Linear plots were obtained,

confirming the similarity of the genomic structures across the entire assembly length (Figure 1B and

C). The matching coordinate data for the dot-plot comparison are available in Files S1 and S2.

We then evaluated the HiFi assemblies of Cflo-J, which had the highest contiguity. K-mer-

based evaluation showed high base accuracy and completeness of primary contigs (a quality value

(QV) = 68.08 and completeness = 97.6%, Table S2). The Merqury spectrum plot showed a single k-

mer peak in the primary contigs, indicating extremely low heterozygosity (Figure S1A, red line).

Because *C. floridanum* males possess a haploid genome, the single k-mer peak of the primary contig

was reasonable. We also confirmed the presence of reads included only in the alternate contigs (Figure

S1A, blue line), suggesting that the genome assembly constructed in this study may contain

heterozygous regions.

Taxonomic assignments of the contigs were performed to reveal the taxonomic origin of the

contigs. The primary and alternate contigs were assigned to Arthropoda, and the remaining contigs

were either no-hit or chordata (Figures S1B and C). The mapping rate of no-hit was remarkably low

(primary contig, 0.27%; alternate contig, 0.51%), and that of chordata in alternate contigs was also

extremely low (0.05%). The assembly of Cflo-US also includes Arthropoda and no-hit contigs as

shown in Cflo-J (ENA https://www.ebi.ac.uk/ena/browser/view/GCA_000648655.2, accessed on

September 20, 2023). In conclusion, an ideal haploid representation was observed in the primary

contigs, and no contig contamination from non-arthropods was observed.

195   Gene annotation differences between Cflo-J and Cflo-US

As repetitive sequences may affect gene prediction accuracy (Hoff et al. 2019), these sequences in the

HiFi primary contig were investigated using RepeatModeler and RepeatMasker. The repeat sequence

was 285.4 Mb, corresponding to 51.65% of the HiFi primary contig, and "Unclassified" accounted for

a substantial percentage of the total (34.97%) (Table S3). Gene prediction was performed using several

RNA-Seq data and protein dataset combinations. The BRAKER3 using RNA-Seq data and the

Arthropoda protein dataset showed the best BUSCO completeness (96.4%, Table 2). The numbers of

predicted genes and transcripts are listed in Table 2. BRAKER2 predicted more genes compared to

BRAKER3 (Table 2). The gene model with the highest BUSCO completeness (96.4%) ("best gene

model") predicted 10,786 protein-coding genes and 13,886 transcripts. The annotated GFF files are

shown in File S3. The best gene model was annotated using Fanflow4Insects (Bono et al. 2022b).

Table 3 shows the results of the searches using Fanflow4Insects for each reference dataset. Of the

10786 gene models, 10585 were annotated using GGSEARCH or HMMSCAN. The full annotation

results for GGSEARCH and HMMSCAN are presented in Table S4.

We evaluated whether annotation differences existed between the Cflo-US assembly and

Cflo-J HiFi assembly. We identified 149 transcripts in 115 gene loci with no corresponding proteins

to Cflo-US and with corresponding proteins to human, mouse, *C.elegans*, and fly (Table S5). The

human gene IDs corresponding to 149 transcripts were used for enrichment analysis (Figure 2A).

"Metabolism of carbohydrates" was the most significantly enriched Gene Ontology (GO) term. In

addition, GO terms for amino acid metabolism and translation were enriched.

Excluding gene loci with at least one isoform corresponding to Cflo-US genes, the number

of genes as mentioned above was limited from 115 to 100. We investigated why these 100 genes did

not hit the Cflo-US gene model. To verify whether corresponding regions of the 100 genes are absent

in the Cflo-US, TBLASTN of 100 genes was performed for the genome of Cflo-US. We manually

examined hit regions of TBLASTN using NCBI Genome Data viewer. TBLASTN results showed that

all 100 genes hit in the Cflo-US genome. Thus, the difference in genome annotation between Cflo-US

and Cflo-J is not due to differences in the extent of the sequenced genomic regions. Among 100 genes,

21 genes had no corresponding gene models in Cflo-US assembly, while 79 genes had corresponding

gene models in Cflo-US (Table S6). The gene model g524 was representative example without

corresponding gene model of Cflo-US (Figure 2B). On both sides of g524 are g523 and g525. Amino

acid sequences of g523 and g525 have sequence similarity to XP_014209029.1 and XP_014209034.1

in Cflo-US genome, respectively (Table S4). Although these genomic position and sequence similarity

indicate that the genomic position around g524 is conserved in both genome assembly, the gene model

corresponding to g524 was absent in Cflo-US.

Among 79 genes with corresponding gene model in Cflo-US, TBLASTN results were

divided into two groups: genes that hit the single gene model or genes that hit more than two gene

model (Table S6). In addition, differences of the amino acid length between Cflo-J and Cflo-US were

also revealed by the TBLASTN (Table S6). g490 was representative example as the gene that hit the

single gene model of Cflo-US. TBLASTN showed that the amino acid sequence of g490.t1 had

similarity to XP_014209022.1 of Cflo-US gene model (Table S6). The exon structures of

XP_014209022.1 were different from g490 of Cflo-J (Figure 2C). Multiple sequence alignment

showed that the amino acid length of XP_014209022.1 of Cflo-US was extremely longer at N-terminal

region than the corresponding protein of other species (Figure S2). When BLASTP was adopted,

g490.t1 showed high sequence similarity to XP_014209022.1 (E-value: 0.0), despite the gene

structures were largely different. This indicates that GGSEARCH adopted in this study can detect the

genes with extremely different structure of genes. g909 was the representative example as the gene

that hit several gene model of Cflo-US (Table S6). TBLASTN showed that g909 had the similarity

XP_023247179.1 and XP_014214160.1, and these two genes of Cflo-US were distributed in different

scaffolds (Figure 2D). Multiple sequence alignment indicated that XP_023247179.1 and

XP_014214160.1 correspond to the first and second half of the amino acid sequence of g909.t1,

respectively (Figure S3). The example of g909 suggested that long-read sequencing contributed to the

accuracy of genome annotation. TBLASTN also detected the genes that hit more than two gene model

on the same scaffold (Table S6, e.g. g1061.t1).

In this study, we found genes with extremely different gene annotations between Cflo-US

and Cflo-J. This success is attributed to the method used in this study. GGSEARCH, which was

adopted in this study, searches for sequence similarity with the subject within 80-120% of the query

length. This contributed to discovery of protein structure differences between Cflo-J and Cflo-US. In

addition, highly contiguous genome assembly obtained by long-read sequencing suggested the

fragmentation of gene model in Cflo-US genome. All the differences in genome annotation revealed

in this study are noteworthy as the discoveries of genomic differences between Cflo-J and Cflo-US.

Common features and differences of *vasa* between Cflo-J and Cflo-US

Ohno et al. (2019) showed that the amino acid sequence of *vasa* differed between Cflo-J and

Cflo-US. The amino acid sequences of Cflo-US (XP_014219851.1) and Cflo-J *vasa* (g8014.t1),

including the amino acid sequence (BBI30140.1) determined by Ohno et al. (2019), were compared.

We confirmed the differences in amino acid sequences at the N-terminal region between Cflo-US *vasa*

(XP_014219851.1) and Cflo-J *vasa* (g8014.t1), as previously reported[13] (Figure 3A, red arrows).

Within the Japanese strain (BBI30140.1 vs. g8014.t1), the amino acid sequences were almost identical

(Figure 3A).

We also investigated the genomic position of *vasa* (Figure 3B). *vasa* was located in an intron

between the seventh and eighth exons of *cap'n'collar* (*cnc*), and the directions of transcription were

opposite (Figure 2C, blue and green arrows). This positional relationship was also confirmed in Cflo-

US (NCBI Genome Browser: LOC106647823, Location: scaffold6|NW_019379589.1:2,626,762 –

2,630,760.). Generally, overlapping genes such as *vasa* and *cnc* are negatively selected evolutionarily,

whereas overlapping genes are sometimes retained in a species-specific manner (Craig et al. 1997). In

addition, overlapping genes are likely to be co-expressed. Furthermore, the *vasa* genomic positions in

other insects were preliminarily examined (Table S7), and the positional relationship between *vasa*

and *cnc* may be conserved in Hymenoptera. Genomic positional relationship of *vasa* and *cnc* may be

related to the oogenesis or embryogenesis in hymenopteran insects. *cnc* is related to the nuclei

arrangement of oocyte in *Drosophila melanogaster* (Guichet et al. 2001), suggesting its involvement

in egg development in *C. floridanum*. The evolution to control the caste development by *vasa* in

Copidosoma may have required a conserved positional relationship between *vasa* and *cnc* in the

Hymenoptera. However, further verifications for relationship between *vasa* and *cnc* are needed.

Conclusion

We obtained a highly contiguous genome assembly of Japanese *C. floridanum*. We showed that long-

read sequencing and functional annotation method adopted in this study can highlight the differences

of genome annotation between Cflo-J and Cflo-US. These regions may be due to the influence of the

annotation and assembly accuracy, but in any case, they are remarkable genes that reflect differences

among the strains. Parasitic wasps can control the physiological state of host insects, and their use

for pest control is expected (Soldà et al. 2008). Comparison of the genome sequences of Cflo-US and

Cflo-J will contribute to the identification of detailed genomic regions relevant to the ecological and

physiological characteristics of *C. flolidanum*, leading to a detail understanding of its parasitic ability

against pests.

Data availability

All sequencing data (assembled sequences and raw sequence reads) were deposited in the DDBJ under

the umbrella BioProject accession number PRJDB16642. Genome assembly from primary contigs was

deposited in the DDBJ under the accession numbers BTPR01000001−BTPR01000149. Alternate

contigs were deposited in DDBJ under the accession numbers BTPQ01000001−BTPQ01001162. Raw

sequence reads were deposited in the DDBJ under accession number DRA017002 (PacBio HiFi reads)

and DRA017125 (PacBio Subreads). Gene annotation and functional annotation of protein-coding

genes are available at figshare (https://doi.org/10.6084/m9.figshare.24145182.v5). Supplementary

Files are available at figshare (https://doi.org/10.6084/m9.figshare.24145182.v5). Scripts of

genome   assembly,   quality   check,   and   gene   annotation   were   also   available

(https://doi.org/10.6084/m9.figshare.24145182.v5)

Acknowledgments

We would like to thank all laboratory members at Hiroshima University for their valuable comments.

Computations were performed on the computers at the Hiroshima University Genome Editing

Innovation Center. Computations were partially performed on the NIG supercomputer at ROIS

National Institute of Genetics.

Authors' contributions

Conceptualization: H.T. and H.B.; methodology: Kouhei T. and H.B.; software: Kouhei T. Formal

analysis, data curation, and visualization: Kouhei T., Keita T., and T.S. Validation: Kouhei T., and

H.B. Investigation: Kouhei T., and H.B. Resources: Kouhei T., H.B., and H.T. Writing—original draft:

Kouhei T. Writing—review and editing: T.S., M.K, Keita T., K.O., H.T., and H.B. Supervision: H.T.,

and H.B. Project administration: H.B. Funding acquisition: H.T., and H.B.

Conflict of interest

The authors declare no conflict of interest.

Funding

This study was supported by the Center of Innovation for Bio-Digital Transformation (BioDX), an

open innovation platform for industry-academia co-creation of JST (COI-NEXT, grant number

JPMJPF2010 to H.T. and H.B.), and JSPS KAKENHI (grant number 23K17418 to H.T. and H.B.).

References

Ardui S, Ameur A, Vermeesch JR, Hestand MS. 2018. Single molecule real-time (SMRT) sequencing
comes of age: applications and utilities for medical diagnostics. Nucleic Acids Res. 46(5):2159–2168.
doi:10.1093/nar/gky066.

Bono H, Sakamoto T, Kasukawa T, Tabunoki H. 2022a. Systematic Functional Annotation Workflow
for Insects. Insects. 13(7):586. doi:10.3390/insects13070586.

Bono H, Sakamoto T, Kasukawa T, Tabunoki H. 2022b. Systematic Functional Annotation Workflow
for Insects. Insects. 13(7):586. doi:10.3390/insects13070586.

Brůna T, Hoff KJ, Lomsadze A, Stanke M, Borodovsky M. 2021. BRAKER2: automatic eukaryotic
genome annotation with GeneMark-EP+ and AUGUSTUS supported by a protein database. NAR
Genom Bioinform. 3(1). doi:10.1093/nargab/lqaa108.

Chen S. 2023. Ultrafast one-pass FASTQ data preprocessing, quality control, and deduplication using
fastp. iMeta. 2(2). doi:10.1002/imt2.107.
340

341   Cheng H, Concepcion GT, Feng X, Zhang H, Li H. 2021. Haplotype-resolved de novo assembly using
342   phased assembly graphs with hifiasm. Nat Methods. 18(2):170–175. doi:10.1038/s41592-020-01056-
343   5.
344

345   Cook JM. 1993. Sex determination in the Hymenoptera: a review of models and evidence.
346

347   Craig SF, Slobodkin LB, Wray GA, Biermann CH. 1997. The 'paradox' of polyembryony: A review
348   of the cases and a hypothesis for its evolution. Evol Ecol. 11(2):127–143.
349   doi:10.1023/A:1018443714917.
350

351   Cruz YP. 1981. A sterile defender morph in a polyembryonic hymenopterous parasite. Nature.
352   294(5840):446–447. doi:10.1038/294446a0.
353

354   Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A, Keane T, McCarthy
355   SA, Davies RM, et al. 2021. Twelve years of SAMtools and BCFtools. Gigascience. 10(2).
356   doi:10.1093/gigascience/giab008.
357

358   Donnell DM, Corley LS, Chen G, Strand MR. 2004. Caste determination in a polyembryonic wasp
359   involves inheritance of germ cells. Proc Natl Acad Sci U S A. 101(27):10095–10100.
360   doi:10.1073/pnas.0403625101.
361

362   Eid J, Fehr A, Gray J, Luong K, Lyle J, Otto G, Peluso P, Rank D, Baybayan P, Bettman B, et al. 2009.
363   Real-Time DNA Sequencing from Single Polymerase Molecules. Science (1979). 323(5910):133–138.
364   doi:10.1126/science.1162986.
365

366   Gabriel L, Brůna T, Hoff KJ, Ebel M, Lomsadze A, Borodovsky M, Stanke M. BRAKER3: Fully
367   Automated Genome Annotation Using RNA-Seq and Protein Evidence with GeneMark-ETP,
368   AUGUSTUS and TSEBRA. doi:10.1101/2023.06.10.544449.
369   https://doi.org/10.1101/2023.06.10.544449.
370

Giron D, Harvey JA, Johnson JA, Strand MR. 2007. Male soldier caste larvae are non-aggressive in
the polyembryonic wasp *Copidosoma floridanum*. Biol Lett. 3(4):431–434.
doi:10.1098/rsbl.2007.0199.

Grbic M, Rivers D, Strand MR. 1997. Caste formation in the polyembryonic wasp Copidosoma
floridanum (Hymenoptera: Encyrtidae): in vivo and in vitro analysis. J Insect Physiol. 43(6):553–565.
doi:10.1016/S0022-1910(97)00004-8.

Guerrieri E, Noyes J. 2005. Revision of the European species of Copidosoma Ratzeburg
(Hymenoptera: Encyrtidae), parasitoids of caterpillars (Lepidoptera). Syst Entomol. 30(1):97–174.
doi:10.1111/j.1365-3113.2005.00271.x.

Guichet A, Peri F, Roth S. 2001. Stable anterior anchoring of the oocyte nucleus is required to establish
dorsoventral polarity of the drosophila egg. Dev Biol. 237(1):93–106. doi:10.1006/dbio.2001.0354.

Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUAST: Quality assessment tool for genome
assemblies. Bioinformatics. 29(8):1072–1075. doi:10.1093/bioinformatics/btt086.

Hay B, Yeh Jan L, Nung Jan Y. 1988. A Protein Component of Drosophila Polar Granules Is Encoded
by vasa and Has Extensive Sequence Similarity to ATP-Dependent Helicases.

Hoff KJ, Lomsadze A, Borodovsky M, Stanke M. 2019. Whole-Genome Annotation with BRAKER.
Methods Mol Biol. 1962:65–95. doi:10.1007/978-1-4939-9173-0_5.

Hotaling S, Sproul JS, Heckenhauer J, Powell A, Larracuente AM, Pauls SU, Kelley JL, Frandsen PB.
2021. Long Reads Are Revolutionizing 20 Years of Insect Genome Sequencing. Genome Biol Evol.
13(8). doi:10.1093/gbe/evab138.

Hotaling S, Wilcox ER, Heckenhauer J, Stewart RJ, Frandsen PB. 2023. Highly accurate long reads
are crucial for realizing the potential of biodiversity genomics. BMC Genomics. 24(1):117.
doi:10.1186/s12864-023-09193-9.

i5K Consortium. 2013. The i5K Initiative: advancing arthropod genomics for knowledge, human
health, agriculture, and the environment. J Hered. 104(5):595–600. doi:10.1093/jhered/est050.

Iwabuchi K. 2019. Polyembryonic Insects. Singapore: Springer Singapore.

Kawahara AY, Storer CG, Markee A, Heckenhauer J, Powell A, Plotkin D, Hotaling S, Cleland TP,
Dikow RB, Dikow T, et al. 2022. Long-read HiFi sequencing correctly assembles repetitive heavy

fibroin silk genes in new moth and caddisfly genomes. GigaByte. 2022:1–14.
doi:10.46471/gigabyte.64.
412

413   Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. 2019. Graph-based genome alignment and
414   genotyping with HISAT2 and HISAT-genotype. Nat Biotechnol. 37(8):907–915. doi:10.1038/s41587-
415   019-0201-4.
416

417   Kolmogorov M, Yuan J, Lin Y, Pevzner PA. 2019. Assembly of long, error-prone reads using repeat
418   graphs. Nat Biotechnol. 37(5):540–546. doi:10.1038/s41587-019-0072-8.
419

420   Kuznetsov D, Tegenfeldt F, Manni M, Seppey M, Berkeley M, Kriventseva EV, Zdobnov EM. 2023.
421   OrthoDB v11: annotation of orthologs in the widest sampling of organismal diversity. Nucleic Acids
422   Res. 51(D1): D445–D451. doi:10.1093/nar/gkac998.
423

424   Laetsch DR, Blaxter ML. 2017. BlobTools: Interrogation of genome assemblies. F1000Res. 6:1287.
425   doi:10.12688/f1000research.12232.1.
426

427   Lasko PF, Ashburner M. 1988. The product of the Drosophila gene vasa is very similar to eukaryotic
428   initiation factor-4A. Oxford University Press.
429

430   Li H. 2018. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics. 34(18):3094–
431   3100. doi:10.1093/bioinformatics/bty191.
432

433   Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. 2021. BUSCO Update: Novel and
434   Streamlined Workflows along with Broader and Deeper Phylogenetic Coverage for Scoring of
435   Eukaryotic, Prokaryotic, and Viral Genomes. Mol Biol Evol. 38(10):4647–4654.
436   doi:10.1093/molbev/msab199.
437

438   Marçais G, Delcher AL, Phillippy AM, Coston R, Salzberg SL, Zimin A. 2018. MUMmer4: A fast
439   and versatile genome alignment system. PLoS Comput Biol. 14(1): e1005944. doi:
440   10.1371/journal.pcbi.1005944.
441

442   Mistry J, Chuguransky S, Williams L, Qureshi M, Salazar GA, Sonnhammer ELL, Tosatto SCE,
443   Paladin L, Raj S, Richardson LJ, et al. 2021. Pfam: The protein families database in 2021. Nucleic
444   Acids Res. 49(D1): D412–D419. doi:10.1093/nar/gkaa913.
445

446   Ode PJ, Strand MR. 1995. Progeny and Sex Allocation Decisions of the Polyembryonic Wasp
447   Copidosoma floridanum.
448

449   Ohno H, Sakamoto T, Okochi R, Nishiko M, Sasaki S, Bono H, Tabunoki H, Iwabuchi K. 2019.
450   Apoptosis-mediated vasa down-regulation controls developmental transformation in Japanese
451   Copidosoma floridanum female soldiers. Dev Biol. 456(2):226–233. doi:
452   10.1016/j.ydbio.2019.09.005.
453

454   Ou S, Su W, Liao Y, Chougule K, Agda JRA, Hellinga AJ, Lugo CSB, Elliott TA, Ware D, Peterson
455   T, et al. 2019. Benchmarking transposable element annotation methods for creation of a streamlined,
456   comprehensive pipeline. Genome Biol. 20(1):275. doi:10.1186/s13059-019-1905-y.
457

458   Pertea G, Pertea M. 2020. GFF Utilities: GffRead and GffCompare. F1000Res. 9:304.
459   doi:10.12688/f1000research.23297.2.
460

461   Pollard MO, Gurdasani D, Mentzer AJ, Porter T, Sandhu MS. 2018. Long reads: their purpose and
462   place. Hum Mol Genet. 27(R2): R234–R241. doi:10.1093/hmg/ddy177.
463

464   Raz E. 2000. The function and regulation of vasa-like genes in germ-cell development. Genome Biol.
465   1(3): REVIEWS1017. doi:10.1186/gb-2000-1-3-reviews1017.
466

467   Rhie A, Walenz BP, Koren S, Phillippy AM. 2020. Merqury: reference-free quality, completeness,
468   and phasing assessment for genome assemblies. Genome Biol. 21(1):245. doi:10.1186/s13059-020-
469   02134-9.
470

471   Rhoads A, Au KF. 2015. PacBio Sequencing and Its Applications. Genomics Proteomics
472   Bioinformatics. 13(5):278–289. doi: 10.1016/j.gpb.2015.08.002.
473

474   Sakamoto T, Nishiko M, Bono H, Nakazato T, Yoshimura J, Tabunoki H, Iwabuchi K. 2020. Analysis
475   of molecular mechanism for acceleration of polyembryony using gene functional annotation pipeline
476   in Copidosoma floridanum. BMC Genomics. 21(1):152. doi:10.1186/s12864-020-6559-3.
477   https://bmcgenomics.biomedcentral.com/articles/10.1186/s12864-020-6559-3.
478

Schupbach T, Wieschaus E. 1986. Maternal-effect mutations altering the anterior-posterior pattern of
the Drosophila embryo. Roux's Archives of Developmental Biology. 195(5):302–317.
doi:10.1007/BF00376063.

Segoli M, Harari AR, Rosenheim JA, Bouskila A, Keasar T. 2010. The evolution of polyembryony in
parasitoid wasps. J Evol Biol. 23(9):1807–1819. doi:10.1111/j.1420-9101.2010.02049.x.

Smith MS, Shirley A, Strand MR. 2017. Copidosoma floridanum (Hymenoptera: Encyrtidae) Rapidly
Alters Production of Soldier Embryos in Response to Competition. Ann Entomol Soc Am.
110(5):501–505. doi:10.1093/aesa/sax056.

Soldà G, Suyama M, Pelucchi P, Boi S, Guffanti A, Rizzi E, Bork P, Tenchini ML, Ciccarelli FD.
2008. Non-random retention of protein-coding overlapping genes in Metazoa. BMC Genomics. 9.
doi:10.1186/1471-2164-9-174.

Strand MR. 2003. Polyembryony. Carde R., Resch V., editors. San Diego: Academic Press.

Szathmáry E, Smith JM. 1995. The major evolutionary transitions. Nature. 374(6519):227–232.
doi:10.1038/374227a0.

Uka D, Takahashi-Nakaguchi A, Yoshimura J, Iwabuchi K. 2013. Male soldiers are functional in the
Japanese strain of a polyembryonic wasp. Sci Rep. 3(1):2312. doi:10.1038/srep02312.

Vladimir Zhurov, Tomislav Terzin, Miodrag Grbic. 2004. Early blastomere determines embryo
proliferation and caste fate in a polyembryonic wasp. Nature. 432(7018):761–764.
doi:10.1038/nature03030.

Whyte BA. 2021. The weird eusociality of polyembryonic parasites. Biol Lett. 17(4).
doi:10.1098/rsbl.2021.0026.

Yamamoto D, Henderson R, Corley LS, Iwabuchi K. 2007. Intrinsic, inter-specific competition
between egg, egg–larval, and larval parasitoids of plusiine loopers. Ecol Entomol. 32(2):221–228.
doi:10.1111/j.1365-2311.2006.00857.x.

Zhou Y, Zhou B, Pache L, Chang M, Khodabakhshi AH, Tanaseichuk O, Benner C, Chanda SK. 2019.
Metascape provides a biologist-oriented resource for the analysis of systems-level datasets. Nat
Commun. 10(1):1523. doi:10.1038/s41467-019-09234-6. http://www.nature.com/articles/s41467-
019-09234-6.

Tables

Table 1 Comparison of assembly statistics with the current reference genomes

	parameter	Cflo-J HiFi (primary)	Cflo-J subread	Cflo-US (Cflo2.0)
assembly statistics	Length	552,713,363	534,638,157	503,815,983
	N50	17,879,143	4,591,336	1,210,516
	L50	9	29	130
	No. of contigs	149	996	4,840
	Largest contig	42,954,611	26,880,362	8,149,722
	GC %	35.62	35.78	35.50
BUSCO results (OrthoDB v10 Insecta database)	Complete (single, duplicated)	97.9 % (96.4%, 1.5%)	98.1% (96.9%,1.2%)	96.7% (95.0%,1.7%)
	Fragmented	0.6%	0.7%	1.5%
	Missing	1.5%	1.2%	1.8%

Table 2 BUSCO analysis for the protein set predicted by BRAKER

	BRAKER3 with RNA seq + Arthropoda protein	BRAKER2 with RNA seq	BRAKER3 with Arthropoda protein	BRAKER2 with Arthropoda protein
Complete (single, duplicated)	96.4 % (79.5%, 16.9%)	94.0% (73.9%, 20.1%)	93.1% (78.8%, 14.3%)	92.5% (82.8%, 9.7%)
Fragmented	0.5%	2.4%	0.7%	1.8%
Missing	3.1%	3.6%	6.2%	5.7%
No. of genes	10786	18531	9913	22781
No. of transcripts	13886	19986	12370	24487

Table 3 Summary of functional annotation results

Annotation category	Reference	The number of gene count
Gene annotation by top hit (GGSEARCH)	Human	8467
	Mouse	8291

	C. elegans	7061
	fly	8403
	Copidosoma floridanum	9850
	UniProtKB/Swiss-Prot	9038
	At least one of the above	10537
Genes annotated only by HMMSCAN	Pfam domain database	48
All genes annotated at protein level	At least one of the above	10585
Hypothetical protein	No hit	201
Total		10786

Figures

Figure 1 Contiguity and sequence similarity of each assembly. (A) Each color shows the cumulative

base pair length and number of contigs for each assembly. The assemblies generated from HiFi reads

contain the fewest contigs. (B, C) Dot-plot between the Japanese *C. floridanum* strain (Cflo-J) and US

C. floridanum (Cflo-US) high-fidelity (HiFi) assemblies and between Cflo-J HiFi and Cflo-J subread

assemblies. The vertical axis indicates Cflo-US or Cflo-J subread assemblies. The horizontal axis

indicates Cflo-J HiFi assembly. The average identity percentages (per query) are displayed in different

colors.

Figure 2

Features of genes with no corresponding proteins to Cflo-US. (A) The results of enrichment analysis

(Metascape) of 149 transcripts with no corresponding proteins to Cflo-US and with corresponding

proteins to human, mouse, *C. elegans*, and fly. Bars are colored differently according to the value of -

log10 (P-value). (B-D) Representative gene model illustrating the genome annotation differences.

Yellow and blue boxes indicate the ORF and UTR, respectively. Red lines indicate the region with

similar exon structures.

A

B

Figure 3 Amino acid sequences genomic position and of *vasa* gene. Comparison of gene annotation

between the Cflo-US and Cflo-J HiFi assemblies. (A) Comparison of amino acid sequences of the

vasa gene in the US (XP_014219851.1) and Japanese strains (g8014.t1 and BBI30140.1). g8014.t1 is

the transcript ID of *vasa* predicted in Cflo-J assembly, and BBI30140.1 is the protein ID of *vasa* in the

Japanese lineage reported by Ohno et al. (2019). Red arrows indicate amino acids that differ between

the US and Japanese strains, as Ohno et al. 2019 revealed. (B) Genomic position of *vasa* with RNA-

551   Seq read coverages. The CDS regions are shown at the bottom, including the *vasa* and *cap'n'collar*

552   (*cnc*) exon structures. Blue and green arrows indicate the direction of transcription.

553