Morphometrics of the preserved post-surgical hemisphere in pediatric drug-resistant epilepsy

Michael C. Granovetter, PhD,[1,2†] Anne Margarette S. Maallo, PhD,[1,†]

Christina Patterson, MD,[3] Daniel Glen, BS,[4] and Marlene Behrmann, PhD[1,5]

[†]These authors contributed equally to this work.

Author affiliations:

1 Department of Psychology and Neuroscience Institute, Carnegie Mellon University, Pittsburgh, PA, USA 15213

2 School of Medicine, University of Pittsburgh, Pittsburgh, PA, USA 15213

3 Department of Pediatrics, University of Pittsburgh, Pittsburgh, PA, USA 15213

4 Scientific and Statistical Computing Core, National Institute of Mental Health, Bethesda, MD, USA 20892

5 Department of Ophthalmology, University of Pittsburgh, Pittsburgh, PA, USA 15213

Correspondence to: Marlene Behrmann, PhD

Department of Ophthalmology, University of Pittsburgh, Pittsburgh, PA, USA 15213

mbehrmann@pitt.edu

Word Count: 2981

Key Points

Question: Following large pediatric cortical resection for the treatment of drug-resistant epilepsy, how does the structural integrity of the preserved hemisphere differ from that of healthy children?

Findings: In this case-control study of 32 patients with childhood epilepsy surgery, left-sided—but not right-sided—resection cases showed significantly reduced cortical volume and thickness and increased surface area relative to 51 non-neurological matched controls.

Meaning: There is reorganization of the right hemisphere that is specific to left hemispheric resection but not vice versa; this is potentially a consequence of unique left-lateralized functions (e.g., language) requiring postoperative remapping to the right hemisphere.

Abstract

Importance: Structural integrity of cortex following cortical resection for epilepsy management has been previously characterized, but only in adult patients.

Objective: This study sought to determine whether morphometrics of the preserved hemisphere in pediatric cortical resection patients differ from non-neurological controls.

Design: This was a case-control study, from 2013-2022.

Setting: This was a single-site study.

Participants: 32 patients with childhood epilepsy surgery and 51 age- and gender-matched controls participated.

Main Measure(s): We quantified morphometrics of the preserved hemisphere at the level of gross anatomy (lateral ventricle size, volume of gray and white matter). Additionally, cortical thickness, volume, and surface area were measured for 34 cortical regions segmented with the Desikan-Killiany atlas, and, last, volumes of nine subcortical regions were also quantified.

Results: 13 patients with left hemisphere (LH) surgery and a preserved right hemisphere (RH) (median age/median absolute deviation of age: 15.7/1.7 yr; 6 females, 7 males) and 19 patients with RH surgery and a preserved LH (15.4/3.7 yr; 11 females, 8 males) were compared to 51 controls (14.8/4.9 yr; 24 females, 27 males). Patient groups had larger ventricles and reduced total white matter volume relative to controls, and only patients with a preserved RH, but not patients with a preserved LH, had reduced total gray matter volume relative to controls. Furthermore, patients with a preserved RH had lower cortical thickness and volume and greater surface area of several cortical regions, relative to controls. Patients with a preserved LH had no differences in thickness, volume, or area, of any of the 34 cortical regions, relative to controls. Moreover, both LH and RH patients showed reduced volumes in select subcortical structures, relative to controls.

Conclusions and Relevance: That left-sided, but not right-sided, resection is associated with more pronounced reduction in cortical thickness and volume and increased cortical surface area relative to typically developing, age-matched controls suggests that the preserved RH undergoes structural plasticity to an extent not observed in cases of right-sided pediatric resection. Future work probing the association of the current findings with neuropsychological outcomes will be necessary to understand the implications of these structural findings for clinical practice.

Introduction

Surgical resection effectively reduces seizures in drug-resistant epilepsy (DRE).[1] Whether there are postoperative changes in the structural integrity of non-resected cortex is unclear. Post-surgically, reduction in cortical thickness (CxT), a proxy for structural integrity,[2] has been identified beyond the epileptogenic zone/lobe, even in the preserved hemisphere. In contrast, in patients with temporal lobe epilepsy, pre-surgical CxT reduction can be reversed to within normal limits, and the extent of reversal is correlated with better seizure outcomes.[3-8] Additionally, reversed cortical volume (CV) loss has been observed postsurgically, albeit only in patients with anterior temporal lobectomy,[9] but postsurgical deceleration of cortical atrophy has also been reported.[3, 5, 10, 11] The discrepant postoperative atrophy versus recovery warrants further study. If the surgical outcome is of progressive—rather than reduced—atrophy, alternative or additional disease-modifying treatments might be needed to halt progression.

Here, we compared postsurgical morphometry of the preserved hemisphere of children with DRE and non-neurological controls on (i) gross volumes of the lateral ventricles (LV), gray matter (GM), and white matter (WM); (ii) CxT, CV, and cortical surface area (CSA) of 34 regions; and (iii) volume of nine subcortical structures. CxT, CV, and CSA are commonly utilized in studying DRE[12] and have distinct genetic profiles and lifespan trajectories.[13-15] CxT linearly decreases with age;[16] CSA and CV follow a curvilinear trajectory with CSA peaking before CV.[14] Complementary information gleaned from these measures can offer insights into structural perturbations. Lastly, although postsurgical reduction in cortical thinning affects both hemispheres equally in adults,[3] here, we also determined whether, in children, postsurgical structural differences depended on which hemisphere is preserved. We focused analyses on the preserved hemisphere, as ipsilesional cortex may have structural abnormalities from surgery itself, confounding postoperative structural plasticity that potentially occurs post-resection.

Materials and methods

Participants

Thirty-two pediatric cortical resection or ablation patients participated: 13 with LH surgery and a preserved RH (median age/median absolute deviation of age: 15.7/1.7 yr; 6 females, 7 males) and 19 with RH surgery and a preserved LH (median age/median absolute deviation of age: 15.4/3.7 yr; 11 females, 8 males; see eTable 1 for details). "RH patients" are those with left-sided resections but a preserved RH, and likewise "LH patients" are those with a

preserved LH, with the labels indicating the hemisphere from which the measurement is taken.

Patient groups were matched on age ($F(1,30)=1.12$; $p=.30$), gender ($z(30)=0.65$; $p=.51$), initial surgery age ($F(1,30)=0.03$; $p=.87$), seizure-onset age ($F(1,29)=0.01$; $p=0.93$), and binned International League Against Epilepsy (ILAE) outcome scales[17] ($z(30)=1.29$; $p=.20$). Patients were recruited from Children's Hospital of Pittsburgh or via the Pediatric Epilepsy Surgery Alliance. Patients were matched with 51 control participants (median age/median deviation of age: 14.8/4.9 yr; 24 females, 27 males) on age ($F(1,81)=1.18$; $p=.28$) and gender ($z(81)=0.54$; $p=.59$).

Carnegie Mellon University and University of Pittsburgh Institutional Review Boards approved the study.

Protocol

T1-weighted images were acquired using a magnetization-prepared rapid acquisition gradient echo sequence (1-mm isotropic resolution, TE=1.97 ms, TR=2300 ms, scan time≅5 min) with either a 3T Siemens Verio scanner (32-channel head coil; 36 controls, 13 patients) or Siemens Prisma 3T scanner (64-channel head coil; 15 controls, 19 patients).

Outcome Measures

Images were motion-corrected, normalized, and segmented with FreeSurfer (v7.1.0)[18-20] before manual inspection. As FreeSurfer often does not accurately segment hemispherectomy/hemispherotomy/hemidecortication brains, for these cases, the intact hemisphere was mirrored using affine/non-linear transformations robust to aberrations, and only the preserved hemisphere was analysed.[21, 22]

Morphometric measures from the preserved hemisphere of patients and each hemisphere of controls were as follows: (i) total GM, WM, and LV volumes; (ii) cortical morphometry for 34 regions, parcellated according to the Desikan-Killiany atlas;[23, 24] and (iii) nine subcortical structures' volumes.[25, 26] We did not normalize CxT (as in Westman et al.[27]), but did so for CSA by dividing by the mean CSA of all of regions in that hemisphere, and for CV as the percentage of total hemisphere volume (sum of GM, WM, and LV volumes).[28-30]

Analysis

Data were analyzed in R[31] (for packages, see eTable 2) and SPSS 29.0.1.0. To harmonize data across the two scanners, data were modelled as a linear combination of group, age, gender, and scanner, assuming scanner effects have both additive and multiplicative factors.[32, 33] For each measure and ROI, data were winsorized: values above the 95th and

below the 5th percentile of the distribution were replaced with approximate corresponding percentiles (separately for controls and patients, and by hemisphere).

Given the small sample sizes from this relatively rare patient population, for each comparison, permutation testing was implemented by randomly shuffling the group label, and a general linear model (GLM) was fit with group as the primary predictor and age and gender as covariates. This was repeated 1,000 times to create a distribution of β-coefficients for the group effect. p-value was the percentage of occurrences in which the absolute value of the β-coefficient from the simulated distribution exceeded the absolute value of the true β-coefficient (α=.05). Per measure, Benjamini-Hochberg correction[34] was applied to p-values across ROIs. Bayes factor (BF) was also computed by comparing the model with the group term to a null model without the group term.[35, 36]

To elucidate which measures, alone or in combination, predicted group membership for each patient group and the matched hemisphere of the controls and between LH and RH patient groups, we conducted forward binary logistic regression analyses with $p<.05$ for entry criterion. We also predicted ILAE outcome (high versus low) for the patients using this multivariate approach. r^2 of models are provided using the Nagelkerke measure, and no other factors are entered when the model change was less than .001.

Results

<u>Gross Morphometrics</u>

Relative to controls' LH, patients' LH had larger LV volume (p=.02, BF=0.57), smaller WM volume (p=.05, BF=1.02), and no GM volume difference (p=.27, BF=4.37). Relative to controls' RH, patients' RH had larger LV volume ($p<.01$, BF=0.01), smaller WM volume ($p<.01$, BF=$5.56*10^{-2}$), and smaller GM volume ($p<0.001$, BF=$1.6*10^{-3}$). LH and RH patients did not differ on LV volume (p=.55, BF=4.54) or WM volume (p=.13, BF=1.39). However, RH patients had smaller GM volume than LH patients ($p<.01$, BF=$1.98*10^{-2}$). See Figure 1 for data and eTable 3 for statistics.

Evaluating whether LV, GM, and WM, singly or in combination predicted group membership, LH patients and LH controls were undifferentiable. RH patients and RH controls' group membership could be predicted by GM and LV (r^2=0.5), in that order, but not by WM. GM alone best predicted hemisphere preserved when comparing the two patient groups (r^2=0.28). No predictor/s differentiated patients' ILAE outcome scales.

<u>Cortical Morphometrics</u>

Thickness

Relative to controls' LH, patients' LH showed no differences in CxT in any region whereas, compared to controls' RH, patients' RH had reduced CxT in 15 of the 34 regions and no region with greater thickness. Relative to LH patients, RH patients had reduced CxT in three regions: the caudal middle frontal, rostral middle frontal, and superior frontal regions. See Figure 2 for data and eTables 4-5 for statistics.

Forward logistic regression analysis with CxT indices of all 34 regions could not differentiate LH patients vs. LH controls. However, a model with CxT of caudal middle frontal, frontal pole, lateral occipital, lingual, pars orbitalis, and transverse temporal regions differentiated RH patients vs. RH controls (r^2=0.8). A model with CxT of entorhinal, rostral middle frontal, and superior temporal regions differentiated LH from RH patients (r^2=0.89). No predictor/s differentiated patients' ILAE outcome scales.

Surface Area

Compared with their matched controls, LH patients had no differences in CSA, whereas RH patients had greater CSA in lateral orbitofrontal, paracentral, and parahippocampal cortices. The patient groups also differed from one another: LH patients had greater CSA in the pars opercularis, rostral anterior cingulate, and transverse temporal regions while RH patients had greater CSA in the frontal pole, inferior parietal, parahippocampal, and pars orbitalis regions. See Figure 2 for data and eTables 6-7 for statistics.

Forward logistic regression analysis with CSA indices of all 34 regions could not differentiate LH patients vs. LH controls. However, a model with CSA of entorhinal, lateral orbitofrontal, paracentral, rostral middle frontal, and superior temporal regions differentiated RH patients vs. RH controls (r^2=0.83). A model with CSA of pars orbitalis differentiated LH from RH patients (r^2=0.95), and a model with CSA of paracentral cortex predicted ILAE outcome score (r^2=0.33).

Volume

Compared with matched controls, LH patients had no differences in CV, whereas RH patients had greater CV in rostral middle frontal cortex. The two patient groups differed from one another: LH patients had greater CV in the inferior parietal, pars opercularis, transverse temporal, and rostral anterior cingulate regions while RH patients had greater CV in the pars orbitalis and superior frontal regions. See Figure 2 for data and eTables 8-9 for statistics.)

Forward logistic regression analysis with CV indices of all 34 regions could not differentiate LH patients vs. LH controls. However, a model with CV of rostral middle frontal and pars orbitalis regions differentiated RH patients vs. RH controls (r^2=0.57). A model with CV of pars opercularis and pars orbitalis differentiated LH from RH patients (r^2=0.9), and a model with CV of the paracentral, pars triangularis, and superior parietal regions predicted ILAE outcome score (r^2=0.71).

For an illustration of group differences by individual regions per the logistic regressions, see Figure 3.

Subcortical Morphometrics

LH patients had lower accumbens, caudate, pallidum, and putamen volumes relative to controls' LH. RH patients had lower accumbens and hippocampus volumes relative to controls' RH. LH patients had lower caudate and putamen volumes than RH patients. See Figure 4 for data and eTables 10-11 for statistics.)

Forward logistic regression analysis yielded a model with putamen and accumbens volumes that could differentiate LH patients vs. LH controls (r^2=0.44). A model with accumbens volume differentiated RH patients vs. RH controls (r^2=0.28). A model with putamen volume differentiated LH from RH patients (r^2=0.52). A regression model that included the putamen volume alone differentiated strongly between the preserved RH vs. LH patients, with r^2=0.52. No predictor/s could differentiate patients' ILAE outcome scales.

Analysis Excluding Ablation Cases

One possible explanation for differences between LH vs. RH patients may be because of differences in the extent of the resection, with more LH than RH patients having an ablation: LH – 7 ablation, 12 resection; RH – 2 ablation, 11 resection. Considering this, we recomputed all forward logistic regressions with just the resection cases (12 LH, 11 RH). Notwithstanding loss of statistical power with this reduction, the findings using the three gross measures (LV, GM, and WM) largely mirrored that of the analysis including all patients: whereas no model could predict group membership between LH patients and LH controls or between LH patients and RH patients, a model with LV and GM discriminated between RH patients and RH controls (r^2 = 0.48).

Additionally, cortical analysis with just the resection patients revealed largely similar results to that conducted with all patients. LH patients were differentiated from LH controls with CxT of fusiform and inferior parietal regions as predictors (r^2=0.27). A stronger association between group membership (patient vs. control) for the RH was observed with CxT of frontal pole and posterior cingulate regions as predictors (r^2=.40), and a model with caudal middle frontal and lateral thickness predicted differentiated the two patient groups (r^2=.77). For CSA,

no model distinguished between LH patients vs. LH controls. However, a model with lateral orbito-frontal, parahippocampal, rostral middle frontal, and superior temporal CSA differentiated RH patients vs. RH controls (r^2=0.76), and a model with pars orbitalis CSA differentiated the two patients groups (r^2=0.90). The findings of differences in CV were also well-substantiated in the no-ablation analysis. For CV, no model distinguished between LH patients vs. LH controls. Nevertheless, a model with pars orbitalis and rostral middle frontal differentiated RH patients vs RH controls (r^2=0.54), and a model with pars opercularis CV differentiated the two patient groups (r^2=0.70),

Last, subcortical analysis with resection patients replicated the analysis with all patients. A model with accumbens and putamen volumes differentiated LH patients from LH controls (r^2=0.56); accumbens volume differentiated RH patients from RH controls (r^2=0.33); and putamen volume differentiated the two patient groups (r^2=0.66).

Discussion

Prior morphometry studies, primarily carried out in adults with temporal lobe epilepsy, have typically studied one variable (e.g., CxT or CV) and have also yielded mixed findings[2-5, 37]. Here, we comprehensively characterized the structural integrity of the preserved hemisphere in DRE patients who underwent surgery in *childhood*, a time of greater plasticity than in adulthood [38].

We recruited 32 patients, with surgical interventions ranging from large resections (e.g., hemispherectomy) to focal procedures (e.g., ablation), and compared their preserved hemisphere to that of matched controls, and also compared between patient groups. Using GLM and binary logistic regression, we examined gross indices of LV, GM, and WM volumes; multiple indices of 34 cortical parcels; and nine subcortical regions' volumes. Several findings emerged, summarized below and in Table 1.

First, both patient groups had larger LVs and lower total WM (and GM in RH patients). GM volume alone predicted whether the LH vs. RH was preserved, with less LH than RH atrophy. Second, we quantified CxT, CSA, and CV of 34 cortical regions. GLM showed that LH patients' measures were not different from controls', and forward logistic regressions showed the groups were only differentiated by superior temporal region CSA. RH patients differed from controls on all measures in both GLM and forward logistic regression analyses. Between patient groups, differences in pars orbitalis and pars opercularis CSA and CV manifested as key sites in both analyses. Of subcortical regions, accumbens, caudate, hippocampus, pallidum, and putamen volumes were lower in patients vs. controls, and putamen volume alone (smaller in LH than RH) differentiated the two patient groups.

In summary, we identified differences between patients' preserved RH and each controls' RH and patients' LH (and nearly no differences between LH patients vs. LH controls). The preserved RH has lower CxT and CV but, surprisingly, greater CSA in several regions.

Because the two patient groups were matched on age, gender, initial surgery age, seizure-onset age, and ILAE scores, and age and gender were included as covariates in all GLM analyses, none of these factors accounted for hemispheric differences. Both patient groups were roughly equally represented on each scanner and data were harmonized across scanners. Gross replication of results with sub-analyses of only resection cases also rules out this potential confound. Hemispheric differences are also not obviously explained by differences in lobar resections: both LH and RH groups had four temporal resections and two frontal resections. The LH and RH groups had one and two occipital resections, and two and one parietal resections, respectively (number of cases exceed 32 as some patients have multi-lobar resections).

We observed LH vs RH patients' regional differences primarily in the pars orbitalis and pars triangularis, but also in caudal middle and superior frontal and middle, superior, and transverse temporal regions. These findings, derived from an unbiased, whole-brain, data-driven analysis appear to converge largely on regions implicated in language functions. Resection of the LH, typically the native dominant language hemisphere, is likely a source of alterations of LH homologues of language areas in the preserved RH. In contrast, RH resection results in remarkably few morphometric LH differences relative to controls' LH, consistent with RH resection children showing better cognitive and language outcomes.[39]

The hypothetical account, then, is that after LH resection, regions of the preserved RH that are homotopic with LH language regions come to assume language functions. The process of accommodating language in the RH (which has some nascent language function[40, 41]) potentially triggers morphometric RH changes. Much evidence attests to the fact that, in early childhood, both hemispheres are predisposed to language function. For example, in contrast to the marked aphasias evident in adults with brain injury, many children with unilateral brain injury have unaffected language function, independent of affected hemisphere.[42] Likewise, adolescents and young adults who suffered a LH perinatal ischemic stroke displayed sentence-processing abilities comparable to controls, presumably due to RH language engagement.[41, 43] The idea of RH plasticity for language has also been observed with neuroimaging: individuals with peri- or prenatal periventricular damage to the LH evince RH blood-oxygen-level-dependent activation during a silent word generation task, and the activation extent equaled that of the LH in controls.[44] These findings attest to the potential for functional reorganization of language to RH homotopic frontotemporal regions.

What remains to be explained is why we see reduction in RH homotopic language regions rather than maintenance or even expansion, which is often associated with the assumption of a new function. The cortical thinning we observe is widespread but, notably, primarily homotopic with or proximal to standard LH language regions (e.g., lateral orbitofrontal, middle temporal, rostral middle frontal, superior frontal, and transverse temporal areas). Presumably, as is also true over the course of normal development, cortical regions are pruned through the removal of inefficient synapses as functions are acquired and the cortical tissue loss reflects improved neural processing by optimizing brain circuits for particular operations.[45] Cortical thinning is coupled with morphological changes,[45] and, in childhood, GM thinning,[46] specifically, is associated with changes in behavior, which may be associated with increases in WM.[47] In a longitudinal study, changes of CxT asymmetry in the inferior frontal gyrus, constituting a portion of Broca's area in the LH, is suggestive of a neural correlate of language improvement in children's brains at roughly ages 5-7 yr.[48] Furthermore, changes to regions proximal to RH homotopic language areas observed here may reflect the imprecision of the changes (for example, in pruning of neurons and associated dendrites).

Given the cross-sectional, rather than longitudinal, design of the current study, we are unable to evaluate changes over time. Whether the profiles of the childhood DRE brains improved specifically over presurgical profiles or not and whether the relationship between these changes over time, hemisphere of resection, and cognition (specifically language) remain unaddressed and serve as rich fodder for future studies.

Data availability

A DOI is reserved on Carnegie Mellon University's KiltHub (Figshare) repository: 10.1184/R1/24153423. Data/code will be published upon article publication.

Acknowledgements

We are grateful to all our study participants, who generously donated their time to make this research possible. We thank the patients, the controls, and their families for their time and cooperation, as well as the Pediatric Epilepsy Surgery Alliance, especially Monika Jones, JD, Chief Executive Officer. We also thank Scott Kurdilla, Dr. Tina Liu, Sophia Robert, Mark Vignone, and Debbie Viszlay for assistance with data acquisition; and Dr. Vladislav Ayzenberg, Nicholas Blauch, Dr. Alison Butler, Dr. Samuel Dienel, Dr. Joel Greenhouse, Dr. Alexander Layden, Dr. Timothy Verstynen, and the CMU Visual Cognition group for their helpful discussions and feedback.

Funding

This research was supported by Award Numbers T32GM081760 and T32GM144300 from the National Institute of General Medical Sciences (M.C.G.), R01EY027018 (M.B. and C.P.) from the National Eye Institute, a fellowship from the American Epilepsy Society (#847556) (M.C.G.), a scholarship from the American Psychological Foundation (M.C.G.), and funding from the CMU-Pitt BRIDGE Center. DRG was supported by the National Institute of Mental Health Intramural Research Program (ZICMH002888) of the National Institutes of Health/Health and Human Services, USA. M.B. also acknowledges support from P30 CORE award EY08098 from the NEI, NIH, and unrestricted supporting funds from The Research to Prevent Blindness Inc, NY, and the Eye & Ear Foundation of Pittsburgh. The content is solely the responsibility of the authors and does not necessarily represent the official views of NIGMS, NEI, NIMH, AES, or APF.

Competing interests

Marlene Behrmann is a founder of Precision Neuroscopics.

Supplementary material

Supplementary material will be available online.

Figures

Figure 1: Volume of (top) lateral ventricles, (middle) grey matter, and (bottom) white matter for patients and controls, separately for the preserved left and right hemispheres. *: $p<.05$; **: $p<.01$; ***: $p<.001$. Each dot represents a single participant, and the solid horizontal line in the box indicates the median of the group.

Figure 2: Median cortical thickness (top), normalized cortical surface area (middle), and region volume (normalized by total volume; bottom) for 34 cortical regions for patients with a preserved left hemisphere or right hemisphere relative to their matched controls. *: *p*<0.05; **: *p*<0.01; ***: *p*<0.001.

Figure 3: Regions that significantly distinguished between patients versus controls in left columns, right hemisphere patients versus left hemisphere patients in right columns per binary logistic regression modelling for (A) cortical thickness, (B) cortical surface area, and (C) cortical volume. LH = left hemisphere; RH = right hemisphere.

Figure 4: Subcortical regional volumes (normalized) for patients with a preserved left hemisphere or right hemisphere relative to their matched controls *: $p < .05$; **: $p < .01$; ***: $p < .001$.

Table 1 Summary of all results for univariate and multivariate analyses

Measures	LH patients/LH controls	RH patients/RH controls	RH patients/LH patients
Gross volumes			
General linear model			
Lateral Ventricle	patients > controls	patients > controls	n.s.
Grey Matter	n.s.	patients < controls	RH < LH
White Matter	patients < controls	patients < controls	n.s.
Logistic regression	LV and WM	GM and LV	GM
Cortical			
Thickness			
General linear model	n.s.	patients < controls: caudal middle frontal, cuneus, frontal pole, fusiform, insula, isthmus, lateral orbitofrontal, lingual, middle temporal, paracentral, posterior cingulate, precuneus, rostral middle frontal, superior frontal, transverse temporal	RH < LH: caudal middle frontal, rostral middle frontal, superior frontal
Logistic regression	n.s.	caudal middle frontal, frontal pole, lateral occipital, lingual, pars orbitalis, transverse temporal	entorhinal, rostral middle frontal, superior temporal
Surface area			
General linear model	n.s.	patients > controls: lateral orbitofrontal, parahippocampal, paracentral	LH > RH: pars opercularis, rostral anterior cingulate, transverse temporal; RH > LH: frontal pole, inferior parietal, parahippocampal, pars orbitalis
Logistic regression	superior temporal	entorhinal, lateral orbitofrontal, paracentral, rostral middle frontal, superior temporal	pars orbitalis
Volume			
General linear model	n.s.	patients < controls: rostral middle frontal area	LH > RH: pars opercularis, rostral anterior cingulate, superior frontal, transverse temporal; RH > LH: inferior parietal, pars orbitalis
Logistic regression	n.s.	rostral middle frontal, pars orbitalis	pars opercularis, pars orbitalis areas
Subcortical volumes			
General linear model	patients < controls; accumbens, caudate, pallidum, putamen	patients < controls: accumbens, hippocampus	RH > LH: caudate, putamen
Logistic regression	accumbens, putamen	accumbens	putamen

n.s.: not significant
LH: left hemisphere
RH: right hemisphere

References

1.   Weil AG, Fallah A, Wang S, Ibrahim GM, Elkaim LM, Jayakar P, et al. Functional hemispherectomy: can preoperative imaging predict outcome? Journal of neurosurgery Pediatrics. 2020;25(6).

2.   Federico P, Wiebe S. Is bad brain worse than no brain? Salvaging the cerebral cortex in epilepsy. Brain. 2020;143(11):3172-5.

3.   Galovic M, de Tisi J, McEvoy AW, Miserocchi A, Vos SB, Borzi G, et al. Resective surgery prevents progressive cortical thinning in temporal lobe epilepsy. Brain. 2020.

4.   McDonald CR. Flattening the Curve: SlowingAge-Accelerated Brain Atrophy With Epilepsy Surgery. Current Literature in Clinical Research: Epilepsy Currents. 2021;https://doi.org/10.1177/15357597211001066.

5.   Galovic M, van Dooren VQH, Postma T, Vos SB, Caciagli L, Borzi G, et al. Progressive Cortical Thinning in Patients With Focal Epilepsy. JAMA Neurol. 2019.

6.   Caciagli L, Bernasconi A, Wiebe S, Koepp MJ, Bernasconi N, Bernhardt BC. A meta-analysis on progressive atrophy in intractable temporal lobe epilepsy: Time is brain? Neurology. 2017;89(5):506-16.

7.   Grinenko O, Li J, Mosher JC, Wang IZ, Bulacio JC, Gonzalez-Martinez J, et al. A fingerprint of the epileptogenic zone in human epilepsies. Brain. 2018;141(1):117-31.

8.   Fonseca E, Saaria-Estrada S, Pareto D, Turon M, Quinatana M, E. S, et al. Relationship between visuoperceptual functions and parietal structural abnormalities in temporal lobe epilepsy. Brain Imaging and Behavior. 2022;in press.

9.   Zhao Y, Zhang C, Yang H, Liu C, Yu T, Lu J, et al. Recovery of cortical atrophy in patients with temporal lobe epilepsy after successful anterior temporal lobectomy. Epilepsy Behav. 2021;123:108272.

10.   Lopez SM, Aksman LM, Oxtoby NP, Vos SB, Rao J, Kaestner E, et al. Event-based modeling in temporal lobe epilepsy demonstrates progressive atrophy from cross-sectional data. Epilepsia. 2022;63(8):2081-95.

11.   Park BY, Lariviere S, Rodriguez-Cruces R, Royer J, Tavakol S, Wang Y, et al. Topographic divergence of atypical cortical asymmetry and atrophy patterns in temporal lobe epilepsy. Brain. 2022;145(4):1285-98.

12.   Azzony S, Moria K, Alghamdi J. Detecting Cortical Thickness Changes in Epileptogenic Lesions Using Machine Learning. Brain Sci. 2023;13(487):doi.org/10.3390/brainsci13030487.

13.   Frangou S, Modabbernia A, Williams SCR, Papachristou E, Doucet GE, Agartz I, et al. Cortical thickness across the lifespan: Data from 17,075 healthy individuals aged 3-90 years. Hum Brain Mapp. 2021.

14.   Wierenga LM, Langen M, Oranje B, Durston S. Unique developmental trajectories of cortical thickness and surface area. Neuroimage. 2014;87:120-6.

15.   Lyall AE, Shi F, Geng X, Woolson S, Li G, Wang L, et al. Dynamic Development of Regional Cortical Thickness and Surface Area in Early Childhood. Cereb Cortex. 2015;25(8):2204-12.

16.   Sanders AFP, Baum GL, Harms MP, Kandala S, Bookheimer SY, Dapretto M, et al. Developmental trajectories of cortical thickness by functional brain network: The roles of pubertal timing and socioeconomic status. Developmental cognitive neuroscience. 2022;57:101145.

17.   Scheffer IE, Berkovic S, Capovilla G, Connolly MB, French J, Guilhoto L, et al. ILAE classification of the epilepsies: Position paper of the ILAE Commission for Classification and Terminology. Epilepsia. 2017;58(4):512-21.

18.   Reuter M, Rosas HD, Fischl B. Highly accurate inverse consistent registration: a robust approach. Neuroimage. 2010;53(4):1181-96.

19.   Sled JG, Zijdenbos AP, Evans AC. A nonparametric method for automatic correction of intensity nonuniformity in MRI data. IEEE Trans Med Imaging. 1998;17(1):87-97.

20.   Ségonne F, Dale AM, Busa E, Glessner M, Salat D, Hahn HK, et al. A hybrid approach to the skull stripping problem in MRI. Neuroimage. 2004;22(3):1060-75.

21.   Maallo AMS, Granovetter MC, Freud E, Kastner S, Pinsk MA, Patterson C, et al. Large-scale resculpting of cortical circuits in children after surgical resection. Scientific reports. 2020;10(1):21589.

22.   Glen D, Levenstein JM, Granovetter MC, Maallo AMS, Behrmann M. LARGE Lesion Brain Alignment with AFNI. Organization of Human Brain Mapping2021.

23.   Desikan RS, Segonne F, Fischl B, Quinn BT, Dickerson BC, Blacker D, et al. An automated labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest. Neuroimage. 2006;31(3):968-80.

24.   Fischl B, van der Kouwe A, Destrieux C, Halgren E, Segonne F, Salat DH, et al. Automatically parcellating the human cerebral cortex. Cereb Cortex. 2004;14(1):11-22.

25.   Fischl B, Salat DH, Busa E, Albert M, Dieterich M, Haselgrove C, et al. Whole brain segmentation: automated labeling of neuroanatomical structures in the human brain. Neuron. 2002;33(3):341-55.

26.   Fischl B, Salat DH, van der Kouwe AJ, Makris N, Segonne F, Quinn BT, et al. Sequence-independent segmentation of magnetic resonance images. Neuroimage. 2004;23 Suppl 1:S69-84.

27.   Westman E, Aguilar C, Muehlboeck JS, Simmons A. Regional magnetic resonance imaging measures for multivariate analysis in Alzheimer's disease and mild cognitive impairment. Brain Topogr. 2013;26(1):9-23.

28.   Fischl B, Dale AM. Measuring the thickness of the human cerebral cortex from magnetic resonance images. Proc Natl Acad Sci U S A. 2000;97(20):11050-5.

29.   Han X, Jovicich J, Salat D, van der Kouwe A, Quinn B, Czanner S, et al. Reliability of MRI-derived measurements of human cerebral cortical thickness: the effects of field strength, scanner upgrade and manufacturer. Neuroimage. 2006;32(1):180-94.

30.   Reuter M, Schmansky NJ, Rosas HD, Fischl B. Within-subject template estimation for unbiased longitudinal image analysis. Neuroimage. 2012;61(4):1402-18.

31.   TeamCore R. R: A Language and Environment for Statistical Computing. Vienna, Austria: R Foundation for Statistical Computing; 2020.

32.   Fortin J-P. NeuroCombat: Harmonization of Multi-Site Imaging Data with ComBat2021.

33.   Fortin JP, Cullen N, Sheline YI, Taylor WD, Aselcioglu I, Cook PA, et al. Harmonization of cortical thickness measurements across scanners and sites. Neuroimage. 2018;167:104-20.

34.   Benjamini Y, Yekutieli D. The control of the false discovery rate in multiple testing under dependency. Ann Statist 2001;29(1165-1188).

35.   Lee MD, Wagenmakers EJ. Bayesian Cognitive Modeling: A Practical Course. Cambridge: Cambridge University Press; 2013.

36.   Wagenmakers EJ. A practical solution to the pervasive problems of p values. Psychon Bull Rev. 2007;14(5):779-804.

37.   Li W, Jiang Y, Qin Y, Li X, Lei D, Zhang H, et al. Cortical remodeling before and after successful temporal lobe epilepsy surgery. Acta Neurol Scand. 2022;146(2):144-51.

38.   Bourne JA. Unravelling the development of the visual cortex: implications for plasticity and repair. Journal of Anatomy. 2010;217(4):449-68.

39.   Puka K, Jones M, Mathern GW. Functional cognitive and language outcomes after cerebral hemispherectomy for hemimegalencephaly. Epilepsia. 2021.

40.   Martin KC, Seydell-Greenwald A, Berl MM, Gaillard WD, Turkeltaub PE, Newport EL. A Weak Shadow of Early Life Language Processing Persists in the Right Hemisphere of the Mature Brain. Neurobiol Lang (Camb). 2022;3(3):364-85.

41.   Newport EL, Seydell-Greenwald A, Landau B, Turkeltaub PE, Chambers CE, Martin KC, et al. Language and developmental plasticity after perinatal stroke. Proc Natl Acad Sci U S A. 2022;119(42):e2207293119.

42.   Bates E, Reilly J, Wulfeck B, Dronkers N, Opie M, Fenson J, et al. Differential effects of unilateral lesions on language production in children and adults. Brain Lang. 2001;79(2):223-65.

43.   Martin KC, Seydell-Greenwald A, Berl MM, Gaillard WDT, P. E., Newport EL. A Weak Shadow of Early Life Language Processing Persists in the Right Hemisphere of the Mature Brain. Neurobiology of Language. 2022;https://doi.org/10.1162/nol_a_00069.

44.   Staudt M, Lidzba K, Grodd W, Wildgruber D, Erb M, Krageloh-Mann I. Right-hemispheric organization of language following early left-sided brain lesions: functional MRI topography. Neuroimage. 2002;16(4):954-67.

45.   Natu VS, Gomez J, Barnett M, Jeska B, Kirilina E, Jaeger C, et al. Apparent thinning of human visual cortex during childhood is associated with myelination. Proc Natl Acad Sci U S A. 2019;116(41):20750-9.

46.   Sowell ER, Thompson PM, Leonard CM, Welcome SE, Kan E, Toga AW. Longitudinal mapping of cortical thickness and brain growth in normal children. J Neurosci. 2004;24(38):8223-31.

47.   Giedd JN, Blumenthal J, Jeffries NO, Castellanos FX, Liu H, Zijdenbos A, et al. Brain development during childhood and adolescence: a longitudinal MRI study. Nat Neurosci. 1999;2(10):861-3.

48.   Qi T, Schaadt G, Friederici AD. Cortical thickness lateralization and its relation to language abilities in children. Developmental cognitive neuroscience. 2019;39:100704.

A.

RH Patients vs RH Controls

Caudal middle frontal
Transverse temporal
Frontal pole
Lateral occipital
Pars orbitalis

RH Patients vs LH Patients

Rostral middle frontal
Superior temporal

LH Patients vs LH Controls

Lingual

Entorhinal

B.

RH Patients vs RH Controls

Rostral middle frontal
Lateral orbitofrontal
Superior temporal

RH Patients vs LH Patients

Pars orbitalis

LH Patients vs LH Controls

Paracentral

Entorhinal

C.

RH Patients vs RH Controls

Rostral middle frontal
Pars orbitalis

RH Patients vs LH Patients

Pars opercularis
Pars orbitalis