Genomic signatures of adaptation in native lizards exposed to human-introduced
fire ants

Braulio A. Assis*#[1,2], Alexis P. Sullivan#[1,3], Stephanie Marciniak[2,4], Christina M. Bergey[1,2,5],

Vanessa Garcia[1], Zachary A. Szpiech[1,6], Tracy Langkilde*[1,7], George H. Perry*[1,2,7]

[1] Department of Biology, Penn State University, University Park, PA, USA
[2] Department of Anthropology, Penn State University, University Park, PA, USA
[3] Institute for Systems Genetics, NYU Langone Health, New York City, NY, USA
[4] Department of Anthropology, McMaster University, Hamilton, Ontario, Canada, L8S 4L9
[5] Department of Genetics, Rutgers, The State University of New Jersey, Piscataway, NJ 08854,
USA
[6] Institute for Computational and Data Sciences, Penn State University, University Park, PA,
USA
[7] Huck Institutes of the Life Sciences, Penn State University, University Park, PA USA

*bmd5458@psu.edu; alexispsullivan@gmail.com; tll30@psu.edu; ghp3@psu.edu

# Authors contributed equally

Keywords: genetic adaptation; anthropogenic; invasive species; rapid adaptation

Abstract

Understanding the process of genetic adaptation in response to human-mediated ecological

change will help elucidate the eco-evolutionary impacts of human activity. Red fire ants

(*Solenopsis invicta*) spread across Southeastern USA since their accidental introduction via Port

Mobile, Alabama in the 1930s, serving today as both novel venomous predator and novel toxic

prey to native eastern fence lizards (*Sceloporus undulatus*). To identify potential signatures of

genetic adaptation in lizards to invasive fire ants, we generated whole genome sequencing data

from 420 native fence lizards sampled across three populations, two of which had not been

invaded by fire ants (in Tennessee and Arkansas) and one which had been invaded for ~70 years

(Alabama). We detected signatures of positive selection exclusive to the exposed Alabama

population for genetic variants overlapping genes related to the membrane attack complex of the

complement immune system, growth factor pathways, and morphological development. Prior

work identified a relationship between increased lizard survival of fire ant attack and longer hind

limbs, which lizards use to remove ants from their bodies. Furthermore, we conducted a genome-

wide association study with 381 Alabama lizards to identify 24 hind limb length-associated

genetic loci. For two loci, positive-effect alleles occur in high frequency and overlap genomic

regions that are highly differentiated from the populations naïve to fire ants. Collectively, these

findings represent plausible genetic adaptations in response to fire ant invasion, whereby

morphological differentiation may increase survival against swarming ants and altered immune

responses may allow the exploitation of a novel, toxic food resource.

Significance statement

Human activity can force interactions between species from distinct ecological backgrounds. These interactions can consequently impose novel selective pressures on endemic populations via predation or disruption of ecological niches through community-wide effects. While some endemic taxa have been able to adapt biologically to these disruptions, we do not have a full understanding of the underlying genetic processes that may allow it. Here we identify genomic signatures of recent adaptation nearby genes involved in morphological and immunological processes in native fence lizards that are consistent with pressures imposed by the venomous, predatory fire ants introduced by humans. These signatures are largely absent from lizard populations that are naïve to fire ants.

Introduction

Endemic species worldwide face rapid environmental change resulting from various types of human activity. For example, human-mediated translocation of species into new environments promotes novel ecological interactions, often with detrimental effects (Saul and Jeschke 2015). The consequences can be dire: species introductions have been identified as an underlying cause of at least 170 animal extinctions (Clavero and García-Berthou 2005). In some invasive-endemic interaction cases, standing genetic variation may provide endemic populations with the raw materials needed for quick adaptative responses (Barrett and Schluter 2008). Yet even then, such adaptations can induce cascading effects on the broader ecosystem (Mooney and Cleland 2001; Schlaepfer et al. 2005; Strauss et al. 2006; Hale et al. 2016). Therefore, understanding the process of genetic adaptation to novel species interactions will better inform us of the scope of potential ecological impacts related to human behavior.

In the 1930s, the red imported fire ant, *Solenopsis invicta*, was accidentally transported, by humans, from South America (presumably northeastern Argentina) to Port Mobile, Alabama, in the United States of America (Ascunce et al. 2011). Since then, fire ants have established and steadily expanded their range in the Southeastern USA, with both economic and public health impacts arising from their potent venom and aggressive nature (Gruber et al. 2022). Furthermore, ecological impacts stemming from fire ant invasion are marked and diverse. In the United States, fire ants outcompete and displace native ants, with cascading effects on the broader invertebrate community (Porter and Savignano 1990; Morrison 2002; Roeder et al. 2021). Experimental studies have confirmed that fire ants directly and indirectly impact various endemic small vertebrates. Specifically, predation by fire ants led to a reduction of up to 66% of *Ambystoma* salamander populations within 48h (Todd et al. 2008), whereas fire ant suppression led to

significant increases in small vertebrate abundance (Stahlschmidt et al. 2018). Meanwhile, fire

ant disruption of arthropod communities led to a 10% decrease in the number of eastern bluebird

fledglings and to the displacement of adult birds (Ligon et al. 2011).

Among impacted vertebrates, the eastern fence lizard *Sceloporus undulatus* has been an

important model system for studying the ecological impacts of this fire ant introduction. First,

despite their toxicity, fire ants have become a novel prey item in the fence lizard's diet (Robbins

et al. 2013). Second, fire ants are a novel predator of fence lizards: swarming worker ants can

envenomate and kill juvenile and adult individuals (Langkilde 2009), as well as prey on their

eggs (Thawley and Langkilde 2016). The predatory impact is likely very high: experimental

removal of fire ants from enclosures increased fence lizard recruitment by ~60% (Darracq et al.

2017), while lizard hatchling survival is negatively associated with fire ant mound density

(Gifford et al. 2017). Even lizards that initially survive fire ant encounters experience a 20%

increase in mortality rate over the 11 weeks post-exposure relative to unexposed lizards

(Langkilde and Freidenfelds 2010). Yet, fence lizards remain abundant in fire ant-invaded

habitats, prompting behavioral, ecological, and evolutionary investigations into the underlying

mechanisms.

Exposure to fire ant venom induce wide-ranged immunological responses in fence lizards

(Tylan et al. 2020; Tylan et al. 2023). Prior to venom exposure, however, behavioral defenses are

also employed by fence lizards against swarming fire ants. These defenses include body-

twitching – using their hind legs to directly remove ants with a flicking motion – and fleeing

from the attack (Langkilde 2009). In behavioral trials, the success of these tactics was positively

linked to hind limb length (relative to body size), with a ~20% longer relative hind limb resulting

in the removal of ~30% more fire ants (Langkilde 2009). A morphological study of fence lizard

museum specimens that predate the fire ant introduction revealed a latitudinal cline where lizards

from the southern extreme of their range had ~5% *smaller* relative hind limb lengths than those

in the northern extreme (a difference of ~14° in latitude), a trend likely driven by temperature

and precipitation gradients (Thawley et al. 2019). However, a different pattern is observed in the

present day, following this introduction. Relative limb length is greater in populations with the

longest history of fire ant invasion; a pattern that could be explained by recent evolutionary

change in response to fire ants (Thawley et al. 2019). A positive, albeit non-significant,

correlation between mother and offspring relative limb lengths suggests that this trait is likely

heritable in fence lizards (Langkilde 2009), as it is for other lizards (Tsuji et al. 1989; Kolbe and

Losos 2005). Together, these observations suggest that relative limb length in fence lizards may

have been and could continue to be a viable target for natural selection in response to novel

interactions with fire ants.

Results

In this study we sought to identify potential fire ant invasion-related genomic signatures

of adaptation in eastern fence lizards. To do so, we analyzed whole-genome sequencing data for

a total of n=420 lizards from three sampling sites (Fig 1A) – two sites were naïve to fire ants (in

Arkansas, n=19, and Tennessee, n=20) and one had a long history of fire ant exposure (66 to 77

years in Alabama, n=381). We first generated high-coverage sequencing data (average of 13.01X

sequence reads per site, per individual) from n=20 individuals from each site (total n=59; one

sample failed QC) for population history and evolutionary genomic analyses. We also generated

low-coverage (4.39X) sequencing data from an additional n=361 individuals from the fire ant-

invaded site in Alabama (total n=381 when including those individuals also in the high-coverage

dataset) to identify any significant genotype associations with relative hind limb length.

The lizards from the fire ant-exposed population were collected between 2006 and 2017

from the Solon Dixon Forestry Education Center, in Andalusia, Alabama (AL). Fire ants were

recorded and this site was quarantined in the early 1940s (Code of Federal Regulations 2006).

Estimates of eastern fence lizard generation time vary from 1.15 to 2.24 years (Rodríguez-

Romero et al. 2011); thus, the approximate 70 years of coexistence with fire ants may have

encompassed 31 to 60 generations of fence lizards. The other two sampled populations had no

record of invasion by fire ants at the time of sampling. These lizards were collected from St.

Francis National Forest, Arkansas (AR) in 2006, and from Edgar Evins State Park, Tennessee

(TN) between 2011 and 2013.

We mapped sequence reads from the high-coverage dataset to the fence lizard 1.0

reference assembly (Westfall et al. 2021). After quality control and filtering (see *Methods*), we

identified a total of 46,934,027 SNPs across the three populations, though only 2,249,567 of

these SNPs were variable in all three populations. The AL population was the most genetically

diverse (AL genome-wide mean pairwise nucleotide distance $\pi = 0.125$; TN $\pi = 0.093$; AR $\pi =$

0.076) and also had the largest number of private variants by far (AL = 24,705,862; TN =

7,360,015; AR =5,985,756). The number of SNPs variable in two of the three populations were

5,975,685 for TN and AL; 3,452,917 for AR and AL; and 4,378,018 for TN and AR. In the AL

population, the presence of such a large proportion of rare genetic variants – including

10,507,067 singleton or doubleton SNPs – is consistent with a recent, large-scale demographic

expansion (Slatkin 1993; Amos and Harwood 1998), even in the face of fire ant invasion.

We used several descriptive approaches to examine genetic relationships among the three

populations. First, we examined patterns of population differentiation using the F_{ST} statistic,

restricting each pairwise comparison to SNPs that were either variable in both populations or for

which the minor allele in one population was fixed in the second. Average F_{ST} values were AL-

TN = 0.210, AL-AR = 0.141, and TN-AR = 0.223 (Fig 1b). The three populations also clustered

independently based on results from a principal components analysis (Fig 1b) when retaining

only the 11,437,455 SNPs variable within or between at least two populations (results were

qualitatively equivalent when using all SNPs; Supp fig S1). We also used this set of SNPs to

construct a genetic distance matrix-based neighbor-joining tree, in which individuals from each

population were distinctly separated (Fig 1c). Lastly, we used ADMIXTURE (Alexander et al.

2009) to compute model-based estimates of individual ancestry. When specifying k=3

populations (which has the lowest cross-validation error; Supp Figure S2), cluster membership

proportions for members of all three populations were distinct (Figure 1d). With k=2, AL and

AR membership proportions were distinct, with TN individuals exhibiting a mix of the two.

Taking all these results together, despite the Mississippi River being a putative biogeographic

barrier for AR fence lizards, we did not observe a clear, strong genetic separation between AR

vs. AL+TN, precluding the definitive assignment of any population as a true outgroup for

downstream evolutionary analyses.

A Fire ant invasion and fence lizard populations

B Genetic differentiation

Fence lizard sample collection sites
- Solon Dixon Forestry Education Center, AL
  n = 20 HC / 381 LC, 2006-2017
- St. Francis National Forest, AR
  n = 19, 2006
- Edgar Evins State Park, TN
  n = 20, 2011-2013

Fire ant dispersal
- 1930s - 1949
- 1950 - 1969
- 1970 - 1989
- 1990 - 2009
- 2010 - present

C Genetic tree **D** Pop. structure

Figure 1. **Invasive fire ant distribution and fence lizard population structure**. *A*: Reported detection and quarantine of the red imported fire ant *Solenopsis invicta* in the Southeastern US since the 1930s (Code of Federal Regulations 2006), along with sampling sites, sampling periods, and number of collected eastern fence lizards, *Sceloporus undulatus*. *B*. Principal Components Analysis and mean Weir & Cockerham F_{ST} values for the three pairwise population comparisons. *C*. Genetic distance-based neighbor-joining tree analysis. *D*. Admixture analysis for K = 2 and K = 3 ancestral groups. K = 3 yielded the lowest rate of cross-validation error (supp figure S2).

Signatures of selection exclusive to Alabama lizards overlap morphology- and immune system-related genes

To identify candidate regions for recent positive selection in each lizard population, we used three different population genetic approaches: Tajima's D (Tajima 1989); saltilassi (DeGiorgio and Szpiech 2022), and LSBL (locus-specific branch length; (Shriver et al. 2004). If selective pressures imposed by fire ants resulted in recent genetic adaptation in the fire ant-exposed AL population, then we would predict the commensurate signatures of selection reflecting such adaptations to be absent in the two northern populations naïve to fire ants.

Tajima's D compares the mean number of pairwise genetic differences to the sample

size-corrected number of variable sites in a population across a given genomic region. Low

Tajima's D values (an excess of rare alleles) may reflect recent population expansion or positive

selection (Slatkin 1993; Amos and Harwood 1998). With genome-wide deviations from D=0

most likely reflecting past demographic history, we considered genomic regions containing

Tajima's D values in the lowest 0.5% of all 100 Kb regions for a given population as positive

selection candidates. For AL we identified n=143 such regions (D < -1.81; Figure 2a). For the

two uninvaded populations, AR and TN, we identified n=98 (D < -2.11) and n=97 (D < -1.9)

Tajima's D candidate selection regions, respectively (Supp fig S3-S4).

The program saltilassi computes likelihood ratio tests on the haplotype frequency

spectrum of a given population to identify haplotypes with high frequencies compared to the

genome-wide expectation as candidate targets of past positive selection. We similarly considered

regions containing one or more variants in the top 0.5% distribution of saltilassi's Λ statistic (see

Methods) as selection candidate regions (for AL: $\Lambda > 92.24$ and n=1582 regions; Figure 2a; For

AR: $\Lambda > 407.56$ and n = 566 regions; for TN: $\Lambda > 252.27$ and n = 764 regions; Supp Figure S5-

S6).

Finally, we used the LSBL statistic (Shriver et al. 2004) to identify variants with

frequencies highly differentiated in one population relative to each of the two others, based on all

pairwise F_{ST} values (LSBL$_A$ = (AB F_{ST} + AC F_{ST} – BC F_{ST}) / 2). Here, we considered regions as

candidates for positive selection if at least 3 SNPs in the top 0.1% of the LSBL distribution were

observed within 50Kb of one another and in high linkage disequilibrium ($r^2 \geq 0.9$). In total, we

identified n=2,210 candidate regions (LSBL > 0.76) for AL (Fig 2x), n=2,960 regions (LSBL >

0.74) for TN, and n=1,686 regions (LSBL > 0.78) for AR (Supplementary Figures S7-S8;

Supplementary Table S9-S11).

We initially focused our analysis on genomic regions that were flagged as candidates for

a history of past selection by at least two of the above approaches for a given population,

resulting in a dataset of 42, 35, and 24 multi-signal selection candidate regions for AL, TN, and

AR, respectively (Supp Table S12). We used g:Profiler (Raudvere et al. 2019) to perform

functional profiling enrichment analyses to identify known biological and molecular functions

and pathways (see *Methods*) significantly overrepresented among the set of genes overlapping or

nearby (± 25Kb) the candidate selection regions for each population.

For the fire ant-invaded population, AL, genes within the multi-signal selection candidate

regions were significantly enriched for multiple anatomical structure and the immune system

functional categories (supp table S13), including Myofibril (GO:0030016; 7 observed vs. 1.21

expected genes; Fisher's one-tailed test, FDR adjusted = 0.011) and Complement and

coagulation cascades (KEGG:04610; 3 observed vs. 0.18 expected genes; FDR = 0.042). One of

the multi-signal candidate regions that contains myofibril-related genes overlaps a myosin gene

cluster on chromosome 2 (Figure 2d). The different myosin proteins are responsible for the

distinct contractile properties across muscle cells (Weiss and Leinwand 1996; Foth et al. 2006).

The complement immune system genes of interest include *C8A* and *C8B* (Figure 2d), which are

directly involved in the membrane attack complex (MAC). Nucleated cells targeted by the MAC

can undergo autoimmune and inflammatory processes through the secretion of proinflammatory

proteins such as IL-β and IL-18 (Morgan 2016; Xie et al. 2020), making genetic variants within

or nearby these genes plausible targets for natural selection by fire ant venom exposure. Lastly,

another multi-selection signature (Tajima's D and lassi) locus of interest is a 411Kb region in

chromosome 1 that overlaps two genes – *COL4A1* and *COL4A2* – involved in the production of

collagen IV (Figure 2). Collagen IV is the main component of the cellular basal membrane and is

responsible for skeletal muscle stability (Koskinen et al. 2001; Csapo et al. 2020). In the green

anole *Anolis carolinensis*, skeletal muscle is the tissue with second highest expression of

COL4A1 and the third highest expression of *COL4A2* (Bastian et al. 2021).

Among regions identified in the Tajima's D analysis (but not necessarily as outliers with

the other two statistics), growth and growth factor-related pathways were significantly

overrepresented in the invaded AL lizard population (supp table S14). Significant enrichments

included regulation of insulin-like growth factor transport and uptake by insulin-like growth

factor binding proteins (REAC:R-MMU-381426; 7 observed vs. 1.12 expected genes; FDR <

0.01) and osteoblast signaling (WP:WP238; 2 observed vs. 0.1 expected genes; FDR = 0.042).

Specific genes involved in these enrichments include the macrophage colony-stimulating factor

gene *CSF1* (chromosome 4), Insulin-like growth factor 1 (*IGF1*; chromosome 5), *IGF2*

(chromosome 1), and the Insulin-like Growth Factor binding protein 1 (*IGFBP1*; chromosome

6), which all serve as primary drivers of embryonic growth (Holt 2002), and fibroblast growth

factor 23 (*FGF23*; chromosome 5), which has significant expression in bone tissue and is

involved in osteoblast differentiation (Wang et al. 2008).

For the TN and AR populations, naïve to fire ants, genes overlapping candidate regions

for positive selection in at least two statistics showed no functional enrichments analogous to

those in the fire ant-invaded AL population. In TN (supp table S16), the most significant

enrichment was for ovarian infertility (WP273; FDR = 0.021), while there were no significant

enrichments for AR. However, when considering only the saltilassi results, there was a

functional enrichment for negative regulation of developmental growth in AR (GO:0048640, 6

observed vs. 0.28 expected genes; FDR < 0.001) and analogous terms (Supp table S20).

Figure 2. **Signatures of genetic adaptation in fence lizards from the fire ant-invaded population in Alabama**.

A. Top: genome-wide Tajima's D in 100Kb windows with a 20Kb step. Candidate selection regions fall below the

0.05th percentile cutoff (D < -1.81). Middle: genome-wide saltilassi statistic for sweeping haplotypes (DeGiorgio

and Szpiech 2022). Candidate selection regions fall above the 99.5th percentile cutoff (Λ > 92.24). Bottom: locus-

specific branch length (Shriver et al. 2004) means for the AL population in 50Kb windows and a 10Kb step. *B.*

Number of genes overlapping or nearby (± 25Kb) candidate selection regions for each of the three selection

statistics. *C.* Functional enrichment analysis for genes overlapping candidate regions under selection for Tajima's D.

Plausible adaptations to selective pressure from fire ants include variants in genes for myosin, collagen, growth

factors, and complement immune response. *D.* Selected multi-signal candidate selection regions, with scores for the

Tajima's D and saltilassi statistics. Horizontal lines represent the same significance thresholds in *A.*

Genomic associations with lizard limb length in the fire ant-invaded population in Alabama

To uncover genomic associations with relative limb length variation, we brought low coverage

(average 4.39X sequence coverage per individual) whole genome sequence data from an

additional n=361 lizards from the same population in Alabama, into our study. We first used the

high-coverage data for the 20 AL individuals (described above) to improve genotyping rates for

the 361 low-coverage genomes via genomic imputation (Browning et al. 2018). Following QC

and filtering, our dataset consisted of 4,245,544 SNP genotypes for n=381 total AL lizards, each

with limb and body length phenotypic data available. We tested for genetic associations with

relative limb length separately for each SNP using linear models, with individual sex and the first

four components of a principal components analysis (to account for population structure) as

covariates (see Methods). Using this procedure, we identified a total of n=24 genomic loci with

significant (P < 1e-6) genotype-phenotype associations (Figure 3a; supp table S21; Q-Q plot

available on supplemental figure S22).

A gene ontology enrichment analysis on the set of genes overlapping or nearby (±25Kb)

these regions (supp table S23) revealed an overrepresentation of signaling by epidermal growth

factor receptor pathway genes (REAC:R-MMU-177929; 2 observed vs. 0.03 expected genes;

FDR = 0.049). The two genes, both members of the protein tyrosine phosphatase family, are

PTPN11 on chromosome 7 and *PTPN12* on chromosome 5. In humans, mutations in *PTPN11*

have a strong association with the Noonan syndrome, characterized by short stature and skeletal

malformations (Tartaglia et al. 2002).

We did not observe more overlaps than expected by chance between the GWAS regions

and our AL lizard candidate positive selection regions (permutation analysis; for Tajima's D: P =

0.31; Supp Figure S24; for saltilassi, no overlaps). However, two loci were both strongly

associated with relative limb length and contained or were in proximity to alleles that were

highly differentiated between AL and the fire ant-naïve AR and TN populations. The first locus

overlaps *PTPN11* (chromosome 7), discussed above as part of the enrichment for signaling by

epithelial growth factor receptor (Supp Fig 25). The second locus overlaps *ARHGAP44*

(chromosome 2), which belongs to the gene family of Rho GTPase-activating proteins. These

proteins interact with insulin-like growth factors and the CREB transcription factor to modulate

body size during embryonic development (Moon and Zheng 2003). Here, three non-coding

SNPs, located between 2747 and 3711 bp upstream of *ARHGAP44* exon 1 (Figure 3b), are

strongly associated with relative limb length (β = 0.11, *P* = 2.75e-8; Figure 3c). For these SNPs,

the alleles with positive effects on relative limb length are abundant (>84% frequency) in the AL

population yet absent in TN and AR (Figure 3d). The orthologous region in humans overlaps an

annotated alternative splicing isoform of *ARHGAP44* (AS-1).

Figure 3. Genome-wide genotype associations with fence lizard hind limb length. *A*. Genome-wide associations

with relative hind limb length (residuals of hind limb length on snout-to-vent length with sex and PCs 1-4 from a

Principal Components Analysis of the genome-wide SNP genotype data as covariates). The green line indicates a

significance cutoff of P < 5e-8 (four genomic loci) and the gray line a significance cutoff of P < 1e-6 (24 total loci).

B. GWAS results for SNPs at the *ARHGAP44* gene locus, with vertical bars representing exons, and arrows

representing transcriptional direction. *C*. Representative SNP immediately upstream of *ARHGAP44* and its

association with relative limb length. D. For this same SNP, the negative-effect allele is fixed in the fire ant-free TN

and AR populations, while the limb length positive-effect allele is abundant in the fire ant-invaded AL population.

Discussion

The adaptive potential of natural populations when faced with the sudden and pervasive impacts of anthropogenic activity is uncertain. Understanding these fundamental drivers of ecosystem change is crucial—especially in an era of unprecedented human activity and climate change. Our study focused on endemic fence lizards, who have co-existed with introduced fire ants for more than 70 years in parts of the Southeastern United States. We sampled lizards from a long-invaded site in Alabama and identified multiple genomic signatures of positive selection that were generally absent from two lizard populations naïve to fire ants, in Arkansas and Tennessee.

Putative adaptations to fire ants

The set of genes contained within or nearby the candidate signatures of positive selection that we identified in lizards from the fire ant-invaded AL population were significantly enriched for morphological development functions. Two additional loci – which respectively overlap and are immediately upstream of genes involved in body growth processes – contain alleles that are simultaneously strongly associated with longer legs in AL lizards and occur in regions of high genetic differentiation relative to the fire ant-unexposed TN and AR populations. These variants may constitute some of the genetic adaptations underlying fence lizard limb length variation, with longer hind limbs associated with increased survival against swarming ants (Langkilde 2009).

We also identified multiple signatures of positive selection for variants within or nearby two genes associated with membrane attack complex inflammatory processes (Morgan 2016; Xie et al. 2020). Inflammatory processes are a key component of the immunological response to cellular damage from venom toxins (Stafford 1996; dos Santos Pinto et al. 2012; Zamith-

Miranda et al. 2018; Liu et al. 2023). Remarkably, *S. undulatus* field and laboratory experiments

have previously detected associations between fire ant venom and complement immune activity

(Tylan et al. 2020; Tylan et al. 2023), which includes the membrane attack complex.

Specifically, field-caught lizards from AL had significantly lower levels of complement activity

relative to northern, unexposed lizards (Tylan et al. 2020). Separately, lizards from a TN

population naïve to fire ants showed higher levels of complement activity immediately after

consuming fire ants, as well as three weeks post fire ant stinging (Tylan et al. 2023). Therefore, it

seems likely that the complement immune system is involved in both routes of exposure to fire

ant venom, albeit with temporally distinct signals for each.

Given this context, the findings from our study lead us to suggest that population-level

differences in complement immune activity may in part be a product of recent evolutionary

processes in response to the introduced fire ant. Specifically, we hypothesize that variants within

the *C8A* and *C8B* membrane attack genes candidate selection region that we detected in the fire

ant-exposed AL population but not the fire ant-naïve TN and AR populations may underlie an

adaptive suppression of the innate complement system. If constant consumption of fire ants

triggers chronic, costly inflammatory responses that are of limited immunological benefit, then it

follows that such inflammatory suppression will be sparing of resources otherwise diverted from

other physiological processes (Straub and Schradin 2016). Such a tradeoff between immune

response and energetic resources has been demonstrated in poultry (Van Der Most et al. 2011),

house sparrows (Martin et al. 2003), and the tobacco hornworm (Adamo et al. 2017).

Community-scale consequences

Across ecosystems, evidence for rapid adaptation to human-induced environmental changes

continues to build (Strauss et al. 2006; Shaw and Etterson 2012; Sullivan et al. 2017; McCulloch

and Waters 2023). A few illustrative examples include the water flea *Daphnia* showing signals

of adaptation for resistance to salinization derived from human activity (Wersebe and Weider

2023), and independent populations of crested anoles (*Anolis cristatellus*) carrying signatures of

convergent genetic adaptation to urbanization (Winchell et al. 2023). Meanwhile, human-

commensal house sparrows (*Passer domesticus*) have signatures of positive selection

overlapping an amylase gene, which encodes an enzyme involved in starch digestion (Ravinet et

al. 2018), and poaching of African elephants for ivory appears to have resulted in rapid

adaptation for a tuskless phenotype (Campbell-Staton et al. 2021).

Although the above examples and our results in fence lizards following the introduction

of fire ants contribute to the growing evidence of successful native species adaptations in

response to anthropogenic change the species-specific and community-wide ramifications of

rapid adaptation in a keystone species remain unknown (Thompson 1999; Mooney and Cleland

2001; Strauss et al. 2006). For fence lizards, while fire ant predation is at least partly

counteracted by longer hind limbs combined with a twitch/flee response for fire ant removal and

escape, such behavior is likely accompanied by a break in crypsis. Field surveys in an Alabama

fence lizard population have demonstrated that close to 50% of male lizards show signs of

injuries – a two-fold increase from fire ant-naïve populations – speculatively due to increased

detection by birds of prey (Thawley and Langkilde 2017). This conundrum likely represents a

delicate balance between context-dependent antipredator responses (Martín et al. 2009) faced by

fence lizards since fire ant introduction, exemplifying the broader community ramifications of

adaptation to human activity.

Meanwhile, if it is true that fence lizards from fire ant invaded sites, such as in Alabama,

have adapted to exploit fire ants as a novel food resource, as our results suggest, then this may

have significant cascading effects on that ecosystem's food web. The voracious predatory

behavior of fire ants has been broadly demonstrated; they can effectively prey on higher-level

consumers such as salamanders (Todd et al. 2008), bird nestlings (Kopachena et al. 2000), cotton

rats (Long et al. 2015), and hatchling sea turtles (Allen et al. 2001). Consequently, with potential

adaptations to include fire ants in their dietary niche, fence lizards may indirectly assimilate

biomass otherwise unavailable. The ability to exploit this nutritional input may be one factor

underlying a recent, rapid population expansion of AL fence lizards, evidenced by an abundance

of rare alleles observed in our analysis. Such food web disruptions by invasive species are to be

expected (Zanden et al. 1999; Miehls et al. 2009). However, less discussed is how standing

genetic variation can capitalize on rapid anthropogenic change for an ecological advantage.

In conclusion, our study identifies genetic signatures of positive selection in fence lizards

exposed to human-introduced fire ants. These plausible adaptations to the fire ant introduction

are observed in conjunction with a recent and large-scale population expansion in fence lizards,

inferred from our population genomic data. Together, our findings highlight the potential of

standing genetic variation in promoting population resilience in the face of anthropogenic

disturbance.

Methods

Animal sampling

Between 2006 and 2017, adult male and female lizards were collected from the three study populations using the lasso method. The AL population was included in the study due to its long history with introduced fire ants. The individuals from the TN and AR populations were included because these sampling locations had not yet been invaded by fire ants at the time of collection. Phenotypic data were collected from the n=381 sexually mature lizards in our AL population sample: hind limb length (HLL) and snout-to-vent length (SVL) were each measured to the nearest 0.5 mm using a ruler following a protocol described in (Langkilde 2009) The phenotypic data are available in supp table S26.

DNA Extraction

All the fence lizard tissue toe and/or tail samples were stored in 70% ethanol at 4°C. We used up to 30 mg of each of the 421 preserved fence lizard tissue samples for E.Z.N.A.® tissue kit (D3396, Omega Bio-Tek, Inc., Norcross, GA, USA) DNA extractions. DNA extractions were performed following the manufacturer's instructions with the following exceptions: each tissue sample was ground with a polypropylene pestle in a 1.5-mL microcentrifuge tube, total digestion time was increased to 14-15 hours in a 600 rpm shaking thermomixer, 1 μL of Pellet Paint® NF Co-Precipitant was added to each sample to increase DNA adherence in the HiBind® DNA Mini Column, and the total elution volume was halved (two 50-μL portions). Each sample's DNA extraction concentration was obtained with a Qubit® 3.0 Fluorometer dsDNA High Sensitivity Assay Kit, and then stored at -20°C until library preparation.

Library Preparation and Sequencing

Portions of each DNA extract were sheared to a target length of 500 bp with a Covaris M220

Focused-ultra sonicator (Peak Incident Power: 50, Duty Factor: 20%, Cycles per Burst: 200,

Temperature: 20°C). Libraries for each sample were prepared from ≥200 ng of sheared DNA

with TruSeq® Nano DNA High Throughput Library Prep Kit (20015965, Illumina Inc., San

Diego, CA, USA) and IDT for Illumina – TruSeq® DNA UD Indexes v1 (Illumina Inc., San

Diego, CA, USA). The libraries were pooled and sequenced with a paired-end 150 bp strategy on

two Illumina NovaSeq 6000 S4 flowcells for 1.3 T of paired-end raw read data each. One pool

(HC-60) had 20 randomly selected AL individuals as well as the 20 lizards from the uninvaded

sites TN and AR. An average of 165.24 million reads were generated for each sample in pool

HC-60. The second pool (LC-381) contained the full set of 381 AL individuals, with an average

27.88 million reads sequenced.

Read Mapping and Quality Filtering

We used a chromosome-level reference genome assembly, "PBJelly," that was recently

developed from two male *S. undulatus* individuals collected at Solon Dixon Forestry Education

Center, AL (English et al. 2012; Westfall et al. 2021)for read mapping. The annotated reference

assembly was indexed with bwa v0.7.16 index and SAMtools v1.5 faidx (Li et al. 2009; Li and

Durbin 2009), and a sequence dictionary was created with picard CreateSequenceDictionary

(Picard Toolkit 2019) for use in read mapping, SNP identification, and downstream analyses.

The LC-381 group reads were sequenced without lanes in their NovaSeq S4 flowcell, but

the HC-60 group reads were sequenced across four lanes and needed to be combined into one

forward and reverse read prior to trimming and mapping to the reference genome. The raw reads

were trimmed with Trimmomatic v0.36 to remove the Illumina TruSeq3-PE-2 adapters and other

reads <36 bases long, as well as leading and trailing low quality or N bases (Bolger et al. 2014).

The trimmed reads were aligned to the reference genome with bwa v0.7.16 mem (default

settings), an alignment tool specialized for large genome sizes that seeds alignments with

maximal exact matches and extends seeds with Smith-Waterman's affine-gap penalty for

insertions or deletions (Li 2013).

SAMtools v1.5 flagstat was used to calculate estimated genome-wide coverage for the

mapped reads, and "-view" was used to convert the mapped .sam files to .bam files (Li et al.

2009). SAMtools BAMtools v2.4.1 was used to sort and filter out unmapped reads and mapped

reads with mapQuality less than 50 (Li et al. 2009; Barnett et al. 2011). The subset of AL (Solon

Dixon) samples subject to both high- and low-coverage sequencing (n=20) were merged after

mapping (using samtools -merge) and were processed as part of the LC-381 pool. Duplicates

were marked in all samples (AL, TN, AR) using Picard MarkDuplicates (Picard Toolkit 2019).

Read groups were added to the mapped read files with Picard AddOrReplaceReadGroups (Picard

Toolkit 2019), then the reads were sorted and indexed with SAMtools v1.5 (Li et al. 2009).

Sequence metrics were collected using Picard CollectWgsMetrics (Picard Toolkit 2019). For the

remaining analyses, we found that several of the more computationally intensive programs

required working iteratively at the chromosome- rather than whole-genome level to finish

processing and within the limits postulated by our computational cluster system; we indicate

such cases accordingly.

SNP Identification

We followed the Genome Analysis Toolkit (GATK, v4.2.0.0) "Best Practices" pipeline for

germline short variant discovery in each of the sequencing pools (McKenna et al. 2010; DePristo

et al. 2011; der Auwera et al. 2013). Even though GATK's pipeline was designed and optimized

for analyzing human genetic data, it has been successfully applied in multiple non-model

systems with available high-quality reference genomes for evolutionary genomic inferences

(Kryvokhyzha et al. 2019; Wright et al. 2019; Bernhardsson et al. 2020; Chen et al. 2020; Wang

et al. 2020) and it has been reported to outperform other variant callers in capability and

accuracy (Liu et al. 2013; Pirooznia et al. 2014).

GATK's pipeline began with HaplotypeCaller calling germline SNPs and indels for each

individual via local *de novo* assembly. In short, HaplotypeCaller defined active regions based on

the presence of evidence for allele variation in each individual's mapped reads, then built a De

Bruijn-like graph to detect overlaps between sequences and reassemble the active region (Poplin

et al. 2018). The possible active regions were realigned against the reference haplotype with the

Smith-Waterman algorithm to identify potential variant sites, i.e. single nucleotide

polymorphisms (SNPs; (Poplin et al. 2018). Likelihoods of alleles were determined using

GATK's PairHMM algorithm, and the most likely genotype per Bayes' rule was assigned to

each potentially variant site.

HaplotypeCaller generated an intermediate GVCF file that contained likelihood data for

every position in each of the top 24 largest chromosomes in every individual's mapped read data.

The per-chromosome GVCFs were merged and then indexed then merged GVCF files with

GATK's IndexFeatureFile program. Following the "Best Practices" pipeline,

GenomicsDBImport was used to import the single-sample GVCFs into a per-chromosome

database (GenomicsDB) before joint genotyping with GenotypeGVCFs (per-chromosome,

subset into six intervals each for the largest chromosomes, scaffolds 1-6). The resulting

chromosome VCFs were then combined with VCFtools v0.1.12 "vcf-concat" (Danecek et al.

2011) into one file for each sequencing pool. There were 59,006,281 possible SNP sites

identified in the HC-60 group and 67,124,902 SNP sites in the LC-381 group (bcftools -stats;

bcftools v.1.12; (Danecek et al. 2011)

SNP Filtering and Quality Control

The raw SNPs were filtered with a series of thresholds recommended by GATK (Poplin et al.

2017). GATK SelectVariants kept only variants that were classified as SNPs, then

VariantFiltration removed SNPs with hard-filters based on the INFO and FORMAT fields of the

VCF files: quality score by depth (QD) <2.0, Phred-scaled p-value using Fisher's exact test (FS)

>60.0, and mapping quality score (MQ) < 40.0. SelectVariants was applied again to only keep

SNPs that were not filtered out by VariantFiltration. After GATK filtering there remained

56,598,888 SNPs in the HC-60 group and 64,214,575 SNPs in the LC-381 group.

The SNPs that remained in each pool after GATK's suggested parameters were

additionally filtered with VCFtools v0.1.12 for analysis (Danecek et al. 2011) Both pools were

filtered to keep only biallelic sites (min-alleles 2, max-alleles 2) and remove sites with insertions

and deletions (Danecek et al. 2011). The HC-60 pool was also filtered for Hardy-Weinberg

Equilibrium with a low enough setting to remove sites that were likely to be erroneous variant

calls (hwe 0.000001), leaving 49,837,059 SNPs for analysis (Danecek et al. 2011). PLINK (v1.9)

was also used to filter the HC-60 VCF with -geno 0.05 and -mind 0.1 flags to filter out variants

with missing call rates prior to downstream population analyses, with 46,934,026 SNPs

remaining (Chang et al. 2015). The LC-381 group was also filtered for Hardy-Weinberg

Equilibrium (hwe 0.001) and to remove sites with a minor allele frequency (MAF, number of

times an allele appears over all individuals at that site divided by the total number of non-missing

alleles at that site) less than 0.05 to prevent inflation in downstream statistical estimates and

during imputation with the remaining 458,533 SNPs (Danecek et al. 2011).

To improve genotyping rates for the LC dataset, we leveraged the 20 HC sequences from

AL as a template for genomic imputation. To that end, we first used Shape-IT version 2.r837

(Delaneau et al. 2008) to phase the each of the 24 scaffolds of the 20 HC sequences and obtain

haplotype files. These were used as reference for the imputation of 381 LC sequences.

Imputation was performed with Beagle 5.2 (Browning et al. 2018)using a window size of 100

and overlap of 10. Prior to the genome-wide genotype-phenotype association analysis that was

conducted with the n=381 AL lizard dataset, we removed genotypes with minor allele

frequencies < 0.05, leaving 4,245,544 SNPs.

Demographics analyses

We first calculated the mean genome-wide F_{ST} between each of the three pairs of populations

using the VCFtools function --weir-fst-pop (Danecek et al. 2011). For this analysis, we filtered

the SNP dataset to remove any SNPs whereby an allele that is fixed in one population is also the

minor allele in the second population (for AL/TN, SNP count n = 16,075,146; for AL/AR, n =

33,225,824; for TN/AR, n = 16,383,584). We then ran three demographics analyses involving

the three populations: a principal components analysis (PCA), admixture analysis, and a genetic

neighbor-joining tree based on a genetic distances matrix. For these, the original set of

46,934,026 SNPs was filtered to remove any SNPs whereby an allele that is fixed in two

populations is also the minor allele in the third population, leaving 11,437,455 SNPs. The PCA

was calculated using PLINK's –pca function (Chang et al. 2015). Population structure was

inferred using ADMIXTURE (Alexander et al. 2009) for a number of ancestral populations K of

2 through 5. The neighbor-joining tree was built using the R package phangorn (Schliep 2011)

with a sample genetic distance matrix generated with PLINK (Chang et al. 2015).

Genome-Wide Estimates of Tajima's D

For each population, Tajima's D was estimated in windows across the genomes using vcf-kit

v0.2.6 (Cook and Andersen 2017) Windows were 100 Kb in length with a 20 Kb step. For each

population, putative windows under recent positive selection were those in the bottom 0.5^{th}

percentile of the genome-wide distribution (*i.e.*, D < -1.81 for AL; D < -2.11 for AR; and D < -

1.9 for TN).

Genome-wide Haplotype Distributions via saltilassi statistic

The filtered VCF files were assessed for signatures of positive selection with lassip v1.1.1 using

the "saltiLassi" method (DeGiorgio and Szpiech 2022) The VCF files for each population were

further separated into a single VCF file for each scaffold. A population ID file was created

containing each individual sample ID and the corresponding population IDs. Each scaffold was

passed to lassip with the following parameters to calculate haplotype statistics and the haplotype

frequency spectrum (HFS): --hapstats –winsize 201 –k 20 –calc-spec –winstep 100. The genome

wide average of the HFS, which functions as the null spectrum, was then determined before

calculating the saltilassi statistic, a likelihood ratio statistic denoted as Λ, for each chromosome.

For each population, haplotypes in the top 0.5% of the genome-wide Λ (*i.e.*, Λ > 92.24 for AL;

Λ > 407.56 for AR; and Λ > 252.27 for TN) distribution were designated as candidate signatures

of positive selection.

Locus-specific Branch Lengths (LSBL)

We used LSBL to identify genomic regions in the AL population that are significantly

differentiated from the two northern populations I to fire ants. We first used VCFtools to

calculate the per-SNP F_{ST} of the three pairwise population comparisons, and then used those

values to calculate LSBL for AL as per Shriver *et al.* (2004): LSBL = (AL-AR F_{ST} + AL-TN F_{ST}

– TN-AR F_{ST}). We identified SNPs in the top 0.1% of the genome-wide distribution (LSBL >

0.76) and combined these outlying SNPs into regions that grouped all outlying SNPs within 50

Kb and in high linkage disequilibrium (LD) with each other (R^2 >= 0.9) into a single region. LD

was calculated using vcftools (--geno-r2) (Danecek et al. 2011). Next, we used the R package *ivs*

(Vaughan 2023) to combine linked SNPs into candidate genomic regions.

Functional profiling

For each population, we investigated functional enrichments in the sets of genes located within

or nearby (±25 Kb) candidate regions for positive selection identified by any two of the three

selection statistics (Tajima's D, saltilassi, or LSBL; supp table S12), and separately for Tajima's

D only and saltilassi only, for each population. To that end, we used the g:GOSt function of the

g:Profiler platform (Raudvere et al. 2019) based on functional annotations for *Mus musculus* and

across all available databases (*i.e.*, GO: molecular function, GO: biological process, GO: cellular

component, KEGG, Reactome, WikiPathways, TRANSFAC, miRTarBase, Human Protein

Atlas, CORUM, HP). The background set of genes were all genes in the SceUnd 1.0 assembly

(Westfall et al. 2021). We corrected for multiple tests using the False Discovery Rate (FDR;

Benjamini and Hochberg 1995).

Genome-wide Association Study

Relative limb length was calculated by extracting the residuals of a linear regression for HLL on

SVL. The residuals were then assigned as the response variable for all models. We used the

imputed genotype dataset (see *SNP Filtering and Quality Control* above) for the GWAS. Each

SNP was numerically coded for each biallelic genotype (0, 1, or 2) and included as a predictor in

its respective model. Covariates were individual sex, to control for sex-specific morphological

differences, along with the eigenvectors of a population principal components analysis of the

unimputed LC genomic dataset, to help control for population structure (Supp figure S27).

Principal components 1 through 4 were included in each model, which in combination explained

a total of 23.24% variation explained. Results for each of the 4,245,544 models are available in

supp table S21.

Permutation Analysis

To test whether candidate regions for positive selection and genomic regions associated with

limb length overlapped each other more often than expected by chance, we performed

permutation analyses with the R package *regioneR* (Gel et al. 2016). For each pairwise

permutation (e.g., pairwise comparisons between GWAS and Tajima's D, saltilassi, LSBL, and

the set of regions highlighted by at least two out of the three statistics), the set of regions for one

statistic was held static while regions of same length as those from the second statistic were

randomly placed across the genome in each of 10,000 permutations. To constrain the available

genomic space wherein regions were to be permuted, the number and length of chromosomes

was specified using the argument *genome* in the *permTest* function. These random permutations

generated a neutral probability distribution, and we assessed whether our observed number of

overlaps significantly deviated from this distribution ($\alpha < 0.05$).

Data availability

Raw sequence data have been deposited in NCBI SRA BioProject: PRJNA656311. Quality-

controlled and filtered VCF's and the full outputs for the GWAS and evolutionary genetics

statistical analysis are available on Dryad https://doi.org/10.5061/dryad.tht76hf50. Scripts for all

statistical analyses are available on https://github.com/braulioassis/sce-sol .

Appendix I: Supplemental figures S1-S8.

Appendix II: Supplemental tables S9-S12.

Appendix III: Supplemental tables S13-S20.

Appendix IV: Supplemental tables and figures S21-S27.

Acknowledgements

Animal collections were approved by the Pennsylvania State University Animal Care and Use Committee, and by the respective States where collection occurred. We thank the Lansdale family for allowing us to collect lizard individuals on their property. We also thank T. Schwartz and the *Sceloporus* Genome Collaboration team, T. Adams for assistance in the curation of lizard specimens, and C. Roberts and K. Hickmann for assistance in laboratory work. We thank L. Perez and N. Grube for bioinformatic support, and S. Giery and J. Schluter for comments on the manuscript. We also acknowledge the NYU Langone Genome Technology Center (RRID: SCR_017929). This shared resource is partially supported by the Cancer Center Support Grant P30CA016087 at the Laura and Isaac Perlmutter Cancer Center. Computations for this research were performed on the Pennsylvania State University's Institute for Computational and Data Sciences' Roar supercomputer. Components of this work were supported by the National Science Foundation Graduate Research Fellowship Program (DGE-1255832, to A.P.S.), the National Science Foundation (BCS-1554834, to G.H.P), and the Penn State University College of the Liberal Arts.

References

Adamo SA, Easy RH, Kovalko I, MacDonald J, McKeen A, Swanburg T, Turnbull KF, Reeve C. 2017. Predator exposure-induced immunosuppression: Trade-off, immune redistribution or immune reconfiguration? Journal of Experimental Biology. 220(5):868–875. doi:10.1242/jeb.153320.

Alexander DH, Novembre J, Lange K. 2009. Fast model-based estimation of ancestry in unrelated individuals. Genome Res. 19(9):1655–1664. doi:10.1101/gr.094052.109.

Allen CR, Forys E a., Rice KG, Wojcik DP. 2001. Effects of Fire Ants (Hymenoptera: Formicidae) on Hatching Turtles and Prevalence of Fire Ants on Sea Turtle Nesting Beaches in Florida. Florida Entomologist. 84(2):250–253.

Amos W, Harwood J. 1998. Factors affecting levels of genetic diversity in natural populations.
Philos Trans R Soc Lond B Biol Sci. 353(1366):177–186. doi:10.1098/rstb.1998.0200.

Ascunce MS, Yang CC, Oakey J, Calcaterra L, Wu WJ, Shih CJ, Goudet J, Ross KG, Shoemaker
DW. 2011. Global invasion history of the fire ant Solenopsis invicta. Science (1979).
331(6020):1066–1068. doi:10.1126/science.1198734.

der Auwera GA, Carneiro MO, Hartl C, Poplin R, del Angel G, Levy-Moonshine A, Jordan T,
Shakir K, Roazen D, Thibault J, et al. 2013. From FastQ Data to High-Confidence Variant Calls:
The Genome Analysis Toolkit Best Practices Pipeline. Curr Protoc Bioinformatics.
43(1):11.10.1-11.10.33. doi:https://doi.org/10.1002/0471250953.bi1110s43.

Barnett DW, Garrison EK, Quinlan AR, Strömberg MP, Marth GT. 2011. BamTools: a C++ API
and toolkit for analyzing and managing BAM files. Bioinformatics. 27(12):1691–1692.
doi:10.1093/bioinformatics/btr174.

Barrett RDH, Schluter D. 2008. Adaptation from standing genetic variation. Trends Ecol Evol.
23(1):38–44. doi:10.1016/j.tree.2007.09.008.

Bastian FB, Roux J, Niknejad A, Comte A, Fonseca Costa SS, de Farias TM, Moretti S,
Parmentier G, de Laval VR, Rosikiewicz M, et al. 2021. The Bgee suite: integrated curated
expression atlas and comparative transcriptomics in animals. Nucleic Acids Res. 49(D1):D831–
D847. doi:10.1093/nar/gkaa793.

Benjamini Y, Hochberg Y. 1995. Controlling the False Discovery Rate: A Practical and
Powerful Approach to Multiple Testing. Journal of the Royal Statistical Society: Series B
(Methodological). 57(1):289–300. doi:https://doi.org/10.1111/j.2517-6161.1995.tb02031.x.

Bernhardsson C, Wang X, Eklöf H, Ingvarsson PK. 2020. Variant calling using NGS and
sequence capture data for population and evolutionary genomic inferences in Norway Spruce
(Picea abies). bioRxiv. doi:10.1101/805994.

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: A flexible trimmer for Illumina sequence
data. Bioinformatics. 30(15):2114–2120. doi:10.1093/bioinformatics/btu170.

Browning BL, Zhou Y, Browning SR. 2018. A One-Penny Imputed Genome from Next-
Generation Reference Panels. The American Journal of Human Genetics. 103(3):338–348.
doi:10.1016/j.ajhg.2018.07.015.

Campbell-Staton SC, Arnold BJ, Gonçalves D, Granli P, Poole J, Long RA, Pringle RM. 2021.
Ivory poaching and the rapid evolution of tusklessness in African elephants. Science (1979).
374(6566):483–487. doi:10.1126/science.abe7389.

Chang CC, Chow CC, Tellier LCAM, Vattikuti S, Purcell SM, Lee JJ. 2015. Second-generation
PLINK: Rising to the challenge of larger and richer datasets. Gigascience. 4(1):1–16.
doi:10.1186/s13742-015-0047-8.

Chen Q, Huang B, Zhan J, Wang J, Qu K, Zhang F, Shen J, Jia P, Ning Q, Zhang J, et al. 2020.
Whole-genome analyses identify loci and selective signals associated with body size in cattle. J
Anim Sci. 98(3). doi:10.1093/jas/skaa068.

Clavero M, García-Berthou E. 2005. Invasive species are a leading cause of animal extinctions.
Trends Ecol Evol. 20(3):110. doi:10.1016/j.tree.2005.01.003.

Code of Federal Regulations. 2006. Subpart-imported fre ant. http://ecfr.gpoacces.gov.

Cook DE, Andersen EC. 2017. VCF-kit: assorted utilities for the variant call format.
Bioinformatics. 33(10):1581–1582. doi:10.1093/bioinformatics/btx011.

Csapo R, Gumpenberger M, Wessner B. 2020. Skeletal Muscle Extracellular Matrix – What Do
We Know About Its Composition, Regulation, and Physiological Roles? A Narrative Review.
Front Physiol. 11(March):1–15. doi:10.3389/fphys.2020.00253.

Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter G,
Marth GT, Sherry ST, et al. 2011. The variant call format and VCFtools. Bioinformatics.
27(15):2156–2158. doi:10.1093/bioinformatics/btr330.

Darracq AK, Smith LL, Oi DH, Conner LM, McCleery RA. 2017. Invasive ants influence native
lizard populations. Ecosphere. 8(1). doi:10.1002/ecs2.1657.

DeGiorgio M, Szpiech ZA. 2022. A spatially aware likelihood test to detect sweeps from
haplotype distributions.

Delaneau O, Coulonges C, Zagury J-F. 2008. Shape-IT: new rapid and accurate algorithm for
haplotype inference. BMC Bioinformatics. 9(1):540. doi:10.1186/1471-2105-9-540.

DePristo MA, Banks E, Poplin R, Garimella K V, Maguire JR, Hartl C, Philippakis AA, del
Angel G, Rivas MA, Hanna M, et al. 2011. A framework for variation discovery and genotyping
using next-generation DNA sequencing data. Nat Genet. 43(5):491–498. doi:10.1038/ng.806.

English AC, Richards S, Han Y, Wang M, Vee V, Qu J, Qin X, Muzny DM, Reid JG, Worley
KC, et al. 2012. Mind the Gap: Upgrading Genomes with Pacific Biosciences RS Long-Read
Sequencing Technology. PLoS One. 7(11):1–12. doi:10.1371/journal.pone.0047768.

Foth BJ, Goedecke MC, Soldati D. 2006. New insights into myosin evolution and classification.
Proc Natl Acad Sci U S A. 103(10):3681–3686. doi:10.1073/pnas.0506307103.

Gel B, Díez-Villanueva A, Serra E, Buschbeck M, Peinado MA, Malinverni R. 2016. regioneR:
an R/Bioconductor package for the association analysis of genomic regions based on permutation
tests. Bioinformatics. 32(2):289–291. doi:10.1093/bioinformatics/btv562.

Gifford ME, Robinson CD, Clay TA. 2017. The influence of invasive fire ants on survival, space
use, and patterns of natural selection in juvenile lizards. Biol Invasions. 19(5):1461–1469.
doi:10.1007/s10530-017-1370-z.

Gruber MAM, Santoro D, Cooling M, Lester PJ, Hoffmann BD, Boser C, Lach L. 2022. A
global review of socioeconomic and environmental impacts of ants reveals new insights for risk
assessment. Ecological Applications. 32(4):1–17. doi:10.1002/eap.2577.

Hale R, Morrongiello JR, Swearer SE. 2016. Evolutionary traps and range shifts in a rapidly
changing world. Biol Lett. 12(6). doi:10.1098/rsbl.2016.0003.

Holt RIG. 2002. Fetal programming of the growth hormone–insulin-like growth factor axis.
Trends in Endocrinology & Metabolism. 13(9):392–397. doi:https://doi.org/10.1016/S1043-
2760(02)00697-5.

Kolbe JJ, Losos JB. 2005. Hind-Limb Length Plasticity in Anolis carolinensis Published By :
The Society for the Study of Amphibians and Reptiles Your use of this PDF , the BioOne Web
site , and all posted and associated content. 39(4):674–678.

Kopachena JG, Buckley A, Potts GA. 2000. Effects of the Red Imported Fire Ant (Solenopsis
invicta) on Reproductive Success of Barn Swallows (Hirundo rustica) in Northeast. Southwest
Nat. 45(4):477–482.

Koskinen SOA, Wang W, Ahtikoski AM, Kjær M, Han XY, Komulainen J, Kovanen V, Takala
TES. 2001. Acute exercise induced changes in rat skeletal muscle mRNAs and proteins
regulating type IV collagen content. Am J Physiol Regul Integr Comp Physiol. 280(5 49-
5):1292–1300. doi:10.1152/ajpregu.2001.280.5.r1292.

Kryvokhyzha D, Salcedo A, Eriksson MC, Duan T, Tawari N, Chen J, Guerrina M, Kreiner JM,
Kent T V, Lagercrantz U, et al. 2019. Parental legacy, demography, and admixture influenced
the evolution of the two subgenomes of the tetraploid Capsella bursa-pastoris (Brassicaceae).
PLoS Genet. 15(2):1–34. doi:10.1371/journal.pgen.1007949.

Langkilde T. 2009. Invasive fire ants alter behavior and morphology of native lizards. Ecology.
90(1):208–217. doi:10.1890/08-0355.1.

Langkilde T, Freidenfelds NA. 2010. Consequences of envenomation: Red imported fire ants
have delayed effects on survival but not growth of native fence lizards. Wildlife Research.
37:566–573. doi:10.1071/WR10098.

Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform.
Bioinformatics. 25(14):1754–1760. doi:10.1093/bioinformatics/btp324.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R.
2009. The Sequence Alignment/Map format and SAMtools. Bioinformatics. 25(16):2078–2079.
doi:10.1093/bioinformatics/btp352.

Ligon RA, Siefferman L, Hill GE. 2011. Invasive fire ants reduce reproductive success and alter
the reproductive strategies of a native vertebrate insectivore. PLoS One. 6(7).
doi:10.1371/journal.pone.0022578.

Liu X, Han S, Wang Z, Gelernter J, Yang B-Z. 2013. Variant Callers for Next-Generation
Sequencing Data: A Comparison Study. PLoS One. 8(9):null.
doi:10.1371/journal.pone.0075619.

Liu Y, Huang J, Zhang J, Xu Y, Li X, Lu Y. 2023. Sensitization of Guinea Pig Skin to Imported
Fire Ant Alkaloids and Establishment of an Inflammatory Model. Int J Environ Res Public
Health. 20(3). doi:10.3390/ijerph20031904.

Long AK, Conner LM, Smith LL, Mccleery RA. 2015. Effects of an invasive ant and native
predators on cotton rat recruitment and survival. J Mammal. 96(6):1135–1141.
doi:10.1093/jmammal/gyv121.

Martín J, Luque-Larena JJ, López P. 2009. When to run from an ambush predator: balancing
crypsis benefits with costs of fleeing in lizards. Anim Behav. 78(4):1011–1018.
doi:10.1016/j.anbehav.2009.07.026.

Martin LB, Scheuerlein A, Wikelski M. 2003. Immune activity elevates energy expenditure of
house sparrows: A link between direct and indirect costs? Proceedings of the Royal Society B:
Biological Sciences. 270(1511):153–158. doi:10.1098/rspb.2002.2185.

McCulloch GA, Waters JM. 2023. Rapid adaptation in a fast-changing world: Emerging insights
from insect genomics. Glob Chang Biol. 29(4):943–954. doi:https://doi.org/10.1111/gcb.16512.

McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, Garimella K,
Altshuler D, Gabriel S, Daly M, et al. 2010. The Genome Analysis Toolkit: a MapReduce
framework for analyzing next-generation DNA sequencing data. Genome Res. 20(9):1297–
1303. doi:10.1101/gr.107524.110.

Miehls ALJ, Mason DM, Frank KA, Krause AE, Peacor SD, Taylor WW. 2009. Invasive species
impacts on ecosystem structure and function: A comparison of Oneida Lake, New York, USA,
before and after zebra mussel invasion. Ecol Modell. 220(22):3194–3209.
doi:10.1016/j.ecolmodel.2009.07.020.

Moon SY, Zheng Y. 2003. Rho GTPase-activating proteins in cell regulation. Trends Cell Biol.
13(1):13–22. doi:10.1016/S0962-8924(02)00004-1.

Mooney HA, Cleland EE. 2001. The evolutionary impact of invasive species. Proc Natl Acad Sci
U S A. 98(10):5446–5451. doi:10.1073/pnas.091093398.

Morgan BP. 2016. The membrane attack complex as an inflammatory trigger. Immunobiology.
221(6):747–751. doi:10.1016/j.imbio.2015.04.006.

Morrison LW. 2002. Long-term impacts of an arthropod-community invasion by the imported
fire ant, *Solenopsis invicta*. Ecology. 83(8):2337–2345.

Van Der Most PJ, De Jong B, Parmentier HK, Verhulst S. 2011. Trade-off between growth and
immune function: A meta-analysis of selection experiments. Funct Ecol. 25(1):74–80.
doi:10.1111/j.1365-2435.2010.01800.x.

Pirooznia M, Kramer M, Parla J, Goes FS, Potash JB, McCombie WR, Zandi PP. 2014.
Validation and assessment of variant calling pipelines for next-generation sequencing. Hum
Genomics. 8(1):14. doi:10.1186/1479-7364-8-14.

Poplin R, Ruano-Rubio V, DePristo MA, Fennell TJ, Carneiro MO, der Auwera GA Van, Kling
DE, Gauthier LD, Levy-Moonshine A, Roazen D, et al. 2018. Scaling accurate genetic variant
discovery to tens of thousands of samples. bioRxiv. doi:10.1101/201178.

Porter SD, Savignano DA. 1990. Invasion of polygyne fire ants decimates native ants and
disrupts arthropod community. Ecology. 71(6):2095–2106. doi:10.2307/1938623.

Raudvere U, Kolberg L, Kuzmin I, Arak T, Adler P, Peterson H, Vilo J. 2019. G:Profiler: A web
server for functional enrichment analysis and conversions of gene lists. Nucleic Acids Res.
47(W1):W191–W198. doi:10.1093/nar/gkz369.

Ravinet M, Elgvin TO, Trier C, Aliabadian M, Gavrilov A, Sætre GP. 2018. Signatures of
human-commensalism in the house sparrow genome. Proceedings of the Royal Society B:
Biological Sciences. 285(1884). doi:10.1098/rspb.2018.1246.

Robbins TR, Freidenfelds NA, Langkilde T. 2013. Native predator eats invasive toxic prey:
Evidence for increased incidence of consumption rather than aversion-learning. Biol Invasions.
15(2):407–415. doi:10.1007/s10530-012-0295-9.

Rodríguez-Romero F, Smith GR, Méndez-Sánchez F, Hernández-Gallegos O, Nava PS, Méndez
De La Cruz FR. 2011. Demography of a semelparous, high-elevation population of Sceloporus
bicanthalis (Lacertilia: Phrynosomatidae) from the Nevado de Toluca Volcano, Mexico.
Southwestern Naturalist. 56(1):71–77. doi:10.1894/GC-193.1.

Roeder KA, Penuela Useche V, Levey DJ, Resasco J. 2021. Testing effects of invasive fire ants
and disturbance on ant communities of the longleaf pine ecosystem. Ecol Entomol. 46(4):964–
972. doi:10.1111/een.13033.

dos Santos Pinto JRA, Fox EGP, Saidemberg DM, Santos LD, da Silva Menegasso AR, Costa-
Manso E, Machado EA, Bueno OC, Palma MS. 2012. Proteomic View of the Venom from the
Fire Ant Solenopsis invicta Buren. J Proteome Res. 11(9):4643–4653. doi:10.1021/pr300451g.

Saul WC, Jeschke JM. 2015. Eco-evolutionary experience in novel species interactions. Ecol
Lett. 18(3):236–245. doi:10.1111/ele.12408.

Schlaepfer MA, Sherman PW, Blossey B, Runge MC. 2005. Introduced species as evolutionary
traps. Ecol Lett. 8(3):241–246. doi:10.1111/j.1461-0248.2005.00730.x.

Schliep KP. 2011. phangorn: phylogenetic analysis in R. Bioinformatics. 27(4):592–593.
doi:10.1093/bioinformatics/btq706.

Shaw RG, Etterson JR. 2012. Rapid climate change and the rate of adaptation: Insight from
experimental quantitative genetics. New Phytologist. 195(4):752–765. doi:10.1111/j.1469-
8137.2012.04230.x.

Shriver MD, Kennedy GC, Parra EJ, Lawson HA, Sonpar V, Huang J, Akey JM, Jones KW.
2004. The genomic distribution of population substructure in four populations using 8,525
autosomal SNPs. Hum Genomics. 1(4):274–286.

Slatkin M. 1993. Isolation by distance in equilibrium and non-equilibrium populations. Evolution
(N Y). 47(1):264–279. doi:10.1111/j.1558-5646.1993.tb01215.x.

Stafford CT. 1996. Hypersensitivity to Fire Ant Venom. Annals of Allergy, Asthma &
Immunology. 77(2):87–99. doi:https://doi.org/10.1016/S1081-1206(10)63493-X.

Stahlschmidt ZR, Walman RM, Mills AM. 2018. Red imported fire ants (Solenopsis invicta) and
seasonality influence community refuge use. Biol Invasions. 20(10):2849–2859.
doi:10.1007/s10530-018-1737-9.

Straub RH, Schradin C. 2016. Chronic inflammatory systemic diseases – an evolutionary trade-
off between acutely beneficial but chronically harmful programs. Evol Med Public
Health.:eow001. doi:10.1093/emph/eow001.

Strauss SY, Lau JA, Carroll SP. 2006. Evolutionary responses of natives to introduced species:
What do introductions tell us about natural communities? Ecol Lett. 9(3):357–374.
doi:10.1111/j.1461-0248.2005.00874.x.

Sullivan AP, Bird DW, Perry GH. 2017. Human behaviour as a long-term ecological driver of
non-human evolution. Nat Ecol Evol. 1(3):65. doi:10.1038/s41559-016-0065.

Tajima F. 1989. Statistical method for testing the neutral mutation hypothesis by DNA
polymorphism. Genetics. 123:585–595.

Tartaglia M, Kalidas K, Shaw A, Song X, Musat DL, Van der Burgt I, Brunner HG, Bertola DR,
Crosby A, Ion A, et al. 2002. PTPN11 mutations in noonan syndrome: Molecular spectrum,
genotype-phenotype correlation, and phenotypic heterogeneity. Am J Hum Genet. 70(6):1555–
1563. doi:10.1086/340847.

Thawley CJ, Goldy-Brown M, McCormick GL, Graham SP, Langkilde T. 2019. Presence of an
invasive species reverses latitudinal clines of multiple traits in a native species. Glob Chang Biol.
25(2):620–628. doi:10.1111/gcb.14510.

Thawley CJ, Langkilde T. 2016. Invasive fire ant (Solenopsis invicta) predation of eastern fence
lizard (Sceloporus undulatus) eggs. J Herpetol. 50(2):284–288. doi:10.1670/15-017.

Thawley CJ, Langkilde T. 2017. Attracting unwanted attention: generalization of behavioural
adaptation to an invasive predator carries costs. Anim Behav. 123:285–291.
doi:10.1016/j.anbehav.2016.11.006.

Thompson JN. 1999. Specific hypotheses on the geographic mosaic of coevolution. American
Naturalist. 153(SUPPL.):1–14. doi:10.1086/303208.

Todd BD, Rothermel BB, Reed RN, Luhring TM, Schlatter K, Trenkamp L, Gibbons JW. 2008.
Habitat alteration increases invasive fire ant abundance to the detriment of amphibians and
reptiles. Biol Invasions. 10(4):539–546. doi:10.1007/s10530-007-9150-9.

Tsuji JS, Huey RB, van Berkum FH, Garland T, Shaw RG. 1989. Locomotor performance of
hatchling fence lizards (Sceloporus occidentalis): Quantitative genetics and morphometric
correlates. Evol Ecol. 3(3):240–252. doi:10.1007/BF02270725.

Tylan C, Engler HI, Villar G, Langkilde T. 2023. Consumption of fire ants, an invasive predator
and prey of native lizards, may enhance immune functions needed to combat envenomation. Biol
Invasions. 25(3):725–740. doi:10.1007/s10530-022-02939-8.

Tylan C, Horvat-Gordon M, Bartell PA, Langkilde T. 2020. Ecoimmune reallocation in a native
lizard in response to the presence of invasive, venomous fire ants in their shared environment. J
Exp Zool A Ecol Integr Physiol. 333(10):792–804. doi:10.1002/jez.2418.

Vaughan D. 2023. ivs: Interval Vectors.

Wang H, Yoshiko Y, Yamamoto R, Minamizaki T, Kozai K, Tanne K, Aubin JE, Maeda N.
2008. Overexpression of Fibroblast Growth Factor 23 Suppresses Osteoblast Differentiation and
Matrix Mineralization In Vitro. Journal of Bone and Mineral Research. 23(6):939–948.
doi:https://doi.org/10.1359/jbmr.080220.

Wang J-J, Zhang T, Chen Q-M, Zhang R-Q, Li L, Cheng S-F, Shen W, Lei C-Z. 2020. Genomic
Signatures of Selection Associated With Litter Size Trait in Jining Gray Goat. Front Genet. 11.
doi:10.3389/fgene.2020.00286.

Weiss A, Leinwand LA. 1996. The mammalian myosin heavy chain gene family. Annu Rev Cell
Dev Biol. 12:417–439. doi:10.1146/annurev.cellbio.12.1.417.

Wersebe MJ, Weider LJ. 2023. Resurrection genomics provides molecular and phenotypic
evidence of rapid adaptation to salinization in a keystone aquatic species. Proceedings of the
National Academy of Sciences. 120(6). doi:https://doi.org/10.1073/pnas.2217276120.

Westfall AK, Telemeco RS, Grizante MB, Waits DS, Clark AD, Simpson DY, Klabacka RL,
Sullivan AP, Perry GH, Sears MW, et al. 2021. A chromosome-level genome assembly for the
eastern fence lizard (Sceloporus undulatus), a reptile model for physiological and evolutionary
ecology. Gigascience. 10(10):1–17. doi:10.1093/gigascience/giab066.

Winchell KM, Campbell-Staton SC, Losos JB, Revell LJ, Verrelli BC, Geneva AJ. 2023.
Genome-wide parallelism underlies contemporary adaptation in urban lizards. Proc Natl Acad
Sci U S A. 120(3). doi:10.1073/pnas.2216789120.

Wright B, Farquharson KA, McLennan EA, Belov K, Hogg CJ, Grueber CE. 2019. From
reference genomes to population genomics: comparing three reference-aligned reduced-
representation sequencing pipelines in two wildlife species. BMC Genomics. 20(1):453.
doi:10.1186/s12864-019-5806-y.

Xie CB, Jane-Wit D, Pober JS. 2020. Complement Membrane Attack Complex: New Roles,
Mechanisms of Action, and Therapeutic Targets. American Journal of Pathology. 190(6):1138–
1150. doi:10.1016/j.ajpath.2020.02.006.

Zamith-Miranda D, Fox EGP, Monteiro AP, Gama D, Poublan LE, de Araujo AF, Araujo MFC,
Atella GC, Machado EA, Diaz BL. 2018. The allergic response mediated by fire ant venom
proteins. Sci Rep. 8(1):14427. doi:10.1038/s41598-018-32327-z.

Zanden MJ Vander, Casselman JM, Rasmussen JB. 1999. Stable isotope evidence for the food
web consequences of species invasions in lakes. Nature. 401(September):1997–2000.

890