Transposition of *HOPPLA* in siRNA-deficient plants suggests a limited effect of the environment on retrotransposon mobility in *Brachypodium distachyon*

Michael Thieme[1*], Nikolaos Minadakis[1], Christophe Himber[2], Bettina Keller[1], Wenbo Xu[1], Kinga Rutowicz[1], Calvin Matteoli[2], Marcel Böhrer[2], Bart Rymen[2], Debbie Laudencia-Chingcuanco[3], John Vogel[4], Richard Sibout[5], Christoph Stritt[6], Todd Blevins[2] and Anne C. Roulin[1*]

[1] Department of Plant and Microbial Biology, University of Zurich, Zurich, Switzerland.

[2] Institut de Biologie Moléculaire des Plantes, CNRS, Université de Strasbourg, Strasbourg, France.

[3] USDA ARS Western Regional Research Center, Albany, CA, USA.

[4] DOE Joint Genome Institute, Lawrence Berkeley National Laboratory, Berkeley, CA, USA.

[5] INRAE Unité BIA - 1268 Biopolymères Interactions Assemblages Equipe Paroi Végétale et Polymères Pariétaux (PVPP), Nantes, France.

[6] Swiss Tropical and Public Health Institute (Swiss TPH), Allschwil, Switzerland.

* Corresponding authors: E-Mail

michael.thieme@botinst.uzh.ch (MT), anne.roulin@botinst.uzh.ch (ACR)

Abstract

Long terminal repeat retrotransposons (LTR-RTs) are powerful mutagens regarded as a major source of genetic novelty and important drivers of evolution. Yet, the uncontrolled and potentially selfish proliferation of LTR-RTs can lead to deleterious mutations and genome instability, with large fitness costs for their host. While population genomics data suggest that an ongoing LTR-RT mobility is common in many species, the understanding of their dual roles in evolution is limited. Here, we harness the genetic diversity of 320 sequenced natural accessions of the Mediterranean grass *Brachypodium distachyon* to characterize how genetic and environmental factors influence plant LTR-RT dynamics in the wild. When combining a coverage-based approach to estimate global LTR-RT copy number variations with mobilome-sequencing of nine accessions exposed to eight different stresses, we find little evidence for a major role of environmental factors in LTR-RT accumulations in *B. distachyon* natural accessions. Instead, we show that loss of RNA polymerase IV (Pol IV), which mediates RNA-directed DNA methylation in plants, results in high transcriptional and transposition activities of RLC_BdisC024 (*HOPPLA*) LTR-RT family elements, and that these effects are not stress-specific. This work supports findings indicating an ongoing mobility in *B. distachyon* and reveals that host RNA-directed DNA methylation rather than environmental factors controls their mobility in this wild grass model.

Introduction

Transposable elements (TEs) are DNA sequences with the ability to form extrachromosomal copies and to reintegrate elsewhere into the host genome. In plants, TE-derived sequences are ubiquitous

and can constitute more than 80 % of the genome (Mirouze & Vitte, 2014). In addition to playing a major role in genome size variation (e.g. Hawkins et al., 2006; Piegu et al., 2006; Wang et al., 2021;

Yang et al., 2023), TEs can alter gene expression by acting as promotors or by providing *cis*-regulatory elements to flanking regions (Butelli *et al.*, 2012; Makarevitch *et al.*, 2015; Roquis *et al.*, 2021). TEs

are therefore a major source of genetic change. Given that they are more likely than classic point mutations to cause extreme changes in gene expression and phenotypes (Uzunović *et al.*, 2019;

Thieme *et al.*, 2022; Raúl *et al.*, 2023), they might be especially useful when the survival of an organism or its descendants depends on a quick response to new or challenging environmental

conditions (for review Kawakatsu et al., 2016; Rey et al., 2016; Lanciano & Mirouze, 2018; Dubin et al., 2018). Paradoxically, while population genomics data have revealed ongoing TE activity in natural

plant populations (e.g. (Stuart *et al.*, 2016; Stritt *et al.*, 2018; Baduel *et al.*, 2021), only a handful of TE families have been reported to transpose in real-time. Therefore, how often or under which

natural conditions TEs are activated in the wild remain open questions. In addition, while ongoing transposition is essential for TEs to survive, the presence of mobile and potentially 'selfish' DNA

sequences requires the host to evolve robust silencing mechanisms to prevent an uncontrolled TE proliferation. TE activity thus remains a major puzzle in the field of evolutionary genomics.

In plants, the defence against TEs is multi-layered, comprising repressive histone modifications, DNA methylation and RNA interference (Saze *et al.*, 2012; Zhang *et al.*, 2018; Liu *et al.*, 2022). One

of the main players of TE silencing is the RNA-directed DNA methylation (RdDM) pathway, which involves two plant specific RNA-polymerases derived from Pol II, namely Pol IV and Pol V. The largest

subunits of each polymerase (NRPB1, NRPD1 and NRPE1, respectively) assemble with other proteins

into enzymes with distinct RNA products and functions (Ream et al., 2009; for review Rymen et al.,

62   2020). As a core component of RdDM, Pol IV (including NRPD1) transcribes TE regions into the

precursors of functionally specialized 24 nt small interfering RNAs (siRNAs) (Sigman & Slotkin, 2016;

64   Rymen *et al.*, 2020; Liu *et al.*, 2022). Upon the base pairing of 24 nt siRNAs to scaffold transcripts

produced by Pol V, the *de novo* DNA methyltransferase DRM2 (Zhong *et al.*, 2014) is recruited to

66   mediate the methylation and subsequent transcriptional repression of TEs. The essential role of

RdDM in TE silencing has been shown in *A. thaliana*, where the knockout of Pol IV and resulting

68   depletion of 24 nt siRNAs leads to a drastically increased heat-dependent transposition of the

ONSEN family (Tittel-Elmer *et al.*, 2010; Ito *et al.*, 2011).

70   The case of the heat-responsive *ONSEN* elements not only illustrates the importance of

epigenetic silencing in regulating TEs but also demonstrates that environmental factors may

72   modulate the dynamics of TEs in plants. Since their discovery by Barbara McClintock, who linked the

mobility of AC/DS elements in maize to the occurrence of a 'genomic shock' (McClintock, 1984), the

74   activity of TEs has been frequently associated with the presence of biotic and abiotic stressors. In

fact, certain TEs can sense specific physiological states of their host and use them to initiate their

76   own life cycle (Negi *et al.*, 2016). Besides *ONSEN* in *A. thaliana* (Cavrak *et al.*, 2014), the cold

inducible *Tcs1* element in blood oranges (Butelli *et al.*, 2012) or *Tos17* that gets activated during

78   tissue culture in rice (Hirochika *et al.*, 1996) are two other prominent cases of stress responsive TEs

in plants. Mechanistically, stress can activate TEs via specific motifs allowing the binding of

80   transcription factors and the subsequent recruitment of the transcription machinery to their

promoter-like sequences (Cavrak *et al.*, 2014; Grandbastien, 2015; Baduel *et al.*, 2021; Zhang *et al.*,

82   2022). The small window of increased activity during well-defined physiological states suggests that

some TEs have evolved a distinct lifestyle or 'niche' to successfully reproduce (Kidwell & Lisch, 1997;

84   Venner *et al.*, 2009; Stritt *et al.*, 2021; Stitzer *et al.*, 2021).

Since transcription constitutes the initial step to transposition, the abundance of TE

86   transcripts is often used as a proxy for TE activity (Lanciano & Cristofari, 2020). However, the life

cycle of TEs is complex (Schulman, 2013) and the fate of TE transcripts depends on many factors. For

88   instance, several transcriptionally active TEs have accumulated mutations that prevent the

production of enzymes needed for their autonomous transposition (Tanskanen *et al.*, 2007). To

90   selectively capture TEs that are not only transcriptionally active but also capable of transposing,

several protocols have been developed, including ALE-seq (Cho *et al.*, 2019), VLP-seq (Lee *et al.*,

92   2020) and mobilome-seq (Lanciano *et al.*, 2017). These recent approaches have been particularly

successful when targeting long terminal repeat retrotransposons (LTR-RTs), which represent the

94   largest fraction of TE-derived sequences in plant genomes (Vitte *et al.*, 2014). Indeed, LTR-RTs

transpose through a copy-and-paste mechanism which involves the reverse transcription of a full-

96   length RNA intermediate (Wicker *et al.*, 2007; Schulman, 2013). As part of their life cycle and

presumably through auto-integration, non-homologous and alternative end-joining, active LTR-RTs

98   can also form extrachromosomal circular DNA (eccDNA) intermediates (Flavell & Ish-horowicz, 1981;

Flavell, 1984; Garfinkel *et al.*, 2006; Wicker *et al.*, 2007; Lee *et al.*, 2020; Yang *et al.*, 2023a), whose

100   detection by mobilome-seq can be used as a proxy for their mobility (Lanciano *et al.*, 2017). For

instance, mobilome-seq has been successfully used to track full-length eccDNA of mobilized

102   autonomous RTs, containing both LTRs (2-LTR circles), in plants such as *A. thaliana* and rice

(Lanciano *et al.*, 2017; Thieme *et al.*, 2017).

104   While the activity of LTR-RTs has been extensively studied in the model *A. thaliana* (Baduel *et*

al., 2021), the interplay between genetic and environmental factors in other wild plant species

106   remains poorly investigated. To clarify these questions, we exploit here the *Brachypodium*

distachyon diversity panel (Minadakis *et al.*, 2023a) and explore LTR-RT activity in a wild monocot.

108   *B. distachyon* is a Mediterranean grass with a compact diploid genome of ~272 Mb (International

Brachypodium Initiative, 2010; Hasterok *et al.*, 2022) harboring 40 LTR-RT families (Stritt *et al.*, 2020)

110   that constitute about 30 % of the genome (International Brachypodium Initiative, 2010). In

B. distachyon, LTR-RTs not only evolved varying insertion site preferences (Stritt *et al.*, 2020) but also

112   significantly differ in terms of transposition dynamics (Stritt *et al.*, 2018, 2020). While we previously

suggested an ongoing transposition of LTR-RTs based on population genomics data (Stritt *et al.*, 2018;

114   Horvath *et al.*, 2023), here we aimed to clarify the contribution of genetic and environmental factors

to LTR-RT activity in *B. distachyon*. To that end, we combined population genomics data available for

116   320 natural accessions with mobilome-seq under different stress conditions and asked: (i) does the

accumulation of LTR-RTs in these natural accessions correlate with environmental variables, (ii) is

118   LTR-RT mobility induced by specific stresses, and (iii) which genetic factors influence the

accumulation of LTR-RTs ?

Results

122   **Abundances of LTR-RT families differ but show limited association with bioclimatic variables**

B. distachyon naturally occurs around the Mediterranean rim (**Fig 1A**) and groups into three main

124   genetic lineages (A, B and C) that further split into five genetic clades (Stritt *et al.*, 2022; Minadakis

et al., 2023a): an early diverged C clade and four clades found in Spain and France (B_West), Italy

126 (A_Italia), the Balkans and coastal Turkey (A_East) and inland Turkey and Lesser Caucasus (B_East).

We first aligned genomic reads of 320 accessions to the LTR-RT consensus sequences of

128 *B. distachyon* obtained from the TRansposable Elements Platform (TREP). We then computed the

abundance of TE-derived sequences, hence a proxy for copy number variation (pCNV), for the 40

130 annotated LTR-RT families using a coverage-based approach accounting for sample sequencing

depth (see Materials and Methods). We favored this approach over an analysis based on transposon

132 insertion polymorphisms (TIPs) because estimates based on TIPs are reference genome-dependent

and biased by the phylogeny in our study system. We have, for instance, previously shown that

134 accessions from the B_East clade harbor significantly less TIPs than accessions from the A_East clade

due to the fact that the reference genome Bd21 belongs the B_East clade (Stritt *et al.*, 2018). In

136 addition, whole-genome *de novo* assembly of 54 *B. distachyon* natural accessions and the

subsequent pangenome analysis revealed that non-reference accessions display large genomic

138 variations (Gordon *et al.*, 2017), which may further bias the estimates of TIP abundances.

The heatmap produced based on pCNV (**Fig 1B**) showed that LTR-RTs underwent different

140 transposon accumulations. We found that *Gypsy* elements (RLG) harbor a higher pCNV than *Copia*

(RLC) elements (Wilcoxon test, W = 6517781, p-value < 2.2e-16; **Fig 1B, C**). Furthermore, a PCA based

142 on RLG pCNVs did not allow us to discriminate accessions based on their phylogenetic relationship,

while a PCA performed with RLC elements separated the samples by genetic lineage (**Fig 1D**). The

144 strongest result was found with a PCA performed with the five youngest and putatively most recently

active LTR-RT families found in the pangenome of *B. distachyon* (the Angela families RLC_BdisC022,

146 RLC_BdisC024, RLC_BdisC030, RLC_BdisC209 and the Tekay family RLG_BdisC004), with the first two

axes together explaining more than 77.8% of the variance. Finally, with the exception of samples

148   from the most recently diverged clades A_East and A_Italia (13 kya), they further allowed us to

discriminate samples based on the genetic clade of origin (**Fig 1 B,D**).

Fig 1. Natural diversity of proximal copy number variation (pCNV) of LTR-RTs in B. distachyon. (**A**) Origin of the 320 natural accessions included in this study. Accessions that were used for the mobilome-seq are labelled in the map and a PCA with the bioclimatic variables of their place of origin is depicted. Colors of points correspond to the genetic clades whose estimated split is shown in the phylogenetic tree. Black points indicate that the accession cannot be clearly assigned to one genetic clade. (**B**) Heatmap with read counts of TE-derived sequences (proxy for the copy number variations, pCNVs) of all 40 annotated LTR-RTs in 320 natural accessions of B. distachyon. Accessions are sorted according to their phylogeny and names of recently active TE-families are highlighted in red. (**C**) Overall estimates of the copy numbers of Copia and Gypsy-type LTR-RTs in 320 natural accessions. (**D**) PCAs of pCNVs of all Gypsy, Copia and all recently active LTR-RT families, highlighted in (**B**) belonging to the Angela & Tekay families (RLG_Bdis004, RLC_BdisC030, RLC_BdisC209, RLC_BdisC024 and RLC_BdisC022). Colors of points indicate the genetic clade of accessions. (**E**) Output of the LMM analyses between pCNVs of LTR-RTs and bioclimatic variables at the accessions` origin. Bubbles indicate a significant association (P-value <0.05). Colors and sizes of bubbles show the part of the variance (marginal R^2) explained by the bioclimatic variables in %.

To test whether the accumulation of LTR-RT sequences correlated with environmental factors, we retrieved bioclimatic variables comprising precipitation, temperature, aridity levels, solar

radiation and atmospheric pressure at each locality. We then ran linear mixed models (LMM) where pCNV per LTR-RT family was entered as the response variable, the bioclimatic variables entered

separately as fixed factors and the clade of origin as random factors to account for population structure. Marginal R^2 extracted for each LMM did not exceed 10% even for the putatively most

recently active LTR-RT families (**Fig 1E**), indicating that albeit significant, the association between pCNV per family and the environment was mild in our study system. We observed similar

associations between pCNVs and bioclimatic variables when not accounting for population structure and running classical linear model analyses (**S1 Fig**). With the exception of RLC_BdisC010 and

RLG_BdisC265, for which more than 40% of the variance in pCNVs was explained by environmental factors, the LTR-RT families showed non-significant to mild associations with environmental variables

(**S1 Fig**).

Recently active LTR-RT families produce eccDNAs

The coverage-based approach for estimating LTR-RT CNVs considers all TE-derived sequences and

166	does not take into account that individual families differ in their age structures, turnover times and proportion of full-length, potentially autonomous mobile elements (Stritt *et al.*, 2020). We therefore

168	complemented our *in silico* analysis by experimentally testing whether LTR-RT families were indeed still active *in planta* and whether the mild but significant correlation we observed between global

170	pCNVs and bioclimatic variables may be due to a stress-specific activity.

To cover a wide range of the genetic, geographic and bioclimatic diversity of *B. distachyon*, we

172	selected nine natural accessions belonging to the five genetic clades and originating from contrasting

habitats (**Fig 1A**). Considering the role of Pol IV in the silencing of TEs in plants, we also included two

174   independent Pol IV mutant lines, a sodium azide mutagenized line (hereafter *Bd nrpd1-1*) and a T-

DNA line (hereafter *Bd nrpd1-2*), both carrying a homozygous mutation in the largest subunit of

176   Pol IV (NRPD1; Bradi2g34876) in the Bd21-3 background. We exposed plants to eight different

stresses, namely cold, drought, heat, salt, submergence, infection with *Magnaporthe oryzae*,

178   treatment with glyphosate and chemical de-methylation and performed mobilome-seq on all

resulting 105 samples (see Material and Methods).

180   Following the removal of organelle-derived reads, we first assembled mobilome reads and

aligned the resulting ten longest contigs of each sample to the reference assembly of *B. distachyon*.

182   We then screened genomic regions for which at least three assembled mobilome contigs of different

samples aligned and further assessed in which genotypes or stresses those contigs occurred. We

184   only retained circle-forming regions with a specificity above 50 % (i.e., regions for which more than

half of the contigs belonged to a certain stress or genotype). In addition, we also kept recurrently

186   active regions, present at a high frequency independently of the stress or the genotype in at least

ten samples. In total, we retained 15 circle-forming regions, all of which contained TE sequences

188   (**Fig 2**). Eight of these corresponded to the Angela family (RLC_BdisC024, RLC_BdisC022,

RLC_BdisC030, RLC_BdisC209), four to CRM elements (RLG_BdisC039, RLG_BdisC102), and one each

190   to the SIRE (RLC_BdisC026), the Alesia (RLC_BdisC010) and the non-autonomous and unclassified

RLG_BdisC152 family. We hereafter refer to RLC_BdisC024 as *HOPPLA* (German allusion for the the

192   surprising finding of a jumping element). We did not find stress specificity in the formation of

eccDNAs (**Fig 2A**). However, our results pointed to a genotype-dependent formation of eccDNAs for

194   the RLC_BdisC209, RLC_BdisC026 and *HOPPLA* families (**Fig 2B**). In particular, two contigs containing

HOPPLA elements were exclusively detected in the two *pol IV* mutants (**Fig 2B**).

Fig 2 Assessment of LTR-RT mobility in B. distachyon. Stress (**A**) and genotype (**B**) specificity of the formation of eccDNA as determined by the alignment of assembled mobilome-seq reads. The color represents the degree of specificity and numbers indicate the count of samples from which one of their ten longest contigs aligns to each of the circle-forming regions. Annotations of regions are indicated on the y-axis. Multiple annotations in the same circle- forming region were concatenated. The two pol IV mutants and the controls Bd21-3 and the outcrossed line BdNRPD1 (+/+) are summarized as RdDM and Bd21-3, respectively

HOPPLA activity is increased in the *pol IV* mutants regardless of the stress applied

Fragmented eccDNAs or circles containing only of the two LTRs (1-LTR circles) can be formed following reverse transcription by auto-integration, alternative end-joining in the virus-like particles (Garfinkel *et al.*, 2006; Yang *et al.*, 2023a) or by a recombination of the two LTRs of genomic copies (Smith & Vinograd, 1972; Gaubatz, 1990). Hence, 1-LTR circles do not necessarily imply LTR-RT mobility. In contrast, recent work indicates that 2-LTR circles are formed following the complete reverse transcription by non-homologous end-joining of an intact full-length linear RT copy that is capable of integrating into the genome (Yang *et al.*, 2023a). Indeed, the detection of full-length 2-LTR circles of well characterized autonomous LTR-RTs such as *EVD* (*ATCOPIA93*) has been directly linked to their actual transposition (Lanciano *et al.*, 2017). As a complement to the assembly-based

analysis, we thus followed a stringent approach to analyse our mobilome-seq data. We aligned reads to a library comprising artificial 3'LTR-5'LTR fusions of all full-length LTR-RTs annotated in the

B. distachyon reference assembly (Stritt *et al.*, 2020). Because such reads are not expected to be present in genomic DNA, this allowed us to speciflangically detect intact 2-LTR circles of

extrachromosomal LTR-RTs capable of integrating into the genome. To control for possible traces of undigested genomic DNA that may subsequently be amplified by the Phi29 enzyme during

mobilome-seq (Silander & Saarela, 2008), we also included publicly available genomic reads of all nine accessions in our analysis.

We found that several LTR-RTs formed eccDNA with 2-LTR junctions. Yet, most of them occurred sporadically and we did not observe a recurring stress-specific formation of 2-LTR circles

for any of the 37 LTR-RT families with annotated full-length copies (**Fig 3A**). For instance, we found a very strong signal for RLC_BdisC031 that was solely detected in glyphosate-treated Bd21-3 plants

and therefore not further considered in the analysis. In contrast, and in accordance with the assembly-based approach, the Angela element *HOPPLA* showed a recurrent formation of 2-LTR

circles. However, this formation was not triggered by a specific stress and only occurred in the two independent *pol IV* mutants (**Fig 3B**). Finally, our attempt to transiently inhibit LTR-RT silencing using

alpha-amanitin and Zebularine (a combination of inhibitors shown to increase the activity of LTR-RTs in *A. thaliana* and rice), did not result in the consistent activation of *HOPPLA* or other LTR-RTs in

multiple accessions (**Fig 3A**). The presence of 2-LTR circles of *HOPPLA* in *Bd nrpd1-2 (-/-)* was confirmed by an inverse PCR on total DNA that was not subjected to a rolling circle amplification,

with outward facing primers specific to the two LTRs (**Fig 3C**). Notably, we also detected a faint signal for the outcrossed line *Bd NRPD1 (+/+)* suggesting a weak activity of *HOPPLA* in wild-type plants.

Since the stress- or tissue-dependent activity of LTR-RTs is mediated by the specific binding

of transcription factor (TF), we screened the consensus sequence of *HOPPLA* for motifs of known TF

232   binding sites. First, we validated this approach by analyzing one of most active copies (AT1G11265)

of the heat-responsive *ONSEN* family of *A. thaliana*. As expected, a GO term analysis indicated a

Fig 3 *HOPPLA* forms 2-LTR circles in the *pol IV* mutants. Relative abundance of 2-LTR-junction spanning reads depending on the stress (**A**) and the genotype (**B**) of individual mobilome-seq samples. (**C**) Inverse PCR using total DNA not subjected to a rolling circle amplification for the confirmation of an increased amount of extrachromosomal 2-LTR circles of *HOPPLA* in the *Bd nrpd1-2* (-/-) mutant compared to the *Bd NRPD1* (+/+) outcrossed line. Three biological replicates are shown. (**D**) GO enrichment analysis of transcription factors for which binding sites have been detected in the consensus sequence of *HOPPLA*. Colors indicate number of TF-binding sites found. GO-terms that occur at least six times are highlighted in the plot. All GO-terms and their number of occurrences are listed in S1 Table.

strong enrichment of heat-responsive TFs for this element (**S2 Fig**). In contrast to the well-known,

stress-responsive *ONSEN* LTR-RT, the GO terms of TFs that could bind to *HOPPLA* indicated that

developmental processes and auxin-activated signaling pathways played a role in its activity, rather

than specific stresses (**Fig 3D**).

Members of the *HOPPLA* family differ in activity

Because individual copies of the same LTR-RT family can differ in their activity (Cavrak *et al.*, 2014),

we al*so ana*d the relative abundance of 2-LTR-spanning reads for each annotated full-length copy of

HOPPLA. This analysis revealed a great diversity of eccDNA formation among individual copies of the

HOPPLA family and confirmed the strongest activity of *HOPPLA* in the two *pol IV* mutants (**Fig 4A**).

We also obtained a few reads spanning the 2-LTR-junction of the two most active *HOPPLA* copies

(Bd3_22992889 and Bd4_25471847) in the *Bd NRPD1 (+/+)* control line (**Fig 4A**), which confirmed

the weak but detectable band for the inverse 2-LTR PCR for *Bd NRPD1 (+/+)* (**Fig 3C**).

Using the meta information of individual *HOPPLA* copies described previously (Stritt *et al.*,

2020), we further assessed which genomic factors (DNA methylation, CG content, distance to the

closest gene or age of the copy) were linked to the 2-LTR circle formation for individual *HOPPLA*

copies. While no clear pattern emerged from this analysis, the most active copies of *HOPPLA* tend

to be rather young (**Fig 4B**).

Fig 4 Members of the *HOPPLA* family differ in activity. (**A**) Relative abundance of 2-LTR-junction spanning reads of individual full-length copies of the *HOPPLA* family. (**B**) Age (age_a), closest distance to gene (d_closest), GC content (GC), and methylation levels in CG, CHG and CHH contexts of all individual genomic full-length copies of *HOPPLA* in percent. The color indicates relative abundance of 2-LTR-junction spanning reads from the mobilome-seq in (**A**).

 ***HOPPLA* is targeted by Pol IV-dependent 24 nt siRNAs in the wild type and transposes in *pol IV* mutant plants**

 The pivotal role of Pol IV in producing TE-specific 24-nt siRNAs for RNA-directed DNA methylation

has been demonstrated in many plant species including *A. thaliana* (Ito *et al.*, 2011), rice (Xu *et al.*,

 2020) and for the Alesia family (RLC_BdisC010) in *B. distachyon* (Böhrer *et al.*, 2020). To confirm that

the increased production of 2-LTR eccDNA circles of *HOPPLA* in mutants deficient in *B. distachyon*

 NRPD1 is correlated with a depletion of 24 nt siRNAs, we performed a small RNA blot including

samples from the two *pol IV* mutants and their respective wild-type controls. Using a hybridization

262   probe specific to the *HOPPLA* LTRs, 24 nt siRNAs were detected in the control lines Bd21-3 and

Bd NRPD1 (+/+), but not in either of the *pol IV* mutant lines (**Fig 5A**). This finding strongly suggests

264   that *HOPPLA* is under control of the Pol IV-RdDM pathway, and that the absence of 24 nt siRNAs

results in the upregulation and increased production of 2-LTR eccDNAs from *HOPPLA*. Furthermore,

266   RNA-seq data from part of the same mutant panel shows that *HOPPLA* is the most upregulated LTR-

RT family in the *Bd nrpd1-2 (-/-)* background compared to the *Bd NRPD1 (+/+)* control line, indicating

268   that the reduction of 24 nt siRNAs is likely associated with an increased expression and subsequent

formation of *HOPPLA* eccDNAs in both *pol IV* mutants (**Fig 5B**).

270   To complete their life cycle, reverse transcribed extrachromosomal copies of LTR-RTs have to

integrate into the host genome (Schulman, 2013). Because all our analyses congruently pointed to

272   the activity of *HOPPLA* in the *Bd nrpd1-2 (-/-)* mutant, we sequenced the genome of seven

Fig 5 Loss of 24-nt siRNAs leads to an increased activity of *HOPPLA* in the Pol IV mutants (**A**) Northern plot for the detection of 24-nt siRNAs specific to the 3` LTR of *HOPPLA* in the *pol IV* mutants *Bd nrpd1-1* and *Bd nrpd1-2* (-/-) and their control lines Bd21-3 and the outcrossed line *Bd NRPD1* (+/+). (**B**) SalmonTE analysis of the expression of LTR-RTs in *Bd nrpd1-2* (-/-) relative to the outcrossed control line *Bd NRPD1* (+/+). LTR-RTs with a log2 fold change of at least two are labeled, three biological replicates were analysed.

individuals of *Bd nrpd1-2 (-/-)*, six *Bd NRPD1 (+/+)* plants and one wild-type Bd21-3 plant to detect

new *HOPPLA* insertions. As TIPs were detected relative to the reference genome Bd21 (an accession

closely related to Bd21-3 but not genetically identical), we first removed all conserved Bd21-3-

specific TIPs detected in multiple lines. We manually curated all filtered candidate TIPs and showed

that *HOPPLA* was the only family for which validated TIPs were identified in one of the re-sequenced

Bd nrpd1-2 (-/-) plants (Bd1 38798495, Bd1 42205987, Bd4 28119639) (**S3-5 Figs**). This confirmed

that the loss of Pol IV function led to an increased production of eccDNA, as well as actual

transposition and accumulation of novel *HOPPLA* copies in the tested *Bd nrpd1-2 (-/-)* mutant. The

presence of reads spanning the insertion site indicated that the detected *HOPPLA* insertions were

heterozygous or probably somatic for the insertion Bd4 28119639, which exhibited a specifically low

proportion of clipped reads. No TIPs were detected for any other LTR-RT family.

Genome-wide association studies for pCNVs do not recover known components of RdDM

To decipher the genetic basis of *HOPPLA* accumulations in natural populations, we first performed a

genome-wide association study (GWAS) using *HOPPLA* pCNVs in the diversity panel of 320 natural

accessions (**Fig 1B**) as a phenotype*.* We identified only one region with two significant peaks (FDR-

adjusted p-value < 0.05, Bd5 6920000-6960000 and 7210000-7240000) obtained for the mobile

HOPPLA family (**Fig 6**). Because inserted copies of *HOPPLA* may themselves lead to significantly

associated regions in the GWAS as shown in *A. thaliana* (Quadrana *et al.*, 2016), we first verified that

there were neither TIPs (Horvath *et al.*, 2023) nor annotated reference insertions of *HOPPLA* in that

region (**Fig 6**). As described above, our data suggested that the loss of 24-nt siRNAs in the Pol IV

mutants was sufficient to mobilize *HOPPLA*. We therefore further tested whether any of the genes

encoding subunits of Pol IV or Pol V would be localized in or near this region (window size 50 kb)

296 (**S3 Table**). We did not detect any known Pol IV or Pol V- related genes, but instead found

Bradi5g05225, an ortholog of the *A. thaliana* ROS1-associated methyl-DNA binding protein 1 (RMB1,

298 AT1g63240) (Liu *et al.*, 2021a), to be co-localized with the peak.

To test whether genomic regions might be recurrently associated with their pCNVs, we finally

300 extended the GWAS analyses to the four other most recently active families (RLC_BdisC022,

RLC_BdisC030, RLC _BdisC209 and RLG_BdisC004) (**S6 Fig**). We extracted the candidate genes for

Fig 6 Genetic regions associated with pCNVs of recently active LTR-RTs (**A**) Manhattan plot depicting the GWAS results of pCNV of *HOPPLA* in 320 accessions of *B. distachyon.* Colored points indicate SNPs linked to TIPs (blue), TAPs (red) or known components of the Pol IV and V holoenzymes (asterisks). Threshold of significance (false discovery rate adjusted p-value <0.05) is marked with dashed lines. A significant region containing the candidate gene Bradi5g05225 (window size 50 kb) is highlighted. (**B**) UpSet plot of genes in 20 kb windows surrounding significant regions with at least two SNPs above the threshold of significance (FDR-adjusted p-value <0.05 for *HOPPLA*, RLC_BdisC209 and RLC_BdisC022 and Bonferroni correction for RLC_BdisC030 and RLG_BdisC004) of the five most recently active LTR-RT families in *B. distachyon.* To visualize potential overlaps, a list of the componenlangts of the Pol IV and Pol V holoenzymes is included in the UpSet plot.

302   each of the five families (see **S3 Table** and Material and Methods). Apart from the two closely related

families RLC_BdisC030 and RLC_BdisC209 that shared the majority of their GWAS candidates, and

304   RLC_BdisC022 and RLC_BdisC209 that shared three genes, we found no overlap of annotated loci

potentially contributing to the pCNVs of recently active families (**Fig 6**).

Discussion

308   Understanding the dynamics of TEs and their role in adaptation is currently one of the major

challenges in the field of evolutionary genomics. The fact that mobile TEs are a source of epi/genetic

310   diversity and potential drivers of evolution has been demonstrated in many organisms including

fungi (Muszewska *et al.*, 2019), insects (Gilbert *et al.*, 2021), mammals (Senft & Macfarlan, 2021)

312   and plants (Lisch, 2013). However, while there are a number of examples showing that certain TE

insertions facilitated the adaptation to changing environments (for review Bourgeois & Boissinot,

314   2019), TEs are generally harmful (Bourgeois *et al.*, 2020; Horvath *et al.*, 2023; Langmüller *et al.*, 2023)

and therefore controlled by complex silencing mechanisms. To foster our understanding of TE

316   activity, we investigated the environmental conditions and genetic factors associated with the

accumulation and mobility of LTR-RTs in plant genomes. By measuring LTR-RT pCNVs in a panel of

318   320 *B. distachyon* natural accessions, we show that the intra-specific variations of pCNVs of RLC

elements, but not the pCNVs of the generally older and more abundant RLG elements (Stritt *et al.*,

320   2020), separate accessions according to their genetic cluster of origin. This is even more striking for

members of the Angela (RLC_BdisC022, RLC_BdisC024, RLC_BdisC030, RLC_BdisC209) and the Tekay

322   (RLG_BdisC004) family, which are the youngest families in *B. distachyon* (Stritt *et al.*, 2020). Highly

polymorphic among natural accessions of *B. distachyon* (Stritt *et al.*, 2018, 2020), they are

324   expectedly the main drivers of lineage-specific expansions of pCNVs in our study system. Hence, we

        do not only confirm that LTR-RT families in *B. distachyon* globally differ in size (Stritt *et al.*, 2020) but

326   also demonstrate that the accumulation of genomic sequences derived from specific families varies

        significantly among natural accessions.

328   The transcriptional activity of LTR-RTs can be triggered by specific environmental stresses

        (Grandbastien, 2015; Negi *et al.*, 2016). Given that *B. distachyon* occurs in a wide range of different

330   habitats in the Mediterranean area (Minadakis *et al.*, 2023b), this characteristic feature of LTR-RTs

        provides a potential explanation for the pCNVs we observed across natural accessions (Quadrana *et*

332   *al.*, 2016; Baduel *et al.*, 2021). Yet, for all LTR-RT families, pCNVs correlate only moderately with

        environmental factors. Consequently, our genomic data do not support a large effect of the

334   environment on LTR-RT activity in *B. distachyon*. While this result could seem startling, it is not

        completely surprising. Indeed, many LTR-RT families, and especially the old RLG elements, do not

336   show signs of increased activity in the recent past in *B. distachyon (Stritt et al., 2020)*. Considering

        that their copy number expansions took place in a climate that has drastically changed following the

338   last glacial maximum (Minadakis *et al.*, 2023b), a limited link between their activity and the current

        environmental conditions is actually expected for most families. In contrast, the lack of correlation

340   between current bioclimatic variables and copy number variation for families with ongoing activity

        (RLC_BdisC022, RLC_BdisC024, RLC_BdisC030, RLC_BdisC209 and RLG_BdisC004; Stritt et al., 2020),

342   suggests a more complex mechanism than their pure dependence on specific stresses in certain

        environments. This hypothesis is supported by previous findings in *A. thaliana*. Indeed, a minor

344   impact of the environment on transpositional activity was also found in this species, where the two

        most associated environmental variables ('seasonality of precipitation' and 'diurnal temperature

346   range') only explained about 9 % of the observed variation (Baduel *et al.*, 2021).

Genetic factors are well-known to be essential in regulating LTR-RT activity (Miura *et al.*, 2001;

348   Tsukahara *et al.*, 2009; Mirouze *et al.*, 2009; Bourque *et al.*, 2018). As the loss of main players of the

RdDM silencing pathway leads to increased TE activity (Ito *et al.*, 2011; Benoit *et al.*, 2019; Xu *et al.*,

350   2020; Baduel *et al.*, 2021), the two *B. distachyon* Pol IV (NRPD1) mutants provided an ideal functional

tool to experimentally validate our *in silico* analysis. Since transcriptionally active LTR-RTs are not

352   necessarily able to transpose (Bajus *et al.*, 2022), we used a mobilome-seq approach to detect TE-

derived eccDNAs and transpositionally active LTR-RT families. We deliberately followed a very

354   stringent approach for analysing the data and by doing so, identified *HOPPLA* as the only highly

active LTR-RT family in *B. distachyon.* Indeed, *HOPPLA* is the only family for which we further detect

356   newly inserted copies in the Pol IV mutant.

The non-stress-specific activity of *HOPPLA* in the two independent Pol IV mutants supported

358   our *in silico* approach and strengthened the idea that genetic, rather than environmental stresses,

are major drivers of LTR-RT activity in *B. distachyon*. These results are also in line with our TF binding

360   sites analysis. In contrast to the heat-responsive *A. thaliana* element *ONSEN*, for which we

predominantly recovered TF-binding sites associated with heat response, *HOPPLA* seemed to be

362   targeted by TFs involved in developmental processes and auxin signaling (Leyser, 2018). Follow-up

studies should therefore address the question of whether the activity of *HOPPLA* or other families

364   differs between tissues or developmental stages, as observed for the endosperm-specific mobility

of *PopRice* in rice, for example (Lanciano *et al.*, 2017). Strikingly, despite the central role of Pol IV in

366   the RdDM pathway, we did not observe bursts of multiple LTR-RT families but instead found that the

loss of 24 nt siRNAs specifically activated individual copies of the *HOPPLA* family. Interestingly, we

368   also detected a weak signal for 2-LTR eccDNAs in the Bd21-3 wt and the outcrossed line *Bd NRPD1*

(+/+) but not in other natural accessions. This suggests that the accession-specific composition of

370　　the mobilome, and hence the genetic background of the *pol IV* mutant line, plays an important role

　　　　in LTR-RT activity. Related to this, we sporadically observed very strong signals for individual samples,

372　　which could indicate an accession-specific response of the mobilome to certain triggers.

　　　　　　Given that pCNVs vary greatly among genetic clades, assessing the effect of a genetic

374　　mutation of major components of the RdDM pathway in a set of genetically diverse natural

　　　　accessions would be timely, yet labor-intensive as transformation works more efficiently in the Bd21-

376　　3 background than in the other accession tested. Our attempt to transiently reduce LTR-RT silencing

　　　　in multiple accessions from different genetic clades using the chemical inhibition of Pol II and DNA

378　　methyltransferases (Thieme *et al.*, 2017) did not result in an increased activity of *HOPPLA* or

　　　　members of any other LTR-RT family. In addition, and despite the differences of activities observed

380　　among individual *HOPPLA* copies, we could not detect, in the present study, a clear link between

　　　　their activity and GC contents or methylation states. Taken together, these findings suggest that the

382　　specific function of the canonical RdDM with Pol IV, rather than generic DNA methylation states are

　　　　regulating *HOPPLA* activity. Yet, our GWAS failed to recover major components of the RdDM

384　　pathway. Instead, the diversity of activity within the *HOPPLA* family may suggests that the presence

　　　　of single active copies could determine the fate of an entire family. In addition, pCNV is also

386　　dependent on the removal rate of LTR-RT families, which varies greatly in *B. distachyon* (Stritt *et al.*,

　　　　2020). This complexity of parameters affecting the dynamics of LTR-RT might explain why none of

388　　the genes known to be involved in silencing LTR-RTs are associated with pCNV for *HOPPLA* or any

　　　　recently active family. Our candidate locus containing Bradi5g05225, a gene related to RMB1 whose

390　　loss of function has been shown to result in DNA hypermethylation　(Liu *et al.*, 2021b) , remains

　　　　nonetheless a great candidate for functional validation.

392   Altogether, our work confirms that LTR-RTs in *B. distachyon* are 'well-behaved' (Stritt *et al.*,

2018) and that the evolutionary consequences of their mobility are hard to study in real-time.

394 Indeed, while mobilome-seq revealed a sporadic activity for other families, we only found recurring

activity and new insertions of *HOPPLA* in the *pol IV* mutant. These results somewhat contrast with

396 our population genomics analyses which clearly indicate an ongoing activity of several LTR-RT

families in natural accessions. We propose that the activity of LTR-RTs is relatively low and might

398 depend on a complex interaction between genetic factors, developmental stages and, more

marginally, the punctual occurrence of stresses.

Material and methods

Estimation of LTR-RT pCNVs

404 We used publicly available genomic reads of 320 sequenced natural accessions of *B. distachyon*

(Gordon *et al.*, 2017, 2020; Skalska *et al.*, 2020; Stritt *et al.*, 2022; Minadakis *et al.*, 2023a) to assess

406 the natural variation of copy numbers of LTR-RTs. Because sequencing depth differed substantially

between accessions (Minadakis *et al.*, 2023a), we first downsampled all fastq files to the read

408 number of the sample with the lowest number of reads (4.230.721 reads) using the reformat.sh

function of BBtools (v 38.75, BBMap Bushnell B., sourceforge.net/projects/bbmap/). Downsampled

410 reads were aligned to the TREP consensus sequences of LTR-RTs and the reference assembly of Bd21

(v 3.0) using BWA-MEM (v 0.7.17-r1188) (Li & Durbin, 2009) with the -M and the -a options set,

412 hence outputting all alignments found. Coverages of LTR-RTs and the reference assembly were

assessed using bedtools (v 2.30.0) (Quinlan & Hall, 2010) genomecov with the -d and -split

parameters set. For each of the LTR-RT families, a proxy for copy number was obtained by normalizing the coverage signals by the coverage of the entire reference assembly and by correcting

for the length of the consensus sequences. We favoured this coverage-based approach over an analysis based on transposon insertion polymorphisms (TIPs) as estimates based on TIPs are

reference-dependent and biased by the phylogeny in our study system. We have for instance previously shown that accessions from the B_East clade harbor significantly less TIPs than accessions

from the A_East clade due to the fact that the reference genome Bd21 belongs the B_East clade (Stritt *et al.*, 2018). In addition, whole-genome *de novo* assembly of 54 *B. distachyon* natural

accessions and the subsequent pangenome analysis revealed that non-reference accessions display large genomic variations (Gordon *et al.*, 2017) which may further bias the estimates of TIP

abundances. pCNV raw data were processed using R (v 3.6.3 and 4.0.2) (R Core Team, 2020) in Rstudio(RStudio Team, 2016).

Variation in pCNVs across the 320 natural accessions was visualized with a heatmap drawn with the heatmap() function natively provided in R version 4.0.2. We computed pairwise genetic

distances between accessions with the R package pvclust v 2.2.0 (Suzuki & Shimodaira, 2006). The resulting tree was used to order accessions phylogenetically on the heatmap. PCAs based on pCNVs

were obtained with the R package ggbiplot v 0.55 (Vu, 2011).

      To test for an association between pCNVs and environmental variables, we retrieved information

432   about climatic variables at each local site from (Minadakis *et al.*, 2023b). Linear mixed model analyses where pCNV per LTR-RT family was entered as the response variable, the bioclimatic

434   variables entered separately as fixed factors and the clade of origin as random factors to account for population structure were ran with the R package lme4 (Bates *et al.*, 2015). The part of the variance

436   explained by the fixed- (marginal R^2) were computed following (Nakagawa & Schielzeth, 2013) and

visualized as bubble plot with the R package ggplot2 (Wickham, 2016). Classical linear models were

438   run in base R.

440   **Plant material, growth conditions and stresses for mobilome-seq**

Brachypodium distachyon natural accessions used in this study comprised Bd21, Bd21-3, Cm18,

442   Cb23, ABR2, Bd29-1 BdTR13c, RON2 and Arn1. Because Pol IV is known to play an important role in

LTR-RT silencing in plants (Stonaker *et al.*, 2009; Ito *et al.*, 2011; Ferrafiat *et al.*, 2019), we also

444   included the sodium azide mutagenized *pol IV* mutant line NaN74 (*Bd nrpd1-1*) (Dalmais *et al.*, 2013;

Böhrer *et al.*, 2020), the T-DNA insertion *pol IV* mutant line JJJ18557 Nr31 (Bragg *et al.*, 2012)

446   *Bd nrpd1-2 (-/-)* and a corresponding sibling, outcrossed control line *Bd NRPD1* (+/+) in the

background of the natural accession Bd21-3. For *in vitro* experiments, seeds were soaked for 4 h in

448   tap water and, without damaging the embryo, the lemma was carefully peeled off. Seeds were then

surface-sterilized for 30 seconds in 100% ethanol and immediately rinsed three times with sterile

450   tap water. Surface-sterilized seeds were placed with the embryo facing down and at an angle of

about 30° towards the side, onto solid ½ MS-medium (2.15 g/L MS basal salt without vitamins

452   (Duchefa Biochemie, Haarlem, NL), 0.5 g/L MES-Monohydrate, 10 g/L sucrose , pH 5.8 (KOH), 0.25 %

Phytagel (Sigma-Aldrich, St. Louis, USA) in 'De Wit' culture tubes (Duchefa Biochemie, Haarlem, NL).

454   Plants were grown at 24 °C (day) / 22 °C (night), 16 h light under controlled conditions in an Aralab

600 growth chamber (Rio de Mouro, PT) for 25 to 29 days until the onset of stresses. For salt stress,

456   seedlings were transplanted to solid ½ MS-medium supplied with 300 mM NaCl and grown for five

days at 24/22 °C, 16h light. A solution of sterile-filtrated Glyphosate (Sintagro AG, Härkingen,CH) (20

458   mM, diluted in water) was applied to leaves using a piece of soaked sterile filter paper and plants

were incubated for four days at 24/22 °C, 16h light. Drought stress was induced by uprooting plants

460   from the medium and incubating them for 2:15 h at 24 °C in the light. Before sampling, plants were

allowed to recover for two hours on fresh ½ MS-medium. For the infection with *Magnaporthe oryzae*

462   (rice blast) six isolates (FR13, Mo15-27, 9475-1-3, IK81, M64 and Mo15-19) with spore

concentrations between 130'000- 200'000 K spores per isolate per mL sterile water, supplied with

464   0.2 % Tween 20 were mixed and applied with a cotton swab to plant leaves. Plants were incubated

for 24 h in the dark (24/22 °C) and then grown for another three days at 24/22 °C, 16h light. For heat

466   stress, plants were incubated for 8 h at 42 °C. Before sampling, heat-stressed plants were allowed to

recover for 16 h at 24/22 °C. Cold stress was induced by incubating plants for 24 h at 2 °C on ice at

468   16 h light. Prior to sampling, plants were allowed to recover for two hours at 24 °C in the light. For

submergence stress, two small holes were drilled just above the growth medium and at the top

470   through the wall of the culture tubes. Tubes were then inverted and submerged upside down for 48

h at 24/22 °C, 16h light using a custom rack in a plastic beaker filled with 2.5 liters of 24 °C tap water.

472   In this way, it is possible to submerge plant leaves without the medium coming into contact with the

water. Chemical de-methylation of DNA was conducted according to (Thieme *et al.*, 2017) by

474   germinating and growing plants for 28 days on ½ MS-medium supplied with a mixture of Zebularine

(Sigma-Aldrich, St. Louis, USA) and alpha-amanitin (Sigma-Aldrich, St. Louis, USA). Because the drug

476   treatment severely affected the growth of seedlings, we omitted a treatment of mutant plants and

used reduced concentrations of 20 uM (Zebularine) and 2.5 mg/ml (alpha-amanitin), respectively

478   for all natural accessions.

480   **Mobilome sequencing and validation of eccDNAs**

DNA was extracted using the DNeasy plant kit (Qiagen, Venlo, Netherlands) according to the protocol

482   of the manufacturer. DNA concentration was measured using the Qubit high sensitivity kit

(Invitrogen, Waltham, USA). Mobilome sequencing was performed according to (Lanciano *et al.*,

484   2017) using pooled DNA of two biological replicates per sample. For this, 50 ng of DNA from both

biological replicates were pooled and diluted to a volume of 58 µL. To enrich eccDNA, DNA was first

486   purified using the GENECLEAN kit (MP Biomedicals, Santa Ana, USA) according to manufactures

recommendations using 5 µL glass milk with an elution volume of 35 µL. Thirty µL of the eluate were

488   digested using the Plasmid-Safe ATP-dependent DNase (Biosearch Technologies, Hoddesdon, UK) for

17 h at 37 °C. The digestion product was then subjected to an ethanolic precipitation and the

490   precipitated eccDNA amplified using the illustraTempliPhi Amplification Kit (Cytiva, Marlborough,

USA) according to (Lanciano *et al.*, 2017) with an extended incubation time of 65 h at 28 °C. The

492   templiphi product was diluted 1:10, quantified using the Qubit high sensitivity kit and 120 ng per

sample were used for library preparation. Sequencing libraries were prepared using the Nextera DNA

494   Flex Library Prep and the Nextera DNA CD Indexes (Illumina, San Diego, USA). Quality of libraries

were assessed using the Tape Station (Agilent Technologies, Santa Clara, USA) with High Sensitivity

496   D1000 screen tapes and concentrations were measured using the Qubit high sensitivity kit. Up to 12

indexed libraries were pooled and sequenced with an Illumina MiSeq sequencer using the MiSeq

498   reagent kit v3 (600 cycles). Raw reads have been uploaded to ENA (accession number PRJEB58186).

The presence of extrachromosomal circular copies of *HOPPLA* (RLC_BdisC024) was validated

500   by an inverse PCR using 7 ng/µL total DNA. Input quantities of DNA were controlled using primers

specific to the S-adenosylmethionine decarboxylase (SamDC) gene (Hong *et al.*, 2008). Sequences of

502   primers are listed in **S5 Table**.

Analysis of Mobilome-seq

Reads were trimmed using the BBDuk tool of BBtools (BBMap (v 38.75, Bushnell B., sourceforge.net/projects/bbmap/) with the parameters qtrim = rl and trimq = 20. Reads originating from organelles were removed by aligning reads to the chloroplast genome (NC_011032.1) (Bortiri *et al.*, 2008) and the mitochondrion genome (v 1.0.0) of *B. distachyon* using BWA-MEM (v 0.7.17-r1188) (Li & Durbin, 2009) with the -M parameter set. Unmapped reads were isolated using samtools (v 1.13) (Danecek *et al.*, 2021) view -b -f 4 and bedtools (v 2.30.0) (Quinlan & Hall, 2010) bamtofastq.

Organelle-filtered mobilome reads were assembled using the SPAdes genome assembler (v 3.13.0) (Prjibelski *et al.*, 2020). From each assembly, the top ten contigs were extracted and jointly aligned to the reference assembly of Bd21 (v 3.0) using BWA-MEM (v 0.7.17-r1188) with the -M parameter set. Bam files were converted into bed files using bedtools (v 2.30.0) bamtobed with the -split option set and overlapping contigs were merged using bedtools merge with the -o distinct, count, count_distinct and -c 4 parameters set. Assembled, circle-forming regions were annotated with bedtools intersect using the version 3.1 annotation of the reference assembly and the annotation of all full-length LTR-RTs (Stritt *et al.*, 2020) of the reference assembly. Annotated regions were extracted with bedtools getfasta and all sequences longer than 2 kb were isolated using SeqKit seq (v 0.11.0) (Shen *et al.*, 2016). Circle-forming regions that occurred in less than three samples were not included in the analysis.

To specifically detect mobilized LTR-RTs, we first extracted all annotated full-length LTR-RTs of the Bd21 reference assembly (Stritt *et al.*, 2020). Using a custom python script, we then merged the last 300 bp of the 3' to the first 300 bp of the 5' LTR to obtain a 'tail-to-head' library containing all annotated full-length LTR-RT copies annotated in Bd21. We then aligned organelle-filtered mobilome reads to the tail-to-head library of LTR-RTs and used bedtools (v 2.30.0) intersect to extract

aligned reads that were spanning the 2-LTR junction and that aligned to at least 5 bp of both LTRs.

530   The coverage of the junction-spanning reads was calculated using deeptools (v 3.5.1) (Ramírez *et al.*,

2016) with the parameters -bs 1, --ignoreDuplicates --outRawCounts set. To account for differences

532   in sequencing depth, the obtained coverage for 2-LTR-junction spanning reads was normalized with

the total coverage obtained with bedtools (v 2.30.0) genomecov with the -d and -split parameters

534   set, from the alignments of filtered reads to the reference assembly of Bd21 (v 3.0) obtained from

Phytozome 12 (International Brachypodium Initiative, 2010) generated by BWA-MEM (v 0.7.17-

536   r1188) (Li & Durbin, 2009). To plot the overall activity per family, normalized signals were summed

up for every individual TE family.

mRNA-sequencing and small RNA northern blotting

540   Leaves of 4-week-old *B. distachyon* plants were ground in liquid nitrogen and 500 µL of this powder

was subjected to TRIzol extraction following the supplier instructions (Invitrogen, CA, USA). 20 µg of

542   total RNA was treated with DNase I for 30 min., then repurified via phenol-chloroform extraction

and ethanol precipitation. DNase-treated total RNA samples were sent to Fasteris/Genesupport

544   (Plan-les-Ouates, Switzerland), subjected to poly(A)-tail selection, and then aliquoted for library

construction via the Illumina TruSeq Stranded mRNA Library Prep kit. Resulting stranded polyA+

546   RNA-seq (mRNA-seq) libraries were sequenced on an Illumina NovaSeq 6000. The raw paired-end

read data were deposited at the NCBI Gene Expression Omnibus (GEO accession: GSE243693).

548   For the small RNA blot analysis, 200 µg of each total RNA were size-fractionated using the RNeasy

Midi Kit (QIAGEN), as described previously (Böhrer *et al.*, 2020). Low molecular weight (LMW, <200

550   nt) RNAs are not bound by the silica membrane of the columns and were isolated from the collected

flow-through and wash aliquots. LMW RNAs were precipitated overnight using isopropanol.

552 Following a centrifugation step (45 min. at 24000 x g, 4°C) and the removal of the supernatant, the

pellet was washed with 75% ethanol, centrifuged (15 min. at 24000 x g, 4°C), dried at RT for 20 min.

554 then at 65°C for 5 min., and resuspended in 41 µL of DEPC-treated MilliQ water. LMW RNAs were

quantified using a Nanodrop device and 12.3 µg of LMW RNAs from each sample were loaded into

556 the 16% polyacrylamide gel (Böhrer *et al.*, 2020). After running, transfer and UV crosslinking,

membrane was prehybridized in PerfectHyb Plus buffer (Merck, Darmstadt, Germany) at 35°C and

558 then and hybridized at 35°C with the Klenow internally-labeled probe (*HOPPLA*), or with the 5'-end

labeled probe (miR160) (Böhrer *et al.*, 2020). After overnight hybridization, washing was performed

560 at 37°C. Signal detection requires 5-7 days exposure for *HOPPLA* and 1-2 days for miR160.

Oligonucleotide sequences for the probes are listed in **S5 Table.**

LTR-RT expression analysis

564 RNA-seq raw reads of *Bd nrpd1-2 (-/-)* and *Bd nrpd1-2 (+/+)* were trimmed for adapters using fastp

(v 0.23.2) (Chen, 2023) with the following options: --qualified_quality_phred 15 --

566 unqualified_percent_limit 4 --n_base_limit 20 --low_complexity_filter --

overrepresentation_analysis --correction --detect_adapter_for_pe. Cleaned reads were then

568 analysed using SalmonTE (v 0.4) (Jeong *et al.*, 2018) to measure global expression of LTR-RTs. LTR-RT

consensus sequences of *B. distachyon* obtained from the TRansposable Elements Platform (TREP,

570 https://trep-db.uzh.ch/) were used to generate the custom library for SalmonTE. Default options of

SalmonTE quant and test function were used to quantify expression and to perform statistical

572 analysis. Expression data were plotted using R (v 3.6.3) in RStudio (v 7d165dcf).

Motif analysis

576 The consensus sequence of *HOPPLA* was screened for known transcription factor binding sites

obtained from the PlantTFDB (Jin *et al.*, 2017) using FIMO (v 5.1.1) (Grant *et al.*, 2011). To

578 functionally annotate transcription factors that could bind to *HOPPLA,* we used GO-terms of the

Gramene (release 50) database (Tello-Ruiz *et al.*, 2021) downloaded from the platform agriGO (v 2.0)

580 (Tian *et al.*, 2017). Generic, TF specific GO terms (GO:0003700, GO:0006355, GO:0005634,

GO:0003677, GO:0043565, GO:0046983, GO:0003682 GO:0045893) such as 'positive regulation of

582 DNA-templated transcription' were removed from the list of GO terms as they would interfere with

the downstream analysis. The remaining GO terms of transcription factors potentially binding to

584 *HOPPLA* were visualized with REVIGO (Supek *et al.*, 2011) using the 'SimRel' semantic similarity

measure, the option 'small' and the GO terms of the *Oryza sativa* Japonica Group. The total number

586 of occurrences of individual GO terms was taken into account with the option 'higher value is better'.

GO terms occurring more than five times were labelled in the plots. As a proof of concept, we

588 followed the exact same approach using the sequence of one of the most active *ONSEN* copies

(AT1G11265) and the GO terms of *Arabidopsis thaliana*.

Detection of novel *HOPPLA* insertions in *Bd nrpd1-2 (-/-)*

592 DNA of adult plants was extracted using the DNeasy plant kit (Qiagen, Venlo, Netherlands) according

to the protocol of the manufacturer subjected to whole genome sequencing. Reads were trimmed

594 using the BBDuk tool of BBtools (BBMap (v 38.75, Bushnell B., sourceforge.net/projects/bbmap/)

with the parameters qtrim = rl and trimq = 20. Trimmed reads were aligned to the reference

596 assembly of Bd21 (v 3.0) using BWA-MEM (v 0.7.17-r1188) (Li & Durbin, 2009) with the -M

parameter set. Samtools (v 1.13) (Danecek *et al.*, 2021) was used to obtain sorted and indexed bam

598   files. TIPs were detected with detettore (v 2.0.3) (https://github.com/cstritt/detettore) with the

options –require_split, -q 30 and using the consensus sequences of LTR-RTs of *B. distachyon* (TREP-

600   database) and the annotation of all full-length LTR-RTs of Bd21 (Stritt *et al.*, 2020). Because both

Bd nrpd1-2 lines were in the Bd21-3 background we were able to exclude all Bd21-3 specific TIPs by

602   removing those insertions that were detected in multiple individuals with more than one genetic

background. Remaining TIPs were manually curated using the genome browser IGV (v 2.15.4.12)

604   (Robinson *et al.*, 2011). *HOPPLA* TIPs were visualized with JBrowse 2 (v 2.6.1) (Diesh *et al.*, 2023) .

Raw genomic reads of the re-sequencing of *Bd nrpd1-2 (-/-), Bd NRPD1 (+/+)* and Bd21-3 have been

606   uploaded to ENA (accession number PRJEB64053).

608   **GWAS for pCNV**

GEMMA 0.98.5 (Zhou & Stephens, 2012) was used to test for associations between SNPs

610   (Minadakis *et al.*, 2023a) and the LTR-RT families pCVNs, while correcting for population

structure (Stritt *et al.*, 2022; Minadakis *et al.*, 2023a). A centered relatedness matrix was first

612   created with the option -gk 1 and association tests were performed using the option -maf 0.05

to exclude rare alleles, and the default SNP missingness threshold applied by GEMMA that

614   excludes SNPs with missing data in more than 5% of the accessions. We selected 20 kb genomic

regions with a 10 kb overlap that contained at least two SNPs above the False Discovery Rate of

616   0.05 or Bonferroni correction threshold as candidate region using the R package rehh (v 3.2.2)

(Gautier & Vitalis, 2012). Genes overlapping with candidate regions were selected with the

618   BEDTOOLS (v 2.26.0) (Quinlan & Hall, 2010) intersect command using the version 3.1 of the *B.*

distachyon annotation file (https://phytozome-next.jgi.doe.gov) that are contained in the

620   significant regions. The UpSetR (Conway *et al.*, 2017) R package was used to visualize the

622 intersections of significant genes between the variables. Protein constituents of the Pol IV and Pol V enzymes (see **S4 Table**) were downloaded from the plant RNA polymerase database http://rna.polymerase.eu/.

Acknowledgements and author contributions

626 We thank the Genetic Diversity Center Zürich for providing the infrastructure for the mobilome-seq. M.T. and A.C.R. conceived the study and planned experiments. M.T., B.K., K.R., C.H., B.R. and M.B.

628 generated data. M.T., A.C.R., N.M. and W.X. analysed data. D.L., J.V. and R.S. provided mutagenized germplasm and online TILLING/T-DNA browsing tools. C.H., B.R. and T.B. identified and screened for

630 the *pol IV* null mutants. M.T and A.C.R wrote the paper with contributions from T.B., C.S. All authors read the manuscript. The authors declare that they have no conflict of interest. Professional editing

632 performed by Emanuelle Botté.

Funding

University of Zurich Research Priority Programs (URPP) Evolution in Action to M.T. and A.C.R.;

636 Schweizerischer Nationalfonds zur Förderung der Wissenschaftlichen Forschung. Grant Number: 31003A_182785 to A.C.R., W.X., N.M. and B.K.; Schweizerischer Nationalfonds zur Förderung der

638 Wissenschaftlichen Forschung. Grant Number: PZ00P3_154724 to C.S.; T.B. was supported by the Interdisciplinary Thematic Institute IMCBio (ITI 2021-2028 program), including funds from IdEx

640 Unistra (ANR-10-IDEX-0002), SFRI-STRAT'US (ANR 20-SFRI-0012) and EUR IMCBio (ANR-17-EURE-0023) in the framework of the French Investments for the Future Program.

References

Baduel P, Leduque B, Ignace A, Gy I, Gil J, Loudet O, Colot V, Quadrana L. **2021**. Genetic and environmental modulation of transposition shapes the evolutionary potential of Arabidopsis thaliana. *Genome Biology* **22**: 138.

Bajus M, Macko-Podgórni A, Grzebelus D, Baránek M. **2022**. A review of strategies used to identify transposition events in plant genomes. *Frontiers in Plant Science* **13**: 1080993.

Bates D, Mächler M, Bolker BM, Walker SC. **2015**. Fitting linear mixed-effects models using lme4. *Journal of Statistical Software* **67**: 1–48.

Benoit M, Drost HG, Catoni M, Gouil Q, Lopez-Gomollon S, Baulcombe D, Paszkowski J. **2019**. Environmental and epigenetic regulation of Rider retrotransposons in tomato. *PLoS Genetics* **15**: e1008370.

Böhrer M, Rymen B, Himber C, Gerbaud A, Pflieger D, Laudencia-Chingcuanco D, Cartwright A, Vogel J, Sibout R, Blevins T. **2020**. Integrated Genome-Scale Analysis and Northern Blot Detection of Retrotransposon siRNAs Across Plant Species. *Methods in molecular biology (Clifton, N.J.)* **2166**: 387–411.

Bortiri E, Coleman-Derr D, Lazo GR, Anderson OD, Gu YQ. **2008**. The complete chloroplast genome sequence of Brachypodium distachyon: Sequence comparison and phylogenetic analysis of eight grass plastomes. *BMC Research Notes* **1**: 61.

Bourgeois Y, Boissinot S. **2019**. On the Population Dynamics of Junk: A Review on the Population Genomics of Transposable Elements. *Genes* **10**.

Bourgeois Y, Ruggiero RP, Hariyani I, Boissinot S. **2020**. Disentangling the determinants of transposable elements dynamics in vertebrate genomes using empirical evidences and simulations. *PLoS genetics* **16**.

Bourque G, Burns KH, Gehring M, Gorbunova V, Seluanov A, Hammell M, Imbeault M, Izsvák Z, Levin HL, Macfarlan TS, *et al.* 2018. Ten things you should know about transposable elements. *Genome Biology 2018 19:1* **19**: 1–12.

Bragg JN, Wu J, Gordon SP, Guttman ME, Thilmony R, Lazo GR, Gu YQ, Vogel JP. **2012**. Generation and Characterization of the Western Regional Research Center Brachypodium T-DNA Insertional Mutant Collection. *PLOS ONE* **7**: e41916.

Butelli E, Licciardello C, Zhang Y, Liu J, Mackay S, Bailey P, Reforgiato-Recupero G, Martin C. **2012**. Retrotransposons control fruit-specific, cold-dependent accumulation of anthocyanins in blood oranges. *Plant Cell* **24**: 1242–1255.

Cavrak V V., Lettner N, Jamge S, Kosarewicz A, Bayer LM, Mittelsten Scheid O. **2014**. How a Retrotransposon Exploits the Plant's Heat Stress Response for Its Activation. *PLOS Genetics* **10**: e1004115.

Chen S. **2023**. Ultrafast one-pass FASTQ data preprocessing, quality control, and deduplication using fastp. *iMeta* **2**: e107.

Cho J, Benoit M, Catoni M, Drost H-G, Brestovitsky A, Oosterbeek M, Paszkowski J. **2019**. Sensitive detection of pre-integration intermediates of long terminal repeat retrotransposons in crop plants. *Nature Plants* **5**: 26–33.

682   **Conway JR, Lex A, Gehlenborg N**. **2017**. UpSetR: an R package for the visualization of intersecting sets and their properties. *Bioinformatics* **33**: 2938–2940.

684   **Dalmais M, Antelme S, Ho-Yue-Kuang S, Wang Y, Darracq O, d'Yvoire MB, Cézard L, Légée F, Blondet E, Oria N, *et al.* 2013**. A TILLING Platform for Functional Genomics in Brachypodium
686   distachyon. *PLoS ONE* **8**: 65503.

       Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A, Keane T,
688   **McCarthy SA, Davies RM**. **2021**. Twelve years of SAMtools and BCFtools. *GigaScience* **10**: 1–4.

       Diesh C, Stevens GJ, Xie P, De Jesus Martinez T, Hershberg EA, Leung A, Guo E, Dider S,
690   **Zhang J, Bridge C, *et al.* 2023**. JBrowse 2: a modular genome browser with views of synteny and structural variation. *Genome Biology* **24**: 1–21.

692   **Dubin MJ, Mittelsten Scheid O, Becker C**. **2018**. Transposons: a blessing curse. *Current Opinion in Plant Biology* **42**: 23–29.

694   **Ferrafiat L, Pflieger D, Singh J, Thieme M, Böhrer M, Himber C, Gerbaud A, Bucher E, Pikaard CS, Blevins T**. **2019**. The NRPD1 N-terminus contains a Pol IV-specific motif that is critical
696   for genome surveillance in Arabidopsis. *Nucleic acids research* **47**: 9037–9052.

       Flavell AJ. **1984**. Role of reverse transcription in the generation of extrachromosomal copia mobile
698   genetic elements. *Nature* **310**: 514–516.

       Flavell AJ, Ish-horowicz D. **1981**. Extrachromosomal circular copies of the eukaryotic transposable
700   element copia in cultured Drosophila cells. *Nature 1981 292:5824* **292**: 591–595.

       Garfinkel DJ, Stefanisko KM, Nyswaner KM, Moore SP, Oh J, Hughes SH. **2006**.
Retrotransposon Suicide: Formation of Ty1 Circles and Autointegration via a Central DNA Flap. *Journal of Virology* **80**: 11920.

Gaubatz JW. **1990**. Extrachromosomal circular DNAs and genomic sequence plasticity in eukaryotic cells *. *Mutation Research* **237**: 271–292.

Gautier M, Vitalis R. **2012**. rehh: an R package to detect footprints of selection in genome-wide SNP data from haplotype structure. *Bioinformatics (Oxford, England)* **28**: 1176–1177.

Gilbert C, Peccoud J, Cordaux R. **2021**. Transposable Elements and the Evolution of Insects. *https://doi.org/10.1146/annurev-ento-070720-074650* **66**: 355–372.

Gordon SP, Contreras-Moreira B, Levy JJ, Djamei A, Czedik-Eysenberg A, Tartaglio VS, Session A, Martin J, Cartwright A, Katz A, *et al.* 2020. Gradual polyploid genome evolution
revealed by pan-genomic analysis of Brachypodium hybridum and its diploid progenitors. *Nature Communications* **11**.

Gordon SP, Contreras-Moreira B, Woods DP, Des Marais DL, Burgess D, Shu S, Stritt C, Roulin AC, Schackwitz W, Tyler L, *et al.* 2017. Extensive gene content variation in the
Brachypodium distachyon pan-genome correlates with population structure. *Nature Communications* **8**: 1–13.

Grandbastien MA. **2015**. LTR retrotransposons, handy hitchhikers of plant regulation and stress response. *Biochimica et Biophysica Acta (BBA) - Gene Regulatory Mechanisms* **1849**: 403–416.

Grant CE, Bailey TL, Noble WS. **2011**. FIMO: scanning for occurrences of a given motif. *Bioinformatics* **27**: 1017–1018.

Hasterok R, Catalan P, Hazen SP, Roulin AC, Vogel JP, Wang K, Mur LAJ. **2022**. Brachypodium: 20 years as a grass biology model system; the way forward? *Trends in Plant Science* **27**: 1002–1016.

Hawkins JS, Kim HR, Nason JD, Wing RA, Wendel JF. **2006**. Differential lineage-specific amplification of transposable elements is responsible for genome size variation in Gossypium. *Genome Research* **16**: 1252.

Hirochika H, Sugimoto K, Otsuki Y, Tsugawa H, Kanda M. **1996**. Retrotransposons of rice involved in mutations induced by tissue culture. *Proceedings of the National Academy of Sciences of the United States of America* **93**: 7783–7788.

Hong SY, Seo PJ, Yang MS, Xiang F, Park CM. **2008**. Exploring valid reference genes for gene expression studies in Brachypodium distachyon by real-time PCR. *BMC Plant Biology* **8**: 112.

Horvath R, Minadakis N, Bourgeois Y, Roulin AC. **2023**. The evolution of transposable elements in Brachypodium distachyon is governed by purifying selection, while neutral and adaptive processes play a minor role. *bioRxiv*: 2023.09.15.557873.

International Brachypodium Initiative. **2010**. Genome sequencing and analysis of the model grass Brachypodium distachyon. *Nature* **463**: 763–768.

Ito H, Gaubert H, Bucher E, Mirouze M, Vaillant I, Paszkowski J. **2011**. An siRNA pathway prevents transgenerational retrotransposition in plants subjected to stress. *Nature* **472**: 115–120.

Jeong H-H, Yalamanchili HK, Guo C, Shulman JM, Liu Z. **2018**. An ultra-fast and scalable quantification pipeline for transposable elements from next generation sequencing data. *Biocomputing 2018*: 168–179.

Jin J, Tian F, Yang DC, Meng YQ, Kong L, Luo J, Gao G. **2017**. PlantTFDB 4.0: Toward a central hub for transcription factors and regulatory interactions in plants. *Nucleic Acids Research* **45**: D1040–D1045.

Kawakatsu T, Huang SS, Jupe F, Sasaki E, Schmitz RJ, Urich MA, Castanon R, Nery JR, Barragan C, He Y, *et al.* 2016. Epigenomic Diversity in a Global Collection of Arabidopsis thaliana Accessions. *Cell* **166**: 492–505.

Kidwell MG, Lisch D. **1997**. Transposable elements as sources of variation in animals and plants. *Proceedings of the National Academy of Sciences of the United States of America* **94**: 7704–7711.

Lanciano S, Carpentier MC, Llauro C, Jobet E, Robakowska-Hyzorek D, Lasserre E, Ghesquiere A, Panaud O, Mirouze M. **2017**. Sequencing the extrachromosomal circular mobilome reveals retrotransposon activity in plants. *PLoS Genet* **13**: e1006630.

Lanciano S, Cristofari G. **2020**. Measuring and interpreting transposable element expression. *Nature Reviews Genetics 2020 21:12* **21**: 721–736.

Lanciano S, Mirouze M. **2018**. Transposable elements: all mobile, all different, some stress responsive, some adaptive? *Current Opinion in Genetics & Development* **49**: 106–114.

Langmüller AM, Langmüller L, Nolte V, Dolezal M, Schï Otterer C. **2023**. The genomic distribution of transposable elements is driven by spatially variable purifying selection. *Nucleic Acids Research* **51**: 9203–9213.

Lee SC, Ernst E, Berube B, Borges F, Parent JS, Ledon P, Schorn A, Martienssen RA. **2020**. Arabidopsis retrotransposon virus-like particles and their regulation by epigenetically activated small RNA. *Genome Research* **30**: 576–588.

Leyser O. **2018**. Auxin Signaling. *Plant Physiology* **176**: 465–479.

Li H, Durbin R. **2009**. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* **25**: 1754–1760.

766 **Lisch D**. **2013**. How important are transposons for plant evolution? *Nat Rev Genet* **14**: 49–61.

Liu P, Cuerda-Gil D, Shahid S, Slotkin RK. **2022**. The Epigenetic Control of the Transposable
768 Element Life Cycle in Plant Genomes and Beyond. *https://doi.org/10.1146/annurev-genet-072920-015534* **56**: 63–87.

770 **Liu P, Nie WF, Xiong X, Wang Y, Jiang Y, Huang P, Lin X, Qin G, Huang H, Niu Q, *et al.* 2021a**.
A novel protein complex that regulates active DNA demethylation in Arabidopsis. *Journal of*
772 *Integrative Plant Biology* **63**: 772–786.

Liu P, Nie WF, Xiong X, Wang Y, Jiang Y, Huang P, Lin X, Qin G, Huang H, Niu Q, *et al.* 2021b.
774 A novel protein complex that regulates active DNA demethylation in Arabidopsis. *Journal of*
Integrative Plant Biology **63**: 772–786.

776 **Makarevitch I, Waters AJ, West PT, Stitzer M, Hirsch CN, Ross-Ibarra J, Springer NM**. **2015**.
Transposable elements contribute to activation of maize genes in response to abiotic stress. *PLoS*
778 *Genet* **11**: e1004915.

McClintock B. **1984**. The significance of responses of the genome to challenge. *Science* **226**: 792–
780 801.

Minadakis N, Williams H, Horvath R, Caković D, Stritt C, Thieme M, Bourgeois Y, Roulin AC.
782 **2023a**. New resources for environmental genomics in the wild Mediterranean grass B. distachyon.
bioRxiv: 2023.06.01.543285.

784 **Minadakis N, Williams H, Horvath R, Caković D, Stritt C, Thieme M, Bourgeois Y, Roulin AC**.
2023b. The demographic history of the wild crop relative Brachypodium distachyon is shaped by
786 distinct past and present ecological niches. *bioRxiv*: 2023.06.01.543285.

Mirouze M, Reinders J, Bucher E, Nishimura T, Schneeberger K, Ossowski S, Cao J, Weigel D,
788 **Paszkowski J, Mathieu O**. **2009**. Selective epigenetic control of retrotransposition in Arabidopsis.
Nature **461**: 427–430.

790 **Mirouze M, Vitte C**. **2014**. Transposable elements, a treasure trove to decipher epigenetic variation:
insights from Arabidopsis and crop epigenomes. *J Exp Bot* **65**: 2801–2812.

792 **Miura A, Yonebayashi S, Watanabe K, Toyama T, Shimada H, Kakutani T**. **2001**. Mobilization
of transposons by a mutation abolishing full DNA methylation in Arabidopsis. *Nature* **411**: 212–214.

794 **Muszewska A, Steczkiewicz K, Stepniewska-Dziubinska M, Ginalski K**. **2019**. Transposable
elements contribute to fungal genes and impact fungal lifestyle. *Scientific Reports 2019 9:1* **9**: 1–10.

796 **Nakagawa S, Schielzeth H**. **2013**. A general and simple method for obtaining R2 from generalized
linear mixed-effects models. *Methods in Ecology and Evolution* **4**: 133–142.

798 **Negi P, Rai AN, Suprasanna P**. **2016**. Moving through the stressed genome: Emerging regulatory
roles for transposons in plant stress response. *Frontiers in Plant Science* **7**: 1448.

800 **Piegu B, Guyot R, Picault N, Roulin A, Sanyal A, Kim H, Collura K, Brar DS, Jackson S, Wing**
RA, *et al.* 2006. Doubling genome size without polyploidization: dynamics of retrotransposition-
802 driven genomic expansions in Oryza australiensis, a wild relative of rice. *Genome Res* **16**: 1262–
1269.

804 **Prjibelski A, Antipov D, Meleshko D, Lapidus A, Korobeynikov A**. 2020. Using SPAdes De Novo
Assembler. *Current Protocols in Bioinformatics* **70**: e102.

806 **Quadrana L, Silveira AB, Mayhew GF, Leblanc C, Martienssen RA, Jeddeloh JA, Colot V**.
2016. The Arabidopsis thaliana mobilome and its impact at the species level.

808 **Quinlan AR, Hall IM**. **2010**. BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics* **26**: 841–842.

810 **R Core Team**. **2020**. R: A Language and Environment for Statistical Computing.

Ramírez F, Ryan DP, Grüning B, Bhardwaj V, Kilpert F, Richter AS, Heyne S, Dündar F, Manke
812 **T**. **2016**. deepTools2: a next generation web server for deep-sequencing data analysis. *Nucleic Acids Research* **44**: W160–W165.

814 **Raúl C, Noemia M-D, Sonal G, Michael P, M. CJ**. **2023**. Transposons are important contributors to gene expression variability under selection in rice populations. *eLife* **12**.

816 **Ream TS, Haag JR, Wierzbicki AT, Nicora CD, Norbeck AD, Zhu JK, Hagen G, Guilfoyle TJ, Pasa-Tolic L, Pikaard CS**. **2009**. Subunit compositions of the RNA-silencing enzymes Pol IV and
818 Pol V reveal their origins as specialized forms of RNA polymerase II. *Mol Cell* **33**: 192–203.

Rey O, Danchin E, Mirouze M, Loot C, Blanchet S. **2016**. Adaptation to Global Change: A
820 Transposable Element-Epigenetics Perspective. *Trends in Ecology and Evolution* **31**: 514–526.

Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G, Mesirov JP.
822 **2011**. Integrative genomics viewer. *Nature Publishing Group*.

Roquis D, Robertson M, Yu L, Thieme M, Julkowska M, Bucher E. **2021**. Genomic impact of
824 stress-induced transposable element mobility in Arabidopsis. *Nucleic Acids Research*.

RStudio Team. **2016**. RStudio: Integrated Development Environment for R.

826 **Rymen B, Ferrafiat L, Blevins T**. **2020**. Non-coding RNA polymerases that silence transposable elements and reprogram gene expression in plants. *Transcription* **11**: 172–191.

828 **Saze H, Tsugane K, Kanno T, Nishimura T**. **2012**. DNA Methylation in Plants: Relationship to Small RNAs and Histone Modifications, and Functions in Transposon Inactivation. *Plant and Cell*
830 *Physiology* **53**: 766–784.

Schulman AH. **2013**. Retrotransposon replication in plants. *Current Opinion in Virology* **3**: 604–614.

832 **Senft AD, Macfarlan TS**. **2021**. Transposable elements shape the evolution of mammalian development. *Nature Reviews Genetics 2021 22:11* **22**: 691–711.

834 **Shen W, Le S, Li Y, Hu F**. **2016**. SeqKit: A Cross-Platform and Ultrafast Toolkit for FASTA/Q File Manipulation.

836 **Sigman MJ, Slotkin RK**. **2016**. The First Rule of Plant Transposable Element Silencing: Location, Location, Location. *The Plant Cell* **28**: 304–313.

838 **Silander K, Saarela J**. **2008**. Whole genome amplification with Phi29 DNA polymerase to enable genetic or genomic analysis of samples of low DNA yield. *Methods in molecular biology (Clifton,*
840 *N.J.)* **439**: 1–18.

Skalska A, Stritt C, Wyler M, Williams HW, Vickers M, Han J, Tuna M, Tuna GS, Susek K,
842 **Swain M, *et al.* 2020**. Genetic and methylome variation in Turkish brachypodium distachyon accessions differentiate two geographically distinct subpopulations. *International Journal of*
844 *Molecular Sciences* **21**: 1–17.

Smith CA, Vinograd J. **1972**. Small polydisperse circular DNA of HeLa cells. *Journal of molecular*
846 *biology* **69**.

Stitzer MC, Anderson SN, Springer NM, RossIbarra J. **2021**. The genomic ecosystem of
848 transposable elements in maize. *PLoS Genetics* **17**: e1009768.

Stonaker JL, Lim JP, Erhard KF, Hollick JB. **2009**. Diversity of Pol IV Function Is Defined by Mutations at the Maize rmr7 Locus. *PLOS Genetics* **5**: e1000706.

Stritt C, Gimmi EL, Wyler M, Bakali AH, Skalska A, Hasterok R, Mur LAJ, Pecchioni N, Roulin AC. **2022**. Migration without interbreeding: Evolutionary history of a highly selfing Mediterranean grass inferred from whole genomes. *Molecular Ecology* **31**: 70–85.

Stritt C, Gordon SP, Wicker T, Vogel JP, Roulin AC. **2018**. Recent Activity in Expanding Populations and Purifying Selection Have Shaped Transposable Element Landscapes across Natural Accessions of the Mediterranean Grass Brachypodium distachyon. *Genome Biology and Evolution* **10**: 304–318.

Stritt C, Thieme M, Roulin AC. **2021**. Rare transposable elements challenge the prevailing view of transposition dynamics in plants. *American journal of botany* **108**: 1310–1314.

Stritt C, Wyler M, Gimmi EL, Pippel M, Roulin AC. **2020**. Diversity, dynamics and effects of long terminal repeat retrotransposons in the model grass *Brachypodium distachyon*. *New Phytologist* **227**: 1736–1748.

Stuart T, Eichten SR, Cahn J, Karpievitch Y V, Borevitz JO, Lister R. **2016**. Population scale mapping of transposable element diversity reveals links to gene regulation and epigenomic variation. *Elife* **5**.

Supek F, Bošnjak M, Škunca N, Šmuc T. **2011**. REVIGO Summarizes and Visualizes Long Lists of Gene Ontology Terms (C Gibas, Ed.). *PLoS ONE* **6**: e21800.

Suzuki R, Shimodaira H. **2006**. Pvclust: an R package for assessing the uncertainty in hierarchical clustering. *Bioinformatics (Oxford, England)* **22**: 1540–1542.

Tanskanen JA, Sabot F, Vicient C, Schulman AH. **2007**. Life without GAG: the BARE-2 retrotransposon as a parasite's parasite. *Gene* **390**: 166–174.

Tello-Ruiz MK, Naithani S, Gupta P, Olson A, Wei S, Preece J, Jiao Y, Wang B, Chougule K, Garg P, *et al.* 2021. Gramene 2021: Harnessing the power of comparative genomics and pathways for plant research. *Nucleic Acids Research* **49**: D1452–D1463.

Thieme M, Brêchet A, Bourgeois Y, Keller B, Bucher E, Roulin AC. **2022**. Experimentally heat-induced transposition increases drought tolerance in Arabidopsis thaliana. *New Phytologist* **236**: 182–194.

Thieme M, Lanciano S, Balzergue S, Daccord N, Mirouze M, Bucher E. **2017**. Inhibition of RNA polymerase II allows controlled mobilisation of retrotransposons for plant breeding. *Genome Biol* **18**: 134.

Tian T, Liu Y, Yan H, You Q, Yi X, Du Z, Xu W, Su Z. **2017**. AgriGO v2.0: A GO analysis toolkit for the agricultural community, 2017 update. *Nucleic Acids Research* **45**: W122–W129.

Tittel-Elmer M, Bucher E, Broger L, Mathieu O, Paszkowski J, Vaillant I. **2010**. Stress-Induced Activation of Heterochromatic Transcription. *PLOS Genetics* **6**: e1001175.

Tsukahara S, Kobayashi A, Kawabe A, Mathieu O, Miura A, Kakutani T. **2009**. Bursts of retrotransposition reproduced in Arabidopsis. *Nature* **461**: 423–426.

Uzunović J, Josephs EB, Stinchcombe JR, Wright SI, Parsch J. **2019**. Transposable Elements Are Important Contributors to Standing Variation in Gene Expression in Capsella Grandiflora. *Molecular Biology and Evolution* **36**: 1734–1745.

890 **Venner S, Feschotte C, Biémont C**. **2009**. Dynamics of transposable elements: towards a community ecology of the genome. *Trends in Genetics* **25**: 317–323.

892 **Vitte C, Fustier MA, Alix K, Tenaillon MI**. **2014**. The bright side of transposons in crop evolution. *Briefings in Functional Genomics and Proteomics* **13**: 276–295.

894 **Vu VQ**. **2011**. ggbiplot: A ggplot2 based biplot. R package version 0.55.

Wang D, Zheng Z, Li Y, Hu H, Wang Z, Du X, Zhang S, Zhu M, Dong L, Ren G, *et al.* 2021. 896 Which factors contribute most to genome size variation within angiosperms? *Ecology and Evolution* **11**: 2660–2668.

898 **Wicker T, Sabot F, Hua-Van A, Bennetzen JL, Capy P, Chalhoub B, Flavell A, Leroy P, Morgante M, Panaud O, *et al.* 2007**. A unified classification system for eukaryotic transposable 900 elements. *Nature Reviews Genetics* **8**: 973–982.

Wickham H. **2016**. ggpolt2 Elegant Graphics for Data Analysis. *Use R! series*: 211.

902 **Xu L, Yuan K, Yuan M, Meng X, Chen M, Wu J, Li J, Qi Y**. **2020**. Regulation of Rice Tillering by RNA-Directed DNA Methylation at Miniature Inverted-Repeat Transposable Elements. *Molecular* 904 *plant* **13**: 851–863.

Yang F, Su W, Chung OW, Tracy L, Wang L, Ramsden DA, Zhang ZZ. **2023a**. Retrotransposons 906 hijack alt-EJ for DNA replication and eccDNA biogenesis. *Nature 2023*: 1–8.

Yang L-L, Zhang X-Y, Wang L-Y, Li Y-G, Li X-T, Yang Y, Su Q, Chen N, Zhang Y-L, Li N, *et* 908 *al.* 2023b. Lineage-specific amplification and epigenetic regulation of LTR-retrotransposons contribute to the structure, evolution, and function of Fabaceae species. *BMC Genomics 2023 24:1* 910 **24**: 1–15.

Zhang H, Lang Z, Zhu JK. **2018**. Dynamics and function of DNA methylation in plants. *Nature* 912 *Reviews Molecular Cell Biology 2018 19:8* **19**: 489–506.

Zhang Y, Li Z, Liu J, Zhang Y, Ye L, Peng Y, Wang H, Diao H, Ma Y, Wang M, *et al.* 2022. 914 Transposable elements orchestrate subgenome-convergent and -divergent transcription in common wheat. *Nature Communications* **13**.

916 **Zhong X, Du J, Hale CJ, Gallego-Bartolome J, Feng S, Vashisht AA, Chory J, Wohlschlegel JA, Patel DJ, Jacobsen SE**. **2014**. Molecular Mechanism of Action of Plant DRM De Novo DNA 918 Methyltransferases. *Cell* **157**: 1050–1060.

Zhou X, Stephens M. **2012**. Genome-wide efficient mixed-model analysis for association studies. 920 *Nature Genetics 2012 44:7* **44**: 821–824.

Supporting Information

S1 Fig. Correlation of bioclimatic variables with pCNV when not correcting for population structure. Colors and sizes
of bubbles show the part of the variance (R^2) explained by the bioclimatic variables in %

956 **S2 Fig. TFs binding to *ONSEN* are heat-stress inducible.** GO-enrichment analysis of transcription factors for which binding sites have been detected in AT1G11265, a member of the heat-responsive *ONSEN* (*ATCOPIA78*) LTR-RT family in
958 *A. thaliana*. Colors indicate number of TF-binding sites found. GO terms that occur at least six times are highlighted in the plot. All GO-terms and their number of occurrences is listed in **S1 Table**.

964   **S3 Fig**

S4 Fig

S3-5 Figs. TIPs of *HOPPLA* in one of the resequenced *Bd nrpd1-2 (-/-)* plants. JBrowse screenshot of the insertion site in *Bd nrpd1-2 (-/-)* (bottom) compared to the Bd21-3 wt (top). The target side duplication (TSD) is annotated and soft clipped parts of reads are coloured.

S6 Fig. Manhattan plots of the GWASs of pCNVs of four recently active LTR-RT families. From top: RLG_BdisC004, RLC_BdisC030 RLC_BdisC209 and RLC_BdisC022. The two significance levels, false discovery rate < 0.05 (dashed line) and Bonferroni correction (solid line) are depicted

990   **S1 Table. REVIGO output of the processing of GO terms of transcription factors for which binding sites have been detected in the *HOPPLA* and *ONSEN* consensus sequences.**

992   **S2 Table. Normalized pCNV of LTR-RTs and bioclimatic variables of the 320 natural accessions of *B. distachyon***

994   **S3 Table. Gene list of pCNV GWAS with different levels of significance (FDR < 0.05, BC) and window sizes (20 kb, 50 kb)**

996   **S4 Table. Components of the Pol IV and Pol IV holoenzymes in *B. distachyon***

       S5 Table. Sequences of oligos used in this study