HCN channels regulate excitability and gate synaptic amplification via NMDA in mouse L2/3 pyramidal neurons

Viktor János Oláh[1], Jing Wu[2], Leonard K. Kaczmarek[2,3] & Matthew JM Rowan[1,4*]

[1]Department of Cell Biology, Emory University School of Medicine, Atlanta, GA, 30322, USA

[2]Department of Pharmacology, Yale University School of Medicine, New Haven, Connecticut 06520

[3]Department of Cellular and Molecular Physiology, Yale University School of Medicine

[4]Center for Neurodegenerative Disease, Emory University School of Medicine

*Address for Correspondence:
Matt Rowan, PhD
Emory University School of Medicine
Department of Cell Biology
Atlanta, GA, 30322
Email: mjrowan@emory.edu

Citations: 72; *Figures*: 7 main figures; 4 supplementary figures

ABSTRACT

Layer 2/3 pyramidal cells (L2/3 PCs) play a crucial role in cortical information transfer. Although the dendritic arbors of L2/3 PCs are impressive, they lack the distinct anatomical compartments characteristic of deeper L5 PCs. For example, many L2/3 PCs do not display an apparent distal tuft region. However, L2/3 PCs receive inputs from both thalamic (bottom-up) and cortical (top-down) inputs, preferentially synapsing onto their proximal and distal dendrites, respectively. Nonuniform organization of channels and NMDA receptors in L2/3 dendrites could serve to independently modulate these information streams to affect learning and behavior, yet whether L2/3 PC dendrites possess this capability has not been established. Here, we found a previously unappreciated non-uniform HCN channel distribution in L2/3 PCs, allowing for pathway-specific gating of NMDA receptor recruitment at bottom-up (proximal) but not top-down (distal) synapses. Interestingly, HCN availability could be regulated via neuromodulation, suggesting that the gain of thalamic and cortical-cortical signals in L2/3 may be independently modified *in vivo*.

INTRODUCTION

Layer 2/3 pyramidal cells (L2/3 PCs) are the most numerous excitatory neurons in the neocortex. As a major component of the canonical circuit, L2/3 PCs are thought to play crucial roles in information transfer and gain control of cortical output(1). Thus, disambiguating the afferent circuit connections of L2/3 PCs has been a significant focus of neuroscience research in the past decade(2-5). However, understanding cellular mechanisms by which L2/3 PCs transform information is equally important (6-9). Like their more extensively studied counterparts (e.g., L5 PCs, hippocampal CA1 PCs), L2/3 PCs are equipped with an array of dendritic nonlinearities(10, 11), enabling complex subcellular computations. Nonetheless, L2/3 PCs exhibit marked disparities from other PC types as well. Their morphology is less segmented, due to the lack of a distinctive apical tuft (except for deep layer 3 cells (12)). Their dendritic protrusions are also shorter, and consequently possess less surface area over which specialized nonlinear operations may be actualized. Therefore, the applicability of currently established neuronal dynamics to L2/3 PCs is questionable.

Dendritic voltage-gated channels and receptors are crucial for diversifying the computational power of neuronal cells, by allowing for nonlinear processing of arriving synaptic inputs(13). Nonlinear processing is essential for proper brain function and has been implicated in critical tasks such as conscious perception(14, 15) and processing of sensory and motor input(16). Therefore, a more comprehensive understanding of subcellular signal processing dynamics in L2/3 PCs is warranted. Dendritic hyperpolarization-activated nonselective cation (HCN) channels are a staple of several PC classes(17, 18), playing essential roles in regulating resting membrane potential, the normalization of synaptic events arriving at spatially mismatched locations, and establishing oscillation frequency-selectivity(19). Beyond exerting control over the general excitability of neurons, HCN channels contribute to synaptic plasticity and regulate working memory(19). In following, HCN dysfunction contributes to mechanisms of epilepsy and pain(20). Electrophysiological detection of this current often relies on identification of a characteristic 'sag' potential, elicited by hyperpolarizing current injections. In L2/3 PCs, this sag potential is negligible in most cortical areas (but see (21)) thus, HCN channels are often stated as missing from L2/3 PCs. Therefore, the functional expression of HCN channels in these cells remains ambiguous.

Here we report that L2/3 PCs throughout the cortex express functionally relevant HCN channels with surprising functions. We found that HCN channels robustly influence the overall excitability of L2/3 PCs. Interestingly, we found that HCN channel availability in L2/3 PCs is biased in a unique proximal somato-dendritic manner, which is inverted with respect to L5 and hippocampal CA1 PCs, where HCN is enriched within their distal apical tufts(18, 22). In cooperation with NMDA receptors, this L2/3 HCN organization could modulate the boosting of proximal 'bottom-up' dendritic

inputs, without affecting distal 'top-down' dendritic inputs. Furthermore, we found that HCN

expression is robustly upregulated during early postnatal development and that neuromodulation in

adulthood can rapidly adjust HCN channel availability. Our results demonstrate that L2/3 PCs not only

express dendritic HCN channels, but also employ these conductances in a previously unappreciated

manner.

RESULTS

HCN channel expression in L2/3 PCs

HCN channels have been extensively documented and rigorously characterized in several cortical PC

types(17), however, whether PCs residing in layer 2/3 of the mouse cortex express functionally relevant

HCN channels remains ambiguous. Nonetheless, open-source snRNAseq (23) shows comparable HCN

subunit mRNA levels in all cortical PC types of the visual cortex (Fig. 1a). Therefore, we investigated

whether this mRNA signature translated to HCN surface expression. Immunohistochemical staining

against HCN1 showed discernible somatic fluorescent signal in many cells residing in superficial

layers of the cortex (Fig. 1b) along with staining of putative dendrites. To examine the potential

functional role for HCN channels in L2/3 PCs, we performed *ex vivo* cell-attached recordings from

visually identified L2/3 PCs in the primary visual cortex from mature mice. Spike output upon

extracellular stimulation (50 Hz) in layer 4 was then measured in control conditions and following

pharmacological HCN channel block (Fig. 1c). We found that bath application of Cs^+ (2mM), a

conventional blocker of HCN channels at this concentration, significantly increased their firing (Fig.

1d). This was unlikely due to presynaptic modulation of these synapses, as the paired-pulse ratio was

unchanged in a set of separate whole-cell recordings coupled with stimulation in layer 4 (Fig. 1e).

Together, these results suggest that L2/3 PCs express a functionally relevant set of HCN channels

which limit excess AP firing following synaptic stimulation.

HCN channels constrain the overall excitability of L2/3 PCs

To investigate the role of HCN channels in regulating the excitability of L2/3 PCs, we next performed

further L2/3 PC whole-cell recordings. A conventionally used assessment of HCN channel function is

the characterization of the sag potential(17). This hallmark of I_h activation is measured as the amount

of rectification detected in response to hyperpolarizing current injections. In line with previous reports,

we found that L2/3 PCs exhibited an unremarkable amount of sag potential at 'typical' current

commands. However, upon larger negative current injections, a discernable rectification was revealed

(Fig. 2a). In addition, the rectification of the steady-state responses was abolished in the presence of

external Cs^+ which could be restored by artificial I_h supplementation by dynamic clamp (Fig. 2b).

Similarly, sag was non-detectable when I_h was blocked, which could then be restored by dynamic

clamp (Fig. 2c,d). Furthermore, we found that passive electrical properties were also significantly

modulated by HCN blockade (Fig. 2e,f). Together, these results explain the dramatic effect of Cs^+ on

excitability (Fig. 1). The non-apparent sag potential at conventional hyperpolarizing current steps is

thus surprising, suggesting a markedly faster occurrence of the I_h effect exists in L2/3 PCs which

occludes the biphasic nature of the sag response. To investigate this possibility, we measured the time-

constant of the immediate voltage responses. We found a strong positive correlation between the

amplitude of a passive current command and the voltage response time constant, which was abolished

by Cs^+ bath application (Fig. 2g). This confirms that I_h has a strong effect during the initial phase of

hyperpolarizing voltage responses which may contribute to the lack of detectable sag potential at

modest current injections. However, other biophysical mechanisms may also contribute to the muted

sag effect in L2/3 PCs.

Next, we examined the voltage dependence and kinetic properties of I_h in L2/3 PCs in voltage clamp

(Fig. 2h). Significant Cs^+-sensitive (1mM) currents were activated at subthreshold membrane

potentials, with half-activation voltages more negative than the resting membrane potential, and with

relatively fast activation kinetics (Fig. 2i,j). Overall the pharmacological profile, effects on passive

electrical properties, and voltage dependence all point to previously published data describing HCN

channels (24). Together these results indicate that L2/3 PCs express a functionally relevant population

of HCN channels, which constrain the excitability of these cells. Similar voltage clamp responses were

observed beyond the visual cortex (e.g., in S1, M1, and A1; Supplementary Fig. 1) suggesting that

HCN channels are a consistent feature in L2/3 PCs across sensory cortices.

Noncanonical subthreshold signal processing in L2/3 PCs by I_h

128   Electrical signals at different dendritic compartments undergo distance-dependent distortion (22, 25),

129   which has a major effect on neuronal function in cells with extensive dendritic branching (26). To

130   counteract these effects, larger L5 PCs and CA1 PCs express HCN channels with nonuniform

131   distributions (18, 24). As L2/3 PCs have a much more compact morphology, synaptic input distortions

might be less severe, raising the possibility of an alternative function of HCN channels in these cells. Thus, to investigate the distance-dependence of HCN on L2/3 synaptic inputs we made whole-cell patch recordings from Alexa-594 filled L2/3 PCs, and performed extracellular minimal stimulation at different dendritic locations guided with 2P microscopy (Fig. 3a). Elicited EPSPs likely reflect the activation of a single axon as stimulus intensity was set to achieve an ~50% failure rate(27). As HCN deactivation has been shown to strongly impact the duration of synaptic events(22, 25), we quantified changes in the half-width of locally evoked EPSPs, here using the HCN-selective blocker ZD-7288 (50 µM). Surprisingly, our results were in contrast to previous reports in L5 and CA1 PCs, in that I_h block had a far smaller effect on the time-course of distally-evoked (>100 µm distance from the soma) synaptic responses. In contrast, proximal inputs (<100 µm) were markedly more broadened after I_h blockade (Fig. 3b,c). These results suggest that L2/3 PCs are equipped with a distinctive subcellular HCN distribution, potentially contributing to noncanonical distance-dependent input modulation along the dendritic axis of L2/3 cells.

Pathway-specific modulation of synaptic information

L2/3 PCs receive information from distinct brain regions, such as "top-down" (cortical-cortical) axons, generally arriving to layer 1(28) and "bottom-up" (thalamocortical/layer 4) axons, terminating on proximal dendrites located in layer 4 and 2/3(29). Due to the stronger proximal effect of HCN on L2/3 PC EPSPs, we explored whether this modulation yielded input-pathway-specific differences. Whole-cell recordings were again obtained from L2/3 PCs, and EPSPs were elicited by minimal stimulation with electrodes situated to preferentially stimulate either layer 4 or layer 1 axons (electrode placement ~200µm laterally from recorded PC in either layer). In a subset of our recordings stimulating layer 4 inputs, evoked EPSPs were substantially larger than spontaneously occurring events (Fig. 3d). We deemed these as inputs arriving directly from the lateral geniculate nucleus axons (putative direct thalamocortical input(30) from the LGN, Supplementary Figure 2.). I_h blockade significantly broadened elicited EPSPs when the stimulation electrode was placed in layer 4, but not in layer 1 (Fig. 3e). Consequently, the voltage integral of EPSPs was significantly larger for L4 inputs compared to those in L1 (Fig. 3f). The effect of Cs^+ on EPSP half-width and voltage integral was similar for both L4 and putative LGN inputs, suggesting intermingled termination zones(29). Together these results suggest that HCN channels preferentially regulate bottom-up nonlinear signaling in proximal dendritic regions in L2/3 PCs.

HCN channels are notable for their ability to restrain temporal integration of synaptic inputs(22).

Therefore, we repeated these experiments using more high-frequency stimulation (5 stimuli at 50 Hz;

Fig. 3g). In accordance with the effect on single EPSPs, summation of L1 (i.e., 'distal) inputs was

unaltered with HCN block. In contrast, HCN blockade led to increased temporal summation of L4

inputs (Fig. 3h,i). However, summation of putative LGN inputs was not affected by HCN channel

block, suggesting further diversification of active properties even between L4 and LGN inputs as well.

I_h channels constrain NMDA receptor-dependent synaptic boosting

In a passive environment, dendritic filtering (31) creates attenuated (i.e., wider) EPSPs when

originating further from the soma. Yet our results show an opposite phenomenon with HCN channels

blocked. It has been described that reduced HCN availability can facilitate dendritic boosting

mechanisms, via activation of either voltage-gated (32) or NDMA (33) channels. To investigate

possible contributions of dendritic voltage-dependent processes, we performed NEURON

computational simulations using a realistic L2/3 PC dendritic morphology (34). Inserting voltage-gated

sodium or potassium channels with various distributions could not recapitulate our distance-dependent

EPSP observations (Supplementary Figure 3.). Thus, we next investigated whether NMDA receptors

might play a role in the observed phenomenon, due to their described overwhelming contribution to *in*

vivo activity in L2/3 PCs (8, 35).

We first tested whether NMDA receptors contribute to spontaneously occurring synaptic inputs

(sEPSPs) in L2/3 PCs(36). Due to potential alterations in spontaneous release and overall circuit

activity, extracellularly applied NMDA blockers may introduce significant confounds for this line of

investigation, especially as spontaneous synaptic potentials show tremendous variability in frequency,

amplitude, and kinetics. To overcome this conundrum, we designed a custom-built pipette attachment

through which the intracellular solution could be stably exchanged during whole-cell recordings (37).

We recorded sEPSPs in L2/3 PCs first using control intracellular solution, which was then exchanged

to a 50 µM MK-801 (non-competitive NMDA receptor antagonist) containing intracellular solution

(Fig. 4a). To also capitalize on the voltage-sensitivity of NMDA receptors, average peak amplitudes

were recorded at two different membrane potentials for each intracellular condition; -80 mV and -55

mV (resting membrane potential, and a somatic membrane potential depolarized enough to partially

unblock NMDA receptors (38)). Due to the small nature of sEPSPs, quantification of other waveform

parameters (i.e., EPSP width) was avoided. After quantifying the mean sEPSP peak at -80 mV, we

calculated the expected (linear) sEPSP peak at -55 mV and compared this prediction to our

observations of -55 mV. We found that sEPSPs were significantly larger than predicted after

depolarization (Fig. 4b), alluding to the presence of a nonlinear boosting mechanism. We found that

intracellular MK-801 solution exchange could abolish this synaptic amplification (Fig. 4c), indicating

that NMDA receptors actively contribute to shaping naturally occurring sEPSPs in L2/3 PCs. The

effect of pipette perfusion itself on our measurements was negligible (Supplementary Figure 4)

Having established that NMDA receptors play an active role in modulating sEPSPs, we next explored

whether differential NMDA recruitment could explain the observed boosting of proximal synaptic

inputs following HCN blockade described earlier. Computational simulations (lacking HCN) (Fig. 4d)

revealed that a proximally-biased NMDA distribution, as seen in other PC types (39), could indeed

recapitulate our experimental findings in L2/3 PCs with high accuracy (Fig. 4e,f). Because HCN

channels can reduce NMDA recruitment during synaptic signaling, we next explored whether proximal

enrichment of HCN allows for bottom-up-specific gating of NMDA (Fig. 4g). To accomplish this, we

recorded L2/3 EPSPs following stimulation of both proximal L4 and distal L1 inputs, as described

earlier, during each recording and varied the background pharmacology. Intriguingly, L4-evoked

EPSPs were slightly narrower than L1 in control conditions, potentially owing to increased HCN in

proximal dendrites. This layer-specific relationship was inverted following extracellular Cs^+ to block

HCN (Fig. 4g,h). To confirm that this L4-specific EPSP broadening after HCN channel block was due

to NMDA-inducted boosting, as predicted in our simulations (Fig. 4f), we next examined the EPSPs

with Cs^+ and MK-801 together. In the presence of internal MK-801, the broadening effect of Cs^+ on

L4-evoked EPSPs was abolished, thereby reverting the L4-L1 EPSP relationship to a control-like state.

(Fig. 4g,h). Together, our modeling and experimental results indicate that NMDA-directed

amplification of L4 inputs is strongly constrained by a proximal HCN bias in L2/3 PCs dendrites,

allowing for pathway-specific modulation of 'bottom-up' inputs without affecting those arriving from

'top-down' L1 axons.

Noncanonical HCN distribution predicted in L2/3 PCs

We next carried out computational simulations for HCN in L2/3 PCs, based on our previous

observations and modeling. To reproduce the native relationship between EPSP halfwidth and

dendritic distance, HCN channels were inserted into the model with different distributions and

conductance densities (Fig 5a). Our simulations revealed that proximal bias in HCN channel

distributions could best recapitulate our observations of HCN on proximally and distally originating
EPSP halfwidths. providing the best fit to our measurements (Fig. 5b,c). According to these
simulations, HCN regulation of local input resistance constrained voltage-dependent NMDA receptor
activation. This is in line with the proximal HCN1 channel staining (Fig. 1) revealed by
immunohistochemistry. These results suggest that L2/3 PCs have a unique proximal HCN channel
bias, as previous experiments show a distal HCN expression in L5 PCs and CA1 PCs(18, 24). Possibly
due to the markedly different morphology, specifically the absence of an independent computational
tuft unit, HCN channels have different functions in L2/3 PCs. Thus, together with experimental
observations, these simulations indicate that HCN channels are well positioned to regulate bottom-up
proximal inputs in L2/3 PCs, contrary to their L5 PC counterparts, which express HCN distally to
counteract distance dependent passive filtering from top-down inputs.

Tight developmental upregulation of Layer 2/3 HCN channels

Several ion channels show postnatal maturation in their biophysical properties and surface
expression(40), raising the possibility of a different proximo-distal HCN bias in young L2/3 PCs
compared to adult animals. To examine this possibility, L2/3 PCs were recorded at three developmental
time points (1^{st}, 2^{nd} and 6^{th} postnatal week) in S1. Our initial assessment of HCN maturation showed
that a rapid developmental upregulation occurs between 1 and 2 weeks old. We opted to perform further
developmental experiments in primary somatosensory cortex, where cells already receive adequate
bottom-up information during the first and second postnatal week.

We found that 1 week old L2/3 PCs displayed markedly different passive properties, while 2-week-
old cells were quite similar to mature cells (Fig. 6a). The non-quantifiable sag potential in 1 week old
animals implies the lack of I_h current at this developmental time point, however, the disproportionately
large input resistance and elevated resting membrane potential also suggests that other background
leak channels may also be missing at this developmental window (Fig. 6b). Subsequent voltage clamp
recordings confirmed that L2/3 PCs in 1 week old animals do not possess a considerable amount of I_h
in contrast to 2 and 6 week old animals (Fig. 6c,d). We found that both resting membrane potential and
peak I_h conductance changes throughout development were tightly (negatively) correlated (Fig. 6e).
This provides additional evidence for parallel maturation of HCN with other leaky channels, as HCN
channel expression alone would be predicted to shift the membrane to more depolarized potentials
(Fig. 2e). Next, we mapped the distance-dependence dendritic EPSP kinetics in 1 week old animals

(Fig. 6f). We found a surprising 'non-Rallian' negative correlation between EPSP halfwidth and

stimulation distance from the soma (Fig. 6g, h), similar to our previous results from mature neurons

with HCN channels blocked (Fig. 3a,b). This suggests the presence of nonlinear boosting mechanisms

in proximal dendrites of developing L2/3 PCs, similar to mature cells, which are unabated due to the

lack of HCN at this developmental stage.

I_h dependent neuromodulation of L2/3 PC activity by serotonergic inputs

If the robust dendritic HCN availability in mature L2/3 PCs is unalterable, the ability of NMDA to

modulate bottom-up inputs might be inadequate. Whether the mature brain possesses mechanisms to

modulate HCN in L2/3 PCs is unknown. Nonetheless, previous work in cortical L5 PCs shows that I_h

currents can be modulated following 5-HT_7 serotonergic receptor activation(41). L2/3 PCs are

extensively innervated by serotonergic axons from the dorsal raphe (42), providing a potential

framework to modify HCN availability. Single-cell RNA-seq (23) showed that L2/3 PCs do express

5-HT_7 receptors, interestingly at higher levels than all other cortical PCs (Fig. 7a). We

pharmacologically evaluated a potential role of serotonergic HCN regulation by bath application of 5-

carboxamidotryptamine (5-CT; 30 nM), a specific 5-HT_7 agonist. 5-CT application significantly

reduced evoked firing of L2/3 PCs during extracellular stimulation in L4 (Fig. 7b). Current clamp

experiments revealed that the rectification of steady-state voltage responses to hyperpolarizing current

injection was amplified with 5-CT (Fig. 7c), likely due to the increased availability of HCN channels

due to a rightward shift in their activation voltage-dependence (Fig. 7d). 5-CT also reduced L2/3 input

resistance potentially via HCN regulation, however, did not yield a depolarizing effect on the resting

membrane potential nor affect AP threshold (Fig. 7e). Importantly, 5-CT had no impact on the firing

responses of L2/3 PCs when I_h currents were blocked with background Cs^+ throughout the experiment

(Fig. 7f). Furthermore, the dampening of neuronal output was not related to an downregulating effect

of 5-HT7 on NMDA receptors (Fig. 7g, (43)). Together these results indicate that neuromodulation

represents an endogenous mechanism by which HCN availability can be altered in L2/3 PCs.

DISCUSSION

Precise communication between cortical neurons relies on discrete pre- and postsynaptic forms of

information transfer at synapses. The majority of cortical synapses terminate along the dendritic tree

(44) where, based on local active and passive electrical properties, various types of nonlinear
transformations occur. The spatially and electrically disjointed nature of dendrites (45) from the rest
of the cell allows for local, soma-independent computation. For example, perturbing active dendritic
elements (i.e., local conductances) can impair several fundamental cortical circuit tasks (36) such as
perception (14), underlining the necessity for compartmentalized information processing. Further,
single-subunit model simulations are unable to reproduce *in vivo* observed activity patterns of L2/3
PCs(9). Here, by combining *ex vivo* current clamp, synaptic stimulation, and ion channel recordings
with empirically informed, biophysically realistic simulations, we demonstrate that L2/3 PCs in
sensory cortex express a previously unappreciated, proximally-biased HCN channel distribution. Due
to an apparent universal HCN expression in L2/3 cells from each region, this arrangement should allow
for modification of synaptic gain across all primary sensory areas of cortex. One of the basic functions
of L2/3 PCs in sensory cortex is to encode first-order information, such as the orientation of visual
input(2, 46, 47). Several investigations propose that arithmetic summation of a number of similarly
tuned synapses will dictate whether these neurons will fire in response to sensory input (48). However,
numerous dendritic nonlinearities have been established in L2/3 PCs, such as NMDA receptors (8)
sodium channels (11) and calcium channels (10, 49) which also contribute to this process. Here we
demonstrate that L2/3 PCs express a functionally relevant population of HCN channels well situated
to modify the gain of cortical sensory and contextual information processing.

We found that HCN channels in L2/3 could constrain fundamental aspects of cellular
excitability, such as input resistance and resting membrane potential. This is likely due to a strong
proximal somato-dendritic distribution of HCN channels, as observed in immunohistochemical
experiments here. Likely due to their influence on input resistance, HCN blockade resulted in greatly
increased AP firing during extracellular stimulation. L2/3 HCN channels strongly dampened excitatory
synaptic inputs, however, only to those arriving at proximal dendritic locations. In line with this result,
HCN channels reduced the timecourse of synaptic inputs following local stimulation in L4, but left
those following L1 stimulation unaffected. Together with additional input-specific pharmacological
experiments, NEURON simulations demonstrated that this input-specific synaptic phenomenon could
indeed be explained by a proximal HCN channel distribution in dendrites putatively situated in L4 and
L2/3. This was apparently only possible with a concurrent proximal NMDA recruitment bias. This
should not be confused with observed differences in NMDA recruitment along the axis of individual
L2/3 basal dendrites (36). Although a proximal-basal dendrite NMDA receptor distribution is in line
with findings from L5 PCs (50), HCN channels are instead enriched distal dendritic compartments in
these PC types (18, 24). Thus the unprecedented proximal HCN recruitment in L2/3 PCs permits

pathway-specific control of synaptic amplification, as bottom-up (i.e., L4 originating) inputs target

proximal dendritic compartments homogeneously (29). This was in opposition to top-down inputs,

whose axons largely target distal dendrites in L2/3 PCs (28, 51), as these synapses were unaffected by

HCN channel blockade. To fully understand how compartment-specific pathways are functionally

regulated, further anatomical and *in vivo* imaging experiments are needed to establish the distribution

and synaptic dynamics of L4 and L1 inputs along the L2/3 PC dendritic tree.

HCN channels appear well suited to regulate the synaptic and intrinsic properties of L2/3 PCs,

for example, if their availability is plastic or developmentally regulated. Previously, we found that

HCN channel availability was upregulated in GABAergic interneurons during development, resulting

in reduced propagation of subthreshold signals in mature axons (52). Interestingly, we found that L2/3

PCs in young animals (1 week old) lacked HCN channels. Together, these results suggest a universal,

cell-nonautonomous developmental HCN upregulation in both dendritic and axonal compartments. In

L2/3 PCs, the HCN channel maturation window was rapid and almost complete by the second postnatal

week. Therefore critical period plasticity (53) and developmental I_h upregulation occur during a similar

developmental window, suggesting a causal relationship. In addition to the robust developmental

regulation of HCN channels, we also found that enhancement of HCN channel availability was also

possible in the adult cortex, following serotonergic ($5\text{-}HT_7$) receptor-mediated biophysical regulation.

This was likely due to a metabotropic process that increases cytosolic cAMP levels by stimulating

adenylate cyclase(54) which in turn modulates HCN channel properties. This raises the intriguing

possibility that neuromodulation could induce nonstationary conditions in mature L2/3 circuits, for

example, in re-biasing of bottom-up vs. top-down information, experience-dependent plasticity (55),

or even for reopening of critical window-like states (56).

As L2/3 PCs are one of the most abundant cell populations of the neocortex, HCN channels in

these cells may exert overwhelming control over cortical excitability and thus cognition and memory.

Cell non-autonomous HCN channel dysregulation manifests in clear behavioral changes, such as

altered spatial learning (57) or impaired working memory (58). Curiously, higher HCN1 protein levels

in the postmortem brain also correlate with cognitive resilience in aging humans (59). Therapeutic

HCN channel modulation aimed at restoring healthy brain function in distinct pathophysiological states

has consequently garnered significant interest (60-62) as both HCN1 and HCN2 are linked to

neuropathic pain (63) and for treating major depressive disorder (61, 62, 64). Here we demonstrate that

activation of the serotonergic receptor $5\text{-}HT_7$ alters HCN channel function in L2/3 PCs, resulting in

reduced PC excitability, similar to L5 PCs (41). $5\text{-}HT_7$ receptor targeting agents are highly effective

psychopharmacological compounds. Agents such as the antidepressant *vortioxetine* has a relevant

therapeutic concentration that functionally suppresses 5-HT$_7$ (65), and the mood-stabilizing
antipsychotic agent *lurasidone* possesses an even higher binding affinity to 5-HT$_7$ (66). Commonly
used anesthetic drugs also modulate HCN channel action (67, 68), implying the need to investigate
anesthesia-related mechanisms and confounds in L2/3 in both therapeutic and experimental settings.
Together, we argue that HCN channels are present and functionally relevant in L2/3 PCs. As L2/3 PCs
are one of the largest cell populations in the brain, and as HCN channels are the target of several current
therapeutics, future research should consider their role in these cells in basic and clinical studies.
360
361 **METHODS**
362
Acute slice preparation

All animal procedures were approved by the Emory University IACUC. Acute slices from cortex were
prepared from mature C57Bl/6J mice (6–10 weeks old). Male and female mice were used for all
experiments with data collected from ≥3 mice per experimental condition. Mice were first anesthetized
and perfused with ice-cold cutting solution (in mM) 87 NaCl, 25 NaHO$_3$, 2.5 KCl, 1.25 NaH$_2$PO$_4$, 7
MgCl$_2$, 0.5 CaCl$_2$, 10 glucose, and 7 sucrose. Thereafter, mice were terminated by decapitation and the
brain immediately removed by dissection. Brain slices (250 μm) were sectioned in the coronal plane
using a vibrating blade microtome (VT1200S, Leica Biosystems) in the same solution. Slices were
transferred to an incubation chamber and maintained at 34°C for ~30 min and then at 23–24°C
thereafter. During whole-cell recordings, slices were continuously perfused with (in mM) 128 NaCl,
26.2 NaHO$_3$, 2.5 KCl, 1 NaH$_2$PO$_4$, 1.5 CaCl$_2$, 1.5MgCl$_2$, and 11 glucose, maintained at 30.0°C ± 0.5°C.
All solutions were equilibrated and maintained with carbogen gas (95% O$_2$/5% CO$_2$) throughout.

Electrophysiology

L2/3 PCs were targeted for somatic whole-cell recording in the layer 2/3 region of primary visual
cortex by gradient-contrast video microscopy. Electrophysiological recordings were obtained using
Multiclamp 700B amplifiers (Molecular Devices). Signals were filtered at 6–10 kHz and sampled at
50 kHz with the Digidata 1440B digitizer (Molecular Devices). For whole-cell recordings, borosilicate
patch pipettes were filled with an intracellular solution containing (in mM) 124 potassium gluconate,
2 KCl, 9 HEPES, 4 MgCl$_2$, 4 NaATP, 3 l-ascorbic acid, and 0.5 NaGTP. Pipette capacitance was

neutralized in all recordings and electrode series resistance was compensated using bridge balance in

current clamp. Liquid junction potentials were uncorrected.

For cell attached recordings seal resistance of >1GΩ was established, for accurate measurement of

action potentials. A stimulating electrode was placed in either layer 4 or layer 1 region of the visual

cortex, at least 200 μm away from the recorded soma. Stimulation intensities were calibrated to reliably

elicit at least 3, but no more than 10 action potentials during the 20 consequent stimuli (50 Hz, 30 s

inter-trace intervals, 10-50 μs stimulus length). Stability of the elicited responses was confirmed by an

initial 15 repeated measurements. Recordings were post hoc filtered at 2 kHz to remove stimulation

artifacts.

Recordings had a series resistance <20 MΩ. Membrane potentials were maintained near –80 mV during

current-clamp recordings using constant current bias. AP trains were initiated by somatic current

injection (300 ms) normalized to the cellular capacitance in each recording measured immediately in

voltage clamp after breakthrough (69). Cellular electrical and firing parameters were established by

recording voltage responses to current steps between -6 pA/pF and 20 pA/pF. Input resistance was

measured by averaging voltage responses to a 20 pA prepulse current step. Sag ratio was determined

as the ratio between the sag amplitude and the steady-state response during increasing -2 pA/pF current

steps.

To establish distance dependent EPSP kinetics, cells were filled with 10 μM Alexa-594. Dendrites

were visualized after cell fill with Alexa 594 (10μM) with two-photon laser scanning microscopy using

a commercial scan head (Ultima; Bruker Corp) fitted with galvanometer mirrors using a 60x, 1.0 NA

objective. The fluorophore was allowed to equilibrate for at least 10 minutes before imaging. The

stimulating electrode, filled with extracellular recording solution, was placed in close proximity (<5

μm) of the two-photon visualized dendrite. Stimulation intensity was calibrated to elicit an EPSP with

<1 ms delay and approximately 50% failure rate (54.21 ± 8.72%, n= 41). The 50% failure rate was

necessary in order to maximize the possibility of stimulating a single input. For layer 1, layer 4, and

putative LGN input stimulations the procedure was the same. Synaptic summation was probed by

eliciting 5 consequent responses at 50 Hz. Stimulation intensity was elevated during these recordings

to elicit synaptic responses every time for the first stimuli as failures introduced by lower stimulation

intensities can yield inconsistent input activation during the 5 stimuli.

For voltage-clamp recordings, cells were patched in normal extracellular solution to first record their
firing patterns. Voltage protocols for I_h current measurements consisted of 300 or 500 ms long voltage
steps between −150 and −60 mV. Activation time constant was measured by fitting the initial 100 ms
of the current response with a single exponential function. In a subset of recordings CsMeSO4 (2 mM)
was bath applied after recording control current responses, to establish cesium-sensitive I_h current
kinetics and voltage dependence by subtraction. Series resistances were between 6–20 MΩ (75–80%
compensated with 53 µs lag) and were constantly monitored. Data were discarded if the series
resistance changed more than 25%.

The dynamic clamp system was built in-house based on a previous publication(70), related online
available materials available here: (http://www.dynamicclamp.com/). The equations governing the
implemented gHCN were identical to those used in the NEURON model construction. Synaptic
conductances were built-in predefined conductances available from http://www.dynamicclamp.com/.

Intracellular solution perfusion

Intracellular solution was perfused by a custom-built tubing and relay system attached to the pipette
holder. Briefly, the pipette was filled with normal intracellular solutions to measure firing patterns and
subsequent control measurements of spontaneous and elicited EPSPs. Next, a second intracellular
solution was introduced to the pipette through a small tube drilled through the pipette holder, by
applying miniscule positive pressure. The intracellular fluid levels were marked before the recording
and through a second tube the intracellular solution was siphoned out, to not alter established pipette
capacitance. During initial trial experiments using fluorophores in the second intracellular solution it
was determined that two minutes is enough for complete exchange of the intracellular solution and
intracellular equilibration as well. Data were discarded if the series resistance or pipette capacitance
changed more than 20%.

Immunohistochemistry

Two-month-old mice were anesthetized with Nembutal (50mg/mL, i.p.) and perfused transcardially
with 4% paraformaldehyde (PFA), 0.05% glutaraldehyde in phosphate buffered saline (PBS). After
perfusion, brains were removed from the skull and immersed in 4% PFA overnight at 4°C. Coronal
100 µm-thick sections were then cut on a Vibratome (Leica V1000). Brain slices were permeabilized
and incubated in blocking buffer for one hour at room temperature. After blocking, sections were

overlaid with primary antibody to HCN1 (1:200, Thermo Fisher Scientific) overnight at 4 °C, then

subjected to a 1 h incubation with secondary antibody (Goat anti-Rabbit IgG, 1:200, Thermo Fisher

Scientific) at room temperature. Stained sections were mounted with DAPI-containing mounting

solution and sealed with glass coverslips. Confocal imaging was carried out on a Leica SP5 MP.

NEURON modeling

Computer simulations were performed using the NEURON simulation environment (versions 7.5 and

7.6, downloaded from http://neuron.yale.edu). A constrained NEURON model with an active somatic

compartment was downloaded from the publicly available Cell Type Database of the Allen Institute

(http://celltypes.brain-map.org/experiment/electrophysiology/487661754). Passive properties (axial

resistance, specific membrane capacitance and leak conductance density) were altered to recreate

passive membrane potential responses of whole-cell recorded L2/3 PCs for given stimulus intensities.

Both transient and persistent potassium channels were taken out of the model to gain simulation speed

as the aim of the simulations was to reproduce subthreshold EPSP dynamics. Final simulations were

rerun with intact sodium currents as well to confirm the lack of alterations introduced by these

channels. The I_h conductance was based on the built-in Hodgkin–Huxley model of NEURON with

freely adjustable sets of parameters(71). I_h model parameters were constrained by our whole-cell

voltage clamp recordings. The steady-state activation was governed by the following equation:

457
$$minf = \frac{1}{1 + e^{\frac{v+21.945}{6.85}}}$$

458 where v is the local membrane potential. Due to the lack of apparent inactivation, the model operated

459 with a single gate (activation).

460 The activation and deactivation time constant was defined as

461
$$mtau = 25.46 * e^{-0.5*\left(\frac{v+100.28}{16.2}\right)^2}$$

To simulate EPSP kinetics, a movable spine (length: 0.6 µm, diameter: 0.6 µm, axial resistance: 150

Ωcm), with spine neck (length: 2 µm, diameter: 0.1 µm, axial resistance: 100 Ωcm) was created. Leak

conductance was inserted into both compartments, along with the fitted conductance model. Synaptic

inputs were supplemented by using NEURON's built-in AlphaSynapse class. To probe the contribution

of specific dendritic conductances to shaping the distance-EPSP halfwidth relationship, selected
conductance models were inserted into both apical and basal dendrites with a Gaussian distribution.
This curve was selected as modulation of the Gaussian curve's width and y-offset can reproduce a wide
variety of feasible distributions such as linear, exponential, or hot-zone-like expression. To model the
contribution of dendritic sodium, potassium, and calcium channels to input modulation, NaTs, Kv3_1
and Ca_LVA models were selected. Distribution approximation was done by grid search to avoid local
fitting minimums. NMDA receptors were implemented based on a previous publication (72). During
the fitting procedure, NMDA activation voltage and AMPA:NMDA ratios were adjusted.

Statistics
Averages of multiple measurements are presented as mean ± SD. Data were statistically analyzed by
ANOVA test using Origin software and custom-written Python scripts. Normality of the data was
analyzed with Shapira-Wilks test.
478
479 **Data and software availability**
480 All code used for simulating single and multicompartmental NEURON models will be made available
481 on Github and public repositories.

Fig 1. HCN expression and effect of Cs$^+$ on AP firing in cortical L2/3 PCs

a. HCN gene subunits show uniform expression pattern among cortical PC types. Data is collected from the online available dataset(23). **b.** HCN1 immunohistochemistry reveals dense somatic expression in layer 2/3 of the visual cortex **c.** HCN channels control the excitability of L2/3 PCs. Schematic illustration of the experimental arrangement of cell attached recordings during extracellular stimulation of L4, and the shape of the elicited action potential (left). Increased spiking of L2/3 PC upon blockade of HCN channels with 2 mM CsMeSO4 (right). **d.** Pharmacological block of I$_h$ significantly increases the firing response of L2/3 PCs (0.22 ± 0.06 Hz vs. 0.48 ± 0.09 Hz for control vs CsMeSO4 bath application, p=0.002, t(9)=-4.42, Student's paired t-test, n=10. **e.** Presynaptic contributions were examined using the paired-pulse ratio (PPR) in whole cell current clamp recordings (1.09 ± 0.12 Hz vs. 1.17 ± 0.15 Hz for control vs CsMeSO4 bath application, p=0.422, t(6)=-0.8612, Student's paired t-test, n=7).

Fig 2. Functionally relevant I_h in L2/3 PCs

a. Whole cell current clamp recordings showing sag response to very negative current commands. **b.** Steady-state voltage response rectification upon hyperpolarizing current injections is mediated by I_h. **c.** Sag voltage is abolished by blocking I_h. **d.** Sag amplitude is Cs^+ sensitive (0.7 ± 0.49 mV vs. 0.05 ± 0.1 mV for control vs CsMeSO4 application, p=2.46^{-4}, t(22)=4.37, Student's two-sample t-test, n=13 and 11). **e.** Resting membrane potential is modulated by I_h (-76.75 $\pm$ 6.37 mV vs. -80.29 $\pm$ 6.48 mV for control vs CsMeSO4 bath application, p=1.98^{-4}, t(9)=6.02, Student's paired t-test, n=10). **f.** Input resistance is altered by I_h (74.78 $\pm$ 42.34 mV vs. 104.99 $\pm$ 52.91 mV for control vs CsMeSO4 bath application, p=0.048, t(6)=-2.48, Student's paired t-test, n=7). **g.** The time course of voltage responses to hyperpolarizing current on the magnitude of the current injection is abolished by Cs^+. **h.** Example voltage clamp recording of cesium sensitive currents in a L2/3 PC. **i.** Voltage dependence of cesium sensitive current activation, **j.** Voltage dependent activation kinetics of cesium sensitive currents.

Fig 3. Noncanonical effect of I_h block on EPSPs in L2/3 PCs

a. Experimental arrangement of extracellular targeted stimulation of a L2/3 PC dendrite. **b.** Application of ZD-7288 revealed a proximal bias for EPSP modulation by I_h (Exponential fit τ=104.7 μm, R^2=0.33, n=18). **c.** EPSP halfwidth is significantly modulated along the dendritic axis by I_h, (proximal dendritic locations: 17.42 ± 1.68 ms vs. 40.95 ± 24.78 ms for control vs 50μm ZD-7288 bath application, p=0.03, t(6)=2.84, Student's paired t-test, n=7, distal dendritic locations: 22.68 ± 1.78 ms vs. 28.65 ± 2.17 ms for control vs 50μm ZD-7288 bath application, p=2.1^{-4}, t(8)=-6.39, Student's paired t-test, n=9**). d.** Schematic illustration of the experimental arrangement. L2/3 PCs were recorded in whole-cell current clamp mode, and a stimulating electrode was placed either in L1 or L4. **e.** I_h modulates EPSP halfwidth for L4 stimulation, but not for L1 (13.87 ± 1.37 ms vs. 15.8 ± 1.67 ms for control vs CsMeSO4 bath application in L1, p=0.38, t(18)=-0.89, Student's two-sample t-test, n=10 each, 10.65 ± 0.64 ms vs.

20.17 ± 2.88 ms for control vs CsMeSO4 bath application in L4, p=0.005, t(18)=-3.22, Student's two-sample t-test, n=10 each, 11.01 ± 0.85 ms vs. 17.05 ± 2.41 ms for control vs CsMeSO4 bath application for putative LGN events, p=0.03, t(18)=-2.36, Student's two-sample t-test, n=10 each). **f.** Pathway-specific I_h effect on unitary EPSP voltage integral (9.35 ± 2.54 mV*ms vs. 12.9 ± 2.04 mV*ms for control vs CsMeSO4 bath application in L1, p=0.3, t(19)=-1.07, Student's two-sample t-test, n=11 and n=10 respectively, 7.85 ± 1.24 mV*ms vs. 17.39 ± 2.5 mV*ms for control vs CsMeSO4 bath application in L4, p=0.002, t(20)=-3.6, Student's two-sample t-test, n=12 and n=10 respectively, 28.01 ± 3.45 mV*ms vs. 44.89 ± 6.42 mV*ms for control vs CsMeSO4 bath application for putative LGN events, p=0.03, t(18)=-2.31, Student's two-sample t-test, n=10 each). **g.** Representative recordings of 50 Hz repeated extracellular stimuli (L1 – grey, L4 – blue, putative LGN -green). **h.** Temporal summation of L4 stimulation in control conditions (black) and during CsMeSO4 application (red). **i.** Pathway-specific I_h modulation of synaptic summation (81.05 ± 14.5 mV*ms vs. 71.07 ± 11.19 mV*ms for control vs CsMeSO4 bath application in L1, p=0.6, t(18)=0.54, Student's two-sample t-test, n=10 each, 60.06 ± 9.03 mV*ms vs. 95.15 ± 11.29 mV*ms for control vs CsMeSO4 bath application in L4, p=0.03, t(18)=-2.42, Student's two-sample t-test, n=10 each, 182.08 ± 23.36 mV*ms vs. 182.05 ± 28.59 mV*ms for control vs CsMeSO4 bath application for putative LGN events, p=0.99, t(18)=8.74*10^{-4}, Student's two-sample t-test, n=10 each).

Fig 4. Pathway-specific NMDA receptor boosting of synaptic information

a. Schematic illustration of intracellular pipette solution perfusion during whole-cell patch clamp recordings. Briefly, the pipette was filled with control recording solution (black trace), which was exchanged to an intracellular solution containing 50 μM MK-801 (purple trace) during the recording. **b.** Mean peak amplitude of sEPSPs measured at two different membrane potentials and two different recording conditions (ctrl -80 mV vs. ctrl prediction: p=4.72*10^{-6}, t(10) = 8.86, ctrl -80 mV vs ctrl -55 mV: p=0.02, t(10) = -2.81, ctrl prediction vs ctrl -55 mV: p=1.63*10^{-5}, t(10) = -7.71, MK-801 -80 mV vs. MK-801 prediction: p=1.15*10^{-6}, t(10) = 10.36, MK-801 -80 mV vs MK-801 -55 mV: p=0.58, t(10) = 0.57, MK-801 prediction vs MK-801 -55 mV: p=2.69*10^{-3}, t(10) = -3.95, Student's paired-sample t-test, n = 11 for each). **c.** Synaptic amplification estimated by dividing the predicted peak

amplitude from the hyperpolarized recordings and the measured peak at depolarized membrane potentials (1.69 ± 0.09 and 1.43 ± 0.1 for ctrl and MK-801 amplification respectively, p=0.01, t(10) = 3.04, Student's paired-sample t-test, n=11). **d.** Schematic illustration of the multiparameter NEURON fitting procedure for dendritic NMDA. To cover feasible conductance distributions, the localization of channels along the somato-dendritic axis was set by a Gaussian curve, which was shifted in space, in peak and in width as well (top). Simulations were carried out using fully reconstructed L2/3 PC dendrite from the Allen Institute open-source database (bottom). **e.** Inserting NMDA receptors into proximally located simulated synapses recreates experimentally determined distance-halfwidth relationship. **f.** Best fit of the distance-dependent EPSP halfwidth changes (without HCN). Red asterisk denotes the best fit parameter-space on panel **e. g.** Amplitude-normalized EPSPs recorded from L2/3 PCs by minimal stimulation in either in L1 (dark color tones) or L4 (light color tones) in different recording conditions (ctrl: *black*, extracellular cesium: *green*, extracellular cesium + intracellular MK-801: *red*, intracellular MK-801: *purple*). **h.** The broadening effect of cesium is reversed by NMDA receptor block (14.25 ± 1.17 ms and 10.11 ± 0.56 ms for L1 vs L4 stimuli in control conditions, p=6.73*10^{-3}, t(9)=3.5, Student's paired-sample t-test, n=10 each, 16.22 ± 1.91 ms and 23.61 ± 4.04 ms for L1 vs L4 stimuli during Cs bath application, p=0.03, t(9)=-2.67, Student's paired-sample t-test, n=10 each, 19.92 ± 2.65 ms and 11.8 ± 1.55 ms for L1 vs L4 stimuli during Cs bath application and MK-801 intracellular application, p=0.02, t(7)=3.01, Student's paired-sample t-test, n=8 each, 13.62 ± 1.43 ms and 8.97 ± 0.72 ms for L1 vs L4 stimuli during MK-801 intracellular application, p=0.025, t(7)=2.84, Student's paired-sample t-test, n=8 each).

Fig 5. Proximal bias in HCN channel expression dampens NMDA receptor dependent synaptic boosting in a NEURON model simulation.

a. Schematic illustration of the multiparameter NEURON fitting procedure for dendritic HCN, analogous to Fig 4. Simulations included the NMDA dendritic parameters established in Fig. 4. **b.** Proximal abundance of HCN channels recapitulates experimental findings in control conditions. Best fit in the parameter space (denoted with red asterisk) not only recapitulated EPSP halfwidth change, but also our observations regarding changes in input resistance and resting membrane potential. **c.** Schematic illustration of the proposed proximal HCN channel abundance in L2/3 PCs is noted.

Fig 6. Developmental regulation of I$_h$ expression

a. Representative firing patterns recorded from one week old (left, blue), two weeks old (middle, grey) and six weeks old L2/3 PCs. **b.** Developmental regulation of sag ratio (0.0 ± 0.0, 0.12 ± 0.01 and 0.07 ± 0.01 for one- (n=8), two- (n=5) and six week old L2/3 PCs (n=6), respectively, p=0 for one week old vs two weeks old cells, p=0.01, t(9)=3.04 for two vs six weeks old cells and p=0 for one vs six weeks old cells, Student's two-sample t-test), input resistance (450.6 ± 57.91 MΩ, 175.66 ± 13.66 MΩ and 98.34 ± 18.86 MΩ for one- (n=8), two- (n=5) and six week old L2/3 PCs (n=7), respectively, p=0.004, t(11)=3.65 for one week old vs two weeks old cells, p=0.01, t(10)=3.06 for two vs six weeks old cells and p=1.11*10^{-4}, t(13)=5.54 for one vs six weeks old cells, Student's two-sample t-test) and resting membrane potential (-56.05 ± 2.33 mV, -73.13 ± 2.55 mV and -79.47 ± 1.33 MΩ for one- (n=8), two- (n=5) and six week old L2/3 PCs (n=7), respectively, p=5.9*10^{-4}, t(11)=4.76 for one week old vs two weeks old cells, p=0.04, t(10)=2.39 for two vs six weeks old cells and p=1.36*10^{-6}, t(13)=8.37 for one vs six weeks old cells, Student's two-sample t-test). **c.** Representative voltage clamp recordings of

hyperpolarization activated currents in one week old (left, blue), two weeks old (middle, gray) and six weeks old L2/3 PCs (right, black). **d.** Voltage dependence of hyperpolarization activated currents in one week old (blue), two weeks old (gray) and six weeks old L2/3 PCs (black). **e.** Negative correlation between measured hyperpolarization activated current and resting membrane potential during development (Linear fit (red) slope: -2.84 mV/nS, intercept: -57.67 mV, R^2=0.71, n=20). **f.** Two-photon microscopy image depicting a proximal and a distal extracellular stimulating location positioned closely to an Alexa-594 filled L2/3 PC dendrite (left), and the resulting EPSPs (proximal – black, distal – green). **g.** EPSP halfwidth decreases with dendritic distance (Linear fit (red) slope: -0.42 ms/μm, intercept: 161.01 ms, R^2=0.4, n=35). **h.** EPSPs were significantly slower in proximal dendritic locations (147.07 ± 12.87 ms and 85.57 ± 6.83 ms for proximal and distal events, respectively, p=$5.31*10^{-5}$, t(33)=4.64, Student's two-sample t-test, n=13 and 22, respectively).

Fig 7. Neuromodulation shifts L2/3 PC excitability via HCN channels

a. 5HT$_7$ receptor mRNA is abundantly expressed in L2/3 PCs. Data is collected from the online available dataset[23]. **b.** Example cell attached recording of a L2/3 PC (left) upon 50 Hz extracellular stimulation in L4 in control conditions (black) and 5-CT bath application (orange). 5-CT significantly reduced L2/3 PC spiking (0.2 ± 0.04 spike/stimuli vs. 0.1 ± 0.04 spike/stimuli for control vs 5-CT application, p=3.57*10^{-6}, t(11)=7.92, Student's paired-sample t-test, n=12 each) **c.** Example current clamp recording of voltage responses to hyperpolarizing current injections (left; control – black, 5-CT – orange). Steady-state voltage response rectification is amplified by 5HT$_7$R blockade (right). **d.** 5-CT shifts the voltage dependence of hyperpolarization activated currents (-109.06 ± 0.47 mV vs -102.35

± 1.09 mV half activation for control vs 5-CT bath application, $R^2 = 0.99$ for both). **e.** 5-CT did not alter resting membrane potential (-82.41 ± 1.43 mV vs -80.14 ± 1.38 mV in control vs 5-CT bath application, respectively, p=0.89, t(5)=-1.41, Student's paired-sample t-test, n=6 each), sag ratio (0.02 ± 2.55*10^{-3} vs 0.03 ± 0.01 in control vs 5-CT bath application, respectively, p=0.82, t(5)=-1.01, Student's paired-sample t-test, n=6 each) and action potential threshold (-41.13 ± 2.6 mV, -40.08 ± 3.05 mV in control vs 5-CT bath application, respectively, p=0.64, t(4)=-0.4, Student's paired-sample t-test, n=5 each), but significantly reduced input resistance (108.21± 10.3 mΩ, 92.29 ± 9.9 MΩ in control vs 5-CT bath application, respectively, p=0.02, t(5)=2.75, Student's paired-sample t-test, n=6 each). **f.** Example recording showing no effect of 5-CT bath application when HCN channels were blocked (left: example recordings, right: 0.3 ± 0.11 spike/stimuli vs 0.34 ± 0.12 spike/stimuli in control vs 5-CT bath application, respectively, p=0.13, t(5)=-1.81, Student's paired-sample t-test, n=6 each). **g.** NMDA independent 5-CT effect on L2/3 PC excitability (0.27 ± 0.03 spike/stimuli vs 0.14 ± 0.04 spike/stimuli in control vs 5-CT bath application, respectively, p=0.01, t(7)=3.66, Student's paired-sample t-test, n=7 each).

Supplementary Figure 1. Similar Ih current and effect in different cortical areas.

a. Steady-state rectification of L2/3 PCs in the primary somatosensory cortex (S1), primary motor cortex (M1), primary auditory cortex (A1) and primary visual cortex (V1). **b.** Similar resting membrane potential (left), sag ratio (middle) and input resistance (right) of L2/3 PCs in different cortical areas. **c.** Example hyperpolarization activated current recordings from four cortical areas (left), with similar voltage dependent activation (right).

Supplementary Figure 2. Putative LGN and L4 inputs are well separated by k-means clustering

k-means clustering results plotted along the first and second principal component. Ellipses denote the confidence interval; edges are connected to the cluster center. Hollow circles represent putative LGN inputs and filled circles represent L4 inputs determined by the experimenter.

Supplementary Figure 3. The presence of dendritic sodium and potassium channels cannot explain wider proximal synaptic inputs

a. Sodium channels cannot explain the distance-EPSP halfwidth relationship observed when I_h was blocked. Color coding denotes the ratio between proximal and distal EPSP halfwidths. Notice the miniscule range of the color coding. **C.** Same as panel **b.** but for voltage dependent potassium channels.

Supplementary Figure 4. Pipette perfusion does not alter spontaneous circuit activity measurements

a. sEPSP frequency is unaltered by pipette perfusion (6.63 ± 0.3 and 6.4 ± 0.44 for ctrl -80 mV and perfused ctrl -80 mV, p=0.57, t(6) = 0.59, 6.1 ± 0.25 and 6.03 ± 0.33 for ctrl -55 mV and perfused ctrl -55 mV, p=0.83, t(6) = 0.23, n=7 for each, Student's paired-sample t-test). **b.** sEPSP peak amplitudes are stable during pipette perfusion experiments (0.23 ± 0.01 and 0.25 ± 0.01 for ctrl -80 mV and perfused ctrl -80 mV, p=0.43, t(6) = -0.83, 0.16 ± 0.01 and 0.17 ± 0.01 for ctrl prediction and perfused prediction, p=0.43, t(6) =-0.83, 0.31 ± 0.03 and 0.34 ± 0.02 for ctrl -55 mV and perfused ctrl -55 mV, p=0.24, t(6) = -1.31, n=7 each, Student's paired-sample t-test). **c.** Pipette perfusion does not alter measured synaptic amplification (1.92 ± 0.07 and 2.01 ± 0.24 for ctrl and perfused ctrl, p=0.65, t(6) = -0.48, n=7 each, Student's paired-sample t-test).

REFERENCES

1.   Quiquempoix M, Fayad SL, Boutourlinsky K, Leresche N, Lambert RC, Bessaih T. Layer 2/3 pyramidal neurons control the gain of cortical output. Cell reports. 2018;24(11):2799-807. e4.

2.   Wilson DE, Whitney DE, Scholl B, Fitzpatrick D. Orientation selectivity and the functional clustering of synaptic inputs in primary visual cortex. Nature neuroscience. 2016;19(8):1003-9.

3.   Hage TA, Bosma-Moody A, Baker CA, Kratz MB, Campagnola L, Jarsky T, et al. Synaptic connectivity to L2/3 of primary visual cortex measured by two-photon optogenetic stimulation. Elife. 2022;11:e71103.

4.   Ding Z, Fahey PG, Papadopoulos S, Wang E, Celii B, Papadopoulos C, et al. Functional connectomics reveals general wiring rule in mouse visual cortex. bioRxiv. 2023:2023.03. 13.531369.

5.   Scholl B, Wilson DE, Jaepel J, Fitzpatrick D. Functional logic of layer 2/3 inhibitory connectivity in the ferret visual cortex. Neuron. 2019;104(3):451-7. e3.

6.   Scholl B, Fitzpatrick D. Cortical synaptic architecture supports flexible sensory computations. Current opinion in neurobiology. 2020;64:41-5.

7.   Goetz L, Roth A, Häusser M. Active dendrites enable strong but sparse inputs to determine orientation selectivity. Proceedings of the National Academy of Sciences. 2021;118(30):e2017339118.

8.   Smith SL, Smith IT, Branco T, Häusser M. Dendritic spikes enhance stimulus selectivity in cortical neurons in vivo. Nature. 2013;503(7474):115-20.

9.   Ujfalussy BB, Makara JK, Lengyel M, Branco T. Global and multiplexed dendritic computations under in vivo-like conditions. Neuron. 2018;100(3):579-92. e5.

10.   Landau AT, Park P, Wong-Campos JD, Tian H, Cohen AE, Sabatini BL. Dendritic branch structure compartmentalizes voltage-dependent calcium influx in cortical layer 2/3 pyramidal cells. Elife. 2022;11:e76993.

11.   Ferrarese L, Jouhanneau J-S, Remme MW, Kremkow J, Katona G, Rózsa B, et al. Dendrite-specific amplification of weak synaptic input during network activity in vivo. Cell reports. 2018;24(13):3455-65. e5.

12.   Luo H, Hasegawa K, Liu M, Song W-J. Comparison of the upper marginal neurons of cortical layer 2 with layer 2/3 pyramidal neurons in mouse temporal cortex. Frontiers in Neuroanatomy. 2017;11:115.

13.   Stuart GJ, Spruston N. Dendritic integration: 60 years of progress. Nature neuroscience. 2015;18(12):1713-21.

14.   Takahashi N, Oertner TG, Hegemann P, Larkum ME. Active cortical dendrites modulate perception. Science. 2016;354(6319):1587-90.

15.   Suzuki M, Larkum ME. General anesthesia decouples cortical pyramidal neurons. Cell. 2020;180(4):666-76. e13.

16.   Xu N-l, Harnett MT, Williams SR, Huber D, O'Connor DH, Svoboda K, et al. Nonlinear dendritic integration of sensory and motor input during an active sensing task. Nature. 2012;492(7428):247-51.

17.   Shah MM. Cortical HCN channels: function, trafficking and plasticity. The Journal of physiology. 2014;592(13):2711-9.

18.   Lörincz A, Notomi T, Tamás G, Shigemoto R, Nusser Z. Polarized and compartment-dependent distribution of HCN1 in pyramidal cell dendrites. Nature neuroscience. 2002;5(11):1185-93.

19.   Wahl-Schott C, Biel M. HCN channels: structure, cellular regulation and physiological function. Cellular and molecular life sciences. 2009;66:470-94.

20.   Benarroch EE. HCN channels: function and clinical implications. Neurology. 2013;80(3):304-10.

21.   Nolan MF, Dudman JT, Dodson PD, Santoro B. HCN1 channels control resting and active integrative properties of stellate cells from layer II of the entorhinal cortex. Journal of Neuroscience. 2007;27(46):12440-51.

22.   Williams SR, Stuart GJ. Site independence of EPSP time course is mediated by dendritic I h in
neocortical pyramidal neurons. Journal of neurophysiology. 2000;83(5):3177-82.
23.   Gouwens NW, Sorensen SA, Baftizadeh F, Budzillo A, Lee BR, Jarsky T, et al. Integrated
morphoelectric and transcriptomic classification of cortical GABAergic cells. Cell. 2020;183(4):935-53. e19.
24.   Harnett MT, Magee JC, Williams SR. Distribution and function of HCN channels in the apical
dendritic tuft of neocortical pyramidal neurons. Journal of Neuroscience. 2015;35(3):1024-37.
25.   Magee JC. Dendritic Ih normalizes temporal summation in hippocampal CA1 neurons. Nature
neuroscience. 1999;2(6):508-14.
26.   Vetter P, Roth A, Häusser M. Propagation of action potentials in dendrites depends on dendritic
morphology. Journal of neurophysiology. 2001;85(2):926-37.
27.   Malinow R. Transmission between pairs of hippocampal slice neurons: quantal levels, oscillations,
and LTP. Science. 1991;252(5006):722-4.
28.   Yang W, Carrasquillo Y, Hooks BM, Nerbonne JM, Burkhalter A. Distinct balance of excitation and
inhibition in an interareal feedforward and feedback circuit of mouse visual cortex. Journal of
Neuroscience. 2013;33(44):17373-84.
29.   Balcioglu A, Gillani R, Doron M, Burnell K, Ku T, Erisir A, et al. Mapping thalamic innervation to
individual l2/3 pyramidal neurons and modeling their 'readout'of visual input. Nature Neuroscience.
2023;26(3):470-80.
30.   Sermet BS, Truschow P, Feyerabend M, Mayrhofer JM, Oram TB, Yizhar O, et al. Pathway-, layer-
and cell-type-specific thalamic input to mouse barrel cortex. elife. 2019;8:e52665.
31.   London M, Hausser M. Dendritic computation. Annual review of neuroscience. 2005;28(1):503-
32.
32.   Berger T, Senn W, Lüscher H-R. Hyperpolarization-activated current I h disconnects somatic and
dendritic spike initiation zones in layer V pyramidal neurons. Journal of neurophysiology.
2003;90(4):2428-37.
33.   Tsay D, Dudman JT, Siegelbaum SA. HCN1 channels constrain synaptically evoked Ca2+ spikes in
distal dendrites of CA1 pyramidal neurons. Neuron. 2007;56(6):1076-89.
34.   Billeh YN, Cai B, Gratiy SL, Dai K, Iyer R, Gouwens NW, et al. Systematic integration of structural
and functional data into multi-scale models of mouse primary visual cortex. Neuron. 2020;106(3):388-
403. e18.
35.   Palmer LM, Shai AS, Reeve JE, Anderson HL, Paulsen O, Larkum ME. NMDA spikes enhance action
potential generation during sensory input. Nature neuroscience. 2014;17(3):383-90.
36.   Branco T, Häusser M. Synaptic integration gradients in single cortical pyramidal cell dendrites.
Neuron. 2011;69(5):885-92.
37.   Bucurenciu I, Kulik A, Schwaller B, Frotscher M, Jonas P. Nanodomain coupling between Ca2+
channels and Ca2+ sensors promotes fast and efficient transmitter release at a cortical GABAergic
synapse. Neuron. 2008;57(4):536-45.
38.   Cornford JH, Mercier MS, Leite M, Magloire V, Häusser M, Kullmann DM. Dendritic NMDA
receptors in parvalbumin neurons enable strong and stable neuronal assemblies. Elife. 2019;8:e49872.
39.   Dodt HU, Frick A, Kampe K, Zieglgänsberger W. NMDA and AMPA receptors on neocortical
neurons are differentially distributed. European journal of Neuroscience. 1998;10(11):3351-7.
40.   Guet-McCreight A, Chameh HM, Mahallati S, Wishart M, Tripathy SJ, Valiante TA, et al. Age-
dependent increased sag amplitude in human pyramidal neurons dampens baseline cortical activity.
Cerebral Cortex. 2023;33(8):4360-73.
41.   Santello M, Nevian T. Dysfunction of cortical dendritic integration in neuropathic pain reversed
by serotoninergic neuromodulation. Neuron. 2015;86(1):233-46.
42.   Dori I, Dinopoulos A, Blue M, Parnavelas J. Regional differences in the ontogeny of the
serotonergic projection to the cerebral cortex. Experimental neurology. 1996;138(1):1-14.

43.   Vasefi MS, Yang K, Li J, Kruk JS, Heikkila JJ, Jackson MF, et al. Acute 5-HT7 receptor activation
increases NMDA-evoked currents and differentially alters NMDA receptor subunit phosphorylation and
trafficking in hippocampal neurons. Molecular brain. 2013;6:1-9.
44.   Megıas M, Emri Z, Freund T, Gulyas A. Total number and distribution of inhibitory and excitatory
synapses on hippocampal CA1 pyramidal cells. Neuroscience. 2001;102(3):527-40.
45.   Nevian T, Larkum ME, Polsky A, Schiller J. Properties of basal dendrites of layer 5 pyramidal
neurons: a direct patch-clamp recording study. Nature neuroscience. 2007;10(2):206-14.
46.   Hubel DH, Wiesel TN. Brain mechanisms of vision. Scientific American. 1979;241(3):150-63.
47.   Hubel DH, Wiesel TN. Brain and visual perception: the story of a 25-year collaboration: Oxford
University Press; 2004.
48.   Scholl B, Thomas CI, Ryan MA, Kamasawa N, Fitzpatrick D. Cortical response selectivity derives
from strength in numbers of synapses. Nature. 2021;590(7844):111-4.
49.   Waters J, Larkum M, Sakmann B, Helmchen F. Supralinear Ca2+ influx into dendritic tufts of layer
2/3 neocortical pyramidal neurons in vitro and in vivo. Journal of Neuroscience. 2003;23(24):8558-67.
50.   Major G, Polsky A, Denk W, Schiller J, Tank DW. Spatiotemporally graded NMDA spike/plateau
potentials in basal dendrites of neocortical pyramidal neurons. Journal of neurophysiology.
2008;99(5):2584-601.
51.   Harris JA, Mihalas S, Hirokawa KE, Whitesell JD, Choi H, Bernard A, et al. Hierarchical organization
of cortical and thalamic connectivity. Nature. 2019;575(7781):195-202.
52.   Rowan MJ, Christie JM. Rapid state-dependent alteration in Kv3 channel availability drives flexible
synaptic signaling dependent on somatic subthreshold depolarization. Cell reports. 2017;18(8):2018-29.
53.   Levelt CN, Hübener M. Critical-period plasticity in the visual cortex. Annual review of
neuroscience. 2012;35:309-30.
54.   Bard JA, Zgombick J, Adham N, Vaysse P, Branchek TA, Weinshank RL. Cloning of a novel human
serotonin receptor (5-HT7) positively linked to adenylate cyclase. Journal of Biological Chemistry.
1993;268(31):23422-6.
55.   Karmarkar UR, Dan Y. Experience-dependent plasticity in adult visual cortex. Neuron.
2006;52(4):577-85.
56.   Hensch TK, Bilimoria PM, editors. Re-opening windows: manipulating critical periods for brain
development. Cerebrum: the Dana forum on brain science; 2012: Dana Foundation.
57.   Nolan MF, Malleret G, Dudman JT, Buhl DL, Santoro B, Gibbs E, et al. A behavioral role for dendritic
integration: HCN1 channels constrain spatial memory and plasticity at inputs to distal dendrites of CA1
pyramidal neurons. Cell. 2004;119(5):719-32.
58.   Wu J, El-Hassar L, Datta D, Thomas M, Zhang Y, Jenkins DP, et al. Interaction Between HCN and
Slack Channels Regulates mPFC Pyramidal Cell Excitability and Working Memory. bioRxiv. 2023:2023.03.
04.529157.
59.   Yu L, Tasaki S, Schneider JA, Arfanakis K, Duong DM, Wingo AP, et al. Cortical proteins associated
with cognitive resilience in community-dwelling older persons. JAMA psychiatry. 2020;77(11):1172-80.
60.   Ramírez D, Zúñiga R, Concha G, Zúñiga L. HCN channels: new therapeutic targets for pain
treatment. Molecules. 2018;23(9):2094.
61.   Ku SM, Han M-H. HCN channel targets for novel antidepressant treatment. Neurotherapeutics.
2017;14(3):698-715.
62.   Kim CS, Johnston D. A possible link between HCN channels and depression. Chronic Stress.
2018;2:2470547018787781.
63.   Postea O, Biel M. Exploring HCN channels as novel drug targets. Nature reviews Drug discovery.
2011;10(12):903-14.

64.   Lyman KA, Han Y, Chetkovich DM. Animal models suggest the TRIP8b-HCN interaction is a
therapeutic target for major depressive disorder. Expert Opinion on Therapeutic Targets. 2017;21(3):235-
7.
65.   Fukuyama K, Motomura E, Okada M. Therapeutic potential and limitation of serotonin type 7
receptor modulation. International Journal of Molecular Sciences. 2023;24(3):2070.
66.   Ishibashi T, Horisawa T, Tokuda K, Ishiyama T, Ogasa M, Tagashira R, et al. Pharmacological profile
of lurasidone, a novel antipsychotic agent with potent 5-hydroxytryptamine 7 (5-HT7) and 5-HT1A
receptor activity. Journal of Pharmacology and Experimental Therapeutics. 2010;334(1):171-81.
67.   Shimizu M, Mi X, Toyoda F, Kojima A, Ding W-G, Fukushima Y, et al. Propofol, an Anesthetic Agent,
Inhibits HCN Channels through the Allosteric Modulation of the cAMP-Dependent Gating Mechanism.
Biomolecules. 2022;12(4):570.
68.   Chen X, Shu S, Kennedy DP, Willcox SC, Bayliss DA. Subunit-specific effects of isoflurane on
neuronal I h in HCN1 knockout mice. Journal of neurophysiology. 2009;101(1):129-40.
69.   Taylor AL. What we talk about when we talk about capacitance measured with the voltage-clamp
step method. Journal of computational neuroscience. 2012;32(1):167-75.
70.   Desai NS, Gray R, Johnston D. A dynamic clamp on every rig. eneuro. 2017;4(5).
71.   Oláh VJ, Tarcsay G, Brunner J. Small size of recorded neuronal structures confines the accuracy in
direct axonal voltage measurements. Eneuro. 2021;8(4).
72.   Testa-Silva G, Rosier M, Honnuraiah S, Guzulaitis R, Megias AM, French C, et al. High synaptic
threshold for dendritic nmda spike generation in human layer 2/3 pyramidal neurons. Cell reports.
2022;41(11).