Host-pathogen-vector continuum in a changing landscape: Drivers of *Bartonella* prevalence and evidence of historic spillover in a multi-host community

B.R. Ansil[1,2]*, Ashwin Viswanathan[3], Vivek Ramachandran[1], H.M. Yeshwanth[1], Avirup Sanyal[1,4], Uma Ramakrishnan[1,*]

[1] National Centre for Biological Sciences, Tata Institute of Fundamental Research, Bangalore, Karnataka, India, 560065

[2] Manipal Academy of Higher Education, Manipal, Karnataka, India, 576104

[3] Nature Conservation Foundation, Mysore, Karnataka, India, 570017

[4]Trivedi School of Biosciences, Ashoka University, Sonipat, Haryana, 131029

* Corresponding author: ansilbr@ncbs.res.in, uramakri@ncbs.res.in

Keywords: Spillover, Western Ghats, Small mammals, Vector-borne pathogens

Abstract

Small mammals and their ectoparasites present a unique system to investigate the eco-epidemiology of multi-host vector-borne pathogens and identify specific bacterial spillover determinants. We applied ecological and evolutionary analyses in a rainforest-human-use mosaic to investigate *Bartonella spp.* across small mammal and ectoparasite communities. We observed substantial overlap among small mammal communities in different habitat types, predominantly driven by habitat generalists. Most ectoparasites were generalists, infecting multiple hosts. We observed high *Bartonella* prevalence at both study sites –a forest-plantation mosaic (47.4%) and a protected area (28.8%). Seven of the ten ectoparasite species sampled were also positive for *Bartonella*, following prevalence trends in their hosts. A generalised linear model revealed an independent association between aggregated ectoparasite load in hosts and *Bartonella* prevalence, implicating ectoparasites in transmission. *Bartonella* lineages from small mammals were host-specific, while ectoparasites carried *Bartonella* associated with other small mammal hosts, indicating the potential for cross-species transmission. Phylogenetic ancestral trait reconstruction of *Bartonella* haplotypes suggest historic spillover events in the small mammal community, validating the potential for contemporary spillover events. These results highlight the necessity to disentangle the complex relationship between hosts, ectoparasites, and pathogens to understand the zoonotic implications of undetected spillover events in such multi-host communities.

1. Background

Large-scale modification of natural habitats is associated with loss of biodiversity and ecosystem services [1]. Other unexpected consequences, such as an elevated risk of zoonotic infections, are poorly acknowledged and understood, especially in biodiverse tropical systems. Both theoretical and empirical studies demonstrate that land-use change and associated habitat perturbations can result in modified communities of zoonotic hosts driven by movement of animals across habitat boundaries [2,3].Consequently, there is potential for pathogen spillover between animals that did not previously co-occur, i.e. from reservoir hosts to incidental hosts, directly or indirectly through intermediate hosts and vectors [4,5].

Spillover between wildlife can lead to the emergence of zoonoses [4], and has been implicated in several past epidemics, including SARS-CoV-1 [6]. Pathogen sharing between animal host communities can lead to the recombination and evolution of new variants with altered host ranges, vector specificities, and pathogenic potential [7]. Spillover events, however, are complex and infrequent, influenced by numerous ecological and evolutionary filters that vary in space and time, including the distribution of reservoir hosts, pathogen prevalence, and infection intensity [8]. For example, a high density of reservoir hosts elevates direct interactions, leading to density-dependent transmission among the reservoir population and to other susceptible members of the community [9]. Additionally, in the case of vector-borne pathogens, ectoparasites also respond to host densities and influence pathogen prevalence in reservoirs and incidental hosts [10].

Bats and small mammals (rodents and shrews) are the best-known reservoir taxa for

zoonotic pathogens [11]. Small mammals, however, harbour a high number of pathogens

[12,13] owing to their disproportionately high diversity within mammalian communities

[14], even more so than bats. Small mammals also differ from bats in that they often

increase (rather than decrease) in abundance in altered landscapes, resulting in the

elevated prevalence and transmission risk of associated zoonoses [15,16]. Small mammal

densities are known to increase locally, for example, when large herbivore densities

decrease [17], leading to an increase in pathogen prevalence [18]. In addition, small

mammals with generalist natural histories (often reservoirs) tend to move across habitat

boundaries where they interact with incidental hosts [4,19], thereby moving pathogens

across these boundaries. We expect such processes to be especially prevalent in tropical

regions, as they harbour some of the richest small mammal communities [20], the highest

rates of habitat modification [21,22], and therefore the highest predicted risk of zoonotic

infections [23,24].

To understand the zoonotic consequence of small mammals and their ectoparasites in

changing landscapes, we must first understand how their communities are structured and

how they have responded (and continue to respond) to land-use change and habitat

modification. In this study, we characterize differences in host community structure,

ectoparasite community structure, and pathogen prevalence in a human-dominated

rainforest-plantation landscape in South India, part of the Western Ghats biodiversity

hotspot. We used *Bartonella*, a generalist zoonotic bacterial pathogen, as a model system to

understand variation in prevalence and transmission dynamics in the landscape. We

sampled small mammal communities and their ectoparasites in this landscape to address

the following questions – i) How are small mammal communities structured in a tropical

rainforest-plantation landscape? ii) How does *Bartonella* prevalence vary across small

mammal species in this human-use landscape? iii) What are the main drivers of *Bartonella*

prevalence and transmission in the landscape? (iv) and finally, do we find evidence for

spillover, detected as shared *Bartonella* haplotypes between members of these

communities? We began this study in a forest-plantation mosaic and expanded to a

relatively less disturbed (currently) protected area. However, the protected area had

significant human-use in the past [25], similar to the forest-plantation mosaic, allowing us to

derive insights that can be broadly applied and generalized. Given the mixed-use nature of

the landscape, we expected similarities between small mammals and ectoparasite

communities across land-use types [26]. We hypothesized that *Bartonella* prevalence would

be high and significantly influenced by the densities of known host species (e.g., *Rattus*

satarae) [9,27]. We also expected that *Bartonella* haplotypes would be specific to host

species [28] with limited spillover (vector-mediated) among hosts that occupy the same

habitats [29].

2. Methods

(a) Field sampling

We carried out this study in two sites in the central Western Ghats: i) Kadamane, a mixed-

use, forest-plantation mosaic in the Karnataka state in India, and ii) Kudremukh National

Park, a protected area in the same state **(Figure 1)**. We sampled small mammals (rodents

and shrews) from different land-use types in Kadamane and Kudremukh between 2016 and

2021. Kadamane was sampled three times during this period (2016, 2017, and 2018) and

Kudremukh was sampled once in 2021. In Kadamane, we sampled in forest fragments,

grasslands, tea plantations, rested plantations (unmanaged cardamon and coffee), and

built-up (human habitation). In Kudremukh, we sampled forests, grasslands, and built-up (a

partially occupied township inside Kudremukh National Park). In both sites, sampling was

conducted during dry months between January- April.

We used medium-sized Sherman traps in a grid framework to sample small mammals. A

trapping grid consisted of 10 trapping lines, each line consisting of 10 traps separated by 10

meters, covering an area of one hectare in total in each habitat type. An additional 20 traps

were placed on trees to capture arboreal species in forests and rested plantations. Trapping

efforts were proportionately divided based on the area available under each land-use

category. Specifically, in Kadamane, this translated to three grids in forests, two grids in

grasslands, two grids in rested plantations, one grid in tea plantations, and one grid in built-

up areas. Because the rested plantations in Kadamane were often indistinguishable from

forests, we grouped them under 'forests' in all analyses. In Kudremukh, three grids were

placed in forests, two in grasslands, and one in built-up areas **(Figure 1)**. Each grid was

sampled for four consecutive days during each trapping session (assuming a closed

population). All traps were baited using peanut butter and locally available seeds. Traps

were checked twice a day (morning and evening) for captured individuals. Captured

individuals were anesthetised using Isoflurane for collecting measurements and samples.

Small mammal species were first identified in field with the help of taxonomic keys and field

guides [30–32]. In addition, ear clippings were collected to confirm species identification

through molecular methods. We collected dried blood spots from every captured individual

in 2018 and 2021 to screen for pathogens. To detect and collect ectoparasites, captured

individuals were combed on a white paper for 2-3 minutes. All individuals were marked and

released at the site of capture after sample collection (except in 2018; individuals were

sacrificed for harvesting organs to screen for various pathogens).

Figure 1: Map of the study area. A) Map of India with Karnataka state, Western Ghats and

study sites marked. (B) Land-use map of Kadamane forest-plantation mosaic with sampling

locations marked. (C) Land-use map of Kudremukh National Park with sampling locations

marked. The legends indicate different land-use types and the number of grids sampled per

land-use type. Note that rested plantations are grouped under forests.

(b) Small mammal community

DNA was extracted from ear clippings using DNeasy Blood & Tissue Kit (Qiagen), following

which 1140 bp of Cytochrome b (*Cytb*) gene was amplified and sequenced [33]. Maximum

likelihood phylogenetic trees were constructed to identify cryptic *Mus* species and validate

morphological species identification. All the other species were identified morphologically

and validated using BLAST (blast.ncbi.nlm.nih.gov) similarity scores of *Cytb* sequences.

Based on species identification, we created a species abundance matrix for each land-use

type, followed by a distance matrix using the Morisita-Horn index, a standard beta diversity

measure to calculate dissimilarity [34]. The beta diversity index was plotted following

hierarchical clustering to visualise the dissimilarity between small mammal communities in

different land-use types and sampling grids. We also calculated the relative abundance of

each small mammal species in each land-use type to understand species-habitat

associations. The community ecology package Vegan [35] in R was used for this analysis.

We used a Capture-Mark-Recapture approach [36] to estimate densities of the four most

common small mammals by creating individual capture histories in forest and grassland. We

assumed a closed population with varying capture probability to estimate population size

following a customised log-linear model [37]. Density (number of individuals per hectare)

for each species was estimated after accounting for the number of grids sampled in each

habitat. We used R package *Rcapture* for this analysis [37].

(c) Ectoparasite community

Ectoparasites were brought back to the laboratory in 100% ethanol. Mites, ticks, and fleas

were segregated using a stereo-zoom microscope. We identified ticks and fleas to the genus

level under a high-resolution microscope with the help of taxonomic keys and monographs

[38–40]. Mites were mounted on a slide for observation under the microscope and

identified through descriptions in a monograph [41]. We serially named and counted

individual ectoparasites that belonged to the same morphotypes (potentially species).

We constructed a distance matrix based on ectoparasite occurrence per individual small

mammal species [42]. We used this matrix for a Principal Coordinate Analysis (PCoA) in R

[43], and plotted the clustering using ggplot2 [44] to understand the overlap in ectoparasite

communities between small mammal species (the first two PCoA axes). We also estimated

ectoparasite density, or the mean number of each morphotype per host species in forests,

grasslands, and built-up areas. The infestation rate, i.e., frequency of ectoparasite (mites,

ticks, and fleas) occurrence in each host species and habitat, was recorded as aggregated

ectoparasite load.

(d) *Bartonella* screening and prevalence estimation

To detect *Bartonella* in small mammals, DNA was extracted using QIAamp DNA Mini Kit

(Qiagen) from the dried blood spots (DBS) collected in 2018 (Kadamane) and 2021

(Kudremukh), and screened for *Bartonella* specific *rpoB* gene (825 bp) using conventional

PCR [45]. *Bartonella* prevalence in the Kadamane samples was reported previously [27]. To

detect *Bartonella* in ectoparasites, ectoparasite were pooled, mechanically homogenised,

and DNA was extracted using DNeasy Blood & Tissue Kit (Qiagen). A pool contained 1-10

individual ectoparasites (mean = 3.06) belonging to the same species (previously called

morphotypes) collected from the same individual small mammal. We used PCR reagents as

follows- 25 µl reaction mixture composed of 6 µl of template DNA, 12.5 µl of 1X HotStarTaq

Master Mix (Qiagen), 1 µl each of 5 µM primers and 4.5 µl of nuclease-free water. PCR

products were visualized on a 1.5% agarose gel stained with GelRed Nucleic Acid Gel Stain

(Biotium Inc) with reference to negative and positive controls set along the test samples.

Positive samples were purified using AMPure XP magnetic beads (Beckman Coulter) and

sequenced. Sequences were verified for *Bartonella* after comparing with the reference

database (NCBI) using BLAST [46]. All *Bartonella* sequences generated in this study have

been deposited in GenBank under accession numbers OR574780-OR574826.

Prevalence was calculated as the proportion of positive individuals against the number of

individuals tested. While calculating the overall prevalence, we included two forest grids

from Kudremukh twice (randomly selected) to compensate for additional trapping effort in

Kadamane (rested plantations) to make the sites comparable. We calculated 95%

confidence intervals (Wilson interval) associated with the prevalence using the R package

Hmisc [47]. Prevalence (positivity) in ectoparasite pools was represented as the proportion

of positive pools among the total number of ectoparasite pools tested.

(e) Ecological correlates of *Bartonella* prevalence

We used a generalized linear model (GLM) to understand drivers of *Bartonella* prevalence

among small mammals in the landscape. We modelled prevalence as the incidence of

Bartonella (presence/absence) in each individual small mammal as a function of the

sampling site, aggregated ectoparasite load (the proportion of all individuals of the species

with any ectoparasite presence - mites, ticks, and fleas - within the relevant habitat-site

combination), and the density of the small mammal species itself. We assumed a binomial

error distribution as our response consisted of 0s and 1s. We restricted this analysis to the

10 species-habitat-site combinations that contained data from more than three individuals

of the species, together comprising 196 individuals of five species. We also examined host

species identity and habitat type as covariates but dropped them because the models did

not converge. The covariate 'aggregated ectoparasite load' is the frequency of ectoparasite

occurrence in each species and habitat, and is not ectoparasite density per individual small

mammal. We considered this a more reliable measure than raw ectoparasite densities since

it is possible to occasionally encounter individuals with heavy infestation in the community

and this might potentially influence our inference.

(f) Phylogenetic analyses

We generated a consensus of forward and reverse sequences for the partial *rpoB* sequence.

These sequences were aligned with known *Bartonella* species sequences from the NCBI

database and sequences reported in Ansil et al 2021 (accession number MT787671-

MT787733) [27]. A 773 bp alignment was created using MUSCLE in MEGA 7.0.26 [48].

Sequences generated from the ectoparasites were also included in this alignment.

Alignment was codon-optimized and tested for the appropriate nucleotide substitution

models using PartitionFinder 2.1.1 [49]. We performed a Bayesian phylogenetic analysis

using this alignment in BEAST 1.8.4 [50]. The phylogeny was run for 10^8 Markov Chain

Monte Carlo (MCMC) cycles. Effective sample sizes (ESS) and parameter convergence of

MCMC sampling were assessed using Tracer 1.6.0 [51]. The resulting trees were summarised

into a consensus tree after 25% burnin, and were visualized and annotated using FigTree

1.4.4 [52]. The phylogenetic tree was rooted using *Brucella melitensis* (AY562181) as an

outgroup. Internal nodes of the tree are marked and coloured based on the Posterior

Probability (PP).

(g) Host- *Bartonella* association

We also constructed a tanglegram (co-phylogeny) using sequence data from the hosts (Cytb)

and *Bartonella* (*rpoB*) in TREEMAP [53] to understand host- *Bartonella* haplotype

association. We visually analysed the resulting co-phylogeny for potential host switching

(spillover) events.

A separate alignment was constructed using unique *Bartonella* haplotypes and considered

for a Bayesian phylogenetic ancestral trait reconstruction. We restricted this analysis to

unique haplotypes recovered from each host species. We included one additional sequence

from *Mus cf. fernandoni* to avoid over-representation and subsequent biases from *Rattus*

satarae sequences. We used six identified host states as traits to estimate host probabilities

for all ancestral nodes in BEAST 1.8.4 [50]. The resultant phylogeny was summarised and

annotated as described before.

3. Results

(a) Small mammal community

We recorded 11 small mammal species, of which nine were rodents, and two were shrews

(Table S1). The rodent species were *Rattus satarae, Rattus rattus, Mus cf. fernandoni, Mus*

cf. famulus, Mus cf. terricolor, Golunda ellioti, Vandeleuria nilagirica, Platacanthomys

lasiurus, and *Funambulus tristriatus*. The two shrew species were *Suncus niger* and

Crocidura horsfieldii. The three *Mus* species were morphologically similar and could only be

differentiated using their phylogenetic clustering **(Figure S1)**. The average genetic distances

(p distances) for *M. cf. fernandoni, M. cf. famulus*, and *M. cf. terricolor* from their respective

parent species (for example, *M. cf. fernandoni* vs *M. fernandoni*) were 0.05, 0.07 and 0.09,

respectively.

Among the various habitats sampled, grasslands had the most diverse small mammal

communities with five species detected in both Kadamane and Kudremukh. Forest

fragments in Kadamane also had five species of small mammals, but forests in Kudremukh

only had three species. Built-up areas were less diverse with just three species captured at

each site, while tea plantations were the most species-poor land-use type, with only one

species (*M. cf. famulus*) captured once in 2016.

Based on the hierarchical clustering of beta diversity, forests (FR16, FR17, FR18, and FR21)

showed similar community assemblages by forming a single cluster with minimal variability

across years (<20%); the forest community from Kudremukh (FR21) was also a part of this

cluster **(Figure 2A left)**. The grassland and tea plantation (TP16) formed a cluster with 40%

dissimilarity with the forest cluster. The grassland communities had minimal variability

(<20%) between years (GL16, GL17, GL18) and sites (GL21). Built-up habitats were the most

dissimilar (60%) to other land-use types, although they retained two species (*S. niger* and *M.*

cf. famulus) detected in the other land-use types **(Figure 2A right)**. Overall, Overall, the

small mammal community composition within each habitat was similar across years.

We were able to estimate densities for four (most abundant of the eleven) common small

mammal species (*Rattus satarae, M. cf. fernandoni, M. cf. famulus,* and *M. cf. terricolor*).

These densities are reported in Supplementary results and **Figure S2**.

Figure 2: Community structure of small mammals and their ectoparasites. A) Left: a dendrogram showing dissimilarity between small mammal communities inhabiting different habitats sampled and sampling years (2016, 2017, 2018, 2021). Abbreviations: FR- Forest, GL-Grassland, TP-Tea plantations, and BT-Built-up areas. Right: Relative abundance of different small mammal species captured in different habitats. B) Principal Coordinate (PCoA) plot showing ectoparasite community structure across different small mammals and habitats. The ellipses represent the 95% confidence level of the respective sample group centroid.

(c) Ectoparasite community

Ten ectoparasite species were recorded from the small mammals; two species of mites belonging to the genus *Laelaps*, three species of ticks belonging to the genus *Rhipicephalus*, *Ixodes* and *Haemaphysalis* and five species of fleas belonging to the genus *Xenopsylla* (**Table S2)**. Since more than one species was detected under *Laelaps* (mites) and *Xenopsylla* (fleas), these morphotypes were named serially *(e.g. Xenopsylla sp 1, Xenopsylla sp 2 and so on).* Only one morphotype each was identified from three tick genera. Interestingly, six ectoparasite species (60%) were detected in multiple hosts (generalists), and six of eight small mammals sampled (75%) were polyparasitic (infested with two or more ectoparasite species).

Our PCoA revealed substantial overlap and variation in ectoparasite communities hosted by small mammals (**Figure 2B**). *R. satarae* showed significant variation in ectoparasite community along the PCoA axis 1 and 2 (13.67 % and 10.68 % explained, respectively), while *M. cf.* fernandoni predominantly displayed limited variation (similar ectoparasite community dominated by *Laelaps* 1). However, individuals from *R. rattus*, *F. tristriatus*, *M. cf. fernandoni*, *M. cf. terricolor*, and *S. niger* showed ectoparasite community similarity with *R. satarae*. We also estimated the mean density of each ectoparasite species recovered from each small mammal species per habitat presented in Supplementary results and **Figure S2**.

(d) *Bartonella* prevalence in small mammals

We detected *Bartonella* in five species of small mammals from both sites; *R. satarae*, *R. rattus*, *M. cf. fernandoni*, *M. cf. famulus,* and *S. niger* (**Table 1**) with varying prevalence (0-

75.8%). In both sites, *R. satarae* showed the highest prevalence of 75.8% (n= 66) in

Kadamane and 72% (n= 25) in Kudremukh, followed by *M. cf. fernandoni* that showed a low

prevalence of 54.5% (n=22) in Kadamane and 40% (n=5) in Kudremukh. *M. cf. famulus*, on

the other hand showed a low prevalence of 3.8% in Kadamane (n=26, only one individual

positive) and zero prevalence in Kudremukh (n=3), while *M. cf. terricolor* showed zero

prevalence at both sites despite good sample size (n =6 and 37 in Kadamane and

Kudremukh, respectively). Other species, such as *R. rattus* and *S. niger* had low prevalence,

but also had low sample sizes, hence we have minimal confidence in their prevalence

estimates.

We observed similar patterns of *Bartonella* prevalence at both sites (*R. satarae*, *M. cf.*

fernandoni, *M. cf. famulus*, and *M. cf. terricolor*) and overall community at both sites **(Figure**

3A). Since samples were collected in different years (2018 in Kadamane and 2021 in

Kudremukh), we did not attempt any comparison between sites, but treated them

separately. Pooled ectoparasite positivity was largely correlated with prevalence in small

mammals, albeit slightly lower in every case. Positivity in specific ectoparasite species is

provided in **Table S3.**

Table 1: *Bartonella* prevalence in small mammal blood and ectoparasites. Prevalence reported for ectoparasites is not true prevalence since ectoparasites were pooled for screening (see methods).

	Bartonella prevalence; number of positives / number of samples tested (%)			
	Dried blood spot (DBS)		Pooled ectoparasites	
	Kadamane	Kudremukh	Kadamane	Kudremukh
Rattus satarae	50/66 (75.8)	18/25 (72)	24/71 (33.8)	6/24 (25)
Rattus rattus	0/10 (0)	1/4 (25)	-	0/3 (0)
Mus cf. fernandoni	12/22 (54.5)	2/5 (40)	9/29 (31)	0/6 (0)
Mus cf. famulus	1/26 (3.8)	0/3 (0)	0/1 (0)	-
Mus cf. terricolor	0/6 (0)	0/37 (0)	-	0/7 (0)
Platacanthomys lasiurus	-	0/1 (0)	-	0/1 (0)
Funambulus tristriatus	-	0/1 (0)	0/2 (0)	0/2 (0)
Suncus niger	0/2 (0)	2/4 (50)	0/1 (0)	1/2 (50)

Figure 3: (A) *Bartonella* prevalence in the overall community and the four most abundant small mammals. (B) *Bartonella* prevalence positively correlated with aggregated ectoparasite densities within a small mammal species at each habitat site. The trend line indicates predicted *Bartonella* prevalence as a function of aggregated ectoparasite load. The shaded area represents the 95% confidence interval of the predicted prevalence estimate.

(e) Factors affecting *Bartonella* prevalence

Bartonella prevalence increased with aggregated ectoparasite load (mites, ticks, and fleas)

within a species at each habitat site (GLM, p<0.05, **Figure 3B**). We did not find evidence that

Bartonella prevalence was influenced by host densities and sampling locations (**Table 2**). A

second GLM with host species identity and habitat type as additional covariates did not

converge; hence we tested these variables separately and found them not significant.

360
361
362 **Table 2:** Results of generalised linear model (GLM) examining drivers of *Bartonella*

363 prevalence. Significant variables are marked with (*)

	Estimate	Std. Error
Intercept	-1.633595	1.05266
Host density	-0.07574	0.06711
Site	-0.47880	0.72288
Aggregated ectoparasite load	5.10933*	0.96562

(f) Phylogenetic relationship

Phylogenetic analysis (*rpoB*) revealed eight distinct *Bartonella* lineages (BL1-BL8) from small

mammals and ectoparasites. We found that five lineages among them (BL1-BL5) were

phylogenetically related to known zoonotic *Bartonella* species (**Figure 4**). We observed

three lineages in *R. satarae* (BL1, BL6, and BL7) and two lineages in *M. cf. fernandoni* (BL4

and BL8). BL1 included *Bartonella* sequences from *R. satarae* and ectoparasites (*Laelaps sp*

1, Rhipicephalus sp, Ixodes sp, Xenopsylla sp 1, and Xenopsylla sp 5). Interestingly,

sequences from *Laelaps sp 1* and *Xenopsylla sp 5* in this lineage were not collected from *R.*

satarae but *M. cf. fernandoni* and *S. niger*, respectively. BL2 consisted of a single sequence

collected from *R. rattus*, and closely related to BL1. These two lineages were

phylogenetically related to *B. queenslandensis*. BL3 consisted of two sequences, one from *S.*

niger and the other from *Laelaps sp 1* collected from *M. cf. fernandoni*. This lineage was

related to *B. tribocorum*. BL4 consisted of *Bartonella* sequences from *M. cf. fernandoni*, *M.*

cf. famulus, and ectoparasites (*Laelaps sp 1, Rhipicephalus sp, Ixodes sp, Haemaphysalis sp,*

and Xenopsylla sp 3). In this lineage, the *Haemaphysalis sp* was recovered from *R. satarae*

and not from any of the *Mus* species. BL5 was a single sequence from *F. tristriatus* (reported

in Ansil et al. 2021 [27], accession number- MT787671), forming a sister lineage to BL4. We

included this sequence in our analysis since it was collected during our study in Kadamane.

The remaining lineages (BL6-BL8) were related to non-pathogenic *Bartonella* species. BL6

comprised sequences from *R. satarae* (second lineage from *R. satarae*) and *Ixodes sp*

recovered from them. This lineage was related to *B. phoceensis*. BL7 and BL8 were two

distinct lineages from *R. satarae* (third lineage from *R. satarae*) and *M. cf. fernandoni*

(second lineage from *M. cf. fernandoni*), respectively. Both these lineages were

phylogenetically related to *B. thailandensis*.

Figure 4: Bayesian phylogenetic inference of *Bartonella* based on 773 bp of *rpoB* sequences.
Nodes are coloured based on posterior probability, and monophyletic clades are collapsed
for easy visualization. Ectoparasites part of collapsed lineages and their hosts are expanded
on the left side.

(g) Host-*Bartonella* association and historic spillover

The co-phylogeny of small mammals and *Bartonella* revealed host-specific lineages **(Figure
5A)**. However, one *Bartonella* lineage from *M. cf. fernandoni* (BL8) was observed to be
phylogenetically related to *Bartonella* sequences from *R. satarae* (BL7). Our phylogenetic
ancestral trait reconstruction revealed a high probability (0.87) of *R. satarae* being the host
state of the common ancestor of these lineages **(Figure 5B)**, suggesting a potential past
spillover event. Similarly, we also observed a moderately high probability (0.73) of *R.*

404   *satarae* being the host state for the *Bartonella* sequence from *R. rattus* and the remote

405   common ancestor (0.8) of the *Bartonella* community level phylogeny. We found no

406   evidence for contemporary *Bartonella* spillover or sharing between small mammal hosts.

Figure 5: A) Co-phylogeny of small mammals and *Bartonella* association. One lineage of *M. cf. fernandoni* (BL8) was observed to have phylogenetic similarity with one of the *R. satarae* lineage (BL7). B) Bayesian phylogenetic inference showing ancestral host state of different *Bartonella* haplotypes. Each colour represents a specific host, and the number at the common ancestor nodes indicates the probability for the corresponding host state.

4. Discussion

We characterized small mammal communities, their ectoparasites, and associations with *Bartonella* in a human-dominated rainforest-plantation landscape in the Western Ghats. We demonstrated that certain small mammal host species occur across habitats, bringing them in contact with specialist hosts with specific *Bartonella* lineages. We also demonstrate that ectoparasites can serve as potential vectors, because *Bartonella* prevalence in the community is influenced by the cumulative ectoparasite load in small mammals. Additionally, we present evidence of a historic spillover event between two major hosts in the community and substantiate the potential for future *Bartonella* sharing between hosts, including other domestic animals and humans, possibly through ectoparasites.

(a) Some small mammals have the potential to carry *Bartonella* across habitat boundaries

We observed strong species-habitat associations in the focal small mammal community, characterised by exclusive occurrences and high densities, mirroring several ecological communities [54,55]. While species like *R. satarae* (a major *Bartonella* host with highest prevalence) showed high abundance and nearly exclusive occurrence in forested habitats, *R. rattus*, a known synanthropic species, was only detected in built-up areas, reaffirming its commensal nature [56]. Similarly, *M. cf. terricolor* was exclusively recorded in grasslands in both sites, albeit with higher density in Kudremukh. Other *Mus* species , *M. cf. fernandoni*

and *M. cf. famulus* were also abundant in grasslands, indicating a generalizable association

of *Mus* species with grassland ecosystems. Significantly from a disease ecology perspective,

few species like *M. cf. famulus* and *S. niger*, however, were recorded in multiple habitats,

including forests, grasslands, and built-up areas. Species like *M. cf. famulus* can also be

incidental hosts (as may be the case with the single positive), connecting potential reservoir

hosts like *R. satarae* and *M. cf. fernandoni*, with *R. rattus* (another seemingly incidental host

with a single positive) and humans. Similarly, *M. cf. fernandoni* was found to occur at low

densities in forests in addition to grasslands, bringing them in contact with *R. satarae*,

potentially paving the way for the historic spillover (discussed later) from *R. satarae* to *M.*

cf. fernandoni.

We did not find any notable difference between small mammal communities within

Kudremukh (protected area) and Kadamane (forest-plantation mosaic). One reason for this

might be that both landscapes (and much of the Western Ghats) have a long history of

human use [25,57], dampening differences that might exist between 'natural' and human

dominated areas. However, inadequate temporal replicates and delay in sampling

Kudremukh diminishes the generality of the pattern observed. Interestingly, all *Mus* species

in Kadamane showed variation in annual densities, indicating annual population oscillations

[58,59]. We could not fully capture this pattern due to the relatively short duration of this

study. However, understanding such variation in host density is crucial as it can be a

determinant of zoonotic hazard and spillover [8,60]. This reiterates the importance of

detailed ecological studies on host communities to understand spillover dynamics.

Our study suggests that small mammals in this part of the Western Ghats are in urgent need

of taxonomic revision. Previous studies from the landscape morphologically identified *M. cf.*

fernandoni as *M. musculus* and *M. cf. famulus* and *M. cf terricolor* as *M. booduga* [27].

Phylogenetic analysis and genetic distances suggest that these are distinct lineages related

to *M. fernandoni*, *M. famulus*, and *M. terricolor*, respectively. We urge researchers to

sample rodents in this landscape and resolve their taxonomy on priority, because pathogen

and habitat associations are often linked with species identity as we show in this study.

(b) *Bartonella* prevalence and spillover may be mediated by ectoparasites

Our results have revealed some broad ectoparasite-host associations that may have

implications for *Bartonella* maintenance and transmission. The observed patterns include

instances of polyparasitism, host sharing, and the influence of ectoparasitism on *Bartonella*

prevalence. In our dataset, ectoparasite diversity and density was centered around medium

to large-bodied, and synanthropic species. Specifically, *R. satarae* and *M. cf. fernandoni*, two

relatively large-bodied and abundant species with high *Bartonella* prevalence, displayed

polyparasitism, often with high densities of specific ectoparasites. This pattern aligns with

the positive correlation between body size and ectoparasitism observed in other studies

[61]. Potentially, habitat sharing of these species with other small mammals facilitates novel

ectoparasite acquisitions and driving overall ectoparasite richness on them [62,63].

Host sharing was another predominant pattern in the ectoparasite community. Several

ectoparasites (e.g., *Rhipicephalus sp*, *Xenopsylla sp 1*) showed a wider host range,

presumably due to their ability to infect multiple locally available hosts [64]. These results

are consistent with previous studies and indicate that host sharing among vectors is more

common than previously believed [65]. Such non-specific host-ectoparasite associations can

facilitate transmission of several pathogens, making their ecology a crucial part of vector-

borne pathogen dynamics [66].

Contrary to our expectations based on host density, our results revealed ectoparasitism

(aggregated ectoparasite load per species) was the most important predictor for

prevalence. While we did not find the patterns we expected for *Bartonella* prevalence, the

correlation between ectoparasitism and *Bartonella* prevalence holds ecological and

epidemiological significance. *Bartonella* is a vector-borne pathogen known for its

transmission through blood-feeding arthropods [67]. Hence, elevated densities of infected

ectoparasites in the environment (and on a specific host) can increase transmission (density-

dependant transmission) and impact prevalence and risk of infection [68]. Polyparasitism

and lack of host specialization suggests high chances of *Bartonella* transmission between

hosts through ectoparasites. In addition, known synanthropic species such as *R. rattus* with

multiple generalist fleas (occur in many hosts as demonstrated in this study) pose risks of

Bartonella transmission to humans and domestic animals [67].

Unfortunately, ecological drivers of aggregated ectoparasite load remain poorly understood.

As discussed earlier, longitudinal sampling targeting larger sample sizes from each

representative species in combination with ecological models incorporating additional

appropriate variables (e.g., host traits, season, etc.) might help explain variance in

ectoparasitism. We caution that other variables previously shown to influence pathogen

prevalence, like seasonality [9] and environmental stressors [69] were unavailable in our

study.

(c) Non-specific ectoparasite-*Bartonella* associations and evidence of historic spillover

suggest future spillover risk

Characterizing *Bartonella* diversity and its association with small mammals and

ectoparasites is crucial from a public health perspective, especially since many of the

Bartonella species are zoonotic and known to associate with small mammals and their

ectoparasites [70]. For instance, endemic *Bartonella* variants such as *B. bacilliformis*,

responsible for Carrion's disease, can be fatal and cause serious mortality [71]. Five (of

eight) *Bartonella* lineages identified in this study share a close evolutionary relationship with

known zoonotic species such as *B. queenslandensis* and *B. tribocorum*. Previous studies have

documented the widespread presence and global circulation of these *Bartonella* species,

primarily facilitated by small mammals and domestic animals [72]. However, it is important

to note that phylogenetic similarity alone does not confirm their ability to cause disease in

humans. Additional validation through molecular and *in vitro* experiments is warranted to

establish their pathogenicity in humans [73,74].

We observed contrasting patterns in *Bartonella* association with small mammals and

ectoparasites. In hosts, *Bartonella* displayed species-specific associations, as we expected

based on previous studies [27,75]. Remarkably, *Bartonella* showed non-specific association

with ectoparasites (e.g*., Xenopsylla sp* 5 collected from *S. niger* carried *Bartonella* associated

with *R. satarae*), leading to a higher overall diversity of haplotypes in specific ectoparasites

compared to their corresponding hosts. These complex patterns are not entirely

unprecedented, as they have been documented elsewhere [76]. Nevertheless, these non-

specific associations between *Bartonella* and ectoparasites have significance for cross-

species transmission, given vector competency.

Our phylogenetic ancestral trait reconstruction also highlighted the possibility for cross-

species transmission in the community. We observed *R. satarae* as the ancestral host for

several *Bartonella* lineages with a medium to high probability, indicating past spillover

events involving this species. This was evident between the BL7 and BL8 lineages, recovered

from *R. satarae* and *M. cf. fernandoni*, respectively. Interestingly, these two species stood

out as the primary *Bartonella* hosts in our dataset, displaying high prevalence, high

haplotype diversity, and co-occurrence in forested habitats. Such pathogen movement

between host species could provide an opportunity to evolve new virulent variants through

recombination [4], similar to those observed in other pathogen systems [6]. Although we

did not find instances of contemporary spillover, our results suggest past spillover events,

potentially through ectoparasites, emphasizing the likelihood of similar events. The

communities we studied include many small mammals and their ectoparasites that overlap

between habitats, increasing the likelihood of such inter-species transmissions [77]. In

addition, the movement of generalist hosts across multiple habitats carrying generalist

ectoparasites with non-specific *Bartonella* associations is likely to further drive pathogen

spillover in the landscape [78].

5. Conclusion

This study has uncovered several significant insights and nuances about the intricate

relationship between small mammal hosts, their ectoparasites, and bacterial agents within a

tropical mixed-use landscape. While our model system and investigation were specific to

Bartonella, the results and inferences extend to a broader spectrum of vector-borne

bacterial pathogens in multi-host communities. Our research highlights the crucial role of

ectoparasitism in driving *Bartonella* prevalence within these small mammal communities

where some hosts move across habitat boundaries. Our results also highlight the non-

specific association of ectoparasites with *Bartonella* diversity compared to their hosts,

underscoring the potential for ectoparasite-driven cross-species transmissions. Using

evolutionary analyses, we further demonstrate the role of historical spillover events in

shaping the current *Bartonella* distribution in the community and highlight the future

prospects of such spillover events. Collectively, our findings underscore the need for

comprehensive ecological studies to advance our understanding of the transmission

dynamics of vector-borne bacterial pathogens within multi-host communities and their

zoonotic implications, particularly in rapidly changing landscapes.

Ethics statement

This study and associated protocols were approved by National Centre for Biological Science

Institutional Animal Ethics Committee (NCBS-IAEC-2016/10-[M], NCBS-IAE-2020/ 02[N]) and

Institutional Biosafety Committee (TFR:NCBS:23_IBSC/2017).

Data accessibility

The manuscript and Supplementary materials present small mammal capture data and

Bartonella prevalence data from small mammals and ectoparasites. *Bartonella* sequence

data generated during this study are deposited in GenBank under the accession numbers

OR574780-OR574826.

Author contributions

ABR: conceptualization, methodology, investigation, data curation, formal analysis,

visualization, project administration, writing -original draft, writing-review and editing; AV:

methodology, data curation, formal analysis, visualization, writing-review and editing; VR:

methodology, investigation, writing-review and editing; HMY: methodology, investigation,

writing-review and editing; AS: investigation, writing-review and editing; UR:

conceptualization, funding acquisition, project administration, supervision, writing-review

and editing.

Acknowledgement

This study was primarily supported by the Department of Atomic Energy, Government of

India (Project Identification RTI 4006) awarded to UR. We are grateful to Mr. K.M. Cariappa

of Kadamane for facilitating the field study; Karnataka Forests and Wildlife Department and

Kudremukh Wildlife Division for granting permission for small mammal sampling;

Kudremukh Iron Ore Company for providing accommodation in Kudremukh. We thank S.

Yashas, Vikrant Jathar, S. Vijayakumar, K. Kamaraj, Tamil Selvan, Subramanian, and C.

Karnan for their help in field sampling. We also acknowledge the support of NCBS Sanger

sequencing facility and museum and field Stations facility. ABR was supported by the SPM

fellowship by CSIR-HRDG, Government of India. ABR also acknowledges the support from

the Tata Institute for Genetics and Society during the manuscript preparation. AS was

supported by fellowship from the Trivedi School of Biosciences, Ashoka University.

References

1. Dirzo R, Young HS, Galetti M, Ceballos G, Isaac NJB, Collen B. 2014 Defaunation in the
Anthropocene. *Science* **345**, 401–406. (doi:10.1126/science.1251817)

2. Gibb R, Redding DW, Chin KQ, Donnelly CA, Blackburn TM, Newbold T, Jones KE. 2020
Zoonotic host diversity increases in human-dominated ecosystems. *Nature* **584**, 398–
402. (doi:10.1038/s41586-020-2562-8)

3.   Hassell JM, Begon M, Ward MJ, Fèvre EM. 2017 Urbanization and Disease Emergence:
Dynamics at the Wildlife–Livestock–Human Interface. *Trends Ecol. Evol.* **32**, 55–67.
(doi:10.1016/j.tree.2016.09.012)

4.   Borremans B, Faust C, Manlove KR, Sokolow SH, Lloyd-Smith JO. 2019 Cross-species
pathogen spillover across ecosystem boundaries: mechanisms and theory. *Phil Trans R*
Soc B **374**, 9. (doi:doi.org/10.1098/rstb.2018.0344)

5.   Faust CL, McCallum HI, Bloomfield LSP, Gottdenker NL, Gillespie TR, Torney CJ, Dobson
AP, Plowright RK. 2018 Pathogen spillover during land conversion. *Ecol. Lett.* **21**, 471–
483. (doi:10.1111/ele.12904)

6.   Wang L-F, Eaton BT. 2007 Bats, Civets and the Emergence of SARS. In *Wildlife and*
Emerging Zoonotic Diseases: The Biology, Circumstances and Consequences of Cross-
Species Transmission, pp. 325–344. Berlin, Heidelberg: Springer Berlin Heidelberg.
(doi:10.1007/978-3-540-70962-6_13)

7.   Pérez-Losada M, Arenas M, Galán JC, Palero F, González-Candelas F. 2015
Recombination in viruses: Mechanisms, methods of study, and evolutionary
consequences. *Infect. Genet. Evol.* **30**, 296–307. (doi:10.1016/j.meegid.2014.12.022)

8.   Plowright RK, Parrish CR, McCallum H, Hudson PJ, Ko AI, Graham AL, Lloyd-Smith JO.
2017 Pathways to zoonotic spillover. *Nat. Rev. Microbiol.* **15**, 502–510.
(doi:10.1038/nrmicro.2017.45)

9.   Rodríguez-Pastor R, Escudero R, Lambin X, Vidal MD, Gil H, Jado I, Rodríguez-Vargas M,
Luque-Larena JJ, Mougeot F. 2019 Zoonotic pathogens in fluctuating common vole
(Microtus arvalis) populations: occurrence and dynamics. *Parasitology* **146**, 389–398.
(doi:10.1017/S0031182018001543)

10.   Stafford KC, Cartter ML, Magnarelli LA, Ertel S-H, Mshar PA. 1998 Temporal Correlations
between Tick Abundance and Prevalence of Ticks Infected with Borrelia burgdorferi and
Increasing Incidence of Lyme Disease. *J. Clin. Microbiol.* **36**, 1240–1244.
(doi:10.1128/JCM.36.5.1240-1244.1998)

11.   Luis AD *et al.* 2013 A comparison of bats and rodents as reservoirs of zoonotic viruses:
are bats special? *Proc. R. Soc. B Biol. Sci.* **280**, 20122753. (doi:10.1098/rspb.2012.2753)

12.   Han BA, Schmidt JP, Bowden SE, Drake JM. 2015 Rodent reservoirs of future zoonotic
diseases. *Proc. Natl. Acad. Sci.* **112**, 7039–7044. (doi:10.1073/pnas.1501598112)

13.   Johnson CK, Hitchens PL, Pandit PS, Rushmore J, Evans TS, Young CCW, Doyle MM. 2020
Global shifts in mammalian population trends reveal key predictors of virus spillover
risk. *Proc. R. Soc. B Biol. Sci.* **287**, 20192736. (doi:10.1098/rspb.2019.2736)

14.   Burgin CJ, Colella JP, Kahn PL, Upham NS. 2018 How many species of mammals are
there? *J. Mammal.* **99**, 1–14. (doi:10.1093/jmammal/gyx147)

15. Andreazzi CS, Martinez-Vaquero LA, Winck GR, Cardoso TS, Teixeira BR, Xavier SCC,
Gentile R, Jansen AM, D'Andrea PS. 2023 Vegetation cover and biodiversity reduce
parasite infection in wild hosts across ecological levels and scales. *Ecography* **2023**,
e06579. (doi:10.1111/ecog.06579)

16. Morand S *et al.* 2015 Assessing the distribution of disease-bearing rodents in human-
modified tropical landscapes. *J. Appl. Ecol.* **52**, 784–794. (doi:10.1111/1365-2664.12414)

17. Young HS *et al.* 2014 Declines in large wildlife increase landscape-level prevalence of
rodent-borne disease in Africa. *Proc. Natl. Acad. Sci.* **111**, 7036–7041.
(doi:10.1073/pnas.1404958111)

18. Allan BF, Keesing F, Ostfeld RS. 2003 Effect of Forest Fragmentation on Lyme Disease
Risk. *Conserv. Biol.* **17**, 267–272. (doi:10.1046/j.1523-1739.2003.01260.x)

19. Estavillo C, Pardini R, Rocha PLB da. 2013 Forest Loss and the Biodiversity Threshold: An
Evaluation Considering Species Habitat Requirements and the Use of Matrix Habitats.
PLoS ONE **8**, e82369. (doi:10.1371/journal.pone.0082369)

20. García-Peña GE, Rubio AV, Mendoza H, Fernández M, Milholland MT, Aguirre AA, Suzán
G, Zambrana-Torrelio C. 2021 Land-use change and rodent-borne diseases: hazards on
the shared socioeconomic pathways. *Philos. Trans. R. Soc. B Biol. Sci.* **376**, 20200362.
(doi:10.1098/rstb.2020.0362)

21. Brooks TM *et al.* 2002 Habitat Loss and Extinction in the Hotspots of Biodiversity.
Conserv. Biol. **16**, 909–923. (doi:10.1046/j.1523-1739.2002.00530.x)

22. Ellis-Cockcroft I, Cotter J. 2014 Tropical Forest Fragmentation; Implications for
Ecosystem Function. , 1–15.

23. Allen T, Murray KA, Zambrana-Torrelio C, Morse SS, Rondinini C, Di Marco M, Breit N,
Olival KJ, Daszak P. 2017 Global hotspots and correlates of emerging zoonotic diseases.
Nat. Commun. **8**, 1124. (doi:10.1038/s41467-017-00923-8)

24. Jones KE, Patel NG, Levy MA, Storeygard A, Balk D, Gittleman JL, Daszak P. 2008 Global
trends in emerging infectious diseases. *Nature* **451**, 990–993.
(doi:10.1038/nature06536)

25. Krishnaswamy J, Bunyan M, Mehta VK, Jain N, Karanth KU. 2006 Impact of iron ore
mining on suspended sediment response in a tropical catchment in Kudremukh,
Western Ghats, India. *For. Ecol. Manag.* **224**, 187–198.
(doi:10.1016/j.foreco.2005.12.018)

26. Baker J, French K, Whelan RJ. 2002 The Edge Effect and Ecotonal Species: Bird
Communities Across a Natural Edge in Southeastern Australia. *Ecology* **83**, 3048–3059.
(doi:10.1890/0012-9658(2002)083[3048:TEEAES]2.0.CO;2)

27. Ansil BR, Mendenhall IH, Ramakrishnan U. 2021 High prevalence and diversity of
Bartonella in small mammals from the biodiverse Western Ghats. *PLoS Negl. Trop. Dis.*
15, e0009178. (doi:10.1371/journal.pntd.0009178)

28. Lei BR, Olival KJ. 2014 Contrasting Patterns in Mammal–Bacteria Coevolution: Bartonella
and Leptospira in Bats and Rodents. *PLoS Negl. Trop. Dis.* **8**, e2738.
(doi:10.1371/journal.pntd.0002738)

29. Neves ES, Mendenhall IH, Borthwick SA, Su YCF, Smith GJD. 2018 Detection and genetic
characterization of diverse Bartonella genotypes in the small mammals of Singapore.
Zoonoses Public Health **65**, e207–e215. (doi:10.1111/zph.12430)

30. Ellerman JR. 1961 *The fauna of India including Pakistan, Burma and Ceylon*. Zoological
Survey of India.

31. Johnsingh A, Manjrekar N, editors. 2015 *Mammals of South Asia*. Universities Press.

32. Menon V. 2014 *Indian mammals: a field guide*. Hachette India.

33. He K, Li Y-J, Brandley MC, Lin L-K, Wang Y-X, Zhang Y-P, Jiang X-L. 2010 A multi-locus
phylogeny of Nectogalini shrews and influences of the paleoclimate on speciation and
evolution. *Mol. Phylogenet. Evol.* **56**, 734–746. (doi:10.1016/j.ympev.2010.03.039)

34. Wolda H. 1981 Similarity indices, sample size and diversity. *Oecologia* **50**, 296–302.
(doi:10.1007/BF00344966)

35. Oksanen J *et al.* 2019 vegan. R package version 2.5-6.

36. Seber G. 1982 *The Estimation of Animal Abundance and Related Parameters*. 2nd edn.
Macmillan, New York.

37. Baillargeon S, Rivest L-P. 2007 Rcapture: Loglinear Models for Capture-Recapture in R. *J.*
Stat. Softw. **19**, 31.

38. Iyengar R. 1973 The Siphonaptera of the Indian Subregion. *Orient. Insects* **7**, 1–91.
(doi:10.1080/00305316.1973.11745225)

39. Geevarghese G, Mishra A C. 2011 *Haemaphysalis Ticks of India*. National Institute of
Virology, Pune, Maharashtra, India.

40. Elango A. 2022 Hands-on Training Module on Medically Important Hard Ticks (Ixodidae)
ICMR- Vector Control Research Centre. , 1–44.

41. Mullen GR, OConnor BM. 2019 Mites (Acari). In *Medical and Veterinary Entomology*, pp.
533–602. Elsevier. (doi:10.1016/B978-0-12-814043-7.00026-1)

42. Bray JR, Curtis JT. 1957 An Ordination of the Upland Forest Communities of Southern
Wisconsin. *Ecol. Monogr.* **27**, 325–349. (doi:10.2307/1942268)

43. R Core Team. 2020 R: A language and environment for statistical computing.

44. Wickham H. 2011 ggplot2: R package Version 3.3.2.

45. Renesto P, Gouvernet J, Drancourt M, Roux V, Raoult D. 2001 Use of rpoB Gene Analysis
for Detection and Identification of Bartonella Species. *J. Clin. Microbiol.* **39**, 430–437.
(doi:10.1128/JCM.39.2.430-437.2001)

46. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990 Basic Local Alignment Search
Tool. *J. Mol. Biol.* **215**, 403–410. (doi:10.1016/S0022-2836(05)80360-2)

47. Harrell Jr F. 2019 'Hmisc'. R package version 4.2-0.

48. Kumar S, Stecher G, Tamura K. 2016 MEGA7: Molecular Evolutionary Genetics Analysis
Version 7.0 for Bigger Datasets. *Mol. Biol. Evol.* **33**, 1870–1874.
(doi:doi:10.1093/molbev/msw054)

49. Lanfear R, Frandsen PB, Wright AM, Senfeld T, Calcott B. 2016 PartitionFinder 2: New
Methods for Selecting Partitioned Models of Evolution for Molecular and Morphological
Phylogenetic Analyses. *Mol. Biol. Evol.* **34**, 260. (doi:10.1093/molbev/msw260)

50. Drummond AJ, Suchard MA, Xie D, Rambaut A. 2012 Bayesian Phylogenetics with BEAUti
and the BEAST 1.7. *Mol. Biol. Evol.* **29**, 1969–1973. (doi:10.1093/molbev/mss075)

51. Rambaut A, Drummond AJ, Xie D, Baele G, Suchard MA. 2018 Posterior Summarization
in Bayesian Phylogenetics Using Tracer 1.7. *Syst. Biol.* **67**, 901–904.
(doi:10.1093/sysbio/syy032)

52. Rambaut A. 2014 FigTree 1.4. 2 Software.

53. Charleston MA, Robertson DL. 2002 Preferential Host Switching by Primate Lentiviruses
Can Account for Phylogenetic Similarity with the Primate Phylogeny. *Syst. Biol.* **51**, 528–
535. (doi:10.1080/10635150290069940)

54. Hill JG, Summerville KS, Brown RL. 2008 Habitat Associations of Ant Species
(Hymenoptera: Formicidae) in a Heterogeneous Mississippi Landscape. *Environ.*
Entomol. **37**, 11. (doi:doi.org/10.1093/ee/37.2.453)

55. Stephens RB, Anderson EM. 2014 Habitat associations and assemblages of small
mammals in natural plant communities of Wisconsin. *J. Mammal.* **95**, 404–420.
(doi:10.1644/13-MAMM-A-025)

56. Ebani VV. 2022 Commensal Rodents: Still a Current Threat. *Pathogens* **11**, 1483.
(doi:10.3390/pathogens11121483)

57. Ramachandra TV, Shivamurthy V, Aithal DrB. 2015 *Environmental Flow Assessment in*
Yettinaholé: Where is 24 TMC to divert? (doi:10.13140/RG.2.1.5136.4323)

58. Lindström J, Ranta E, Kokko H, Lundberg P, Kaitala V. 2001 From arctic lemmings to
adaptive dynamics: Charles Elton's legacy in population ecology. *Biol. Rev. Camb. Philos.*
Soc. **76**, 129–158. (doi:10.1017/S1464793100005637)

59. Myers JH. 2018 Population cycles: generalities, exceptions and remaining mysteries.
Proc. R. Soc. B Biol. Sci. **285**, 20172841. (doi:10.1098/rspb.2017.2841)

60. Gibb R, Franklinos LHV, Redding DW, Jones KE. 2020 Ecosystem perspectives are needed
to manage zoonotic risks in a changing climate. *BMJ* , m3389. (doi:10.1136/bmj.m3389)

61. Esser HJ, Foley JE, Bongers F, Herre EA, Miller MJ, Prins HHT, Jansen PA. 2016 Host body
size and the diversity of tick assemblages on Neotropical vertebrates. *Int. J. Parasitol.*
Parasites Wildl. **5**, 295–304. (doi:10.1016/j.ijppaw.2016.10.001)

62. Bordes F *et al.* 2015 Habitat fragmentation alters the properties of a host-parasite
network: rodents and their helminths in South-East Asia. *J. Anim. Ecol.* **84**, 1253–1263.
(doi:10.1111/1365-2656.12368)

63. Dáttilo W, Barrozo-Chávez N, Lira-Noriega A, Guevara R, Villalobos F, Santiago-Alarcon
D, Neves FS, Izzo T, Ribeiro SP. 2020 Species-level drivers of mammalian ectoparasite
faunas. *J. Anim. Ecol.* **89**, 1754–1765. (doi:10.1111/1365-2656.13216)

64. McCoy KD, Léger E, Dietrich M. 2013 Host specialization in ticks and transmission of tick-
borne diseases: a review. *Front. Cell. Infect. Microbiol.* **3**.
(doi:10.3389/fcimb.2013.00057)

65. Goldberg AR, Conway CJ, Biggins DE. 2020 Flea sharing among sympatric rodent hosts:
implications for potential plague effects on a threatened sciurid. *Ecosphere* **11**.
(doi:10.1002/ecs2.3033)

66. Sotomayor-Bonilla J, Tolsá-García MJ, García-Peña GE, Santiago-Alarcon D, Mendoza H,
Alvarez-Mendizabal P, Rico-Chávez O, Sarmiento-Silva RE, Suzán G. 2019 Insights into
the Host Specificity of Mosquito-Borne Flaviviruses Infecting Wild Mammals. *EcoHealth*
16, 726–733. (doi:10.1007/s10393-019-01442-2)

67. Moreno Salas L, Espinoza-Carniglia M, Lizama Schmeisser N, Torres LG, Silva-de la
Fuente MC, Lareschi M, González-Acuña D. 2019 Fleas of black rats (Rattus rattus) as
reservoir host of Bartonella spp. in Chile. *PeerJ* **7**, e7371. (doi:10.7717/peerj.7371)

68. Kantsø B, Svendsen CB, Jensen PM, Vennestrøm J, Krogfelt KA. 2010 Seasonal and
habitat variation in the prevalence of Rickettsia helvetica in Ixodes ricinus ticks from
Denmark. *Ticks Tick-Borne Dis.* **1**, 101–103. (doi:10.1016/j.ttbdis.2010.01.004)

69. McMichael L, Edson D, Smith C, Mayer D, Smith I, Kopp S, Meers J, Field H. 2017
Physiological stress and Hendra virus in flying-foxes (Pteropus spp.), Australia. *PLoS ONE*
12. (doi:10.1371/journal.pone.0182171)

70. Saisongkorh W, Rolain J-M, Suputtamongkol Y, Raoult D. 2009 Emerging Bartonella in
Humans and Animals in Asia and Australia. *J Med Assoc Thai* **92**, 25.

71. Garcia-Quintanilla M, Dichter AA, Guerra H, Kempf VAJ. 2019 Carrion's disease: more
than a neglected disease. *Parasit. Vectors* **12**, 141. (doi:10.1186/s13071-019-3390-2)

72. Frank HK, Boyd SD, Hadly EA. 2018 Global fingerprint of humans on the distribution of
Bartonella bacteria in mammals. *PLoS Negl. Trop. Dis.* **12**, e0006865.
(doi:10.1371/journal.pntd.0006865)

73. Saenz HL, Dehio C. 2005 Signature-tagged mutagenesis: technical advances in a negative
selection method for virulence gene identification. *Curr. Opin. Microbiol.* **8**, 612–619.
(doi:10.1016/j.mib.2005.08.013)

74. Che D, Hasan M, Chen B. 2014 Identifying Pathogenicity Islands in Bacterial
Pathogenomics Using Computational Approaches. *Pathogens* **3**, 36–56.
(doi:10.3390/pathogens3010036)

75. Withenshaw SM, Devevey G, Pedersen AB, Fenton A. 2016 Multihost Bartonella
parasites display covert host specificity even when transmitted by generalist vectors. *J.*
Anim. Ecol. **85**, 1442–1452. (doi:10.1111/1365-2656.12568)

76. Silaghi C, Pfeffer M, Kiefer D, Kiefer M, Obiegala A. 2016 Bartonella, Rodents, Fleas and
Ticks: a Molecular Field Study on Host-Vector-Pathogen Associations in Saxony, Eastern
Germany. *Microb. Ecol.* **72**, 965–974. (doi:10.1007/s00248-016-0787-8)

77. Boyard C, Vourc'h G, Barnouin J. 2008 The relationships between Ixodes ricinus and
small mammal species at the woodland–pasture interface. *Exp. Appl. Acarol.* **44**, 61–76.
(doi:10.1007/s10493-008-9132-3)

78. Becker DJ, Streicker DG, Altizer S. 2018 Using host species traits to understand the
consequences of resource provisioning for host-parasite interactions. *J. Anim. Ecol.* **87**,
511–525. (doi:10.1111/1365-2656.12765)

799