Unravelling arthropod movement in natural landscapes: small-scale effects of body size and weather conditions

Garben Logghe[1,2]* (ORCID: 0000-0003-2000-7982)

Charlotte Taelman[1,2]* (ORCID: 0000-0003-2701-8297)

Florian Van Hecke[1,2] (ORCID: 0009-0004-3499-2279)

Femke Batsleer[1] (ORCID: 0000-0001-5893-5966)

Dirk Maes[2,3] (ORCID: 0000-0002-7947-3788)

Dries Bonte[1] (ORCID: 0000-0002-3320-7505)

1 Ghent University, Department of Biology, Terrestrial Ecology Unit, Karel Lodewijk Ledeganckstraat
35, 9000 Ghent, Belgium

2 Research Institute for Nature and Forest (INBO), Herman Teirlinckgebouw, Havenlaan 88 Bus 73,
1000 Brussels, Belgium

3 Radboud Institute for Biological and Environmental Sciences (RIBES), Radboud University, PO Box
9010, NL-6500 GL Nijmegen, The Netherlands

Corresponding author: G. Logghe, Ghent University, Department of Biology, Terrestrial Ecology Unit,
K. L. Ledeganckstraat 35, 9000 Ghent, Belgium. E-mail: garben.logghe@ugent.be

Keywords: insects, movement speed, movement direction, habitat boundaries, matrix

*These two authors share first authorship

Abstract

20   1. Arthropod movement has been noticeably understudied compared to vertebrates. A crucial knowledge gap pertains to the factors influencing arthropod movement at habitat boundaries, which has direct implications for population dynamics and gene flow. While larger arthropod species generally achieve greater dispersal distances and large-scale movements are affected by weather conditions, the applicability of these relationships at a local scale remains uncertain. Existing studies on this subject are not only scarce but often limited to a few species or laboratory conditions.

27   2. To address this knowledge gap, we conducted a field study in two nature reserves in Belgium, focusing on both flying and cursorial (non-flying) arthropods. Over 200 different arthropod species were captured and released within a circular setup placed in a resource-poor environment, allowing quantification of movement speed and direction. By analysing the relationship between these movement variables and morphological (body size) as well as environmental factors (temperature and wind), we aimed to gain insights into the mechanisms driving arthropod movement at natural habitat boundaries.

34   3. For flying species, movement speed was positively correlated with both body size and tailwind speed. In contrast, movement speed of cursorial individuals was solely positively related with temperature. Notably, movement direction was biased towards the vegetated areas where the arthropods were originally caught, suggesting an internal drive to move towards suitable habitat. This tendency was particularly strong in larger flying individuals and under tailwind conditions. Furthermore, both flying and cursorial taxa were hindered from moving towards the habitat by strong upwind.

41   4. In conclusion, movement speed and direction at patch boundaries are dependent on body size and prevailing weather conditions, and reflect an active decision-making process.

1. Background

Movement is a crucial component of the life history of many organisms. Across small and large spatial scales, it mitigates the finding of resources and mates, ultimately shaping species distributions and interactions (Bowler & Benton, 2005; Gilbert et al., 1998; McIntyre & Wiens, 1999). Studies involving animal movement tend to focus on vertebrates, leaving other groups such as arthropods severely underrepresented, despite the latter constituting a substantial portion of the global biodiversity (Clark & May, 2002; Kissling et al., 2014). Admittedly, studying arthropod movement involves many inherent challenges, such as their small size and vast species diversity. Nevertheless, understanding the ecological dynamics of arthropods is crucial, as they play pivotal roles in ecosystem functions such as pollination, nutrient cycling and trophic interactions (Yang & Gratton, 2014).

Natural systems are often composed of mosaic landscapes, in which the area is heterogeneously divided into habitats that widely differ in type and suitability (Forman, 2014). Within such landscapes, arthropods may encounter challenging environments while seeking resources or evading unfavourable conditions (Lima & Zollner, 1996; Zollner & Lima, 1999). These unfavourable zones are known as "matrix", whereas suitable areas are considered "habitat" (Ricketts, 2001; Shreeve & Dennis, 2011). Movement and orientation behaviours at habitat-matrix boundaries will influence the fate of each individual: drifting further in the matrix and losing any connection with suitable environment, such as the home patch or any other potential habitat, or returning to the home patch and remaining philopatric. Unlike routine movements within a habitat, compensatory movement (Peterman et al., 2014; Schtickzelle et al., 2006) through the matrix tends to be rapid with a higher degree of directionality (Hein et al., 2003; Mola et al., 2020; Turlure et al., 2011; Van Dyck & Baguette, 2005). Consequently, it can be expected that crossing a hostile matrix will impose substantial dispersal costs (Bonte et al., 2012), particularly for small-bodied organisms like arthropods (Hillaert et al., 2018). Behaviours at boundaries are therefore expected to have pervasive

effects on population dynamics, gene flow and hence the adaptive capacities of species in

fragmented landscapes (Batsleer et al., 2023; Driscoll et al., 2013).

Body size stands out as one of the most relevant factors influencing arthropod movement and their

capacity to orientate, due to its association with various morphological, physiological and life-history

traits (White et al., 2007; Woodward et al., 2005). Numerous arthropod taxa, ranging from cursorial

groups (i.e. individuals running on the ground) like ants and beetles, to flying taxa such as butterflies,

bugs and dragonflies, seem to consequently exhibit an allometric relationship between body size and

mobility (Hirt et al., 2017; Hurlbert et al., 2008; Kuussaari et al., 2014; Sekar, 2012; Wootton, 2020).

Larger arthropod individuals, both within and across species, generally display greater mobility,

resulting in higher movement speeds and larger travelling distances.

Besides species-specific characteristics such as body size, arthropod movement can also be

influenced by external conditions, such as weather. The impact of local weather conditions should be

most apparent for small-scale movement bouts, because of the high variability of weather conditions

at small temporal and spatial scales. A first component of local weather conditions is the prevailing

temperature. Higher ambient temperatures have been shown to positively impact the activity and

running speed of invertebrate ectotherms, owing to improved oxygenation rates and enhanced

kinetic muscle properties (Frizzi, 2018; Hurlbert et al., 2008). Moreover, arthropod species often

adjust their movement behaviour according to the local thermal conditions, with some even

requiring a minimum ambient temperature to initiate movement (Frizzi, 2018; Knight et al., 2019;

Knoblauch et al., 2021). However, movement responses to increased temperatures are also

constrained by critical thermal upper limits, leading to impaired movement capacity when

temperatures exceed these thresholds (Chown & Nicolson, 2004; Dudley, 2000; Heinrich, 1995). A

second component of local weather is wind. The direction and force of the wind exert considerable

influence on movement distances and decisions, especially for flying insects undertaking long-

distance dispersal (Knoblauch et al., 2021). Additionally, many arthropods possess anemotactic

reception, allowing them to accurately estimate the direction of prevailing winds by sensing air

currents, and consequently, make informed choices regarding movement trajectories (Buehlmann et

al., 2020; Knoblauch et al., 2021).

Most of the previously mentioned research on the relationships between arthropod movement,

body size and weather conditions focuses on large-scale distances, and usually involves only a limited

number of large and highly mobile species (Frick et al., 2013; Knight et al., 2019; Knoblauch et al.,

2021). Small-scale movement studies investigating biometeorological relationships in arthropod

movement are scarce (Aikman & Hewitt, 1972; Eguagie, 1974; Merckx et al., 2008) and especially

lacking when considering behaviour at habitat boundaries. Traditional methods, such as capture-

mark-recapture or tracking devices, have limitations in achieving large sample sizes, often leading to

a narrow focus on individual species and hindering generalizations across taxa and life history traits

(Goodwin & Fahrig, 2002; Jonsen et al., 2001). Moreover, concerns have been raised about the use of

such tracking devices, as the impact of these tags on arthropods is often neglected, ultimately biasing

data (Batsleer et al., 2020). While some notable efforts have been made to study arthropod

exploratory movement through image-based tracking under laboratory conditions, the applicability

of these findings to natural field conditions remains uncertain (Hirt et al., 2017; Terlau et al., 2023).

This study aims to understand the interplay between movement, body size and weather conditions

at habitat boundaries by conducting a field release experiment on a diverse range of arthropod

species. In contrast to macro-ecological approaches that treat dispersal as colonisation at

biogeographical scales (Weil et al., 2023), our methodology focuses on individual departure and

settlement behaviour. Multi-species experiments under similar environmental settings have the

potential to yield generalisations on trait-environment correlations across and within species, and to

understand potential evolutionary determinants of these relationships (Felsenstein, 1985).

Two different nature reserves consisting of dune and heathland habitats were used as field sites,

which differed in spatial heterogeneity. This allowed us to assess whether arthropods choose to

move towards suitable habitat areas when venturing in the matrix. The realised movement speed

and direction of arthropods at the patch boundary—without relying on tracking devices—were

measured and correlated with individual body size and prevailing weather conditions. Movement

behaviours after release represent movements that we consider to be relevant for short-term

displacements when organisms leave their habitat and enter the matrix. We expect (i) a positive

correlation between movement and body size, (ii) the strengthening of this effect under warmer

temperatures and stronger tailwinds, (iii) an influence of wind conditions on the movement direction

of flying insects, and (iv) a biased movement direction towards suitable environments due to the

perception (correct interpretation by the individual) of the environment and avoidance of risks when

crossing the matrix.

2. Methods

129   Permits to access the field sites and capture protected species were granted by the Agency of Nature

130   and Forest. This study did not require ethical approval.

2.1 Experimental fieldwork

The fieldwork for this study was conducted in the summer of 2021 at two locations in Belgium. The

first site was a coastal dune grassland in De Panne, while the second site consisted of heathland

patches in Kalmthout (Appendix S.1). Both sites differed in structural heterogeneity of the landscape.

In the coastal dune site at De Panne, large areas of open sand (representing the matrix) bordered

large areas of shrubland, thus creating a "binary" landscape. In Kalmthout, patches of open sand

were considerably smaller, and were embedded within a dense heath landscape, thereby creating a

patchier landscape where matrix and habitat surrounded each other. By selecting these two distinct

sites, we aimed to determine whether arthropod movement direction at habitat boundaries is either

random or biased towards vegetated areas. If arthropods indeed show an internal drive to move

towards habitat patches, we should find a strong bias towards one direction in the "binary"

landscape at De Panne, while movement should be more random in the patchy landscape at

Kalmthout (Appendix S.1).

Within the vegetated areas at both sites, arthropod individuals from various species were randomly

captured either by hand or by sweep netting. Each captured individual was assigned a unique ID and

photographed in the field with a graph paper (1 mm squares) as background. To assess the

movement behaviour of these arthropods, separate set-ups were designed for flying and cursorial

(i.e. running on the ground) individuals, as we anticipated different movement speeds and

perceptual abilities. In a large area of open sand, which functions as matrix for species living in

vegetated areas, two circular set-ups were demarcated by placing coloured flags at five different

radii with uniform interval distances (2 metres for flying individuals and 20 centimetres for cursorial

individuals). The shortest distance from the point of release to habitat measured 1 m and 10 m for

respectively the experiments on cursorial and flying movement (west orientation for De Panne and

northwest orientation for Kalmthout). In accordance with the earlier described habitat

heterogeneity, distance to habitat was more than 100 m in other cardinal directions for De Panne,

while for Kalmthout these distances were less than 50 m.

Movement speed was measured using a chronometer, whereby the movement time was recorded as

the duration an individual took to traverse either 10 meters (for flying locomotion) or 1 meter (for

cursorial locomotion), or when an individual came to stop at a specific distance. The directions of

movement and wind were categorised using cardinal directions (N, NE, E, SE, S, SW, W or NW), and

then converted into angles in degrees, with west serving as the zero degrees reference (see further).

By comparing the movement and wind angles, we were able to determine whether the movement

was aligned with the wind or not, and to differentiate between movement upwind and movement

assisted by tailwinds. Specifically, we defined tailwind movements when the direction of movement

deviated by a maximum of 45 degrees from both sides of the prevailing wind direction, while upwind

movements deviated by at least 135 degrees from both sides of the prevailing wind direction (or 45

degrees from both sides of the direction of the tail; Figure 1). West was chosen as the zero-degree

reference, as the vegetated areas at De Panne were located in this direction relative to the

experimental set-up (Figure 1). This decision assumed that the arthropods would exhibit a preference

for returning to the location where they were initially caught, since they were released in a matrix

with very few available resources. The surface area of the matrix was sufficiently large to assume

that moving through it would bring a high energetic cost and uncertainty for the released arthropod.

By adopting the orientation as shown in Figure 1, we aimed to gain insights in navigation behaviour,

as movement patterns would likely reflect the tendency and ability to return to the habitat of origin.

During measurements, the captured arthropods were temporarily housed in dark boxes to minimize

stress and prevent mortality due to overheating. Each individual spent a maximum of thirty minutes

in these boxes. Subsequently, the arthropods were taken out one by one and released in the open

sand. To mitigate the likelihood of an escaping movement response and directional bias arising from

the researchers' presence, cursorial animals were released at a distance of 1 meter from the

observer, whereas flying animals were released above the head of the observer. Individuals were not

chased but followed by eye, and timed from the moment of release until the moment they reached

the edge of the experimental arena. Every individual had complete freedom of where to move and

when to stop. By doing this, we aimed to minimize stress and present the option to actively choose

movement directions.

Every half-hour of the release experiments, wind direction, wind speed and temperature were

recorded at an average height of 1.5 to 1.8 metres, using a weather station equipped with a built-in

thermometer and anemometer. Wind speed was calculated as the average of the minimum and

maximum values measured within a half-minute interval. Sampling days were carefully selected to

ensure suitable weather conditions for representative arthropod movement: a cloud cover below

60% and temperatures between 15°C and 34°C.

a)

b)

Figure 1: a) Schematic representation of how the cardinal directions were converted into angles. The black arrow indicates the direction of the movement. The blue area indicates all wind directions that can be classified as tailwind in this specific example, while the yellow area indicates upwind. b) Map of the experimental set-up at De Panne, with the vegetated areas to the west. Circular setups were drawn at a realistic scale. The illustrations in this figure were created by author Charlotte Taelman, aerial photograph (winter 2022) obtained from Agency for Information Flanders (geopunt.be).

Due to practical constraints, determining the dry weight of individual arthropods was not feasible.

Instead, body length was used as an alternative proxy for body size (Ganihar, 1997; Martin et al.,

2014; Moretti et al., 2017). The pictures of the arthropods taken in the field were analysed using

ImageJ version 1.52a (Schneider et al., 2012). The grid pattern in the background of the photos was

used as reference to measure the length of each individual's body, starting from the cephalon

(excluding the antennae) and extending to the tip of the abdomen.

2.2 Statistical analyses

All data analyses were conducted using R version 4.2.2 (R Core Team, 2021). The "dplyr" package

(Wickham et al., 2022) was used for general data manipulation and organization. All visuals were

created with the "ggplot2" (Wickham, 2016), "RcolorBrewer" (Neuwirth, 2022) and "showtext" (Qiu,

2023) packages.

2.2.1 Movement speed

As a first explorative analysis, a Principal Component Analysis (PCA) was conducted, separately for

flying and cursorial species to understand how species movement behaviours group according to

their traits, phylogeny and the prevalent environmental conditions.

To confirm any observed relationships between movement speed, body size and weather conditions

in flying and cursorial arthropods, a distinct Bayesian Generalized Linear model was employed for

each group ("brms" package (Bürkner, 2021)). The dependent variable in these models was

movement speed, while the independent variables were body size, temperature and wind speed (all

standardized to a mean of zero and a standard deviation of one). The models were fitted with a

Gaussian distribution using 4 chains and 4000 iterations to ensure reliable results. Non-informative

flat priors were used for the model parameters. As we expected wind speed to have opposing effects

depending on its orientation relative to an individual's movement, we constructed additional models

by splitting the dataset into individuals that moved either tail- or upwind and ran two separate

models. This approach was used to test potential differences in responses and proved to be more

efficient than the incorporation of an interaction between two circular variables to the Bayesian

models.

Since the models did not account for the potential influence of phylogenetic relationships between

species, additional models were constructed to address this issue. For this purpose, phylogenetic

trees were created by linking the species names to the Open Tree Taxonomy database, establishing a

tree based on the known phylogenetic relationships between species ("rotl" package (Michonneau et

al., 2016)). Branch lengths were calculated with the "ape" package (Paradis & Schliep, 2019) using

the "Grafen" method (Grafen, 1989), leading to the construction of a phylogenetic covariance matrix

("geiger" package (Matthew et al., 2014)). First of all, this tree was used to calculate Pagel's lambda,

a measure for the phylogenetic signal in a specific trait, for movement speed and direction of both

flying and cursorial species ("phytools" package (Revell, 2012)). Secondly, the phylogenetic distances

between species were incorporated into all Bayesian Generalized Linear models (similar to those

above) as blocking factors. By doing so, the models considered the phylogenetic relationships

between species when analysing the correlation between movement speed and the other variables.

We contrast insights from the non-corrected models (ecological responses of the sampled arthropod

community) and phylogenetically corrected models (ecological responses of arthropod species,

corrected for the shared, and hence non-independent evolutionary history of life histories and

morphologies). As such, we can understand whether the community-wide movement responses are

a direct outcome from the specific reactions of the sampled species, or whether they have evolved as

a general pattern across species.

2.2.2 Movement direction

The analysis of arthropod movement behaviour in different wind scenarios involved several circular

statistics tests ("circular" package (Agostinelli & Lund, 2022)). These analyses were performed

separately for both sampling sites to analyse the effect of structural heterogeneity within the

landscapes.

To assess the correlation between wind direction and movement direction, the Jammalamadaka-

Sarma correlation coefficient (Jammalamadaka & Sarma, 1988) was calculated for both flying and

cursorial arthropods within different taxonomic orders, limited to orders represented by 8 or more

samples. Additionally, the Rayleigh's test of Uniformity (Pewsey et al., 2013) was performed to

examine whether there was a unimodal directional bias in the distribution of arthropod movement

directions. The significance of the results from these tests was evaluated by correcting for multiple

testing using the Benjamini-Hochberg method (Benjamini & Hochberg, 1995), addressing potential

issues of false positives.

Furthermore, several Bayesian Generalized Linear Models were fitted to explore the relationship

between body size and movement direction. For these analyses, only data from De Panne was used

since the difference in direction between habitat and matrix was much clearer compared to the site

at Kalmthout. A binary value (0 or 1) was assigned to indicate whether or not an individual had

moved towards the habitat from where they were initially caught. Deviations greater than 45

degrees from the zero-degree point (west) were considered as not moving towards the habitat and

were assigned a value of 0. Body size and wind speed were used as independent variables, while this

binary value was the dependent variable, modelled using a binomial distribution with a logit link

function. Temperature was omitted from these models because we considered it irrelevant in the

context of navigation. As with movement speed, the datasets were again split to test whether

different responses were present under upwind or tailwind conditions. As for movement speed, each

model was also corrected for phylogenetic relationships, which involved constructing phylogenetic

trees using Open Tree Taxonomy and incorporating phylogenetic distances as blocking factors in the

models.

3. Results

In total, 1355 arthropod individuals from 255 different species were tested (Appendix S.2). Among these individuals, 762 exhibited flying locomotion, while the remaining 593 displayed cursorial locomotion. Measured temperatures ranged from 15°C to 40°C, while wind speed varied between 0 and 6.4 metres per second.

3.1 Movement speed

The PCA shows Odonata, having larger body sizes, to be separated from the other studied arthropod orders (Figure 2). For both cursorial and flying arthropod groups, PC1 was correlated with temperature and windspeed and PC2 with body size (Appendix S.3). That the first axis explains respectively 36% of the variation in flying species and 34% of the variation in cursorial species, shows that weather conditions are indeed an important determinant of movement. We further explore interspecific variation in more detail by means of (phylogenetically corrected) linear models.

In flying species, large species generally exhibited higher movement speeds compared to smaller ones (Figure 3a, left). However, this relationship appears to reach an optimal point around 15 mm body size, beyond which a further increase in body size does not lead to higher movement speeds (Appendix S.4.2). The overall positive impact of body size and wind speed on movement speed of flying species remains when only tailwind movements are considered, but disappears when movement occurs upwind (Figure 3b, c, left). For tailwind speed, the correlation also demonstrated a saturation point, which occurred around 3 m/s wind speed, beyond which movement speed did not increase further (Appendix S.4.1). The overall positive effect of wind speed on movement speed is thus primarily driven by the effects found under tailwind conditions (Figure 3, left). Temperature had no specific effect on the movement speed of the studied flying insects (Figure 3, left). Pagel's lambda of movement speed was notably high (λ=0.733), indicating a strong phylogenetic signal. Nevertheless, as expected from the overlap among taxa in the PCA, phylogenetic corrections had no substantial qualitative impact on the posterior distributions of the effect sizes and their

interpretation (Figure 3). Species evolutionary history is thus an important driver of movement

speed, independent of the species' attained body sizes. Since a phylogenetic correction did not affect

any of the patterns, responses to weather are shared by all species present in the study area,

independent of their phylogenetic background.

The movement speed of cursorial species was, in contrast, only affected by temperature (Figure 3,

right), with the positive effect indicating higher speeds reached under warmer conditions. The

Pagel's lambda of movement speed for the cursorial species was considerably smaller compared to

flying species (λ=0.357). Closely related cursorial arthropods are therefore more variable in their

movement speed relative to their flying counterparts. For a detailed overview of the results from

these models, see Appendix S.5.1.

Figure 2: PCA results for flying (top) and cursorial (bottom) individuals. Dots represent individuals with symbol colour and shape according to their taxonomic order. A convex polygon is drawn around individuals of the same taxonomic order, by connecting the most extreme values. The arrows represent the direction and magnitude of each variable's contribution to the principal components. Longer arrows indicate stronger influence, while the direction shows the correlation between variables and components. The angle between the arrows reflects the correlation between traits; very small angles indicate strong positive correlation, while very large angles indicate strong negative correlation.

Figure 3: Posterior distributions of the effect sizes between movement speed as dependent variable and body size, temperature and wind speed as independent variables. For a) all wind conditions, b) tailwind conditions (n=527) and c) upwind conditions (n=281). Posterior distributions that are marked with a phylogenetic tree resulted from the phylogenetically corrected models. Effect sizes were deemed clearly determined when 0 was not included in the 95% credibility intervals of the posteriors.

3.2 Movement direction

Arthropod movement exhibited a noticeable directional bias, indicating that individuals did not move randomly (Figure 4). The R-values, representing concentration of movement and thus directional bias, were significant after correction for multiple testing for most taxonomic orders (Appendix S.6). In addition, the Jammalamadaka-Sarma correlation coefficients between wind and movement direction did not show a general pattern across taxa, with some orders following the wind more than others (Appendix S.7). However, these correlations where only significant for Lepidoptera (Kalmthout, r=0.36), Araneae (De Panne, r=0.34) and Diptera (De Panne, r=0.19).

At De Panne, where the habitat of the arthropod species was clearly located in one direction (west), movement direction was mostly oriented towards habitat areas across all observations (Figure 4.1). The lowest R-value at this site (R=0.19; p=0.05) was associated with a southwest wind opposite to the habitat direction, indicating lack of any directionality of movements under these wind conditions. In contrast, the highest R-value and hence strongest directionality prevailed under southeast winds, which largely aligned with the orientation towards the habitat (R=0.71, p<0.01). Furthermore, the Jammalamadaka-Sarma coefficient across all observations was positive (r=0.07), indicating a minor effect of wind direction on movement direction. In contrast, at the Kalmthout site, which was surrounded by suitable habitat in multiple directions, movement directional bias was only significant under NE winds (R=0.44; p<0.01). This indicates that arthropod movement was more random compared to the site at De Panne. Additionally, the Jammalamadaka-Sarma coefficient was larger (r=0.14) compared to De Panne, suggesting that movement direction was more positively aligned with wind direction (Figure 4.2).

Movement direction did not only correlate to wind direction, but additionally to wind speed as well. We ran Bayesian generalized linear models for the experiments in De Panne only, which showed that higher wind speeds reduce the efficiency of flying individuals to move towards habitat (Figure 5a, left). This effect was mainly steered by individuals tested under upwind conditions (Figure 5c, left).

Similarly, cursorial species only showed a negative correlation between wind speed and movement

direction under upwind conditions (Figure 5, right). This indicates that strong upwinds lower the

propensity of both groups to navigate back to the habitat. These patterns remained conserved under

phylogenetic correction ($\lambda < 0.001$), indicating generally similar arthropod-wide responses. Habitat-

directed movements also differed with body size. Larger flying species are more prone to move

towards habitat, while the pattern is reversed for cursorial species (Fig 5a, right). Additionally, the

effect of wind speed on cursorial navigation only holds under upwind conditions when

phylogenetically corrected ($\lambda = 0.210$). Our observed community-wide responses were thus biased by

the specific sample of tested species. For a detailed overview of the results from these models, see

Appendix S.5.2.

Figure 4.1: Circular bar plots of movement directions of flying species at both field sites. Each concentric slice on the

bar represents an individual and is coloured according to its body size. Thus, larger bars indicate a higher number of

individuals that moved in that direction. Similar figures for specific wind directions can be found in Appendix S.8.

349
350 *Figure 4.2: Circular bar plots of dominant wind directions at both field sites. The larger the slices, the more observations*

351 *were made under that specific wind direction. Wind speed data for each wind direction can be found in Appendix S.9.*

Figure 5: Posterior distributions of the effect sizes between movement direction as dependent variable and body size and wind speed as independent variables. For a) all wind conditions, b) tailwind conditions (n=440) and c) upwind conditions (n=248). Only data from the De Panne site was used for this analysis. Posterior distributions that are marked with a phylogenetic tree resulted from the phylogenetically corrected models. Effect sizes were deemed clearly determined when 0 was not included in the 95% credibility intervals of the posteriors.

4. Discussion

We used release experiments to gain insights into the condition-dependency of small-scale arthropod movement at their patch boundary. Movements at this spatiotemporal scale can determine the future trajectories within or back from the matrix in fragmented habitats. These fundamental movement elements are anticipated to eventually shape large-scale canonical and lifetime movement patterns (Getz & Saltz, 2008), and thus patterns of gene flow, colonisation-extinction dynamics and species assembly in fragmented landscapes (Batsleer et al., 2023; Jeltsch et al., 2013). Using a standardised approach under natural conditions, we demonstrate the importance of phylogeny, body size and weather conditions on arthropod movement in the hostile matrix close to the patch boundary. Larger flying arthropods exhibited greater movement speeds compared to their smaller counterparts, while the presence of strong tailwinds further increased their movement speed. In contrast, the movement speed of cursorial arthropods appeared unrelated to body size but increased with temperature. Notably, our observations revealed a clear directional bias towards suitable habitat at both sites, indicating an internal drive among arthropods to move towards suitable environments. However, species showed varying levels of navigational proficiency, with larger flying and smaller cursorial arthropods displaying a greater tendency to move in the direction of suitable habitat. Furthermore, our study identified strong upwinds as hindering the movement of both groups, impeding the capacity for navigation towards suitable habitat.

The positive correlation between movement speed and body size in flying arthropods was in alignment with our expectations. Previous studies already established that—within a taxon—larger arthropods tend to move faster (Kuussaari et al., 2014; Wootton, 2020). Our study emphasizes that this effect remains consistent across multiple taxa, even after accounting for phylogenetic relationships. Consequently, there is a difference in movement speed between taxa, but the positive relationship between movement speed and body size remains conserved. For example, larger bees

are generally faster than smaller bees or smaller butterflies, but do not necessarily surpass the pace

of a larger dragonfly.

In contrast, the absence of a correlation between movement speed and body size in cursorial

arthropods does not match the positive relationship retrieved in previous studies (Foellmer et al.,

2011; Hirt et al., 2017; Hurlbert et al., 2008). A first possible explanation is that a relatively limited

range of body sizes was tested, spanning from 1.21 mm to 21.31 mm, preventing us from detecting a

signal when effects are mainly determined by very large species. Secondly, our proxy for body size

could obscure true relationships. For instance, harvestmen may appear small in our dataset due to

their compact body length, but their long legs could enable faster running compared to species with

similar body length (Arnold et al., 2017; Kaspari & Weiser, 1999; Smith et al., 2012). The use of a

relative measure (e.g. leg or wing length relative to body length) remains, however, hard to

generalize across taxa and thus limits the applicability for cross-taxa comparisons. A third potential

explanation is linked to the interaction between the body mass of an individual and the texture of

the surface on which it moves. Sand is a porous substance, in which heavier individuals might sink

deeper than smaller ones, hence slowing their movement.

In addition to body size, wind speed also positively influenced the movement speed of flying

individuals. Conversely, when individuals faced upwind, the effect reversed. This implies that

movements away from habitat boundaries may be more pronounced when arthropods fly with

tailwind, which can assist them in crossing the matrix (DiLeo et al., 2022). However, under natural

conditions, the prospect of wind blowing arthropods out of their habitat might deter them from

departing, as for instance found in ballooning spiders (Bonte et al., 2007). In our study, however,

individuals initiated movement within the local matrix after being placed there involuntarily. Release

experiments, like those we conducted with many species, or similar work on single species (Bonte et

al., 2004; Kallioniemi et al., 2014; Merckx & Van Dyck, 2007; Schultz et al., 2012), represent

movement behaviour at habitat boundaries. However, these experiments are less likely

representative of routine behaviours such as foraging within suitable habitat, or special dispersal

behaviours when crossing the matrix (Bonte et al., 2012; Clobert et al., 2009).

Surprisingly, temperature did not impact the movement speed of flying individuals. This can

potentially be attributed to the capacity of many flying arthropods to elevate their body temperature

through wing movement (Heinrich, 1974). Moreover, our dataset lacked extreme (hot or cold)

temperatures and we used an approximative measure for temperature, measured at the height of

the observer, not in the direct surroundings of the individuals. Both can potentially mask any trend

between temperature and movement of flying arthropods. In contrast, cursorial individuals

demonstrated the opposite pattern: temperature exerted a positive effect and wind did not. Given

that cursorial species remain in direct contact with the bare sand, they experience higher

temperatures compared to flying species due to the fast and more extreme heating of sandy

surfaces. However they do not have specific mechanisms in order to thermoregulate body

temperature (Maes et al., 2006; Mutiibwa et al., 2015) and it is thus expected that their movement is

more directly linked to the temperatures within their close environment. Our measurements reflect

this, on the condition that temperature measured at observer height is directly related to surface

temperature. Nevertheless, extremely high temperatures can potentially become detrimental when

a certain thermal threshold is reached, and consequently impede any movement (Heinrich, 1995).

However, we did not observe such conditions.

The movement direction of both flying and cursorial species exhibited a clear bias towards habitat

areas, indicating that the intention of moving towards suitable environments is a more important

driver for orientation than wind direction (Acevedo & Fletcher, 2017). This was supported by the

data from Kalmthout, where the movement direction should be relatively random due to the

proximity of suitable habitats in every direction. At this site, we indeed observed that movement was

more random and correlated with wind direction, indicating that the arthropods at De Panne actively

chose to move towards suitable habitat. Homing behaviour and site fidelity have been shown in

Hymenoptera, Lepidoptera and Blattodea, indicating that insects are indeed able to actively navigate

the landscape and determine where suitable areas are located (Boeddeker et al., 2015; Moura et al.,

2022; Rivault & Durier, 2004).

Although the direction of movement is clearly related to the location of habitat, movement direction

was not completely decoupled from wind conditions. For both flying and cursorial species, strong

upwind negatively impacted the returning rates towards the habitat patch, indicating the wind's

ability to divert individuals away from patch boundaries, even if they are able to navigate.

Nonetheless, the preference for a specific direction persisted despite wind resistance, implying that

arthropods are willing to endure, at least at small spatial scales, the energetic costs associated with

navigating towards hospitable patches. Similarly, tailwind assistance could minimize energy

expenditure during movement, which is in line with much larger-scale observations from migratory

species (Knight et al., 2019; Knoblauch et al., 2021). Only when winds were aligned with the habitat

direction, a clear-cut positive correlation between movement and wind direction was observed,

indicating that arthropods opportunistically harness winds to facilitate movement towards their

intended destinations. Tailwinds are known to drive large-scale migratory events (Åkesson, 2016),

but are thus equally important for small-scale movement behaviours at patch boundaries.

Larger flying arthropods moved more towards the habitat compared to smaller individuals and

species. A large body size provides advantages in handling upwind movement (Kuenen & Cardé,

1993) and is potentially related to a better-developed perceptual range (i.e. the distance at which an

individual can sense its suitable environment). For cursorial arthropods, only phylogenetically

uncorrected models showed that smaller taxa are more inclined to move towards suitable habitats.

This is most likely related to the abundance of taxa dominated by small species with more extensive

perceptual ranges in our study area. For instance, small-bodied arthropod taxa like spiders and ants

are known to mainly navigate based on social, chemical and/or tactical cues, which might increase

their perceptual range compared to species that solely rely on visual cues (Foelix, 2011).

The tendency of actively reaching a suitable habitat patch (often interpreted as success when costs of

entering the matrix are high) depends heavily on both body size and prevailing weather conditions.

Larger body sizes and the ability to fly provide advantages for arthropods navigating through the

matrix, enhancing movement efficiency. Moreover, ambient temperatures and wind speeds can

either augment or impede small-scale movement efficiency, depending on factors like wind direction

and thermal thresholds. Thus, movement behaviours at patch boundaries are condition dependent,

non-random (Goicolea et al., 2021), and reflect an active decision-making process (Bonte et al., 2012;

Clobert et al., 2009; Poethke et al., 2011).

Such non-random movements and dispersal patterns have the strong potential to impact

metapopulation dynamics and long-term persistence (Mortier et al., 2019). Therefore, in the context

of arthropod metacommunity dynamics, we expect spatial processes and environmental variation to

have a greater impact on communities of large-sized flying and smaller, cursorial species (De Bie et

al., 2012). Our study provides a mechanistic behavioural perspective on movement at habitat

boundaries, which can be integrated in predictive biodiversity models to better understand future

metacommunity processes (Urban et al., 2016). This is particularly important given the impending

effects of climate change, which are anticipated to alter body sizes (Daly et al., 2024; Davies, 2019;

Polidori et al., 2020), temperatures and wind patterns (Greene et al., 2010). This suggests that global

warming will not only impact population sizes by increasing stress on a local scale but that it may also

directly influence behaviours at habitat boundaries. Therefore, shifting weather patterns could

facilitate or constrain movement among habitat patches, directly affecting the composition and

future viability of arthropod metacommunities.

Declarations

Availability of data and materials

Data available from Zenodo https://doi.org/10.5281/zenodo.8232574 (Logghe et al., 2024).

Competing interests

The authors declare that they have no competing interests.

Funding

Garben Logghe was funded by FWO (grantnr: 1130223N). Charlotte Taelman was funded by BOF

(grantnr: BOF.24Y.2021.0012.01) and FWO (grantnr: 1SH0I24N).

Author's contributions

Garben Logghe, Charlotte Taelman, Florian Van Hecke and Dries Bonte designed the experiments.

Charlotte Taelman, Florian Van Hecke and Garben Logghe conducted the practical work. Garben

Logghe and Charlotte Taelman analysed the data and wrote the first draft of the manuscript. All

authors contributed substantially to the interpretation of the results and revision of the manuscript.

Acknowledgements

We would like to thank everyone who assisted with the fieldwork: Margaux Pottiez, Ruben Van De

Walle, Arthur Van Roey and Elias Van Den Broeck. Special thanks to Johan Lamaire (Agency for

Nature and Forests—Flemish Government) and Rudi Delvaux (Grenspark Kalmthoutse Heide) for

allowing us access to restricted areas in nature reserves De Westhoek (De Panne) and Kalmthoutse

Heide respectively. Finally, we would like to thank three anonymous reviewers for their valuable

comments on a previous version of the manuscript.

497 References

Acevedo, M. A., & Fletcher, R. J. (2017). The proximate causes of asymmetric movement across

heterogeneous landscapes. *Landscape Ecology*, *32*(6), 1285–1297.

https://doi.org/10.1007/s10980-017-0522-y

Agostinelli, C., & Lund, U. (2022). *R package "circular": Circular Statistics (version 0.4-95)* [Computer

software]. https://r-forge.r-project.org/projects/circular/

Aikman, D., & Hewitt, G. (1972). An Experimental Investigation of the Rate and Form of Dispersal in

Grasshoppers. *The Journal of Applied Ecology*, *9*(3), 807. https://doi.org/10.2307/2401906

Åkesson, S. (2016). Flying with the winds: Differential migration strategies in relation to winds in

moth and songbirds. *Journal of Animal Ecology*, *85*(1), 1–4. https://doi.org/10.1111/1365-

2656.12450

Arnold, P. A., Cassey, P., & White, C. R. (2017). Functional traits in red flour beetles: The dispersal

phenotype is associated with leg length but not body size nor metabolic rate. *Functional*

Ecology, *31*(3), 653–661. https://doi.org/10.1111/1365-2435.12772

Batsleer, F., Bonte, D., Dekeukeleire, D., Goossens, S., Poelmans, W., Van der Cruyssen, E., Maes, D.,

& Vandegehuchte, M. L. (2020). The neglected impact of tracking devices on terrestrial

arthropods. *Methods in Ecology and Evolution*, *11*(3), 350–361.

https://doi.org/10.1111/2041-210X.13356

Batsleer, F., Duez, F., Maes, D., & Bonte, D. (2023). *Strong gene flow across an urbanized coastal*

landscape in a dune specialist digger wasp [Preprint]. Ecology.

https://doi.org/10.1101/2023.04.15.537020

Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and Powerful

Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B*

(Methodological), *57*(1), 289–300. https://doi.org/10.1111/j.2517-6161.1995.tb02031.x

Boeddeker, N., Mertes, M., Dittmar, L., & Egelhaaf, M. (2015). Bumblebee Homing: The Fine

Structure of Head Turning Movements. *PLOS ONE*, *10*(9), e0135020.

https://doi.org/10.1371/journal.pone.0135020

Bonte, D., Bossuyt, B., & Lens, L. (2007). Aerial dispersal plasticity under different wind velocities in a

salt marsh wolf spider. *Behavioral Ecology*, *18*(2), 438–443.

https://doi.org/10.1093/beheco/arl103

527 Bonte, D., Lens, L., & Maelfait, J. (2004). Lack of homeward orientation and increased mobility result

528   in high emigration rates from low-quality fragments in a dune wolf spider. *Journal of Animal*

529   *Ecology*, *73*(4), 643–650. https://doi.org/10.1111/j.0021-8790.2004.00838.x

530 Bonte, D., Van Dyck, H., Bullock, J. M., Coulon, A., Delgado, M., Gibbs, M., Lehouck, V., Matthysen, E.,

531   Mustin, K., Saastamoinen, M., Schtickzelle, N., Stevens, V. M., Vandewoestijne, S., Baguette,

532   M., Barton, K., Benton, T. G., Chaput-Bardy, A., Clobert, J., Dytham, C., … Travis, J. M. J.

533   (2012). Costs of dispersal. *Biological Reviews*, *87*(2), 290–312.

534   https://doi.org/10.1111/j.1469-185X.2011.00201.x

535 Bowler, D. E., & Benton, T. G. (2005). Causes and consequences of animal dispersal strategies:

536   Relating individual behaviour to spatial dynamics. *Biological Reviews*, *80*(2), 205–225.

537   https://doi.org/10.1017/S1464793104006645

538 Buehlmann, C., Mangan, M., & Graham, P. (2020). Multimodal interactions in insect navigation.

539   *Animal Cognition*, *23*(6), 1129–1141. https://doi.org/10.1007/s10071-020-01383-2

540 Bürkner, P.-C. (2021). *Bayesian Item Response Modeling in R with brms and Stan. 100*(5), 1–54.

541   https://doi.org/doi:10.18637/jss.v100.i05

542 Chown, S. L., & Nicolson, S. W. (2004). *Insect phyisological ecology: Mechanisms and patterns*. Oxford

543   University Press.

544 Clark, J. A., & May, R. M. (2002). *Taxonomic bias in conservation research. 297*(5579), 191–192.

545 Clobert, J., Le Galliard, J.-F., Cote, J., Meylan, S., & Massot, M. (2009). Informed dispersal,

546   heterogeneity in animal dispersal syndromes and the dynamics of spatially structured

547   populations. *Ecology Letters*, *12*(3), 197–209. https://doi.org/10.1111/j.1461-

548   0248.2008.01267.x

549 Daly, K. M., Sikes, D. S., Mann, D. H., & Breed, G. A. (2024). Wing lengths of three Arctic butterfly

550   species decrease as summers warm in Alaska. *Ecography*, e07075.

551   https://doi.org/10.1111/ecog.07075

Davies, W. J. (2019). Multiple temperature effects on phenology and body size in wild butterflies

predict a complex response to climate change. *Ecology*, *100*(4).

https://doi.org/10.1002/ecy.2612

De Bie, T., De Meester, L., Brendonck, L., Martens, K., Goddeeris, B., Ercken, D., Hampel, H., Denys, L.,

Vanhecke, L., Van der Gucht, K., Van Wichelen, J., Vyverman, W., & Declerck, S. A. J. (2012).

Body size and dispersal mode as key traits determining metacommunity structure of aquatic

organisms. *Ecology Letters*, *15*(7), 740–747. https://doi.org/10.1111/j.1461-

0248.2012.01794.x

DiLeo, M. F., Nonaka, E., Husby, A., & Saastamoinen, M. (2022). Effects of environment and genotype

on dispersal differ across departure, transfer and settlement in a butterfly metapopulation.

Proceedings of the Royal Society B: Biological Sciences, *289*(1976), 20220322.

https://doi.org/10.1098/rspb.2022.0322

Driscoll, D. A., Banks, S. C., Barton, P. S., Lindenmayer, D. B., & Smith, A. L. (2013). Conceptual domain

of the matrix in fragmented landscapes. *Trends in Ecology & Evolution*, *28*(10), 605–613.

https://doi.org/10.1016/j.tree.2013.06.010

Dudley, R. (2000). *The biomechanics of insect flight—Form, function, evolution*. Princeton University

Press.

Eguagie, W. E. (1974). An Analysis of Movement of Adult Tingis ampliata H.-S. (Heteroptera: Tingidae)

in a Natural Habitat. *The Journal of Animal Ecology*, *43*(2), 521. https://doi.org/10.2307/3381

Felsenstein, J. (1985). Phylogenies and the Comparative Method. *The American Naturalist*, *125*(1), 1–

15. https://doi.org/10.1086/284325

Foelix, R. F. (2011). *Biology of Spiders* (3rd ed.). Oxford University Press.

Foellmer, M. W., Marson, M., & Moya-Laraño, J. (2011). *Running performance as a function of body*

size, leg length, and angle of incline in male orb-web spiders, Argiope aurantia. *13*, 513–526.

Forman, R. T. (2014). *Land mosaics: The ecology of landscapes and regions*. Cambridge University

Press.

Frick, W. F., Chilson, P. B., Fuller, N. W., Bridge, E. S., & Kunz, T. H. (2013). Aeroecology. In R. A.

Adams & S. C. Pedersen (Eds.), *Bat Evolution, Ecology, and Conservation* (pp. 149–167).

Springer New York. https://doi.org/10.1007/978-1-4614-7397-8_8

Frizzi, F. (2018). Complexity of searching movement in the European harvester ant Messor wasmanni:

Effect of temperature and body size. *Insectes Sociaux*, *65*(2), 263–273.

https://doi.org/10.1007/s00040-018-0609-8

Ganihar, S. R. (1997). Biomass estimates of terrestrial arthropods based on body length. *Journal of*

Biosciences, *22*(2), 219–224. https://doi.org/10.1007/BF02704734

Getz, W. M., & Saltz, D. (2008). A framework for generating and analyzing movement paths on

ecological landscapes. *Proceedings of the National Academy of Sciences*, *105*(49), 19066–

19071. https://doi.org/10.1073/pnas.0801732105

Gilbert, F., Gonzalez, A., & Evans-Freke, I. (1998). Corridors maintain species richness in the

fragmented landscapes of a microecosystem. *Proceedings of the Royal Society of London.*

Series B: Biological Sciences, *265*(1396), 577–582. https://doi.org/10.1098/rspb.1998.0333

Goicolea, T., Gastón, A., Cisneros-Araujo, P., García-Viñas, J. I., & Mateo-Sánchez, M. C. (2021).

Deterministic, random, or in between? Inferring the randomness level of wildlife

movements. *Movement Ecology*, *9*(1), 33. https://doi.org/10.1186/s40462-021-00273-7

Goodwin, B. J., & Fahrig, L. (2002). Effect of landscape structure on the movement behaviour of a

specialized goldenrod beetle, *Trirhabda borealis*. *Canadian Journal of Zoology*, *80*(1), 24–35.

https://doi.org/10.1139/z01-196

Grafen, A. (1989). *The phylogenetic regression*. *326*, 119–157.

Greene, S., Morrissey, M., & Johnson, S. E. (2010). Wind Climatology, Climate Change, and Wind

Energy: Wind climatology and wind energy. *Geography Compass*, *4*(11), 1592–1605.

https://doi.org/10.1111/j.1749-8198.2010.00396.x

Hein, S., Gombert, J., Hovestadt, T., & Poethke, H.-J. (2003). Movement patterns of the bush cricket

Platycleis albopunctata in different types of habitat: Matrix is not always matrix: Movement

patterns of P. albopunctata. *Ecological Entomology*, *28*(4), 432–438.

https://doi.org/10.1046/j.1365-2311.2003.00531.x

Heinrich, B. (1974). Thermoregulation in Endothermic Insects: Body temperature is closely attuned to

activity and energy supplies. *Science*, *185*(4153), 747–756.

https://doi.org/10.1126/science.185.4153.747

Heinrich, B. (1995). *Insect thermoregulation*. *19*(1), 28–33. https://doi.org/10.1016/0160-

9327(95)98891-I

Hillaert, J., Hovestadt, T., Vandegehuchte, M. L., & Bonte, D. (2018). Size-dependent movement

explains why bigger is better in fragmented landscapes. *Ecology and Evolution*, *8*(22), 10754–

10767. https://doi.org/10.1002/ece3.4524

Hirt, M. R., Lauermann, T., Brose, U., Noldus, L. P. J. J., & Dell, A. I. (2017). The little things that run: A

general scaling of invertebrate exploratory speed with body mass. *Ecology*, *98*(11), 2751–

2757. https://doi.org/10.1002/ecy.2006

Hurlbert, A. H., Ballantyne, F., & Powell, S. (2008). Shaking a leg and hot to trot: The effects of body

size and temperature on running speed in ants. *Ecological Entomology*, *33*(1), 144–154.

https://doi.org/10.1111/j.1365-2311.2007.00962.x

Jammalamadaka, S. R., & Sarma, Y. R. (1988). A correlation coefficient for angular variables. In

Statistical Theory and Data Analysis II (pp. 349–363). Elsevier Science Publishers B. V.

Jeltsch, F., Bonte, D., Pe'er, G., Reineking, B., Leimgruber, P., Balkenhol, N., Schröder, B., Buchmann,

C. M., Mueller, T., Blaum, N., Zurell, D., Böhning-Gaese, K., Wiegand, T., Eccard, J. A., Hofer,

H., Reeg, J., Eggers, U., & Bauer, S. (2013). Integrating movement ecology with biodiversity

research—Exploring new avenues to address spatiotemporal biodiversity dynamics.

Movement Ecology, *1*(1), 6. https://doi.org/10.1186/2051-3933-1-6

Jonsen, I. D., Bourchier, R. S., & Roland, J. (2001). The influence of matrix habitat on Aphthona flea

beetle immigration to leafy spurge patches. *Oecologia*, *127*(2), 287–294.

https://doi.org/10.1007/s004420000589

Kallioniemi, E., Zannese, A., Tinker, J. E., & Franco, A. M. A. (2014). Inter- and intra-specific

631   differences in butterfly behaviour at boundaries. *Insect Conservation and Diversity*, *7*(3),

632   232–240. https://doi.org/10.1111/icad.12046

633   Kaspari, M., & Weiser, M. D. (1999). The size–grain hypothesis and interspecific scaling in ants.

634   *Functional Ecology*, *13*(4), 530–538. https://doi.org/10.1046/j.1365-2435.1999.00343.x

635   Kissling, W. D., Pattemore, D. E., & Hagen, M. (2014). Challenges and prospects in the telemetry of

636   insects: Insect telemetry. *Biological Reviews*, *89*(3), 511–530.

637   https://doi.org/10.1111/brv.12065

638   Knight, S. M., Pitman, G. M., Flockhart, D. T. T., & Norris, D. R. (2019). Radio-tracking reveals how

639   wind and temperature influence the pace of daytime insect migration. *Biology Letters*, *15*(7),

640   20190327. https://doi.org/10.1098/rsbl.2019.0327

641   Knoblauch, A., Thoma, M., & Menz, M. H. M. (2021). Autumn southward migration of dragonflies

642   along the Baltic coast and the influence of weather on flight behaviour. *Animal Behaviour*,

643   *176*, 99–109. https://doi.org/10.1016/j.anbehav.2021.04.003

644   Kuenen, L. P. S., & Cardé, R. T. (1993). *Effects of moth size on velocity and steering during upwind*

645   *flight toward a sex pheromone source by Lymantria dispar (Lepidoptera: Lymantriidae). 6*,

646   177–193.

647   Kuussaari, M., Saarinen, M., Korpela, E., Pöyry, J., & Hyvönen, T. (2014). Higher mobility of butterflies

648   than moths connected to habitat suitability and body size in a release experiment. *Ecology*

649   *and Evolution*, *4*(19), 3800–3811. https://doi.org/10.1002/ece3.1187

650   Logghe, G., Taelman, C., Van Hecke, F., Batsleer, F., Maes, D., & Bonte, D. (2024). U*nravelling*

651   *arthropod movement in natural landscapes: small-scale effects of body size and weather*

652   *conditions.* [dataset on Zenodo]. https://doi.org/10.5281/zenodo.8232574

653   Lima, S. L., & Zollner, P. A. (1996). Towards a behavioral ecology of ecological landscapes. *Trends in*

654   *Ecology & Evolution*, *11*(3), 131–135. https://doi.org/10.1016/0169-5347(96)81094-9

Maes, D., Ghesquiere, A., Logie, M., & Bonte, D. (2006). Habitat Use and Mobility of Two Threatened

Coastal Dune Insects: Implications for Conservation. *Journal of Insect Conservation*, *10*(2),

105–115. https://doi.org/10.1007/s10841-006-6287-2

Martin, C. A., Proulx, R., & Magnan, P. (2014). The biogeography of insects' length-dry mass

relationships. *Insect Conservation and Diversity*, *7*(5), 413–419.

https://doi.org/10.1111/icad.12063

Matthew, P., Eastman, J. M., Slater, G. J., Brown, J. W., Uyeda, J. C., FitzJohn, R. G., Alfaro, M. E., &

Harmon, L. J. (2014). *geiger v2.0: An expanded suite of methods for fitting macroevolutionary*

models to phylogenetic trees. *30*, 2216–2218.

McIntyre, N. E., & Wiens, J. A. (1999). *Interactions between landscape structure and animal behavior:*

The roles of heterogeneously distributed resources and food deprivation on movement

patterns. *14*, 437–447.

Merckx, T., Van Dongen, S., Matthysen, E., & Van Dyck, H. (2008). Thermal flight budget of a

woodland butterfly in woodland versus agricultural landscapes: An experimental assessment.

Basic and Applied Ecology, *9*(4), 433–442. https://doi.org/10.1016/j.baae.2007.03.009

Merckx, T., & Van Dyck, H. (2007). Habitat fragmentation affects habitat-finding ability of the

speckled wood butterfly, Pararge aegeria L. *Animal Behaviour*, *74*(4), 1029–1037.

https://doi.org/10.1016/j.anbehav.2006.12.020

Michonneau, F., Brown, J. W., & Winter, D. J. (2016). *rotl: An R package to interact with the Open*

Tree of Life data. *7*(12), 1476–1481. https://doi.org/doi:10.1111/2041-210X.12593

Mola, J. M., Miller, M. R., O'Rourke, S. M., & Williams, N. M. (2020). Forests do not limit bumble bee

foraging movements in a montane meadow complex. *Ecological Entomology*, *45*(5), 955–965.

https://doi.org/10.1111/een.12868

Moretti, M., Dias, A. T. C., Bello, F., Altermatt, F., Chown, S. L., Azcárate, F. M., Bell, J. R., Fournier, B.,

Hedde, M., Hortal, J., Ibanez, S., Öckinger, E., Sousa, J. P., Ellers, J., & Berg, M. P. (2017).

680   Handbook of protocols for standardized measurement of terrestrial invertebrate functional

681   traits. *Functional Ecology*, *31*(3), 558–567. https://doi.org/10.1111/1365-2435.12776

682 Mortier, F., Jacob, S., Vandegehuchte, M. L., & Bonte, D. (2019). Habitat choice stabilizes

683   metapopulation dynamics by enabling ecological specialization. *Oikos*, *128*(4), 529–539.

684   https://doi.org/10.1111/oik.05885

685 Moura, P. A., Corso, G., Montgomery, S. H., & Cardoso, M. Z. (2022). *True site fidelity in pollen-*

686   *feeding butterflies. 36*, 572–582.

687 Mutiibwa, D., Strachan, S., & Albright, T. (2015). Land Surface Temperature and Surface Air

688   Temperature in Complex Terrain. *IEEE Journal of Selected Topics in Applied Earth*

689   *Observations and Remote Sensing*, *8*(10), 4762–4774.

690   https://doi.org/10.1109/JSTARS.2015.2468594

691 Neuwirth, E. (2022). *RColorBrewer: ColorBrewer Palettes. R package version 1.1-3* [Computer

692   software]. https://CRAN.R-project.org/package=RColorBrewer

693 Paradis, E., & Schliep, K. (2019). *ape 5.0: An environment for modern phylogenetics and evolutionary*

694   *analyses in R. 35*, 526–528.

695 Peterman, W. E., Connette, G. M., Semlitsch, R. D., & Eggert, L. S. (2014). *Ecological resistance*

696   *surfaces predict fine-scale genetic differentiation in a terrestrial woodland salamander.*

697   *23*(10), 2402–2413. https://doi.org/10.1111/mec.12747

698 Pewsey, A., Neuhäuser, M., & Ruxton, G. D. (2013). *Circular Statistics in R*. Oxford University Press.

699 Poethke, H. J., Gros, A., & Hovestadt, T. (2011). The ability of individuals to assess population density

700   influences the evolution of emigration propensity and dispersal distance. *Journal of*

701   *Theoretical Biology*, *282*(1), 93–99. https://doi.org/10.1016/j.jtbi.2011.05.012

702 Polidori, C., Gutiérrez-Cánovas, C., Sánchez, E., Tormos, J., Castro, L., & Sánchez-Fernández, D. (2020).

703   Climate change-driven body size shrinking in a social wasp. *Ecological Entomology*, *45*(1),

704   130–141. https://doi.org/10.1111/een.12781

Qiu, Y. (2023). *Showtext: Using fonts more easily in R graphs. R package version 0.9-6* [Computer

software]. https://CRAN.R-project.org/package=showtext

R Core Team. (2021). *R: A language and environment for statistical computing* [Computer software].

https://www.r-project.org/

Revell, L. J. (2012). *phytools: An R package for phylogenetic comparative biology (and other things). 3*,

217–223. https://doi.org/doi:10.1111/j.2041-210X.2011.00169.x

Ricketts, T. H. (2001). The Matrix Matters: Effective Isolation in Fragmented Landscapes. *The*

American Naturalist, 158(1), 87–99. https://doi.org/10.1086/320863

Rivault, C., & Durier, V. (2004). *Homing in German cockroaches, Blatella germanica (L.) (Insecta:*

Dictyoptera): Multi-channeled orientation cues. 110, 761–777.

https://doi.org/ttps://doi.org/10.1111/j.1439-0310.2004.01018.x

Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years of image

analysis. *Nature Methods, 9*(7), 671–675. https://doi.org/10.1038/nmeth.2089

Schtickzelle, N., & Baguette, M. (2003). Behavioural responses to habitat patch boundaries restrict

dispersal and generate emigration–patch area relationships in fragmented landscapes.

Journal of Animal Ecology, 72(4), 533–545. https://doi.org/10.1046/j.1365-

2656.2003.00723.x

Schtickzelle, N., Mennechez, G., & Baguette, M. (2006). *Dispersal depression with habitat*

fragmentation in the bog fritillary butterfly. 87(4), 1057–1065. https://doi.org/10.1890/0012-

9658(2006)87[1057:DDWHFI]2.0.CO;2

Schultz, C. B., Franco, A. M. A., & Crone, E. E. (2012). Response of butterflies to structural and

resource boundaries. *Journal of Animal Ecology, 81*(3), 724–734.

https://doi.org/10.1111/j.1365-2656.2011.01947.x

Sekar, S. (2012). A meta-analysis of the traits affecting dispersal ability in butterflies: Can wingspan

be used as a proxy?: Factors affecting dispersal ability in butterflies. *Journal of Animal*

Ecology, 81(1), 174–184. https://doi.org/10.1111/j.1365-2656.2011.01909.x

Shreeve, T. G., & Dennis, R. L. H. (2011). Landscape scale conservation: Resources, behaviour, the

matrix and opportunities. *Journal of Insect Conservation*, *15*(1–2), 179–188.

https://doi.org/10.1007/s10841-010-9336-9

Smith, A. T., Cook, D. R., Johnson, M. B., Townsend, V. R., & Proud, D. N. (2012). Comparative study

of walking and climbing speeds among Neotropical harvestmen from Costa Rica. *Journal of*

Arachnology, *40*(3), 304–308. https://doi.org/10.1636/Hi10-103.1

Terlau, J. F., Brose, U., Boy, T., Pawar, S., Pinsky, M., & Hirt, M. R. (2023). Predicting movement speed

of beetles from body size and temperature. *Movement Ecology*, *11*(1), 27.

https://doi.org/10.1186/s40462-023-00389-y

Turlure, C., Baguette, M., Stevens, V. M., & Maes, D. (2011). Species- and sex-specific adjustments of

movement behavior to landscape heterogeneity in butterflies. *Behavioral Ecology*, *22*(5),

967–975. https://doi.org/10.1093/beheco/arr077

Urban, M. C., Bocedi, G., Hendry, A. P., Mihoub, J.-B., Pe'er, G., Singer, A., Bridle, J. R., Crozier, L. G.,

De Meester, L., Godsoe, W., Gonzalez, A., Hellmann, J. J., Holt, R. D., Huth, A., Johst, K., Krug,

C. B., Leadley, P. W., Palmer, S. C. F., Pantel, J. H., … Travis, J. M. J. (2016). Improving the

forecast for biodiversity under climate change. *Science*, *353*(6304), aad8466.

https://doi.org/10.1126/science.aad8466

Van Dyck, H., & Baguette, M. (2005). Dispersal behaviour in fragmented landscapes: Routine or

special movements? *Basic and Applied Ecology*, *6*(6), 535–545.

https://doi.org/10.1016/j.baae.2005.03.005

Weil, S.-S., Gallien, L., Nicolaï, M. P. J., Lavergne, S., Börger, L., & Allen, W. L. (2023). Body size and life

history shape the historical biogeography of tetrapods. *Nature Ecology & Evolution*, *7*(9),

1467–1479. https://doi.org/10.1038/s41559-023-02150-5

White, E. P., Ernest, S. K. M., Kerkhoff, A. J., & Enquist, B. J. (2007). Relationships between body size

and abundance in ecology. *Trends in Ecology & Evolution*, *22*(6), 323–330.

https://doi.org/10.1016/j.tree.2007.03.007

Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis* [Computer software]. Springer-

Verlag.

Wickham, H., François, R., Henry, L., & Müller, K. (2022). *Dplyr: A Grammar of Data Manipulation. R*

package version 1.0.10 [Computer software]. https://CRAN.R-project.org/package=dplyr

Woodward, G., Ebenman, B., Emmerson, M., Montoya, J., Olesen, J., Valido, A., & Warren, P. (2005).

Body size in ecological networks. *Trends in Ecology & Evolution*, *20*(7), 402–409.

https://doi.org/10.1016/j.tree.2005.04.005

Wootton, R. (2020). Dragonfly flight: Morphology, performance and behaviour. *International Journal*

of Odonatology, *23*(1), 31–39. https://doi.org/10.1080/13887890.2019.1687991

Yang, L. H., & Gratton, C. (2014). Insects as drivers of ecosystem processes. *Current Opinion in Insect*

Science, *2*, 26–32. https://doi.org/10.1016/j.cois.2014.06.004

Zollner, P. A., & Lima, S. L. (1999). *Search strategies for landscape-level interpatch movements. 80*(3),

1019–1030. https://doi.org/10.1890/0012-9658(1999)080[1019:SSFLLI]2.0.CO;2