Multiscale mechanisms driving tissue rupture by invading cells.

Selwin K. Wu[1,2,12,14*], Fuqiang Sun[3,12], Celestine Z. Ho[1,4,13], Yuting Lou[1,13], Christina Bao-
Xian Huang[1], Mui Hoon Nai[4], Jingwei Xiao[1], Murat Shagirov[1], Jasmine Fei Li Chin[1], Diana
Lim[5], Suzie Verma[6], David SP Tan[7], Philippe Marcq[8], Alpha S. Yap[6], Chwee Teck Lim[1,4],
Tetsuya Hiraiwa[1,9], Yuan Lin[3,10*], Boon Chuan Low[1,2,11*]

[1]Mechanobiology Institute, National University of Singapore, Singapore

[2]Department of Biological Sciences, National University of Singapore, Singapore

[3]Department of Mechanical Engineering, University of Hong Kong, Hong Kong, China

[4]Department of Biomedical Engineering, National University of Singapore, Singapore

[5]Department of Pathology, National University Health System and National University of
Singapore, Singapore, Singapore

[6]Centre of Cell Biology and Chronic Disease, Institute for Molecular Bioscience, University
of Queensland, Brisbane, Australia

[7]Cancer Science Institute of Singapore & Department of Medical Oncology, National
University Cancer Institute Singapore, National University Health Systems, Singapore

[8]Physique et Mécanique des Milieux Hétérogènes, CNRS, ESPCI Paris, PSL University,
Sorbonne Université, Université de Paris, 75005, Paris, France

[9]Present address: Institute of Physics, Academia Sinica, Taipei, Taiwan

[10]Advanced Biomedical Instrumentation Centre, Hong Kong Science Park, Shatin, New
Territories, Hong Kong, China

[11]NUS College, National University of Singapore, Singapore

[12, 13]Contributed Equally

[14]Lead contact

*Correspondence: ylin@hku.hk & dbslowbc@nus.edu.sg &
selwin_wu@mail.dfci.harvard.edu

Summary

Cells migrate and invade tissues during development, immune responses, and cancer
metastasis. Generally, invading cells are thought to be the main drivers of this process,
pushing through mechanical barriers such as the extracellular matrix and tissue. However,
whether these mechanical barriers actively contribute to tissue invasion remains unclear.

Using tumour spheroids invading the mesothelium as an experimental model of tissue invasion with theoretical and computational models, we uncovered the interplay between mechanics and biological processes, ranging from molecular to multicellular scales. For the first time, we visualised strands of intercellular integrin connections between the invasive leader cell and the tissue barrier, which triggers the active constriction of the cells within the tissue barrier. This active constriction shrinks the adhesive contacts between cells of the tissue barrier, eventually causing them to rupture. Instead of the established models of invading cells pushing through the tissue barrier or jamming transition-driven cell invasion, we show that a coordinated patterning of subcellular contractility by the invading spheroid and the tissue barrier drives cell invasion and barrier rupture. Our data reveals a mechanically active role for the tissue barrier in collective invasion, thus providing a new model for understanding tissue invasion and morphogenesis.

Introduction

Cell migration and invasion into tissues and the extracellular matrix are fundamental for the development and maintenance of organisms and are rampant in diseases such as cancer[1, 2]. During cancer, tumour cells traverse into foreign tissues during metastasis[2], while in development, undifferentiated cells move into tissues to form different tissue types[3].

A variety of microenvironments contain the extracellular matrix and closely packed cells adherent to one another[4, 5]. Whether these mechanical barriers actively contribute to mammalian tissue invasion and whether cell-matrix adhesion receptors, such as integrins, are involved remains unclear. Since tumour cell clusters are more capable of forming secondary metastases than single cells[6]. Another key challenge is to identify the changes in subcellular dynamics that link molecular mechanisms to cellular and multicellular rearrangements in collective invasion of tissues.

Our understanding of how cells collectively invade tissues stems from the study of angiogenesis, neural crest migration and border cell migration in Drosophila. Landmark studies in Drosophila showed that the microtopography of barrier tissue provides an energetically favourable path of least resistance for border cell cluster migration, and cell-cell adhesion contributes to the traction of border cells[7]. Notably, mechanical feedback by E-cadherin-based Rac signalling promotes direction sensing of border cells through nurse cells[8].

However, a deep understanding of collective invasion into mammalian tissues is lacking. An

interdisciplinary approach to achieve a multiscale understanding[9, 10] of collective tissue

invasion in cancer remains largely unexplored.

Three-dimensional co-cultures mimicking ovarian tumour invasion of the mesothelium were

used as a model system in our experiments. Previously, it was suggested that integrin and

Talin-dependent activation of non-muscle myosin-II was required for spheroids to displace

the mesothelial cells apart during invasion[11, 12]. Here, we study the interplay between fracture

mechanics and cell invasion, ranging from molecular to multicellular scales. First, we have

identified a new phenomenon of tissue rupture by collective invasion and the patterns of

motion driving invasion. Then, we uncover the cellular and molecular mechanisms

underlying the patterns of motion driving invasion and the role of multicellular dynamics

during spheroid invasion. Finally, our theoretical modelling predicts the quantitative basis of

the invasion process. We conclude that instead of the widely accepted model of invading

cells pushing through the tissue barrier or jamming transition-mediated cell invasion, our data

reveals that a coordinated patterning of subcellular contractility by invading cells and the

tissue barrier drives tissue invasion.

Results

We study how cells migrate and invade the ECM and tissue barrier through experiments and

theory. Our experiments used three-dimensional co-cultures that mimic ovarian tumour

invasion of the mesothelium[11-14]. The mesothelium is a monolayer of simple squamous

epithelium[15] covering the human peritoneum's stroma (Fig. 1a). In peritoneal metastasis[16],

the mesothelium is cleared by ovarian tumours, eventually leading to the absence of the

mesothelium underneath the tumour and indentation of the stroma (Fig. 1b,b').

Our modified 3D culture systems can phenocopy our clinical observations (Fig. 1c). Human

ovarian adenocarcinoma OVCA433 epithelial cells were cultured to form spheroids in

suspension, bearing a grape cluster-like morphology (Fig. 1d)[17]. These spheroids were

introduced onto the ZT mesothelia[13] that form a laminin-rich basement membrane when

grown on Matrigel (Fig. 1c). The E-cad GFP expressing spheroid invaded and cleared the

underlying mesothelium (Fig. 1c), mimicking our clinical observations. Additionally,

multicellular assemblies migrating into tissues can also occur as a general morphogenetic

process that is not just specific to tumour invasion. This is because spheroids formed from

mesothelial cells can also migrate into mesothelium grown on Matrigel (Supplementary Fig.

1a).

Rupture initiation of the mesothelium during spheroid invasion

We first study tissue invasion at high resolution. As spheroids invade the mesothelium, a

phenomenon resembling crazing is observed at mesothelial cell-cell junctions (Fig. 1d' and

Video 1). Crazing is a unique feature when thermoplastic polymer materials rupture[18],

characterized by a network of fine cracks within the material. It occurs when excessive

tensile stress or stretching is applied to a polymer, producing gaps termed microvoids

between fibrillar bridges that are thought to help sustain some stress (Supplementary Fig. 1b).

At the mesothelium, we observed the pulling of cell-cell junctions leading to the elongation

of fibrillar bridges linked by the intercellular adhesion molecule, E-cadherin, and the

formation of microvoids before mesothelial rupture (Fig. 1d'' and Video 1). Mesothelial

rupture initiation was also characterised by the elongation of E-cadherin puncta, and we did

not observe any dramatic reduction of E-cadherin at intact regions of the mesothelial cell-cell

contacts during rupture initiation (Fig. 1d''', Video 1). These E-cadherin fibrillar bridges are

not specific to mesothelium cultured on glass, as we observe fibrillar bridges at sites of

spheroid invasion when mesothelium is grown on Matrigel (Video 2). Notably, fibrillar

bridges and microvoids in polymer crazing are ~10-20 nanometres, much smaller than the E-

cadherin fibrillar bridges and microvoids between cells we observed (Fig. 1d'' and Video 1).

Thus, instead, we termed the formation of E-cadherin fibrillar bridges and microvoids during

the initiation of cell-cell junction rupture as cell-cell contact crazing.

We weakened cell-cell contacts to test if these E-cadherin fibrillar bridges help resist rupture.

(Fig. 1d, Video 1 & 2). RhoA-Myosin-II mediated contractility supports E-cadherin cell-cell

adhesion strength[19-22]. We reduced E-cadherin adhesiveness of the mesothelium by

expressing a dominant negative RhoA mutant T19N fused to E-cadherin (Fig. 1e and

Supplementary Fig. 1c-f). Strikingly, the E-cadherin fibrillar bridges formed during spheroid

invasion were reduced in the mesothelium expressing E-cad-RhoA-T19N mutant (Fig. 1f,g).

Although the rate of mesothelial rupture remains unchanged in mesothelium expressing E-

cad-RhoA-T19N mutant (Supplementary Fig. 1g), more extensive mesothelial clearance by

spheroid invasion was observed (Fig. 1h). Thus, these mesothelial E-cadherin linked fibrillar

bridges may resist invasion. Taken together, we observe new features of tissue rupture during

cell invasion.

Tensile fracture of the mesothelium by spheroid engagement

Next, we identify the underlying patterns of motion and mode of mesothelium rupture

involved in invasion. To analyze the mesothelium fracturing process, we tagged its apical

surface with fluorescent microbeads[23] and imaged it undergoing spheroid invasion. 500nm

beads were coated with wheat-germ-agglutinin for cell surface binding[23].

A material's fracture response can be classified as brittle or more ductile according to the

extent of its deformation during fracture[24]. A ductile fracture is seen in readily deformable

materials and is characterized by extensive plastic deformation before the material finally

breaks apart[25]. By contrast, a brittle fracture exhibits little or no plastic deformation before its

fracture[25]. Velocity magnitude quantification from particle image velocimetry (PIV) of the

WGA-coated beads on the mesothelium apical surface revealed significant deformation of the

mesothelium (Fig. 1i, Supplementary Fig. 1h). This suggests that the tearing of the

mesothelium is akin to ductile rupture.

Generally, there are three principal modes by which fracture/rupture occurs, namely tensile

fracture[26, 27] (mode I), shear fracture (mode II) and out-of-plane tearing (mode III) (Fig. 1j).

Since cell-cell contact crazes typically form at the plane normal to tensile stress, spheroid

invasion should likely induce mesothelial tensile fracture.

Indeed, PIV of the beads on the mesothelium apical surface after spheroid invasion revealed a

constriction zone next to the region where the crack was initiated (Fig. 1i, Supplementary

Fig. 1h). After being pulled from both sides, the small crack opened a hole in the middle of

the mesothelium, an indication of a tensile fracture.

Given that cells did not slide past each other upon fracture, it is unlikely that in-plane shear is

an important driver of fracture initiation (Fig. 1d). To quantify the deformation nature of the

mesothelium, we examined both the normal strain rate and shear strain rate of the

mesothelium undergoing fracture (Fig. 1i,k, Supplementary Fig. 1h). Here, the normal strain

rate represents the strength of the velocity gradient in the direction of the local velocity, and

the shear strain rate represents the strength of the velocity gradient perpendicular to the local

velocity[28]. We did not observe a high shear rate near the initial fracturing site (Fig. 1i,k,

Supplementary Fig. 1h). Instead, we observed a low shear rate with high expansion velocity,

thus corresponding to tensile fracture (Fig. 1i,k, Supplementary Fig. 1h, Methods).

Finally, using lattice light sheet microscopy to achieve a clear cross-section view, we did not

observe any out-of-plane tearing (mode III) of the mesothelium caused by the spheroid (Fig.

1l). Protrusions penetrating the mesothelium E-cadherin junctions preceding mesothelial

fracture were only observed in ~6% of the invading spheroids (Supplementary Fig. 1i). While

in most of the spheroids, their protrusions were formed after mesothelial fracture. Indeed,

lattice light sheet imaging revealed that mesothelial cells were forced apart from the apical

side first, and such separation then propagated to the basal side, leading to a complete rupture

of the mesothelium. Thus, spheroid invasion induces tensile rupture, and we have ruled out

other modes of tissue rupture.

Spheroid leader cell attachment initiates apical constriction and tensile fracture of the

mesothelium

We then investigated the cellular mechanism underlying tension-induced mesothelial fracture

by spheroids. A large displacement at one side of the microvoid (Fig. 1i) may reflect an

apical constriction starting from a mesothelial cell beneath the spheroid as the mesothelium

begins to fracture (Fig. 2a). By contrast, such aggressive apical constriction is not detected in

other areas of the intact mesothelium away from the fracture site as shown by a more constant

cell apical area (Fig. 2b). It is conceivable that apical constriction of a cell could stretch its

junctions with neighbouring cells and eventually result in the aforementioned tensile fracture

(Fig. 1i). Apical constriction is a cell shape deformation mechanism, promoting tissue

remodelling during development[29-32], that is triggered by increased activation of myosin-II at

the apical cortex[21, 31, 33, 34]. Structured illumination super-resolution imaging of myosin-

regulatory-light-chain (MRLC)[35] revealed a myosin-II network extending across the apical

region of the mesothelial cells (Fig. 2c, c'). In quantitative live-imaging, MRLC intensifies at

the junctional and apical cortex region of a mesothelial cell beneath the interacting spheroid

(Fig. 2a,d and Video 3). Such intensification of MRLC precedes junction constriction and

fracture (Fig. 2d). These observations support the notion that constriction of apical

mesothelial myosin-II cortex pulls on E-cadherin junctions, eventually leading to initiation of

mesothelial fracture.

Besides the apical constriction mechanism, we use traction force microscopy (TFM) to

analyse the contribution of in-plane traction-mediate-migration of mesothelial cells during

fracture[23, 26]. First, we find that the spheroid leader cell tends to contract the

Polydimethylsiloxane (PDMS) substrate in the absence of mesothelium (Fig. 2e). Such

inward contraction from spheroid cells is well poised to activate the inward contraction/apical

constriction on contacting mesothelial cells.

Next, we introduced a spheroid on the mesothelium grown on PDMS but failed to detect an

enhancement in the mesothelial cell traction force on the PDMS substrate (Fig. 2f,g) or any

change in radial traction stress of mesothelial cells upon fracture initiation (Fig. 2f,

Supplementary Fig. 2a). Unexpectedly, in contrast to previous qualitative observations, we

did not observe traction-mediated lateral migration of mesothelial cells away from the

invading spheroid during fracture initiation[11]. Notably, our finding reinforces the conclusion

that mesothelial fracture ductility has absorbed and dissipated forces exerted by the spheroid

(Fig. 1j,k,l), preventing any obvious changes in the patterns of mesothelium traction forces.

Although obvious changes in the patterns of mesothelium traction forces are undetected,

Bayesian inversion stress microscopy analysis[36] reveals an increase in supracellular

mesothelial tensile stresses surrounding the spheroid before invasion (Fig. 2h,i,

Supplementary Fig. 2b). The increase in mesothelial tensile stresses be due to the

reorganisation of apical actomyosin filaments (Supplementary Fig. 2c-g). Taken together, this

reinforces that tensile stresses, likely due to apical constriction of the mesothelium, are

responsible for invasion-induced fracture.

Intercellular and apical integrins trigger mesothelial fracture and collective cell

invasion.

We then look deeper into the molecular mechanisms underlying the forces driving invasion.

We hypothesize that the spheroid-mesothelium interactions trigger mesothelial apical

constriction. It was shown that mesothelial clearance is initiated by ovarian tumour

attachment to the mesothelium via integrin alpha-5[11, 12], but the organization of intercellular

integrin-matrix complexes in a 3D microenvironment remains unclear. We imaged Paxillin -

a key component of the integrin matrix adhesion at the spheroid-mesothelium contact region

to visualise the intercellular integrin-matrix complexes. Strikingly, when they attached to the

mesothelium, spheroids expressing Paxillin-GFP exhibited strands of Paxillin at the spheroid-

mesothelium interface (Fig. 3a, Supplementary Fig. 2h), which grew from smaller initial

spots (Fig. 3b). The mesothelium tends to rupture after a spheroid's leader cell spreads on a

mesothelial cell (Supplementary Fig. 2i). These strands of Paxillin morphologically resemble

the integrin adhesion complexes seen when cells spread on glass[37]. We observed that

mesothelial rupture occurred after spreading of a spheroid's leader cell on the mesothelium in

~70% of invasion cases (Supplementary Fig. 2j).

As E-cadherin interactions between the epithelial assemblies[8, 19, 22, 33, 38-43] were undetected

with co-localization analysis between E-cad GFP spheroid and E-cad tdT mesothelium (Fig.

3c,d), we continued to analyse the role of intercellular integrin instead. Knocking down α5

integrin and using a thrombospondin-1 (TSP-1) peptide[44] to specifically block fibronectin-

based integrin adhesion of cells successfully blocked mesothelial fracture by the spheroid

(Fig. 3e-g, Supplementary Fig. 2j,k), validating the role of α5 integrin in the spheroid-

mesothelium attachment.

Adhesion receptors such as integrins could be present at the apical surface of mesothelial

cells, mediating their attachment to the spheroid. This is supported by our finding that

mesothelial cells detached from the adhesive (i.e. fibronectin-coated) round micropattern can

be associated with the spheroid at non-adhesive region of the substrate (Fig. 3h). Whereas in

controls, mesothelial cells remained attached on the adhesive round micropattern

(Supplementary Fig. 2l). Additionally, this observation was consistent with our TFM data,

together suggesting that mesothelial cell's outward migration away from the spheroid by

basal traction is dispensable for mesothelial fracture.

Strikingly, strands of Paxillin were localized at the apical side of mesothelial cells and

exhibited co-localization with immunostained fibronectin (Fig. 3i). Even in the absence of

spheroid attachment to the apical side of the mesothelium, α5 integrin and other binding

partners of the integrin complex, including Paxillin, kank2, tensin1, fibronectin and vinculin,

co-localized at sites of apically localized integrin-based adhesions (Fig. 3j). It was suggested

that integrin and Talin-dependent activation of non-muscle myosin-II was required for an

uncoupling mechanism to separate the mesothelial cells from fibronectin fibrils to drive

mesothelial cell migration away from the spheroid during invasion[11, 12]. However, instead of

such an uncoupling mechanism, the mesothelial apical fibronectin network undergoes

fracture as the spheroid invades (Video 4). Thus, our data shows that ECM and mesothelium

barrier fracturing mechanism is the dominant invasion mechanism.

Furthermore, mesothelial α5 integrin or mesothelial/spheroid fibronectin knockdown was

sufficient to block spheroid-induced fracture (Fig. 3k,l, Supplementary Fig. 2m-p and Video

5). Whereas fibronectin from the media is not required for invasion, as invasion remains

robust in serum-free media experimental condition (Fig. 3k, Supplementary Fig. 2n). Of note,

knockdown of α5 integrin primarily depleted apical integrins with a subtle reduction of basal

integrins (Supplementary Fig. 2o). Thus, the mesothelium's apical integrin adhesion complex

and spheroid fibronectin are required for spheroid invasion.

Our evidence suggests that mesothelial apical constriction from intercellular integrin

engagement initiates the mesothelial tensile fracture. To directly test this notion, we blocked

spheroid-induced mesothelial apical constriction by linking a dominant-negative RhoA

mutant to α5 integrin (Fig. 3m and Supplementary Fig. 2q-t). This significantly reduced

spheroid-induced mesothelial fractures (Fig. 3n,o, Supplementary Fig. 2u,v, and Video 5).

Although RhoA inhibition reduces mesothelial apical fibronectin fibril assembly

(Supplementary Fig. 2w,x), the restoration of fibronectin fibrils by introducing exogenous

fibronectin-1 fails to rescue the reduction of invasion caused by RhoA inhibition (Fig 3n,o,

Video 6). Thus, intercellular integrin engagement induces mesothelial fracture, in part by

apical myosin-II activation, is the key molecular mechanism driving invasion.

Spheroid invasion of the mesothelium did not involve induction of jamming transition

We then addressed how spheroid multicellular dynamics affect invasion. Invading spheroids

break through the mesothelium and then attach to the submesothelial substrate to prevent

anoikis of spheroid cells[45](Fig. 4a). Thus, we assess the critical relationship between spheroid

cell dynamics and mesothelial fracture. We quantitatively compared the cellular movement

pattern in (1) invading spheroids with (2) non-invading spheroids due to ITGA5 knockdown

in the mesothelium. We obtained tracking data of the spheroid's cell nuclei by deep-learning

segmentation[46, 47] and IMARIS tracking (Fig. 4b,c, Supplementary Fig. 3a). In invading

spheroids, there was a spreading of cells at the bottom layer, as well as an overall bias in cell

movement towards the mesothelium base even before mesothelial fracture (Fig. 4c,d, Video

7). By contrast, the cell movement in non-invading spheroids has no particular direction (Fig.

4d,e). We then assess whether directed cell motion is involved in spheroid invasion by

quantifying the persistency of cell velocity (Methods). Our results suggest that cells in non-

invading spheroids move less persistent than those in invading spheroids (Fig. 4f).

Jamming transition is a common trigger of collective cell migration, involving the transition

of multicellular assemblies from a jammed to fluid state[48, 49]. To quantify the involvement of

jamming transition, we then quantify the mean-square displacement (MSD) of cell nucleus

centroids obtained from AI segmentation and cell tracking. To exclude the effect of a global

drift of the spheroids, we adopted pair-wise MSD for our system (see Methods). We found

that at a small timescale of about 10 mins, both non-invading and invading spheroids show

roughly diffusive movement with MSD linearly growing with lag time, while on the

timescale larger than 50 minutes, MSD grows with lag time super-linearly for invading and

free-spreading spheroids and sub-linearly in non-invading spheroid (Fig. 4g,h). The short-

time diffusive movement of cells suggests that the spheroid is in a fluid state[36,37,38] even when

the spheroid cannot invade due to the lack of ITGA5 expression in the mesothelium. Our 3D

cell tracking and MSD analysis indicate that cancer cells are in an unjammed state even when

confined as a spheroid with intact E-cadherin adhesion. Thus, the triggering of jamming

transition is not a key driver for cancer spheroid invasion and spreading.

Instead, the whole spheroid could be considered as a multicellular droplet held by an

effective tensile surface. The long-time sub-linearity of MSD with non-invading spheroids

indicates the caging effect brought by the boundary of this effective surface, whereas, in

invading spheroids, the super-linearity of MSD on the larger timescale implies the existence

of some external force that guides cells in persistent and directed movement. Thus, this

external force likely originates from wetting at the intercellular integrin cell-cell contact

between the spheroid leader cell and the mesothelium. Indeed, α5 integrin and fibronectin are

required for the invasion. Moreover, the spreading of the spheroid leader cell coincides with

the shrinkage of the underlying mesothelial cell-cell junction preceding mesothelium fracture

(Fig 4i). Thus, spheroid invasion is independent of jamming transition induction.

Theoretical modelling of migration and tissue invasion

To elucidate the underlying physical mechanisms behind the observed experimental

phenomena, we developed a theoretical framework grounded in the concept of active wetting

(Methods). Specifically, we modelled the spreading and invasion of leader tumor cells as a

wetting-driven interfacial displacement process, in which the system's free energy is

determined by interfacial energy and polarization work (Eq. 1). The interfacial energy, in turn,

evolves dynamically according to interfacial adhesion and cortical tension, as governed by

Eqs. (2~4).

Our theory predicts the critical conditions under which tumor cell invasion occurs. A physical

process can only proceed spontaneously when the change in free energy is negative (Fig. 5a),

thereby providing a clear boundary between invasive and non-invasive regimes (Fig. 5b,c).

The theoretical analysis indicates that the critical condition for spontaneous invasion is

characterized by a 1/2 power law (Eq. 8) between the tumor-mesothelial integrin adhesion

strength (W_{I0}, Methods) and the activated contractility of mesothelial cells (described by the

coupling strength β, Methods). Moreover, due to the apical constriction of mesothelial cells,

the effective adhesion length between the two cell types is substantially reduced, which in

turn increases the required integrin-mediated adhesion strength for effective invasion. As

expected, the critical condition also reveals that polarization of the tumor cell toward the

mesothelial layer lowers the energy barrier and facilitates invasion, while an increase in the

thickness of the barrier cell layer impedes it (Fig. 5b,c).

To directly compare theoretical predictions with experimental results, we conducted

numerical simulations of the invasion process using the proposed theoretical framework in

conjunction with the active foam model [50](Supplementary Fig. 3b, Methods). Key simulation

parameters were informed by experimental observations and cortical tension measurements

by atomic force microscopy (AFM)(Supplementary Fig. 3c,d, Methods).

The simulations reproduced all key experimental observations, including the integrin-

mediated extension of the tumor-mesothelial interface, enhanced cortical tension due to

active mesothelial contraction, E-cadherin junction fracture between mesothelial cells in

contact with tumor cells, and internal cellular rearrangements within the invading tumor cell

(Fig. 5d,e, Video 8). The accuracy of our model was further validated by comparing

simulated and experimentally observed apical contraction areas and apical tension in

mesothelial cells (Fig. 5f,g). In particular, by tracking the changes in tumor cell shape

(quantified by the shape index I_s, a key descriptor of cell morphology and intercellular

junction lengths (Fig. 5h), our simulation revealed that tumor cells undergo prominent T1

transition during initiation of mesothelial invasion. This leads to super-diffusive motion of

the tumor mass, consistent with experimental observations (Fig.4d,e,g, Supplementary Fig.

3e,f).

Numerical modelling enabled us to quantitatively control the parameters of invasion; thus, we

seek to identify key factors affecting the invasion process. As a direct driver of invasion,

increased integrin adhesion promoted rapid internal rearrangement of tumor cells and rapid

rupture between mesothelial cells, collectively resulting in rapid tumor cell invasion

(Supplementary Fig. 3m-q; Video 9). Conversely, weakening integrin adhesion between cells

slowed down tumor rearrangement and ultimately abolished the tumor's ability to invade the

barrier layer (Supplementary Fig. 3h-l; Video 10). We did not observe an obvious effect of

reduced E-cadherin strength in mesothelial cells on collective invasion (Supplementary Fig.

4a-d, Video 11). Nonetheless, this reduction caused greater deformation of neighboring

mesothelial cells during invasion (Supplementary Fig. 3g, Supplementary Fig. 4e), which

may explain the experimentally observed increased mesothelial clearance upon E-cadherin

adhesion weakening (Fig. 1h).

Finally, we performed simulations to evaluate the impact of mesothelial cell contractility and

summarized the results alongside those involving intercellular adhesion strength in phase

diagrams (Fig. 5i,j,k, Supplementary Fig. 4f). Specifically, moderate enhancement of apical

contraction in barrier cells accelerated invasion, whereas overtly high contraction effectively

suppressed both tumor invasion and junctional rupture (Fig. 5i). Therefore, invasion occurs

only within a defined region of the phase diagram characterized by high integrin adhesion

strength and low-to-moderate barrier cell contractility. Moreover, consistent with apical

constriction without tissue fracture during embryonic development, our theory and numerical

simulations predict that when the intercellular integrin-mediated adhesion strength is weak

(or absent), apical constriction alone does not drive the rupture of the barrier cell monolayer

(Fig. 5b,c,i,j). The invasion–non-invasion boundary predicted by simulations aligned closely

with theoretical predictions (Fig. 5i,j), further confirming that tumor cell invasion is governed

by an interfacial displacement process driven by active tissue wetting. Similarly, due to the

competition between integrin adhesion–mediated wetting and mesothelial apical constriction–

driven dewetting, the spheroid cells exhibited morphological transitions between wetting and

dewetting states, as evidenced by a significant difference in shape index between the invasion
and non-invasion regions in the phase figures (Fig. 5k, Supplementary Fig. 4f).
408

Discussion
Our data supports an alternative model to the prevailing models of tumor cells pushing into
tissues. We uncover a violent mode of cell-cell contact fracturing. We observed that this
fracturing morphologically resembles 'crazing', a material science phenomenon originally
studied in the failure of thermoplastics[25]. Furthermore, 1. prevailing models indicate that
spheroid cell protrusion and dissociation of surface fibronectin fibrils from the mesothelium
forcefully drive mesothelial cell migration outward, achieving invasion[11]. 2. Additionally, the
spatiotemporal control of fluid-like and solid-like tissue states by jamming transition is a key
trigger for driving collective migration and invasion in a wide variety of models.
418

419 However, our 3D cell tracking and MSD analysis indicate that cancer cells are in an
420 unjammed state even when confined as a spheroid with intact E-cadherin adhesion. Thus, the
421 triggering of jamming transition is not a key driver for cancer spheroid invasion and
422 spreading.
423

By contrast, we present a new model of collective invasion, independent of jamming
transition induction. Our data demonstrate that RhoA-dependent mechanical forces are
transmitted across the spheroid integrins to the mesothelium integrins via the fibronectin
ECM sandwiched in between them (Fig 4a). In summary, intercellular integrin-mediated
adhesion between spheroids and mesothelial cells triggers and augments mesothelial cell
apical constriction. This drives the shrinkage and eventual tensile rupture of mesothelial cell-
cell junctions. Simultaneously, the growth of intercellular integrin adhesion induces striking
deformation and directional movement of spheroid cells, facilitating rupture of the
mesothelium. Importantly, our data reveal a mechanically active role for the tissue barrier in
collective invasion, thus providing a new model for understanding tissue invasion and
morphogenesis. Our findings of the tissue barrier playing an active role in invasion is
consistent with a previous report showing that the myoepithelial barrier cells restraining and
recapturing invasive breast cancer spheroid cells[51].
437

Additionally, the mesothelial apical integrin that we have identified here, although not

visualised previously, is reported to functionally mediate apical constriction in the embryonic

Drosophila epithelium [52, 53].

In our simulations, we use the theory of wetting to model the spheroid invasion of tissue. In

classical physics, wetting typically refers to the process by which a liquid displaces a gas to

maintain contact with a solid surface, governed by the relative strengths of interfacial

energies [54]. Contrary to this classical description, our study reveals that, beyond the passive

interfacial energy fluctuations induced by the formation and rupture of integrin- and E-

cadherin-mediated intercellular junctions, the active behavior of barrier cells, particularly

apical contraction, can actively modulate interfacial energy. This, in turn, regulates wetting

and plays a critical role in tumor cell invasion.

Specifically, we demonstrate that active apical contraction in barrier cells, in coordination

with intercellular adhesion (mediated by integrins and E-cadherins), can exert dual and

opposing effects on tumor invasion. Moderate enhancement of contractility accelerates the

invasion process, whereas excessive contractility significantly suppresses it. Our theoretical

framework attributes this inhibitory effect to a negative feedback mechanism in interfacial

energy competition: overactivated contractility in barrier cells, triggered by integrin

engagement, effectively restrains further growth of integrin-mediated adhesion, thereby

impeding invasion. This finding show that barrier cells actively participate in the invasion

process and play a critical regulatory role.

While active wetting behaviour at the cellular and multicellular level has been extensively

studied, no prior study has demonstrated that active wetting theory can predict interfacial

invasion dynamics between different cells. Our results reveal that cell cluster invasion can be

quantitatively predicted using the active wetting, driven interfacial displacement framework

that we have established. Moreover, because the interfacial energy in this model depends on

fluctuations in cellular mechanical properties, including cortical tension and adhesion

strength, the proposed framework is applicable to other multicellular reorganization systems

as well.

In conclusion, instead of the widely accepted model of invading cells pushing through the

tissue barrier or jamming transition-driven spheroid invasion, we show that a coordinated

patterning of subcellular contractility by the invading cells and the tissue barrier is the

dominant mechanism driving cell invasion and barrier rupture.

Limitations of the study

First, whether the model that we have identified the key mechanism in driving metastatic

colonisation in cancer patients remains to be determined. Second, consistent with the inward

contraction observed on the surface of mesothelium prior to invasion, we have shown that the

spheroid leader cell spreading induces inward contraction of the TFM PDMS substrate.

However, the TFM PDMS substrate is not necessarily reflecting the behaviour of the

mesothelial apical surface.

<u>Figures</u>

Figure 1. Spheroid-induced mesothelial tensile fracture.

(a,b) Ovarian tumour metastasis. Hematoxylin and eosin-stained human peritoneal surface

without metastatic tumour implantation (a) and a microscopic tumour implant on the omental

surface from a patient with metastatic serous ovarian carcinoma (b). Mesothelium (green

arrow, a) lines the abdominal surface but is not detected beneath the invading ovarian

tumour (b & b'). Representative images from 4 ovarian cancer patients with different layers

of tissue are indicated. Inset from b' with arrows indicating the invading tumour/stroma

interface without the mesothelium.

(c) An E-cadherin GFP-expressing spheroid invades and clears the mesothelium with a

basement membrane.

(d) Mesothelial cell-cell contact forms E-cadherin fibrillar bridges and microvoids upon

spheroid invasion. Representative time-lapse images of the mesothelium under a spheroid

(colour-coded in hot orange). d'-d''' with detailed views. d''' shows a cadherin fibrillar

bridge being extended as crazing proceeds. Red arrows point towards a fibrillar extension,

and green arrows point towards a microvoid.

(e) Schematic of E-cad-GFP-RhoA-T19N fusion protein.

(f,g) Reduction of E-cad fibrillar bridges during mesothelial fracture when E-cad-based

contractility is reduced. Representative images with arrows indicating fibrillar bridges (f) and

quantification (g) of the percentage of mesothelium E-cad fibrillar bridges when either RhoA-

T19N mutant is fused to E-cad or in controls.

(h) Increased size of mesothelial clearance when E-cadherin adhesion is compromised.

Quantitative comparison of the mesothelial clearance size when either RhoA-T19N mutant is

fused to mesothelial E-cad or in controls.

(i) Tension-induced fracture. PIV analysis of fluorescent microbeads on top of the

mesothelium expressing E-cad tdTomato. The heatmaps of shear strain rate (s^{-1}) or normal

strain rate (s^{-1}) or velocity(µm/s) from the PIV analysis. Right magnified view of the velocity

PIV heatmap shows outward vectors (red dotted circles) away from the fracture initiation site

(grey dotted circle).

(j) Diagram of three different modes of fracture which the mesothelium may undergo

(modified from modes of material fracture). The modes are Mode I: Opening (stress

orthogonal to the local plane of the crack surface), Mode II: In-plane-shear (stress parallel to

the crack surface but orthogonal to the crack front), and Mode III: Out-of-plane shear (stress

parallel to the crack surface and the crack front). The arrows indicate the direction of stress,

and asterisks indicate the crack point.

(k) Tension-induced fracture is the dominant initiating mode of fracture. Comparison

between the fold changes of the shear strain rate and normal strain rate with velocity

magnitude analysis during fracture initiation and fracture.

(l) Mode III: Out-of-plane shear fracture is not detected when the spheroid begins to invade

the mesothelium. XZ views of time-lapse lattice light-sheet microscopy images of E-cad GFP

spheroid invading E-cad tdT mesothelium. Grey arrows indicate the opening of mesothelial

cell-cell contacts.

All data are means ± SEM from at least n=2 independent experiments, except for (g), which

are from n=8 independent experiments, ns, not significant, *$p < 0.05$, **$p < 0.01$, ****$p <$

0.0001; analysed with unpaired t-test. Scale bars, 5µm.

Figure 2. Initiation of fracture from apical constriction of the mesothelial cell.

(a,b,c,d) Apical constriction precedes fracture initiation of the mesothelium. (a) Time-lapse

images of the spheroid (brightfield, top right) invading mesothelium expressing E-cad tdT

and MRLC-GFP. Intensification of MRLC filaments is observed as apical constriction of the

mesothelial cell led to subsequent fracture initiation of the mesothelium. (b) Variance

quantification of mesothelial apical cell area beneath spheroid during fracture compared to

areas not undergoing fracture. (c) Network of myosin-II filaments spans the apical region of

the mesothelium. Structured illumination microscopy of mesothelium expressing MRLC-
GFP, with a detailed view in (c') (d) Myosin-II filaments intensification precedes apical
constriction. Representative temporal changes in myosin-II filaments intensity at the apical
cortex and size of mesothelial cell apical area.

(e) Spheroid leader cell contracts basal substrate inwards. Time-lapse images of radial
traction stress of E-cad GFP expressing spheroid leader cell attaching onto the PDMS
substrate.

Heat maps in (e,f) shows the direction and magnitude of radial traction stress. The colour
scale bar indicates the magnitude of the outward (red) and inward (blue) radial traction stress
from the mesothelial fracture site.

(f,g) Traction force of mesothelial cells did not exhibit obvious changes upon fracture. Time-
lapse images (f) and quantification of fracture site average cellular traction stress (g) of E-cad
tdT expressing mesothelium when invaded by spheroid expressing E-cad GFP. The
discontinuous line in (f) indicates the area of the fracture.

(h,i) An increase in mesothelial tensile stresses at regions surrounding the invading spheroid
prior to invasion. Comparison of isotropic tensile stress at a 10-micron boundary (h, red)
surrounding invading spheroids prior to invasion and control which are spheroids that fail to
invade (i). For green regions highlighted in (h), refer to Supplementary Fig. 2b.

All data are means ± SEM from n=7 cells from at least 3 independent experiments, except for
(b), which is from 4 independent experiments, *p < 0.05, analysed with Mann Whitney U
test. Scale bars, 5µm.

Figure 3. Intercellular integrin adhesion and fibronectin matrix complexes connect the
spheroid to the mesothelium.

(a) Strands of integrin adhesion clusters at the spheroid-mesothelium interface from spheroid
expressing Paxillin GFP with indicated XZ representative views (bottom).

(b) The spheroid leader cell's Paxillin GFP clusters grow gradually at the spheroid-
mesothelium interface. The cell that is first attached to the mesothelium is defined as the
invasion leader cell. Time-lapse imaging of Paxillin GFP at the spheroid-mesothelium
interface.

(c) Cadherin at the interface between spheroid (top left, brightfield) expressing E-cad GFP
and mesothelium expressing E-cad tdTomato does not co-localize, with detailed views of c'
in c'' and c'''.

(d) Representative fluorescence line-scan analysis of spheroid E-cad GFP and mesothelium

E-cad tdTomato in c''' reveals a lack of homophilic E-cadherin interactions.

(e) Thrombospondin-1 peptide treatment, blocking integrin-based fibronectin attachment,

prevents spheroid-induced mesothelial fracture. Rate of spheroid-induced fracture in control

compared to thrombospondin-1 peptide treated cultures.

(f,g) Spheroid integrin knockdown blocks fracture. Representative images (f) and

quantification of the spheroid (g) invading the mesothelium in control compared to ITGA5

knockdown in the spheroid.

(h) Mesothelial cells associating with spheroids at non-adhesive regions of micropatterned

glass. Time-lapse images of spheroid displacing mesothelial doublets from round adhesive

fibronectin patterns into non-adhesive zones.

(i) Co-localization of fibronectin with apical mesothelial integrin clusters.

Immunofluorescence images of immunostained fibronectin and mesothelium expressing

Paxillin-GFP with detailed views (bottom).

(j) Mesothelial apical integrin adhesion complex without spheroid engagement.

Immunofluorescence images of mesothelium apical region immunostained for Paxillin,

ITGA5, FN1, Kank-2 and Vinculin.

(k,l) FN1 and ITGA5 from the mesothelium and spheroid are required for spheroid-induced

mesothelial fracture. Rate of spheroid-induced fracture in control compared to either

mesothelium FN1 knockdown, spheroid FN1 knockdown, serum-free media condition (k) or

ITGA5 knockdown (l).

(m) Design of ITGA5-GFP fused to RhoA dominant negative T19N mutant for expression in

the mesothelium.

(n,o) Spheroid-induced fracture requires activation of contractility at sites of mesothelial

ITGA5 adhesion. Representative images (n) and quantification (o) of spheroid-induced

mesothelial fracture rate in control mesothelium expressing ITGA5 knockdown with

reconstitution of ITGA5-GFP compared to the reconstitution of ITGA5-GFP fused to RhoA

dominant negative T19N mutant in the mesothelium.

All data are means ± SEM from at least with ~50 spheroid from n=3 independent

experiments, *$p < 0.05$, **$p < 0.01$; analysed with unpaired t-test. Scale bars, 5µm.

Figure 4. Jamming transition did not induce the spheroid invasion of the mesothelium.

(a) Model for spheroid-induced mesothelial tensile fracture

(b) Representative images of an OVCA433 spheroid labeled with H2B-mNeon segmented by

3D StarDist (left) and a 3D rendering of the segmented nuclei.

(c) Representative 2D projection of a 3D cell trajectories in spheroid invading the

mesothelium over time: (left) in XY axis, projected from the top, (middle) map in XZ axis

projected from the side and (right) map in Radial Z (RZ) axis, a radial projection of spheroid

assuming the rotational symmetry around the axis at the in-plane center of the spheroid.

(d) Spheroid cells exhibit motility bias towards the mesothelium prior to mesothelial fracture.

Representative XZ and RZ side view velocity maps of a spheroid 6 and 3 minutes before

mesothelial fracture compared to spheroids that will not induce fracture due to ITGA5

knockdown of the mesothelium.

(e) Mesothelium ITGA5 is required for spheroid cells motility bias towards the mesothelium.

The ratio of spheroid cells Z velocity mean over standard deviation of XY velocities

distribution of spheroid cells on control mesothelium compared to spheroid cells on ITGA5

knockdown mesothelium.

(f) Persistency of directed spheroid cell migration (see Methods) is higher in spheroid

invading the mesothelium compared to spheroid without invasion due to ITGA5 knockdown

in the mesothelium. Quantification of spheroid cells velocity auto-correlation persistency in

spheroid with or without invasion.

(g, h) Pair-wise mean square displacement (MSD) (g) (see Methods) of representative

samples for spheroid with or without invasion. (h) Local slope of MSD value obtained from

the pair-wise mean square displacement graph for six selected regions from top to base of

spheroids (see Methods) for two different conditions.

(i) Shrinkage of mesothelial cell-cell junctions and spreading of leader cell on mesothelium

precedes fracture. Representative temporal changes in leader cell spreading area at

mesothelial and sub-mesothelial regions is in green, and mesothelial junction length is in red.

Dotted line indicates the onset of mesothelial cell-cell contact rupture.

(j,k) Spheroids expressing higher levels of E-cadherin have reduced rate of invasion.

Representative E-cad GFP expressing spheroid images (j) and quantification of the spheroid

(k) invading the mesothelium in control compared to higher E-cadherin levels expressing

spheroid.

All data are means ± SEM from at least n=4 independent experiments, except for (n), which

are means ± SD from n=133 spheroid cells from 2 independent experiments, *p < 0.05, **p <

0.01, ****p < 0:0001; analysed with Mann-Whitney U test. Scale bars, 5µm.

Figure 5. Theoretical predictions and numerical simulations of wetting-induced invasion.

(a-c) Theoretical predictions for spontaneous fracture initiation and spheroid invasion. (a)

Schematics for the relationship between active wetting (increasing δL_{in}) and free energy

change, where a spontaneous invasion sets the condition as $\Delta G_{\text{m}} \leq 0$. (b) Increasing

mesothelial monolayer thickness (represented by increasing L_{m0}) significantly pushes the

critical invasion conditions toward higher intercellular integrin adhesion. (c) Spheroid

polarization promotes cell invasion and lowers the requirements of W_{I0} for complete fracture

initiation.

(d-h) Simulated invasion process agrees with experimental results when the intercellular

adhesion is comparable to the mesothelial E-cadherin adhesion, i.e. when $W_{\text{I0}}/W_0 = 0.9$,

$W_0/T_0 = 1.2, \beta/T_0 = 1.5$. (d) Intercellular integrin adhesion drives spheroid invasion and

mesothelial fracture. Stages of collective invasion include [i] initial state, [ii] establishment of

contact between the leader spheroid cell and the 2$^{\text{nd}}$ mesothelium cell, [iii] invasion, and [iv]

complete mesothelial fracture. To make a direct comparison with experimental results, we set

the initial time as -3 mins. (e) Complete fracture occurs ~10 mins after the spheroid leader

cell comes into contact with the 2$^{\text{nd}}$ mesothelial cell. The right interface refers to the interface

between the leader spheroid cell #1 and the #3 mesothelial cell, the left interface refers to the

interface between the leader spheroid cell #1 and the #2 mesothelial cell, and the fracturing

junction refers to the junction between #2 and #3 mesothelial cells labelled in (d). (f)

Simulated apical tension of the #2 mesothelial cell is in qualitative agreement with the

observed MRLC results. The computed tension is normalized by the static mesothelial

tension T_0. (g) Simulated apical area of the #2 mesothelial cell fits well with the

experimentally measured apical cell area. The apical area is calculated by the square of the

simulated apical cell length and is normalized by the initial area. (h) Spheroid cells show T1

transition during the invasion process. The shape index (individual and average) of the

spheroid cells is used to quantitively evaluate the rearrangement of the spheroid.

(i-k) Mesothelial fracture and spheroid rearrangement are strongly interrelated during

invasion. The simulated critical boundary for the occurrence of mesothelial fracture agrees

well with the prediction (black curves) given by Eq. (5). Complete mesothelial fracture and

spheroid rearrangement only occur at low to medium apical contractility and high integrin

adhesion. Low integrin adhesion strength or high apical contractility leads to lower spheroid

T1 transition and incomplete fracture. (i) Complete fracture time as a function of the integrin

adhesion strength and apical contractility. Rapid invasion occurs at medium apical

contractility and strong integrin adhesion strength. No data is at the bottom right corner since

no complete mesothelial fracture occurs in this region. The time taken for fracture is

calculated as the period between the contact of the leader spheroid cell with the 2nd

mesothelial cell and the complete rupture of the mesothelial junction. (j) High apical

contractility with low intercellular integrin adhesion strength leads to incomplete fracture, i.e.,

a non-vanishing minimal junction length (in the lower right region) between #2 and #3

mesothelial cells labeled in (d). The minimal fracturing junction length is normalized by its

initial length. (k) Spheroid cell rearrangement is regulated by intercellular integrin adhesion

and apical contractility. The shape index reflects the deformation of the leader spheroid cell

with respect to different intercellular integrin adhesion strength and apical contractility. Here,

the shape indices are taken when the fracturing junction length reaches 0 (for cases where

mesothelium is completely fractured) or when the fracturing junction length reaches its

minimum for cases where there is no complete mesothelial fracture.

Supplementary Figure 1. Spheroid cell protrusions were rarely observed during the initial

stages of invasion.

(a) Representative time-lapse images of mesothelial spheroid (brightfield, top right)

migrating into mesothelium expressing E-cad GFP grown on Matrigel. Invasion initiation

is observed at 15 mins.

(b) Schematic diagram of craze formation prior to thermoplastic fracture. Formation of

microvoids and fibrillar bridges precedes complete cracking of the material. Green arrows

indicate microvoids and red arrows indicate fibrillar bridges.

(c-e) E-cad GFP fused with RhoA-T19N reduces phosphoMLC intensity specifically at

mesothelial cell-cell junctions. Representative image (c) and quantification (d) of

mesothelium expressing E-cad GFP wild-type and E-cad GFP fused with RhoA-T19N,

immunostained for phospho-myosin light chain(red). E-cad GFP fused with RhoA-T19N

does not reduce medioapical phosphoMLC intensity levels. Representative images of

mesothelial cells expressing E-cad GFP wild-type and E-cad GFP fused with RhoA-T19N

immunostained for medioapical phospho-myosin light chain(green). White arrows point

towards medioapical phosphoMLC (e).

(f) Immunoblot of GFP, RhoA and α-Tubulin loading control of mesothelial cells expressing

E-cad-GFP and E-cad-GFP fused with RhoA-T19N. MW = apparent molecular weight.

(g) E-cad GFP fused with RhoA-T19N expressed in the mesothelium did not affect the

invasion rate. Spheroid's rate of invading mesothelium expressing either wild-type E-

cadherin or E-cadherin fused with RhoA-T19N.

(h) Control for Fig. 1i. PIV analysis of fluorescent microbeads on the apical region of E-cad

tdTomato–expressing mesothelium, in the absence of spheroid invasion. The heatmaps of

velocity (µm/s) from the PIV analysis show no significant variation among 0 minutes, 81

minutes, and 321 minutes.

(i) Spheroid cell protrusions are rare during and prior to fracture. Quantification of spheroid

protrusions before and after fracture.

All data are means ± SEM from n=8 independent experiments, ns, not significant; analysed

with unpaired t-test. Scale bars, 5µm.

Supplementary Figure 2. Intercellular integrin complex activation of mesothelial apical

constriction contributes to invasion.

(a) Mesothelial fracture induced by spheroid was not accompanied by changes in radial

traction stress. Fracture site radial traction stress quantification of E-cad tdT expressing

mesothelium when invaded by spheroid expressing E-cad GFP. Data are means ± SEM from

n=2 independent experiments, ns, not significant; analysed with unpaired t-test.

(b) No significant difference observed between average mesothelial cell heights at BISM

analysis region and control region where cells are further away from the spheroid. Green

regions highlighted in Fig. 2h indicate the control boundary region where mesothelial cell

heights were measured.

(c-d) Mesothelial MRLC filaments are reoriented parallel in the cell undergoing apical

constriction (marked by gray arrows, Figure 2a), relative to the site of mesothelial fracture.

Graphs indicating that apical MRLC filaments show an increase of parallel filament bundles

to the crack boundary (0°) at (d) 21 mins than that at (c) 9 mins.

(e) Mesothelial apical F-actin bundles are organised parallel to the crack boundary at the late

stage of invasion. The F-actin bundles near the crack boundary (marked by the yellow

rectangle) are aligned (f) along the direction of the crack boundary (0°, white arrow), while

fibers far from the crack boundary (marked by the red rectangle) are more uniformly

distributed (g).

(h) Additional representative image of XZ views of a spheroid cell's Paxillin-GFP clusters on

a mesothelium expressing E-cadherin tdTomato.

(i) Spreading of a spheroid leader cell on a mesothelial cell is sufficient for mesothelial

fracture. Quantifying the number of spheroid cells spreading on the mesothelium before

spheroid-induced mesothelial fracture.

(j) Blocking of integrin to fibronectin interactions with TSP-1 inhibits spheroid invasion.

Representative image of spheroid invading mesothelium in control compared to when TSP-1

is introduced.

(k) Immunoblot of α5 integrin knockdown in OVCA433 spheroids with α-Tubulin as loading

control. MW = apparent molecular weight.

(l) Mesothelial cells remained attached to the adhesive fibronectin round pattern in the

absence of invading spheroids. Time-lapse imaging of mesothelial cells expressing E-

cadherin tdTomato and cytoplasmic GFP on fibronectin round patterns.

(m) Immunoblot of fibronectin knockdown in mesothelium with α-Tubulin as loading

control. MW = apparent molecular weight.

(n) Knockdown of mesothelium fibronectin or spheroid fibronectin inhibits spheroid

invasion, whereas spheroid invasion persists in the absence of serum. Representative image

of spheroid invading mesothelium in control compared to when fibronectin-1 is knocked

down in the mesothelium or knocked down in the spheroid or serum-free condition.

(o) Knockdown of α5 integrin is most prominent at the apical region of mesothelium.

Representative images of mesothelium with α5 integrin knockdown at the apical and basal

regions, immunostained with α5 integrin and Paxillin.

(p) Knockdown of mesothelial α5 integrin inhibits spheroid invasion. Representative image

of spheroid invading mesothelium in control compared to when α5 integrin is knocked down

in the mesothelium.

(q,r) Immunoblot of GFP, RhoA and GAPDH loading control of mesothelial cells expressing

ITGA5-GFP and ITGA5-GFP fused with RhoA-T19N. MW = apparent molecular weight.

(s,t) ITGA5-GFP fused with RhoA-T19N reduces apical phosphoMLC. Mesothelial cells

expressing ITGA5-GFP and ITGA5-GFP fused with RhoA-T19N immunostained (s) and

quantified (t) for phosphoMLC fluorescence.

(u,v) Paxillin intensity per integrin adhesion is unaffected by ITGA5-GFP fusion with

RhoA-T19N. Representative images (u) and quantification (v) of mesothelium expressing

ITGA5-GFP and ITGA5-GFP fused with RhoA-T19N immunostained for medioapical

paxillin (green). White arrows point toward integrin adhesions.

(w,x) Addition of exogenous fibronectin to mesothelium restores the level of apical

fibronectin fibrils that are reduced by RhoA inhibition. Apical mesothelial FN1 of Control

ITGA5-GFP, ITGA5-GFP+RhoA T19N and ITGA5-GFP+RhoA T19N with exogenous

FN1 immunostaining(w) and their corresponding intensity quantification(x).

Supplementary Figure 3.

(a) Representative XY top view, XZ and RZ side views of spheroid cells tracks that did not

invade the mesothelium over time.

(b) Schematic of the model formulation described in Eq.5, M2 & M4, where the intercellular

vertex motion is driven by polarization force and the difference in cortical tension and

osmotic pressure.

(c,d) Atomic Force Microscopy analysis of spheroid and mesothelium for deriving the

tension measurements used in computational modelling.[i] Representative approach (blue) and

retraction (orange) force-indentation curves when a maximum force of 3nN is applied to the

mesothelial cell (c) and cancer spheroid cell (d).

(e,f) Spheroid cells rearrange and move collectively towards the mesothelium during invasion

in the simulation ($W_{I0}/W_0 = 0.9$). (e) Simulated collective spheroid cell motion towards the

direction of the mesothelium (negative Z direction). (f) Central spheroid junction shrinkage

indicates striking deformation and intercalation of spheroid cells.

(g) Longitudinal strain of the mesothelial cells ($W_{I0}/W_0 = 0.9$). Longitudinal strain is

computed by the dynamic change of cell length.

(h-l) Low integrin adhesion strength ($W_{I0}/W_0 = 0.3$) prevents spheroid T1 transition and

leads to incomplete mesothelial fracture. (h) Process of incomplete fracture includes [i] initial

state, [ii] contacting between the leader spheroid cell and the 2[nd] mesothelium cell, [iii]

further invasion, and [iv] complete mesothelial fracture nucleation. The initial time is set as -

3 mins. (i) Fracturing junction reaches its minimal length ~25 minutes after contacting the 2[nd]

mesothelial cell, indicating an incomplete fracture at low integrin adhesion strength. The right

interface refers to the interface between the leader spheroid cell (#1) and the #3 mesothelial

cell, the left interface refers to the interface between the leader spheroid cell (#1) and the #2

mesothelial cell, and the fracturing junction refers to the junction between #2 and #3

mesothelial cells. (j) Low integrin adhesion strength prevents T1 transition of spheroid cells

during invasion. (k) Low integrin adhesion strength results in slow central junction shrinkage

and thus prevents T1 transition in the spheroid. (l) Low integrin adhesion strength changes

the distribution of mesothelial strain in adjacent cells.

(m-q) High integrin adhesion strength ($W_{I0}/W_0 = 1.5$) results in striking spheroid T1

transition and mesothelial fracture. (m) Dynamic process of rapid fracture includes [i] initial

state, [ii] contacting between the leader spheroid cell and the 2[nd] mesothelium cell, [iii]

further invasion, and [iv] complete mesothelial fracture nucleation. The initial time is set as -

3 mins. (n) Complete fracture occurs ~6 mins after the leader cancer cell comes into contact

with the 2[nd] mesothelial cell. The right interface refers to the interface between the leader

spheroid cell #1 and the #3 mesothelial cell, the left interface refers to the interface between

the leader spheroid cell #1 and the #2 mesothelial cell, and the fracturing junction refers to

the junction between #2 and #3 mesothelial cells. (o) High intercellular integrin adhesion

strength leads to striking T1 transition of spheroid cells during invasion. (p) High integrin

adhesion strength results in quick central junction shrinkage and, thus, striking T1 transition

in the spheroid. (q) Slightly smaller mesothelial deformation was observed by increasing

intercellular integrin adhesion strength.

Supplementary Figure 4. Low mesothelial E-cadherin adhesion strength ($W_0/T_0 = 0.6$, static

mesothelial tension $0.5T_0$) results in large mesothelial deformation during invasion-induced

fracture.

(a) The fracture process under low mesothelial E-cadherin adhesion includes [i] initial state,

[ii] contacting between the leader spheroid cell and the 2[nd] mesothelium cell, [iii] further

invasion, and [iv] complete mesothelial junction rupture. The initial time is set as -3 mins.

(b) No significant change in fracturing time by lowering mesothelial cell-cell adhesion.

Complete fracture occurs ~10 mins after the leader cancer cell comes into contact with the 2[nd]

mesothelial cell. The right interface refers to the interface between the leader spheroid cell

(#1) and the #3 mesothelial cell, the left interface refers to the interface between the leader

spheroid cell (#1) and the #2 mesothelial cell, and the fracturing junction refers to the

junction between #2 and #3 mesothelial cells.

(c,d) Low mesothelial cell-cell adhesion does not significantly affect the T1 transition of

spheroid cells. (c) The spheroid cells have no significant change in the computed shape index

(for both the leader cell and the average of all cells) compared to the reference scenario

($W_0/T_0 = 1.2$, high mesothelial cell-cell adhesion). (d) No significant change in central

spheroid junction length in comparison to the reference scenario ($W_0/T_0 = 1.2$, high

mesothelial cell-cell adhesion).

(e) Higher longitudinal strain of the mesothelial cells induced by lower mesothelial cell-cell

adhesion. Longitudinal strain is computed by the dynamic change of cell length.

(f) Average shape index of cells representing spheroid T1 transition is influenced by both

intercellular integrin adhesion and apical contractility.

(g,h) Spheroids expressing higher levels of E-cadherin have reduced rate of invasion.

Immunoblot (g) and the corresponding quantification of (h) of GFP for E-cadherin-GFP and

α-Tubulin loading control of spheroid cells expressing a low (control) or high amount of E-

cadherin-GFP. MW = apparent molecular weight.

Supplementary Videos

Video 1. Formation of fibrillar bridges and microvoids at mesothelial E-cadherin (hot orange)

cell-cell contacts during spheroid invasion.

Video 2. Formation of fibrillar bridges (E-cad tdTomato, red) and microvoids at mesothelium

(GFP, green) grown on matrigel during spheroid invasion.

Video 3. Apical constriction of a mesothelial cell expressing myosin regulatory light chain

(MRLC-GFP, green) and E-cadherin tdTomato (red) during the spheroid invasion.

Video 4. Mesothelial apical fibronectin GFP and mesothelium (E-cad tdTomato, red) undergo

fracture induced by an invading spheroid.

Video 5. Representative multipanel videos of spheroid invasion of the mesothelium in the

three following experimental conditions. 1) Fibronectin-1 and 2) α5 integrin siRNA mediated

knockdown in mesothelium compared to controls. 3) Mesothelium expressing α5 integrin

GFP fused with RhoA-T19N compared to control mesothelium expressing wild-type α5

integrin GFP. Arrows indicate four ends of the mesothelial fracture propagation.

Video 6. Representative multipanel videos of spheroid invasion of the mesothelium in the

three following experimental conditions. 1) Control mesothelium expressing wild-type α5

integrin GFP, 2) mesothelium expressing α5 integrin GFP fused with RhoA-T19N, and 3)

mesothelium expressing α5 integrin GFP fused with RhoA-T19N with addition of exogenous

FN1. Arrows indicate four ends of the mesothelial fracture propagation.

Video 7. Representative multipanel video of spheroid expressing H2B-mNeon invading E-

cad tdT expressing mesothelium and the corresponding XZ and RZ velocity maps on the

right.

Video 8. Computational vertex results of spheroid cell invasion and mesothelial fracture with

medium intercellular integrin adhesion ($W_{I0}/W_0 = 0.9$).

Video 9. Computational vertex results of spheroid cell invasion and mesothelial fracture with

low intercellular integrin adhesion ($W_{I0}/W_0 = 0.3$).

Video 10. Computational vertex results of spheroid cell invasion and mesothelial fracture

with high intercellular integrin adhesion ($W_{I0}/W_0 = 1.5$).

Video 11. Computational vertex results of spheroid cell invasion and mesothelial fracture

with low mesothelial E-cadherin adhesion ($W_0/T_0 = 0.6$).

<u>Acknowledgments</u>

We gratefully thank Ernest Poon, Phoebe Lim, Daniella Lawther, Shiwei Beh, Lee Kang

Wei, Hui Ting Ong, Chaoyu Fu, Ti Weng Chew, Xingyu Xia, T Thivakar, Soumya Bunk,

Thuan Beng Saw, Jennifer Young, Erik Sahai, Robert Parton and Peter Friedl for their input

and discussions during this project. Ivan Yow for providing PDMS dishes for TFM and

helping with TFM setup and discussions. Raymond Woo and Samuel Ko for Zeiss lattice

light sheet microscopy setup and imaging support. Wei Jia Goh for discussions and

suggestions to use 3D StarDist. Tay Shu Chian for editing the manuscript. Diego Pitta de

Araujo for making the illustration diagrams. Ince Tan for providing the ZT-GFP cells.

Funding: This research was supported by a Young Individual Research Grant (MOH Grant

No: MOH-000653) by the National Medical Research Council of Singapore to S.K.W. A

Research Scholarship Block Research Fellow Scheme and MOE Tier 1 by the Singapore

Ministry of Education to S.K.W and B.C.L. Yuan Lin thanks the financial support from the

Research Grants Council (Project No.: GRF/17210618, GRF/17210520) of the Hong Kong

SAR Government and the National Natural Science Foundation of China (Project No. 12272332). We also thank the Singapore National Research Foundation NRF for the Mid-Sized Grant (NRF-MSG-2023-0001) to B.C.L. and Mechanobiology Institute's Seed Funds to B.C.L and T.H. A.S.Y was supported by the National Health and Medical Research Council of Australia (1136592 and 1163462); the Australian Research Council (DP220103951); the US Department of Defence (Discovery Grant HT94252310088). **Competing Interests:** Authors declare no competing interests.

Author Contributions

S.K.W conceived the project with inputs from Yuting Lou and A.S.Y. SKW, and C.Z.H. and performed all experiments except for traction force microscopy and immunoblotting, which were shared with J.X and C.B.H.; D.L and D.S.T. provided cancer pathology analysis. F.Q.S and Yuan Lin developed the theoretical model and conducted the simulations and quantification. T.H and Yuting Lou conducted spheroid jamming transition analysis; P.M. provided the BISM code and helped with the analysis, M.H Nai, Yuting Lou and C.T Lim contributed to the AFM tension analysis; C.B.H, I.Y and M.S contributed to traction force analysis and 3D StarDist A.I based nucleus segmentation. S.V and A.S.Y provided input on cadherin and myosin experiments. S.K.W and F.Q.S wrote the manuscript. B.C.L, C.B.H, Yuan Lin and Lou Yuting, A.S.Y edited the manuscript. J.C contributed to imaging setup. All analyses were performed with S.K.W assisted by C.Z.H. All authors feedback on the manuscript.

Declaration of interests

A.S.Y. is a member of the Advisory Board of Developmental Cell.

Methods

Cell culture OVCA433 ovarian cancer cell lines and the ZT mesothelial cells were cultured in a 1:1 ratio of medium MCDB 105 (Cell Applications Inc.) and Medium 199 (GIBCO) supplemented with 10% FBS (GIBCO). The OVCA433 cell lines were obtained from Michael Sheetz's laboratory. ZT mesothelial cells were obtained from a benign pleural effusion, a gift from Ince Tan's laboratory. These cells were immortalized by ectopic expression of SV40 T antigen and overexpression of human telomerase (fused to GFP), as previously described [13]. Non-GFP expressing ZT cells were obtained by sorting out non-

fluorescence ZT cells, and they were subsequently immortalized by transfecting them with
SV40 T-antigen expressing plasmid (Addgene).

Human tissue sections
Hematoxylin and eosin (H&E)-staining sections of human ovarian cancer sections were
obtained from archived patients in the National University Hospital, Singapore. All samples
were obtained with approval by Singapore National Healthcare Group's Domain Specific
Review Board under the reference number 2013/00705. A senior pathologist reviewed the
specimens to identify the ovarian tumour invading the peritoneum.

Heterotypic spheroid-monolayer coculture IWAKI 35mm dishes (12mm glass diameter)
were coated with 1:100 Human Fibronectin plasma (Sigma-Aldrich) in PBS and incubated at
4°C overnight. For Matrigel conditions, 20µl of Geltrex was added to the glass base and
evenly spread. Subsequently, ZT mesothelial cells were seeded and incubated overnight at
37°C to form a confluent monolayer.

Spheroids of OVCA433 were prepared by dissociating cells with trypsin-free Cell
Dissociation Buffer, PBS (Gibco), and incubated in suspension in 2ml of 1:1 ratio of medium
MCDB 105 (Cell Applications Inc.) and Medium 199 (Gibco) with 10% FBS (Gibco) for 30
mins twice at 37°C to remove trace amounts of cell dissociation buffer and promote spheroid
formation. Spheroids of approximately 50µm size were imaged and analysed. Spheroids were
transferred to a dish containing a confluent monolayer and incubated for 15 min to allow
spheroids to settle before live imaging. Imaging was performed using Nikon W1 Spinning
Disk confocal microscope with the integrated Perfect Focus System on a 60x water-
immersion magnification objective. Brightfield, GFP, and RFP images were captured every 3
min for 12 h. After live imaging, spheroids that invaded the monolayer were quantified by
viewing data obtained using MetaMorph Microscopy Automation and Image Analysis
Software. To quantify the mesothelial clearance area, the non-fluorescent area created by the
invading spheroid in the GFP mesothelial monolayer images was measured at 24 h and
divided by the initial area of the cancer spheroid 2D size at time 0. All measurements were
taken using ImageJ software.

Protein coupling of microbeads PolyLink Protein Coupling Kit (Polysciences, Inc) was
used to prepare 50µl of fluorescence beads solution. 200µg of Wheat Germ Agglutinin was

added to the activated beads suspension and incubated for 2 hr at room temperature.
Subsequently, the beads suspension was sonicated for 15 min and immediately added to a
dish containing a confluent monolayer and centrifuged until no floating beads could be
observed. Prepared OVCA433 spheroids were transferred to the dish and incubated for 15
min to allow spheroids to settle before live imaging.

Multicellular spheroids of OVCA433 were prepared by dissociating cells with trypsin-free
Cell Dissociation Buffer, PBS (Gibco) and incubated in suspension in 2ml of 1:1 ratio of
medium MCDB 105 (Cell Applications Inc.) and Medium 199 (Gibco) with 10% FBS
(Gibco) for 30 mins twice at 37°C to remove trace amounts of cell dissociation buffer and
promote spheroid formation. Spheroids were transferred to a dish containing a confluent
monolayer and incubated for 15 min to allow spheroids to settle before live imaging. Imaging
was performed using Nikon W1 Spinning Disk confocal microscope with the integrated
Perfect Focus System on the 60x water-immersion magnification objective. Brightfield, GFP,
and RFP images were captured every 3 min for 12 h. Spheroids that invaded the monolayer
were quantified with MetaMorph.

Plasmids, siRNA and shRNA For the expression of fluorescence transgenes we subcloned
shRNA-resistant transgene fused with the fluorescence tag into the multiple cloning site of
either lentiviral pLL3.7 or pLL5.0 vectors by recombination. The shRNA targeting the
corresponding endogenous proteins is cloned into the lentiviral pLL vectors between the
XhoI and HpaI restriction sites to prevent overexpression artefacts. The fluorescent
transgenes are designed to be shRNA resistant by using a non-human transgene or codon
change with site-directed mutagenesis. The lentiviral constructs generated are 1. pLL3.7
human Paxillin shRNA knockdown reconstituted with gallus gallus Paxillin GFP. pLL3.7
Paxillin shRNA/gallus gallus Paxillin-GFP, 2. pLL5.0 human ITGA5 shRNA knockdown
reconstituted with shRNA resistant human ITGA5-GFP -- pLL5.0 human ITGA5
shRNA/shRNA resistant human ITGA5-GFP. Additionally, a linker (5′ -
GGGTCTACATCTGGATCTGGGAAGCCGGGTTCTGGTGAGGGTTCT-3′) is fused
between fluorescence transgenes and the RhoA dominant negative mutants for the following
constructs. 3. pLL5.0 human ITGA5 shRNA knock down reconstituted with siRNA resistant
human ITGA5-GFP fused to a linker and RhoA-T19N pLL5.0 human ITGA5 shRNA/shRNA
resistant human ITGA5-GFP-RhoA-T19N, 4. pLL5.0 human E-cadherin shRNA knockdown

reconstituted with mE-cadherin-GFP fused to a linker and RhoA-T19N pLL5.0 E-cadherin
shRNA/mEcad–GFP-RhoA-T19N.

The following constructs were published previously. 1. pLL5.0 Myosin Regulatory Light
Chain-GFP, 2. pLL5.0 human E-cadherin shRNA knockdown reconstituted with mE-
cadherin-tdTomato and 3. pLL5.0 E-cadherin shRNA/mEcad–GFP used was as previously
described[19], 4. Lentiviral H2B-mNeon is a gift from Rong Li's lab.

Fibronectin 1 and Integrin alpha-5 were silenced by siRNA directed against human ITGA5
5'-GGACCAGGATGGCTACAATGA-3' and Dharmacon Smartpool siRNA directed against
human FN1(5'-GCAGCACAACUUCGAAUUA-3' ; 5'-GAAAUAGAUGCAACGAUCA-
3'; 5'-GAGGAAAUCUGCACAACCA-3'; 5'-CCUAAAGACUCCAUGAUCU-3')
respectively.

Computational fracture analysis of microbeads Analysis of microbeads at regions of
mesothelial fracture was performed as previously described with minor modifications and
additional quantifications[23]. Particle image velocimetry (PIV) analysis (PIVlab package[55] in
MATLAB) was used to quantify the flow fields in the top-viewed cell layer microbeads time-
lapse datasets. The PIV analysis was carried out for specific duration segments of these
datasets where the field of view was fixed. The time interval of each slice is 30 seconds. We
then carry out the PIV analysis by dividing the images into 64×64 pixel interrogation
windows with 50% overlap on the first pass, 32×32 pixel interrogation windows with 50%
overlap for the second pass, and 16×16 pixel interrogation windows with 50% overlap for the
third pass. Next, we select velocity threshold limits for post-processing the resulting velocity
vector fields, smooth the data, and use bicubic interpolation for missing vectors. After
calibrating the results, velocity magnitude ($v = \sqrt{v_x^2 + v_y^2}$), normal strain rate [$\dot{\varepsilon} = \partial_x v_x -$
$\partial_y v_y$][28] and shear strain rate [$\dot{\gamma} = \partial_x v_y + \partial_y v_x$] were calculated, where x and y are the
directions of vectors while v_x and v_y represent the corresponding mesothelium velocity
components[23].

Traction force microscopy TFM substrate was prepared as previously described [56] with
minor modifications. Briefly, PDMS with an average stiffness of 15 kPa was prepared by
dispensing 40 µL of 1:1 CyA and CyB mixture on the 14 mm well of the 35 mm glass bottom

dish (Cellvis), spin-coating the dish at 1000 rpm for 1 min and letting it cure at 80°C for 2 h.

PDMS surface was then silanised with 5% APTES in absolute ethanol, functionalised with

filtered and sonicated 0.2% (v/v) 0.2 µm dark red fluorescent carboxylate-modified

microspheres (FluoSpheres). Beads were inactivated by 0.1M Tri's treatment for 10 min. The

substrate was coated with 100 µg/mL of poly-D-lysine for 1 hour, followed by the adding 50

µg/ml (1:100) of human plasma fibronectin (Sigma-Aldrich) for overnight incubation at 4°C.

Subsequently, ZT mesothelial cells were seeded and incubated at 37°C to form a confluent

monolayer. Multicellular spheroids of OVCA433 were prepared and transferred to the

confluent monolayer and incubated for 15 min to allow spheroids to settle before live

imaging. Imaging was performed using Nikon W1 Spinning Disk confocal microscope with

the integrated Perfect Focus System on the x60 water magnification objectives. After 12 h,

10% SDS was added to remove cells, and the substrate was left alone to recover to the resting

state for an hour before acquiring reference images.

TFM Analysis The fluorescent bead images were maximum projected for each timepoint and

subsequently contrast-enhanced and registered to the maximum projected reference images

using MultiStackReg. PIVlab[57] add-on for MATLAB was used to calculate the displacement

field from the registered bead images using three passes of cross-correlation with the

interrogation windows as follows: 128(64), 64(32), and 32(16). Outlier vectors with

abnormally large values were removed post-processing. FTTC, an ImageJ plugin for a

Fourier transform-based, semi-infinite half-space method, was used to reconstruct the force

field from the displacement field[58]. Customised codes were written to generate the force map

or superimpose the traction force over the cell image. For downstream analysis requiring

quantification for each cell, a code was customised to import ImageJ ROIs into MATLAB to

calculate the parameters for each ROI. Radial traction stress is analysed as previously

described, with the middle of initially spreading cells or fracture site being the point of

reference[59]. Statistical analysis was performed using one-way ANOVA in GraphPad Prism.

Bayesian inversion stress microscopy and spheroid surrounding border analysis of

tensile stresses BISM analysis was performed on traction force microscopy data as

previously described[26, 36]. Next, we identified the spheroid's outline from brightfield live

imaging and created masks for a 10-micron border around spheroids that invaded or did not

invade with a MATLAB script. Stress values from the BISM analysis in each 10-micron

region bordering the spheroid are obtained to calculate the average stress.

Immunofluorescence and microscopy Cells were fixed at 4 ∘ C with either −20 °C methanol

on ice for 5 min or with 4% paraformaldehyde in cytoskeleton stabilization buffer (10 mM

PIPES at pH 6.8, 100 mM KCl, 300 mM sucrose, 2 mM EGTA and 2 mM MgCl2) on ice for

20 min and subsequently permeabilized with 0.2% Triton-X in PBS for 5 min at 4 °C[60-64].

For structured illumination microscopy, spinning-disc confocal microscopy (Yokogawa)

coupled with the Live SR module (GATACA Systems), Nikon Eclipse Ti-E inverted

microscope with Perfect Focus System, controlled by MetaMorph software (Molecular

device) supplemented with a 100x oil 1.45 NA CFI Plan Apo Lambda oil immersion

objective and sCMOS camera (Prime 95B, Photometrics)

1066

1067   For lattice light sheet microscopy, Zeiss Lattice Lightsheet 7 is equipped with a 10x 0.4 NA

1068   illumination objective at a 30-angle to cover glass and a 48x 1.0 NA detection objective at a

1069   60-angle to cover glass. Pre-defined Sinc3 beams with 30mm length x 1000nm thickness

1070   were used for image acquisition. Images were captured with Pco.edge 4.2 CLHS sCMOS

1071   camera and processed with Lattice Lightsheet 7 Processing Module.

1072

Antibodies and blocking peptide Primary antibodies in this study were: mouse monoclonal

antibody against human Fibronectin (1:100; BD Transduction Laboratories; catalogue

number 610077); mouse monoclonal antibody against human CD49e/Integrin α5 Chain

(1:1000 for western blotting; BD Pharmingen; catalogue number 555614); mouse monoclonal

antibody against human Integrin alpha-5 (1:100 for immunofluorescence; EMD Millipore;

catalogue number MABT201, clone number SNAKA51); rabbit monoclonal antibody [Y113]

against mouse and human Paxillin alpha, beta, and gamma isoforms (1:200 for

immunofluorescence; Abcam; catalogue number ab32084); mouse monoclonal antibody

against GAPDH (1:1000 for western blotting; Santa Cruz Biotechnology; catalogue number

sc-32233); rabbit polyclonal antibody against non-muscle myosin heavy chain IIA (1:200 for

immunofluorescence; Sigma-Aldrich; catalogue number M8064); rabbit polyclonal antibody

against laminin (1:100 for immunofluorescence; Novus Biologicals; catalogue number

NB300-144); Mouse monoclonal antibody against native and denatured forms of GFP

(1:1000 for western blotting; Invitrogen; catalogue number MA5-15256, clone number

GF28R).

Generation of micropatterned substrates Micropatterned substrate was fabricated by

stencil patterning. Polydimethylsiloxane (PDMS) (Sylgard 184 kit, Dow Corning) was

applied onto a photoresist mould, containing circular micropatterns. A mixture of elastomer

to crosslinker in a 10:1 ratio (w/w) was used, followed by curing at 80°C for 2 hours. The

resulting crosslinked PDMS layer was carefully detached, and stamps were manually crafted.

Inverting the PDMS stamp, it was positioned on an uncoated 35mm μ-dish (ibidi GmbH)

with hydrophobic properties. Next, Norland Optical Adhesive 73 (NOA-73, Norland Inc.)

was applied along one edge of the stamp. Through capillary action, NOA-73 flowed into the

gaps between the PDMS stamp and the dish, effectively sealing the stamp on all sides. Curing

the NOA-73 stencil took place under ultraviolet light for 15 seconds. Following removal of

the PDMS stamp, the stencil and dish were subjected to incubation with fluorescence

fibrinogen mixed fibronectin at 50 μg/ml concentration in PBS. This incubation occurred

overnight at 4°C following a brief degassing at 10 mbar. After the incubation, the fibronectin

solution was removed, and the stencil was detached. The dish's printed bottom was treated

with 0.2% Pluronic acid-H2O for 30 minutes as a passivation step. Finally, the passivated

dishes underwent three washes with PBS before cell seeding.

Image Processing and Analysis The images presented were processed with ImageJ

(http://rsb.info.nih.gov/ij/). For quantification of apical fluorescent density and line-scan

analysis, the mean intensity of the selected apical region was measured from the microscopy

images. Line-scan analysis was performed as described previously[62].

For quantification of the phosphoMLC intensity at E-cadherin cell-cell contacts and apical

integrin adhesions, segmented lines were drawn on immunostained E-cadherin cell-cell

contacts (line width 20) and apical integrin adhesions to obtain multiple regions of interest

with FIJI. The corresponding phosphoMLC intensities were measured at these regions of

interest.

For quantification of apical mesothelial paxillin intensity, an image mask was created by

subtracting the background intensity with the rolling ball radius set to 50 pixels with FIJI,

followed by the Gaussian blur with a sigma of 2. Particles with an area larger than 0.1 μm^2

were considered and segmented for the final mask to measure Paxillin intensity. The area of
each focal adhesion was recorded as a region of interest (ROI) in ImageJ, and the mean
intensity of the focal adhesions was calculated by subtracting the background intensity
obtained from an area of the image without paxillin.
1120
For quantifying mesothelial apical F-actin orientation, we measured the alignment of F-actin
bundles in cell fluorescence images at 10-degree intervals, using the Directionality Function
in Fiji (v. 1.54p).

For quantification of the abundance of fibronectin fibrils at the apical regions of the
mesothelium, an intensity threshold mask was created to outline the fibronectin fibrils with
FIJI. Their intensity multiplied by the area of these fibronectin fibrils are then compared
between the different experimental groups.

For quantification of cell-cell contact crazing, time-lapse image stacks of mesothelial
monolayers were analysed using Metamorph with Z-projection. Mesothelial monolayers were
first visually verified to be intact before tearing. Timeframes showing the initial cell-cell
contact breakdown were determined after playing through each movie and visually assessed
for multiple fibrillar bridges and microvoids indicative of crazing.

Spheroid nucleus segmentation with StarDist 3D deep learning model and cell tracking
by IMARIS The spheroids' image stacks underwent manual labelling using the Labkit plugin
in ImageJ. These labels were subsequently employed as the ground truth masks for the
StarDist 3D segmentation. A subset of the labelled stacks, selected randomly, was allocated
for training (N_train=19) and testing (N_test=2) datasets. To augment the data, various
techniques, including random flips, rotations, and intensity modifications, were applied
during training to expand the pool of training examples. The initial model parameters were
set using the pre-trained StarDist model weights.

Throughout the training process, the last five Conv3D layers of the model were reinitialized
and trained over 100 epochs. A learning rate of 10^{-4} was employed, along with a patch size of
8x96x96. All other training parameters were kept consistent as previously described.
Following this, the best-performing model, determined by the lowest test dataset loss,
underwent further fine-tuning for an additional 10 epochs. During this fine-tuning phase, a

reduced learning rate of 10^{-6} was employed. All training and fine-tuning procedures were

executed on a shared Jupyter notebook server equipped with an NVIDIA RTX 6000 GPU.

After model training, the trained model was deployed for predictions on new spheroid

images. This inference step was carried out using a Google Colab notebook with GPU

acceleration. After executing the prediction code, the resultant labels were compared with the

original image stack to assess segmentation accuracy. Following this evaluation, the

autoregressive motion algorithm was applied to enable automated tracking of the segmented

H2B mNeon nuclei time-lapse images in IMARIS to generate nuclei tracks. These tracks

were then subjected to manual refinement across all frames to accurately capture the

movement of spheroid cells.

Pair-wise mean square displacement (MSD) analysis of cell nucleus Given a time t, the

distance of two nuclei i and j is denoted as $Dx_{i,j}(t)$. The displacement of this distance over a

time interval dt is $Dx_{i,j}(t, dt) = Dx_{i,j}(t+dt) - Dx_{i,j}(t)$. Then, the pair-wise MSD $\langle Dx^2 \rangle (dt)$ is the

squared distance averaged over all time points and all nuclei pairs. MSD increases with dt

with a power $\langle Dx^2 \rangle \mu\, dt^s$, where $s(dt) = \partial \ln\langle Dx^2 \rangle / \partial \ln dt$, is referred to as local slope in the

main text. This local slope is 1 across all lag time dt if the particles move by random

diffusion.

Cell migration persistency analysis Let x_i be the position of cell nuclei i in the XYZ

(Cartesian) coordinate space, where the origin of the space $x_0 = \langle x \rangle$ is the averaged positions

of all cells in spheroids. Then, the velocity of nucleus i is calculated as the distance of the

nucleus between two consecutive frames: $v_i(t) = x_i(t+dt) - x_i(t)$, where $dt = 3$ minutes, the time

resolution of tracking. The velocity direction is $\hat{v}_i(t) = v_i(t)/|v_i(t)|$, from which the auto-

correlation of velocity directions $C(dt) = \langle \hat{v}_i(t)\hat{v}_i(t + dt) \rangle$ could be obtained. The decay of C

with the lag time dt follows an exponential form as $C \mu \exp(-dt/t)$, where t is the

characteristic timescales of persistency in the cell velocity direction.

Z-stacks of spheroid Let S be the total number of stacks to be sliced, and each stack is

enumerated as integer s ranging from 1 (top) to S (base). We first calculate the cumulative

distribution M of the cell nuclei in the spheroid along the Z direction, then $M=0$ corresponds

to the top surface and $M=1$ at the bottom surface. Next, we determine the upper bound z_s and

lower bound z'_s of a stack s in such a way that $M(z_s)= (s\text{-}1)/S$ and $M(z'_s)= \min(1, (s\text{-}1)/S + h\,)$,

where h is a predefined effective thickness $(0 < h < 1)$. Note that a fixed h imposes a similar

number of cells within each stack and excludes the bias from the population number in the

statistics. Finally, we select the cells with their vertical position z within the range of $[z_s, z'_s]$

for a stack s. In the main text, the total number of stacks S is 6, and the effective thickness h

is 0.3.

Interface Energy Formulation During spheroid wetting and invasion into the cell

monolayer, the free energy change (ΔG) of the system is due to variations in the interface

energies and cell polarization energy, i.e.,

$$\Delta G = \Delta\gamma_i + \Delta\gamma_m + \Delta\gamma_c + \phi_p, \tag{1}$$

in which $\Delta\gamma_i$, $\Delta\gamma_m$ and $\Delta\gamma_c$ represent changes in the cadherin-mediated spheroid cell-cell,

mesothelial cell-cell, and integrin-based spheroid-mesothelium interface energies,

respectively. ϕ_p is the work done by the polarization force in spheroid cells. Physically, these

aforementioned interface energies depend on mesothelial cell cortical tension T_0, spheroid

cell tension T_c, mesothelial adhesion strength W_0, spheroid adhesion strength W_c and integrin-

based adhesion energy W_I between spheroid leader cell and mesothelial cells.

1191

1192   During wetting, i.e. when the spheroid leader cell is in contact with the mesothelium, integrin

1193   is recruited to the interface (Fig. 3a,b) and increases the adhesion strength W_I. Here, such

1194   gradual enhancement of adhesion is described by

$$\tau_I \partial_t W_I = W_{I0} - W_I, \tag{2}$$

1196   where τ_I represents the timescale for integrin recruitment, W_{I0} is the maximum interface

1197   adhesion energy density that can be achieved.

1198

1199   It is widely reported that active contraction of cells could be triggered by mechanical signals

1200   such as local stretching and indentation [53, 65]. To incorporate this important feature into our

1201   model, a first-order coupling between the cortical tension T and the length L_{in} of spheroid-

1202   mesothelium adhesive interface [66, 67] was adopted as

$$\tau_A \partial_t T = T_0 - T + \frac{\beta L_{in}}{L_0}, \tag{3}$$

where T_0 is the static mesothelial cortical tension, L_0 is a reference length, τ_A characterizes

the reaction timescale for myosin recruitment and activation, and β is the tension provided by

the myosin recruitment per unit length of integrin adhesion. Evidently, under a fixed interface

length, the maximum mesothelium contractility that can be induced by myosin recruitment is

simply $T_0 + \frac{\beta L_{\text{in}}}{L_0}$.

During wetting, formation or elongation of intercellular junctions would bring in free energy

change in the system. Since active constriction is only found at the apical side of the

mesothelium, the interface energy changes are $\Delta\gamma_{\text{m}} = -W_0\delta L_{\text{m}}$, $\Delta\gamma_{\text{c}} = -W_{\text{c}}\delta L_{\text{c}}$ and

$$\Delta\gamma_{\text{i}} = (T_{\text{act}} - T_0 - W_{\text{I0}})\delta L_{\text{in}} \tag{4}$$

where T_{act} is the increased mesothelium apical tension in the equilibrium state. Considering

the steady-state solution of Eq. (3), the apical tension change is simply $T_{\text{act}} - T_0 = \frac{\beta L_{\text{in}}}{L_0}$. At

the same time, as spheroid cells move towards the mesothelium, free energy change induced

by polarization work is $\phi_{\text{p}} = -F_{\text{p}}\delta L_{\text{m}}$.

1217

At steady-state, the change of system free energy according to Eqs. (1-4) is

$$\Delta G = \frac{\beta}{L_0}\delta L_{\text{in}}^2 - W_{\text{I0}}\delta L_{\text{in}} + F_{\text{p}}\delta L_{\text{m}} - W_0\delta L_{\text{m}} - W_{\text{c}}\delta L_{\text{c}}, \tag{5}$$

where δL_{in}, δL_{m} and δL_{c} are the length change of spheroid-mesothelial integrin junction,

mesothelial E-cad junction and spheroid E-cad junction, respectively, F_{p} is the polarization

force in spheroid cells (Supplementary Fig. 3b). For the spheroid itself, the free energy

change reduces to $\Delta G_{\text{c}} = -W_{\text{I0}}\delta L_{\text{in}} - W_{\text{c}}\delta L_{\text{c}}$, which shows a competition between the

integrin interface energy and spheroid E-cad interface energy. This implies that the shortening

of spheroid E-cad junction due to the favoring of spheroid-mesothelial junctions lengthening,

promotes the wetting of spheroid on the mesothelium[68]. Indeed, experiments with spheroids

sorted for higher E-cadherin expression, which prevents cell-cell contact shrinkage[69]and

reduces spheroid invasion (Supplementary Fig. 4g,h). In the scenario of single-cell wetting,

we have $\delta L_{\text{c}} = 0$. Under such circumstances, the mesothelial free energy change is $\Delta G_{\text{m}} =$

$\frac{\beta}{L_0}\delta L_{\text{in}}^2 - W_{\text{I0}}\delta L_{\text{in}} + f_{\text{p}}\delta L_{\text{m}} - W_0\delta L_{\text{m}}$, indicating the fracture of mesothelial E-cad junction

(represented by the vanishing mesothelial junction length due to δL_{m}) is regulated by

mesothelial apical constriction, spheroid-mesothelium adhesion, and spheroid cell

polarization. Specifically, considering the complete rupturing condition $\delta L_{\text{m}} \leq -L_{\text{m0}}$ (with

L_{m0} being the initial length of the mesothelial junction), the critical condition for spontaneous

mesothelium fracturing (Fig. 5a) is

$$\Delta G_{\text{m}} \leq \frac{\beta}{L_0}\delta L_{\text{in}}^2 - W_{\text{I0}}\delta L_{\text{in}} - F_{\text{p}}L_{\text{m0}} + W_0 L_{\text{m0}} \leq 0. \tag{6}$$

For this equation to have a solution, we must have

1238
$$W_{\text{I}0}{}^2 - \frac{4\beta(W_0 - F_{\text{p}})L_{\text{m}0}}{L_0} \geq 0. \tag{7}$$

1239   Then, the critical condition for mesothelium fracture (Fig. 5a) can be obtained as

1240
$$\frac{W_{\text{I}0}}{W_0} \geq \left[4\left(1 - \frac{F_{\text{p}}}{\zeta T_0}\right)\frac{L_{\text{m}0}}{\zeta L_0}\right]^{1/2}\left(\frac{\beta}{T_0}\right)^{1/2}, \tag{8}$$

1241   where $\zeta = \frac{W_0}{T_0}$ defines the initial mesothelial tissue state.

1242

1243   **Active Foam Formulation** In our model, cell movement and deformation are captured by the

1244   position change of a set of vertices, describing the shape of each cell as

1245
$$\eta_R \frac{dR_\alpha}{dt} = \sum_{i,j \in F(\alpha)}\left(T_{ij} + N_{ij} + P_{ij}\right) \tag{9}$$

where R_α is the position vector of vertex α, t represents the time, η_R is the friction

coefficient associated with vertex movement, $F(\alpha)$ represents the set of junctions sharing

vertex α, N_{ij} is the normal force (due to osmotic pressure [50]) acting on the junction ij, T_{ij} is

the active cortical tension along the tangential direction of junction ij (Supplementary Fig.

3b). P_{ij} is the polarization force within the spheroid leader cell. Note that the effective

tension on adherent junctions is affected by the strength of cell-cell adhesion. For example,

the magnitude of effective junctional tension ij between cell a and cell b is expressed as

$T_{ij}^a + T_{ij}^b - W_{ij}^{ab}$, where W_{ij}^{ab} is the adhesion strength on segment ij between two cells. The

asymmetry of surface forces induced by cell adhesion elongates the interface and drives the

spreading of the spheroid leader cell on the mesothelium. The physical meaning and adopted

value of different parameters are summarized in Table S1.

1257

Initial Configuration Four tumor cells that formed a cell cluster and were designated near a

mesothelium cell layer were used to study the dynamic invasion process. Periodic boundary

conditions are applied to the left and right sides of the mesothelium monolayer during the

simulation. As the cancer cells are supposed to form a spheroid, a relatively high adhesion

energy among themselves is adopted (Table S1).

On the other hand, a very weak interface adhesion between cancer spheroid and

mesothelial cells (i.e., $W_{\text{I}0}/T_0 = 0.05$) is applied at the beginning of the simulation, given

that stable spheroid-mesothelial attachment takes time to form (after their initial contact). A

slightly lower E-cadherin adhesion energy (i.e., $0.8W_0$) between mesothelial cells is also

adopted. Before the actual simulation, the system was allowed to relax the steady state.

Once the simulation starts, a dynamic spheroid-mesothelium adhesion strength (given by

Eq. (3)) is invoked. Driven by the increased spheroid-mesothelial adhesion, the spheroid
leader cell spreads more on the mesothelium, triggering its apical contraction and junction
rupture and eventually leading to spheroid invasion into the mesothelial layer.

Based on our experimental observations, L_0 and T_0 are taken to be 20 mm and 5 nN,
respectively. According to previous studies [67, 70, 71], the friction coefficient η_R could vary from
0.01 to 0.1 N·s·m^{-1}; here, we take a high friction coefficient (0.1 N·s·m^{-1}) for mesothelial cells
as they are attached to the substrate and a low friction coefficient (0.05 N·s·m^{-1}) for the
floating spheroid cells. Since the integrin adhesion timescale is believed to be significantly
shorter than that for apical constriction [70, 72, 73], we choose $\tau_I = 20$ s and $\tau_A = 100$ s in the
simulation. The values of all dimensionless parameters are listed in Table S1.

Table S1 Description of dominating parameters

Parameter	Physical description	Range	Data Estimation
$\dfrac{L_c}{L_0}$	Characteristic spheroid size	1	Measurement
$\dfrac{T_c}{T_0}$	Spheroid cortical tension	0.5	Measurement
$\dfrac{\beta}{T_0}$	Integrin-based apical tension strengthening	0.5~5	Literature[66, 67]
$\dfrac{W_0}{T_0}$	Mesothelial adhesion strength	1.2	Shape estimation
$\dfrac{W_c}{T_c}$	Spheroid adhesion strength	1.1	Shape estimation
$\dfrac{P_0 L_0}{T_0}$	Static osmotic pressure	8000	Literature[74]
$\dfrac{F_p}{T_0}$	Cell polarization force	0.72	Literature[75]

1280

Onset of Tissue Fracture At the early stage of spheroid cell invasion, the mesothelial
junction below spheroid cells is shortened by activated apical contraction. In our simulation,
we set a mesothelial junction to be completely ruptured when its length is below a small
threshold value (i.e., $\leq L_0/100$), corresponding to the onset of tissue fracture in our
experiments. In this situation, the remaining central vertices are copied and divided equally
between the cells on both sides.

1287

Vertex Motion and Contact Formation Movements of vertices in our model are governed
by Eq. (1). To maintain the accuracy of the simulation; automatic re-meshing is adopted
where a new vertex is added to the middle of the boundary segment when its deformed length

is above a critical value (i.e., $L_0/20$).

To capture the growth of adhesive contact between cells in our simulation, vertices in front of

each contact tip are checked at every time step. Once a segment from one cell is close enough

to the other cell boundary, the two segments (as well as vertices) are combined. In this way,

elongation of the contacting spheroid-mesothelial interface is realized.

The shape index of the cell is calculated as $I_s = \frac{L_P}{\sqrt{S}}$ where L_P is the perimeter of the cell

and S is the area.

1298

Cell stiffness and surface tension measurement by atomic force microscopy (AFM) The

stiffness of the mesothelial cells and spheroids was measured using a Nanowizard IV

BioAFM system (JPK Instruments, Germany). Confluent mesothelial cell monolayers were

cultured on a glass bottom petri dish while the spheroids were adhered to a poly-d-lysine

(PDL) coated glass bottom petri dish. Indentations for surface tension measurements were

performed on randomly selected cells with a silicon nitride pyramidal tip (MLCT-D, Bruker;

nominal cantilever spring constant k = 0.03 N/m; nominal tip radius 20 nm) using a force of 3

nN at a speed of 5 µm/s. The force curves from more than nine mesothelial cells and multiple

cells from nine ovarian cancer spheroids were analysed to evaluate the surface tension. The

calculation of cell surface tension was performed as recently described[76-79]. In brief, the cell

membrane is assumed to be a thin tensile shell. With an AFM indentation range much larger

than cell membrane thickness, the formula used is T = K/2pi, where K is fitted from the linear

coefficient of force vs. indentation ranging from 50nm to 300 nm.

Data availability

The authors declare that the data supporting this study's findings are available within the

paper and its supplementary files. The raw data generated in this study are available from the

corresponding authors upon reasonable request.

Code availability The scripts generated in this study are available from

https://github.com/selwinwu/morphogenesis.git

1318

REFERENCES

1.   Friedl, P. & Alexander, S. Cancer invasion and the microenvironment: plasticity and
reciprocity. *Cell* **147**, 992-1009 (2011).
2.   Naora, H. & Montell, D.J. Ovarian cancer metastasis: integrating insights from
disparate model organisms. *Nat Rev Cancer* **5**, 355-366 (2005).
3.   Kulesa, P.M. & Gammill, L.S. Neural crest migration: patterns, phases and signals.
Dev Biol **344**, 566-568 (2010).
4.   Akhmanova, M. *et al.* Cell division in tissues enables macrophage infiltration. *Science*
376, 394-396 (2022).
5.   Stewart, T.A., Hughes, K., Hume, D.A. & Davis, F.M. Developmental Stage-Specific
Distribution of Macrophages in Mouse Mammary Gland. *Front Cell Dev Biol* **7**, 250
(2019).
6.   Cheung, K.J. *et al.* Polyclonal breast cancer metastases arise from collective
dissemination of keratin 14-expressing tumor cell clusters. *Proc Natl Acad Sci U S A*
113, E854-863 (2016).
7.   Dai, W. *et al.* Tissue topography steers migrating Drosophila border cells. *Science*
370, 987-990 (2020).
8.   Cai, D. *et al.* Mechanical feedback through E-cadherin promotes direction sensing
during collective cell migration. *Cell* **157**, 1146-1159 (2014).
9.   Wu, S.K. *et al.* Editorial: Forces in biology - Cell and developmental mechanobiology
and its implications in disease - Volume II. *Front Cell Dev Biol* **10**, 1082857 (2022).
10.   Wu, S.K., Gomez, G.A., Stehbens, S.J. & Smutny, M. Editorial: Forces in Biology - Cell
and Developmental Mechanobiology and Its Implications in Disease. *Front Cell Dev*
Biol **8**, 598179 (2020).
11.   Iwanicki, M.P. *et al.* Ovarian cancer spheroids use myosin-generated force to clear
the mesothelium. *Cancer Discov* **1**, 144-157 (2011).
12.   Kenny, H.A. *et al.* Mesothelial cells promote early ovarian cancer metastasis through
fibronectin secretion. *J Clin Invest* **124**, 4614-4628 (2014).
13.   Davidowitz, R.A. *et al.* Mesenchymal gene program-expressing ovarian cancer
spheroids exhibit enhanced mesothelial clearance. *J Clin Invest* **124**, 2611-2625
(2014).
14.   Naffar-Abu Amara, S. *et al.* Transient commensal clonal interactions can drive tumor
metastasis. *Nat Commun* **11**, 5799 (2020).
15.   Mutsaers, S.E. Mesothelial cells: their structure, function and role in serosal repair.
Respirology **7**, 171-191 (2002).
16.   Birbeck, M.S. & Wheatley, D.N. An Electron Microscopic Study of the Invasion of
Ascites Tumor Cells into the Abdominal Wall. *Cancer Res* **25**, 490-497 (1965).
17.   Kenny, P.A. *et al.* The morphologies of breast cancer cell lines in three-dimensional
assays correlate with their profiles of gene expression. *Mol Oncol* **1**, 84-96 (2007).
18.   Rottler, J., Barsky, S. & Robbins, M.O. Cracks and crazes: on calculating the
macroscopic fracture energy of glassy polymers from molecular simulations. *Phys*
Rev Lett **89**, 148304 (2002).
19.   Smutny, M. *et al.* Multicomponent analysis of junctional movements regulated by
myosin II isoforms at the epithelial zonula adherens. *PLoS One* **6**, e22458 (2011).
20.   Shewan, A.M. *et al.* Myosin 2 is a key Rho kinase target necessary for the local
concentration of E-cadherin at cell-cell contacts. *Mol Biol Cell* **16**, 4531-4542 (2005).
21.   Wu, S.K. & Priya, R. Spatio-Temporal Regulation of RhoGTPases Signaling by Myosin
II. *Front Cell Dev Biol* **7**, 90 (2019).

22. Acharya, B.R. *et al.* Mammalian Diaphanous 1 Mediates a Pathway for E-cadherin to
Stabilize Epithelial Barriers through Junctional Contractility. *Cell Rep* **18**, 2854-2867
(2017).
23. Prakash, V.N., Bull, M.S. & Prakash, M. Motility-induced fracture reveals a ductile-to-
brittle crossover in a simple animal's epithelia. *Nature Physics* **17**, 504-511 (2021).
24. Greenhalgh, E.S. *Failure analysis and fractography of polymer composites.*
(Woodhead Publishing Limited : CRC Press, Cambridge, England ; New Delhi, India ;
Boca Raton, Florida; 2009).
25. Anderson, T.L. *Fracture mechanics : fundamentals and applications*, Edn. Fourth
edition. (CRC Press/Taylor & Francis, Boca Raton; 2017).
26. Sonam, S. *et al.* Mechanical stress driven by rigidity sensing governs epithelial
stability. *Nature Physics* **19**, 132-141 (2023).
27. Lv, J.-Q. *et al.* Active hole formation in epithelioid tissues. *Nature Physics* **20**, 1313-
1323 (2024).
28. Stamhuis, E.J. Basics and principles of particle image velocimetry (PIV) for mapping
biogenic and biologically relevant flows. *Aquatic Ecology* **40**, 463-479 (2006).
29. Sawyer, J.M. *et al.* Apical constriction: a cell shape change that can drive
morphogenesis. *Dev Biol* **341**, 5-19 (2010).
30. Martin, A.C. & Goldstein, B. Apical constriction: themes and variations on a cellular
mechanism driving morphogenesis. *Development* **141**, 1987-1998 (2014).
31. Martin, A.C., Kaschube, M. & Wieschaus, E.F. Pulsed contractions of an actin-myosin
network drive apical constriction. *Nature* **457**, 495-499 (2009).
32. Martin, A.C., Gelbart, M., Fernandez-Gonzalez, R., Kaschube, M. & Wieschaus, E.F.
Integration of contractile forces during tissue invagination. *J Cell Biol* **188**, 735-749
(2010).
33. Wu, S.K. & Yap, A.S. Patterns in space: coordinating adhesion and actomyosin
contractility at E-cadherin junctions. *Cell Commun Adhes* **20**, 201-212 (2013).
34. Mason, F.M., Xie, S., Vasquez, C.G., Tworoger, M. & Martin, A.C. RhoA GTPase
inhibition organizes contraction during epithelial morphogenesis. *J Cell Biol* **214**, 603-
617 (2016).
35. Wu, S.K., Budnar, S., Yap, A.S. & Gomez, G.A. Pulsatile contractility of actomyosin
networks organizes the cellular cortex at lateral cadherin junctions. *Eur J Cell Biol* **93**,
396-404 (2014).
36. Nier, V. *et al.* Inference of Internal Stress in a Cell Monolayer. *Biophys J* **110**, 1625-
1635 (2016).
37. Parsons, J.T., Horwitz, A.R. & Schwartz, M.A. Cell adhesion: integrating cytoskeletal
dynamics and cellular tension. *Nat Rev Mol Cell Biol* **11**, 633-643 (2010).
38. Wu, S.K. *et al.* Cortical F-actin stabilization generates apical-lateral patterns of
junctional contractility that integrate cells into epithelia. *Nat Cell Biol* **16**, 167-178
(2014).
39. Ladoux, B. & Mege, R.M. Mechanobiology of collective cell behaviours. *Nat Rev Mol
Cell Biol* **18**, 743-757 (2017).
40. Wu, S.K., Lagendijk, A.K., Hogan, B.M., Gomez, G.A. & Yap, A.S. Active contractility at
E-cadherin junctions and its implications for cell extrusion in cancer. *Cell Cycle* **14**,
315-322 (2015).

41. Gomez, G.A. *et al.* An RPTPalpha/Src family kinase/Rap1 signaling module recruits
myosin IIB to support contractile tension at apical E-cadherin junctions. *Mol Biol Cell*
26, 1249-1262 (2015).
42. Han, S.P. *et al.* Cortactin scaffolds Arp2/3 and WAVE2 at the epithelial zonula
adherens. *J Biol Chem* **289**, 7764-7775 (2014).
43. Leerberg, J.M. *et al.* Tension-sensitive actin assembly supports contractility at the
epithelial zonula adherens. *Curr Biol* **24**, 1689-1699 (2014).
44. Sipes, J.M. *et al.* Inhibition of fibronectin binding and fibronectin-mediated cell
adhesion to collagen by a peptide from the second type I repeat of thrombospondin.
J Cell Biol **121**, 469-477 (1993).
45. Liotta, L.A. & Kohn, E. Anoikis: cancer and the homeless cell. *Nature* **430**, 973-974
(2004).
46. Weigert, M., Schmidt, U., Haase, R., Sugawara, K. & Myers, G. in Proceedings of the
IEEE/CVF winter conference on applications of computer vision 3666-3673 (2020).
47. Ho, C.Z. *et al.* Protocol for AI-assisted quantitative analysis and setup of tumor
spheroid invasion into tissue. *STAR Protoc* **6**, 104140 (2025).
48. Ilina, O. *et al.* Cell-cell adhesion and 3D matrix confinement determine jamming
transitions in breast cancer invasion. *Nat Cell Biol* **22**, 1103-1115 (2020).
49. Mongera, A. *et al.* A fluid-to-solid jamming transition underlies vertebrate body axis
elongation. *Nature* **561**, 401-405 (2018).
50. Kim, S., Pochitaloff, M., Stooke-Vaughan, G.A. & Campas, O. Embryonic Tissues as
Active Foams. *Nat Phys* **17**, 859-866 (2021).
51. Sirka, O.K., Shamir, E.R. & Ewald, A.J. Myoepithelial cells are a dynamic barrier to
epithelial dissemination. *J Cell Biol* **217**, 3368-3381 (2018).
52. Munster, S. *et al.* Attachment of the blastoderm to the vitelline envelope affects
gastrulation of insects. *Nature* **568**, 395-399 (2019).
53. Bailles, A. *et al.* Genetic induction and mechanochemical propagation of a
morphogenetic wave. *Nature* **572**, 467-473 (2019).
54. de Gennes, P.G. Wetting: statics and dynamics. *Reviews of Modern Physics* **57**, 827-
863 (1985).
55. Thielicke, W. & Stamhuis, E.J. PIVlab – Towards User-friendly, Affordable and
Accurate Digital Particle Image Velocimetry in MATLAB. *Journal of Open Research*
Software **2** (2014).
56. Teo, J.L., Lim, C.T., Yap, A.S. & Saw, T.B. A Biologist's Guide to Traction Force
Microscopy Using Polydimethylsiloxane Substrate for Two-Dimensional Cell Cultures.
STAR Protoc **1**, 100098 (2020).
57. Thielicke, W. & Sonntag, R. Particle Image Velocimetry for MATLAB: Accuracy and
enhanced algorithms in PIVlab. *Journal of Open Research Software* (2021).
58. Tseng, Q. (Université de Grenoble, 2011).
59. Kawaue, T. *et al.* Inhomogeneous mechanotransduction defines the spatial pattern
of apoptosis-induced compensatory proliferation. *Dev Cell* **58**, 267-277 e265 (2023).
60. Liu, O.X. *et al.* A ZO-2 scaffolding mechanism regulates the Hippo signalling pathway.
FEBS J **292**, 1587-1601 (2025).
61. Greenlees, R. *et al.* Mutations in SIPA1L3 cause eye defects through disruption of cell
polarity and cytoskeleton organization. *Hum Mol Genet* **24**, 5789-5804 (2015).
62. Mangold, S. *et al.* Hepatocyte growth factor acutely perturbs actin filament
anchorage at the epithelial zonula adherens. *Curr Biol* **21**, 503-507 (2011).

63.   Moore, T. *et al.* Self-organizing actomyosin patterns on the cell cortex at epithelial
1460   cell-cell junctions. *Biophys J* **107**, 2652-2661 (2014).

1461   64.   Lin, D. *et al.* The Senescence-Associated-Secretory Phenotype constituting HIF-1α
1462   activation is independent of micronuclei induction and associated cGAS-mediated
1463   interferon response. *bioRxiv*, 2024.2009.2025.615081 (2024).

1464   65.   Pouille, P.A., Ahmadi, P., Brunet, A.C. & Farge, E. Mechanical signals trigger Myosin II
1465   redistribution and mesoderm invagination in Drosophila embryos. *Sci Signal* **2**, ra16
1466   (2009).

1467   66.   Recho, P., Hallou, A. & Hannezo, E. Theory of mechanochemical patterning in
1468   biphasic biological tissues. *Proc Natl Acad Sci U S A* **116**, 5344-5349 (2019).

1469   67.   Sun, F., Fang, C., Shao, X., Gao, H. & Lin, Y. A mechanism-based theory of cellular and
1470   tissue plasticity. *Proc Natl Acad Sci U S A* **120**, e2305375120 (2023).

1471   68.   Perez-Gonzalez, C. *et al.* Active wetting of epithelial tissues. *Nat Phys* **15**, 79-88
1472   (2019).

1473   69.   Maitre, J.L. & Heisenberg, C.P. Three functions of cadherins in cell adhesion. *Curr Biol*
1474   **23**, R626-633 (2013).

1475   70.   Sun, F., Fang, C., Shao, X., Gao, H. & Lin, Y. A mechanism-based theory of cellular and
1476   tissue plasticity. *Proc. Natl. Acad. Sci. USA* **120**, e2305375120 (2023).

1477   71.   Lin, S.Z., Li, B., Lan, G. & Feng, X.Q. Activation and synchronization of the oscillatory
1478   morphodynamics in multicellular monolayer. *Proc. Natl. Acad. Sci. USA* **114**, 8157-
1479   8162 (2017).

1480   72.   Hu, Y. *et al.* DNA-based ForceChrono probes for deciphering single-molecule force
1481   dynamics in living cells. *Cell* **187**, 3445-3459.e3415 (2024).

1482   73.   Boocock, D., Hino, N., Ruzickova, N., Hirashima, T. & Hannezo, E. Theory of
1483   mechanochemical patterning and optimal migration in cell monolayers. *Nat. Phys.*
1484   **17**, 267-274 (2020).

1485   74.   Guo, M. *et al.* Cell volume change through water efflux impacts cell stiffness and
1486   stem cell fate. *Proc Natl Acad Sci U S A* **114**, E8618-E8627 (2017).

1487   75.   Messi, Z., Bornert, A., Raynaud, F. & Verkhovsky, A.B. Traction Forces Control Cell-
1488   Edge Dynamics and Mediate Distance Sensitivity during Cell Polarization. *Curr Biol*
1489   **30**, 1762-1769 e1765 (2020).

1490   76.   Pontes, B., Monzo, P. & Gauthier, N.C. Membrane tension: A challenging but
1491   universal physical parameter in cell biology. *Semin Cell Dev Biol* **71**, 30-41 (2017).

1492   77.   Couturier, E., Vella, D. & Boudaoud, A. Compression of a pressurized spherical shell
1493   by a spherical or flat probe. *Eur Phys J E Soft Matter* **45**, 13 (2022).

1494   78.   Vella, D., Ajdari, A., Vaziri, A. & Boudaoud, A. The indentation of pressurized elastic
1495   shells: from polymeric capsules to yeast cells. *J R Soc Interface* **9**, 448-455 (2012).

1496   79.   Biswas, A. *et al.* Theca cell mechanics and tissue pressure regulate mammalian
1497   ovarian folliculogenesis. *bioRxiv*, 2024.2005.2006.592641 (2024).

1498

---

i

Figure 1

a Without metastatic tumour implantation

Mesothelium

Peritoneum Space

Stroma

Adipose tissue

b Invading ovarian tumour

b'

c Mesothelium   Ecad Spheroid   Laminin basement membrane

d Spheroid   Mesothelium   Mesothelial E-cadherin tdT

d' 0 mins   45 mins   93 mins   300 mins

d''' 0 mins   6 mins   12 mins   18 mins   24 mins

High   Intensity   Low

d'' Cell 1   Cell 3   Cell 2   Cell 4   27 mins   45 mins   93 mins   300 mins   1 2 3 4

e E-cad   GFP   Linker   RhoA T19N

f Mesothelium Ecad   Spheroid   WT   +RhoA T19N

g % of 'Craze' at microfracture (%)   Ecad   Ecad +RhoA T19N   ****

h Mesothelial Clearance (a.u)   Ecad   Ecad +RhoA T19N   ****

i Mesothelial   Ecad   0 mins   Spheroid   81 mins

Shear strain rate (s^{-1})   −0.04   0.04

Normal strain rate (s^{-1})   −0.04   0.04

Velocity (µm/s)   0.01   0.05

0 mins   81 mins   j

Initial fracture   Constriction   Beads displacement

j Mesothelium   Fracture Modes   I: Opening   II: In-plane shear   III: Out-of-plane shear

k Fold change   Initial fracture   Shear Strain rate   Normal Strain rate   Velocity magnitude   *   **

l Spheroid Ecad GFP   Mesothelial Ecad tdT   XZ view   0 min   12 mins   24 mins   63 mins

Figure 2

a Mesothelial E-cad tdT MRLC GFP

Spheroid

0 min   9 mins   12 mins   21 mins   60 mins

b

Mesothelial apical cell area variance

***

W/O Invasion   With Invasion

c Mesothelial medioapical MRLC-GFP

High

Intensity

Low

c'

d

Onset of Mesothelial E-cad contact Rupture

Apical cortex MRLC mean intensity(a.u)

Mesothelial cell apical area (µm²)

Time (minutes)

e

Spheroid   Traction Vectors

Ecad-GFP spheroid cell

(Pa)

Radial Traction

0 min   10 mins   20 mins

f Spheroid E-cad GFP Basal Mesothelial E-cad tdT

Spheroid

Traction vectors

Radial Traction

(Pa)

0 min   15 mins   30 mins

g

Average Traction Stress (Pa)

-9mins -6mins -3mins Initial Fracture 3mins 6mins 9mins

h

Spheroid

Region Quantified (BISM)

i Bayesian Inversion Stress Microscopy

Tension (kPa µm)

*

Control   3 minutes before Spheroid Invasion

Figure 3

a. Spheroid Paxillin-GFP Mesothelial E-cad tdT

XY

Spheroid

XZ

XY

XZ view of integrin adhesion on mesothelium

Spheroid Paxillin-GFP

b. Spheroid cell Pax-GFP

0 min

5 mins

10 mins

15 mins

20 mins

c. Spheroid E-cad GFP Mesothelial E-cad tdT

Spheroid

c'

c''

c'''

c''

c'''

d.

e. Rate of Invasion (%)

Cont   TSP-1

f. Mesothelial E-cad tdT

Control   Spheroid

Spheroid ITGA5 KD

g. Rate of Invasion (%)

Control   Spheroid ITGA5 KD

h. Mesothelial E-cad tdT

Spheroid

0 min   240 mins   339 mins   387 mins   603 mins

i. Mesothelial Pax-GFP FN1

Spheroid

j. Mesothelial cells apical region

Paxillin   ITGA5   Paxillin   FN1   Tensin-1   Vinculin   Kank-2   Vinculin

k. Rate of Invasion (%)

Control   Mesothelial FN1 KD   Spheroid FN1 KD   Serum free

l. Rate of Invasion (%)

Control   Mesothelial ITGA5 KD

m. ITGA5   GFP   Linker   RhoA T19N

n. Mesothelial ITGA5 KDR E-cad tdT

Spheroid

ITGA5 KDR + RhoA T19N

o. Rate of Invasion (%)

ITGA5 KDR   ITGA5 KDR+RhoA T19N   ITG5A KDR with FN1   ITG5A KDR+RhoA T19N with FN1

Figure 4

a OVCA433 suspension spheroid

Intercellular integrin-matrix engagement

Leader cell

Integrin alpha-5 complex

Fibronectin fibrils

Mesothelium

2. Mesothelial apical constriction & tensile fracture initation

3. Spheroid accesses submesothelial substrate and spreads

b Control Invading spheroid
H2B-mNeon
3D Segmentation

3D Segmented nuclei

c Invading spheroid cell tracks

XY, top view XZ, side view RZ, side view

TIME

d Velocity maps

Control: 6 mins prior to fracture

Top Base

XZ side Spheroid RZ side

Control: 3 mins prior to fracture

Top Base

XZ side Spheroid RZ side

Mesothelium ITGA5 KD

Top Base

XZ side Spheroid RZ side

e Ratio of Vz mean/ Vxy Std Dev (Spheroid cells prior to Fracture)

Control ITGA5 KD

Mesothelium

f Spheroid cells directed migration persistency (mins)

No invasion (ITG5A KD mesothelium) Invasion (control mesothelium)

g Spheroid cells Pairwise MSD (μm^2)

non-invasion invasion

0.87 1.31

dt (min)

i Leader cell spreading area (μm^2) Mesothelial junction length (μm)

Onset of Mesothelial E-cad contact Rupture

Mesothelial spreading Sub-mesothelial spreading

Time (mins)

h Spheroid w/o invasion (ITGA5 KD mesothelium)

Local slope

$dt \sim 50$ min

Top Z region Base
1 2 3 4 5 6

Spheroid invasion (control mesothelium)

Top Z region Base
1 2 3 4 5 6

j OVCA433 E-cad GFP

High E-cad expression Spheroid

Control

k Rate of Invasion (%)

Control High E-cad expression

Figure 5