Title: Automatic learning mechanisms for flexible human locomotion.

Authors: Cristina Rossi[a,b], Kristan A. Leech[c,d], Ryan T. Roemmich[b,e], Amy J. Bastian[a,b,*]

[a] Department of Neuroscience, The Johns Hopkins University School of Medicine, Baltimore, MD, 21205,
USA; [b] Center for Movement Studies, Kennedy Krieger Institute, Baltimore, MD, 21205, USA; [c] Division of
Biokinesiology and Physical Therapy, University of Southern California, Los Angeles, CA, 90033, USA; [d]
Neuroscience Graduate Program, University of Southern California, Los Angeles, CA, 90007, USA; [e]
Department of Physical Medicine and Rehabilitation, The Johns Hopkins University School of Medicine,
Baltimore, MD, 21205, USA

*Corresponding author: Amy J. Bastian.

Email: bastian@kennedykrieger.org

Author Contributions: Study design & implementation: C.R., K.A.L., R.T.R., A.J.B.; Data
acquisition & analysis: C.R., K.A.L.; Interpretation of results: C.R., K.A.L., R.T.R., A.J.B.; Writing
& revision: C.R., K.A.L., R.T.R., A.J.B.

Competing Interest Statement: The authors declare no competing interests.

Classification: Biological Sciences: Neuroscience.

Keywords: Adaptation. Motor control. Motor learning. Human locomotion. Perception.

This PDF file includes:

22
23
24

Abstract

Movement flexibility and automaticity are necessary to successfully navigate different environments. When encountering difficult terrains such as a muddy trail, we can change how we step almost immediately so that we can continue walking. This flexibility comes at a cost since we initially must pay deliberate attention to how we are moving. Gradually, after a few minutes on the trail, stepping becomes automatic so that we do not need to think about our movements. Canonical theory indicates that different adaptive motor learning mechanisms confer these essential properties to movement: explicit control confers flexibility, while forward model recalibration confers automaticity. Here we uncover a distinct mechanism of treadmill walking adaptation – an automatic stimulus-response mapping - that confers both properties to movement. The mechanism is flexible as it learns stepping patterns that can be rapidly changed to suit a range of treadmill configurations. It is also automatic as it can operate without deliberate control or explicit awareness by the participants. Our findings reveal a tandem architecture of forward model recalibration and automatic stimulus-response mapping mechanisms for walking, reconciling different findings of motor adaptation and perceptual realignment.

Abbreviations
CI, confidence interval; CI_{LB}, lower bound of CI; CI_{UB}, upper bound of CI; PSE, point of subjective equality; norm, normalized; SD, standard deviation; SE, standard error

Main Text

Introduction

Flexibility and automaticity are essential features of human movement, so much so that we rarely
think about the details of how we move. While walking, we do not think about the bend of the
ankle or how quickly to swing the leg forward to step. Yet, we easily adjust walking to
accommodate many situations – a muddy trail, a grassy slope, or a snowy path. These abilities
are often taken for granted until something goes awry – a sprained ankle quickly brings the details
of movement execution to awareness and deliberate control is often used to avoid pain or further
injury. It is only then that we truly appreciate how much the sensorimotor system is doing without
our conscious awareness.

We currently do not understand how different learning mechanisms confer both flexibility and
automaticity to human movement. One well-studied process of motor learning is sensorimotor
adaptation, which occurs in response to errors and helps adjust motor commands to correct for
the perturbations to our movements that are introduced by the changed environmental demands
(Bastian, 2008; Huberdeau et al., 2015). Adaptation is traditionally thought to rely on the
cerebellar-dependent recalibration of a forward model that associates motor commands with
expected sensory consequences (Ito, 1989; Wolpert et al., 2001). For example, to walk on an icy
sidewalk, we may need to recalibrate our prediction of how firmly our feet will grip the ground.
The process of forward model recalibration is automatic and implicit, as it has been shown to
operate without intention or awareness (Huberdeau et al., 2015; Taylor et al., 2014). However,
forward model recalibration is not as flexible because it cannot make immediate changes in
movement – it can only adjust movement gradually trial-by-trial or step-by-step, a process that
can take several minutes (Bastian, 2008; Huberdeau et al., 2015). For this reason, forward model
recalibration leads to lasting movement errors (called "aftereffects") when the environment returns
to its original state after adaptation, as the newly acquired sensorimotor recalibration must be
unlearned over time to restore a normal movement (Bastian, 2008; Martin et al., 1996).

Adaptation can rely on more than forward model recalibration. For example, adaptation of
reaching movements consistently involves the use of explicit strategies, where people deliberately
change where they are aiming to reach (Taylor et al., 2014; Taylor and Ivry, 2011). It can also
involve memory-based caching (Huberdeau et al., 2019; McDougle and Taylor, 2019), where

people learn motor responses in association with the respective environmental sensory stimuli,
and cache them in memory for future retrieval, as in a lookup table. Finally, adaptation of reaching
movements can involve structural learning, where people learn the general relationships or
equations that link environmental sensory stimuli and appropriate motor responses, and later use
them to compute motor responses anew (Bond and Taylor, 2017; Braun et al., 2009). Aiming
strategies, memory-based caching, and structural learning differ from forward model recalibration
in that they do not lead to aftereffects (Huberdeau et al., 2015; Taylor et al., 2014; Taylor and Ivry,
2011). This is thought to be because the motor responses are mapped to sensory information
about the environment and are therefore flexible – they can be promptly abandoned or changed
for different environmental stimuli. Hence, these mechanisms are collectively called "stimulus-
response mapping mechanisms". In reaching, stimulus-response mapping mechanisms do not
operate automatically, and rely instead on explicit control (Bond and Taylor, 2017; Huberdeau et
al., 2019; Taylor and Ivry, 2011). However, explicit control is poorly suited for automatic,
continuous movements like walking (Clark, 2015; Paul et al., 2005) and can even interfere with
them (Uiga et al., 2020; Wong et al., 2008). Unlike discrete reaching, individuals do not
consciously aim their feet while engaged in continuous walking on surfaces that are typically
encountered (e.g. floors, sidewalks) (Long et al., 2016; Roemmich et al., 2016). For these
reasons, we speculated that other stimulus-response mapping mechanisms that are automatic
and not explicit may contribute to walking adaptation.

Studying the automatic mechanisms of walking adaptation has proven challenging, because it is
still unknown how to measure each mechanism in isolation. Here, we propose leveraging the
phenomenon known as "perceptual realignment" to dissect the mechanisms underlying walking
adaptation. When we adapt, we learn not only a new *motor* calibration, but also a new *perceptual*
calibration of the action we are performing ('t Hart and Henriques, 2016; Haith et al., 2008; Jensen
et al., 1998; Vazquez et al., 2015). For example, as people adapt their reaching movements, they
also realign their perception of movement direction or force production ('t Hart and Henriques,
2016; Haith et al., 2008). Similarly, as people adapt their walking movements on a split-belt
treadmill (one foot made to move faster than the other), they also realign their perception of leg
speed (Jensen et al., 1998; Kristan A. Leech et al., 2018; Vazquez et al., 2015). Despite
suggestions that perceptual realignment may also stem from recalibration of a forward model ('t
Hart and Henriques, 2016; Izawa et al., 2012; Rossi et al., 2021a; Synofzik et al., 2008), we do
not yet know whether *the same* forward model recalibrates both movement and perception in

parallel. It remains unclear whether there exists a direct relationship between perceptual
realignment and the portion of motor adaptation achieved specifically by a mechanism of forward
model recalibration. Establishing such a relationship would provide compelling evidence that
perceptual realignment can be used as a measure of the forward model recalibration mechanism
of motor adaptation. In turn, this would allow us to quantify the contribution of other, stimulus-
response mapping mechanisms to adaptation.

In this study, we aimed to identify the learning mechanisms involved in split-belt treadmill walking
adaptation. In Experiment 1, we uncovered a stimulus-response mapping mechanism that
contributes to walking adaptation. We also isolated forward model recalibration and show that it
changes movement and perception together. In Experiment 2, we found that the stimulus-
response mapping mechanism is automatic and not under deliberate or explicit control; rather, it
operates via either memory-based caching or structural learning.

Results

<u>Experiment 1</u>

We asked whether walking adaptation involves both forward model recalibration and a stimulus-
response mapping mechanism learned in tandem. We hypothesized that we could assess this by
carefully studying the motor and perceptual aftereffects that occur after split-belt walking. Note
that "Recalibration" and "mapping" are at times used as shorthand for "forward model
recalibration" and "stimulus-response mapping".

Motor paradigm and hypotheses

We used established motor measurements to quantify split-belt walking adaptation (Reisman et
al., 2005) (Finley et al., 2015). The "Perturbation" measures the effect of the split-belt treadmill on
the stepping pattern — the total movement error we would see in the absence of adaptation. "Δ
Motor output" measures the extent that individuals compensate for the perturbation by changing
their stepping pattern — how much they alter when (time) and where (position) they step on the

treadmill with each foot. "Step length asymmetry" measures the remaining movement error — the
difference between the perturbation and Δ motor output (see Methods).

Figure 1A illustrates the standard paradigm used in studies of split-belt adaptation, with abrupt
transitions between tied-belt to split-belt phases, and Figure 1B illustrates the standard motor
response. In the "baseline" phase, the perturbation, Δ motor output, and step length asymmetry
are zero reflecting equal belt speeds and symmetric walking (red, blue, purple, respectively). In
the "adaptation" phase, the belt speeds are different and the perturbation is positive. Initially, this
leads to movement errors observed as the negative step length asymmetry. The Δ motor output
gradually adapts to compensate for the perturbation, so that the step length asymmetry returns
to zero. In "post-adaptation", the belt speeds are tied and the perturbation is again zero.
Individuals exhibit initial movement errors called "aftereffects": the Δ motor output mismatches the
perturbation (it remains elevated) and step length asymmetry is positive. Importantly, these
aftereffects are smaller than the total motor learning: the Δ motor output is *less* elevated than in
late adaptation, and the step length asymmetry magnitude is *smaller* than in initial adaptation
(Kristan A. Leech et al., 2018; Rossi et al., 2021a, 2019; Statton et al., 2018).

Why are aftereffects smaller than the perturbation that was learned? We hypothesize that people
may learn to compensate for the perturbation using a combination of forward model recalibration
and stimulus-response mapping mechanisms. The key distinction between these mechanisms is
that forward model recalibration leads to aftereffects (Bastian, 2008; Martin et al., 1996; Reisman
et al., 2010), while stimulus-response mapping mechanisms do not (Huberdeau et al., 2015;
Taylor et al., 2014; Taylor and Ivry, 2011) (SI Appendix, Experiment 1 mechanisms characteristics
and Fig. S1). Hence, this hypothesis would account for the consistent finding that aftereffects are
"partial" even when motor learning is "complete", because only the portion learnt by recalibration
would lead to aftereffects.

Standard paradigms with abrupt transitions from adaptation to post-adaptation phases are
insufficient for assessing the hypothesis that people learn through both forward model
recalibration and stimulus response mapping. This is because aftereffects on tied-belts may
inaccurately appear "partial" due to testing methodologies (e.g., unlearning or forgetting due to
momentary stops in the treadmill between different testing conditions) (Malone and Bastian,

2016), leaving open the possibility that people exclusively learn through forward model
recalibration.

To obtain a robust measure of the aftereffects, we devised a "Ramp Down" paradigm with a
gradual transition from adaptation to post-adaptation. Specifically, the right belt speed was
ramped down for ~80 seconds until it matched the left belt speed (Fig. 1C), so that the perturbation
returned to zero gradually. We asked whether participants exhibit aftereffects (Δ motor output
mismatching perturbation, and positive step length asymmetry) for these intermediate
perturbations. No aftereffects for a range of perturbations would be indicative of a stimulus-
response mapping mechanism, because only Δ motor outputs learnt by mapping can flexibly
change to match a range of perturbations.

We used this paradigm to whether adaptation is achieved by 1) forward model recalibration only,
2) stimulus-response mapping only, or 3) a combination of the two mechanisms. The three
competing hypotheses make different predictions about when in the Ramp Down the aftereffects
should emerge (Fig. 1D).

If adaptation involves forward model recalibration only, aftereffects should emerge immediately in
the Ramp Down. The Δ motor output would remain elevated, because adjustments would take a
longer time to change. Hence, Δ motor output (blue) would immediately diverge from the
perturbation (red) and step length asymmetry (purple) would immediately grow (Fig. 1D top; also
see SI Appendix, Fig. S1B).

If adaptation involves stimulus-response mapping only, there would be no aftereffects during the
Ramp Down. Specifically, the Δ motor output would match the perturbation throughout the task,
because adjustments to the Δ motor output by mapping are flexible and can be promptly scaled
down. In turn, step length asymmetry would remain zero (Fig. 1D middle).

If adaptation involves both recalibration and mapping, aftereffects should emerge at some mid-
point during the Ramp Down. The Δ motor output would initially match the perturbation and step
length asymmetry would remain zero (Fig. 1D bottom). At some point in the ramp down, Δ motor
output would diverge from the perturbation and step length asymmetry would begin to grow. This
occurs when adjustments to the Δ motor output by mapping are zero, and adjustments by
recalibration remain elevated (Fig. 1D bottom, second ~half of the Ramp Down task).

 Motor results – step length asymmetry

The paradigm used for Experiment 1 is shown in Figure 2A, with the time course of step length
asymmetry (group mean ± SE) in Figure 2B. The Ramp Down task corresponds to the
manipulation illustrated in Figure 1C and is the focus of our analysis for Experiment 1. A similar
but shorter Ramp Up task was performed prior to adaptation as a control to see how participants
responded to a gradually ramped perturbation at baseline. Participants performed perceptual
tests during the Ramp Up and Ramp Down tasks, which will be discussed in a later section (see
"*Perceptual results*"). Patterns of step length asymmetry during other portions of the paradigm
was consistent with a large body of previous work (e.g., (Kristan A Leech et al., 2018; Kristan A.
Leech et al., 2018; Reisman et al., 2005; Roemmich et al., 2016; Rossi et al., 2019; Vazquez et
al., 2015)) and will not be discussed further.

Step length asymmetry during the baseline Ramp Up and post-adaptation Ramp Down tasks are
displayed in Figure 2C-D. For each speed configuration in the Ramp Down task, we statistically
compared step length asymmetry to zero. We evaluated the emergence of aftereffects by looking
for when step length asymmetry became significantly positive. We found that step length
asymmetry was *not* statistically different from zero during the first half of the Ramp Down task, for
right belt speeds ranging from 1m/s to 0.5m/s faster than the left speed (these speed
configurations are highlighted in green in Fig. 2D; all CI_{LB} (confidence intervals lower bounds) ≤ -
0.001 and CI_{UB} (upper bounds) ≥ 0.009), SI Appendix, Fig. S2 and Table S1). This result indicates
that aftereffects do not emerge immediately in the Ramp Down task, which negates the
recalibration only prediction (Fig. 2D inset). Step length asymmetry was instead significantly
positive for the second half of the task, for right belt speeds ranging from 0.45m/s to 0m/s faster
than the left speed (all CI_{LB} > 0.01). These results indicate that aftereffects emerge at a mid-point
during the Ramp Down task and align with behavioral predictions from the recalibration + mapping
hypothesis (Fig. 2D inset).

As a control, we repeated the same analysis for the baseline Ramp Up task (Fig. 2C). We found
that there was only one speed configuration for which step length asymmetry was not statistically
different from zero (right speed 0.05m/s slower than left; SL asym. = 0.006 [-0.017, 0.028], mean
[CI]). For all other configurations, step length asymmetry was significantly positive (CI_{LB} ≥ 0.048
for right speed 0.1 to 0.35m/s slower than left) or negative (CI_{UB} ≤ -0.019 for right speed 0 to
0.35m/s faster than left). This is in stark contrast to the wide range of speed configurations with

near-zero step length asymmetry observed in the Ramp Down period. Hence, the ability to walk
symmetrically at different speed configurations is not innate but dependent on the adaptation
process, corroborating the hypothesis that the "no aftereffects" region of the Ramp Down reflects
the operation of a mapping mechanism.

Motor results – perturbation and Δ motor output

To further test the competing hypotheses, we examined the perturbation and Δ motor output
during the Ramp Down (Figure 3A). As the belt speed difference decreased, so did the
perturbation component (red). In the first half of the task, the Δ motor output (blue) appeared to
match the perturbation – consistent with the lack of step length asymmetry aftereffects.
Additionally, in the second half of the task, the Δ motor output appeared larger than the
perturbation – consistent with the positive step length asymmetry aftereffects observed in Fig. 2D.

We formally contrasted predictions made by the competing hypotheses by developing
mathematical models of the Δ motor output as a function of perturbation. In the simplest
framework, the Δ motor output behavior illustrated in Figure 1D can be formalized as follows:
237

238   Forward Model Recalibration: $u(p) = r$ (1)
239

240   Stimulus-Response Mapping: $u(p) = p$ (2)
241

242   Recalibration + Mapping: $u(p) = \begin{cases} p & , & \text{for } p \geq r \\ r & , & \text{otherwise} \end{cases}$ (3)
243

Where u is the modelled Δ motor output, p is the perturbation, and r is a free parameter
representing the portion of the Δ motor output related to recalibration. The recalibration + mapping
model describes the scenario where participants modulate the Δ motor output to match the
perturbation ($u(p) = p$) for the first portion of the Ramp Down task. The Δ motor output remains
constant during the second portion of the task ($u(p) = r$) because participants are unable to
reduce the Δ motor output to values smaller than "r" (the amount achieved by recalibration).

Parameters were estimated by fitting each model to individual participants' Δ motor output data

from the Ramp Down (see Methods).

The fits of the Ramp Down Δ motor output for the recalibration only, mapping only, and

recalibration + mapping models are displayed in Figures 3B, C, and D, respectively (mean ± SE

of fits across participants; individual fits for recalibration + mapping are shown in SI Appendix,

Fig. S3). As expected, only the recalibration + mapping fit captured the matching-then-divergent

behavior of Δ motor output in response to the changing perturbation. The BIC statistic revealed

that the recalibration + mapping function explained the data better than either the recalibration

only or mapping only functions (BIC differences: 36.171 [24.868, 47.901] and 58.698 [43.708,

74.215], mean [CI]).

We additionally considered the dual state model of motor adaptation from Smith et al. (Smith et

al., 2006) as an additional framework to represent the recalibration only hypothesis (see Methods

and SI Appendix, Fig. S1B). In contrast to the simpler recalibration only model illustrated above,

the dual state model has four parameters and can account for potential forgetting or unlearning

of the Δ motor output that may occur during the Ramp Down. It can also account for the possibility

that adaptation involves two recalibration mechanisms – i.e., two distinct mechanisms that both

learn via a process of forward model recalibration and both contribute to aftereffects, but that learn

and forget at different rates. We chose this model because it is a well-established model that is

widely used to capture the Δ motor output time-course for traditional motor adaptation paradigms

(i.e., those consisting of adaptation and post-adaptation phases). Yet, performance of this model

for a manipulation like the Ramp Down task performed here has not yet been tested. As such, if

the recalibration + mapping model fit the Ramp Down data better than the dual state model, this

would provide robust evidence for the presence of a stimulus-response mapping mechanism.

We show the Δ motor output fit by the dual state model in Figure 3E (group mean ± SE; individual

fits are shown in SI Appendix, Fig. S4). Similar to its simpler recalibration only model counterpart,

the dual state model was not able to capture the matching-then-divergent behavior of Δ motor

output. The BIC statistic confirmed that the recalibration + mapping model fitted the data

significantly better than the dual state (BIC difference = 8.455 [3.557, 13.597], mean [CI]). In sum,

the modeling analysis of the Δ motor output further support the recalibration + mapping
hypothesis.

Perceptual test and results

The second goal of Experiment 1 was to evaluate the hypothesis that "perceptual realignment" (a
phenomenon leading to altered perception following adaptation) results from the operation of the
same forward model recalibration mechanism involved in adaptation of the Δ motor output (Rossi
et al., 2021a).

Previous work shows that perception realigns following gait adaptation: after adapting to a
perturbation where the right treadmill belt is *faster* than the left, return to tied belts results in
perception of the opposite asymmetry (i.e., right speed feels *slower* than left) (Jensen et al., 1998;
Vazquez et al., 2015). We therefore expected that people would perceive the belt speeds as equal
at some point in the Ramp Down and eventually perceive the opposite asymmetry on tied belts.
We measured this by asking participants to press a keyboard button at two separate occasions
during the Ramp Down task: (1) when the belts first felt equal, and (2) when they no longer felt
equal (see Methods).

Figure 4A depicts the button presses for the Ramp Down perceptual test (top panel, group mean)
overlaid onto the Ramp Down motor data (recalibration + mapping fit). Note that the task
captures a *range* of belt speed configurations that participants perceive as "equal speeds", with
button presses (1) and (2) corresponding to the upper and lower bounds of this range. This is
expected because perception is known to be noisy and may not be sensitive enough to
discriminate between belt speed configurations that are too similar.

We quantified perceptual realignment using the established measure of *point of subjective*
equality (PSE), defined as the belt speed difference perceived as "equal speeds". A PSE of zero
would indicate no perceptual realignment (accurate perception) and a PSE of 1m/s (i.e., a
magnitude equivalent to the difference between the belt speeds during adaptation) would indicate

complete perceptual realignment such that the belt speeds feel equal during the adaptation
phase.

We measured belt speed difference at the time of each button press. We computed PSE as the
range of belt speed difference values between these two measurements:

308
$$PSE_{\text{upper bound}} = [\,right\ belt\ speed - left\ belt\ speed\,]_{\text{time of button press 1}} \qquad (4)$$

310
$$PSE_{\text{lower bound}} = [\,right\ belt\ speed - left\ belt\ speed\,]_{\text{time of button press 2}} \qquad (5)$$

We found that $PSE_{\text{upper bound}}$ was 0.64±0.03m/s and $PSE_{\text{lower bound}}$ was 0.39±0.05m/s (Figure 4A,
bottom panel, group mean ± SE). This was consistent with previous work (Kristan A. Leech et al.,
2018). This perceptual realignment was not present during baseline testing, as illustrated in SI
Appendix, Fig. S5.

We aimed to evaluate the hypothesis that perceptual realignment arises from the forward model
mechanism of motor adaptation and is unaffected by the stimulus-response mapping mechanism.
This hypothesis predicts that the extent of perceptual realignment should be: 1) ~equal to the
extent of motor adaptation achieved by recalibration, and 2) less than the total extent of motor
adaptation (which also includes mapping). We quantified 1) motor adaptation by recalibration as
the fitted parameter "*r*" from the recalibration + mapping model, and 2) total motor adaptation as
the Δ motor output at adaptation plateau (Fig. 4A, right panel, see previous section "*Motor results*
– perturbation and Δ motor output"). We expressed these motor measures and perceptual
realignment as "percent compensation for the perturbation" (i.e., normalized to the perturbation
magnitude in the respective units) so that they could be compared:

$$compensation_{\text{perceptual}} = \frac{PSE}{1\,\text{m/s}} \qquad (6)$$

$$compensation_{\text{motor recalibration}} = \frac{r}{p_{\text{plateau}}} \qquad (7)$$

$$compensation_{\text{motor total}} = \frac{u_{\text{plateau}}}{p_{\text{plateau}}} \qquad (8)$$

Where r is the fitted parameter from the recalibration + mapping model, p_{plateau} is the mean

perturbation over the last 30 strides of adaptation, and u_{plateau} is the mean Δ motor output over

the last 30 strides of adaptation (for Eq. 6, note that 1m/s is the belt speed difference in

adaptation).

We show group level compensation measures in Fig. 4B. We found that

$compensation_{\text{motor recalibration}}$ (56±4%, group mean ± SE) fell within the $compensation_{\text{perceptual}}$

range (39±5% to 64±3%): it was significantly smaller than the upper bound (difference = -8 [-14,

-2]%) and significantly larger than the lower bound (difference = 18 [8, 28]%, mean [CI]). This

supports our first prediction that perceptual realignment is comparable to the extent of motor

adaptation achieved by recalibration. Furthermore, $compensation_{\text{perceptual}}$ was significantly

smaller than $compensation_{\text{motor total}}$ (95±2%, mean ± SE; difference = 31 [25, 38]% or 57 [48,

66]%, upper or lower perceptual bounds, mean [CI]). This supports our second prediction that

perceptual realignment is less than the total extent of motor adaptation.

We show individual $compensation_{\text{motor recalibration}}$ and $compensation_{\text{perceptual}}$ measures for each

participant in Figure 4C-D. We evaluated Pearson's correlation coefficients between these

measures to test whether there is a direct relationship between perceptual realignment and the

motor adaptation achieved by recalibration. We found that $compensation_{\text{motor recalibration}}$ was

significantly correlated with the upper bound of $compensation_{\text{perceptual}}$ – the value computed using

the "speeds feel equal" button press (r=0.64, p=0.002). This supports the hypothesized

relationship between motor recalibration and changes to leg speed perception. Instead,

$compensation_{\text{motor recalibration}}$ was not correlated with the lower bound of $compensation_{\text{perceptual}}$

(r=0.30, p=0.195) – this value is computed using the "right feels slower than left" button press,

suggesting the true PSE may lay closer to the first button press. Together, our results support the

hypothesis that perceptual realignment can be used as a proxy measure for the extent of motor adaptation achieved by forward model recalibration.

Control experiments

We performed six control experiments and reanalyzed previously-published data (Kristan A. Leech et al., 2018) to replicate the findings of Experiment 1 across different paradigm conditions (SI Appendix, Control experiments, Fig. S6-8, and Table S2-6). This provided additional support for the recalibration + mapping hypothesis. We first checked that the motor and perceptual behaviors observed in the Ramp Down could be replicated using a different method of assessment. A "speed match" task was given where participants use a joystick to increase the speed of the right belt (initially stationary) until they feel it matches the left belt (SI Appendix, Fig. S6B, "Ascend") (Kristan A. Leech et al., 2018; Rossi et al., 2019; Statton et al., 2018; Vazquez et al., 2015). This approximates the second half of the Ramp Down (Fig. 4A, right leg feels equal or slower than left). Consistent with this, we observed step length asymmetry aftereffects early in the speed match task (initial SL asym. = 0.433 [0.271, 0.612], mean [CI]). Asymmetry decreased and was eventually ~0 when the belt speeds felt equal (final SL asym. = 0.002 [-0.040, 0.045]; SI Appendix, Fig. S7B).

We also used a "descend" speed match task where participants *decreased* the speed of the right belt (initially fast as in adaptation) until they felt it matched the left (SI Appendix, Fig. S6B). This approximated the first half of the Ramp Down. As expected, step length asymmetry was close to zero for entire task (initial: -0.017 [-0.069, 0.039], final: 0.043 [-0.0004, 0.090], mean [CI]; SI Appendix, Fig. S7B). The PSE for both speed match groups was comparable to that of Experiment 1 ($compensation_{\mathrm{perceptual}}$ difference between Ramp Down lower bound and ascend or descend Control = -3 [-16, 10]% or 4 [-12, 18]%, mean [CI]), and smaller than motor adaptation ($compensation_{\mathrm{motor\ total}} - compensation_{\mathrm{perceptual}}$ = 48 [39, 57]% or 56 [42, 70]% for ascend or descend, mean [CI]; SI Appendix, Fig. S7A).

We replicated our findings in additional speed match experiments that varied in adaptation duration (3, 15, or 30 minutes), perturbation magnitude (1m/s or 0.4m/s speed difference) and schedule (abrupt or gradual) (SI Appendix, Control experiments and Fig. S6). Regardless of the

paradigm, the step length asymmetry and PSE were consistent with the recalibration + mapping

hypothesis (SI Appendix, Fig. S7 and Table S2-3).

<u>Experiment 2</u>

In Experiment 2, we considered two hypotheses for how stimulus-response mapping may

operate, that could both explain the results of Ramp Down task of Experiment 1. First, the

mechanism may be a memory-based mapping that caches prior Δ motor outputs in memory, and

later retrieves them in response to a matching perturbation (Huberdeau et al., 2019; McDougle

and Taylor, 2019; Poggio and Bizzi, 2004; Wolpert et al., 2001). In adaptation, the Δ motor output

changes gradually from zero to a value matching the perturbation (see Fig. 1B), and all

intermediate values may be stored in memory. Therefore, a memory-based mapping can produce

Δ motor outputs that match perturbations *smaller* than the adaptation perturbation (Kristan A

Leech et al., 2018) – such as those of the Ramp Down task of Experiment 1. However, a memory-

based mapping would not be able to account for perturbations *larger* than the adaptation

perturbation: the largest Δ motor output available in memory would still undershoot such

perturbations, resulting in negative step length asymmetry.

Alternatively, the mechanism may be a structure-based mapping that learns the algorithm to

compute Δ motor outputs from sensory stimuli (Bond and Taylor, 2017; Braun et al., 2009;

McDougle and Taylor, 2019; Wolpert et al., 2001). In adaptation, the mechanism may learn the

relationship between perturbation and appropriate Δ motor output, and later use it to generate Δ

motor outputs anew. Therefore, a structure-based mechanism can also *extrapolate* Δ motor

outputs beyond what was experienced in adaptation, accounting for perturbations *larger* than the

adaptation perturbation.

Thus, we tested a Ramp Up & Down condition where the speed of the right belt was both gradually

increased and then decreased after adaptation (Fig. 5A). The memory-based hypothesis predicts

that step length asymmetry will become negative for perturbations *larger* than adaptation, while

the structure-based hypothesis predicts it will remain close to zero (Fig. 5A inset, "predictions").

The rest of the paradigm was analogous to that of Experiment 1 except there was no perceptual
assessment.

Figure 5B shows step length asymmetry throughout the paradigm, and Figure 5C shows a close-
up of the Ramp Up & Down task performance (group mean ± SE). On average, participants' step
length asymmetry patterns did not remain zero for speed differences larger than adaptation (teal).
However, we observed that individual participants exhibited markedly different patterns of step
length asymmetry during this phase (SI Appendix, Fig. S9). We quantified this observation by
evaluating, for each participant, the number of strides in this phase with step length asymmetry
below their own baseline CI (SI Appendix, Table S7). We used a density-based analysis to
formally assess whether there were separate clusters in our data (see Methods). Indeed, the
algorithm detected two separate clusters of participants: for 12 participants, between 38 to 60
strides were asymmetric (out of 60 total strides); for the other 8 participants, only 3 to 21 strides
were asymmetric (Fig. 5D-E insets, and SI Appendix Fig. S10; difference in strides between
subgroups = 36.083 [29.917, 42.250], mean [CI]). As a control, we performed the same clustering
analysis for Experiment 1, and did not find separate clusters for any of our measures of interest
(SI Appendix, Experiment 1 clustering analysis, Fig. S10-11, and Table S8-9).

This result indicates that 12 of 20 participants could not account for belt speed differences larger
than that of adaptation, suggesting that they used a memory-based mapping mechanism (Fig.
5D). In contrast, 8 of 20 participants could account for these speeds, suggesting that they
engaged a structure-based mapping mechanism (Fig. 5E).

We next aimed to exclude the possibility that participants in the structure-based subgroup may
simply be faster at adapting to new perturbations than those in the memory-based subgroup, and
may be adapting to the new perturbations of the Ramp Up and Down rather than generating Δ
motor output using a previously learnt structure. To this end, we evaluated learning rates during
adaptation. We found that participants in the two subgroups adapted at similar rates (strides to
plateau difference, structure – memory: 135.875 [-53.208, 329.708], mean [CI]; SI Appendix, Fig.
S12-13), confirming that different pattern of step length asymmetry in the Ramp Up and Down
task reflects different types of mapping in the subgroups.

We considered that the mapping adjustments described here may or may not be *deliberate* (i.e.,
participants are trying to correct for the perturbation) and done with an *explicit strategy* (i.e.,

participants can accurately report a relevant strategy that would counter the perturbation) (Long
et al., 2016; Roemmich et al., 2016). An example of an explicit strategy in visuomotor reaching
adaptation is when participants report that they aimed to offset a visual rotation (Bond and Taylor,
2017; Taylor et al., 2014); conceivable examples in split-belt treadmill adaptation may be reports
of "taking steps of similar length", "stepping further ahead with the right foot" or "standing on the
left foot for longer". We tested whether participants could explicitly report changes to the gait
pattern that specifically correct for the split-belt perturbation.

At the end of the experiment, participants were asked to report (in writing) if/how they had changed
the way they walked during adaptation (note that this was a later addition to the protocol and was
only collected in 16 of the 20 participants). We assessed reports by categorizing them in three
steps: 1) did the report mention *any* deliberate changes? 2) were the changes *relevant* to
adaptation? (i.e., did the report mention any gait metric contributing to the overall Δ motor output
or step length asymmetry adaptation in any amount?) 3) was the response *accurate?,* as
participants often reported strategies that they did not actually execute.

We summarize results from the questionnaire in Figure 6 (original responses are reported in SI
Appendix, Table S10). We found that, while 13 participants reported *deliberate* changes, only 6
people mentioned *relevant* aspects of the walking pattern (in particular, participants mentioned
"limping" or temporal coordination). Furthermore, only one of these participants reported an
accurate gait parameter ("I tried to spend as much time leaning on my left leg as possible"), while
the remainder of the relevant responses were inaccurate (e.g., "matched duration of standing on
each foot") or vague (e.g., "I adjusted as if I was limping"). The aspect of gait most frequently
reported across participants was stability (also reported as balance, not falling, or controlling
sway). This suggests that, while participants may deliberately adjust their body to feel more stable,
they do not seem to explicitly strategize how to offset the perturbation. Thus, mapping tends to
adjust aspects of the walking pattern that participants are not explicitly aware of controlling,
suggesting that this differs from explicit strategies often observed in reaching.

In sum, Experiment 2 suggests that the mapping mechanism involved in split-belt adaptation is
memory-based in some people and structure-based in others. Neither mechanism seems to rely
on explicit strategies.

Discussion

In this study, we showed that locomotor adaptation involves two learning mechanisms: forward model recalibration and a stimulus-response mapping mechanism. The recalibration mechanism changes movement and relates to the perceptual changes observed in locomotor adaptation; the mapping mechanism is flexible and, once learned, can change our motor output immediately to account for a range of belt speed configurations. Our data suggest it can be memory-based or structure-based. We propose a comprehensive conceptual model where the same output from the forward model (Fig. 7, light blue) may project to both sensory integration and motor control areas (green and blue boxes), where it may be used to recalibrate perception and movement respectively. We also propose that the same *recalibrated perception* (green arrow) serves both as signal for conscious perception and as an input to the mapping mechanism (dark blue box), where it is used to compute Δ motor output adjustments that will be summed to the recalibration-related adjustments (Δ motor output, blue box).

Forward model recalibration of movement and perception

Our results support the hypothesis that changes in movement perception with adaptation are also mediated by the recalibration of a forward model ('t Hart and Henriques, 2016; Izawa et al., 2012; Sombric et al., 2019; Synofzik et al., 2008; Yavari et al., 2016) (for a review, see (Rossi et al., 2021a)). In line with a variety of previous studies (e.g., (Haith et al., 2008; Harris, 1963; Jensen et al., 1998; Moidell and Bedell, 1988; Sombric et al., 2019)), we find that people recalibrate their movement perception in adaptation and feel less perturbed than they really are – here, they feel that their right leg moves slower than its true speed (Jensen et al., 1998; Kristan A. Leech et al., 2018; Rossi et al., 2019; Statton et al., 2018; Vazquez et al., 2015). Consistent with the operation of a forward model thought to predict sensory consequences from efference copies (Haruno et al., 2001; Ito, 1989; Wolpert et al., 2001), perceptual changes are expressed during active and not passive movements (Sombric et al., 2019; Vazquez et al., 2015). Perceptual changes with adaptation are consistently affected by damage to (Izawa et al., 2012; Statton et al., 2018; Synofzik et al., 2008) or stimulation of the cerebellum (Yavari et al., 2016), the structure imputed with forward models. We suggest that perceptual changes may reflect cerebellar recalibration (Fig. 7, "recalibration", light blue) processed downstream by cortical sensory areas ("perception", green box). Indeed, activity in both cerebellum and parietal lobe is shown to increase with split-belt walking adaptation (Hinton et al., 2019).

We suggest participants may filter out the portion of the belt speed perturbation that is predictively

accounted for by the forward model, leading to the perceptual changes observed here (Fig. 7,

green box: the forward model recalibration, x_r, is cancelled from the speed difference sensory

signal, p, to produce the perceived speed difference, $\tilde{p}$). The parietal lobe receives input from the

cerebellum (Judd et al., 2021; Sultan et al., 2012) and has the capacity to selectively cancel

sensory prediction signals (Blakemore et al., 1998; Brooks and Cullen, 2019). This is important

for filtering out self-generated sensory stimuli from those externally produced (Blakemore et al.,

1998; Brooks and Cullen, 2019; Rondi-Reig et al., 2014), or predictable environmental stimuli

from those that are novel (Anderson et al., 2012; Rondi-Reig et al., 2014). While we think this is

a parsimonious explanation, cerebellar predictions may also be integrated with and sharpen

proprioceptive estimates (Bhanpuri et al., 2013; Weeks et al., 2017), raising the possibility that a

more complex integration of predicted and actual sensory signals contribute to perceptual

changes.

We demonstrated that perceptual and motor recalibrations are directly related. Motor recalibration

is approximately constant when the belt speed difference is changed after adaptation, as captured

by the parameter "*r*". We find "*r*" is correlated with the "belt speeds feel equal" button press,

suggesting that output from the same forward model (Fig. 7, x_r, light blue) may be used to change

both movement (u, blue) and perception ($\tilde{p}$, green) during adaptation (SI Appendix,

Supplementary Discussion 1). There is evidence to suggest that single regions of the cerebellar

cortex (Pisano et al., 2021) and possibly single output cells from the cerebellar nuclei (Judd et al.,

2021; Sultan et al., 2012) may project downstream to both motor and sensory cortical areas.

Furthermore, the cerebellum has been shown to modulate activity in sensory and motor cortices

in a coordinated manner (Lindeman et al., 2020; Popa et al., 2013), and both regions show

evidence of sensory cancellation during active movement (Seki and Fetz, 2012). Hence, it is

possible that the motor and perceptual recalibrations observed here reflect the differential

processing of cerebellar output (Fig. 7, "recalibration", light blue) by motor and sensory (blue and

green, respectively) cortical areas.

Automatic yet flexible stimulus-response mapping

We identified a stimulus-response mapping mechanism that does not conform with currently

known mechanisms of walking adaptation. When an explicit goal is given, adaptation mechanisms

deployed in addition to forward model recalibration are consistently found to operate under

deliberate control (Codol et al., 2018; Taylor et al., 2014; Taylor and Ivry, 2011) – including

memory-based caching (Huberdeau et al., 2019; McDougle and Taylor, 2019) and structural

learning mechanisms (Bond and Taylor, 2017; McDougle and Taylor, 2019). Mounting evidence

indicates that explicit strategies do not play a strong role in split-belt walking adaptation where an

explicit movement goal is not provided to the participants; for example, people do not adapt faster

even after watching someone else adapt (Song et al., 2020). Strategic adjustments to the walking

pattern can be temporarily elicited by providing additional visual feedback of the legs and an

explicit goal of how to step. Yet, these adjustments disappear immediately upon removal of the

visual feedback (Roemmich et al., 2016), and have no effect on the underlying adaptive learning

process (Long et al., 2016; Malone and Bastian, 2010; Roemmich et al., 2016). Furthermore,

performing a secondary cognitive task during walking adaptation does not affect the amount of

motor learning (Hinton et al., 2020; Malone and Bastian, 2010; Rossi et al., 2021b; Vervoort et

al., 2019). Here, we confirmed that explicit strategies are not systematically used to adapt step

length asymmetry and Δ motor output: only one person accurately reported making relevant

changes appropriate to walk with symmetry; the rest either did not know what they did, reported

changes that did not actually occur or would not lead symmetry (SI Appendix, Supplementary

Discussion 2).

Despite not relying on explicit strategies, we find that the mapping mechanism scales down

immediately to account for the amount of perturbation not corrected by recalibration. To the best

of our knowledge, this is the first study to show that such flexible mechanism plays a role in

walking adaptation. Previous work from our group used a manipulation that involved ramping

down the speed of the right belt after adaptation (Roemmich and Bastian, 2015), similar to the

Ramp Down used here. However, this was done slowly over 10 minutes, so that when participants

walked with near-zero step length asymmetry for a portion of the ramp, this was attributed to them

having unlearnt the adapted pattern. We here eliminated this confounder by ramping down the

right speed quickly – on average, the ramp down took 1 minute and 20 seconds. Importantly, this

time scale was short enough that participants' aftereffect at the end of the ramp down was

comparable to that observed in studies without the ramp down (e.g., (Reisman et al., 2005)). This

manipulation allowed us to discover a sharp change in Δ motor output behavior half-way through

the ramp down that supports the presence of a mapping mechanism. That is, the Δ motor output

can rapidly change and match the perturbation but only for speed differences that are ~50-100%

the adaptation speed difference, remaining mostly constant for smaller speed differences. Hence,

the flexible behavior of the Δ motor output cannot be explained by the unlearning process, and it

is instead indicative of the operation of a mapping mechanism.

As such, the mapping mechanism combines the advantages of automaticity and flexibility

(Huberdeau et al., 2015; Taylor et al., 2014; Taylor and Ivry, 2011). This is ecologically important

for both movement accuracy (Uiga et al., 2020; Wong et al., 2008) and for walking safely in real-

world situations, where we walk while talking or doing other tasks, and terrains are uneven (Clark,

2015; Paul et al., 2005; Wong et al., 2008) (SI Appendix, Supplementary Discussion 3).

Recalibrated perception maps to Δ motor output

The flexible properties of the mapping mechanism indicate that its adjustments to Δ motor output

(Fig. 7, x_m, dark blue) are computed from the *recalibrated* perception of speed difference ($\tilde{p}$).

During the Ramp Down task (Figs. 3-4), mapping-related adjustment *complements* recalibration-

related adjustment to produce the overall Δ motor output that *tracks* the perturbation. This

indicates that mapping-related adjustment is computed as *total* belt speed difference perturbation

minus recalibration-related adjustment, which corresponds to the *perceived* perturbation ($x_m = $

$\tilde{p} = p - x_r$). As further evidence, we observe that the sharp "corner" in the Δ motor output data of

the Ramp Down occurs when the *perceived,* not actual, perturbation changes sign (Fig. 4,

mapping stops complementing recalibration and remains zero when the right leg *feels* slower than

the left even though it is *actually* faster). Features of the mapping mechanism are further

discussed in SI Appendix, Supplementary Discussion 4.

We propose the following series of computations (Fig. 7). First, the recalibration is used to

compute the perceived perturbation (green box). Second, the perceived perturbation is relayed to

the mapping mechanism and used to compute the mapping-related motor adjustment (dark blue

box). Third, recalibration-related and mapping-related motor adjustments are summed to produce

the Δ motor output (blue box). Interconnectivity between sensory and motor cortical areas, and

the cerebellum (Judd et al., 2021; Kandel et al., 2013; Sultan et al., 2012), is consistent with the

architecture proposed here. In contrast to the fixed adjustments typically attributed to predictive

models in the cerebellum, flexible control of the walking pattern has traditionally been associated

with cortical motor areas (Drew and Marigold, 2015; Reisman et al., 2010). Cortical involvement

appears to go beyond voluntary control (Delval et al., 2020; Petersen et al., 2012) and may

specifically contribute to split-belt adaptation (Hinton et al., 2019). More generally, cortical motor

areas receive extensive sensory input from primary and association areas of the parietal lobe

(Kandel et al., 2013), and use it to guide movement in a variety of sensorimotor mappings tasks

(Dassonville et al., 2001; Georgopoulos, 2000). As such, we suggest cortical motor areas are the

ideal substrate to receive perceptual signals (such as the perceived perturbation; Fig. 7, $\tilde{p}$, green)

and map it to motor adjustments ($x_m = \tilde{p}$, "mapping" dark blue box).

Finally, our framework suggests that mapping-related motor adjustments are combined with

forward model recalibration downstream ($u = x_m + x_r$, "Δ motor output" blue box). Signals may

be combined at the cortical level, as suggested by cortical correlates of adaptation in monkeys

that resemble the combined operation of recalibration-like and mapping-like mechanisms (Chase

et al., 2012; Sun et al., 2022) (SI Appendix, Supplementary Discussion 5). Alternatively, signals

may be combined subcortically, as the cerebellum modulates movement through both cortical

and subcortical projections (Kandel et al., 2013), and cerebral damage does not block walking

adaptation (Choi et al., 2009; Reisman et al., 2007).

Mapping operates as memory-based in some people, structure-based in others

Results from Experiment 2 highlight individual differences in the learning mechanisms underlying

generalization to unexperienced belt speed differences (usage of the term "generalization" is

discussed in SI Appendix, Supplementary Discussion 6). The generalization to novel perturbation

sizes observed here is in line with previous suggestions of "meta-learning" in the savings of

walking adaptation (i.e., faster relearning when exposed to a different perturbation) (Kristan A

Leech et al., 2018; Malone et al., 2011). Generalization to *larger* perturbations after reaching

adaptation was shown to be incomplete (Abeele and Bock, 2001; Lazar and Van Laer, 1968), and

it was unclear whether the Δ motor output had been *extrapolated* beyond what had been

experienced, or rather was just the same as in adaptation. Surprisingly, we found 40%-60% divide

in our participants regarding the capacity to extrapolate Δ motor output to account for larger

perturbations.

We suggest that structural learning may underly the ability to extrapolate the Δ motor output and

walk symmetrically for belt speed differences larger than adaptation, as seen in 8 of 20

participants. Indeed, generalization in reaching adaptation may rely on a process of structural

learning (Bond and Taylor, 2017; Braun et al., 2009) (note however that this was tied to explicit

aiming, (Bond and Taylor, 2017)). Our participants may have learned the scaling relationship

between belt speed perturbation and Δ motor output needed to walk symmetrically and use this

scaling to produce new Δ motor outputs matching larger perturbations. In contrast, memory-based

theories of mapping (Dassonville et al., 2001; Poggio and Bizzi, 2004; Van Vugt and Ostry, 2018;

Wolpert et al., 2001) may explain why 12 participants walked asymmetrically for perturbations

larger than adaptation, despite generalizing to smaller perturbations. Here, Δ motor outputs may

be stored in memory during adaptation in association with the amount of perturbation they correct,

and then retrieved during the ramp tasks. People transition through a range of Δ motor outputs in

adaptation (see gradual transition from baseline to adapted output in Fig. 1B), so storage of

intermediate outputs may reflect generalization to smaller perturbations (while larger Δ motor

outputs are not experienced). SI Appendix, Supplementary Discussion 7, discusses potential

interpolation computations.

In sum, consistent with findings that reaching adaptation strategies can be memory or structure-

based (McDougle and Taylor, 2019), we show that walking adaptation can involve either memory-

based or structure-based mappings.

Conclusions and future directions

We here characterized two distinct learning mechanisms involved in walking adaptation, both of

which are not under explicit control: recalibration - which leads to motor and perceptual

aftereffects, and mapping – which only changes movement, does not contribute to aftereffects,

and shows meta-learning to different perturbation sizes. Future work should explore whether the

mapping mechanism described here could be of clinical significance: given the combined flexibility

and automaticity, mapping has the potential to ameliorate known problems such as transitioning

between walking environments (Sombric et al., 2017) and dual-tasking (Clark, 2015).

Furthermore, our findings that different people adapt using different learning mechanisms (i.e.,

memory or structure-based mapping) highlights the importance of assessing individual

characteristics of adaptation for both understanding of the neural mechanisms as well as

translation to rehabilitation. Finally, as detailed in SI Appendix, Supplementary Discussions 8-10,

the framework proposed here has important implications for the development of computational

models of adaptation, and may help reconcile different findings related to savings, energetics,

and sources of errors.

Materials and Methods

<u>Participants</u>

We recruited one-hundred adults (66 females, 23.6 ± 3.9 years old - mean ± SD) for this study, and we reanalyzed data from ten additional adults collected in a previous study (Kristan A. Leech et al., 2018) (9 female, 21.3 ± 2.9 years old). The protocol was approved by the Johns Hopkins Institutional Review Board and participants provided written informed consent. Participants had no known neurological or musculoskeletal disorders, were naïve to split-belt walking, and participated in only one of the nine experiments.

<u>Data collection</u>

Participants walked on a split-belt treadmill (Woodway, Waukesha, WI, USA) and belt speeds were controlled using a custom Vizard program (WorldViz). Kinematic data were collected using infrared-emitting markers (Optotrak, Northern Digital, Waterloo, ON, Canada) at 100Hz (SI Appendix, Supplementary Methods).

<u>Ramp tasks</u>

In the Ramp Up, Ramp Down, and Ramp Up and Down tasks of Experiments 1 and 2, we changed the speed of the right belt by 0.05m/s every 3 strides during right leg swing. The Ramp Up consisted of 7 increasing speeds from 0.35m/s to 0.65m/s. The Ramp Down consisted of 21 decreasing speeds from 1.5m/s to 0.5m/s. The Ramp Up and Down consisted of 41 total speeds: 11 increasing speeds from 1.5m/s to 2m/s, followed by 30 decreasing speeds from 1.95m/s to 0.5m/s. The left belt speed was constant at 0.5 m/s for all ramp tasks. The treadmill was not stopped before the ramp tasks (people transitioned directly from the preceding walking blocks into the ramp) but was briefly stopped after each ramp task. We occluded vision and sound of the belt speeds using a cloth drape and headphones playing white noise.

In Experiment 1, a keyboard was placed on the treadmill handrail and participants were asked to press a button the first time they perceived the right belt to be (1) as fast as the left, and (2) faster/slower than the left (for the Ramp Up and Ramp Down, respectively).

<u>Questionnaire</u>

At the end of Experiment 2, participants answered this question on a computer: *Did you*
deliberately change how you walked to account for how fast the belts were moving? If so, describe
how. Note: deliberately means that you thought about and decided to move that way. The question
refers to the entire central ~20min walking block. To ensure participants remembered what phase
they were asked about, before adaptation we told participants that the following block will be
called "central ~20min walking block".

<u>Motor measures</u>

We defined a stride as the period between two consecutive left heel strikes (LHS_1 to LHS_2), and
computed the motor measures for each stride as in previous work (Finley et al., 2015; Sombric et
al., 2017):

693
$$step\ length\ asymmetry\ =\ \frac{1}{stride\ length}*\Delta step\ length \tag{9}$$

695
$$perturbation\ =\ \frac{1}{stride\ length}*\overline{step\ time}*\Delta step\ velocity \tag{10}$$

697
$$\Delta\ motor\ output\ =\ \frac{1}{stride\ length}*\left(\Delta step\ position\ -\ \overline{step\ velocity}*\Delta step\ time\right) \tag{11}$$

699 The terms in the equations above are defined as follows:

701
$$\Delta step\ length\ =\ R\ step\ length\ -\ L\ step\ length \tag{12}$$

703
$$stride\ length\ =\ R\ step\ length\ +\ L\ step\ length \tag{13}$$

705 Where right (R) and left (L) step lengths are the anterior-posterior distance between the ankle
706 markers of the two legs at right heel strike (RHS) and left heel strike (LHS_2) respectively.

$$\Delta step\ position\ = \left(R\ step\ position_{RHS} - L\ step\ position_{LHS_1}\right)$$
$$- \left(L\ step\ position_{LHS_2} - R\ step\ position_{RHS}\right) \qquad (14)$$

Where step position is the anterior-posterior position of the ankle marker, relative to average of the two hip markers, at heel strike of the same leg.

$$\Delta step\ time\ = R\ step\ time - L\ step\ time \qquad (15)$$

$$\overline{step\ time}\ = \frac{R\ step\ time + L\ step\ time}{2} \qquad (16)$$

Where left and right step times are the times from LHS_1 to RHS, and from RHS to LHS_2 respectively.

$$\Delta step\ velocity\ = R\ step\ velocity - L\ step\ velocity \qquad (17)$$

$$\overline{step\ velocity}\ = \frac{R\ step\ velocity + L\ step\ velocity}{2} \qquad (18)$$

Where step velocity is the average anterior-posterior velocity of the ankle marker relative to average of the two hip markers, over the duration of the step (left: LHS_1 to RHS, right: RHS to LHS_2).

In Experiment 2, we also computed the measure of "strides to plateau" using individual step length asymmetry data from the adaptation phase only. We first smoothed the data with a 5-point moving average filter. We then calculated the number of strides till five consecutive strides fell within the mean ± 1SD of the last 30 strides (Malone and Bastian, 2010; Rossi et al., 2019).

 Clustering analysis

 We tested to see if there were separate clusters of participants in our dataset. For Experiment 1,

 the measure used for clustering was the number of strides in the Ramp Down with step length

 asymmetry above the baseline CI. For Experiment 2, it was the number of strides in the first

 portion of the Ramp Up and Down (first 60 strides) with step length asymmetry below the baseline

 CI. For both experiments, "baseline CI" was computed separately for each participant as the 95%

 CI of the mean of step length asymmetry data in the second baseline tied-belt block (after the

 Ramp Up; see Statistical Analysis).

 We used the MATLAB "dbscan" density-based clustering algorithm (Ester et al., 1996) and

 automated the choice of parameters adapting previously published procedures (Naik Gaonkar

 and Sawant, 2013; Rahmah and Sitanggang, 2016) (SI Appendix, Supplementary Methods and

 pseudocode). The resulting algorithm does not require any user input; it automatically assigns

 participants to clusters (whose number is not predefined), or labels them as outliers, based solely

 on each participant's measure of interest.

 Data fitting

 In Experiment 1, we fitted the recalibration only (Eq. 1), mapping only (Eq. 2), recalibration +

 mapping (Eq. 3) models to individual participants' Δ motor output data in the Ramp Down. A

 reasonable upper limit for the parameter r would be the magnitude of the perturbation in

 adaptation, but this motor measure differs across participants, which would lead to

 inconsistencies in the data fitting analysis. We therefore reformulated the recalibration and

 recalibration + mapping models to use a normalized parameter r_{norm}:

 $$r = r_{norm} * p_{plateau} \tag{19}$$

 Forward Model Recalibration: $u(p) = r_{norm} * p_{plateau}$ (20)

 Recalibration + mapping: $u(p) = \begin{cases} p & , \text{ for } p \geq r_{norm} * p_{plateau} \\ r_{norm} * p_{plateau} , & \text{otherwise} \end{cases}$ (21)

Where p_{plateau} is mean perturbation over the last 30 strides of adaptation, computed separately

for each participant, p is the perturbation in the Ramp Down, and u is the modelled Δ motor

output. We fitted the normalized parameter r_{norm} using initial parameter value = 0.5 and bounds

= [0, 1] for all participants. We then used Eq. 19 to compute the unnormalized value r.

We additionally fitted the dual state model defined as in Smith et al. (Smith et al., 2006):

Δ motor output: $\quad u(i) = u_f(i) + u_s(i)$ (22)

fast state: $\quad u_f(i+1) = A_f * u_f(i) + B_f * \left(p(i) - u(i)\right)$ (23)

slow state: $\quad u_s(i+1) = A_s * u_s(i) + B_s * \left(p(i) - u(i)\right)$ (24)

Where p is the perturbation, i is the stride in the task (0 to 62), u is the fitted Δ motor output, u_f

and u_s are the contribution of the fast and slow states to changing the Δ motor output, and A_f, A_s,

B_f, B_s are free parameters representing forgetting and learning rates, with $0<A_f<A_s<1$ and

$0<B_f>B_s<1$. We set the initial values as A_f=0.92, A_s=0.99, B_f=0.1, and B_s=0.01 (as in (Roemmich

et al., 2016)).

We used the *fmincon* MATLAB function to find the parameters of each model that minimize the

residual sum of squares between data and model fits. We tightened constraint and optimality

tolerances to 10^-20 (as in (Roemmich et al., 2016)). For the dual state model, we fitted the model

to both adaptation and Ramp Down data (this was necessary as the initial value of the fast and

slow states in the Ramp Down was otherwise unknown), but computed the residual sum of

squares solely on the Ramp Down (ensuring fair model comparison).

<u>Statistical analysis</u>

Statistical analyses were performed in MATLAB with significance level $\alpha = 0.05$ corrected for

multiple comparisons using the False Discovery Rate procedure (FDR) (Benjamini and Yekutieli,

2005) (corrected significance level $\alpha_{\text{corr}} = \frac{\alpha R}{m}$, with m = number of total tests for that comparison

group, reported below, and R = number of significant tests). We used a bootstrap procedure
(10,000 samples drawn with replacement) (DiCiccio and Efron, 1996; Efron and Tibshirani, 1994)
to generate confidence intervals (CI) for the following measures:

• Group mean step length asymmetry for each speed in the Ramp Up (m=7) and Ramp
Down (m=21) tasks of Experiment 1 (pre-averaged within-participant over the 3 strides
taken at that speed).

• Group mean value for the BIC difference between the recalibration + mapping model fit,
and each of the other model fits (dual state, recalibration only, mapping only) of
Experiment 1 (m=3).

• Group mean value for the difference between $compensation_{\text{motor total}}$ or
$compensation_{\text{motor recalibration}}$, minus each $compensation_{\text{perceptual}}$ (Experiment 1, m=4).

• Difference of the means between Structure and Memory subgroups (Experiment 2) for the
measures of strides to plateau (m=1), and number of asymmetric strides in Ramp Up and
Down (teal portion; m=1)

• Individual means for baseline step length asymmetry (2-min block) for each participant in
Experiment 1 and 2 (all m=1).

We report FDR-corrected CIs and consider statistical tests significant if the CI does not overlap
zero (Cumming and Finch, 2005; DiCiccio and Efron, 1996; Efron and Tibshirani, 1994).

Data Availability

Note to reviewers: the dataset cited below is currently "private for peer review", and can be
accessed by the alternative "reviewer URL" provided by the Dryad database:

https://datadryad.org/stash/share/Qa0pD7rMFT7kVgDeP-8PMYKBaCNuy4yx1VGIQIpVgvM

All data and code used for the study have been deposited in: Rossi, Cristina (2023), Dataset for
automatic learning mechanisms for flexible human locomotion, Dryad, Dataset,
https://doi.org/10.5061/dryad.18931zd27.

Acknowledgments

Supported by NIH grant 5 R37 NS090610 to AJB, American Heart Association predoctoral

fellowship 20PRE35180131 to CR, and NIH grant K01 AG073467 to KAL. We thank Daniel

Wolpert, Adrian Haith, Jonathan Tsay, and Amanda Therrien for helpful discussions and

comments on data analysis.

References
818
't Hart BM, Henriques DYP. 2016. Separating predicted and perceived sensory consequences of motor
learning. *PLoS One* **11**:e0163556. doi:10.1371/journal.pone.0163556
Abeele S, Bock O. 2001. Mechanisms for sensorimotor adaptation to rotated visual input. *Exp Brain Res*
139:248–253. doi:10.1007/S002210100768
Anderson SR, Porrill J, Pearson MJ, Pipe AG, Prescott TJ, Dean P. 2012. An Internal Model Architecture
for Novelty Detection: Implications for Cerebellar and Collicular Roles in Sensory Processing. *PLoS*
One **7**:e44560. doi:10.1371/journal.pone.0044560
Bastian AJ. 2008. Understanding sensorimotor adaptation and learning for rehabilitation. *Curr Opin*
Neurol **21**:628–633. doi:10.1097/WCO.0b013e328315a293
Benjamini Y, Yekutieli D. 2005. False discovery rate-adjusted multiple confidence intervals for selected
parameters. *J Am Stat Assoc* **100**:71–81. doi:10.1198/016214504000001907
Bhanpuri NH, Okamura AM, Bastian AJ. 2013. Predictive modeling by the cerebellum improves
proprioception. *J Neurosci* **33**:14301–14306. doi:10.1523/JNEUROSCI.0784-13.2013
Blakemore S-J, Wolpert DM, Frith CD. 1998. Central cancellation of self-produced tickle sensation. *Nat*
Neurosci **1**:635–640.
Bond KM, Taylor JA. 2017. Structural learning in a visuomotor adaptation task is explicitly accessible.
eNeuro **4**:e0122-17. doi:10.1523/ENEURO.0122-17.2017
Braun DA, Aertsen A, Wolpert DM, Mehring C. 2009. Motor Task Variation Induces Structural Learning.
Curr Biol **19**:352–7. doi:10.1016/j.cub.2009.01.036
Brooks JX, Cullen KE. 2019. Predictive Sensing: The Role of Motor Signals in Sensory Processing. *Biol*
psychiatry Cogn Neurosci neuroimaging **4**:842. doi:10.1016/J.BPSC.2019.06.003
Chase SM, Kass RE, Schwartz AB. 2012. Behavioral and neural correlates of visuomotor adaptation
observed through a brain-computer interface in primary motor cortex. *J Neurophysiol* **108**:624–644.
doi:10.1152/jn.00371.2011
Choi JT, Vining EPG, Reisman DS, Bastian AJ. 2009. Walking flexibility after hemispherectomy: Split-belt
treadmill adaptation and feedback control. *Brain* **132**:722–733. doi:10.1093/brain/awn333
Clark DJ. 2015. Automaticity of walking: Functional significance, mechanisms, measurement and
rehabilitation strategies. *Front Hum Neurosci* **9**:246. doi:10.3389/fnhum.2015.00246
Codol O, Holland PJ, Galea JM. 2018. The relationship between reinforcement and explicit control during
visuomotor adaptation. *Sci Rep* **8**:9121. doi:10.1038/s41598-018-27378-1
Cumming G, Finch S. 2005. Inference by eye: confidence intervals and how to read pictures of data. *Am*
Psychol. doi:10.1037/0003-066X.60.2.170
Dassonville P, Lewis SM, Zhu XH, Ugurbil K, Kim SG, Ashe J. 2001. The effect of stimulus-response
compatibility on cortical motor activation. *Neuroimage* **13**:1–14. doi:10.1006/nimg.2000.0671
Delval A, Bayot M, Defebvre L, Dujardin K. 2020. Cortical oscillations during gait: Wouldn't walking be so
automatic? *Brain Sci*. doi:10.3390/brainsci10020090
DiCiccio TJ, Efron B. 1996. Bootstrap Confidence Intervals. *Stat Sci* **11**:189–228. doi:10.1007/978-3-319-
01931-4_24
Drew T, Marigold DS. 2015. Taking the next step: Cortical contributions to the control of locomotion. *Curr*

Opin Neurobiol **33**:25–33. doi:10.1016/j.conb.2015.01.011

Efron B, Tibshirani RJ. 1994. An Introduction to the Bootstrap, An Introduction to the Bootstrap.
doi:10.1201/9780429246593

Ester M, Kriegel H-P, Sander J, Xu X. 1996. A Density-Based Algorithm for Discovering Clusters in Large
Spatial Databases with NoiseProceedings of the 2nd International Conference on Knowledge
Discovery and Data Mining. pp. 226–231.

Finley JM, Long A, Bastian AJ, Torres-Oviedo G. 2015. Spatial and Temporal Control Contribute to Step
Length Asymmetry during Split-Belt Adaptation and Hemiparetic Gait. *Neurorehabil Neural Repair*
29:786–795. doi:10.1177/1545968314567149

Georgopoulos AP. 2000. Neural aspects of cognitive motor control. *Curr Opin Neurobiol* **10**:238–241.
doi:10.1016/S0959-4388(00)00072-6

Haith A, Jackson C, Miall C, Vijayakumar S. 2008. Unifying the sensory and motor components of
sensorimotor adaptationAdvances in Neural Information Processing Systems 21 - Proceedings of
the 2008 Conference. pp. 593–600.

Harris CS. 1963. Adaptation to Displaced Vision: Visual, Motor, or Proprioceptive Change? *Science (80-)*
140:812–813. doi:10.1017/CBO9781107415324.004

Haruno M, Wolpert DM, Kawato M. 2001. MOSAIC Model for Sensorimotor Learning and Control. *Neural*
Comput **13**:2201–2220. doi:10.1162/089976601750541778

Hinton DC, Conradsson D, Bouyer L, Paquette C. 2020. Does dual task placement and duration affect
split-belt treadmill adaptation? *Gait Posture* **75**:115–120. doi:10.1016/j.gaitpost.2019.10.005

Hinton DC, Thiel A, Soucy JP, Bouyer L, Paquette C. 2019. Adjusting gait step-by-step: Brain activation
during split-belt treadmill walking. *Neuroimage* **202**:116095. doi:10.1016/j.neuroimage.2019.116095

Huberdeau DM, Krakauer JW, Haith AM. 2019. Practice induces a qualitative change in the memory
representation for visuomotor learning. *J Neurophysiol* **122**:1050–1059. doi:10.1152/jn.00830.2018

Huberdeau DM, Krakauer JW, Haith AM. 2015. Dual-process decomposition in human sensorimotor
adaptation. *Curr Opin Neurobiol* **33**:71–77. doi:10.1016/j.conb.2015.03.003

Ito M. 1989. Long-term depression. *Annu Rev Neurosci* **12**:85–102.

Izawa J, Criscimagna-Hemminger SE, Shadmehr R. 2012. Cerebellar contributions to reach adaptation
and learning sensory consequences of action. *J Neurosci* **32**:4230–4239.
doi:10.1523/JNEUROSCI.6353-11.2012

Jensen L, Prokop T, Dietz V. 1998. Adaptational effects during human split-belt walking: influence of
afferent input. *Exp Brain Res* **118**:126–130. doi:10.1007/s002210050262

Judd EN, Lewis SM, Person AL. 2021. Diverse inhibitory projections from the cerebellar interposed
nucleus. *Elife* **10**:e66231. doi:10.7554/eLife.66231

Kandel E, Schwartz JH, Jessell TM, Siegelbaum SA, Hudspeth AJ. 2013. Principles of Neural Science,
Fifth Edit. ed. McGraw-Hill, New York.

Lazar G, Van Laer J. 1968. Adaptation to displaced vision after experience with lesser displacements.
Percept Mot Skills **26**:579–582. doi:10.2466/pms.1968.26.2.579

Leech Kristan A., Day KA, Roemmich RT, Bastian AJ. 2018. Movement and perception recalibrate
differently across multiple days of locomotor learning. *J Neurophysiol* **120**:2130–2137.
doi:10.1152/jn.00355.2018.-Learning

Leech Kristan A, Roemmich RT, Bastian AJ. 2018. Creating flexible motor memories in human walking.
Sci Rep **8**:94. doi:10.1038/s41598-017-18538-w

Lindeman S, Hong S, Kros L, Mejias JF, Romano V, Oostenveld R, Negrello M, Bosman LWJ, de Zeeuw
CI. 2020. Cerebellar purkinje cells can differentially modulate coherence between sensory and
motor cortex depending on region and behavior. *Proc Natl Acad Sci U S A* **118**:2015292118.
doi:10.1073/pnas.2015292118

Long AW, Roemmich RT, Bastian AJ. 2016. Blocking trial-by-trial error correction does not interfere with
motor learning in human walking. *J Neurophysiol* **115**:2341–2348. doi:10.1152/jn.00941.2015

Malone LA, Bastian AJ. 2016. Age-related forgetting in locomotor adaptation. *Neurobiol Learn Mem*
128:1–6. doi:10.1016/j.nlm.2015.11.003

Malone LA, Bastian AJ. 2010. Thinking about walking: Effects of conscious correction versus distraction
on locomotor adaptation. *J Neurophysiol* **103**:1954–1962. doi:10.1152/jn.00832.2009

Malone LA, Vasudevan EVL, Bastian AJ. 2011. Motor adaptation training for faster relearning. *J Neurosci*

31:15136–15143. doi:10.1523/JNEUROSCI.1367-11.2011

Martin TA, Keating JG, Goodkin HP, Bastian AJ, Thach WT. 1996. Throwing while looking through prisms

II. Specificity and storage of multiple gaze-throw calibrations. *Brain* **119**:1199–1211.

doi:10.1093/brain/119.4.1183

McDougle SD, Taylor JA. 2019. Dissociable cognitive strategies for sensorimotor learning. *Nat Commun*

10:1–11. doi:10.1038/s41467-018-07941-0

Moidell BG, Bedell HE. 1988. Changes in oculocentric visual direction induced by the recalibration of

saccades. *Vision Res* **28**:329–336. doi:10.1016/0042-6989(88)90161-7

Naik Gaonkar M, Sawant K. 2013. AutoEpsDBSCAN : DBSCAN with Eps Automatic for Large Dataset. *Int*

J Adv Comput Theory Eng **2**:2319–2526.

Paul SS, Ada L, Canning CG. 2005. Automaticity of walking–implications for physiotherapy practice. *Phys*

Ther Rev **10**:15–23. doi:10.1179/108331905X43463

Petersen TH, Willerslev-Olsen M, Conway BA, Nielsen JB. 2012. The motor cortex drives the muscles

during walking in human subjects. *J Physiol* **590**:2443–2452. doi:10.1113/jphysiol.2012.227397

Pisano TJ, Dhanerawala ZM, Kislin M, Bakshinskaya D, Engel EA, Hansen EJ, Hoag AT, Lee J, de Oude

NL, Venkataraju KU, Verpeut JL, Hoebeek FE, Richardson BD, Boele HJ, Wang SSH. 2021.

Homologous organization of cerebellar pathways to sensory, motor, and associative forebrain. *Cell*

Rep **36**:109721. doi:10.1016/j.celrep.2021.109721

Poggio T, Bizzi E. 2004. Generalization in vision and motor control. *Nature* **431**:768–774.

doi:10.1038/nature03014

Popa D, Spolidoro M, Proville RD, Guyon N, Belliveau L, Léna C. 2013. Functional role of the cerebellum

in gamma-band synchronization of the sensory and motor cortices. *J Neurosci* **33**:6552–6556.

doi:10.1523/JNEUROSCI.5521-12.2013

Rahmah N, Sitanggang IS. 2016. Determination of Optimal Epsilon (Eps) Value on DBSCAN Algorithm to

Clustering Data on Peatland Hotspots in SumatraIOP Conference Series: Earth and Environmental

Science. Institute of Physics Publishing. p. 012012. doi:10.1088/1755-1315/31/1/012012

Reisman DS, Bastian AJ, Morton SM. 2010. Neurophysiologic and rehabilitation insights from the split-

belt and other locomotor adaptation paradigms. *Phys Ther* **90**:187–195. doi:10.2522/ptj.20090073

Reisman DS, Block HJ, Bastian AJ. 2005. Interlimb coordination during locomotion: what can be adapted

and stored? *J Neurophysiol* **94**:2403–2415. doi:10.1152/jn.00089.2005

Reisman DS, Wityk R, Silver K, Bastian AJ. 2007. Locomotor adaptation on a split-belt treadmill can

improve walking symmetry post-stroke. *Brain* **130**:1861–1872. doi:10.1093/brain/awm035

Roemmich RT, Bastian AJ. 2015. Two ways to save a newly learned motor pattern. *J Neurophysiol*

113:3519–3530. doi:10.1152/jn.00965.2014

Roemmich RT, Long AW, Bastian AJ, Roemmich RT, Long AW, Bastian AJ. 2016. Seeing the Errors You

Feel Enhances Locomotor Performance but Not Learning. *Curr Biol* **26**:1–10.

doi:10.1016/j.cub.2016.08.012

Rondi-Reig L, Paradis AL, Lefort JM, Babayan BM, Tobin C. 2014. How the cerebellum may monitor

sensory information for spatial representation. *Front Syst Neurosci* **8**:205.

doi:10.3389/fnsys.2014.00205

Rossi C, Bastian AJ, Therrien AS. 2021a. Mechanisms of proprioceptive realignment in human motor

learning. *Curr Opin Physiol* **20**:186–197. doi:10.1016/j.cophys.2021.01.011

Rossi C, Chau CW, Leech KA, Statton MA, Gonzalez AJ, Bastian AJ. 2019. The capacity to learn new

motor and perceptual calibrations develops concurrently in childhood. *Sci Rep* **9**:9322.

doi:10.1038/s41598-019-45074-6

Rossi C, Roemmich RT, Schweighofer N, Bastian AJ, Leech KA. 2021b. Younger and Late Middle-Aged

Adults Exhibit Different Patterns of Cognitive-Motor Interference During Locomotor Adaptation, With

No Disruption of Savings. *Front Aging Neurosci* **13**:789. doi:10.3389/fnagi.2021.729284

Seki K, Fetz EE. 2012. Gating of sensory input at spinal and cortical levels during preparation and

execution of voluntary movement. *J Neurosci* **32**:890–902. doi:10.1523/JNEUROSCI.4958-11.2012

Smith MA, Ghazizadeh A, Shadmehr R. 2006. Interacting adaptive processes with different timescales

underlie short-term motor learning. *PLoS Biol* **4**:e179. doi:10.1371/journal.pbio.0040179

Sombric C, Gonzalez-Rubio M, Torres-Oviedo G. 2019. Split-Belt walking induces changes in active, but

not passive, perception of step length. *Sci Rep* **9**:16442. doi:10.1038/s41598-019-52860-9

Sombric CJ, Harker HM, Sparto PJ, Torres-Oviedo G. 2017. Explicit action switching interferes with the
context-specificity of motor memories in older adults. *Front Aging Neurosci* **9**:40.
doi:10.3389/fnagi.2017.00040
Song CN, Stenum J, Leech KA, Keller CK, Roemmich RT. 2020. Unilateral step training can drive faster
learning of novel gait patterns. *Sci Reports 2020 101* **10**:1–13. doi:10.1038/s41598-020-75839-3
Statton MA, Vazquez A, Morton SM, Vasudevan EVL, Bastian AJ. 2018. Making sense of cerebellar
contributions to perceptual and motor adaptation. *Cerebellum* **17**:111–121. doi:10.1007/s12311-017-
0879-0
Sultan F, Augath M, Hamodeh S, Murayama Y, Oeltermann A, Rauch A, Thier P. 2012. Unravelling
cerebellar pathways with high temporal precision targeting motor and extensive sensory and parietal
networks. *Nat Commun* **3**:1–10. doi:10.1038/ncomms1912
Sun X, O'Shea DJ, Golub MD, Trautmann EM, Vyas S, Ryu SI, Shenoy K V. 2022. Cortical preparatory
activity indexes learned motor memories. *Nature* **602**:274–279. doi:10.1038/s41586-021-04329-x
Synofzik M, Lindner A, Thier P. 2008. The cerebellum updates predictions about the visual consequences
of one's behavior. *Curr Biol* **18**:814–818. doi:10.1016/J.CUB.2008.04.071
Taylor JA, Ivry RB. 2011. Flexible cognitive strategies during motor learning. *PLoS Comput Biol*
7:e1001096. doi:10.1371/journal.pcbi.1001096
Taylor JA, Krakauer JW, Ivry RB. 2014. Explicit and implicit contributions to learning in a sensorimotor
adaptation task. *J Neurosci* **34**:3023–3032. doi:10.1523/JNEUROSCI.3619-13.2014
Uiga L, Capio CM, Ryu D, Young WR, Wilson MR, Wong TWL, Tse ACY, Masters RSW. 2020. The Role
of Movement-Specific Reinvestment in Visuomotor Control of Walking by Older Adults. *Journals*
Gerontol - Ser B Psychol Sci Soc Sci **75**:282–292. doi:10.1093/geronb/gby078
Van Vugt FT, Ostry DJ. 2018. The structure and acquisition of sensorimotor maps. *J Cogn Neurosci*
30:290–306. doi:10.1162/jocn_a_01204
Vazquez A, Statton MA, Busgang SA, Bastian AJ. 2015. Split-belt walking adaptation recalibrates
sensorimotor estimates of leg speed but not position or force. *J Neurophysiol* **114**:3255–3267.
doi:10.1152/jn.00302.2015
Vervoort D, Rob Den Otter A, Buurke TJW, Vuillerme N, Hortobágyi T, Lamoth CJC. 2019. Effects of
aging and task prioritization on split-belt gait adaptation. *Front Aging Neurosci* **11**:10.
doi:10.3389/fnagi.2019.00010
Weeks HM, Therrien AS, Bastian AJ. 2017. The cerebellum contributes to proprioception during motion. *J*
Neurophysiol **118**:693–702. doi:10.1152/jn.00417.2016
Wolpert DM, Ghahramani Z, Flanagan JR. 2001. Perspectives and problems in motor learning. *Trends*
Cogn Sci. doi:10.1016/S1364-6613(00)01773-3
Wong WL, Masters RSW, Maxwell JP, Abernethy AB. 2008. Reinvestment and falls in community-
dwelling older adults. *Neurorehabil Neural Repair* **22**:410–414. doi:10.1177/1545968307313510
Yavari F, Mahdavi S, Towhidkhah F, Ahmadi-Pajouh MA, Ekhtiari H, Darainy M. 2016. Cerebellum as a
forward but not inverse model in visuomotor adaptation task: a tDCS-based and modeling study.
Exp Brain Res **234**:997–1012. doi:10.1007/s00221-015-4523-2
1005
1006

 Figure Legends

 Figure 1. Experiment 1, Hypotheses and Predictions. (A-B) Treadmill belt speeds for the
 standard split-belt paradigm (A) and schematic time course of standard motor measures of
 walking adaptation (B): step length asymmetry – a measure of error (solid purple), Δ motor output
 – a measure of compensatory spatial and temporal asymmetries (dotted blue), and perturbation
 – the effect of the speed asymmetry on the walking pattern (dashed red). In baseline, the belts
 are tied, and perturbation, Δ motor output, and step length asymmetry are all ~0. In adaptation,
 the right leg is faster than the left such that the perturbation is positive. The Δ motor output is still
 ~0 in early adaptation, causing step length asymmetry errors (negative values). By late adaptation
 the Δ motor output is adapted to match the perturbation, and step length asymmetry returns to
 ~0. Changes to Δ motor output persist in tied-belts post-adaptation, but the perturbation is ~0,
 causing step length asymmetry aftereffects (positive error values, smaller magnitude than early
 adaptation). **(C)** Schematic of our paradigm with the Ramp Down task: after adaptation, the right
 belt speed is gradually ramped down to match the left. **(D)** Predictions for the Ramp Down motor
 measures made by three competing hypotheses. **Recalibration only:** changes to Δ motor output
 by recalibration are fixed, predicting that step length asymmetry aftereffects will emerge
 immediately as the perturbation decreases. **Mapping only:** changes by mapping are flexible and
 are predicted to track the perturbation, with no step length asymmetry aftereffect. **Recalibration**
 + mapping: changes involving both recalibration and mapping predict an intermediate result. At
 first, changes by mapping would be gradually abandoned to track the perturbation, such that step
 length asymmetry would remain zero. When only recalibration-related changes are left, the Δ
 motor output would be fixed and step length asymmetry aftereffects would begin to emerge.
 Aftereffect magnitude on tied belts would be smaller than the initial adaptation asymmetry.

 Figure 2. Experiment 1, Step length asymmetry. (A) Experimental protocol. The Ramp Down
 task (purple) is that used for predictions in Fig. 1D. **(B)** Step length asymmetry time course.
 Background shading darkness increases with belt speed difference (color bar). Phases (except
 Ramp tasks) are truncated to the participant with fewest strides. **(C-D)** Zoomed-in Ramp Up (C,
 baseline) and Ramp Down (D, post-adaptation) tasks. Speed differences for which step length
 asymmetry is not significantly different from zero are indicated by the green shade. Inset depicts
 predictions made by the competing hypotheses as in Fig. 1D. All curves show group mean ± SE.

 Figure 3. Experiment 1, Perturbation and Δ motor output. (A) Perturbation (red) and Δ motor
 output (blue) data for the Ramp Down task. Inset depicts predictions made by the competing
 hypotheses as in Fig. 1D. **(B-E)** Perturbation data (red) and model fit for the Δ motor output (blue)
 for recalibration only, mapping only, recalibration + mapping, and dual state models. The blue
 bracket in (D) points to the value on the y-axis of the model parameter "r". All curves show group
 mean ± SE.

 Figure 4. Experiment 1, Perceptual Results. (A) Top: perturbation data (red) and recalibration
 + mapping fit (blue). Bottom: perceptual task button presses (green, group mean ± SE), as a
 function of belt speed difference. Right: measures of motor recalibration ("r") and total motor
 adaptation ("u_{plateau}"). **(B)** Perturbation compensation (normalized perceptual and motor
 measures of adaptation): $compensation_{\text{perceptual}}$ bounds (green), $compensation_{\text{motor total}}$ (dark
 blue), and $compensation_{\text{motor recalibration}}$ (light blue). **(C-D)** Individual participants'

1053 $compensation_{\text{motor recalibration}}$ versus $compensation_{\text{perceptual}}$ (first or second button press). Solid
1054 black: least squares line. Dashed gray: unity line.
1055
Figure 5. Experiment 2, Step length asymmetry. (A) Experimental protocol, equivalent to that
of Experiment 1 except for the Ramp Up & Down part shaded in teal, where the right speed was
faster than in adaptation (ramped up to 2m/s and back down to 1.5m/s). **Inset:** Predictions for the
step length asymmetry during the teal portion of the Ramp Up & Down task, for the memory-
based (top) or structure-based (bottom) mapping hypotheses. **(B)** step length asymmetry time
course (entire group mean ± SE). Background shade represents belt speed difference. Phases
(except Ramp tasks) are truncated to the participant with fewest strides. **(C)** Zoomed-in Ramp Up
& Down task (entire group mean ± SE). Step length asymmetry for strides taken at right speeds
larger than adaptation is shown in teal. **(D-E)** Separate plots of the step length asymmetry in the
Ramp Up & Down task for the subgroups of participants that walked asymmetrically (D, "memory-
based") versus symmetrically (E, "structure-based") in the teal portion of the task (subgroups
mean ± SE). **Insets:** circles represent individual participants' number of strides, in the teal portion
of the task, with step length asymmetry below their own baseline CI. Error bars depict subgroup
mean ± SE. Subgroup assignment was performed by clustering on this measure.
1070
1071 **Figure 6.** Summary of self-reported deliberate changes to the walking pattern in adaptation. Only
1072 one participant accurately described changes to the walking pattern that related to adaptation,
1073 while other responses were negative (i.e. no deliberate changes, 3 participants), irrelevant (7
1074 participants), or inaccurate (5 participants).
1075
Figure 7. Schematic model of adaptation. Body movement depends on environment
perturbations (red) and Δ motor output (blue). The Δ motor output is adjusted by recalibration
(light blue) and mapping (dark blue) mechanisms, which perform different operations and are
arranged in tandem. We propose the following architecture and flow: **(1. Recalibration)** The
recalibration mechanism produces adjustment x_r that is fixed regardless of perturbation size (light
blue box, x_r is constant for varying p). The *same* recalibration adjustment x_r serves as an input
to *both* areas responsible for conscious perception (green box) and Δ motor output (blue box). **(2.**
Perception) Conscious perception is computed by cancelling out the recalibration adjustment
from the actual sensory feedback (green box, perception of the belt speed difference perturbation
$\tilde{p}$ is the difference between the actual speed difference p and recalibration x_r). The perceived
perturbation $\tilde{p}$ serves as an input to the mapping mechanism (dark blue box). **(3. Mapping)** The
mapping mechanism produces adjustment x_m that can vary in magnitude to appropriately
account for the perceived perturbation $\tilde{p}$ (dark blue box, x_m scales with $\tilde{p}$ and matches its
magnitude). **(4. Δ Motor output)** The overall adjustment to Δ motor output is computed by adding
the mapping adjustment x_m and recalibration adjustment x_r. The corner in the Δ motor output
versus perturbation profile arises because mapping is computed based on the perceived
perturbation $\tilde{p}$ (not the actual perturbation p) and is only learnt for positive $\tilde{p}$ (the experienced
direction). When the perturbation is perceived to be opposite than adaptation, even if it is not,
mapping is zero and the Δ motor output is constant, reflecting recalibration adjustments only (blue
box, when $\tilde{p} < 0$ and $p \geq 0$ the mapping adjustment x_m is zero and $u = x_r$).
1096

standard paradigm

A

baseline · adaptation · post-adaptation

speeds

- - - right
—— left

standard response

B

baseline · adaptation · post-adaptation

motor measures

stride

step length asymmetry =

····· Δ **motor output**
(asymmetry in step time and position)

−

- - - **perturbation**
(asymmetry in foot velocity)

Ramp Down paradigm

C

baseline · adaptation · Ramp Down

right

- - - right
—— left

predictions

D

recalibration only

or

mapping only

or

mapping + **recalibration**

stride

Data

A

predictions

recalibration only

mapping only

recalibration + mapping

— Δmotor output
— perturbation

right - left speed [m/s]

Model fits

B recalibration only

C mapping only

stride

right - left speed [m/s]

D recalibration + mapping

r

stride

right - left speed [m/s]

E dual state
(recalibration only with forgetting)

stride

right - left speed [m/s]

— model fit of Δmotor output   — perturbation

A

right leg feels...

faster   equal   slower

① ②

motor

u_plateau

r

PSE_lower bound ②

PSE_upper bound ①

perceptual

right – left speed [m/s]

B

perturbation compensation [%]

① upper bound

② lower bound

perceptual total   motor recalibration   motor total

C

motor recalibration [%]

perceptual upper bound [%]

r=0.64
p=0.002

(① "speeds feel equal")

D

motor recalibration [%]

perceptual lower bound [%]

r=0.30
p=0.195

(② "right feels slower")