Title: Automatic learning mechanisms for flexible human locomotion.

Authors: Cristina Rossi[a,b], Kristan A. Leech[c,d], Ryan T. Roemmich[b,e], Amy J. Bastian[a,b,*]

[a] Department of Neuroscience, The Johns Hopkins University School of Medicine, Baltimore, MD, 21205,
USA; [b] Center for Movement Studies, Kennedy Krieger Institute, Baltimore, MD, 21205, USA; [c] Division of
Biokinesiology and Physical Therapy, University of Southern California, Los Angeles, CA, 90033, USA; [d]
Neuroscience Graduate Program, University of Southern California, Los Angeles, CA, 90007, USA; [e]
Department of Physical Medicine and Rehabilitation, The Johns Hopkins University School of Medicine,
Baltimore, MD, 21205, USA. *Corresponding author: Amy J. Bastian, bastian@kennedykrieger.org.

This PDF file includes:

Abstract

Movement flexibility and automaticity are necessary to successfully navigate different environments. When encountering difficult terrains such as a muddy trail, we can change how we step almost immediately so that we can continue walking. This flexibility comes at a cost since we initially must pay deliberate attention to how we are moving. Gradually, after a few minutes on the trail, stepping becomes automatic so that we do not need to think about our movements. Canonical theory indicates that different adaptive motor learning mechanisms confer these essential properties to movement: explicit control confers rapid flexibility, while forward model recalibration confers automaticity. Here we uncover a distinct mechanism of treadmill walking adaptation – an automatic stimulus-response mapping – that confers both properties to movement. The mechanism is flexible as it learns stepping patterns that can be rapidly changed to suit a range of treadmill configurations. It is also automatic as it can operate without deliberate control or explicit awareness by the participants. Our findings reveal a tandem architecture of forward model recalibration and automatic stimulus-response mapping mechanisms for walking, reconciling different findings of motor adaptation and perceptual realignment.

Abbreviations
CI, confidence interval; CI_{LB}, lower bound of CI; CI_{UB}, upper bound of CI; PSE, point of subjective equality; norm, normalized; SD, standard deviation; SE, standard error

Main Text

Introduction

Flexibility and automaticity are essential features of human movement, so much so that we rarely think about the details of how we move. While walking, we do not think about the bend of the ankle or how quickly to swing the leg forward to step. Yet, we easily adjust walking to accommodate many situations – a muddy trail, a grassy slope, or a snowy path. These abilities are often taken for granted until something goes awry – a sprained ankle quickly brings the details of movement execution to awareness, and deliberate control is often used to avoid pain or further injury. It is only then that we truly appreciate how much the sensorimotor system is doing without our conscious awareness.

We currently do not understand how different motor learning mechanisms confer both flexibility and automaticity to human movement. One well-studied process of motor learning is sensorimotor adaptation, which occurs in response to errors between the expected and actual sensory consequences of our movements. Accordingly, this motor learning mechanism helps adjust motor commands to correct for perturbations to our movements caused by altered environmental demands (Bastian, 2008; Huberdeau et al., 2015).

Adaptation is traditionally thought to rely on the cerebellum-dependent recalibration of a forward model that associates motor commands with expected sensory consequences (Ito, 1989; Wolpert et al., 2001). For example, to walk on an icy sidewalk, we may need to recalibrate our prediction of how firmly our feet will grip the ground. The process of forward model recalibration is automatic and implicit, as it has been shown to operate without intention or awareness (Huberdeau et al., 2015; Taylor et al., 2014). However, forward model recalibration does not confer rapid flexibility because it cannot make immediate changes in movement – it can only adjust movement gradually trial-by-trial or step-by-step, a process that can take several minutes (Bastian, 2008; Huberdeau et al., 2015). The newly acquired sensorimotor recalibration must be unlearned over time to restore a normal movement when the environment returns to its original state after adaptation (Bastian, 2008; Martin et al., 1996a). For this reason, forward model recalibration leads to lasting movement errors (called "aftereffects").

Sensorimotor adaptation can rely on more than forward model recalibration. We know that adaptation of goal-directed reaching movements can involve rapid flexible learning mechanisms

sometimes called "stimulus-response mapping mechanisms". Different types of stimulus-
response mapping mechanisms have been characterized in previous studies: First, explicit
strategies – where people deliberately change where they are aiming to reach (Taylor et al., 2014;
Taylor and Ivry, 2011). Second, memory-based caching – where people learn motor responses
in association with the respective environmental sensory stimuli, and cache them in memory for
future retrieval (like a lookup-table) (Huberdeau et al., 2019; McDougle and Taylor, 2019). Third,
structural learning – where people learn general relationships between environmental sensory
stimuli and motor responses and use them to produce novel responses (like when we learn to
map the movement of a computer mouse to that of the cursor) (Bond and Taylor, 2017; Braun et
al., 2009). Stimulus-response mapping mechanisms differ from forward model recalibration in that
they confer rapid flexibility – novel responses can be promptly abandoned or changed for different
environmental stimuli – and therefore do not lead to aftereffects (Huberdeau et al., 2015; Taylor
et al., 2014; Taylor and Ivry, 2011).

It is unclear if stimulus-response mapping mechanisms are involved in adapting movement types
that are continuous like walking versus discrete movements like reaching. The goal of this study
was to understand whether walking adaptation involves any stimulus-response mapping
mechanism and, if so, how it operates and how it can be dissected from forward model
recalibration. We focused on adaptation of walking movements on a split-belt treadmill: it is well
established that when people walk with one foot faster than the other, they adapt the timing and
location of their step to restore symmetry (Box 1).

In Experiment 1A we tested the presence of stimulus-response mapping during gait adaptation
by evaluating whether people develop the ability to modify their walking pattern immediately for
different split-belt magnitudes. Preliminary evidence suggests that people may be able to switch
between at least two different walking patterns more rapidly than we would expect from forward
model recalibration alone (Box 2) (Leech et al., 2018a). Based on this evidence, we hypothesized
that walking adaptation involves stimulus-response mapping in addition to forward model
recalibration.

As our findings corroborated this hypothesis, in Experiment 1B we aimed to develop a measure
to dissect individual contributions of the two mechanisms to adaptation. We based this on the
well-known phenomenon of "perceptual realignment", where perception of the belt speed
difference diminishes over time during adaptation (Box 2) (Jensen et al., 1998; Leech et al.,

2018a; Rossi et al., 2019; Vazquez et al., 2015). Previous studies show that perceptual realignment is only partial – the belt speeds feel similar but not completely equal at the end of adaptation, despite complete motor adaptation (Leech et al., 2018a; Vazquez et al., 2015). Studies also suggest that perceptual realignment may stem from forward model recalibration processes ('t Hart and Henriques, 2016; Izawa et al., 2012; Rossi et al., 2021a; Synofzik et al., 2008). Based on these studies, we hypothesized that the extent of perceptual realignment corresponds to the contribution of forward model recalibration to motor adaptation, with the remainder attributed to stimulus-response mapping.

Finally, in Experiment 2 we began to explore characteristics of stimulus-response mapping. We asked whether the stimulus-response mapping mechanism is automatic, or instead under deliberate or explicit control. Stimulus-response mapping mechanisms for reaching require explicit control (Bond and Taylor, 2017; Huberdeau et al., 2019; Taylor and Ivry, 2011), but explicit control is poorly suited for automatic, continuous movements like walking (Clark, 2015; Paul et al., 2005; Uiga et al., 2020), and can even lead to falls (Wong et al., 2008). Unlike reaching, adaptation of walking is unaffected by explicit goals or instructions given to the participants on where to aim their feet (Long et al., 2016; Roemmich et al., 2016). Hence, we hypothesized that participants would not be able to describe how they changed walking, in contrast to what has been previously reported in reaching. We also asked if participants could use the mapping mechanism to produce novel motor outputs akin to what has been interpreted as structural learning in reaching (Bond and Taylor, 2017; Braun et al., 2009).

Results

Experiment 1

Motor paradigm and hypotheses

We asked whether walking adaptation involves both forward model recalibration and stimulus-response mapping mechanisms learned in tandem ("recalibration + mapping hypothesis"). We contrasted this with the alternative hypothesis that walking adaptation may involve only forward model recalibration mechanisms ("recalibration only hypothesis").

To test this, we devised the "Ramp Down" paradigm depicted in Figure 1A. After adaptation, we

gradually decreased the speed of the right belt every 3 strides (see Methods). This enabled us to

obtain reliable measurements of aftereffects across 21 predetermined speed configurations,

spanning from the full split-belt perturbation to tied-belts. We measure aftereffects as the

magnitude of step length asymmetry.

As depicted in Figure 1B, the recalibration only hypothesis predicts that aftereffects will be present

for all speed configurations in the Ramp Down. This is because the Δ motor output (blue) can

only change gradually, through unlearning or forgetting of the forward model recalibration (SI

Appendix, Fig. S1). This change is too slow to track the perturbation (red), which is ramped down

to zero rapidly over ~80 seconds. Therefore, step length asymmetry (purple) becomes positive

immediately in the Ramp Down.

In contrast, as depicted in Figure 1C, the recalibration + mapping hypothesis predicts that

aftereffects will be present for some, but not all speed configurations in the Ramp Down.

Specifically, there would be no aftereffects (~zero step length asymmetry) in the first portion of

the Ramp Down, because the Δ motor output (blue) changes rapidly and matches the perturbation

(red). This reflects the flexibility of the stimulus-response mapping (SI Appendix, Fig. S1).

Aftereffects would emerge in the second portion of the task (positive step length asymmetry),

reflecting forward model recalibration.

We include forgetting effects for the recalibration only hypothesis to account for previous findings

from Leech et al. (Leech et al., 2018a) (Box 2). The inset of Figure 1 (right column) illustrates how

both hypotheses may account for near-zero step length asymmetry at the end of the speed match

task. The Δ motor output decreases to match the perturbation either because the mapping

mechanism flexibly scales down (recalibration + mapping hypothesis) or, alternatively, because

of forgetting/unlearning (recalibration only hypothesis).

Figure 1. Experiment 1, Hypotheses and Predictions. (A) Conceptual schematic of our paradigm with
the Ramp Down task: after adaptation, the right belt speed is gradually ramped down to match the left. **(B-
C)** Predictions for the Ramp Down motor measures made by two competing hypotheses. **(B) Recalibration
only:** recalibration can only change movement gradually. The Δ motor output (dotted blue line) changes
slowly and does not track the rapidly decreasing perturbation (dashed red line), so that step length
asymmetry aftereffects emerge immediately (solid purple line, magnitude is positive). **(C) Recalibration +
mapping:** mapping can change movement immediately. In the first part of the task (highlighted in green),
the mapping contribution to Δ motor output (dark blue shade) is scaled down immediately as the
perturbation decreases. Hence, the Δ motor output (dotted blue line) changes rapidly and tracks the
perturbation (dashed red line), so that there are no step length asymmetry aftereffects (solid purple line,
magnitude is ~zero). In the second part of the task, the mapping contribution to Δ motor output is zero, and
the recalibration contribution to Δ motor output (light blue shade) does not change significantly. Hence, the
Δ motor output (dotted blue line) does not track the perturbation (red dashed line), and step length
asymmetry aftereffects emerge (solid purple line, magnitude is positive). **Right column inset:** conceptual
explanation of how both hypotheses may account for the speed match results from Leech at al. (Leech et
al., 2018a). In the first post-adaptation speed match task, participants increase the speed of the right belt
from zero to a value that is smaller than adaptation but larger than the left belt (top panel). The perturbation
increases until a value that is positive but smaller than adaptation (dashed red line, middle and bottom
panels). Leech et al. observed symmetric step lengths at the end of the task, indicating that the Δ motor
output (dotted blue line) is smaller than it was in adaptation and matches the perturbation. The decrease in
Δ motor output can be explained by the recalibration only hypothesis as forgetting/unlearning (middle
panel), or by the recalibration + mapping hypothesis as flexible scaling of the mapping contribution (bottom
panel).

Motor results – step length asymmetry

Figure 2A shows the paradigm used for Experiment 1, and the time course of step length

asymmetry (group mean ± SE). The Ramp Down task corresponds to the manipulation illustrated

in Figure 1 and is the focus of our analysis for Experiment 1. A similar ramp task was performed

prior to adaptation as a control to see how participants responded to a gradually ramped

perturbation at baseline (see Methods, "Ramp tasks"). Participants performed perceptual tests

during the ramp tasks, which will be discussed in a later section (see "Perceptual results*").

Patterns of step length asymmetry during other portions of the paradigm were consistent with a

large body of previous work (e.g., Leech et al., 2018a, 2018b; Reisman et al., 2005; Roemmich

et al., 2016; Rossi et al., 2019; Vazquez et al., 2015) and will not be discussed further.

Step length asymmetry during the baseline ramp and post-adaptation Ramp Down tasks are

displayed in Figure 2B. For each speed configuration in the Ramp Down task, we statistically

compared step length asymmetry to zero. We evaluated the emergence of aftereffects by

examining when step length asymmetry became significantly positive. We found that step length

asymmetry was *not* statistically different from zero during the first half of the Ramp Down task, for

right belt speeds ranging from 1m/s to 0.5m/s faster than the left speed (these speed

configurations are highlighted in green in Fig. 2B right; all CI_{LB} [confidence intervals lower bounds]

≤ -0.001 and CI_{UB} [upper bounds] ≥ 0.009, SI Appendix, Fig. S2 and Table S1). This result

indicates that aftereffects do not emerge immediately in the Ramp Down task, a finding at odds

with the recalibration only prediction (Fig. 2B 'predictions' inset, top). Step length asymmetry was

instead significantly positive for the second half of the task, for right belt speeds ranging from

0.45m/s to 0m/s faster than the left speed (all CI_{LB} > 0.01). These results indicate that aftereffects

emerge at a mid-point during the Ramp Down task and align with behavioral predictions from the

recalibration + mapping hypothesis (Fig. 2B 'predictions' inset, bottom).

As a control, we repeated the same analysis for the baseline ramp task (Fig. 2B, left). We found

that there was only one speed configuration for which step length asymmetry was not statistically

different from zero (right speed 0.05m/s slower than left; SL asym. = 0.006 [-0.017, 0.028], mean

[CI]). For all other configurations, step length asymmetry was significantly positive (CI_{LB} ≥ 0.048

for right speed 0.1 to 0.35m/s slower than left) or negative (CI_{UB} ≤ -0.019 for right speed 0 to

0.35m/s faster than left). This is in stark contrast to the wide range of speed configurations with

near-zero step length asymmetry observed in the Ramp Down period. Hence, the ability to walk

symmetrically at different speed configurations is not innate but dependent on the adaptation
process, corroborating the hypothesis that the "no aftereffects" region of the Ramp Down reflects
the operation of a mapping mechanism.

Figure 2. Experiment 1, Step length asymmetry. (A) Top: Experimental protocol. The Ramp Down task
(purple) is used to test the predictions illustrated in Fig. 1. Bottom: Step length asymmetry time course.
Background shading darkness increases with belt speed difference (color bar). Phases (except Ramp
tasks) are truncated to the participant with fewest strides. **(B)** Zoomed-in baseline ramp and post-adaptation
Ramp Down tasks. Speed differences for which step length asymmetry is not significantly different from
zero are indicated by the green shade. Inset depicts predictions made by the competing hypotheses as in
Fig. 1. All curves show group mean ± SE.
219

Motor results – perturbation and Δ motor output

To further test the competing hypotheses, we examined the perturbation and Δ motor output
during the Ramp Down (Figure 3A). As the belt speed difference decreased, so did the
perturbation component (red). In the first half of the task, the Δ motor output (blue) appeared to
match the perturbation – consistent with the lack of step length asymmetry aftereffects (green
shaded portion, matching Fig. 2B). Additionally, in the second half of the task, the Δ motor output
appeared larger than the perturbation – consistent with the positive step length asymmetry
aftereffects observed in Fig. 2B.

Figure 3. Experiment 1, Perturbation and Δ motor output. (A) Perturbation (red) and Δ motor output
(blue) data for the Ramp Down task. **(B-C)** Perturbation data (red) and model fit for the Δ motor output
(blue) for recalibration only and recalibration + mapping models. All curves show group mean ± SE. Green
shade corresponds to speeds with symmetric step lengths as in Fig. 2.
235

We formally contrasted predictions made by the competing hypotheses by developing
mathematical models of the Δ motor output as a function of perturbation (Fig. 3B-C). In the
simplest framework, the Δ motor output behavior for the recalibration and mapping mechanisms
can be formalized as follows:

Forward Model Recalibration: $u(p) = r$ (1)

Stimulus-Response Mapping: $u(p) = p$ (2)

Recalibration + Mapping: $u(p) = \begin{cases} p & , & \text{for } p \geq r \\ r & , & \text{otherwise} \end{cases}$ (3)

Where u is the modelled Δ motor output, p is the perturbation, and r is a free parameter
representing the portion of the Δ motor output related to recalibration. We used equation (3) as
our model for the recalibration + mapping hypothesis. This describes the scenario where
participants modulate the Δ motor output to match the perturbation ($u(p) = p$) for the first portion
of the Ramp Down task. The Δ motor output remains constant during the second portion of the
task ($u(p) = r$) because participants are unable to reduce the Δ motor output to values smaller
than "r" (the amount achieved by recalibration). Parameters were estimated by fitting the model
to individual participants' Δ motor output data from the Ramp Down (see Methods).

The fitted Δ motor output is displayed in Figure 3C (mean ± SE of fits across participants;
individual fits are shown in SI Appendix, Fig. S3). As expected, the recalibration + mapping fit
captured the matching-then-divergent behavior of Δ motor output in response to the changing
perturbation.

We used the dual state model of motor adaptation from Smith et al. (Smith et al., 2006) as our
model for the recalibration only hypothesis (see Methods). The model has four parameters and
can account for potential forgetting or unlearning of the Δ motor output that may occur during the
Ramp Down. It can also account for the possibility that adaptation involves two recalibration
mechanisms – i.e., two distinct 'fast' and 'slow' mechanisms that both learn via a process of

forward model recalibration, and both contribute to aftereffects, but that learn and forget at
different rates. We chose this model because it is a well-established model that is widely used to
capture the Δ motor output time-course for traditional motor adaptation paradigms (i.e., those
consisting of adaptation and post-adaptation phases). Yet, performance of this model for a
manipulation like the Ramp Down task performed here has not yet been tested. As such, if the
recalibration + mapping model fit the Ramp Down data better than the dual state model, this would
provide robust evidence for the presence of a stimulus-response mapping mechanism.

We show the Δ motor output fit by the recalibration only (dual state) model in Figure 3B (group
mean ± SE; individual fits are shown in SI Appendix, Fig. S4). In contrast to the recalibration +
mapping model, the dual state model was not able to capture the matching-then-divergent
behavior of Δ motor output. The BIC statistic confirmed that the recalibration + mapping model
fitted the data significantly better than the dual state (BIC difference = 8.422 [3.532, 13.573], mean
[CI]).

In a supplementary analysis, we considered two additional prominent models for motor
adaptation: optimal feedback control (Izawa and Shadmehr, 2011; Shadmehr and Krakauer,
2008; Todorov, 2004) and memory of errors (Herzfeld et al., 2014). Similar to the dual state, these
models could not capture the matching-then-divergent behavior of Δ motor output and fitted the
Ramp Down data significantly worse than the recalibration + mapping (see SI Appendix,
"Experiment 1 additional models" and Fig. S5; memory of errors minus recalibration+mapping BIC
difference = 11.610 [6.220, 17.558], optimal feedback control minus recalibration+mapping BIC
difference = 19.272 [14.378, 24.399], mean [CI]). In sum, the modeling analysis of the Δ motor
output further supports the recalibration + mapping hypothesis.

Perceptual test and results

The second goal of Experiment 1 was to evaluate the hypothesis that "perceptual realignment" (a
phenomenon leading to altered perception following adaptation) results from the operation of the
same forward model recalibration mechanism involved in adaptation of the Δ motor output (Rossi
et al., 2021a).

Previous work shows that perception realigns following gait adaptation: after adapting to a
perturbation where the right treadmill belt is faster than the left, return to tied belts results in
perception of the opposite asymmetry (i.e., right speed feels slower than left) (Jensen et al., 1998;

Vazquez et al., 2015). We therefore expected that people would perceive the belt speeds as equal
at some point in the Ramp Down and eventually perceive the opposite asymmetry on tied belts.
We measured this by asking participants to press a keyboard button at two separate occasions
during the Ramp Down task: (1) when the belts first felt equal, and (2) when they no longer felt
equal (see Methods).

Figure 4A depicts the button presses for the Ramp Down perceptual test (top panel, group mean)
overlayed onto the Ramp Down motor data (recalibration + mapping fit). Note that the task
captures a *range* of belt speed configurations that participants perceive as "equal speeds", with
button presses (1) and (2) corresponding to the upper and lower bounds of this range. This is
expected because perception is known to be noisy and may not be sensitive enough to
discriminate between belt speed configurations that are too similar.

We quantified perceptual realignment using the established measure of *point of subjective*
equality (PSE), defined as the belt speed difference perceived as "equal speeds". A PSE of zero
would indicate no perceptual realignment (accurate perception) and a PSE of 1m/s (i.e., a
magnitude equivalent to the difference between the belt speeds during adaptation) would indicate
complete perceptual realignment such that the belt speeds feel equal during the adaptation
phase.

We measured belt speed difference at the time of each button press. We computed PSE as the
range of belt speed difference values between these two measurements:

$$PSE_{\text{upper bound}} = [\, right\ belt\ speed - left\ belt\ speed\,]_{\text{time of button press 1}} \qquad (4)$$

$$PSE_{\text{lower bound}} = [\, right\ belt\ speed - left\ belt\ speed\,]_{\text{time of button press 2}} \qquad (5)$$

We found that $PSE_{\text{upper bound}}$ was 0.64±0.03m/s and $PSE_{\text{lower bound}}$ was 0.39±0.05m/s (Figure 4A,
bottom panel, group mean ± SE). This was consistent with previous work (Leech et al.,
2018a).This perceptual realignment was not present during baseline testing, as illustrated in SI
Appendix, Fig. S6.

Figure 4. Experiment 1, Perceptual Results. (A) Top: perturbation data (red) and recalibration + mapping
fit (blue); this is the same as Fig. 3C. Bottom: perceptual task button presses (green, group mean ± SE),
as a function of belt speed difference. Right: measures of motor recalibration ("r") and total motor adaptation
("u_{plateau}"). **(B)** Perturbation compensation (normalized perceptual and motor measures of adaptation):
$compensation_{\text{perceptual}}$ bounds (green), $compensation_{\text{motor total}}$ (dark blue), and $compensation_{\text{motor recalibration}}$
(light blue). **(C-D)** Individual participants' $compensation_{\text{motor recalibration}}$ versus $compensation_{\text{perceptual}}$ (first or
second button press). Solid black: least squares line. Dashed gray: unity line.

We aimed to evaluate the hypothesis that perceptual realignment arises from the forward model

mechanism of motor adaptation and is unaffected by the stimulus-response mapping mechanism.

This hypothesis predicts that the extent of perceptual realignment should be: 1) approximately

equal to the extent of motor adaptation achieved by recalibration, and 2) less than the total extent

of motor adaptation (which also includes mapping). We quantified 1) motor adaptation by

recalibration as the fitted parameter "r" from the recalibration + mapping model, and 2) total motor

adaptation as the Δ motor output at adaptation plateau (Fig. 4A, right panel, see previous section

"*Motor results – perturbation and Δ motor output*"). We expressed these motor measures and

perceptual realignment as "percent compensation for the perturbation" (i.e., normalized to the

perturbation magnitude in the respective units) so that they could be compared:

$$compensation_{\text{perceptual}} = \frac{PSE}{1\,\text{m/s}} \tag{6}$$

$$compensation_{\text{motor recalibration}} = \frac{r}{p_{\text{plateau}}} \tag{7}$$

$$compensation_{\text{motor total}} = \frac{u_{\text{plateau}}}{p_{\text{plateau}}} \tag{8}$$

Where r is the fitted parameter from the recalibration + mapping model, p_{plateau} is the mean

perturbation over the last 30 strides of adaptation, and u_{plateau} is the mean Δ motor output over

the last 30 strides of adaptation (for Eq. 6, note that 1m/s is the belt speed difference in

adaptation).

We   show   group   level   compensation   measures   in   Fig.   4B.   We   found   that

$compensation_{\text{motor recalibration}}$ (56±4%, group mean ± SE) fell within the $compensation_{\text{perceptual}}$

range (39±5% to 64±3%): it was significantly smaller than the upper bound (difference = -8 [-14,

-2]%) and significantly larger than the lower bound (difference = 18 [8, 28]%, mean [CI]). This

supports our first prediction that perceptual realignment is comparable to the extent of motor

adaptation achieved by recalibration. Furthermore, $compensation_{\text{perceptual}}$ was significantly

smaller than $compensation_{\text{motor total}}$ (95±2%, mean ± SE; difference = 31 [25, 38]% or 57 [48,

66]%, upper or lower perceptual bounds, mean [CI]). This supports our second prediction that

perceptual realignment is less than the total extent of motor adaptation.

We show individual $compensation_{\text{motor recalibration}}$ and $compensation_{\text{perceptual}}$ measures for each

participant in Figure 4C-D. We evaluated Pearson's correlation coefficients between these

measures to test whether there is a direct relationship between perceptual realignment and the

motor adaptation achieved by recalibration. We found that $compensation_{\text{motor recalibration}}$ was

significantly correlated with the upper bound of $compensation_{\text{perceptual}}$ – the value computed using

the "speeds feel equal" button press (r=0.64, p=0.002). This supports the hypothesized

relationship between motor recalibration and changes to leg speed perception. Instead,

$compensation_{\text{motor recalibration}}$ was not correlated with the lower bound of $compensation_{\text{perceptual}}$

(r=0.30, p=0.195) – this value is computed using the "right feels slower than left" button press,

suggesting the true PSE may lay closer to the first button press. Together, our results support the

hypothesis that perceptual realignment can be used as a proxy measure for the extent of motor

adaptation achieved by forward model recalibration.

Control experiments

We performed six control experiments and reanalyzed previously-published data (Leech et al.,

2018a) to replicate the findings of Experiment 1 across different paradigm conditions (SI

Appendix, "Control experiments", Fig. S7-9, and Table S2-6). This provided additional support for

the recalibration + mapping hypothesis. We first checked that the motor and perceptual behaviors

observed in the Ramp Down could be replicated using a different method of assessment. A "speed

match" task was given where participants use a joystick to increase the speed of the right belt

(initially stationary) until they feel it matches the left belt (SI Appendix, Fig. S7B, "Ascend") (Leech

et al., 2018a; Rossi et al., 2019; Statton et al., 2018; Vazquez et al., 2015). This approximates

the second half of the Ramp Down (Fig. 4A, right leg feels equal or slower than left). Consistent

with this, we observed step length asymmetry aftereffects early in the speed match task (initial

SL asym. = 0.433 [0.271, 0.612], mean [CI]). Asymmetry decreased and was eventually near-

zero when the belt speeds felt equal (final SL asym. = 0.002 [-0.040, 0.045]; SI Appendix, Fig.

S8B).

We also used a "descend" speed match task where participants *decreased* the speed of the right

belt (initially fast as in adaptation) until they felt it matched the left (SI Appendix, Fig. S7B). This

approximated the first half of the Ramp Down. As expected, step length asymmetry was close to

zero for entire task (initial: -0.017 [-0.069, 0.039], final: 0.043 [-0.0004, 0.090], mean [CI]; SI

Appendix, Fig. S8B). The PSE for both speed match groups was comparable to that of Experiment

1 ($compensation_\text{perceptual}$ difference between Ramp Down lower bound and ascend or descend

Control = -3 [-16, 10]% or 4 [-12, 18]%, mean [CI]), and smaller than motor adaptation

($compensation_\text{motor total} - compensation_\text{perceptual}$ = 48 [39, 57]% or 56 [42, 70]% for ascend or

descend, mean [CI]; SI Appendix, Fig. S8A).

We replicated our findings in additional speed match experiments that varied in adaptation

duration (3, 15, or 30 minutes), perturbation magnitude (1m/s or 0.4m/s speed difference) and

schedule (abrupt or gradual) (SI Appendix, "Control experiments" and Fig. S7). Regardless of the

paradigm, the step length asymmetry and PSE were consistent with the recalibration + mapping

hypothesis (SI Appendix, Fig. S8 and Table S2-3).

<u>Experiment 2</u>

Experiment 1 demonstrated the presence of a stimulus-response mapping mechanism that can

produce Δ motor outputs that match a range of perturbations *smaller* than the adaptation

perturbation (in the Ramp Down task). In Experiment 2, we asked whether it can also produce Δ

motor outputs that match perturbations *larger* than the adaptation perturbation.

This sheds light on whether the stimulus-response mapping of walking adaptation operates akin

to memory-based or structure-based mechanisms observed in reaching adaptation. Failure to

account for larger perturbations would suggest that it is memory-based. That is, the Δ motor

outputs produced in adaptation may be cached in memory and later retrieved (Huberdeau et al.,

2019; McDougle and Taylor, 2019; Poggio and Bizzi, 2004; Wolpert et al., 2001). These Δ motor

outputs span a range of magnitudes that are smaller than the adaptation perturbation (see Box

1), so that they could match smaller but not larger perturbations.

Conversely, success in accounting for larger perturbations would suggest that the stimulus-

response mapping may be structure-based. That is, it may learn the general relationship between

perturbation and appropriate Δ motor output in adaptation and later use it to generate Δ motor

outputs anew (Bond and Taylor, 2017; Braun et al., 2009; McDougle and Taylor, 2019; Wolpert

et al., 2001). Hence, it would be able to generate Δ motor outputs to match either smaller or larger

perturbations.

Thus, we tested a "Ramp Up & Down" condition where the speed of the right belt was both gradually increased and then decreased after adaptation (Fig. 5A). The memory-based hypothesis predicts that step length asymmetry will become negative for perturbations *larger* than adaptation, while the structure-based hypothesis predicts it will remain close to zero (Fig. 5B inset, "predictions"). The rest of the paradigm was analogous to that of Experiment 1 except there was no perceptual assessment. Figure 5B shows step length asymmetry throughout the paradigm, and Figure 5C shows a close-up of the Ramp Up & Down task performance (group mean ± SE). The magenta portion of the task corresponds to the Ramp Down of Experiment 1, but the step length asymmetry differs because of the exposure to larger speed differences in the teal portion (a supplementary analysis confirmed that this is consistent with the recalibration + mapping hypothesis; see SI Appendix, "Ramp Down comparison between Experiments 1 and 2", Fig. S10-11 and Table S7).

On average, participants' step length asymmetry patterns did not remain zero for speed differences larger than adaptation (Fig. 5C, teal). However, we observed that individual participants exhibited markedly different patterns of step length asymmetry during this phase (SI Appendix, Fig. S12). We quantified this observation by evaluating, for each participant, the number of strides in this phase with step length asymmetry below their own baseline CI (SI Appendix, Table S8). We used a density-based analysis to formally assess whether there were separate clusters in our data (see Methods). Indeed, the algorithm detected two separate clusters of participants: for 12 participants, between 38 to 60 strides were asymmetric (out of 60 total strides); for the other 8 participants, only 3 to 21 strides were asymmetric (Fig. 5D-E insets, and SI Appendix, Fig. S13; difference in strides between subgroups = 36.083 [29.917, 42.250], mean [CI]). As a control, we performed the same clustering analysis for Experiment 1, and did not find separate clusters for any of our measures of interest (SI Appendix, "Experiment 1 clustering analysis", Fig. S13-14, and Table S9-10).

This result indicates that 12 of 20 participants could not account for belt speed differences larger than that of adaptation, suggesting that they used a memory-based mapping mechanism (Fig. 5D). In contrast, 8 of 20 participants could account for these speeds, suggesting that they engaged a structure-based mapping mechanism (Fig. 5E).

Figure 5. Experiment 2, Step length asymmetry. (A) Experimental protocol, equivalent to that of Experiment 1 except for the Ramp Up & Down part shaded in teal, where the right speed was faster than in adaptation (ramped up to 2m/s and back down to 1.5m/s). **(B)** Step length asymmetry time course (entire group mean ± SE). Background shade represents belt speed difference. Phases (except ramp tasks) are truncated to the participant with fewest strides. **Inset:** Predictions for the step length asymmetry during the teal portion of the Ramp Up & Down task, for the memory-based (top) or structure-based (bottom) mapping hypotheses. **(C)** Zoomed-in Ramp Up & Down task (entire group mean ± SE). Step length asymmetry for strides taken at right speeds larger than adaptation is shown in teal. **(D-E)** Separate plots of the step length asymmetry in the Ramp Up & Down task for the subgroups of participants that walked asymmetrically (D, "memory-based") versus symmetrically (E, "structure-based") in the teal portion of the task (subgroups mean ± SE). **Insets:** circles represent individual participants' number of strides, in the teal portion of the task, with step length asymmetry below their own baseline CI. Error bars depict subgroup mean ± SE. Subgroup assignment was performed by clustering on this measure.

We next aimed to exclude the possibility that participants in the structure-based subgroup may
simply be faster at adapting to new perturbations than those in the memory-based subgroup and
may be adapting to the new perturbations of the Ramp Up & Down rather than generating Δ motor
output using a previously learnt structure. To this end, we evaluated learning rates during
adaptation. We found that participants in the two subgroups adapted at similar rates (strides to
plateau difference, structure – memory: 135.875 [-53.208, 329.708], mean [CI]; SI Appendix, Fig.
S15-16), confirming that the different pattern of step length asymmetry in the Ramp Up & Down
task reflects different types of mapping in the subgroups.

We considered that the mapping adjustments described here may or may not be *deliberate* (i.e.,
participants are trying to correct for the perturbation) or done with an *explicit strategy* (i.e.,
participants can accurately report a relevant strategy that would counter the perturbation) (Long
et al., 2016; Roemmich et al., 2016). An example of an explicit strategy in visuomotor reaching
adaptation is when participants report that they aimed to offset a visual rotation (Bond and Taylor,
2017; Taylor et al., 2014); conceivable examples in split-belt treadmill adaptation may be reports
of "taking steps of similar length", "stepping further ahead with the right foot" or "standing on the
left foot for longer". We tested whether participants could explicitly report changes to the gait
pattern that specifically correct for the split-belt perturbation.

At the end of the experiment, participants were asked to report (in writing) if/how they had changed
the way they walked during adaptation (note that this was a later addition to the protocol and was
only collected in 16 of the 20 participants). We assessed reports by categorizing them in three
steps: 1) did the report mention *any* deliberate changes? 2) were the changes *relevant* to
adaptation? (i.e., did the report mention any gait metric contributing to the overall Δ motor output
or step length asymmetry adaptation in any amount?) 3) was the response *accurate?,* as
participants often reported strategies that they did not actually execute.

We summarize results from the questionnaire in Figure 6 (original responses are reported in SI
Appendix, Table S11). We found that, while 13 participants reported *deliberate* changes, only 6
people mentioned *relevant* aspects of the walking pattern (in particular, participants mentioned
"limping" or temporal coordination). Furthermore, only one of these participants reported an
accurate gait parameter ("I tried to spend as much time leaning on my left leg as possible"), while
the remainder of the relevant responses were inaccurate (e.g., "matched duration of standing on
each foot") or vague (e.g., "I adjusted as if I was limping"). The aspect of gait most frequently

reported across participants was stability (also reported as balance, not falling, or controlling
sway). This suggests that, while participants may deliberately adjust their body to feel more stable,
they do not seem to explicitly strategize how to offset the perturbation. Thus, mapping tends to
adjust aspects of the walking pattern that participants are not explicitly aware of controlling,
suggesting that this differs from explicit strategies often observed in reaching.

In sum, in Experiment 2 we found that 1) participants are largely unable to explicitly describe
adaptations to their walking pattern, and 2) some but not all participants can extrapolate their
walking pattern to account for larger perturbations. This sheds light on how the stimulus-response
mapping mechanism of walking adaptation may align with established mechanisms: it may differ
from explicit strategies and resemble memory-based caching in some people and structural
learning in others.

Figure 6. Summary of self-reported deliberate changes to the walking pattern in adaptation. Only one
participant accurately described changes to the walking pattern that related to adaptation, while other
responses were negative (i.e. no deliberate changes, 3 participants), irrelevant (7 participants), or
inaccurate (5 participants).

Discussion

In this study, we showed that locomotor adaptation involves two learning mechanisms: forward model recalibration and a stimulus-response mapping mechanism. The recalibration mechanism changes movement gradually and relates to the perceptual changes observed in locomotor adaptation. The mapping mechanism is flexible and, once learned, can change movement immediately to account for a range of belt speed configurations. Our data suggest that this mapping operates independently of explicit strategies, and that it can be memory-based or structure-based.

Forward model recalibration of movement and perception

In line with various adaptation studies, we showed that people recalibrated their perception in a way that reduced how perturbed they felt (Haith et al., 2008; Harris, 1963; Jensen et al., 1998; Moidell and Bedell, 1988; Sombric et al., 2019) – i.e., their perception of the leg speed difference diminishes (Jensen et al., 1998; Leech et al., 2018a; Rossi et al., 2019; Statton et al., 2018; Vazquez et al., 2015). The extent of the perceptual change correlated with the extent of the motor change that was achieved through forward model recalibration. Although previous research suggested that perceptual changes may be mediated by recalibration processes, the specific link to motor adaptation remained unclear ('t Hart and Henriques, 2016; Izawa et al., 2012; Sombric et al., 2019; Synofzik et al., 2008; Yavari et al., 2016; for a review, see Rossi et al., 2021a). Our findings demonstrate a direct relationship between perceptual and motor recalibrations, offering a novel approach to dissect forward model recalibration from stimulus-response mapping contributions to adaptation (SI Appendix, Supplementary Discussion 1).

Based on our findings, we suggest that forward model recalibration counters the perturbation in the motor and perceptual domains simultaneously – it adapts the walking pattern while realigning perception of leg speed. In support of this idea, perceptual realignment resembles a well-studied "sensory cancellation" phenomenon – where forward model predictions are used to filter redundant sensory information (Anderson et al., 2012; Blakemore et al., 1998). Forward model predictions may also be integrated with and sharpen proprioceptive estimates (Bhanpuri et al., 2013; Weeks et al., 2017), and may mediate a more complex integration of predicted and actual sensory signals contributing to perceptual changes. Consistent with the operation of a forward model thought to predict sensory consequences from efferent copies of motor commands (Haruno

et al., 2001; Ito, 1989; Wolpert et al., 2001), perceptual changes are expressed during active and

not passive movements (Sombric et al., 2019; Vazquez et al., 2015).

The cerebellum is thought to house forward models (Tanaka et al., 2020; Wolpert et al., 2001),

and plays a role in motor adaptation (Bastian, 2011; Martin et al., 1996b; Morton and Bastian,

2006), perceptual realignment (Izawa et al., 2012; Statton et al., 2018; Synofzik et al., 2008;

Yavari et al., 2016), as well as coordinating motor and sensory cortical activities (Lindeman et al.,

2020; Popa et al., 2013). Hence, motor and perceptual changes with adaptation may reflect

recalibration of forward models in the cerebellum processed downstream by motor and sensory

cortices. Neural substrates are further discussed in SI Appendix, Supplementary Discussion 2.

Flexible yet automatic stimulus-response mapping

To the best of our knowledge, this is the first study to show that a stimulus-response mapping

mechanism plays a role in walking adaptation. Unlike previous work employing a slow (10-minute)

post-adaptation perturbation ramp down (Roemmich and Bastian, 2015), our "Ramp Down" task

was quick (averaging 1 minute and 20 seconds) so that participants did not undergo washout (the

aftereffect magnitude after the Ramp Down was comparable to that measured post-adaptation in

prior work) (Reisman et al., 2005). This ruled out the possibility that changes in walking patterns

were due to washout, reaffirming the presence of a flexible mapping mechanism that

complemented the adjustments made by recalibration to produce a walking pattern well-suited to

the current treadmill configuration (SI Appendix, Supplementary Discussion 3).

Our results also suggest that this mapping mechanism may operate independently of explicit

control. In previous work where an explicit goal is given, adaptation mechanisms deployed in

addition to forward model recalibration are consistently found to operate under deliberate control

(Codol et al., 2018; Taylor et al., 2014; Taylor and Ivry, 2011) – including memory-based caching

(Huberdeau et al., 2019; McDougle and Taylor, 2019) and structural learning mechanisms (Bond

and Taylor, 2017; McDougle and Taylor, 2019). Mounting evidence indicates that explicit

strategies do not play a strong role in split-belt walking adaptation where an explicit movement

goal is not provided to the participants; for example, people do not adapt faster even after

watching someone else adapt (Song et al., 2020). Strategic adjustments to the walking pattern

can be temporarily elicited by providing additional visual feedback of the legs and an explicit goal

of how to step. Yet, these adjustments disappear immediately upon removal of the visual
feedback (Roemmich et al., 2016), and have no effect on the underlying adaptive learning process
(Long et al., 2016; Malone and Bastian, 2010; Roemmich et al., 2016). Furthermore, performing
a secondary cognitive task during walking adaptation does not affect the amount of motor learning
(Hinton et al., 2020; Malone and Bastian, 2010; Rossi et al., 2021b; Vervoort et al., 2019). Here,
we show that explicit strategies are not systematically used to adapt step length asymmetry and
Δ motor output: the participants in our study either did not know what they did, reported changes
that did not actually occur or would not lead symmetry. Only one person reported "leaning" on the
left (slow) leg for as much time as possible, which is a relevant but incomplete description for how
to walk with symmetry (SI Appendix, Supplementary Discussion 4).

Recalibration-adjusted perceptual information on the treadmill configuration may be relayed via
sensory cortical areas and cerebellum to motor cortical areas, where it may be mapped to motor
adjustments. This is supported by the interconnectivity between these substrates (Judd et al.,
2021; Kandel et al., 2013; Sultan et al., 2012), and by the cortical contribution to the flexible control
of walking (Drew and Marigold, 2015; Reisman et al., 2010) beyond voluntary control (Delval et
al., 2020; Petersen et al., 2012). Mapping may also involve spinal control – which is rapid and
automatic (Takakusaki, 2013) and can play a role in split-belt walking (Ogawa et al., 2014;
Vasudevan et al., 2011) – but likely through connection with supraspinal structures due to the
extensive training needed for plasticity within the spinal cord (Iturralde and Torres-Oviedo, 2019;
Thompson et al., 2009) (SI Appendix, Supplementary Discussion 5).

In sum, the mapping mechanism combines the advantages of automaticity and flexibility
(Huberdeau et al., 2015; Taylor et al., 2014; Taylor and Ivry, 2011). This is ecologically important
for both movement accuracy (Uiga et al., 2020; Wong et al., 2008) and for walking safely in real-
world situations, where we walk while talking or doing other tasks, and terrains are uneven (Clark,
2015; Paul et al., 2005; Wong et al., 2008) (SI Appendix, Supplementary Discussion 6).

Mapping operates as memory-based in some people, structure-based in others

Results from Experiment 2 highlight individual differences in the learning mechanisms underlying
generalization to unexperienced belt speed differences (usage of the term "generalization" is
discussed in SI Appendix, Supplementary Discussion 7). The generalization to novel perturbation

sizes observed here is in line with previous suggestions of "meta-learning" in the savings of
walking adaptation (i.e., faster relearning when exposed to a different perturbation) (Leech et al.,
2018b; Malone et al., 2011). Generalization to *larger* perturbations after reaching adaptation was
shown to be incomplete (Abeele and Bock, 2001; Lazar and Van Laer, 1968), and it was unclear
whether the movement patterns had been *extrapolated* beyond what had been experienced, or
rather was just the same as in adaptation. Surprisingly, we found 40%-60% divide in our
participants regarding the capacity to extrapolate walking patterns to account for larger
perturbations.

We suggest that structural learning may underlie the ability to extrapolate walking patterns and
walk symmetrically for belt speed differences larger than adaptation, as seen in 8 of 20
participants. Indeed, generalization in reaching adaptation may rely on a process of structural
learning (Bond and Taylor, 2017; Braun et al., 2009) (note, however, that this was tied to explicit
aiming; Bond and Taylor, 2017). Our participants may have learned the scaling relationship
between belt speed perturbation and the walking pattern needed to walk symmetrically and use
this scaling to produce new walking patterns matching larger perturbations. In contrast, memory-
based theories of mapping (Dassonville et al., 2001; Poggio and Bizzi, 2004; Van Vugt and Ostry,
2018; Wolpert et al., 2001) may explain why 12 participants walked asymmetrically for
perturbations larger than adaptation, despite generalizing to smaller perturbations. Here, the
walking patterns may be stored in memory during adaptation in association with the amount of
perturbation they correct, and then retrieved during the ramp tasks. People transition through a
range of walking patterns in adaptation (see gradual transition from baseline to adapted Δ motor
outputs in Box 1 and Fig. 1), so storage of intermediate outputs may reflect generalization to
smaller perturbations (while larger Δ motor outputs are not experienced). SI Appendix,
Supplementary Discussion 8, discusses potential *interpolation* computations.

630
Conceptual model

We propose a comprehensive conceptual model of walking adaptation that captures key
behavioral properties uncovered in our study. We build upon standard models (Shadmehr and
Krakauer, 2008; Smith et al., 2006) and introduce a mechanism and architecture for adaptive
learning that accounts for the flexibility of the walking pattern and the relationship between motor
and perceptual changes. This conceptual model is illustrated in Figure 7.

The perturbation p represents the effect of the belt speed difference on movement or perception.

Note that motor and perceptual quantities are represented in the same relative units (relative to

the perturbation).

The light blue "recalibration" box represents the operation of the forward model recalibration

mechanism: this mechanism is not flexible, so that it produces the same output x_r for any

perturbation p. The recalibration x_r may project to both sensory integration and motor control

areas (green and blue boxes), where it may be used to recalibrate perception and movement.

Figure 7. Schematic model of adaptation. Body movement depends on environment perturbations (red)
and Δ motor output (blue). The Δ motor output is adjusted by recalibration (light blue) and mapping (dark
blue) mechanisms, which perform different operations and are arranged in tandem. We propose the
following architecture and flow: **(1. Recalibration)** The recalibration mechanism produces adjustment x_r
that is fixed regardless of perturbation size (light blue box, x_r is constant for varying p). The *same*
recalibration adjustment x_r serves as an input to *both* areas responsible for conscious perception (green
box) and Δ motor output (blue box). **(2. Perception)** Conscious perception is computed by cancelling out
the recalibration adjustment from the actual sensory feedback (green box, perception of the belt speed
difference perturbation $\tilde{p}$ is the difference between the actual speed difference p and recalibration x_r). The
perceived perturbation $\tilde{p}$ serves as an input to the mapping mechanism (dark blue box). **(3. Mapping)** The
mapping mechanism produces adjustment x_m that can vary in magnitude to appropriately account for the
perceived perturbation $\tilde{p}$ (dark blue box, x_m scales with $\tilde{p}$ and matches its magnitude). **(4. Δ Motor output)**
The overall adjustment to Δ motor output is computed by adding the mapping adjustment x_m and
recalibration adjustment x_r. The corner in the Δ motor output versus perturbation profile arises because
mapping is computed based on the perceived perturbation $\tilde{p}$ (not the actual perturbation p) and is only
learnt for positive $\tilde{p}$ (the experienced direction). When the perturbation is perceived to be opposite than
adaptation, even if it is not, mapping is zero and the Δ motor output is constant, reflecting recalibration
adjustments only (blue box, when $\tilde{p} < 0$ and $p \geq 0$ the mapping adjustment x_m is zero and $u = x_r$).

The green "perception" box represents the process of perceptual realignment. It receives sensory

information regarding the perturbation, p, and the forward model recalibration output, x_r. It

cancels out the portion of the perturbation predicted by the forward model, so that it is output is

the perceived perturbation $\tilde{p} = p - x_r$. The perceived perturbation $\tilde{p}$ may serves both as signal

for conscious speed perception (that captured by perceptual tests), and as an input to the

stimulus-response mapping mechanism (dark blue box).

The dark blue "mapping" box represents the operation of the stimulus-response mapping

mechanism. It receives the perceived perturbation signal $\tilde{p}$ and transforms into a motor

adjustment appropriate to counter the perceived perturbation, $x_m = \tilde{p}$.

The blue "Δ motor output" box represent the computation of the Δ motor output. It receives the

recalibration x_r and mapping x_m adjustments as inputs, and sums them to compute the Δ motor

output $u = x_m + x_r$.

We propose the following series of computations. First, the recalibration is used to compute the

perceived perturbation (green box). Second, the perceived perturbation is relayed to the mapping

mechanism and used to compute the mapping-related motor adjustment (dark blue box). Third,

recalibration-related and mapping-related motor adjustments are summed to produce the Δ motor

output (blue box).

In conclusion, our model proposes three key features of the mapping mechanism that distinguish

it from standard adaptation models. First, the mapping produces near-immediate changes to

motor output in response to varying perturbations – contrasting the gradual changes proposed by

state-space models (Herzfeld et al., 2014; Roemmich et al., 2016; Smith et al., 2006). Second, it

accesses the perceived perturbation signal which integrates external sensory information with

internal forward model state – contrasting the unimodal input proposed by optimal control

(Shadmehr and Krakauer, 2008; Todorov, 2005). Third, the mapping operates automatically and

is learnt with adaptation – contrasting the readily implementable explicit strategies modeled for

reaching (Taylor and Ivry, 2011). These features enable automatic motor adjustments that

complement recalibration to match varying perturbations, allowing our model to account for the

observed symmetric walking across different treadmill speed configurations (SI Appendix,

Supplementary Discussion 9).

Limitations

We aim to propose a framework for motor adaptation that is both parsimonious and based on a
simple model. However, this simplicity inevitably comes with certain limitations. First, our model
is descriptive and focused on capturing the novel features of the data revealed by the Ramp Down
task, rather than the entire motor adaptation time series. Additionally, there may be alternative
interpretations to our results, and a few mechanisms may underly the stimulus-response mapping
observed here. As described in SI Appendix, Supplementary Discussion 10, there some evidence
to suggest that walking adaptation involves optimizing motor commands to minimize energy costs
(Finley et al., 2013; Sánchez et al., 2021, 2019, 2017). Therefore, our results may be explained
by extending current optimal control models to account for our finding that motor commands are
computed using internal state estimates and sensory feedback from the environment in tandem.
Future work is needed to develop a generative model capable of capturing the features of
adaptation presented here, and further characterize the nature of the stimulus-response mapping
mechanism of automatic adaptation.

Conclusions and future directions

We here characterized two distinct learning mechanisms involved in walking adaptation, both of
which are not under explicit control: recalibration – which leads to motor and perceptual
aftereffects, and mapping – which only changes movement, does not contribute to aftereffects,
and shows meta-learning to different perturbation sizes. Future work should explore whether the
mapping mechanism described here could be of clinical significance: given the combined flexibility
and automaticity, mapping has the potential to ameliorate known problems such as transitioning
between walking environments (Sombric et al., 2017) and dual-tasking (Clark, 2015).
Furthermore, our findings that different people adapt using different learning mechanisms (i.e.,
memory or structure-based mapping) highlights the importance of assessing individual
characteristics of adaptation for both understanding of the neural mechanisms as well as
translation to rehabilitation. Finally, as detailed in SI Appendix, Supplementary Discussions 9-11,
the framework proposed here has important implications for the development of computational
models of adaptation, and may help reconcile different findings related to savings, energetics,
and sources of errors.

Methods

<u>Participants</u>

We recruited one-hundred adults (66 females, 23.6 ± 3.9 years old, mean ± SD) for this study, and we reanalyzed data from ten additional adults collected in a previous study (Leech et al., 2018a) (9 female, 21.3 ± 2.9 years old). The protocol was approved by the Johns Hopkins Institutional Review Board and participants provided written informed consent. Participants had no known neurological or musculoskeletal disorders, were naïve to split-belt walking, and participated in only one of the nine experiments.

<u>Data collection</u>

Participants walked on a split-belt treadmill (Woodway, Waukesha, WI, USA) and belt speeds were controlled using a custom Vizard program (WorldViz). Kinematic data were collected using infrared-emitting markers (Optotrak, Northern Digital, Waterloo, ON, Canada) at 100Hz (SI Appendix, Supplementary Methods).

<u>Ramp tasks</u>

For all ramp tasks of Experiments 1 and 2, the speed of the right belt was changed by 0.05m/s every 3 strides (defined below) during right leg swing. The baseline ramp of both experiments consisted of 7 increasing right speeds from 0.35m/s to 0.65m/s. The post-adaptation Ramp Down of Experiment 1 consisted of 21 decreasing right speeds from 1.5m/s to 0.5m/s. The post-adaptation Ramp Up & Down of Experiment 2 consisted of 41 total right speeds: 11 increasing speeds from 1.5m/s to 2m/s, followed by 30 decreasing speeds from 1.95m/s to 0.5m/s. The left belt speed was constant at 0.5 m/s for all ramp tasks. The treadmill was not stopped before the ramp tasks (people transitioned directly from the preceding walking blocks into the ramp) but was briefly stopped after each ramp task. We decided to not pause the treadmill between adaptation and the post-adaptation ramp tasks, and to make the tasks relatively quick – the Ramp Down was ~80 seconds on average – to limit unwanted forgetting and/or unlearning. We occluded vision and sound of the belt speeds using a cloth drape and headphones playing white noise.

In Experiment 1, a keyboard was placed on the treadmill handrail. In the post-adaptation Ramp Down task, participants were asked to press a button the first time they perceived the right belt to be (1) as fast as the left, and (2) faster than the left. In the baseline ramp, they press to report the right belt feeling (1) as fast as the left, and (2) slower than the left. This was explained prior to the experiment and the above prompts were displayed during the tasks on a television screen placed in front of the treadmill.

Questionnaire

At the end of Experiment 2, participants answered this question on a computer: *Did you deliberately change how you walked to account for how fast the belts were moving? If so, describe how. Note: deliberately means that you thought about and decided to move that way. The question refers to the entire central ~20min walking block.* To ensure participants remembered what phase they were asked about, before adaptation we told participants that the following block will be called "central ~20min walking block".

Motor measures

We defined a stride as the period between two consecutive left heel strikes (LHS_1 to LHS_2), and computed the motor measures for each stride as in previous work (Finley et al., 2015; Sombric et al., 2017):

$$step\ length\ asymmetry\ =\ \frac{1}{stride\ length} * \Delta step\ length \tag{9}$$

$$perturbation\ =\ \frac{1}{stride\ length} * \overline{step\ time} * \Delta step\ velocity \tag{10}$$

$$\Delta\ motor\ output\ =\ \frac{1}{stride\ length} * \left(\Delta step\ position - \overline{step\ velocity} * \Delta step\ time\right) \tag{11}$$

The terms in the equations above are defined as follows:

$\Delta step\ length\ =\ R\ step\ length\ -\ L\ step\ length$ (12)

$stride\ length\ =\ R\ step\ length\ +\ L\ step\ length$ (13)

Where right (R) and left (L) step lengths are the anterior-posterior distance between the ankle
markers of the two legs at right heel strike (RHS) and left heel strike (LHS_2) respectively.

$\Delta step\ position\ =\ \left(R\ step\ position_{RHS}\ -\ L\ step\ position_{LHS_1}\right)$
784                   $-\left(L\ step\ position_{LHS_2}\ -\ R\ step\ position_{RHS}\right)$ (14)

Where step position is the anterior-posterior position of the ankle marker, relative to average of
the two hip markers, at heel strike of the same leg.

$\Delta step\ time\ =\ R\ step\ time\ -\ L\ step\ time$ (15)

$\overline{step\ time}\ =\ \dfrac{R\ step\ time\ +\ L\ step\ time}{2}$ (16)

Where left and right step times are the times from LHS_1 to RHS, and from RHS to LHS_2
respectively.

$\Delta step\ velocity\ =\ R\ step\ velocity\ -\ L\ step\ velocity$ (17)

$\overline{step\ velocity}\ =\ \dfrac{R\ step\ velocity\ +\ L\ step\ velocity}{2}$ (18)

Where step velocity is the average anterior-posterior velocity of the ankle marker relative to

average of the two hip markers, over the duration of the step (left: LHS_1 to RHS, right: RHS to

LHS_2).

In Experiment 2, we also computed the measure of "strides to plateau" using individual step length

asymmetry data from the adaptation phase only. We first smoothed the data with a 5-point moving

average filter. We then calculated the number of strides till five consecutive strides fell within the

"plateau range", defined as the mean ± 1SD of the last 30 strides of adaptation (Malone and

Bastian, 2010; Rossi et al., 2019).

<u>Clustering analysis</u>

We tested to see if there were separate clusters of participants in our dataset. For Experiment 1,

the measure used for clustering was the number of strides in the Ramp Down with step length

asymmetry above the baseline CI. For Experiment 2, it was the number of strides in the first

portion of the Ramp Up & Down (first 60 strides, teal in Fig. 5) with step length asymmetry below

the baseline CI. For both experiments, "baseline CI" was computed separately for each participant

as the 95% CI of the mean of step length asymmetry data in the second baseline tied-belt block

(after the baseline ramp task; see Statistical Analysis).

We used the MATLAB "dbscan" density-based clustering algorithm (Ester et al., 1996) and

automated the choice of parameters adapting previously published procedures (Naik Gaonkar

and Sawant, 2013; Rahmah and Sitanggang, 2016) (SI Appendix, Supplementary Methods and

pseudocode). The resulting algorithm does not require any user input; it automatically assigns

participants to clusters (whose number is not predefined), or labels them as outliers, based solely

on each participant's measure of interest.

<u>Data fitting</u>

In Experiment 1, we fitted different models to individual participants' Δ motor output data in the

Ramp Down. We developed and fitted our own recalibration + mapping model, presented in the

Results section in Eq. 3. A reasonable upper limit for the parameter r would be the magnitude of

the perturbation in adaptation, but this motor measure differs across participants, which would

lead to inconsistencies in the data fitting analysis. We therefore reformulated the recalibration and

recalibration + mapping models to use a normalized parameter r_{norm}:

829   $r = r_{\text{norm}} * p_{\text{plateau}}$              (19)

831   Forward Model Recalibration: $u(p) = r_{\text{norm}} * p_{\text{plateau}}$      (20)

833   Recalibration + mapping: $u(p) = \begin{cases} p & , \text{ for } p \geq r_{\text{norm}} * p_{\text{plateau}} \\ r_{\text{norm}} * p_{\text{plateau}} & , \text{ otherwise} \end{cases}$  (21)

835 Where p_{plateau} is mean perturbation over the last 30 strides of adaptation, computed separately

836 for each participant, p is the perturbation in the Ramp Down, and u is the modelled Δ motor

837 output. We fitted the normalized parameter r_{norm} using initial parameter value = 0.5 and bounds

838 = [0, 1] for all participants. We then used Eq. 19 to compute the unnormalized value r.

839 For the recalibration only hypothesis, we fitted the dual state model defined as in Smith et al.

840 (Smith et al., 2006):

842   net internal estimate: $x(k) = x_f(k) + x_s(k)$      (22)

844   fast state:     $x_f(k+1) = A_f * x_f(k) + B_f * \big(p(k) - x(k)\big)$   (23)

846   slow state:     $x_s(k+1) = A_s * x_s(k) + B_s * \big(p(k) - x(k)\big)$   (24)

848   Δ motor output:  $u(k) = x(k)$        (25)

850 Where k is the stride number, p is the perturbation, x is the net internal estimate of the

851 perturbation, x_f and x_s are the fast and slow states contributing to the estimate. Participants

852 generate Δ motor output u that equals the internal estimate of the perturbation x to minimize the

853 predicted step length asymmetry. The model has four free parameters: A_f, A_s are retention

854 factors controlling forgetting rate, and B_f, B_s are error sensitivities controlling learning rate. We

855 fitted the model parameters using initial values A_f=0.92, A_s=0.99, B_f=0.1, and B_s=0.01 (as in

Roemmich et al., 2016), and constraints $0 < A_f < A_s < 1$ and $0 < B_f > B_s < 1$ (as defined by the model,

Smith et al., 2006).

We also fitted two additional models that are prominent in literature: optimal feedback control

(Izawa and Shadmehr, 2011; Shadmehr and Krakauer, 2008; Todorov, 2004) and memory of

errors (Herzfeld et al., 2014). The details for these models are provided in SI Appendix,

"Experiment 1 additional models".

We fitted each model to individual participants' Δ motor output data in the Ramp Down.

Specifically, we found parameter values that minimize the residual sum of squares between the

Ramp Down Δ motor output data and the modelled u. Dual state, optimal control, and memory of

errors models involve hidden states that depend on the history of perturbations and are not

expected to be zero at the start of the Ramp Down. To account for this, we simulated these models

over the entire paradigm, but computed the residual sum of squares solely on the Ramp Down to

ensure fair model comparison with the recalibration + mapping model. We used the *fmincon*

MATLAB function with constraint and optimality tolerances tightened to 10^-20 (as in Roemmich

et al., 2016).

<u>Statistical analysis</u>

Statistical tests were performed in MATLAB with significance level α=0.05 (two-sided). For

Experiment 1, we performed within-group statistical analyses to compare the following measures

to zero. For each participant and for each speed in the ramp tasks, we first averaged step length

asymmetry over the 3 strides taken at that speed to obtain the measures listed in a) and b):

a) Step length asymmetry for each of the 7 speeds in the baseline ramp task (m=7)

b) Step length asymmetry for each of the 21 speeds in the Ramp Down task (m=21)

c) Difference in BIC between our model (recalibration + mapping) and each of the alternative

models (dual state, optimal feedback control, memory of errors) (m=3)

d) Difference between $compensation_\text{motor total}$ or $compensation_\text{motor recalibration}$, and the upper

and lower bounds of $compensation_\text{perceptual}$ (m=4)

Each item on the list represents a "family of related tests" (measures considered together for
multiple comparisons), where "m" is the number of tests in the family. We computed bootstrap
distributions of each measure by generating 10,000 bootstrapped samples of 20 participants
(resampled with replacement from Experiment 1) and averaging the measure over participants in
each sample (Efron and Tibshirani, 1994). We then computed confidence intervals (CI) corrected
for multiple comparisons using the False Discovery Rate procedure (FDR) (Benjamini and
Yekutieli, 2005). That is, for each family of related tests, we adjusted the significance level to
$\alpha_{corr} = \alpha * \frac{R}{m} = 0.05 * \frac{R}{m}$, where "m" is the total number of tests and "R" is number of tests deemed
significant for $\alpha = 0.05$. A test is significant if the CI does not overlap zero, indicating that the
measure is significantly different from zero (Cumming and Finch, 2005; Efron and Tibshirani,
1994).

For both experiments, we computed within-participant CIs for the mean baseline step length
asymmetry. For each participant, we generated 10,000 bootstrapped samples of N strides,
resampled with replacement from 2-min baseline block following the ramp task, where N is the
number of strides in this phase. We averaged step length asymmetry over strides in each sample
and computed the 95% CI.

For Experiment 2, we performed between-group statistical analyses to assess whether the
following measures differed between the memory-based and structure-based subgroups:

a) Number of strides to plateau (m=1)

b) Number of strides in teal portion of the Ramp Up & Down (first 60 strides) with step length
asymmetry below the within-participant baseline CI (as described above) (m=1)

We generated 10,000 bootstrapped samples, each comprising 20 participants: 12 from the
memory-based subgroup and 8 from the structure-based subgroup (resampled with
replacement). For each sample "b", we averaged the measure of interest over participants
resampled from each subgroup to obtain $\mu_{memory}(b)$ and $\mu_{structure}(b)$, and evaluated the
difference of the means between the subgroups: $\Delta\mu(b) = \mu_{memory}(b) - \mu_{structure}(b)$. We computed
the CI for this difference, correcting for multiple comparisons using FDR as explained for
Experiment 1. Note that we report FDR-corrected CIs in the Results sections.

Data Availability

Note to reviewers: the dataset cited below is currently "private for peer review", and can be accessed by the alternative "reviewer URL" provided by the Dryad database:

https://datadryad.org/stash/share/Qa0pD7rMFT7kVgDeP-8PMYKBaCNuy4yx1VGlQlpVgvM

All data and code used for the study have been deposited in: Rossi, Cristina (2023), Dataset for automatic learning mechanisms for flexible human locomotion, Dryad, Dataset, https://doi.org/10.5061/dryad.18931zd27.

Acknowledgments

Supported by NIH grant 5 R37 NS090610 to AJB, American Heart Association predoctoral fellowship 20PRE35180131 to CR, and NIH grant K01 AG073467 to KAL. We thank Daniel Wolpert, Adrian Haith, Jonathan Tsay, and Amanda Therrien for helpful discussions and comments on data analysis.

Author Contributions: Study design & implementation: C.R., K.A.L., R.T.R., A.J.B.; Data acquisition & analysis: C.R., K.A.L.; Interpretation of results: C.R., K.A.L., R.T.R., A.J.B.; Writing & revision: C.R., K.A.L., R.T.R., A.J.B.

Competing Interest Statement: The authors declare no competing interests.

Classification: Biological Sciences: Neuroscience.

Keywords: Adaptation. Motor control. Motor learning. Human locomotion. Perception.

References

't Hart BM, Henriques DYP. 2016. Separating predicted and perceived sensory consequences of motor learning. *PLoS One* **11**:e0163556. doi:10.1371/journal.pone.0163556

Abeele S, Bock O. 2001. Mechanisms for sensorimotor adaptation to rotated visual input. *Exp Brain Res* **139**:248–253. doi:10.1007/S002210100768

Anderson SR, Porrill J, Pearson MJ, Pipe AG, Prescott TJ, Dean P. 2012. An Internal Model Architecture for Novelty Detection: Implications for Cerebellar and Collicular Roles in Sensory Processing. *PLoS One* **7**:e44560. doi:10.1371/journal.pone.0044560

Bastian AJ. 2011. Moving, sensing and learning with cerebellar damage. *Curr Opin Neurobiol* **21**:596–601. doi:10.1016/j.conb.2011.06.007

Bastian AJ. 2008. Understanding sensorimotor adaptation and learning for rehabilitation. *Curr Opin Neurol* **21**:628–633. doi:10.1097/WCO.0b013e328315a293

Benjamini Y, Yekutieli D. 2005. False discovery rate-adjusted multiple confidence intervals for selected parameters. *J Am Stat Assoc* **100**:71–81. doi:10.1198/016214504000001907

Bhanpuri NH, Okamura AM, Bastian AJ. 2013. Predictive modeling by the cerebellum improves proprioception. *J Neurosci* **33**:14301–14306. doi:10.1523/JNEUROSCI.0784-13.2013

Blakemore S-J, Wolpert DM, Frith CD. 1998. Central cancellation of self-produced tickle sensation. *Nat Neurosci* **1**:635–640.

Bond KM, Taylor JA. 2017. Structural learning in a visuomotor adaptation task is explicitly accessible. *eNeuro* **4**:e0122-17. doi:10.1523/ENEURO.0122-17.2017

Braun DA, Aertsen A, Wolpert DM, Mehring C. 2009. Motor Task Variation Induces Structural Learning. *Curr Biol* **19**:352–7. doi:10.1016/j.cub.2009.01.036

Clark DJ. 2015. Automaticity of walking: Functional significance, mechanisms, measurement and rehabilitation strategies. *Front Hum Neurosci* **9**:246. doi:10.3389/fnhum.2015.00246

Codol O, Holland PJ, Galea JM. 2018. The relationship between reinforcement and explicit control during visuomotor adaptation. *Sci Rep* **8**:9121. doi:10.1038/s41598-018-27378-1

Cumming G, Finch S. 2005. Inference by eye: confidence intervals and how to read pictures of data. *Am Psychol* **60**:170–180. doi:10.1037/0003-066X.60.2.170

Dassonville P, Lewis SM, Zhu XH, Ugurbil K, Kim SG, Ashe J. 2001. The effect of stimulus-response compatibility on cortical motor activation. *Neuroimage* **13**:1–14. doi:10.1006/nimg.2000.0671

Delval A, Bayot M, Defebvre L, Dujardin K. 2020. Cortical oscillations during gait: Wouldn't walking be so automatic? *Brain Sci*. doi:10.3390/brainsci10020090

DiCiccio TJ, Efron B. 1996. Bootstrap Confidence Intervals. *Stat Sci* **11**:189–228. doi:10.1007/978-3-319-01931-4_24

Drew T, Marigold DS. 2015. Taking the next step: Cortical contributions to the control of locomotion. *Curr Opin Neurobiol* **33**:25–33. doi:10.1016/j.conb.2015.01.011

Efron B, Tibshirani RJ. 1994. An Introduction to the Bootstrap, An Introduction to the Bootstrap. doi:10.1201/9780429246593

Ester M, Kriegel H-P, Sander J, Xu X. 1996. A Density-Based Algorithm for Discovering Clusters in Large Spatial Databases with NoiseProceedings of the 2nd International Conference on Knowledge Discovery and Data Mining. pp. 226–231.

Finley JM, Long A, Bastian AJ, Torres-Oviedo G. 2015. Spatial and Temporal Control Contribute to Step Length Asymmetry during Split-Belt Adaptation and Hemiparetic Gait. *Neurorehabil Neural Repair* **29**:786–795. doi:10.1177/1545968314567149

Haith A, Jackson C, Miall C, Vijayakumar S. 2008. Unifying the sensory and motor components of sensorimotor adaptationAdvances in Neural Information Processing Systems 21 - Proceedings of the 2008 Conference. pp. 593–600.

Harris CS. 1963. Adaptation to Displaced Vision: Visual, Motor, or Proprioceptive Change? *Science (80-)* **140**:812–813. doi:10.1017/CBO9781107415324.004

Haruno M, Wolpert DM, Kawato M. 2001. MOSAIC Model for Sensorimotor Learning and Control. *Neural Comput* **13**:2201–2220. doi:10.1162/089976601750541778

Herzfeld DJ, Vaswani PA, Marko MK, Shadmehr R. 2014. A memory of errors in sensorimotor learning. *Science (80-)* **345**:1349–1353. doi:10.1126/science.1253138

Hinton DC, Conradsson D, Bouyer L, Paquette C. 2020. Does dual task placement and duration affect
split-belt treadmill adaptation? *Gait Posture* **75**:115–120. doi:10.1016/j.gaitpost.2019.10.005
Huberdeau DM, Krakauer JW, Haith AM. 2019. Practice induces a qualitative change in the memory
representation for visuomotor learning. *J Neurophysiol* **122**:1050–1059. doi:10.1152/jn.00830.2018
Huberdeau DM, Krakauer JW, Haith AM. 2015. Dual-process decomposition in human sensorimotor
adaptation. *Curr Opin Neurobiol* **33**:71–77. doi:10.1016/j.conb.2015.03.003
Ito M. 1989. Long-term depression. *Annu Rev Neurosci* **12**:85–102.
Iturralde PA, Torres-Oviedo G. 2019. Corrective Muscle Activity Reveals Subject-Specific Sensorimotor
Recalibration. *eNeuro* **6**. doi:10.1523/ENEURO.0358-18.2019
Izawa J, Criscimagna-Hemminger SE, Shadmehr R. 2012. Cerebellar contributions to reach adaptation
and learning sensory consequences of action. *J Neurosci* **32**:4230–4239.
doi:10.1523/JNEUROSCI.6353-11.2012
Izawa J, Shadmehr R. 2011. Learning from sensory and reward prediction errors during motor adaptation.
PLoS Comput Biol **7**:e1002012. doi:10.1371/journal.pcbi.1002012
Jensen L, Prokop T, Dietz V. 1998. Adaptational effects during human split-belt walking: influence of
afferent input. *Exp Brain Res* **118**:126–130. doi:10.1007/s002210050262
Judd EN, Lewis SM, Person AL. 2021. Diverse inhibitory projections from the cerebellar interposed
nucleus. *Elife* **10**:e66231. doi:10.7554/eLife.66231
Kandel E, Schwartz JH, Jessell TM, Siegelbaum SA, Hudspeth AJ. 2013. Principles of Neural Science,
Fifth Edit. ed. McGraw-Hill, New York.
Lazar G, Van Laer J. 1968. Adaptation to displaced vision after experience with lesser displacements.
Percept Mot Skills **26**:579–582. doi:10.2466/pms.1968.26.2.579
Leech, Day KA, Roemmich RT, Bastian AJ. 2018a. Movement and perception recalibrate differently
across multiple days of locomotor learning. *J Neurophysiol* **120**:2130–2137.
doi:10.1152/jn.00355.2018
Leech, Roemmich RT, Bastian AJ. 2018b. Creating flexible motor memories in human walking. *Sci Rep*
8:94. doi:10.1038/s41598-017-18538-w
Lindeman S, Hong S, Kros L, Mejias JF, Romano V, Oostenveld R, Negrello M, Bosman LWJ, de Zeeuw
CI. 2020. Cerebellar purkinje cells can differentially modulate coherence between sensory and
motor cortex depending on region and behavior. *Proc Natl Acad Sci U S A* **118**:2015292118.
doi:10.1073/pnas.2015292118
Long AW, Roemmich RT, Bastian AJ. 2016. Blocking trial-by-trial error correction does not interfere with
motor learning in human walking. *J Neurophysiol* **115**:2341–2348. doi:10.1152/jn.00941.2015
Malone LA, Bastian AJ. 2010. Thinking about walking: Effects of conscious correction versus distraction
on locomotor adaptation. *J Neurophysiol* **103**:1954–1962. doi:10.1152/jn.00832.2009
Malone LA, Vasudevan EVL, Bastian AJ. 2011. Motor adaptation training for faster relearning. *J Neurosci*
31:15136–15143. doi:10.1523/JNEUROSCI.1367-11.2011
Martin TA, Keating JG, Goodkin HP, Bastian AJ, Thach WT. 1996a. Throwing while looking through
prisms II. Specificity and storage of multiple gaze-throw calibrations. *Brain* **119**:1199–1211.
doi:10.1093/brain/119.4.1183
Martin TA, Keating JG, Goodkin HP, Bastian AJ, Thach WT. 1996b. Throwing while looking through
prisms I. Focal olivocerebellar lesions impair adaptation. *Brain* **119**:1183–1198.
doi:10.1093/brain/119.4.1183
McDougle SD, Taylor JA. 2019. Dissociable cognitive strategies for sensorimotor learning. *Nat Commun*
10:1–11. doi:10.1038/s41467-018-07941-0
Moidell BG, Bedell HE. 1988. Changes in oculocentric visual direction induced by the recalibration of
saccades. *Vision Res* **28**:329–336. doi:10.1016/0042-6989(88)90161-7
Morton SM, Bastian AJ. 2006. Cerebellar contributions to locomotor adaptations during splitbelt treadmill
walking. *J Neurosci* **26**:9107–9116. doi:10.1523/JNEUROSCI.2622-06.2006
Naik Gaonkar M, Sawant K. 2013. AutoEpsDBSCAN : DBSCAN with Eps Automatic for Large Dataset. *Int*
J Adv Comput Theory Eng **2**:2319–2526.
Ogawa T, Kawashima N, Ogata T, Nakazawa K. 2014. Predictive control of ankle stiffness at heel contact
is a key element of locomotor adaptation during split-belt treadmill walking in humans. *J*
Neurophysiol **111**:722–732. doi:10.1152/jn.00497.2012

Paul SS, Ada L, Canning CG. 2005. Automaticity of walking–implications for physiotherapy practice. *Phys*
Ther Rev **10**:15–23. doi:10.1179/108331905X43463
Petersen TH, Willerslev-Olsen M, Conway BA, Nielsen JB. 2012. The motor cortex drives the muscles
during walking in human subjects. *J Physiol* **590**:2443–2452. doi:10.1113/jphysiol.2012.227397
Poggio T, Bizzi E. 2004. Generalization in vision and motor control. *Nature* **431**:768–774.
doi:10.1038/nature03014
Popa D, Spolidoro M, Proville RD, Guyon N, Belliveau L, Léna C. 2013. Functional role of the cerebellum
in gamma-band synchronization of the sensory and motor cortices. *J Neurosci* **33**:6552–6556.
doi:10.1523/JNEUROSCI.5521-12.2013
Rahmah N, Sitanggang IS. 2016. Determination of Optimal Epsilon (Eps) Value on DBSCAN Algorithm to
Clustering Data on Peatland Hotspots in SumatraIOP Conference Series: Earth and Environmental
Science. Institute of Physics Publishing. p. 012012. doi:10.1088/1755-1315/31/1/012012
Reisman DS, Bastian AJ, Morton SM. 2010. Neurophysiologic and rehabilitation insights from the split-
belt and other locomotor adaptation paradigms. *Phys Ther* **90**:187–195. doi:10.2522/ptj.20090073
Reisman DS, Block HJ, Bastian AJ. 2005. Interlimb coordination during locomotion: what can be adapted
and stored? *J Neurophysiol* **94**:2403–2415. doi:10.1152/jn.00089.2005
Roemmich RT, Bastian AJ. 2015. Two ways to save a newly learned motor pattern. *J Neurophysiol*
113:3519–3530. doi:10.1152/jn.00965.2014
Roemmich RT, Long AW, Bastian AJ, Roemmich RT, Long AW, Bastian AJ. 2016. Seeing the Errors You
Feel Enhances Locomotor Performance but Not Learning. *Curr Biol* **26**:1–10.
doi:10.1016/j.cub.2016.08.012
Rossi C, Bastian AJ, Therrien AS. 2021a. Mechanisms of proprioceptive realignment in human motor
learning. *Curr Opin Physiol* **20**:186–197. doi:10.1016/j.cophys.2021.01.011
Rossi C, Chau CW, Leech KA, Statton MA, Gonzalez AJ, Bastian AJ. 2019. The capacity to learn new
motor and perceptual calibrations develops concurrently in childhood. *Sci Rep* **9**:9322.
doi:10.1038/s41598-019-45074-6
Rossi C, Roemmich RT, Schweighofer N, Bastian AJ, Leech KA. 2021b. Younger and Late Middle-Aged
Adults Exhibit Different Patterns of Cognitive-Motor Interference During Locomotor Adaptation, With
No Disruption of Savings. *Front Aging Neurosci* **13**:789. doi:10.3389/fnagi.2021.729284
Shadmehr R, Krakauer JW. 2008. A computational neuroanatomy for motor control. *Exp Brain Res*
185:359–381. doi:10.1007/S00221-008-1280-5/FIGURES/9
Smith MA, Ghazizadeh A, Shadmehr R. 2006. Interacting adaptive processes with different timescales
underlie short-term motor learning. *PLoS Biol* **4**:e179. doi:10.1371/journal.pbio.0040179
Sombric C, Gonzalez-Rubio M, Torres-Oviedo G. 2019. Split-Belt walking induces changes in active, but
not passive, perception of step length. *Sci Rep* **9**:16442. doi:10.1038/s41598-019-52860-9
Sombric CJ, Harker HM, Sparto PJ, Torres-Oviedo G. 2017. Explicit action switching interferes with the
context-specificity of motor memories in older adults. *Front Aging Neurosci* **9**:40.
doi:10.3389/fnagi.2017.00040
Song CN, Stenum J, Leech KA, Keller CK, Roemmich RT. 2020. Unilateral step training can drive faster
learning of novel gait patterns. *Sci Reports 2020 101* **10**:1–13. doi:10.1038/s41598-020-75839-3
Statton MA, Vazquez A, Morton SM, Vasudevan EVL, Bastian AJ. 2018. Making sense of cerebellar
contributions to perceptual and motor adaptation. *Cerebellum* **17**:111–121. doi:10.1007/s12311-017-
0879-0
Sultan F, Augath M, Hamodeh S, Murayama Y, Oeltermann A, Rauch A, Thier P. 2012. Unravelling
cerebellar pathways with high temporal precision targeting motor and extensive sensory and parietal
networks. *Nat Commun* **3**:1–10. doi:10.1038/ncomms1912
Synofzik M, Lindner A, Thier P. 2008. The cerebellum updates predictions about the visual consequences
of one's behavior. *Curr Biol* **18**:814–818. doi:10.1016/J.CUB.2008.04.071
Takakusaki K. 2013. Neurophysiology of gait: From the spinal cord to the frontal lobe. *Mov Disord*
28:1483–1491. doi:10.1002/MDS.25669
Tanaka H, Ishikawa T, Lee J, Kakei S. 2020. The Cerebro-Cerebellum as a Locus of Forward Model: A
Review. *Front Syst Neurosci* **14**:19. doi:10.3389/FNSYS.2020.00019/BIBTEX
Taylor JA, Ivry RB. 2011. Flexible cognitive strategies during motor learning. *PLoS Comput Biol*
7:e1001096. doi:10.1371/journal.pcbi.1001096

Taylor JA, Krakauer JW, Ivry RB. 2014. Explicit and implicit contributions to learning in a sensorimotor
adaptation task. *J Neurosci* **34**:3023–3032. doi:10.1523/JNEUROSCI.3619-13.2014
Thompson AK, Xiang YC, Wolpaw JR. 2009. Acquisition of a simple motor skill: Task-dependent
adaptation plus long-term change in the human soleus H-reflex. *J Neurosci* **29**:5784–5792.
doi:10.1523/JNEUROSCI.4326-08.2009
Todorov E. 2005. Stochastic Optimal Control and Estimation Methods Adapted to the Noise
Characteristics of the Sensorimotor System. *Neural Comput* **17**:1084–108.
Todorov E. 2004. Optimality principles in sensorimotor control. *Nat Neurosci*. doi:10.1038/nn1309
Uiga L, Capio CM, Ryu D, Young WR, Wilson MR, Wong TWL, Tse ACY, Masters RSW. 2020. The Role
of Movement-Specific Reinvestment in Visuomotor Control of Walking by Older Adults. *Journals*
Gerontol - Ser B Psychol Sci Soc Sci **75**:282–292. doi:10.1093/geronb/gby078
Van Vugt FT, Ostry DJ. 2018. The structure and acquisition of sensorimotor maps. *J Cogn Neurosci*
30:290–306. doi:10.1162/jocn_a_01204
Vasudevan EVL, Torres-Oviedo G, Morton SM, Yang JF, Bastian AJ. 2011. Younger is not always better:
development of locomotor adaptation from childhood to adulthood. *J Neurosci* **31**:3055–3065.
doi:10.1523/JNEUROSCI.5781-10.2011
Vazquez A, Statton MA, Busgang SA, Bastian AJ. 2015. Split-belt walking adaptation recalibrates
sensorimotor estimates of leg speed but not position or force. *J Neurophysiol* **114**:3255–3267.
doi:10.1152/jn.00302.2015
Vervoort D, Rob Den Otter A, Buurke TJW, Vuillerme N, Hortobágyi T, Lamoth CJC. 2019. Effects of
aging and task prioritization on split-belt gait adaptation. *Front Aging Neurosci* **11**:10.
doi:10.3389/fnagi.2019.00010
Weeks HM, Therrien AS, Bastian AJ. 2017. The cerebellum contributes to proprioception during motion. *J*
Neurophysiol **118**:693–702. doi:10.1152/jn.00417.2016
Wolpert DM, Ghahramani Z, Flanagan JR. 2001. Perspectives and problems in motor learning. *Trends*
Cogn Sci. doi:10.1016/S1364-6613(00)01773-3
Wong WL, Masters RSW, Maxwell JP, Abernethy AB. 2008. Reinvestment and falls in community-
dwelling older adults. *Neurorehabil Neural Repair* **22**:410–414. doi:10.1177/1545968307313510
Yavari F, Mahdavi S, Towhidkhah F, Ahmadi-Pajouh MA, Ekhtiari H, Darainy M. 2016. Cerebellum as a
forward but not inverse model in visuomotor adaptation task: a tDCS-based and modeling study.
Exp Brain Res **234**:997–1012. doi:10.1007/s00221-015-4523-2
1124
1125

Boxes

Box 1. Split-belt walking adaptation paradigm and motor responses.

There are three established motor measurements that are used to quantify split-belt walking
adaptation (Finley et al., 2015; Reisman et al., 2005): "Perturbation" measures the effect of the
split-belt treadmill on the stepping pattern — the total movement error we would see in the
absence of adaptation. "Δ motor output" measures the extent that individuals compensate for the
perturbation by changing their stepping pattern — how much they alter when (time) and where
(position) they step on the treadmill with each foot. "Step length asymmetry" measures the
remaining movement error — the difference between Δ motor output and perturbation (see
Methods).

Box 1-Figure 1A illustrates the standard paradigm used in studies of split-belt adaptation, with
abrupt transitions between tied-belt to split-belt phases, and Box 1-Figure 1B illustrates the
standard motor response. In the "baseline" phase, the perturbation (red), Δ motor output (blue),
and step length asymmetry (purple) are zero, reflecting equal belt speeds and symmetric walking.
In the "adaptation" phase, the belt speeds are different and the perturbation is positive. Initially,
this leads to movement errors observed as the negative step length asymmetry. The Δ motor
output gradually adapts to compensate for the perturbation, so that the step length asymmetry
returns to zero. In "post-adaptation", the belt speeds are tied and the perturbation is again zero.
Individuals exhibit initial movement errors called "aftereffects": the Δ motor output mismatches the
perturbation (it remains elevated) and step length asymmetry is positive (Leech et al., 2018a;
Rossi et al., 2021a, 2019; Statton et al., 2018).

1147

Box 1 - Figure 1. Standard paradigm and measures. (A) Treadmill belt speeds for the standard split-belt
paradigm. **(B)** Schematic time course of standard motor measures of walking adaptation: step length
asymmetry – a measure of error (solid purple), Δ motor output – a measure of compensatory spatial and
temporal asymmetries (dotted blue), and perturbation – the effect of the speed asymmetry on the walking
pattern (dashed red). In baseline, the belts are tied, and perturbation, Δ motor output, and step length
asymmetry are all ~0. In adaptation, the right leg is faster than the left such that the perturbation is positive.
The Δ motor output is still ~0 in early adaptation, causing step length asymmetry errors (negative purple
line). By late adaptation the Δ motor output is adapted to match the perturbation, and step length asymmetry
returns to ~0. Changes to Δ motor output persist in tied-belts post-adaptation, but the perturbation is ~0,
causing step length asymmetry aftereffects (positive purple line).

1158

Box 2. Relevant methodologies and results from prior work on perceptual recalibration during
locomotor learning.

Box 2-Figure 1 illustrates selected portions of the "speed match" paradigm manipulation used in
prior work and the respective step length asymmetry data (Leech et al., 2018a). The paradigm
and data shown in Box 2-Figure 1A are consistent with that explained in Box 1 (with the addition
of a brief tied-belt catch trial in adaptation). Note that participants walk with near-zero step length
asymmetry by the end of adaptation and exhibit large aftereffects post-adaptation.

The study investigated "perceptual realignment", where perception of speed becomes biased in
adaptation, partially compensating for the speed difference so that it feels smaller. They measured
perception of speed with "speed match" tasks: participants control the speed of the right belt and
try to match it to that of the left belt (as described later in our Control experiments).

Left and right panels Box 2-Figure 1B depict the tasks performed before and after adaptation –
top row shows belt speeds and bottom row shows step length asymmetry. Before adaptation,
participants can accurately match the speeds and walk symmetrically at this near-tied-belt
configuration (right and left speeds are ~equal and step length asymmetry is ~zero at the end of
the "before adapt" task). After adaptation, participants overshoot the speed of the right belt and
select a speed configuration that is biased towards that experienced in adaptation, perceiving this
as "equal speeds" (the right speed is intermediate between the adaptation right speed and the
target left speed at the end of the "after adapt" task). The key result is that participants walk with
near-zero step length asymmetry at this configuration ("after adapt", bottom row).

This suggests that participants may have learned to walk symmetrically at two distinct speed
configurations: the adaptation configuration, and the configuration that they perceive as "equal
speeds". As illustrated in a later section (Results: Experiment 1 – Motor paradigm and
hypotheses), this flexible behavior may be indicative of stimulus-response mapping mechanisms.

1183

1184

1185

Box 2 - Figure 1. Relevant results from Leech et al., 2018. (A) Treadmill belt speeds and step length asymmetry time course, similar to that described in Box 1 – Figure 1. Vertical dashed gray lines indicate iterations of the speed match task, where participants adjust the speed of the right belt with a keypad to match it to the left. **(B)** Time courses of the belt speeds (top; orange = right, black = left) and step length asymmetry (bottom) in selected iterations of the speed match task. Left, "before adapt": last baseline task. Right, "after adapt": first post-adaptation task. Dotted horizontal lines depict the right speed (orange, top) and step length asymmetry magnitude (purple, bottom) at adaptation plateau (average over the last 30 strides). **(C)** Belt speed difference (top) and step length asymmetry (bottom) magnitudes at the end of the tasks shown in B. All curves show group mean ± SE, and all data is collected in the Leech at al. study (Leech et al., 2018a).

1196

 Figure Legends

Figure 1. Experiment 1, Hypotheses and Predictions. (A) Conceptual schematic of our paradigm with the Ramp Down task: after adaptation, the right belt speed is gradually ramped down to match the left. **(B-C)** Predictions for the Ramp Down motor measures made by two competing hypotheses. **(B) Recalibration only:** recalibration can only change movement gradually. The Δ motor output (dotted blue line) changes slowly and does not track the rapidly decreasing perturbation (dashed red line), so that step length asymmetry aftereffects emerge immediately (solid purple line, magnitude is positive). **(C) Recalibration + mapping:** mapping can change movement immediately. In the first part of the task (highlighted in green), the mapping contribution to Δ motor output (dark blue shade) is scaled down immediately as the perturbation decreases. Hence, the Δ motor output (dotted blue line) changes rapidly and tracks the perturbation (dashed red line), so that there are no step length asymmetry aftereffects (solid purple line, magnitude is ~zero). In the second part of the task, the mapping contribution to Δ motor output is zero, and the recalibration contribution to Δ motor output (light blue shade) does not change significantly. Hence, the Δ motor output (dotted blue line) does not track the perturbation (red dashed line), and step length asymmetry aftereffects emerge (solid purple line, magnitude is positive). **Right column inset:** conceptual explanation of how both hypotheses may account for the speed match results from Leech at al. (Leech et al., 2018a). In the first post-adaptation speed match task, participants increase the speed of the right belt from zero to a value that is smaller than adaptation but larger than the left belt (top panel). The perturbation increases until a value that is positive but smaller than adaptation (dashed red line, middle and bottom panels). Leech et al. observed symmetric step lengths at the end of the task, indicating that the Δ motor output (dotted blue line) is smaller than it was in adaptation and matches the perturbation. The decrease in Δ motor output can be explained by the recalibration only hypothesis as forgetting/unlearning (middle panel), or by the recalibration + mapping hypothesis as flexible scaling of the mapping contribution (bottom panel).

Figure 2. Experiment 1, Step length asymmetry. (A) Top: Experimental protocol. The Ramp Down task (purple) is used to test the predictions illustrated in Fig. 1. Bottom: Step length asymmetry time course. Background shading darkness increases with belt speed difference (color bar). Phases (except Ramp tasks) are truncated to the participant with fewest strides. **(B)** Zoomed-in baseline ramp and post-adaptation Ramp Down tasks. Speed differences for which step length asymmetry is not significantly different from zero are indicated by the green shade. Inset depicts predictions made by the competing hypotheses as in Fig. 1. All curves show group mean ± SE.

Figure 3. Experiment 1, Perturbation and Δ motor output. (A) Perturbation (red) and Δ motor output (blue) data for the Ramp Down task. **(B-C)** Perturbation data (red) and model fit for the Δ motor output (blue) for recalibration only and recalibration + mapping models. All curves show group mean ± SE. Green shade corresponds to speeds with symmetric step lengths as in Fig. 2.

Figure 4. Experiment 1, Perceptual Results. (A) Top: perturbation data (red) and recalibration + mapping fit (blue); this is the same as Fig. 3C. Bottom: perceptual task button presses (green, group mean ± SE), as a function of belt speed difference. Right: measures of motor recalibration ("r") and total motor adaptation ("$u_{plateau}$"). **(B)** Perturbation compensation (normalized perceptual and motor measures of adaptation): $compensation_{\text{perceptual}}$ bounds (green), $compensation_{\text{motor total}}$ (dark blue), and $compensation_{\text{motor recalibration}}$ (light blue). **(C-D)** Individual

1245 participants' $compensation_{\text{motor recalibration}}$ versus $compensation_{\text{perceptual}}$ (first or second button
1246 press). Solid black: least squares line. Dashed gray: unity line.

1247

Figure 5. Experiment 2, Step length asymmetry. (A) Experimental protocol, equivalent to that
of Experiment 1 except for the Ramp Up & Down part shaded in teal, where the right speed was
faster than in adaptation (ramped up to 2m/s and back down to 1.5m/s). **(B)** Step length
asymmetry time course (entire group mean ± SE). Background shade represents belt speed
difference. Phases (except ramp tasks) are truncated to the participant with fewest strides. **Inset:**
Predictions for the step length asymmetry during the teal portion of the Ramp Up & Down task,
for the memory-based (top) or structure-based (bottom) mapping hypotheses. **(C)** Zoomed-in
Ramp Up & Down task (entire group mean ± SE). Step length asymmetry for strides taken at right
speeds larger than adaptation is shown in teal. **(D-E)** Separate plots of the step length asymmetry
in the Ramp Up & Down task for the subgroups of participants that walked asymmetrically (D,
"memory-based") versus symmetrically (E, "structure-based") in the teal portion of the task
(subgroups mean ± SE). **Insets:** circles represent individual participants' number of strides, in the
teal portion of the task, with step length asymmetry below their own baseline CI. Error bars depict
subgroup mean ± SE. Subgroup assignment was performed by clustering on this measure.

1262

1263 **Figure 6.** Summary of self-reported deliberate changes to the walking pattern in adaptation. Only
1264 one participant accurately described changes to the walking pattern that related to adaptation,
1265 while other responses were negative (i.e. no deliberate changes, 3 participants), irrelevant (7
1266 participants), or inaccurate (5 participants).

1267

Figure 7. Schematic model of adaptation. Body movement depends on environment
perturbations (red) and Δ motor output (blue). The Δ motor output is adjusted by recalibration
(light blue) and mapping (dark blue) mechanisms, which perform different operations and are
arranged in tandem. We propose the following architecture and flow: **(1. Recalibration)** The
recalibration mechanism produces adjustment x_r that is fixed regardless of perturbation size (light
blue box, x_r is constant for varying p). The *same* recalibration adjustment x_r serves as an input
to *both* areas responsible for conscious perception (green box) and Δ motor output (blue box). **(2.**
Perception) Conscious perception is computed by cancelling out the recalibration adjustment
from the actual sensory feedback (green box, perception of the belt speed difference perturbation
$\tilde{p}$ is the difference between the actual speed difference p and recalibration x_r). The perceived
perturbation $\tilde{p}$ serves as an input to the mapping mechanism (dark blue box). **(3. Mapping)** The
mapping mechanism produces adjustment x_m that can vary in magnitude to appropriately
account for the perceived perturbation $\tilde{p}$ (dark blue box, x_m scales with $\tilde{p}$ and matches its
magnitude). **(4. Δ Motor output)** The overall adjustment to Δ motor output is computed by adding
the mapping adjustment x_m and recalibration adjustment x_r. The corner in the Δ motor output
versus perturbation profile arises because mapping is computed based on the perceived
perturbation $\tilde{p}$ (not the actual perturbation p) and is only learnt for positive $\tilde{p}$ (the experienced
direction). When the perturbation is perceived to be opposite than adaptation, even if it is not,
mapping is zero and the Δ motor output is constant, reflecting recalibration adjustments only (blue
box, when $\tilde{p} < 0$ and $p \geq 0$ the mapping adjustment x_m is zero and $u = x_r$).

Box 1 - Figure 1. Standard paradigm and measures. (A) Treadmill belt speeds for the standard
split-belt paradigm. **(B)** Schematic time course of standard motor measures of walking adaptation:
step length asymmetry – a measure of error (solid purple), Δ motor output – a measure of
compensatory spatial and temporal asymmetries (dotted blue), and perturbation – the effect of
the speed asymmetry on the walking pattern (dashed red). In baseline, the belts are tied, and

perturbation, Δ motor output, and step length asymmetry are all ~0. In adaptation, the right leg is
faster than the left such that the perturbation is positive. The Δ motor output is still ~0 in early
adaptation, causing step length asymmetry errors (negative purple line). By late adaptation the Δ
motor output is adapted to match the perturbation, and step length asymmetry returns to ~0.
Changes to Δ motor output persist in tied-belts post-adaptation, but the perturbation is ~0, causing
step length asymmetry aftereffects (positive purple line).

1299

Box 2 - Figure 1. Relevant results from Leech et al., 2018. (A) Treadmill belt speeds and step
length asymmetry time course, similar to that described in Box 1 – Figure 1. Vertical dashed gray
lines indicate iterations of the speed match task, where participants adjust the speed of the right
belt with a keypad to match it to the left. **(B)** Time courses of the belt speeds (top; orange = right,
black = left) and step length asymmetry (bottom) in selected iterations of the speed match task.
Left, "before adapt": last baseline task. Right, "after adapt": first post-adaptation task. Dotted
horizontal lines depict the right speed (orange, top) and step length asymmetry magnitude (purple,
bottom) at adaptation plateau (average over the last 30 strides). **(C)** Belt speed difference (top)
and step length asymmetry (bottom) magnitudes at the end of the tasks shown in B. All curves
show group mean ± SE, and all data is collected in the Leech at al. study (Leech et al., 2018a).
1310
1311

A

baseline | adaptation | Ramp Down | speed match

left — | right ---

speeds

B recalibration only

recalibration

forgetting

aftereffects

forgetting in break | forgetting in task

symmetry

motor measures

stride

C recalibration + mapping

mapping

recalibration

no aftereffects | aftereffects

mapping scaled down

symmetry

motor measures

stride

— step length asym | ···· Δ motor output | --- perturbation

model fits

A Ramp Down

B recalibration only

C recalibration + mapping

motor measures

right − left speed [m/s]

right − left [m/s]

— Δmotor output
— perturbation

A

baseline | adaptation | post-adaptation
4min | 15min | 13.5min

1.5m/s
0.5m/s

10s

— — right
— left

SL asym.

0.4

-0.4

100 strides

speed match

B

before adapt

speeds (m/s)

1.5
0.5

30 sec

SL asym.

0.4

5 strides

after adapt

1.5
0.5

adapt right
right
left

30 sec

0.4

adapt plateau

5 strides

C

measures at plateau

1.5
0.5

before adapt | after adapt

0.4

-0.4