**Title:** Automatic learning mechanisms for flexible human locomotion.

**Authors:** Cristina Rossi[a,b], Kristan A. Leech[c,d], Ryan T. Roemmich[b,e], Amy J. Bastian[a,b,*]

[a] Department of Neuroscience, The Johns Hopkins University School of Medicine, Baltimore, MD, 21205,
USA; [b] Center for Movement Studies, Kennedy Krieger Institute, Baltimore, MD, 21205, USA; [c] Division of
Biokinesiology and Physical Therapy, University of Southern California, Los Angeles, CA, 90033, USA; [d]
Neuroscience Graduate Program, University of Southern California, Los Angeles, CA, 90007, USA; [e]
Department of Physical Medicine and Rehabilitation, The Johns Hopkins University School of Medicine,
Baltimore, MD, 21205, USA. *Corresponding author: Amy J. Bastian, bastian@kennedykrieger.org.

**This PDF file includes:**

## **Abstract**

Movement flexibility and automaticity are necessary to successfully navigate different environments. When encountering difficult terrains such as a muddy trail, we can change how we step almost immediately so that we can continue walking. This flexibility comes at a cost since we initially must pay deliberate attention to how we are moving. Gradually, after a few minutes on the trail, stepping becomes automatic so that we do not need to think about our movements. Canonical theory indicates that different adaptive motor learning mechanisms confer these essential properties to movement: explicit control confers rapid flexibility, while forward model recalibration confers automaticity. Here we uncover a distinct mechanism of treadmill walking adaptation – an automatic stimulus-response mapping – that confers both properties to movement. The mechanism is flexible as it learns stepping patterns that can be rapidly changed to suit a range of treadmill configurations. It is also automatic as it can operate without deliberate control or explicit awareness by the participants. Our findings reveal a tandem architecture of forward model recalibration and automatic stimulus-response mapping mechanisms for walking, reconciling different findings of motor adaptation and perceptual realignment.

## **Abbreviations**
CI, confidence interval; $CI_{LB}$, lower bound of CI; $CI_{UB}$, upper bound of CI; PSE, point of subjective equality; norm, normalized; SD, standard deviation; SE, standard error

**Main Text**

**Introduction**

Flexibility and automaticity are essential features of human movement, so much so that we rarely think about the details of how we move. While walking, we do not think about the bend of the ankle or how quickly to swing the leg forward to step. Yet, we easily adjust walking to accommodate many situations – a muddy trail, a grassy slope, or a snowy path. These abilities are often taken for granted until something goes awry – a sprained ankle quickly brings the details of movement execution to awareness, and deliberate control is often used to avoid pain or further injury. It is only then that we truly appreciate how much the sensorimotor system is doing without our conscious awareness.

We currently do not understand how different motor learning mechanisms confer both flexibility and automaticity to human movement. One well-studied process of motor learning is sensorimotor adaptation, which occurs in response to errors between the expected and actual sensory consequences of our movements. Accordingly, this motor learning mechanism helps adjust motor commands to correct for perturbations to our movements caused by altered environmental demands (Bastian, 2008; Huberdeau et al., 2015).

Adaptation is traditionally thought to rely on the cerebellum-dependent recalibration of a forward model that associates motor commands with expected sensory consequences (Ito, 1989; Wolpert et al., 2001). For example, to walk on an icy sidewalk, we may need to recalibrate our prediction of how firmly our feet will grip the ground. The process of forward model recalibration is automatic and implicit, as it has been shown to operate without intention or awareness (Huberdeau et al., 2015; Taylor et al., 2014; Tsay et al., 2024). However, forward model recalibration does not confer rapid flexibility because it cannot make immediate changes in movement – it can only adjust movement gradually trial-by-trial or step-by-step, a process that can take several minutes (Bastian, 2008; Huberdeau et al., 2015; Tsay et al., 2024). The newly acquired sensorimotor recalibration must be unlearned over time to restore a normal movement when the environment returns to its original state after adaptation (Bastian, 2008; Martin et al., 1996a). For this reason, forward model recalibration leads to lasting movement errors (called "aftereffects").

Sensorimotor adaptation can rely on more than forward model recalibration. We know that adaptation of goal-directed reaching movements can involve rapid flexible learning mechanisms

sometimes called "stimulus-response mapping mechanisms". Different types of stimulus-
response mapping mechanisms have been characterized in previous studies: First, explicit
strategies – where people deliberately change where they are aiming to reach (Taylor et al., 2014;
Taylor and Ivry, 2011). Second, memory-based caching – where people learn motor responses
in association with the respective environmental sensory stimuli, and cache them in memory for
future retrieval (like a lookup-table) (Huberdeau et al., 2019; McDougle and Taylor, 2019). Third,
structural learning – where people learn general relationships between environmental sensory
stimuli and motor responses and use them to produce novel responses (like when we learn to
map the movement of a computer mouse to that of the cursor) (Bond and Taylor, 2017; Braun et
al., 2009). Stimulus-response mapping mechanisms differ from forward model recalibration in that
they confer rapid flexibility – novel responses can be promptly abandoned or changed for different
environmental stimuli – and therefore do not lead to aftereffects (Huberdeau et al., 2015; Taylor
et al., 2014; Taylor and Ivry, 2011; Tsay et al., 2024).

It is unclear if stimulus-response mapping mechanisms are involved in adapting movement types
that are continuous like walking versus discrete movements like reaching. The goal of this study
was to understand whether walking adaptation involves any stimulus-response mapping
mechanism and, if so, how it operates and how it can be dissected from forward model
recalibration. We focused on adaptation of walking movements on a split-belt treadmill: it is well
established that when people walk with one foot faster than the other, they adapt the timing and
location of their step to restore symmetry (Box 1).

In Experiment 1A we tested the presence of stimulus-response mapping during gait adaptation
by evaluating whether people develop the ability to modify their walking pattern immediately for
different split-belt magnitudes. Preliminary evidence suggests that people may be able to switch
between at least two different walking patterns more rapidly than we would expect from forward
model recalibration alone (Box 2) (Leech et al., 2018a). Based on this evidence, we hypothesized
that walking adaptation involves stimulus-response mapping in addition to forward model
recalibration.

As our findings corroborated this hypothesis, in Experiment 1B we aimed to develop a measure
to dissect individual contributions of the two mechanisms to adaptation. We based this on the
well-known phenomenon of "perceptual realignment", where perception of the belt speed
difference diminishes over time during adaptation (Box 2) (Jensen et al., 1998; Leech et al.,

2018a; Rossi et al., 2019; Vazquez et al., 2015). Previous studies show that perceptual
realignment is only partial – the belt speeds feel similar but not completely equal at the end of
adaptation, despite complete motor adaptation (Leech et al., 2018a; Vazquez et al., 2015).
Studies also suggest that perceptual realignment may stem from forward model recalibration
processes ('t Hart and Henriques, 2016; Izawa et al., 2012; Rossi et al., 2021a; Synofzik et al.,
2008). Based on these studies, we hypothesized that the extent of perceptual realignment
corresponds to the contribution of forward model recalibration to motor adaptation, with the
remainder attributed to stimulus-response mapping.

Finally, in Experiment 2 we began to explore characteristics of stimulus-response mapping. We
asked whether the stimulus-response mapping mechanism is automatic, or instead under
deliberate or explicit control. Stimulus-response mapping mechanisms for reaching require
explicit control (Bond and Taylor, 2017; Huberdeau et al., 2019; Taylor and Ivry, 2011; Tsay et
al., 2024), but explicit control is poorly suited for automatic, continuous movements like walking
(Clark, 2015; Paul et al., 2005; Uiga et al., 2020), and can even lead to falls (Wong et al., 2008).
Unlike reaching, adaptation of walking is unaffected by explicit goals or instructions given to the
participants on where to aim their feet (Long et al., 2016; Roemmich et al., 2016). Hence, we
hypothesized that participants would not be able to describe how they changed walking, in
contrast to what has been previously reported in reaching. We also asked if participants could
use the mapping mechanism to produce novel motor outputs akin to what has been interpreted
as structural learning in reaching (Bond and Taylor, 2017; Braun et al., 2009).

**Results**

<u>Experiment 1</u>

*Motor paradigm and hypotheses*

We asked whether walking adaptation involves both forward model recalibration and stimulus-
response mapping mechanisms learned in tandem ("recalibration + mapping hypothesis"). We
contrasted this with the alternative hypothesis that walking adaptation may involve only forward
model recalibration mechanisms ("recalibration only hypothesis").

To test this, we devised the "Ramp Down" paradigm depicted in Figure 1A. After adaptation, we

gradually decreased the speed of the right belt every 3 strides (see Methods). This enabled us to

obtain reliable measurements of aftereffects across 21 predetermined speed configurations,

spanning from the full split-belt perturbation to tied-belts.

We specifically measured aftereffects in step length asymmetry, the movement error that arises

when the compensatory adjustments to step timing and position developed in adaptation (Δ motor

output) fail to match the current treadmill speed difference (perturbation):

$$\text{step length asymmetry} = \Delta \text{ motor output} - \text{perturbation} = \frac{1}{\text{stride length}} * \Delta \text{step length} \quad (1)$$

$$\text{perturbation} = \frac{\text{mean time}}{\text{stride length}} * \Delta \text{step velocity} \quad (2)$$

$$\Delta \text{ motor output} = \frac{1}{\text{stride length}} * \Delta \text{step position} - \frac{\text{mean velocity}}{\text{stride length}} * \Delta \text{step time} \quad (3)$$

where "Δ" represents differences between right and left steps and "mean" represents their

average (see Methods and Box 1).

As depicted in Figure 1B, the recalibration only hypothesis predicts that aftereffects will be present

for all speed configurations in the Ramp Down. This is because the Δ motor output (blue) can

only change gradually, through unlearning or forgetting of the forward model recalibration (SI

Appendix, Fig. S1). This change is too slow to track the perturbation (red), which is ramped down

to zero rapidly over ~80 seconds. Therefore, step length asymmetry (purple) becomes positive

immediately in the Ramp Down.

In contrast, as depicted in Figure 1C, the recalibration + mapping hypothesis predicts that

aftereffects will be present for some, but not all speed configurations in the Ramp Down.

Specifically, there would be no aftereffects (~zero step length asymmetry) in the first portion of

the Ramp Down, because the Δ motor output (blue) changes rapidly and matches the perturbation

(red). This reflects the flexibility of the stimulus-response mapping (SI Appendix, Fig. S1).

Aftereffects would emerge in the second portion of the task (positive step length asymmetry),

reflecting forward model recalibration.

We include forgetting effects for the recalibration only hypothesis to account for previous findings

from Leech et al. (Leech et al., 2018a) (Box 2). The inset of Figure 1 (right column) illustrates how

both hypotheses may account for near-zero step length asymmetry at the end of the speed match

task. The Δ motor output decreases to match the perturbation either because the mapping

mechanism flexibly scales down (recalibration + mapping hypothesis) or, alternatively, because

of forgetting/unlearning (recalibration only hypothesis).

*Motor results – step length asymmetry*

Figure 2A shows the paradigm used for Experiment 1, and the time course of step length

asymmetry (group mean ± SE). The Ramp Down task corresponds to the manipulation illustrated

in Figure 1 and is the focus of our analysis for Experiment 1. A similar ramp task was performed

prior to adaptation as a control to see how participants responded to a gradually ramped

perturbation at baseline (see Methods, "Ramp tasks"). Participants performed perceptual tests

during the ramp tasks, which will be discussed in a later section (see "Perceptual results*")*.

Patterns of step length asymmetry during other portions of the paradigm were consistent with a

large body of previous work (e.g., Leech et al., 2018a, 2018b; Reisman et al., 2005; Roemmich

et al., 2016; Rossi et al., 2019; Vazquez et al., 2015) and will not be discussed further.

Step length asymmetry during the baseline ramp and post-adaptation Ramp Down tasks are

displayed in Figure 2B. For each speed configuration in the Ramp Down task, we statistically

compared step length asymmetry to zero. We evaluated the emergence of aftereffects by

examining when step length asymmetry became significantly positive. We found that step length

asymmetry was *not* statistically different from zero during the first half of the Ramp Down task, for

right belt speeds ranging from 1m/s to 0.5m/s faster than the left speed (these speed

configurations are highlighted in green in Fig. 2B right; all $CI_{LB}$ [confidence intervals lower bounds]

≤ -0.001 and $CI_{UB}$ [upper bounds] ≥ 0.009, SI Appendix, Fig. S2 and Table S1). This result

indicates that aftereffects do not emerge immediately in the Ramp Down task, a finding at odds

with the recalibration only prediction (Fig. 2B 'predictions' inset, top). Step length asymmetry was

instead significantly positive for the second half of the task, for right belt speeds ranging from

0.45m/s to 0m/s faster than the left speed (all $CI_{LB}$ > 0.01). These results indicate that aftereffects

emerge at a mid-point during the Ramp Down task and align with behavioral predictions from the

recalibration + mapping hypothesis (Fig. 2B 'predictions' inset, bottom).

As a control, we repeated the same analysis for the baseline ramp task (Fig. 2B, left). We found

that there was only one speed configuration for which step length asymmetry was not statistically

different from zero (right speed 0.05m/s slower than left; SL asym. = 0.006 [-0.017, 0.028], mean

[CI]). For all other configurations, step length asymmetry was significantly positive ($CI_{LB} \geq 0.048$

for right speed 0.1 to 0.35m/s slower than left) or negative ($CI_{UB} \leq -0.019$ for right speed 0 to

0.35m/s faster than left). This is in stark contrast to the wide range of speed configurations with

near-zero step length asymmetry observed in the Ramp Down period. Hence, the ability to walk

symmetrically at different speed configurations is not innate but dependent on the adaptation

process. We finally tested whether the pattern of motor variability during adaptation aligns with

predictions for learning new stimulus-response maps. In contrast to recalibration, mapping

mechanisms are predicted to be highly variable and erratic during early learning, and stabilize as

learning progresses (Tsay et al., 2024). Consistent with these predictions, the step length

asymmetry residual variance (around a double exponential fit) decreased significantly between

the start and end of adaptation (residual variance at start minus end of adaptation = 0.005 [0.004,

0.007], mean [CI]; SI Appendix, Fig. S3). These control analyses corroborate the hypothesis that

the "no aftereffects" region of the Ramp Down reflects the operation of a mapping mechanism.

*Motor results – perturbation and Δ motor output*

To further test the competing hypotheses, we examined the perturbation and Δ motor output

during the Ramp Down (Figure 3A). As the belt speed difference decreased, so did the

perturbation component (red). In the first half of the task, the Δ motor output (blue) appeared to

match the perturbation – consistent with the lack of step length asymmetry aftereffects (green

shaded portion, matching Fig. 2B). Additionally, in the second half of the task, the Δ motor output

appeared larger than the perturbation – consistent with the positive step length asymmetry

aftereffects observed in Fig. 2B.

We formally contrasted predictions made by the competing hypotheses by developing
mathematical models of the Δ motor output as a function of perturbation (Fig. 3B-C). In the
simplest framework, the Δ motor output behavior for the recalibration and mapping mechanisms
can be formalized as follows:

Forward Model Recalibration:       $u(k) = r$                                    (4)

Stimulus-Response Mapping:       $u(k) = p(k)$                                  (5)

Recalibration + Mapping:       $u(k) = \begin{cases} p(k) & , & \text{if } p(k) \geq r \\ r & , & \text{otherwise} \end{cases}$       (6)

Where $u$ is the modelled Δ motor output, $k$ is the stride number, $p$ is the perturbation, and $r$ is a
free parameter representing the portion of the Δ motor output related to recalibration. We used
Equation (6) as our model for the recalibration + mapping hypothesis. This describes the scenario
where participants modulate the Δ motor output to match the perturbation $\left(u(k) = p(k)\right)$ for the
first portion of the Ramp Down task. The Δ motor output remains constant during the second
portion of the task $(u(k) = r)$ because participants are unable to reduce the Δ motor output to
values smaller than "$r$" (the amount achieved by recalibration). Parameters were estimated by
fitting the model to individual participants' Δ motor output data from the Ramp Down (see
Methods).

The fitted Δ motor output is displayed in Figure 3B (mean ± SE of fits across participants;
individual fits are shown in SI Appendix, Fig. S4). As expected, the recalibration + mapping fit
captured the matching-then-divergent behavior of Δ motor output in response to the changing
perturbation.

We used the dual state model of motor adaptation from Smith et al. (Smith et al., 2006) as our
model for the recalibration only hypothesis (see Methods). The model has four parameters and
can account for potential forgetting or unlearning of the Δ motor output that may occur during the
Ramp Down. It can also account for the possibility that adaptation involves two recalibration
mechanisms – i.e., two distinct 'fast' and 'slow' mechanisms that both learn via a process of
forward model recalibration, and both contribute to aftereffects, but that learn and forget at
different rates. We chose this model because it is a well-established model that is widely used to
capture the Δ motor output time-course for traditional motor adaptation paradigms (i.e., those

consisting of adaptation and post-adaptation phases). Yet, performance of this model for a
manipulation like the Ramp Down task performed here has not yet been tested. As such, if the
recalibration + mapping model fit the Ramp Down data better than the dual state model, this would
provide robust evidence for the presence of a stimulus-response mapping mechanism.

We show the Δ motor output fit by the recalibration only (dual state) model in Figure 3C (left panel,
group mean ± SE; individual fits are shown in SI Appendix, Fig. S5). In contrast to the recalibration
+ mapping model, the dual state model was not able to capture the matching-then-divergent
behavior of Δ motor output. The BIC statistic confirmed that the recalibration + mapping model
fitted the data significantly better than the dual state (BIC difference = 8.422 [3.386, 13.778], mean
[CI]).

We considered two additional prominent models for motor adaptation: optimal feedback control
(Izawa and Shadmehr, 2011; Shadmehr and Krakauer, 2008; Todorov, 2004) and memory of
errors (Herzfeld et al., 2014). Similar to the dual state, these models could not capture the
matching-then-divergent behavior of Δ motor output and fitted the Ramp Down data significantly
worse than the recalibration + mapping (Fig. 3C; memory of errors minus recalibration+mapping
BIC difference = 11.610 [6.068, 17.749], optimal feedback control minus recalibration+mapping
BIC difference = 19.272 [14.216, 24.563], mean [CI]). In sum, the modeling analysis of the Δ
motor output further supports the recalibration + mapping hypothesis.

*Perceptual test and results*

The second goal of Experiment 1 was to evaluate the hypothesis that "perceptual realignment" (a
phenomenon leading to altered perception following adaptation) results from the operation of the
same forward model recalibration mechanism involved in adaptation of the Δ motor output (Rossi
et al., 2021a).

Previous work shows that perception realigns following gait adaptation: after adapting to a
perturbation where the right treadmill belt is faster than the left, return to tied belts results in
perception of the opposite asymmetry (i.e., right speed feels slower than left) (Jensen et al., 1998;
Vazquez et al., 2015). We therefore expected that people would perceive the belt speeds as equal
at some point in the Ramp Down and eventually perceive the opposite asymmetry on tied belts.
We measured this by asking participants to press a keyboard button at two separate occasions

during the Ramp Down task: (1) when the belts first felt equal, and (2) when they no longer felt
equal (see Methods).

Figure 4A depicts the button presses for the Ramp Down perceptual test (top panel, group mean)
overlayed onto the Ramp Down motor data (recalibration + mapping fit). Note that the task
captures a *range* of belt speed configurations that participants perceive as "equal speeds", with
button presses (1) and (2) corresponding to the upper and lower bounds of this range. This is
expected because perception is known to be noisy and may not be sensitive enough to
discriminate between belt speed configurations that are too similar.

We quantified perceptual realignment using the established measure of *point of subjective*
*equality* (PSE), defined as the belt speed difference perceived as "equal speeds". A PSE of zero
would indicate no perceptual realignment (accurate perception) and a PSE of 1m/s (i.e., a
magnitude equivalent to the difference between the belt speeds during adaptation) would indicate
complete perceptual realignment such that the belt speeds feel equal during the adaptation
phase.

We measured belt speed difference at the time of each button press. We computed PSE as the
range of belt speed difference values between these two measurements:

$$PSE_{\text{upper bound}} = [\ right\ belt\ speed - left\ belt\ speed\ ]_{\text{time of button press 1}} \qquad (7)$$

$$PSE_{\text{lower bound}} = [\ right\ belt\ speed - left\ belt\ speed\ ]_{\text{time of button press 2}} \qquad (8)$$

We found that $PSE_{\text{upper bound}}$ was 0.64±0.03m/s and $PSE_{\text{lower bound}}$ was 0.39±0.05m/s (Figure 4A,
bottom panel, group mean ± SE). This was consistent with previous work (Leech et al.,
2018a).This perceptual realignment was not present during baseline testing, as illustrated in SI
Appendix, Fig. S6.

We aimed to evaluate the hypothesis that perceptual realignment arises from the forward model
mechanism of motor adaptation and is unaffected by the stimulus-response mapping mechanism.
This hypothesis predicts that the extent of perceptual realignment should be: 1) approximately
equal to the extent of motor adaptation achieved by recalibration, and 2) less than the total extent
of motor adaptation (which also includes mapping). We quantified 1) motor adaptation by
recalibration as the fitted parameter "$r$" from the recalibration + mapping model, and 2) total motor
adaptation as the Δ motor output at adaptation plateau (Fig. 4A, right panel, see previous section
"*Motor results – perturbation and Δ motor output*"). We expressed these motor measures and
perceptual realignment as "percent compensation for the perturbation" (i.e., normalized to the
perturbation magnitude in the respective units) so that they could be compared:

$$compensation_{\text{perceptual}} = \frac{PSE}{1\,\text{m/s}} \tag{9}$$

$$compensation_{\text{motor recalibration}} = \frac{r}{p_{\text{plateau}}} \tag{10}$$

$$compensation_{\text{motor total}} = \frac{u_{\text{plateau}}}{p_{\text{plateau}}} \tag{11}$$

Where $r$ is the fitted parameter from the recalibration + mapping model, $p_{\text{plateau}}$ is the mean
perturbation over the last 30 strides of adaptation, and $u_{\text{plateau}}$ is the mean Δ motor output over
the last 30 strides of adaptation (for Equation (9), note that 1m/s is the belt speed difference in
adaptation).

We show group level compensation measures in Fig. 4B. We found that
$compensation_{\text{motor recalibration}}$ (56±4%, group mean ± SE) fell within the $compensation_{\text{perceptual}}$
range (39±5% to 64±3%): it was significantly smaller than the upper bound (difference = -8 [-14,
-2]%) and significantly larger than the lower bound (difference = 18 [8, 28]%, mean [CI]). This
supports our first prediction that perceptual realignment is comparable to the extent of motor
adaptation achieved by recalibration. Furthermore, $compensation_{\text{perceptual}}$ was significantly
smaller than $compensation_{\text{motor total}}$ (95±2%, mean ± SE; difference = 31 [25, 38]% or 57 [48,
66]%, upper or lower perceptual bounds, mean [CI]). This supports our second prediction that
perceptual realignment is less than the total extent of motor adaptation.

We show individual $compensation_{\text{motor recalibration}}$ and $compensation_{\text{perceptual}}$ measures for each

participant in Figure 4C-D. We evaluated Pearson's correlation coefficients between these

measures to test whether there is a direct relationship between perceptual realignment and the

motor adaptation achieved by recalibration. We found that $compensation_{\text{motor recalibration}}$ was

significantly correlated with the upper bound of $compensation_{\text{perceptual}}$ – the value computed using

the "speeds feel equal" button press (r=0.64, p=0.002). This supports the hypothesized

relationship between motor recalibration and changes to leg speed perception. Instead,

$compensation_{\text{motor recalibration}}$ was not correlated with the lower bound of $compensation_{\text{perceptual}}$

(r=0.30, p=0.195) – this value is computed using the "right feels slower than left" button press,

suggesting the true PSE may lay closer to the first button press. Together, our results support the

hypothesis that perceptual realignment can be used as a proxy measure for the extent of motor

adaptation achieved by forward model recalibration.

*Modelling analysis for perceptual realignment*

We asked whether the Ramp Down results could be explained by two recently developed

frameworks designed to capture the relationship between perceptual and motor changes with

reaching adaptation: the proprioceptive re-alignment model (PReMo) (Tsay et al., 2022, 2021)

and the Perceptual Error Adaptation model (PEA) (Zhang et al., 2024). To apply these models to

walking adaptation, we translated reaching to walking adaptation variables using the conceptual

equivalence outlined by Tsay et al., which extends the model to different perturbation types in

reaching adaptation (from visual-proprioceptive to force-field perturbations – which are

mechanical like split-belt; SI Appendix, "Evaluation and development of perceptual models").

Figure 5A-B shows the fitted Δ motor output and predicted perception of speed difference for

PReMo and PEA. The models could not capture the matching-then-divergent behavior of Δ motor

output, performing significantly worse than the recalibration + mapping model (PReMo minus

recalibration+mapping    BIC    difference    =    24.591    [16.483,    32.037],    PEA    minus

recalibration+mapping BIC difference = 6.834 [1.779, 12.130], mean [CI]). Furthermore, they

could not capture the perceptual realignment and instead predicted that the right leg would feel

faster than the left throughout the entire Ramp Down.

While the mathematical formulations of the models failed to account for the observed patterns of

motor recalibration and perceptual realignment, we suspected that their underlying principles

remain valid for walking adaptation. We therefore developed a new model – the perceptuomotor

recalibration + mapping model (PM-ReMap) – that preserved the main structure of PReMo while

addressing its limitations, as identified through iterative simulations (SI Appendix, Fig. S7-8 and

"Evaluation and development of perceptual models"). The final model retains PReMo's core

principle that both motor adaptation and perceptual realignment are driven by the altered

perception of the Δ motor output, reflecting the Bayesian integration of predicted and actual Δ

motor output further shifted by recalibration (see Methods). The PM-ReMap model also maintains

the key operational properties of the mechanisms described by the recalibration + mapping model,

but includes additional variables for forgetting or unlearning of recalibration during the Ramp

Down.

As shown Figure 5C, the PM-ReMap model captured the Δ motor output in the Ramp Down with

performance comparable to that of the recalibration + mapping model (BIC difference = 2.381 [-

0.739, 5.147], mean [CI]). It also captured perceptual realignment, predicting that some

intermediate belt speed difference in the Ramp Down is perceived as "equal speeds" ($\widehat{PSE}$, Fig.

5C). We compared this $\widehat{PSE}$ estimate to the actual PSE measured by the perceptual task,

expressed as $compensation_{\text{perceptual}}$ in normalized units as before. At group level, $\widehat{PSE}$ was

comparable to the upper bound of $compensation_{\text{perceptual}}$ (difference = -7 [-15, 1]%, mean [CI]),

but significantly larger than the lower bound (difference = 19 [8, 31]%, mean [CI]). Furthermore,

we found a significant correlation between individual participants' $\widehat{PSE}$ and their upper bound of

$compensation_{\text{perceptual}}$ (r=0.63, p=0.003), but not their lower bound (r=0.30, p=0.203). Both sets

of results are consistent with those observed for the recalibration + mapping model. We also

confirmed that the significant correlation was driven by the model parameter $\eta_p$, which captures

the extent of perceptual realignment at adaptation plateau (correlation of $\eta_p$ with

$compensation_{\text{perceptual}}$ upper bound: r=0.57, p=0.009, lower bound: r=0.37, p=0.110). In sum, the

PM-ReMap model accurately captures the observed patterns of motor recalibration and

perceptual realignment, predicting that both reflect the same proportion of the perturbation size.

As such, it extends the recalibration + mapping model by incorporating the ability to account for

forgetting – typical of state space models – while still effectively capturing both recalibration and

mapping mechanisms. However, performance of the PM-ReMap model does not exceed that of

the simpler recalibration + mapping model, suggesting that forgetting and unlearning do not have

a substantial impact on the Ramp Down.

<u>Control experiments</u>

*Replication of Experiment 1 results*

We performed six control experiments and reanalyzed previously-published data (Leech et al.,
2018a) to replicate the findings of Experiment 1 across different paradigm conditions (Fig. 6). This
provided additional support for the recalibration + mapping hypothesis.

First, we checked that the motor and perceptual behaviors observed in the Ramp Down could be
replicated using a different method of assessment. A "speed match" task was given where
participants use a joystick to increase the speed of the right belt (initially stationary) until they feel
it matches the left belt (Fig. 6B, "Ascend") (Leech et al., 2018a; Rossi et al., 2019; Statton et al.,
2018; Vazquez et al., 2015). This approximates the second half of the Ramp Down (Fig. 4A, right
leg feels equal or slower than left). Consistent with this, we observed step length asymmetry
aftereffects early in the speed match task (initial SL asym. = 0.433 [0.271, 0.612], mean [CI]).
Asymmetry decreased and was eventually near-zero when the belt speeds felt equal (final SL
asym. = 0.002 [-0.040, 0.045]; Fig. 7B).

We also used a "descend" speed match task where participants *decreased* the speed of the right
belt (initially fast as in adaptation) until they felt it matched the left (Fig. 6B). This approximated
the first half of the Ramp Down. As expected, step length asymmetry was close to zero for entire
task (initial: -0.017 [-0.069, 0.039], final: 0.043 [-0.0004, 0.090], mean [CI]; Fig. 7B). The PSE for
both speed match groups was comparable to that of Experiment 1 ($compensation_{\text{perceptual}}$
difference between Ramp Down lower bound and ascend or descend Control = -3 [-16, 10]% or
4 [-12, 18]%, mean [CI]), and smaller than motor adaptation ($compensation_{\text{motor total}} -$
$compensation_{\text{perceptual}}$ = 48 [39, 57]% or 56 [42, 70]% for ascend or descend, mean [CI]; Fig. 7A).

Second, we replicated these findings in additional speed match experiments that varied in
adaptation duration (3, 15, or 30 minutes), perturbation magnitude (1m/s or 0.4m/s speed
difference) and schedule (abrupt or gradual). Across all conditions, $compensation_{\text{perceptual}}$ was
significantly smaller than $compensation_{\text{motor total}}$ (Fig. 7A; statistical results in SI Appendix, Table
S2-3). Moreover, all conditions exhibited the expected pattern of step length asymmetry in the
first post-adaptation speed match task (Fig. 7B). Specifically, step length asymmetry was: 1) ~0
at the end of the task (when belt speeds felt equal) for all conditions, 2) significantly positive at

the start of the task for Ascend conditions (Short, Medium, and Long Ascend), and 3) ~0 at the

start of task for Medium Descend and Small Abrupt Descend conditions. While step length

asymmetry was initially negative for Short Descend and Small Gradual Descend conditions, this

direction aligns with that expected from incomplete adaptation rather than aftereffects, thus

supporting the recalibration + mapping hypothesis.

In sum, these control experiments corroborate the recalibration + mapping hypothesis by

confirming that, across various conditions of perturbation duration, magnitude, and schedule, the

step length asymmetry and PSE patterns are consistent with the presence of both mechanisms.

*Relative effect of time, perturbation magnitude and schedule, and task repetition*

We examined how the relative contributions of recalibration and mapping evolve over time and

are influenced by perturbation magnitude and schedule (Fig. 8A). We compared Short, Medium,

and Long Ascend (adapting for 3, 15, or 30 minutes), and observed progression from incomplete

to complete adaptation: $compensation_{\text{motor total}}$ significantly increased with each successive

duration, reaching full adaptation only in Long Ascend (SI Appendix, Tables S4-5). Despite this

progression, the relative contributions of recalibration and mapping did not change over time (the

ratio $compensation_{\text{perceptual}}$ / $compensation_{\text{motor total}}$ did not differ between conditions; Fig. 8A left

and SI Appendix, Table S6). We next compared Medium Descend and Small Abrupt (1m/s or

0.4m/s perturbation), and found that recalibration contributed significantly more for the smaller

perturbation (larger $compensation_{\text{perceptual}}$ / $compensation_{\text{motor total}}$ in Small Abrupt than Medium

Descend, Fig. 8A middle and Table S6). Comparing Small Abrupt and Gradual (differing in

adaptation schedule), recalibration appeared larger for Small Gradual, but this effect was not

significant (Fig. 8A right and Table S6). Finally, we confirmed no differences between the "ascend"

and "descend" speed match tasks (Table S6).

We finally examined how motor aftereffects and perceptual realignment changed across the six

iterations of the speed match task post-adaptation (Fig. 8B; tasks are interleaved with varying

intervals of tied-belt walking). We quantified motor aftereffects as the step length asymmetry

expressed at the tied-belt configuration in each task (interpolated, see Methods), and perceptual

realignment as the final speed difference, normalized to the respective adaptation perturbation

like before. We focus on "Ascending" groups because no tied-belt configuration is expected in

"Descending" tasks. At the end of adaptation, motor aftereffects were comparable to perceptual

realignment in all groups (first task in Fig. 8B; SI Appendix, Table S7). However, they decayed

faster with repeated iterations of the task, becoming significantly smaller than perceptual

realignment after 1 or 2 minutes (significant difference in the second task at 1 minute for Short

and Long Ascend, and in the third task at 2 minutes for Medium Ascend, shaded in Fig. 8B; Table

S7). Motor aftereffects remained significantly smaller than perceptual realignment for the

remainder of the post-adaptation phase is Medium and Long Ascend; however, the difference

dissipated by 8 minutes for Short Ascend (Table S7).

In sum, we show that recalibration and mapping learn at similar rates during adaptation, resulting

in comparable time courses, but their relative contributions are modulated by error size. Moreover,

motor aftereffects and perceptual realignment decay differently with repeated post-adaptation

tasks, potentially suggesting a differential effect on the mechanisms.

<u>Experiment 2</u>

Experiment 1 demonstrated the presence of a stimulus-response mapping mechanism that can

produce Δ motor outputs that match a range of perturbations *smaller* than the adaptation

perturbation (in the Ramp Down task). In Experiment 2, we asked whether it can also produce Δ

motor outputs that match perturbations *larger* than the adaptation perturbation.

This sheds light on whether the stimulus-response mapping of walking adaptation operates akin

to memory-based or structure-based mechanisms observed in reaching adaptation. Failure to

account for larger perturbations would suggest that it is memory-based. That is, the Δ motor

outputs produced in adaptation may be cached in memory and later retrieved (Huberdeau et al.,

2019; McDougle and Taylor, 2019; Poggio and Bizzi, 2004; Wolpert et al., 2001). These Δ motor

outputs span a range of magnitudes that are smaller than the adaptation perturbation (see Box

1), so that they could match smaller but not larger perturbations.

Conversely, success in accounting for larger perturbations would suggest that the stimulus-

response mapping may be structure-based. That is, it may learn the general relationship between

perturbation and appropriate Δ motor output in adaptation and later use it to generate Δ motor

outputs anew (Bond and Taylor, 2017; Braun et al., 2009; McDougle and Taylor, 2019; Wolpert

et al., 2001). Hence, it would be able to generate Δ motor outputs to match either smaller or larger

perturbations. Thus, we tested a "Ramp Up & Down" condition where the speed of the right belt
was both gradually increased and then decreased after adaptation (Fig. 9A). The memory-based
hypothesis predicts that step length asymmetry will become negative for perturbations *larger* than
adaptation, while the structure-based hypothesis predicts it will remain close to zero (Fig. 9B inset,
"predictions"). The rest of the paradigm was analogous to that of Experiment 1 except there was
no perceptual assessment. Figure 9B shows step length asymmetry throughout the paradigm,
and Figure 9C shows a close-up of the Ramp Up & Down task performance (group mean ± SE).
The magenta portion of the task corresponds to the Ramp Down of Experiment 1, but the step
length asymmetry differs because of the exposure to larger speed differences in the teal portion
(a supplementary analysis confirmed that this is consistent with the recalibration + mapping
hypothesis; see SI Appendix, "Ramp Down comparison between Experiments 1 and 2", Fig. S9-
10 and Table S8).

On average, participants' step length asymmetry patterns did not remain zero for speed
differences larger than adaptation (Fig. 9C, teal). However, we observed that individual
participants exhibited markedly different patterns of step length asymmetry during this phase (SI
Appendix, Fig. S11). We quantified this observation by evaluating, for each participant, the
number of strides in this phase with step length asymmetry below their own baseline CI (SI
Appendix, Table S9). We used a density-based analysis to formally assess whether there were
separate clusters in our data (see Methods). Indeed, the algorithm detected two separate clusters
of participants: for 12 participants, between 38 to 60 strides were asymmetric (out of 60 total
strides); for the other 8 participants, only 3 to 21 strides were asymmetric (Fig. 9D-E insets, and
SI Appendix, Fig. S12; difference in strides between subgroups = 36.083 [29.917, 42.250], mean
[CI]). A silhouette analysis confirmed strong evidence for these clusters: the average silhouette
score was 0.90, with 19 of 20 participants scoring above 0.7 – considered strong evidence – and
one scoring between 0.5 and 0.7 – considered reasonable evidence (Dalmaijer et al., 2022;
Kaufman and Rousseeuw, 1990; Rousseeuw, 1987). As a control, we performed the same
clustering analysis for Experiment 1, and did not find separate clusters for any of our measures
of interest (SI Appendix, "Clustering analysis", Fig. S12-13, and Table S10-11). This result
indicates that 12 of 20 participants could not account for belt speed differences larger than that of
adaptation, suggesting that they used a memory-based mapping mechanism (Fig. 9D). In

contrast, 8 of 20 participants could account for these speeds, suggesting that they engaged a
structure-based mapping mechanism (Fig. 9E).

We next aimed to exclude the possibility that participants in the structure-based subgroup may
simply be faster at adapting to new perturbations than those in the memory-based subgroup and
may be adapting to the new perturbations of the Ramp Up & Down rather than generating Δ motor
output using a previously learnt structure. To this end, we evaluated learning rates during
adaptation. We found that participants in the two subgroups adapted at similar rates (strides to
plateau difference, structure – memory: 135.875 [-53.208, 329.708], mean [CI]; SI Appendix, Fig.
S14-15). Furthermore, the pattern of step length asymmetry variability was similar between the
subgroups (structure – memory difference in residual variance relative to double exponential
during initial adaptation = -0.0052 [-0.0161, 0.0044], adaptation plateau = -0.0007 [-0.0021,
0.0003], difference in variance decay = -0.0045 [-0.0155, 0.0052], mean [CI]; SI Appendix, Fig.
S16). This confirms that the distinct performance clusters in the Ramp Up & Down task are not
driven by natural variations in learning ability, such as differences in learning speed or variability.
Rather, these findings indicate that the subgroups employ different types of mapping
mechanisms, which perform similarly during initial learning but differ fundamentally in how they
encode, retrieve, and generalize relationships between perturbations and Δ motor outputs.

We considered that the mapping adjustments described here may or may not be *deliberate* (i.e.,
participants are trying to correct for the perturbation) or done with an *explicit strategy* (i.e.,
participants can accurately report a relevant strategy that would counter the perturbation) (Long
et al., 2016; Roemmich et al., 2016). A recent framework for motor learning by Tsay et al. defines
explicit strategies as motor plans that are both intentional and reportable (Tsay et al., 2024).
Within this framework, Tsay et al. clarify that "intentional" means participants deliberately perform
the motor plan, while "reportable" means they are able to clearly articulate it. An example of an
explicit strategy in visuomotor reaching adaptation is when participants report that they aimed to
offset a visual rotation (Bond and Taylor, 2017; Taylor et al., 2014); conceivable examples in split-
belt treadmill adaptation may be reports of "taking steps of similar length", "stepping further ahead
with the right foot" or "standing on the left foot for longer". We tested whether participants could
explicitly report changes to the gait pattern that specifically correct for the split-belt perturbation.

At the end of the experiment, participants were asked to report (in writing) if/how they had changed
the way they walked during adaptation (note that this was a later addition to the protocol and was

only collected in 16 of the 20 participants). We assessed reports by categorizing them in three
steps: 1) did the report mention *any* deliberate changes? 2) were the changes *relevant* to
adaptation? (i.e., did the report mention any gait metric contributing to the overall Δ motor output
or step length asymmetry adaptation in any amount?) 3) was the response *accurate?,* as
participants often reported strategies that they did not actually execute.

We summarize results from the questionnaire in Figure 10 (original responses are reported in SI
Appendix, Table S12). We found that, while 13 participants reported *deliberate* changes, only 6
people mentioned *relevant* aspects of the walking pattern (in particular, participants mentioned
"limping" or temporal coordination). Furthermore, only one of these participants reported an
*accurate* gait parameter ("I tried to spend as much time leaning on my left leg as possible"), while
the remainder of the relevant responses were inaccurate (e.g., "matched duration of standing on
each foot") or vague (e.g., "I adjusted as if I was limping"). The aspect of gait most frequently
reported across participants was stability (also reported as balance, not falling, or controlling
sway). This suggests that, while participants may deliberately adjust their body to feel more stable,
they do not seem to explicitly strategize how to offset the perturbation. Thus, mapping tends to
adjust aspects of the walking pattern that participants are not explicitly aware of controlling,
suggesting that this differs from explicit strategies often observed in reaching.

In sum, in Experiment 2 we found that 1) participants are largely unable to explicitly describe
adaptations to their walking pattern, and 2) some but not all participants can extrapolate their
walking pattern to account for larger perturbations. This sheds light on how the stimulus-response
mapping mechanism of walking adaptation may align with established mechanisms: it may differ
from explicit strategies and resemble memory-based caching in some people and structural
learning in others.

**Discussion**

In this study, we showed that locomotor adaptation involves two learning mechanisms: forward model recalibration and a stimulus-response mapping mechanism. The recalibration mechanism changes movement gradually and relates to the perceptual changes observed in locomotor adaptation. The mapping mechanism is flexible and, once learned, can change movement immediately to account for a range of belt speed configurations. Our data suggest that this mapping operates independently of explicit strategies, and that it can be memory-based or structure-based.

*Forward model recalibration of movement and perception*

In line with various adaptation studies, we showed that people recalibrated their perception in a way that reduced how perturbed they felt (Haith et al., 2008; Harris, 1963; Jensen et al., 1998; Moidell and Bedell, 1988; Sombric et al., 2019) – i.e., their perception of the leg speed difference diminishes (Jensen et al., 1998; Leech et al., 2018a; Rossi et al., 2019; Statton et al., 2018; Vazquez et al., 2015). This perceptual change aligned with the motor change achieved through forward model recalibration. Specifically, we found a significant correlation between motor aftereffects and the speed difference participants perceived as equal. Motor aftereffects did not correlate with the speed difference later reported as no longer feeling equal, likely because participants are biased to delay responding until they are sufficiently confident, reflecting individual factors like perceptual noise (Camacho et al., 2015; Gescheider, 1997). Although previous research suggested that perceptual changes may be mediated by recalibration processes, the specific link to motor adaptation remained unclear ('t Hart and Henriques, 2016; Izawa et al., 2012; Sombric et al., 2019; Synofzik et al., 2008; Yavari et al., 2016; for a review, see Rossi et al., 2021a). Our findings demonstrate a direct relationship between perceptual and motor recalibrations, offering a novel approach to dissect forward model recalibration from stimulus-response mapping contributions to adaptation.

Based on our findings, we suggest that forward model recalibration counters the perturbation in the motor and perceptual domains simultaneously – it adapts the walking pattern while realigning perception of leg speed. In support of this idea, perceptual realignment resembles a well-studied "sensory cancellation" phenomenon – where forward model predictions are used to filter redundant sensory information (Anderson et al., 2012; Blakemore et al., 1998). Forward model

predictions may also be integrated with and sharpen proprioceptive estimates (Bhanpuri et al.,
2013; Weeks et al., 2017), and may mediate a more complex integration of predicted and actual
sensory signals contributing to perceptual changes. Consistent with the operation of a forward
model thought to predict sensory consequences from efferent copies of motor commands (Haruno
et al., 2001; Ito, 1989; Wolpert et al., 2001), perceptual changes are expressed during active and
not passive movements (Sombric et al., 2019; Vazquez et al., 2015).

*Stimulus-response mapping is flexible but requires learning*

To the best of our knowledge, this is the first study to show that a stimulus-response mapping
mechanism plays a role in walking adaptation. We isolated this mechanism by designing a novel
"Ramp Down" task, in which the belt speed difference perturbation was ramped down to zero
gradually but rapidly after adaptation (averaging 1 minute and 20 seconds). This approach
contrasted prior work that employed either abrupt or slow (10-minute) transitions to tied belts post-
adaptation (Leech et al., 2018b; Reisman et al., 2005; Roemmich and Bastian, 2015). By
implementing this rapid ramp-down, we were able to dissect the operation of a flexible mapping
mechanism that complements recalibration, producing walking patterns suited to the changing
treadmill configuration.

The mapping mechanism observed in our study aligns with the corrective responses described
by Iturralde and Torres-Oviedo, which operate relative to a recalibrated "new normal" rather than
relying solely on environmental cues (Iturralde and Torres-Oviedo, 2019). Accordingly, our
findings suggest a tandem architecture: forward model recalibration adjusts the nervous system's
"normal state," while stimulus-response mapping computes motor adjustments relative to this
"new normal." This architecture explains the sharp transition from flexible to rigid motor
adjustments observed in our Ramp Down task. The transition occurs at the configuration
perceived as "equal speeds" (~0.5 m/s speed difference) because this corresponds to the
recalibrated "new normal".

In the first half of the Ramp Down, participants adequately modulated their walking pattern to
accommodate the gradually diminishing perturbation, achieving symmetric step lengths. Due to
the recalibrated "new normal", perturbations within this range are perceived as congruent with the
direction of adaptation but reduced in magnitude. This allows the mapping mechanism to flexibly
modulate the walking pattern by using motor adjustments previously learned during adaptation.

Importantly, the rapid duration of the Ramp Down task rules out the possibility that the observed
modulation may instead reflect washout, as confirmed by the fact the aftereffects measured post-
Ramp-Down were comparable to previous work (Kambic et al., 2023; Reisman et al., 2005).

In the second half of the Ramp Down, aftereffects emerged as participants failed to accommodate
perturbations smaller than the recalibrated "new normal". These perturbations were perceived as
opposite to the adaptation perturbation and, therefore, novel. Accordingly, the mapping
mechanism responded as it would to a newly introduced perturbation, rather than leveraging
previously learned adjustments (Iturralde and Torres-Oviedo, 2019). Due to the rapid nature of
the Ramp Down, the mapping mechanism lacked sufficient time to learn the novel motor
adjustments required for these perturbations – a process that typically takes several minutes, as
shown by our baseline ramp tasks and control experiments. As mapping-related learning was
negligible, the rigid recalibration adjustments dominated during this phase. Consequently, the
walking pattern did not change to accommodate the gradually diminishing perturbation, leading
to the emergence of aftereffects.

*Mapping operates independently of explicit control*

Our results suggest that stimulus-response mapping may operate independently of explicit control
in walking adaptation. In previous work where an explicit goal is given, adaptation mechanisms
deployed in addition to forward model recalibration are consistently found to operate under
deliberate control (Codol et al., 2018; Taylor et al., 2014; Taylor and Ivry, 2011) – including
memory-based caching (Huberdeau et al., 2019; McDougle and Taylor, 2019) and structural
learning mechanisms (Bond and Taylor, 2017; McDougle and Taylor, 2019). While the mapping
described here shares some characteristics with explicit mechanisms, such as flexibility and
modulation by error size (Kagerer et al., 1997; Modchalingam et al., 2019; Neville and Cressman,
2018; Saijo and Gomi, 2010), it diverges in critical ways. Unlike explicit strategies, which are
rapidly acquired and diminish over time, this mapping mechanism exhibits prolonged learning
beyond 15 minutes, with a rate comparable to recalibration (Huberdeau et al., 2015; McDougle et
al., 2016, 2015; Taylor et al., 2014; Taylor and Ivry, 2011).

In line with these distinctions, mounting evidence indicates that explicit strategies do not play a
strong role in split-belt walking adaptation where an explicit movement goal is not provided to the
participants; for example, people do not adapt faster even after watching someone else adapt

(Song et al., 2020). Strategic adjustments to the walking pattern can be temporarily elicited by

providing additional visual feedback of the legs and an explicit goal of how to step. Yet, these

adjustments disappear immediately upon removal of the visual feedback (Roemmich et al., 2016),

and have no effect on the underlying adaptive learning process (Long et al., 2016; Malone and

Bastian, 2010; Roemmich et al., 2016). Furthermore, performing a secondary cognitive task

during walking adaptation does not affect the amount of motor learning (Hinton et al., 2020;

Malone and Bastian, 2010; Rossi et al., 2021b; Vervoort et al., 2019).

Here, we show that explicit strategies are not systematically used to adapt step length asymmetry

and Δ motor output: the participants in our study either did not know what they did, reported

changes that did not actually occur or would not lead symmetry. Only one person reported

"leaning" on the left (slow) leg for as much time as possible, which is a relevant but incomplete

description for how to walk with symmetry. Four reports mentioned pressure or weight, which may

indirectly influence symmetry (Hirata et al., 2019; Lauzière et al., 2014), but they were vague and

conflicting (e.g., "making heavy steps on the right foot" or "put more weight on my left foot"). All

other responses were null, explicitly wrong or irrelevant, or overly generic, like wanting to "stay

upright" and "not fall down". We acknowledge that our testing methodology has limitations. First,

it may introduce biases related to memory recall or framing of the questionnaire. Second, while it

focuses on participants' intentional use of explicit strategies to control walking, it does not rule out

the possibility of passive awareness of motor adjustments or treadmill configurations. Despite

these limitations, the motor adjustments reported by participants consistently fail to meet the

criteria for explicit strategies as outlined by Tsay et al.: reportability and intentionality (Tsay et al.,

2024). Together with existing literature, this supports the interpretation that stimulus-response

mapping operates automatically.

In sum, the mapping mechanism combines the advantages of automaticity and flexibility

(Huberdeau et al., 2015; Taylor et al., 2014; Taylor and Ivry, 2011). This is ecologically important

for both movement accuracy (Uiga et al., 2020; Wong et al., 2008) and for walking safely in real-

world situations, where we walk while talking or doing other tasks, and terrains are uneven (Clark,

2015; Paul et al., 2005; Wong et al., 2008).

*Mapping operates as memory-based in some people, structure-based in others*

Results from Experiment 2 highlight individual differences in the learning mechanisms underlying
generalization to unexperienced belt speed differences. The generalization to novel perturbation
sizes observed here is in line with previous suggestions of "meta-learning" in the savings of
walking adaptation (i.e., faster relearning when exposed to a different perturbation) (Leech et al.,
2018b; Malone et al., 2011). Generalization to *larger* perturbations after reaching adaptation was
shown to be incomplete (Abeele and Bock, 2001; Lazar and Van Laer, 1968), and it was unclear
whether the movement patterns had been *extrapolated* beyond what had been experienced, or
rather was just the same as in adaptation. Surprisingly, we found 40%-60% divide in our
participants regarding the capacity to extrapolate walking patterns to account for larger
perturbations.

We suggest that structural learning may underlie the ability to extrapolate walking patterns and
walk symmetrically for belt speed differences larger than adaptation, as seen in 8 of 20
participants. Indeed, generalization in reaching adaptation may rely on a process of structural
learning (Bond and Taylor, 2017; Braun et al., 2009) (note, however, that this was tied to explicit
aiming; Bond and Taylor, 2017). Our participants may have learned the scaling relationship
between belt speed perturbation and the walking pattern needed to walk symmetrically and use
this scaling to produce new walking patterns matching larger perturbations.

In contrast, memory-based theories of mapping (Dassonville et al., 2001; Poggio and Bizzi, 2004;
Van Vugt and Ostry, 2018; Wolpert et al., 2001) may explain why 12 participants walked
asymmetrically for perturbations larger than adaptation, despite generalizing to smaller
perturbations. Consistent with error-correction mappings, the walking patterns may be stored in
memory during adaptation in association with the amount of perturbation they correct, and then
retrieved during the ramp tasks (Van Vugt and Ostry, 2018). As people transition through a range
of walking patterns in adaptation (see gradual adaptation of Δ motor output in Box 1 and Fig. 1),
they may store these intermediate walking patterns and later use them to generalize to smaller
perturbations. Alternatively, people may store a limited number of walking patterns in memory,
such as those for baseline and adaptation plateau, and produce intermediate patterns by
interpolating between these memories (Poggio and Bizzi, 2004). While interpolation has
sometimes been used as a marker of structural learning (Braun et al., 2009; Haruno et al., 2001),
we argue that it fundamentally relies on stored memories. Moreover, distinguishing interpolation

from intermediate memory storage during adaptation is challenging, so we adopted extrapolation
as a more robust approach.

Both memory- and structure-based operations of mapping align with Tsay et al.'s framework for
motor learning: first, action–outcome relationships are learned through exploration; second, motor
control policies are refined to optimize rewards or costs, such as reducing error; and finally,
learned mappings or policies are retrieved based on contextual cues (Tsay et al., 2024).
Consistent with the proposed stages of exploration followed by refinement, we found that motor
behavior during adaptation was initially erratic but became less variable at later stages of learning.
Similarly, consistent with the retrieval stage, the generalization observed in the ramp tasks
indicates that learned motor outputs are flexibly retrieved based on belt speed cues.

*Mapping may underly savings upon re-exposure to the same or different perturbation*

Our control experiments demonstrate that repeated exposure to perturbations opposite to
adaptation – induced by the ascending speed match tasks post-adaptation – reduces motor and
perceptual aftereffects at a different rate. Specifically, the magnitudes of motor and perceptual
aftereffects were comparable at the end of adaptation, but diverged within 2 minutes of washout
as motor aftereffects became smaller than perceptual aftereffects. This pattern aligns with that
from Leech et al. showing that savings reduces motor aftereffects to a greater extent than
perceptual aftereffects (Leech et al., 2018a). Notably, the study involved repeated exposure to
both the adaptation perturbation and the opposite perturbation through ascending speed match
tasks structured like our paradigm. We suggest that savings – traditionally defined as the smaller
initial errors, faster relearning, and smaller aftereffects when re-adapting to the same perturbation
– also occurs for the opposite perturbation (Day et al., 2018; Krakauer and Shadmehr, 2006;
Leech et al., 2018a; Malone et al., 2011; Martin et al., 1996a; Reisman et al., 2013; Roemmich
and Bastian, 2015; Rossi et al., 2021b; Shadmehr and Brashers-Krug, 1997). This may explain
why studies have found savings after exposure to perturbations of different magnitudes (Bond
and Taylor, 2017; Leech et al., 2018b).

Specifically, we suggest that savings operates via the mapping mechanism. Mapping is ideal for
savings because, in contrast to recalibration, it can take a value of zero during post-adaptation
without having to unlearn. This means that, upon readaptation, the mapping adjustment may be
able to immediately "jump" to a positive value, rather than adapting de novo from zero like

recalibration. This may give mapping a competitive advantage such that, with repeated exposure
to split-belt walking, it may contribute relatively more than recalibration to the overall adaptation.
Indeed, savings has opposite effects on recalibration and mapping-like explicit mechanisms –
reducing the former and increasing the latter contribution – at least in reaching adaptation
(Avraham et al., 2021). In walking adaptation, Roemmich and Bastian show that savings is larger
when participants recall experiencing a larger perturbation (Roemmich and Bastian, 2015). While
their test differs from the perceptual tasks in our study, it is plausible that participants with smaller
perceptual realignment may have perceived and then recalled the perturbation more accurately,
relating the magnitude of savings to the extent learnt by mapping.

The different decay of motor and perceptual aftereffects observed by us and Leech et al. indicates
that the contribution of mapping to reducing aftereffects through savings is twofold (Leech et al.,
2018a). First, upon repeated exposure to the adaptation perturbation, mapping accounts for a
larger proportion of the learning, thereby reducing recalibration and hence both motor and
perceptual aftereffects. Second, upon repeated transitions from the adaptation perturbation to
tied-belts, mapping learns motor adjustments in the direction opposite to those induced by
adaptation, using these adjustments to actively counteract recalibration. As adjustments by
mapping do not change perception, this process only reduces motor aftereffects, explaining why
they are smaller than perceptual aftereffects. Two findings directly support the interpretation that
the return to tied belts post-adaptation induces opposite learning in mapping. First, abrupt
transitions from split belts to tied belts after adaptation are treated as perturbations in the opposite
direction in terms of muscle activity (Iturralde and Torres-Oviedo, 2019). Second, previous work
demonstrated savings following adaptation to the opposite split-belt perturbation (Malone et al.,
2011). In traditional paradigms with abrupt removal of the perturbation post-adaptation, mapping
may mask the true magnitude of motor aftereffects, leading to an apparent lack of relationship
with perceptual aftereffects (Rossi et al., 2019; Statton et al., 2018).

*Conceptual model*

We propose a comprehensive conceptual model of walking adaptation that captures key
behavioral properties uncovered in our study. We build upon standard models (Shadmehr and
Krakauer, 2008; Smith et al., 2006) and introduce a mechanism and architecture for adaptive
learning that accounts for the flexibility of the walking pattern and the relationship between motor
and perceptual changes. This conceptual model is illustrated in Figure 11.

The perturbation $p$ represents the effect of the belt speed difference on movement or perception.
Note that motor and perceptual quantities are represented in the same relative units (relative to
the perturbation).

The light blue "recalibration" box represents the operation of the forward model recalibration
mechanism: this mechanism is not flexible, so that it produces the same output $x_r$ for any
perturbation $p$. The recalibration $x_r$ may project to both sensory integration and motor control
areas (green and blue boxes), where it may be used to recalibrate perception and movement.

The green "perception" box represents the process of perceptual realignment. It receives sensory
information regarding the perturbation, $p$, and the forward model recalibration output, $x_r$. It
cancels out the portion of the perturbation predicted by the forward model, so that it is output is
the perceived perturbation $\tilde{p} = p - x_r$. The perceived perturbation $\tilde{p}$ may serves both as signal
for conscious speed perception (that captured by perceptual tests), and as an input to the
stimulus-response mapping mechanism (dark blue box).

The dark blue "mapping" box represents the operation of the stimulus-response mapping
mechanism. It receives the perceived perturbation signal $\tilde{p}$ and transforms into a motor
adjustment appropriate to counter the perceived perturbation, $x_m = \tilde{p}$.

The blue "Δ motor output" box represent the computation of the Δ motor output. It receives the
recalibration $x_r$ and mapping $x_m$ adjustments as inputs, and sums them to compute the Δ motor
output $u = x_m + x_r$.

We propose the following series of computations. First, the recalibration is used to compute the
perceived perturbation (green box). Second, the perceived perturbation is relayed to the mapping
mechanism and used to compute the mapping-related motor adjustment (dark blue box). Third,
recalibration-related and mapping-related motor adjustments are summed to produce the Δ motor
output (blue box).

In conclusion, our model proposes three key features of the mapping mechanism that distinguish
it from standard adaptation models. First, the mapping produces near-immediate changes to
motor output in response to varying perturbations – contrasting the gradual changes proposed by
state-space models (Herzfeld et al., 2014; Roemmich et al., 2016; Smith et al., 2006). Second, it
accesses the perceived perturbation signal which integrates external sensory information with

internal forward model state – contrasting the unimodal input proposed by optimal control
(Shadmehr and Krakauer, 2008; Todorov, 2005). Third, the mapping operates automatically and
is learnt with adaptation – contrasting the readily implementable explicit strategies modeled for
reaching (Taylor and Ivry, 2011). These features enable automatic motor adjustments that
complement recalibration to match varying perturbations, allowing our model to account for the
observed symmetric walking across different treadmill speed configurations.

*Implications for models of adaptation*

We found that prominent computational models for motor adaptation could not account for the
flexible properties of the stimulus-response mapping mechanism. State-space models, such as
the dual state or memory of error models, could only capture recalibration-like mechanisms
because their states store a single motor adjustment at any given time and change only gradually
(Coltman et al., 2019; Diedrichsen et al., 2010; Herzfeld et al., 2014; Inoue et al., 2015; Lee and
Schweighofer, 2009; Roemmich et al., 2016; Smith et al., 2006; Tanaka et al., 2012). Despite
promising concepts involving external sensory stimuli, current implementations of optimal control
models for adaptation cannot capture mapping because they operate like state-space models,
computing motor output linearly from a state that holds a single value and is updated gradually
(Izawa and Shadmehr, 2011; Shadmehr and Krakauer, 2008; Todorov, 2005, 2004). Similarly,
despite their potential to account for parallel changes in movement and perception, models like
Proprioceptive Re-alignment Model (PReMo) or Perceptual Error Adaptation also operate akin to
state-space models and could not capture mapping (Tsay et al., 2022; Zhang et al., 2024).

We developed two models that account for the post-learning properties of mapping mechanism
in the Ramp Down task, with or without unlearning of the recalibration mechanism. Future work
is needed to expand these models to also account for the learning process in adaptation. Despite
not being as thoroughly formalized as serial or parallel models (Lee and Schweighofer, 2009),
tandem models integrate the features necessary to effectively account for the properties of
mapping (Honda et al., 2018). Like parallel models, tandem models allow both mechanisms to
access external error signals, which is necessary for mapping to respond quickly to changes in
the perturbation. This cannot occur in serial models, where mapping only receives input from the
recalibration mechanism and has no access to external signals. Instead, mapping would only
change as a response to changes in recalibration, limiting its rate of change to be equal to or
slower than that of recalibration. Like serial models, tandem models use the output of one

mechanism (recalibration) as input to the other (mapping), which is necessary for mapping to
compute complementary adjustments that achieve symmetric walking during the initial portion of
the Ramp Down. This cannot occur in parallel models, where mapping only receives external
inputs and has no access to the state of recalibration. Instead, mapping would need to operate
solely relative to its own portion of the learning, scaling linearly from its magnitude at adaptation
plateau to zero throughout the Ramp Down. Since the magnitude of mapping is smaller than the
perturbation at the adaptation plateau, mapping has less to adjust over the Ramp Down.
Consequently, it would decay slower than the perturbation and aftereffects would emerge
immediately.

Existing models offer key insight into further development of a comprehensive framework for
mapping. Taylor and Ivry (Taylor and Ivry, 2011) was able to capture flexible adjustments by
strategic aiming and incorporate it into state-space models – though assuming these adjustments
are immediately available upon exposure to the adaptation perturbation. Modifications could be
explored to capture the adaptation learning process of mechanisms that are not immediate, like
the mapping mechanism for walking adaptation and uninstructed aiming strategies (Bond and
Taylor, 2015; McDougle et al., 2015; Taylor et al., 2014). Additionally, optimal control frameworks
are promising for modeling mapping mechanisms across paradigms because they can account
for how paradigm-specific stimuli – implicit and explicit rewards and costs – influence movement
(Todorov, 2004; Todorov and Jordan, 2002). They may also be instrumental to account for the
potentially multiple sources of errors contributing to adaptation – which may include a combination
of motor (Haith and Krakauer, 2013; Shadmehr and Krakauer, 2008; Smith et al., 2006),
perceptual (Tsay et al., 2022; Zhang et al., 2024), sensory prediction, (Haith and Krakauer, 2013;
Lee et al., 2018; Shadmehr and Krakauer, 2008; Tseng et al., 2007), and energy cost signals
(Finley et al., 2013; Sánchez et al., 2019, 2017). While we use kinematic measures for our
modelling analysis to align with established procedures (McDougle et al., 2015; Roemmich et al.,
2016; Smith et al., 2006), recalibration can at least partially proceed when kinematic error is
clamped (Gonzalez-Rubio et al., 2019; Long et al., 2016), suggesting that other sources of error
may also be at play. Building on preliminary frameworks that integrate sensory predictions and
external sensory stimuli for motor control (Cheng and Sabes, 2006; Honda et al., 2018), future
models may effectively account for mapping across paradigms.

*Neural substrates*

Operation of the recalibration mechanism may rely on the cerebellum and its protections to
sensorimotor cortices. It is well known that the cerebellum houses forward models (Tanaka et al.,
2020; Wolpert et al., 2001) and is involved in motor adaptation (Bastian, 2011; Martin et al., 1996b;
Morton and Bastian, 2006) and perceptual realignment (Izawa et al., 2012; Statton et al., 2018;
Synofzik et al., 2008; Yavari et al., 2016). Moreover, its neural architecture and functional
organization are highly optimized for integrating multiple input signals into a unified learning
process: First, the error input from climbing fibers to individual Purkinje cells can itself be
multisensory (Diedrichsen et al., 2019; Ju et al., 2019; Tanaka et al., 2020). Second, processing
within the cerebellum can integrate multisensory signals to produce a coordinated sensory
cancellation response, either directly within individual Purkinje cells (Lai et al., 2021) or through
their convergence in the nuclei (Ito, 1984; Tanaka et al., 2020). Third, cerebellar outputs carrying
distinct signals, such as different movement parameters, can be integrated in the premotor cortex
to adapt movement in a coordinated manner (Herzfeld et al., 2018). These characteristics support
the suggestion that adaptation is driven by multiple sources of error. Cerebellar connectivity also
supports the opposite process, whereas the same forward model recalibration in the cerebellum
may have diverse downstream effects depending on the specific targets of its output projections
(Welniarz et al., 2021). Cerebellar circuits are anatomically homogeneous and may perform
analogous computations, supporting the idea that their differing functional roles depend largely
on their inputs and downstream output structures (Ito, 1984; Schmahmann, 1996; Tanaka et al.,
2020).

We hypothesize that perceptual and recalibration-related motor changes with adaptation reflect
recalibration of the same forward models in the cerebellum, processed downstream by motor and
sensory cortices. Cerebellar output from single regions of the cerebellar cortex (Pisano et al.,
2021), and possibly single output cells from the cerebellar nuclei (Judd et al., 2021; Sultan et al.,
2012), projects to both sensory and motor cortical areas, modulating their activity in a coordinated
manner (Lindeman et al., 2020; Popa et al., 2013). We suggest that this connection mediates
sensory cancellation during active movement, as this cancellation is observed in both sensory
and motor cortices (Seki and Fetz, 2012) and relies on cerebellar predictions (Blakemore et al.,
1998; Brooks and Cullen, 2019; Rondi-Reig et al., 2014). Perceptual realignment may operate by
filtering out self-generated sensory stimuli from the adapted movement (Blakemore et al., 1998;
Brooks and Cullen, 2019; Rondi-Reig et al., 2014), or the predictable environmental stimuli from

the treadmill speeds (Anderson et al., 2012; Rondi-Reig et al., 2014). This may explain the
increased activity in the parietal lobe, alongside cerebellum and frontal lobe, with split-belt walking
adaptation (Hinton et al., 2019).

Operation of the mapping mechanism may rely on a similar network. Recalibration-adjusted
perceptual information on the treadmill configuration may be relayed via sensory cortical areas
and cerebellum to motor cortical areas, where it may be mapped to motor adjustments. This is
supported by the interconnectivity between these substrates (Judd et al., 2021; Kandel et al.,
2013; Sultan et al., 2012), and by the cortical contribution to the flexible control of walking (Drew
and Marigold, 2015; Reisman et al., 2010) beyond voluntary control (Delval et al., 2020; Petersen
et al., 2012). Mapping may also involve spinal control – which is rapid and automatic (Takakusaki,
2013) and can play a role in split-belt walking (Ogawa et al., 2014; Vasudevan et al., 2011) – but
likely through connection with supraspinal structures due to the extensive training needed for
plasticity within the spinal cord (Iturralde and Torres-Oviedo, 2019; Thompson et al., 2009). In
contrast, we hypothesize that mapping does not involve the prefrontal cortex, which is instead
associated with explicit control (Goldman-Rakic, 1987; Miller and Cohen, 2001). Indeed, the
prefrontal cortex becomes less active with split-belt adaptation, and greater deactivation is
associated with more symmetric strides (Hinton et al., 2019).

Motor adjustments resulting from recalibration and mapping may be integrated either cortically or
subcortically. Cortical integration is supported by adaptation studies in primates, demonstrating
cortical correlates consistent with the combined operation of recalibration-like and mapping-like
mechanisms (Chase et al., 2012; Sun et al., 2022). The recalibration-like mechanism modulates
cortical motor cells uniformly via a shared upstream input, leading to fixed motor adjustments
across environmental conditions. The mapping-like mechanism changes the weight or tuning
direction of individual cortical cells, leading to flexible motor adjustments that differ across
environmental conditions. Nevertheless, studies in humans show that cerebral damage does not
block walking adaptation (Choi et al., 2009; Reisman et al., 2007), potentially supporting the
alternative hypothesis that recalibration and mapping signals may be combined subcortically.

*Limitations*

We aim to propose a framework for motor adaptation that is both parsimonious and based on a
simple model. However, this simplicity inevitably comes with certain limitations. First, our model
is descriptive and focused on capturing the novel features of the data revealed by the Ramp Down
task, rather than the entire motor adaptation time series. Second, there may be alternative
interpretations to our results, and a few mechanisms may underly the stimulus-response mapping
observed here. While mapping differs from explicit strategies as they are currently defined, we
still lack a comprehensive framework to capture the varying levels and nuanced characteristics of
intentionality and awareness of different mechanisms (Tsay et al., 2024). Additionally, evidence
suggests that walking adaptation involves optimizing motor commands to minimize energy costs
(Finley et al., 2013; Sánchez et al., 2021, 2019, 2017). Therefore, our results may be explained
by extending current optimal control models to account for our finding that motor commands are
computed using internal state estimates and sensory feedback from the environment in tandem.
Future work is needed to develop a generative model capable of capturing the features of
adaptation presented here, and further characterize the nature of the stimulus-response mapping
mechanism of automatic adaptation.

*Conclusions and future directions*

We here characterized two distinct learning mechanisms involved in walking adaptation, both of
which are not under explicit control: recalibration – which leads to motor and perceptual
aftereffects, and mapping – which only changes movement, does not contribute to aftereffects,
and shows meta-learning to different perturbation sizes. Future work should explore whether the
mapping mechanism described here could be of clinical significance: given the combined flexibility
and automaticity, mapping has the potential to ameliorate known problems such as transitioning
between walking environments (Sombric et al., 2017) and dual-tasking (Clark, 2015).
Furthermore, our findings that different people adapt using different learning mechanisms (i.e.,
memory or structure-based mapping) highlights the importance of assessing individual
characteristics of adaptation for both understanding of the neural mechanisms as well as
translation to rehabilitation. Finally, the framework proposed here has important implications for
the development of computational models of adaptation, and may help reconcile different findings
related to savings, energetics, and sources of errors.

**Methods**

<u>Participants</u>

We recruited one-hundred adults (66 females, 23.6 ± 3.9 years old, mean ± SD) for this study, and we reanalyzed data from ten additional adults collected in a previous study (Leech et al., 2018a) (9 female, 21.3 ± 2.9 years old). The protocol was approved by the Johns Hopkins Institutional Review Board and participants provided written informed consent. Participants had no known neurological or musculoskeletal disorders, were naïve to split-belt walking, and participated in only one of the nine experiments.

<u>Data collection</u>

Participants walked on a split-belt treadmill (Woodway, Waukesha, WI, USA) with a thin wooden divider between the belts to avoid stepping on the opposite belt. We controlled the belt speeds with a custom Vizard program (WorldViz), and briefly stopped the treadmill between each phase of the paradigm with the following exceptions: the treadmill was not stopped before the ramp tasks of Experiments 1 and 2 or the first post-adaptation speed match task of the Small Gradual control experiment (people transitioned directly from the preceding walking blocks into the tasks, avoiding abrupt speed changes). We occluded vision and sound of the belt speeds using a cloth drape and headphones that played white noise during the ramp or speed match tasks. A television screen was placed in front of the treadmill and used for these tasks as described later.

Participants wore a non-weight-bearing safety harness, and held on to the treadmill handrail when the treadmill was started or stopped, but were instructed to let go and cross their arms as soon as they began walking. Kinematic data were collected using infrared-emitting markers (Optotrak, Northern Digital, Waterloo, ON, Canada) at 100Hz, placed bilaterally over the toe (fifth metatarsal head), ankle (lateral malleolus), knee (lateral femoral epicondyle), hip (greater trochanter), pelvis (iliac crest), and shoulder (acromion process).

<u>Ramp tasks</u>

For all ramp tasks of Experiments 1 and 2, the speed of the right belt was changed by 0.05m/s every 3 strides (defined below) during right leg swing. The baseline ramp of both experiments consisted of 7 increasing right speeds from 0.35m/s to 0.65m/s. The post-adaptation Ramp Down

of Experiment 1 consisted of 21 decreasing right speeds from 1.5m/s to 0.5m/s. The post-
adaptation Ramp Up & Down of Experiment 2 consisted of 41 total right speeds: 11 increasing
speeds from 1.5m/s to 2m/s, followed by 30 decreasing speeds from 1.95m/s to 0.5m/s. The left
belt speed was constant at 0.5 m/s for all ramp tasks.

In Experiment 1, a keyboard was placed on the treadmill handrail. In the post-adaptation Ramp
Down task, participants were asked to press a button the first time they perceived the right belt to
be (1) as fast as the left, and (2) faster than the left. In the baseline ramp, they press to report the
right belt feeling (1) as fast as the left, and (2) slower than the left. This was explained prior to the
experiment and the above prompts were displayed during the tasks on the TV screen.

As described in SI Appendix, "Ramp Down design", the Ramp Down task was specifically
designed to measure the pattern of aftereffects in a way that ensured reliable and robust
measurements with sufficient resolution across speeds, and that minimized washout to prevent
confounding the results. To balance time constraints with a measurement resolution adequate for
capturing perceptual realignment, we used 0.05 m/s speed decrements, matching the perceptual
sensitivity estimated from our re-analysis of the baseline data from Leech et al. (Leech et al.,
2018a). To obtain robust motor aftereffect measurements, we collected three strides at each
speed condition, as averaging over three strides represents the minimum standard for consistent
and reliable aftereffect estimates in split-belt adaptation (typically used in catch trials) (Leech et
al., 2018a; Rossi et al., 2019; Vazquez et al., 2015). To minimize unwanted washout by forgetting
and/or unlearning, we did not pause the treadmill between adaptation and the post-adaptation
ramp tasks, and ensured the Ramp Down was relatively quick, lasting approximately 80 seconds
on average. Of note, the Ramp Down design ensures that even in cases of partial forgetting, the
emergence pattern of aftereffects remains consistent with the underlying hypotheses.

Questionnaire

At the end of Experiment 2, participants answered this question on a computer: *Did you*
*deliberately change how you walked to account for how fast the belts were moving? If so, describe*
*how. Note: deliberately means that you thought about and decided to move that way. The question*
*refers to the entire central ~20min walking block.* To ensure participants remembered what phase
they were asked about, before adaptation we told participants that the following block will be
called "central ~20min walking block".

 Speed match tasks and control experiments

The protocol and tasks for controls experiments, shown in Fig. 6, were based on previous work
(Leech et al., 2018a; Rossi et al., 2019; Statton et al., 2018; Vazquez et al., 2015). The left belt
was fixed at 0.4m/s for Small Abrupt or Gradual and at 0.5m/s otherwise. The right belt speed
equaled the left during baseline, post-adaptation, and catch trial (administer two-third into
adaptation to all but Small Gradual). Small Gradual adapted for 15 minutes, with the right belt at
0.4 m/s for the first 30 seconds, 0.8 m/s for the last 30 seconds, and increasing linearly in between.
Small Abrupt adapted for 15 minutes with the right belt at 0.8m/s. Short Ascend & Descend,
Medium Ascend & Descend, and Long Ascend adapted for 3, 15, or 30 minutes with the right belt
at 1.5m/s.

Speed match tasks lasted 30 seconds with a time bar displayed on the TV screen. Participants
controlled the right speed with a keypad with 4 buttons: 1) large increment varying between 50,
55 or 65 mm/s as in Vazquez et al. (Vazquez et al., 2015), 2) large -50mm/s decrement, 3) small
5mm/s increment, 4) small -5mm/s decrement. "Ascend" tasks, beginning with the right belt
stationary, were used in baseline for all experiments, and post-adaptation for Short Ascend,
Medium Ascend and Long Ascend (Fig. 6). "Descend" tasks, beginning at the adaptation speed,
were used for post-adaptation of the other experiments.

Motor measures

We defined a stride as the period between two consecutive left heel strikes ($LHS_1$ to $LHS_2$). We
computed motor measures for each stride using equations (1) – (3) (see Results), as in previous
work (Finley et al., 2015; Sombric et al., 2017). The equations terms are defined as follows:

$$\Delta \text{step length} = \text{R step length} - \text{L step length}$$

$$\text{stride length} = \text{R step length} + \text{L step length}$$

$$\Delta \text{step position} = \left( \text{R step position}_{RHS} - \text{L step position}_{LHS_1} \right)$$
$$- \left( \text{L step position}_{LHS_2} - \text{R step position}_{RHS} \right)$$

$$\Delta \text{step time} = \text{R step time} - \text{L step time}$$

$$\text{mean time} \; = \; \frac{\text{R step time} + \text{L step time}}{2}$$

$$\Delta\text{step velocity} \; = \; \text{R step velocity} - \text{L step velocity}$$

$$\text{mean velocity} \; = \; \frac{\text{R step velocity} + \text{L step velocity}}{2}$$

Right ($R$) and left ($L$) step lengths are the anterior-posterior distance between the ankle markers
of the two legs at right heel strike ($RHS$) and left heel strike ($LHS_2$) respectively. Step position is
the anterior-posterior position of the ankle marker, relative to average of the two hip markers, at
heel strike of the same leg. Left and right step times are the times from $LHS_1$ to $RHS$, and from
$RHS$ to $LHS_2$ respectively. Step velocity is the average anterior-posterior velocity of the ankle
marker relative to average of the two hip markers, over the duration of the step (left: $LHS_1$ to $RHS$,
right: $RHS$ to $LHS_2$).

In Experiment 2, we also computed the measure of "strides to plateau" using individual step length
asymmetry data from the adaptation phase only. We first smoothed the data with a 5-point moving
average filter. We then calculated the number of strides till five consecutive strides fell within the
"plateau range", defined as the mean ± 1SD of the last 30 strides of adaptation (Malone and
Bastian, 2010; Rossi et al., 2019).

In both experiments, we assessed how adaptation influences variability by evaluating within-
participant residual variance in step length asymmetry around a double exponential model fit
during adaptation. The double exponential model equation is $A_1 \exp\left\{-\frac{stride}{\tau_1}\right\} + A_2 \exp\left\{-\frac{stride}{\tau_2}\right\}$,
where stride is the stride number in adaptation, and $A_1$, $\tau_1$, $A_2$, $\tau_2$, are model parameters. We
fitted this model to individual participants' step length asymmetry data in adaptation, using the
MATLAB *lsqcurvefit* function. Initial parameters values were set to [-0.177, 322.072 -0.216,
29.662] for Experiment 1 and [-0.177, 418.725, -0.203, 44.307] for Experiment 2, derived from
fitting the model to group mean data (with initial parameters [−0.2,300,−0.2,30]). We computed
residuals as the difference between individual step length asymmetry data and the corresponding
double exponential model fit. We finally computed variability as the variance of these residuals
over (1) the initial 30 strides of adaptation and (2) the final 30 strides of adaptation.

In the control experiments, we interpolated step length asymmetry as a function of speed in the
task. We used locally weighted linear regression (MATLAB '*smooth*', method = 'lowess', span =

20), with query speeds ranging from minimum to maximum speed at which a stride was taken

(average across participants) in steps of 5 mm/s. The speed of a stride refers to the average

speed for the duration of that stride; similarly, when multiple strides were taken at the same speed,

the average step length asymmetry over those strides was used for our analysis.

<u>Clustering analysis</u>

We tested to see if there were separate clusters of participants in our dataset. For Experiment 1,

the measure used for clustering was the number of strides in the Ramp Down with step length

asymmetry above the baseline CI. For Experiment 2, it was the number of strides in the first

portion of the Ramp Up & Down (first 60 strides, teal in Fig. 9) with step length asymmetry below

the baseline CI. For both experiments, "baseline CI" was computed separately for each participant

as the 95% CI of the mean of step length asymmetry data in the second baseline tied-belt block

(after the baseline ramp task; see Statistical Analysis).

We used the MATLAB "dbscan" density-based clustering algorithm (Ester et al., 1996) and

automated the choice of parameters adapting previously published procedures (Naik Gaonkar

and Sawant, 2013; Rahmah and Sitanggang, 2016) (SI Appendix, "Clustering analysis" and

Pseudocode). The resulting algorithm does not require any user input; it automatically assigns

participants to clusters (whose number is not predefined), or labels them as outliers, based solely

on each participant's measure of interest. We also performed a silhouette analysis to assess the

validity of the clusters identified in Experiment 2, using the MATLAB "silhouette" function

(Dalmaijer et al., 2022; Kaufman and Rousseeuw, 1990; Rousseeuw, 1987).

<u>Data fitting</u>

*Recalibration + mapping.*

We developed and fitted our own recalibration + mapping model, presented in the Results section

in Equation (6). The model has one parameter $r$ representing the magnitude of forward model

recalibration. We fitted $r$ to individual participants' Ramp Down data using initial parameter value

$= \frac{1}{2} * p_{\text{plateau}}$, lower bound $= 0$, and upper bound $= p_{\text{plateau}}$, respectively corresponding to half, no,

or total compensation of the adaptation perturbation by recalibration ($p_{\text{plateau}}$ is defined in

Experiment 1 Results).

*Dual state.*

For the recalibration only hypothesis, we fitted the dual state model defined as in Smith et al.

(Smith et al., 2006):

net internal estimate: $x(k) = x_f(k) + x_s(k)$

fast state: $\qquad\qquad x_f(k+1) = A_f * x_f(k) + B_f * \big(p(k) - x(k)\big)$

slow state: $\qquad\qquad x_s(k+1) = A_s * x_s(k) + B_s * \big(p(k) - x(k)\big)$

Δ motor output: $\qquad u(k) = x(k)$

Where $k$ is the stride number, $p$ is the perturbation, $x$ is the net internal estimate of the

perturbation, $x_f$ and $x_s$ are the fast and slow states contributing to the estimate. Participants

generate Δ motor output $u$ that equals the internal estimate of the perturbation $x$ to minimize the

predicted step length asymmetry. The model has four free parameters: $A_f$, $A_s$ are retention

factors controlling forgetting rate, and $B_f$, $B_s$ are error sensitivities controlling learning rate. We

fitted the model parameters using initial values $A_f$=0.92, $A_s$=0.99, $B_f$=0.1, and $B_s$=0.01 (as in

Roemmich et al., 2016), and constraints $0<A_f<A_s<1$ and $0<B_f>B_s<1$ (as defined by the model,

Smith et al., 2006).

*Optimal control.*

We implemented the optimal control theory model using the original equations provided by

Todorov (Todorov, 2005), and defined our state and observation systems to be consistent with

previous implementations of optimal control theory models for motor adaptation (Izawa and

Shadmehr, 2011). Given $k$ = stride number, $p$ = perturbation, $s$ = step length asymmetry, and $u$ =

motor output, the equation provided in Box 1 (capturing the relationship between these standard

parameters of walking adaptation) is rewritten as follows:

$$s(k) = u(k) - p(k)$$

Participants maintain an internal estimate of the perturbation, $\hat{p}$, and update it similar to other
state space models:

$$\hat{p}(k+1) = a\,\hat{p}(k)$$

1105

1106   where $0 \leq a \leq 1$ is a parameter capturing forgetting rate. They use this estimate to predict the
1107   step length asymmetry that will result from a motor command:

$$\hat{s}(k+1) = u(k) - \hat{p}(k)$$

1108

According to optimal control theory, participants select motor commands $u(k)$ that minimize the
cost associated with the motor task – including the base energetic cost of executing motor
commands, plus additional cost associated with the observable state or performance of the motor
system – such as accuracy, stability, state-dependent energy expenditure... (Todorov, 2005). In
walking adaptation, more asymmetric step lengths are linked with worsen stability (Darter et al.,
2018) and energy expenditure (Sánchez et al., 2019). Therefore, we set the observable state $y$
to represent step length asymmetry.

Our system can be written in matrix form in the following way:

Hidden state: $\boldsymbol{x}(k) = \begin{bmatrix} p(k) \\ s(k) \end{bmatrix}$

Observable state: $y(k) = s(k)$.

Hidden state update equation: $\boldsymbol{x}(k+1) = A\boldsymbol{x}(k) + Bu(k) + \boldsymbol{\varepsilon}_x(k)$

Observable state equation: $y(k) = H\boldsymbol{x}(k) + \boldsymbol{\varepsilon}_y(k)$

The system dynamics and observation matrices are $A = \begin{bmatrix} a & 0 \\ -1 & 0 \end{bmatrix}, B = \begin{bmatrix} 0 \\ 1 \end{bmatrix}, H = \begin{bmatrix} 0 & 1 \end{bmatrix}$.

The noise terms are $\boldsymbol{\varepsilon}_x \sim N(0, Q_x), Q_x = diag\left(\sigma_p^2, \sigma_s^2\right), \boldsymbol{\varepsilon}_y \sim N\left(0, Q_y\right), Q_y = \sigma_y^2$.

$a, \sigma_p^2, \sigma_s^2$, and $\sigma_y^2$ are parameters that vary across participants.

Motor output generation: $u(k) = -G(k)\,\hat{\boldsymbol{x}}(k)$

State estimate update: $\hat{x}(k+1) = A\hat{x}(k) + Bu(k) + AK(k)\big(-p(k) + \hat{p}(k)\big)$

Where $u$ is the motor output, $p$ is the perturbation, $\hat{p}$ is the internal estimate of the perturbation,

$\hat{x} = \begin{bmatrix} \hat{p} \\ \hat{s} \end{bmatrix}$ is the internal estimate of the full state in vector form, $G$ is the "feedback gain" matrix

computed to minimize the cost $J$, $K$ is the "Kalman gain" matrix defining the rate of learning from

the sensory prediction error, and $A$ and $B$ are matrices defined as above. Note that we defined

the sensory prediction error as the difference between the estimated and actual perturbation

$\big(-p(k) + \hat{p}(k)\big)$ to be consistent with previous implementations of optimal control theory for the

case of motor adaptation (Izawa and Shadmehr, 2011). Specifically, this is derived as follows:

Sensory prediction error = $y(k) - \hat{y}(k)$

$= s(k) - \hat{s}(k) = \big(u(k) - p(k)\big) - \big(u(k) - \hat{p}(k)\big) = -p(k) + \hat{p}(k)$

The feedback and Kalman gain are computed recursively using equations provided by Todorov

(Todorov, 2005). We fit this model to individual participants' motor output data similarly to how we

performed the dual state fits. The optimal control theory model implemented here has 6

parameters: $a$ = forgetting factor for the perturbation estimate, $\sigma_p^2$ = variance of the perturbation

estimate, $\sigma_s^2$ = variance of the step length asymmetry estimate, $\sigma_y^2$ = observation variance, $Q$ =

cost associated with step length asymmetry, $L$ = cost associated with motor output. We set the

initial parameter values to [0.5, 0.1, 0.1, 0.1, 1000, 1000], lower bounds to [0, 0, 0, 0, 0, 0], and

upper bounds to [1, 10, 10, 10, 10^6, 10^6].

*Memory of errors*

We implemented the memory of errors model as defined by Herzfeld et al. (Herzfeld et al., 2014).

Similar to the dual state, the model updates an internal estimate of the perturbation, and uses it

to produce a motor output equal to this perturbation. However, the learning rate is not fixed, but

varies on each trial depending on the error and the history of errors:

$u(k+1) = au(k) + \eta(k, e(k))\, e(k)$

$e(k) = p(k) - u(k)$

Where $k$ is the stride number, $u$ is the motor output, $p$ is the perturbation, $e$ is the sensory
prediction error, $a$ is a parameter capturing forgetting, and $\eta$ is the learning rate. As mentioned,
the learning rate depends on the error and history of errors. Specifically, it is computed on each
trial as follows:

1154
$$\eta\big(k, e(k)\big) = \boldsymbol{w}(k)^T \boldsymbol{g}(e(k))$$

1155
$$\boldsymbol{g}\big(e(k)\big) = \exp\left\{\frac{-(e(k) - \breve{e})^2}{2\sigma^2}\right\}$$

1156
$$\boldsymbol{w}(k+1) = \boldsymbol{w}(k) + \beta \operatorname{sign}\{e(k-1)e(k)\} \frac{\boldsymbol{g}\big(e(k-1)\big)}{\boldsymbol{g}\big(e(k-1)\big)^T \boldsymbol{g}\big(e(k-1)\big)}$$

Each $\boldsymbol{g}\big(e(k)\big)$ is a basis element with preferred error $\breve{e}$, so that $\boldsymbol{g}\big(e(k)\big)$ has maximum value when
$e(k) = \breve{e}$, and the value progressively decays, similar to a normal distribution, for $e(k)$ smaller or
larger than $\breve{e}$. $\sigma^2$ corresponds to the variance in a normal distribution, and here captures how
sharply does $\boldsymbol{g}\big(e(k)\big)$ decay as the difference between $e(k)$ and $\breve{e}$ increases.

$\boldsymbol{w}(k)$ is a vector that defines error sensitivity, and is updated each trial to increase sensitivity for
errors that are experienced most consistently. Specifically, it is a vector that stores a scaling factor
for each basis error center contained in the vector $\breve{e}$: the learning rate is higher in response to
errors associated with larger $\boldsymbol{w}$ scaling factor. $\boldsymbol{w}$ is updated such that if the error on one trial has
the same sign as the error on the previous trial, the sensitivity to that error increases, otherwise it
decreases. The model updates the entire $\boldsymbol{w}$ vector, rather than just the element corresponding to
experienced error. The amount of the update is proportional to $\boldsymbol{g}\big(e(k-1)\big)$, meaning that
sensitivities to errors similar to that experienced on this trial are updated the most, while
sensitivities to errors different from that experienced on this trial are updated the least. The update
magnitude is additionally scaled by the learning rate $\beta$.

We set the initial value of $\boldsymbol{w}$ as follows:

1172
$$w_i = \frac{\eta_0}{\sum_j g(0)_j}, \quad for\ all\ w_i\ elements\ of\ \boldsymbol{w}$$

1173
$$g(0) = \exp\left\{\frac{-(0 - \breve{e})^2}{2\sigma^2}\right\}$$

Where $\sum_j g(0)_j$ is the sum of all elements of $g(0)$, and $\eta_0$ is a parameter representing the naïve
learning rate.

In our study, this model captures the scenario where the learning rate is largest for step length
asymmetry values that have been experienced the most and most consistently. We fitted the
model parameters, $\sigma^2$, $\beta$, $\eta_0$, and $a$ using initial values [0.5, 0.001, 0.001, 0.9] and bounds [0, 0,
0, 0] (lower) and [10, 1, 1, 1] (upper).

*Proprioceptive re-alignment model (PReMo)*

We evaluated the PReMo model developed by Tsay et al. (Tsay et al., 2022, 2021), using the
following equivalencies with the walking adaptation variables (SI Appendix, "Evaluation and
development of perceptual models"): $x_p$= step length asymmetry, $\sigma_p^2, \sigma_u^2$ = proprioceptive and
sensory prediction uncertainties, $K$ = learning rate. For the model fits, we set the goal $G = 0$, as it
represents explicit strategies (but alternative values are considered in the SI Appendix).

The model operates by first integrating proprioceptive and predictive estimates for step length
asymmetry (where prediction = goal):

*Integrated estimate*:   $x_p^I(k) = \frac{\sigma_u^2}{\sigma_u^2+\sigma_p^2} x_p(k) + \frac{\sigma_p^2}{\sigma_u^2+\sigma_p^2} G(k)$

In the original model, perception is further shifted by the visual-proprioceptive mismatch.
However, this proprioceptive shift is zero in the absence of visual feedback, and step length
asymmetry perception depends only on the integrated estimate:

*Proprioceptive shift*:   $\beta_p(k) \propto \left(visual\ information - x_p^I(k)\right) = 0$

*Perceived step length asymmetry*:   $x_p^{per}(k) = x_p^I(k) + \beta_p(k) \equiv x_p^I(k)$

Perceptual error drives implicit adaptation of step length asymmetry:

*Implicit adaptation*:   $x_p(k + 1) = x_p(k) + K\left(G(k) - x_p^{per}(k)\right)$

The original model does not contain a perturbation term, which is needed to fit the model to the
Ramp Down data. To introduce this, we replace the implicit adaptation equation above with

separate observation and hidden state update equations, where the hidden state $x(k)$ represents

Δ motor output and $p(k)$ is the perturbation:

*Step length asymmetry observation:* $\qquad x_p(k) = x(k) - p(k)$

*Hidden state update:* $\qquad x(k+1) = x(k) + K\left(G(k) - x_p^{per}(k)\right)$

While the original model does not explicitly define the perceived perturbation, we can infer this

equation using the relationship between step length asymmetry, perturbation, and Δ motor output:

$$p(k) = x(k) - x_p(k)$$

1204

$$\{p^{per}(k) - p(k)\} = \{x^{per}(k) - x(k)\} - \{x_p^{per}(k) - x_p(k)\}$$

1205

We assume no shift in the perception of Δ motor output as this is not defined by the model:

*Perceived perturbation:* $\qquad p^{per}(k) = p(k) - \{x_p^{per}(k) - x_p(k)\}$

We used $p^{per}$ to obtain model predictions for the perception of belt speed difference, setting the

prediction to "right leg feels faster" when $p^{per} > 0$, "feels equal" when $p^{per} = 0$, and "feels slower"

when $p^{per} < 0$.

To limit redundancy during model fitting, we substitute individual uncertainty terms with the

relative weight variable $W_p = \frac{\sigma_u^2}{\sigma_u^2 + \sigma_p^2}$, where $1 - W_p = \frac{\sigma_p^2}{\sigma_u^2 + \sigma_p^2}$. We then fitted the two model

parameters $K$ and $W_p$ to individual data. The initial value of $W_p$ was set to $\frac{1}{3}$ based on results from

Zhang et al. showing that $\sigma_p^2 \approx 2\sigma_u^2$ (Zhang et al., 2024). The initial value of the learning rate $K$

was set to 0.03; this value was selected to ensure that the time course of learning of $x_p$ would

match that of recalibration observed in control experiments (learning of $x_p$ after 3 minutes reached

78% of the learning observed after 15 minutes; similarly, recalibration in the Short groups was

78% of that in the Medium groups – both averaged across Ascend and Descend group means).

We set lower bounds = [0, 0] and upper bounds = [1, 1], and simulated the entire paradigm but

computed error minimization on the Ramp Down task only as explained before.

*Perceptual Error Adaptation model (PEA)*

We evaluated the PEA model developed by Zhang et al., which models perception as the

Bayesian integration of any available sensory modalities as well as sensory predictions (Zhang

et al., 2024). We set the following variable equivalencies: $x_p$= step length asymmetry, $\sigma_p^2, \sigma_u^2 =$

proprioceptive and sensory prediction uncertainties, $A$ = retention rate, $B$ = learning rate. We set

target $T$ = 0. As split-belt adaptation does not involve vision, we do not have the variables $x_v$ and

$\sigma_v^2$ and simplify the model accordingly.

The perceived movement is the Bayesian integration of proprioceptive and predictive information:

*Perceived movement:* $\hat{x}_p(k) = W_p\, x_p(k) + (1 - W_p)\, T(k)$

*With* $\quad W_p = \dfrac{\frac{1}{\sigma_p^2}}{\frac{1}{\sigma_u^2} + \frac{1}{\sigma_p^2}}$

Motor adaptation is driven by the perceptual error – the discrepancy between perceived and target

movement:

*Adaptation:* $\quad x_p(k + 1) = A\, x_p(k) + B\left(T(k) - \hat{x}_p(k)\right)$

1234

Perceived and actual step length asymmetry are further combined in a Bayesian manner to

produce the step length asymmetry report used for the perceptual task:

*Reported step length asymmetry:* $x_{report}(k) = W_R\, x_p(k) + (1 - W_R)\, \hat{x}_p(k)$

*With* $\quad W_R = \dfrac{W_p}{1 + W_p}$

We finally introduced variables for hidden state $x(k)$, perturbation $p(k)$, and reported perceived

perturbation $p^{per}(k)$ (used to predict speed difference perception) as we did for PReMo:

*Step length asymmetry observation:* $\quad x_p(k) = x(k) - p(k)$

*Hidden state update:* $\quad x(k + 1) = A\, x(k) + B\left(T(k) - \hat{x}_p(k)\right)$

*Reported perceived perturbation:* $\quad p^{per}(k) = p(k) - \left\{x_{report}(k) - x_p(k)\right\}$

We fitted the model parameters $A, B$, and $W_p$ to individual data using initial values = [0.97, 0.2,

1/3], selected approximately based on the average parameter values reported by Zhang et al.

across experiments. We set lower bounds = [0, 0, 0], upper bounds = [1, 1, 1].

*Perceptuomotor recalibration + mapping (PM-ReMap)*

As described in SI Appendix, "Evaluation and development of perceptual models", PM-ReMap

was formulated as an extension of PReMo that addressed its key limitations. The first limitation

was that PReMo assumes perceptual realignment arises from mismatches between sensory

modalities, such as vision and proprioception. We addressed this by accounting for perceptual

realignment driven by mismatches between predicted and actual sensory outcomes of motor

commands. The second limitation was that PReMo assumes perceptual realignment changes

immediately to reflect a proportion of the perturbation on each stride – causing it to return to zero

in the Ramp Down. We addressed this by incorporating a learning rate that mediates gradual

changes in perceptual realignment. The third limitation was that PReMo lacks an automatic

mapping mechanism capable of driving immediate changes in motor output. While it includes a

variable that may function as a mapping mechanism, this variable relies on an explicit strategy,

adjusts motor output gradually through a learning rate, and lacks a defined equation to determine

its value. We addressed this limitation by defining an equation for this variable consistent with an

automatic mapping mechanism and removing the learning rate to enable immediate changes in

motor output.

The final model equations are as follows:

*Actual step length asymmetry*:   $x_v(k) = x_p(k) - p(k)$

*Integrated estimate for motor output*:   $x_p^I(k) = \frac{\sigma_u^2}{\sigma_u^2 + \sigma_p^2} x_p(k) + \frac{\sigma_p^2}{\sigma_u^2 + \sigma_p^2} G(k)$

*Integrated estimate for step length asymmetry*:   $x_v^I(k) = \frac{\sigma_u^2}{\sigma_u^2 + \sigma_v^2} x_v(k) + \frac{\sigma_v^2}{\sigma_u^2 + \sigma_v^2} G(k)$

*Perceptual shift for motor output:*   $\beta_p^* = \eta_p \left( x_v^I(k) - x_p^I(k) \right)$

$$\beta_p(k+1) = \beta_p(k) + K \left( \beta_p^* - \beta_p(k) \right)$$

*Perceptual shift for step length asymmetry:*   $\beta_v^* = \eta_v \left( x_p^I(k) - x_v^I(k) \right)$

$$\beta_v(k+1) = \beta_v(k) + K \left( \beta_v^* - \beta_v(k) \right)$$

*Perceived motor output:*   $x_p^{per}(k) = x_p^I(k) + \beta_p(k)$

*Perceived step length asymmetry:*   $x_v^{per}(k) = x_v^I(k) + \beta_v(k)$

*Perceived perturbation:*   $p^{per}(k) = x_p^{per} - x_v^{per}$

| 1274 | *Adaptation of motor output:* | $x_p(k+1) = x_p(k) + G(k) - x_p^{per}(k)$ |
|---|---|---|

*Adaptation of motor output:*      $x_p(k+1) = x_p(k) + G(k) - x_p^{per}(k)$

*Goal for implicit adaptation simulations:*      $G(k) = 0$

*Goal for mapping simulations (Ramp Down only):*      $G^* = \frac{\beta_p(k)}{W_p} + p(k+1)$

$$G(k+1) = \begin{cases} G^* & if \quad G^* \geq 0 \\ 0 & otherwise \end{cases}$$

We used $p^{per}$ to obtain model predictions for the perception of belt speed difference as explained

for PReMo. To obtain the estimated point of subjective equality $\widehat{PSE}$, we first determined the stride

at which $p^{per}$ crossed zero (sign flips from positive to negative). We then computed $\widehat{PSE}$ as the

perturbation at that stride normalized to the perturbation at adaptation plateau: $\widehat{PSE} = \frac{p(k_{\text{sign-flip}})}{p_{\text{plateau}}}$.

To limit redundancy during model fitting, we substitute individual uncertainty with a relative weight

variable $W_p = \frac{\sigma_u^2}{\sigma_u^2 + \sigma_p^2}$, and $1 - W_p = \frac{\sigma_p^2}{\sigma_u^2 + \sigma_p^2}$. We fitted the model parameters $K$ and $W_p$ to individual

data using initial values = [0.5, 0.01, 1/3], lower bounds = [0, 0, 0], upper bounds = [1, 1, 1]. Similar

to the recalibration + mapping model, we only simulated and fitted the Ramp Down task.

*General fitting procedure*

We fitted each model to individual participants' Δ motor output data in the Ramp Down.

Specifically, we found parameter values that minimize the residual sum of squares between the

Ramp Down Δ motor output data and the modelled $u$. Dual state, optimal control, memory of

errors, PReMo, and PEA models involve hidden states that depend on the history of perturbations

and are not expected to be zero at the start of the Ramp Down. To account for this, we simulated

these models over the entire paradigm, but computed the residual sum of squares solely on the

Ramp Down to ensure fair model comparison with the recalibration + mapping model. We used

the *fmincon* MATLAB function with constraint and optimality tolerances tightened to 10^-20 (as in

Roemmich et al., 2016).

<u>Statistical analysis</u>

Statistical tests were performed in MATLAB with significance level α=0.05 (two-sided). For

Experiment 1, we performed within-group statistical analyses to compare the following measures

to zero. For each participant and for each speed in the ramp tasks, we first averaged step length
asymmetry over the 3 strides taken at that speed to obtain the measures listed in a) and b):

a) Step length asymmetry for each of the 7 speeds in the baseline ramp task (m=7)

b) Step length asymmetry for each of the 21 speeds in the Ramp Down task (m=21)

c) Difference in BIC between our model (recalibration + mapping) and each of the alternative
models (dual state, optimal feedback control, memory of errors, PEA, PReMo, PM-
ReMap) (m=6)

d) Difference between $compensation_{\text{motor total}}$ or $compensation_{\text{motor recalibration}}$, and the upper
and lower bounds of $compensation_{\text{perceptual}}$ (m=4)

Each item on the list represents a "family of related tests" (measures considered together for
multiple comparisons), where "m" is the number of tests in the family. We computed bootstrap
distributions of each measure by generating 10,000 bootstrapped samples of 20 participants
(resampled with replacement from Experiment 1) and averaging the measure over participants in
each sample (Efron and Tibshirani, 1994). We then computed confidence intervals (CI) corrected
for multiple comparisons using the False Discovery Rate procedure (FDR) (Benjamini and
Yekutieli, 2005). That is, for each family of related tests, we adjusted the significance level to
$\alpha_{\text{corr}} = \alpha * \frac{R}{m} = 0.05 * \frac{R}{m}$, where "m" is the total number of tests and "R" is number of tests deemed
significant for $\alpha = 0.05$. Note that we report FDR-corrected CIs in the Results sections. A test is
significant if the corrected CI does not overlap zero, indicating that the measure is significantly
different from zero (Cumming and Finch, 2005; Efron and Tibshirani, 1994).

For both experiments, we computed within-participant CIs for the mean baseline step length
asymmetry. For each participant, we generated 10,000 bootstrapped samples of N strides,
resampled with replacement from 2-min baseline block following the ramp task, where N is the
number of strides in this phase. We averaged step length asymmetry over strides in each sample
and computed the 95% CI.

For Experiment 2, we performed between-group statistical analyses to assess whether the
following measures differed between the memory-based and structure-based subgroups:

a) Number of strides to plateau (m=1)

b) Number of strides in teal portion of the Ramp Up & Down (first 60 strides) with step length
asymmetry below the within-participant baseline CI (as described above) (m=1)

We generated 10,000 bootstrapped samples, each comprising 20 participants: 12 from the
memory-based subgroup and 8 from the structure-based subgroup (resampled with
replacement). For each sample "b", we averaged the measure of interest over participants
resampled from each subgroup to obtain $\mu_{memory}(b)$ and $\mu_{structure}(b)$, and evaluated the
difference of the means between the subgroups: $\Delta\mu(b) = \mu_{memory}(b) - \mu_{structure}(b)$. We computed
the CI for this difference, correcting for multiple comparisons using FDR as explained for
Experiment 1.

For the control experiments, we performed within-group statistical analyses to compare the
following measures to zero (or 100% as defined), using the same procedure as Experiment 1:

a) $compensation_{perceptual}$ (m=7, counting one comparison per group), computed using the
first post-adaptation task (normalized to 0.4m/s for Small Gradual)

b) $compensation_{motor\ total}$ (compared to 0 and to 100%, each with m=7)

c) Step length asymmetry in the first stride of the first post-adaptation task (m=7)

d) Step length asymmetry in the last stride of the first post-adaptation task (m=7)

e) Motor aftereffects minus perceptual realignment, both normalized as defined in
Experiment 1 *"Perceptual test and results"*, for each of the six speed match tasks post-
adaptation in the Ascend groups (m=6 for each group)

We compared the following measures between groups using the same procedure as Experiment
2:

a) PSE (m=2, comparing Medium Ascend or Descend to Experiment 1)

b) $compensation_{motor\ total}$ (m=6)

c) recalibration contribution to adaptation $= \dfrac{compensation_{perceptual}}{compensation_{motor\ total}}$ (m=6)

1351

**Boxes**

*Box 1. Split-belt walking adaptation paradigm and motor responses.*

There are three established motor measurements that are used to quantify split-belt walking
adaptation (Finley et al., 2015; Reisman et al., 2005): "Perturbation" measures the effect of the
split-belt treadmill on the stepping pattern — the total movement error we would see in the
absence of adaptation. "Δ motor output" measures the extent that individuals compensate for the
perturbation by changing their stepping pattern — how much they alter when (time) and where
(position) they step on the treadmill with each foot. "Step length asymmetry" measures the
remaining movement error — the difference between Δ motor output and perturbation (see
Methods).

Box 1-Figure 1A illustrates the standard paradigm used in studies of split-belt adaptation, with
abrupt transitions between tied-belt to split-belt phases, and Box 1-Figure 1B illustrates the
standard motor response. In the "baseline" phase, the perturbation (red), Δ motor output (blue),
and step length asymmetry (purple) are zero, reflecting equal belt speeds and symmetric
walking. In the "adaptation" phase, the belt speeds are different and the perturbation is positive.
Initially, this leads to movement errors observed as the negative step length asymmetry. The Δ
motor output gradually adapts to compensate for the perturbation, so that the step length
asymmetry returns to zero. In "post-adaptation", the belt speeds are tied and the perturbation is
again zero. Individuals exhibit initial movement errors called "aftereffects": the Δ motor output
mismatches the perturbation (it remains elevated) and step length asymmetry is positive (Leech
et al., 2018a; Rossi et al., 2021a, 2019; Statton et al., 2018).

*Box 2. Relevant methodologies and results from prior work on perceptual recalibration during*
*locomotor learning.*

Box 2-Figure 1 illustrates selected portions of the "speed match" paradigm manipulation used in
prior work and the respective step length asymmetry data (Leech et al., 2018a). The paradigm
and data shown in Box 2-Figure 1A are consistent with that explained in Box 1 (with the addition

of a brief tied-belt catch trial in adaptation). Note that participants walk with near-zero step length asymmetry by the end of adaptation and exhibit large aftereffects post-adaptation.

The study investigated "perceptual realignment", where perception of speed becomes biased in adaptation, partially compensating for the speed difference so that it feels smaller. They measured perception of speed with "speed match" tasks: participants control the speed of the right belt and try to match it to that of the left belt (as described later in our Control experiments).

Left and right panels Box 2-Figure 1B depict the tasks performed before and after adaptation – top row shows belt speeds and bottom row shows step length asymmetry. Before adaptation, participants can accurately match the speeds and walk symmetrically at this near-tied-belt configuration (right and left speeds are ~equal and step length asymmetry is ~zero at the end of the "before adapt" task). After adaptation, participants overshoot the speed of the right belt and select a speed configuration that is biased towards that experienced in adaptation, perceiving this as "equal speeds" (the right speed is intermediate between the adaptation right speed and the target left speed at the end of the "after adapt" task). The key result is that participants walk with near-zero step length asymmetry at this configuration ("after adapt", bottom row).

This suggests that participants may have learned to walk symmetrically at two distinct speed configurations: the adaptation configuration, and the configuration that they perceive as "equal speeds". As illustrated in a later section (Results: Experiment 1 – Motor paradigm and hypotheses), this flexible behavior may be indicative of stimulus-response mapping mechanisms.

**Data Availability**

*Note to reviewers: the dataset cited below is currently "private for peer review", and can be accessed by the alternative "reviewer URL" provided by the Dryad database:*

*https://datadryad.org/stash/share/Qa0pD7rMFT7kVgDeP-8PMYKBaCNuy4yx1VGIQlpVgvM*

All data and code used for the study have been deposited in: Rossi, Cristina (2023), Dataset for automatic learning mechanisms for flexible human locomotion, Dryad, Dataset, https://doi.org/10.5061/dryad.18931zd27.

**Acknowledgments**

Supported by NIH grant 5 R37 NS090610 to AJB, American Heart Association predoctoral
fellowship 20PRE35180131 to CR, and NIH grant K01 AG073467 to KAL. We thank Daniel
Wolpert, Adrian Haith, Jonathan Tsay, and Amanda Therrien for helpful discussions and
comments on data analysis.

**Author Contributions:** Study design & implementation: C.R., K.A.L., R.T.R., A.J.B.; Data
acquisition & analysis: C.R., K.A.L.; Interpretation of results: C.R., K.A.L., R.T.R., A.J.B.; Writing
& revision: C.R., K.A.L., R.T.R., A.J.B.

**Competing Interest Statement:** The authors declare no competing interests.

**Classification:** Biological Sciences: Neuroscience.

**Keywords:** Adaptation. Motor control. Motor learning. Human locomotion. Perception.

1416

1417

**References**

't Hart BM, Henriques DYP. 2016. Separating predicted and perceived sensory consequences of motor learning. *PLoS One* **11**:e0163556. doi:10.1371/journal.pone.0163556

Abeele S, Bock O. 2001. Mechanisms for sensorimotor adaptation to rotated visual input. *Exp Brain Res* **139**:248–253. doi:10.1007/S002210100768

Anderson SR, Porrill J, Pearson MJ, Pipe AG, Prescott TJ, Dean P. 2012. An Internal Model Architecture for Novelty Detection: Implications for Cerebellar and Collicular Roles in Sensory Processing. *PLoS One* **7**:e44560. doi:10.1371/journal.pone.0044560

Avraham G, Morehead JR, Kim HE, Ivry RB. 2021. Reexposure to a sensorimotor perturbation produces opposite effects on explicit and implicit learning processes. *PLoS Biol* **19**:e3001147. doi:10.1371/journal.pbio.3001147

Bastian AJ. 2011. Moving, sensing and learning with cerebellar damage. *Curr Opin Neurobiol* **21**:596–601. doi:10.1016/j.conb.2011.06.007

Bastian AJ. 2008. Understanding sensorimotor adaptation and learning for rehabilitation. *Curr Opin Neurol* **21**:628–633. doi:10.1097/WCO.0b013e328315a293

Benjamini Y, Yekutieli D. 2005. False discovery rate-adjusted multiple confidence intervals for selected parameters. *J Am Stat Assoc* **100**:71–81. doi:10.1198/016214504000001907

Bhanpuri NH, Okamura AM, Bastian AJ. 2013. Predictive modeling by the cerebellum improves proprioception. *J Neurosci* **33**:14301–14306. doi:10.1523/JNEUROSCI.0784-13.2013

Blakemore S-J, Wolpert DM, Frith CD. 1998. Central cancellation of self-produced tickle sensation. *Nat Neurosci* **1**:635–640.

Bond KM, Taylor JA. 2017. Structural learning in a visuomotor adaptation task is explicitly accessible. *eNeuro* **4**:e0122-17. doi:10.1523/ENEURO.0122-17.2017

Bond KM, Taylor JA. 2015. Flexible explicit but rigid implicit learning in a visuomotor adaptation task. *J Neurophysiol* **113**:3836–3849. doi:10.1152/jn.00009.2015

Braun DA, Aertsen A, Wolpert DM, Mehring C. 2009. Motor Task Variation Induces Structural Learning. *Curr Biol* **19**:352–7. doi:10.1016/j.cub.2009.01.036

Brooks JX, Cullen KE. 2019. Predictive Sensing: The Role of Motor Signals in Sensory Processing. *Biol psychiatry Cogn Neurosci neuroimaging* **4**:842. doi:10.1016/J.BPSC.2019.06.003

Camacho S, Dop M, de Graaf C, Stieger M. 2015. Just Noticeable Differences and Weber Fraction of Oral Thickness Perception of Model Beverages. *J Food Sci* **80**:S1583–S1588. doi:10.1111/1750-3841.12922

Chase SM, Kass RE, Schwartz AB. 2012. Behavioral and neural correlates of visuomotor adaptation observed through a brain-computer interface in primary motor cortex. *J Neurophysiol* **108**:624–644. doi:10.1152/jn.00371.2011

Cheng S, Sabes PN. 2006. Modeling Sensorimotor Learning with Linear Dynamical Systems. *Neural Comput* **18**:760–793. doi:10.1162/089976606775774651

Choi JT, Vining EPG, Reisman DS, Bastian AJ. 2009. Walking flexibility after hemispherectomy: Split-belt treadmill adaptation and feedback control. *Brain* **132**:722–733. doi:10.1093/brain/awn333

Clark DJ. 2015. Automaticity of walking: Functional significance, mechanisms, measurement and rehabilitation strategies. *Front Hum Neurosci* **9**:246. doi:10.3389/fnhum.2015.00246

Codol O, Holland PJ, Galea JM. 2018. The relationship between reinforcement and explicit control during visuomotor adaptation. *Sci Rep* **8**:9121. doi:10.1038/s41598-018-27378-1

Coltman SK, Cashaback JGA, Gribble PL. 2019. Both fast and slow learning processes contribute to savings following sensorimotor adaptation. *J Neurophysiol* **121**:1575–1583. doi:10.1152/jn.00794.2018

Cumming G, Finch S. 2005. Inference by eye: confidence intervals and how to read pictures of data. *Am Psychol* **60**:170–180. doi:10.1037/0003-066X.60.2.170

Dalmaijer ES, Nord CL, Astle DE. 2022. Statistical power for cluster analysis. *BMC Bioinformatics* **23**:1–28. doi:10.1186/S12859-022-04675-1/TABLES/5

Darter BJ, Labrecque BA, Perera RA. 2018. Dynamic stability during split-belt walking and the relationship with step length symmetry. *Gait Posture* **62**:86–91. doi:10.1016/j.gaitpost.2018.03.006

Dassonville P, Lewis SM, Zhu XH, Ugurbil K, Kim SG, Ashe J. 2001. The effect of stimulus-response

compatibility on cortical motor activation. *Neuroimage* **13**:1–14. doi:10.1006/nimg.2000.0671

Day KA, Leech KA, Roemmich RT, Bastian AJ. 2018. Accelerating locomotor savings in learning: Compressing four training days to one. *J Neurophysiol* **119**:2100–2113. doi:10.1152/jn.00903.2017

Delval A, Bayot M, Defebvre L, Dujardin K. 2020. Cortical oscillations during gait: Wouldn't walking be so automatic? *Brain Sci*. doi:10.3390/brainsci10020090

Diedrichsen J, King M, Hernandez-Castillo C, Sereno M, Ivry RB. 2019. Universal Transform or Multiple Functionality? Understanding the Contribution of the Human Cerebellum across Task Domains. *Neuron* **102**:918–928. doi:10.1016/j.neuron.2019.04.021

Diedrichsen J, White O, Newman D, Lally N. 2010. Use-dependent and error-based learning of motor behaviors. *J Neurosci* **30**:5159–5166. doi:10.1523/JNEUROSCI.5406-09.2010

Drew T, Marigold DS. 2015. Taking the next step: Cortical contributions to the control of locomotion. *Curr Opin Neurobiol* **33**:25–33. doi:10.1016/j.conb.2015.01.011

Efron B, Tibshirani RJ. 1994. An Introduction to the Bootstrap, An Introduction to the Bootstrap. doi:10.1201/9780429246593

Ester M, Kriegel H-P, Sander J, Xu X. 1996. A Density-Based Algorithm for Discovering Clusters in Large Spatial Databases with NoiseProceedings of the 2nd International Conference on Knowledge Discovery and Data Mining. pp. 226–231.

Finley JM, Bastian AJ, Gottschall JS. 2013. Learning to be economical: The energy cost of walking tracks motor adaptation. *J Physiol* **591**:1081–1095. doi:10.1113/jphysiol.2012.245506

Finley JM, Long A, Bastian AJ, Torres-Oviedo G. 2015. Spatial and Temporal Control Contribute to Step Length Asymmetry during Split-Belt Adaptation and Hemiparetic Gait. *Neurorehabil Neural Repair* **29**:786–795. doi:10.1177/1545968314567149

Gescheider GA. 1997. The Classical Psychophysical MethodsPsychophysics: The Fundamentals. Mahwah, NJ, US: Lawrence Erlbaum Associates Publishers. pp. 45–72.

Goldman-Rakic PS. 1987. Motor control function of the prefrontal cortex. *Ciba Found Symp* **132**:187–200. doi:10.1002/9780470513545.ch12

Gonzalez-Rubio M, Velasquez NF, Torres-Oviedo G. 2019. Explicit control of step timing during split-belt walking reveals interdependent recalibration of movements in space and time. *Front Hum Neurosci* **13**:207. doi:10.3389/fnhum.2019.00207

Haith A, Jackson C, Miall C, Vijayakumar S. 2008. Unifying the sensory and motor components of sensorimotor adaptationAdvances in Neural Information Processing Systems 21 - Proceedings of the 2008 Conference. pp. 593–600.

Haith AM, Krakauer JW. 2013. Model-based and model-free mechanisms of human motor learning. *Adv Exp Med Biol* **782**:1–21. doi:10.1007/978-1-4614-5465-6_1

Harris CS. 1963. Adaptation to Displaced Vision: Visual, Motor, or Proprioceptive Change? *Science (80- )* **140**:812–813. doi:10.1017/CBO9781107415324.004

Haruno M, Wolpert DM, Kawato M. 2001. MOSAIC Model for Sensorimotor Learning and Control. *Neural Comput* **13**:2201–2220. doi:10.1162/089976601750541778

Herzfeld DJ, Kojima Y, Soetedjo R, Shadmehr R. 2018. Encoding of error and learning to correct that error by the Purkinje cells of the cerebellum. *Nat Neurosci* **21**:736–743. doi:10.1038/s41593-018-0136-y

Herzfeld DJ, Vaswani PA, Marko MK, Shadmehr R. 2014. A memory of errors in sensorimotor learning. *Science (80- )* **345**:1349–1353. doi:10.1126/science.1253138

Hinton DC, Conradsson D, Bouyer L, Paquette C. 2020. Does dual task placement and duration affect split-belt treadmill adaptation? *Gait Posture* **75**:115–120. doi:10.1016/j.gaitpost.2019.10.005

Hinton DC, Thiel A, Soucy JP, Bouyer L, Paquette C. 2019. Adjusting gait step-by-step: Brain activation during split-belt treadmill walking. *Neuroimage* **202**:116095. doi:10.1016/j.neuroimage.2019.116095

Hirata K, Kokubun T, Miyazawa T, Yokoyama H, Kubota K, Sonoo M, Hanawa H, Kanemura N. 2019. Contribution of Lower Limb Joint Movement in Adapting to Re-establish Step Length Symmetry During Split-Belt Treadmill Walking. *J Med Biol Eng* **39**:693–701. doi:10.1007/s40846-018-0456-0

Honda T, Nagao S, Hashimoto Y, Ishikawa K, Yokota T, Mizusawa H, Ito M. 2018. Tandem internal models execute motor learning in the cerebellum. *Proc Natl Acad Sci U S A* **115**:7428–7433. doi:10.1073/pnas.1716489115

Huberdeau DM, Krakauer JW, Haith AM. 2019. Practice induces a qualitative change in the memory

representation for visuomotor learning. *J Neurophysiol* **122**:1050–1059. doi:10.1152/jn.00830.2018

Huberdeau DM, Krakauer JW, Haith AM. 2015. Dual-process decomposition in human sensorimotor adaptation. *Curr Opin Neurobiol* **33**:71–77. doi:10.1016/j.conb.2015.03.003

Inoue M, Uchimura M, Karibe A, O'Shea J, Rossetti Y, Kitazawa S. 2015. Three timescales in prism adaptation. *J Neurophysiol* **113**:328–338. doi:10.1152/jn.00803.2013

Ito M. 1989. Long-term depression. *Annu Rev Neurosci* **12**:85–102.

Ito M. 1984. The cerebellum and neural control, Raven, New York. Raven Press. doi:10.1016/0090-3019(85)90320-9

Iturralde PA, Torres-Oviedo G. 2019. Corrective Muscle Activity Reveals Subject-Specific Sensorimotor Recalibration. *eNeuro* **6**. doi:10.1523/ENEURO.0358-18.2019

Izawa J, Criscimagna-Hemminger SE, Shadmehr R. 2012. Cerebellar contributions to reach adaptation and learning sensory consequences of action. *J Neurosci* **32**:4230–4239. doi:10.1523/JNEUROSCI.6353-11.2012

Izawa J, Shadmehr R. 2011. Learning from sensory and reward prediction errors during motor adaptation. *PLoS Comput Biol* **7**:e1002012. doi:10.1371/journal.pcbi.1002012

Jensen L, Prokop T, Dietz V. 1998. Adaptational effects during human split-belt walking: influence of afferent input. *Exp Brain Res* **118**:126–130. doi:10.1007/s002210050262

Ju C, Bosman LWJ, Hoogland TM, Velauthapillai A, Murugesan P, Warnaar P, van Genderen RM, Negrello M, De Zeeuw CI. 2019. Neurons of the inferior olive respond to broad classes of sensory input while subject to homeostatic control. *J Physiol* **597**:2483–2514. doi:10.1113/JP277413

Judd EN, Lewis SM, Person AL. 2021. Diverse inhibitory projections from the cerebellar interposed nucleus. *Elife* **10**:e66231. doi:10.7554/eLife.66231

Kagerer FA, Contreras-Vidal JL, Stelmach GE. 1997. Adaptation to gradual as compared with sudden visuo-motor distortions. *Exp Brain Res* **115**:557–561. doi:10.1007/PL00005727

Kambic RE, Roemmich RT, Bastian AJ. 2023. Joint-level coordination patterns for split-belt walking across different speed ratios. *J Neurophysiol* **129**:969–983. doi:10.1152/JN.00323.2021

Kandel E, Schwartz JH, Jessell TM, Siegelbaum SA, Hudspeth AJ. 2013. Principles of Neural Science, Fifth Edit. ed. McGraw-Hill, New York.

Kaufman L, Rousseeuw PJ. 1990. Finding Groups in Data: An Introduction to Cluster Analysis, Wiley series in probabilities and statistics, Wiley Series in Probability and Statistics. Wiley. doi:10.1002/9780470316801

Krakauer JW, Shadmehr R. 2006. Consolidation of motor memory. *Trends Neurosci* **29**:58–64. doi:10.1016/j.tins.2005.10.003

Lai NY, Bell JM, Bodznick D. 2021. Multiple behavior-specific cancellation signals contribute to suppressing predictable sensory reafference in a cerebellum-like structure. *J Exp Biol* **224**:240143. doi:10.1242/jeb.240143

Lauzière S, Miéville C, Betschart M, Duclos C, Aissaoui R, Nadeau S. 2014. Plantarflexion moment is a contributor to step length after-effect following walking on a split-belt treadmill in individuals with stroke and healthy individuals. *J Rehabil Med* **46**:849–857. doi:10.2340/16501977-1845

Lazar G, Van Laer J. 1968. Adaptation to displaced vision after experience with lesser displacements. *Percept Mot Skills* **26**:579–582. doi:10.2466/pms.1968.26.2.579

Lee JY, Schweighofer N. 2009. Dual adaptation supports a parallel architecture of motor memory. *J Neurosci* **29**:10396–10404. doi:10.1523/JNEUROSCI.1294-09.2009

Lee K, Oh Y, Izawa J, Schweighofer N. 2018. Sensory prediction errors, not performance errors, update memories in visuomotor adaptation. *Sci Rep* **8**:1–9. doi:10.1038/s41598-018-34598-y

Leech, Day KA, Roemmich RT, Bastian AJ. 2018a. Movement and perception recalibrate differently across multiple days of locomotor learning. *J Neurophysiol* **120**:2130–2137. doi:10.1152/jn.00355.2018

Leech, Roemmich RT, Bastian AJ. 2018b. Creating flexible motor memories in human walking. *Sci Rep* **8**:94. doi:10.1038/s41598-017-18538-w

Lindeman S, Hong S, Kros L, Mejias JF, Romano V, Oostenveld R, Negrello M, Bosman LWJ, de Zeeuw CI. 2020. Cerebellar purkinje cells can differentially modulate coherence between sensory and motor cortex depending on region and behavior. *Proc Natl Acad Sci U S A* **118**:2015292118. doi:10.1073/pnas.2015292118

Long AW, Roemmich RT, Bastian AJ. 2016. Blocking trial-by-trial error correction does not interfere with motor learning in human walking. *J Neurophysiol* **115**:2341–2348. doi:10.1152/jn.00941.2015

Malone LA, Bastian AJ. 2010. Thinking about walking: Effects of conscious correction versus distraction on locomotor adaptation. *J Neurophysiol* **103**:1954–1962. doi:10.1152/jn.00832.2009

Malone LA, Vasudevan EVL, Bastian AJ. 2011. Motor adaptation training for faster relearning. *J Neurosci* **31**:15136–15143. doi:10.1523/JNEUROSCI.1367-11.2011

Martin TA, Keating JG, Goodkin HP, Bastian AJ, Thach WT. 1996a. Throwing while looking through prisms II. Specificity and storage of multiple gaze-throw calibrations. *Brain* **119**:1199–1211. doi:10.1093/brain/119.4.1183

Martin TA, Keating JG, Goodkin HP, Bastian AJ, Thach WT. 1996b. Throwing while looking through prisms I. Focal olivocerebellar lesions impair adaptation. *Brain* **119**:1183–1198. doi:10.1093/brain/119.4.1183

McDougle SD, Bond KM, Taylor JA. 2015. Explicit and implicit processes constitute the fast and slow processes of sensorimotor learning. *J Neurosci* **35**:9568–9579. doi:10.1523/JNEUROSCI.5061-14.2015

McDougle SD, Ivry RB, Taylor JA. 2016. Taking Aim at the Cognitive Side of Learning in Sensorimotor Adaptation Tasks. *Trends Cogn Sci* **20**:535–544. doi:10.1016/j.tics.2016.05.002

McDougle SD, Taylor JA. 2019. Dissociable cognitive strategies for sensorimotor learning. *Nat Commun* **10**:1–11. doi:10.1038/s41467-018-07941-0

Miller EK, Cohen JD. 2001. An integrative theory of prefrontal cortex function. *Annu Rev Neurosci* **24**:167–202. doi:10.1146/annurev.neuro.24.1.167

Modchalingam S, Vachon CM, 't Hart BM, Henriques DYP. 2019. The effects of awareness of the perturbation during motor adaptation on hand localization. *PLoS One* **14**:e0220884. doi:10.17605/OSF.IO/MX5U2

Moidell BG, Bedell HE. 1988. Changes in oculocentric visual direction induced by the recalibration of saccades. *Vision Res* **28**:329–336. doi:10.1016/0042-6989(88)90161-7

Morton SM, Bastian AJ. 2006. Cerebellar contributions to locomotor adaptations during splitbelt treadmill walking. *J Neurosci* **26**:9107–9116. doi:10.1523/JNEUROSCI.2622-06.2006

Naik Gaonkar M, Sawant K. 2013. AutoEpsDBSCAN : DBSCAN with Eps Automatic for Large Dataset. *Int J Adv Comput Theory Eng* **2**:2319–2526.

Neville K-MM, Cressman EK. 2018. The influence of awareness on explicit and implicit contributions to visuomotor adaptation over time. *Exp Brain Res* **236**:2047–2059. doi:10.1007/s00221-018-5282-7

Ogawa T, Kawashima N, Ogata T, Nakazawa K. 2014. Predictive control of ankle stiffness at heel contact is a key element of locomotor adaptation during split-belt treadmill walking in humans. *J Neurophysiol* **111**:722–732. doi:10.1152/jn.00497.2012

Paul SS, Ada L, Canning CG. 2005. Automaticity of walking–implications for physiotherapy practice. *Phys Ther Rev* **10**:15–23. doi:10.1179/108331905X43463

Petersen TH, Willerslev-Olsen M, Conway BA, Nielsen JB. 2012. The motor cortex drives the muscles during walking in human subjects. *J Physiol* **590**:2443–2452. doi:10.1113/jphysiol.2012.227397

Pisano TJ, Dhanerawala ZM, Kislin M, Bakshinskaya D, Engel EA, Hansen EJ, Hoag AT, Lee J, de Oude NL, Venkataraju KU, Verpeut JL, Hoebeek FE, Richardson BD, Boele HJ, Wang SSH. 2021. Homologous organization of cerebellar pathways to sensory, motor, and associative forebrain. *Cell Rep* **36**:109721. doi:10.1016/j.celrep.2021.109721

Poggio T, Bizzi E. 2004. Generalization in vision and motor control. *Nature* **431**:768–774. doi:10.1038/nature03014

Popa D, Spolidoro M, Proville RD, Guyon N, Belliveau L, Léna C. 2013. Functional role of the cerebellum in gamma-band synchronization of the sensory and motor cortices. *J Neurosci* **33**:6552–6556. doi:10.1523/JNEUROSCI.5521-12.2013

Rahmah N, Sitanggang IS. 2016. Determination of Optimal Epsilon (Eps) Value on DBSCAN Algorithm to Clustering Data on Peatland Hotspots in SumatraIOP Conference Series: Earth and Environmental Science. Institute of Physics Publishing. p. 012012. doi:10.1088/1755-1315/31/1/012012

Reisman DS, Bastian AJ, Morton SM. 2010. Neurophysiologic and rehabilitation insights from the split-belt and other locomotor adaptation paradigms. *Phys Ther* **90**:187–195. doi:10.2522/ptj.20090073

Reisman DS, Block HJ, Bastian AJ. 2005. Interlimb coordination during locomotion: what can be adapted

1634  and stored? *J Neurophysiol* **94**:2403–2415. doi:10.1152/jn.00089.2005

1635 Reisman DS, McLean H, Keller J, Danks KA, Bastian AJ. 2013. Repeated split-belt treadmill training
1636  improves poststroke step length asymmetry. *Neurorehabil Neural Repair* **27**:460–468.
1637  doi:10.1177/1545968312474118

1638 Reisman DS, Wityk R, Silver K, Bastian AJ. 2007. Locomotor adaptation on a split-belt treadmill can
1639  improve walking symmetry post-stroke. *Brain* **130**:1861–1872. doi:10.1093/brain/awm035

1640 Roemmich RT, Bastian AJ. 2015. Two ways to save a newly learned motor pattern. *J Neurophysiol*
1641  **113**:3519–3530. doi:10.1152/jn.00965.2014

1642 Roemmich RT, Long AW, Bastian AJ, Roemmich RT, Long AW, Bastian AJ. 2016. Seeing the Errors You
1643  Feel Enhances Locomotor Performance but Not Learning. *Curr Biol* **26**:1–10.
1644  doi:10.1016/j.cub.2016.08.012

1645 Rondi-Reig L, Paradis AL, Lefort JM, Babayan BM, Tobin C. 2014. How the cerebellum may monitor
1646  sensory information for spatial representation. *Front Syst Neurosci* **8**:205.
1647  doi:10.3389/fnsys.2014.00205

1648 Rossi C, Bastian AJ, Therrien AS. 2021a. Mechanisms of proprioceptive realignment in human motor
1649  learning. *Curr Opin Physiol* **20**:186–197. doi:10.1016/j.cophys.2021.01.011

1650 Rossi C, Chau CW, Leech KA, Statton MA, Gonzalez AJ, Bastian AJ. 2019. The capacity to learn new
1651  motor and perceptual calibrations develops concurrently in childhood. *Sci Rep* **9**:9322.
1652  doi:10.1038/s41598-019-45074-6

1653 Rossi C, Roemmich RT, Schweighofer N, Bastian AJ, Leech KA. 2021b. Younger and Late Middle-Aged
1654  Adults Exhibit Different Patterns of Cognitive-Motor Interference During Locomotor Adaptation, With
1655  No Disruption of Savings. *Front Aging Neurosci* **13**:789. doi:10.3389/fnagi.2021.729284

1656 Rousseeuw PJ. 1987. Silhouettes: A graphical aid to the interpretation and validation of cluster analysis. *J*
1657  *Comput Appl Math* **20**:53–65. doi:10.1016/0377-0427(87)90125-7

1658 Saijo N, Gomi H. 2010. Multiple motor learning strategies in visuomotor rotation. *PLoS One* **5**:e9399.
1659  doi:10.1371/journal.pone.0009399

1660 Sánchez N, Park S, Finley JM. 2017. Evidence of Energetic Optimization during Adaptation Differs for
1661  Metabolic, Mechanical, and Perceptual Estimates of Energetic Cost OPEN. *Sci Rep* **7**:7682.
1662  doi:10.1038/s41598-017-08147-y

1663 Sánchez N, Simha SN, Donelan JM, Finley JM. 2021. Using asymmetry to your advantage: Learning to
1664  acquire and accept external assistance during prolonged split-belt walking. *J Neurophysiol* **125**:344–
1665  357. doi:10.1152/jn.00416.2020

1666 Sánchez N, Simha SN, Donelan JM, Finley JM. 2019. Taking advantage of external mechanical work to
1667  reduce metabolic cost: the mechanics and energetics of split-belt treadmill walking. *J Physiol*
1668  **597**:4053–4068. doi:10.1113/JP277725

1669 Schmahmann JD. 1996. From movement to thought: Anatomic substrates of the cerebellar contribution to
1670  cognitive processing. *Hum Brain Mapp* **4**:174–198. doi:10.1002/(SICI)1097-
1671  0193(1996)4:3<174::AID-HBM3>3.0.CO;2-0

1672 Seki K, Fetz EE. 2012. Gating of sensory input at spinal and cortical levels during preparation and
1673  execution of voluntary movement. *J Neurosci* **32**:890–902. doi:10.1523/JNEUROSCI.4958-11.2012

1674 Shadmehr R, Brashers-Krug T. 1997. Functional stages in the formation of human long-term motor
1675  memory. *J Neurosci* **17**:409–419. doi:10.1523/jneurosci.17-01-00409.1997

1676 Shadmehr R, Krakauer JW. 2008. A computational neuroanatomy for motor control. *Exp Brain Res*
1677  **185**:359–381. doi:10.1007/S00221-008-1280-5/FIGURES/9

1678 Smith MA, Ghazizadeh A, Shadmehr R. 2006. Interacting adaptive processes with different timescales
1679  underlie short-term motor learning. *PLoS Biol* **4**:e179. doi:10.1371/journal.pbio.0040179

1680 Sombric C, Gonzalez-Rubio M, Torres-Oviedo G. 2019. Split-Belt walking induces changes in active, but
1681  not passive, perception of step length. *Sci Rep* **9**:16442. doi:10.1038/s41598-019-52860-9

1682 Sombric CJ, Harker HM, Sparto PJ, Torres-Oviedo G. 2017. Explicit action switching interferes with the
1683  context-specificity of motor memories in older adults. *Front Aging Neurosci* **9**:40.
1684  doi:10.3389/fnagi.2017.00040

1685 Song CN, Stenum J, Leech KA, Keller CK, Roemmich RT. 2020. Unilateral step training can drive faster
1686  learning of novel gait patterns. *Sci Reports 2020 101* **10**:1–13. doi:10.1038/s41598-020-75839-3

1687 Statton MA, Vazquez A, Morton SM, Vasudevan EVL, Bastian AJ. 2018. Making sense of cerebellar

contributions to perceptual and motor adaptation. *Cerebellum* **17**:111–121. doi:10.1007/s12311-017-0879-0

Sultan F, Augath M, Hamodeh S, Murayama Y, Oeltermann A, Rauch A, Thier P. 2012. Unravelling cerebellar pathways with high temporal precision targeting motor and extensive sensory and parietal networks. *Nat Commun* **3**:1–10. doi:10.1038/ncomms1912

Sun X, O'Shea DJ, Golub MD, Trautmann EM, Vyas S, Ryu SI, Shenoy K V. 2022. Cortical preparatory activity indexes learned motor memories. *Nature* **602**:274–279. doi:10.1038/s41586-021-04329-x

Synofzik M, Lindner A, Thier P. 2008. The cerebellum updates predictions about the visual consequences of one's behavior. *Curr Biol* **18**:814–818. doi:10.1016/J.CUB.2008.04.071

Takakusaki K. 2013. Neurophysiology of gait: From the spinal cord to the frontal lobe. *Mov Disord* **28**:1483–1491. doi:10.1002/MDS.25669

Tanaka H, Ishikawa T, Lee J, Kakei S. 2020. The Cerebro-Cerebellum as a Locus of Forward Model: A Review. *Front Syst Neurosci* **14**:19. doi:10.3389/FNSYS.2020.00019/BIBTEX

Tanaka H, Krakauer JW, Sejnowski TJ. 2012. Generalization and multirate models of motor adaptation. *Neural Comput* **24**:939–966. doi:10.1162/NECO_a_00262

Taylor JA, Ivry RB. 2011. Flexible cognitive strategies during motor learning. *PLoS Comput Biol* **7**:e1001096. doi:10.1371/journal.pcbi.1001096

Taylor JA, Krakauer JW, Ivry RB. 2014. Explicit and implicit contributions to learning in a sensorimotor adaptation task. *J Neurosci* **34**:3023–3032. doi:10.1523/JNEUROSCI.3619-13.2014

Thompson AK, Xiang YC, Wolpaw JR. 2009. Acquisition of a simple motor skill: Task-dependent adaptation plus long-term change in the human soleus H-reflex. *J Neurosci* **29**:5784–5792. doi:10.1523/JNEUROSCI.4326-08.2009

Todorov E. 2005. Stochastic Optimal Control and Estimation Methods Adapted to the Noise Characteristics of the Sensorimotor System. *Neural Comput* **17**:1084–108.

Todorov E. 2004. Optimality principles in sensorimotor control. *Nat Neurosci*. doi:10.1038/nn1309

Todorov E, Jordan MI. 2002. Optimal feedback control as a theory of motor coordination. *Nat Neurosci* **5**:1226–1235. doi:10.1038/nn963

Tsay JS, Kim H, Haith AM, Ivry RB. 2022. Understanding implicit sensorimotor adaptation as a process of proprioceptive re-alignment. *Elife* **11**:e76639. doi:10.7554/eLife.76639

Tsay JS, Kim HE, McDougle SD, Taylor JA, Haith A, Avraham G, Krakauer JW, Collins AG, Ivry RB. 2024. Fundamental processes in sensorimotor learning: Reasoning, refinement, and retrieval. *Elife* **12**:e91839. doi:10.7554/eLife.91839

Tsay JS, Kim HE, Parvin DE, Stover AR, Ivry RB. 2021. Individual differences in proprioception predict the extent of implicit sensorimotor adaptation. *J Neurophysiol* **125**:1307–1321. doi:10.1152/jn.00585.2020

Tseng YW, Diedrichsen J, Krakauer JW, Shadmehr R, Bastian AJ. 2007. Sensory prediction errors drive cerebellum-dependent adaptation of reaching. *J Neurophysiol* **98**:54–62. doi:10.1152/jn.00266.2007

Uiga L, Capio CM, Ryu D, Young WR, Wilson MR, Wong TWL, Tse ACY, Masters RSW. 2020. The Role of Movement-Specific Reinvestment in Visuomotor Control of Walking by Older Adults. *Journals Gerontol - Ser B Psychol Sci Soc Sci* **75**:282–292. doi:10.1093/geronb/gby078

Van Vugt FT, Ostry DJ. 2018. The structure and acquisition of sensorimotor maps. *J Cogn Neurosci* **30**:290–306. doi:10.1162/jocn_a_01204

Vasudevan EVL, Torres-Oviedo G, Morton SM, Yang JF, Bastian AJ. 2011. Younger is not always better: development of locomotor adaptation from childhood to adulthood. *J Neurosci* **31**:3055–3065. doi:10.1523/JNEUROSCI.5781-10.2011

Vazquez A, Statton MA, Busgang SA, Bastian AJ. 2015. Split-belt walking adaptation recalibrates sensorimotor estimates of leg speed but not position or force. *J Neurophysiol* **114**:3255–3267. doi:10.1152/jn.00302.2015

Vervoort D, Rob Den Otter A, Buurke TJW, Vuillerme N, Hortobágyi T, Lamoth CJC. 2019. Effects of aging and task prioritization on split-belt gait adaptation. *Front Aging Neurosci* **11**:10. doi:10.3389/fnagi.2019.00010

Weeks HM, Therrien AS, Bastian AJ. 2017. The cerebellum contributes to proprioception during motion. *J Neurophysiol* **118**:693–702. doi:10.1152/jn.00417.2016

Welniarz Q, Worbe Y, Gallea C. 2021. The Forward Model: A Unifying Theory for the Role of the

Cerebellum in Motor Control and Sense of Agency. *Front Syst Neurosci* **15**:22.
doi:10.3389/FNSYS.2021.644059/BIBTEX

Wolpert DM, Ghahramani Z, Flanagan JR. 2001. Perspectives and problems in motor learning. *Trends*
*Cogn Sci*. doi:10.1016/S1364-6613(00)01773-3

Wong WL, Masters RSW, Maxwell JP, Abernethy AB. 2008. Reinvestment and falls in community-
dwelling older adults. *Neurorehabil Neural Repair* **22**:410–414. doi:10.1177/1545968307313510

Yavari F, Mahdavi S, Towhidkhah F, Ahmadi-Pajouh MA, Ekhtiari H, Darainy M. 2016. Cerebellum as a
forward but not inverse model in visuomotor adaptation task: a tDCS-based and modeling study.
*Exp Brain Res* **234**:997–1012. doi:10.1007/s00221-015-4523-2

Zhang Z, Wang H, Zhang T, Nie Z, Wei K. 2024. Perceptual error based on Bayesian cue combination
drives implicit motor adaptation. *Elife* **13**:RP94608. doi:10.7554/ELIFE.94608

1753
1754

**Figure Legends**

**Figure 1. Experiment 1, Hypotheses and Predictions. (A)** Conceptual schematic of our paradigm with the Ramp Down task: after adaptation, the right belt speed is gradually ramped down to match the left. **(B-C)** Predictions for the Ramp Down motor measures made by two competing hypotheses. **(B) Recalibration only:** recalibration can only change movement gradually. The Δ motor output (dotted blue line) changes slowly and does not track the rapidly decreasing perturbation (dashed red line), so that step length asymmetry aftereffects emerge immediately (solid purple line, magnitude is positive). **(C) Recalibration + mapping:** mapping can change movement immediately. In the first part of the task (highlighted in green), the mapping contribution to Δ motor output (dark blue shade) is scaled down immediately as the perturbation decreases. Hence, the Δ motor output (dotted blue line) changes rapidly and tracks the perturbation (dashed red line), so that there are no step length asymmetry aftereffects (solid purple line, magnitude is ~zero). In the second part of the task, the mapping contribution to Δ motor output is zero, and the recalibration contribution to Δ motor output (light blue shade) does not change significantly. Hence, the Δ motor output (dotted blue line) does not track the perturbation (red dashed line), and step length asymmetry aftereffects emerge (solid purple line, magnitude is positive). **Right column inset:** conceptual explanation of how both hypotheses may account for the speed match results from Leech at al. (Leech et al., 2018a). In the first post-adaptation speed match task, participants increase the speed of the right belt from zero to a value that is smaller than adaptation but larger than the left belt (top panel). The perturbation increases until a value that is positive but smaller than adaptation (dashed red line, middle and bottom panels). Leech et al. observed symmetric step lengths at the end of the task, indicating that the Δ motor output (dotted blue line) is smaller than it was in adaptation and matches the perturbation. The decrease in Δ motor output can be explained by the recalibration only hypothesis as forgetting/unlearning (middle panel), or by the recalibration + mapping hypothesis as flexible scaling of the mapping contribution (bottom panel).

**Figure 2. Experiment 1, Step length asymmetry. (A)** Top: Experimental protocol. The Ramp Down task (purple) is used to test the predictions illustrated in Fig. 1. Bottom: Step length asymmetry time course. Background shading darkness increases with belt speed difference (color bar). Phases (except Ramp tasks) are truncated to the participant with fewest strides. **(B)** Zoomed-in baseline ramp and post-adaptation Ramp Down tasks. Speed differences for which step length asymmetry is not significantly different from zero are indicated by the green shade. Inset depicts predictions made by the competing hypotheses as in Fig. 1. All curves show group mean ± SE.

**Figure 3. Experiment 1, Perturbation and Δ motor output. (A)** Perturbation (red) and Δ motor output (blue) data for the Ramp Down task. **(B-C)** Perturbation data (red) and model fit for the Δ motor output (blue) for the recalibration + mapping model and three recalibration only models. Timeseries curves show group mean ± SE, and green shade corresponds to speeds with symmetric step lengths as in Fig. 2. **Bar insets in (C):** BIC difference between recalibration + mapping and each recalibration only model (bar ± error bar shows group mean ± CI, circles show individual participants' data).

**Figure 4. Experiment 1, Perceptual Results. (A)** Top: perturbation data (red) and recalibration + mapping fit (blue); this is the same as Fig. 3B. Bottom: perceptual task button presses (green, group mean ± SE of button press stride depicted as a function of belt speed difference). Right: measures of motor recalibration ("$r$") and total motor adaptation ("$u_{\text{plateau}}$"). **(B)** Perturbation

1804     compensation (normalized perceptual and motor measures of adaptation): $compensation_{\text{perceptual}}$
1805     bounds (green), $compensation_{\text{motor total}}$ (dark blue), and $compensation_{\text{motor recalibration}}$ (light blue).
1806     **(C-D)** Individual participants' $compensation_{\text{motor recalibration}}$ versus $compensation_{\text{perceptual}}$ (first or
1807     second button press). Solid black: least squares line. Dashed gray: unity line.
1808
**Figure 5. Experiment 1, Perceptual Models. (A-C)** Perturbation data (red) and model fits for the
Δ motor output (dark blue) for the proprioceptive re-alignment model (PReMo), Perceptual Error
Adaptation model (PEA), and perceptuomotor recalibration + mapping model (PM-ReMap). **Dark**
**blue bar insets:** BIC difference between recalibration + mapping and each perceptual model (bar
± error bar shows group mean ± CI, circles show individual participants' data). **Light blue arrows:**
model predictions for the perception of belt speed difference. **Bottom bar in (C):** group mean ±
SE of the stride at which belts are predicted to feel equal, with $\widehat{PSE}$ reflecting the perturbation at
this stride. **(D)** Individual participants' $\widehat{PSE}$ versus $compensation_{\text{perceptual}}$ (upper or lower bound).
Solid black: least squares line. Dashed gray: unity line.
1818
**Figure 6. Control experiments, protocols. (A)** General protocol for all Control experiments (tied
belts = at the same speed, split belts = right faster than left). **(B)** Sample participants performing
the ascend and descend versions of the speed match task (first iteration after adaptation).
Participants respectively increased or decreased the speed of the right belt (dashed orange) using
up/down buttons with the goal of matching it to the reference speed of the left belt (solid black).
Vision and sound were occluded as shown. The PSE is the belt speed difference at the end of
the task (green). **(C)** Adaptation belt speeds and post-adaptation speed match task version for all
Control experiments. All groups except for Small Gradual experienced a catch-trial (tied belts)
after two-thirds of the adaptation phase.
1828
**Figure 7. Control experiments, validation of Experiment 1. (A)** $compensation_{\text{motor total}}$ (filled
bars) and $compensation_{\text{perceptual}}$ (open bars) at the end of adaptation for all groups (group mean
± SE). **(B)** Step length asymmetry as a function of belt speed difference during the first post-
adaptation speed match task (interpolated, group mean ± SE). Error bars depict step length
asymmetry during the first (depicted to the right or left) and last (depicted in inset) strides in the
task (actual data, group mean ± CI). Asterisks represent significant differences.
1835
**Figure 8. Control experiments, effect of paradigm manipulation and repetition on relative**
**recalibration contribution. (A)** Comparison of the relative contribution of recalibration to motor
adaptation across groups that vary in adaptation duration (left panel, Short, Medium, and Long
Ascend), perturbation magnitude (middle panel, Medium Descend and Small Abrupt), or schedule
(right panel, Small Abrupt and Gradual). Bars depict the ratio $\dfrac{compensation_{\text{perceptual}}}{compensation_{\text{motor total}}}$ (group mean ±
SE)**;** asterisks represent significant differences. **(B)** Post-adaptation time course of motor (solid
lines) and perceptual (dashed lines) aftereffects for Short, Medium, and Long Ascend groups
(group mean ± SE). Aftereffects are computed as the tied-belt step length asymmetry (motor) or
the final speed difference (perceptual) in each post-adaptation speed match task, normalized to
the adaptation perturbation. Gray shades indicate time points for which motor aftereffects are
significantly smaller than perceptual aftereffects.
1847
1848     **Figure 9. Experiment 2, Step length asymmetry. (A)** Experimental protocol, equivalent to that
1849     of Experiment 1 except for the Ramp Up & Down part shaded in teal, where the right speed was
1850     faster than in adaptation (ramped up to 2m/s and back down to 1.5m/s). **(B)** Step length
1851     asymmetry time course (entire group mean ± SE). Background shade represents belt speed

difference. Phases (except ramp tasks) are truncated to the participant with fewest strides. **Inset:** Predictions for the step length asymmetry during the teal portion of the Ramp Up & Down task, for the memory-based (top) or structure-based (bottom) mapping hypotheses. **(C)** Zoomed-in Ramp Up & Down task (entire group mean ± SE). Step length asymmetry for strides taken at right speeds larger than adaptation is shown in teal. **(D-E)** Separate plots of the step length asymmetry in the Ramp Up & Down task for the subgroups of participants that walked asymmetrically (D, "memory-based") versus symmetrically (E, "structure-based") in the teal portion of the task (subgroups mean ± SE). **Insets:** circles represent individual participants' number of strides, in the teal portion of the task, with step length asymmetry below their own baseline CI. Error bars depict subgroup mean ± SE. Subgroup assignment was performed by clustering on this measure.

**Figure 10.** Summary of self-reported deliberate changes to the walking pattern in adaptation. Only one participant accurately described changes to the walking pattern that related to adaptation, while other responses were negative (i.e. no deliberate changes, 3 participants), irrelevant (7 participants), or inaccurate (5 participants).

**Figure 11. Schematic model of adaptation.** Body movement depends on environment perturbations (red) and Δ motor output (blue). The Δ motor output is adjusted by recalibration (light blue) and mapping (dark blue) mechanisms, which perform different operations and are arranged in tandem. We propose the following architecture and flow: **(1. Recalibration)** The recalibration mechanism produces adjustment $x_r$ that is fixed regardless of perturbation size (light blue box, $x_r$ is constant for varying $p$). The *same* recalibration adjustment $x_r$ serves as an input to *both* areas responsible for conscious perception (green box) and Δ motor output (blue box). **(2. Perception)** Conscious perception is computed by cancelling out the recalibration adjustment from the actual sensory feedback (green box, perception of the belt speed difference perturbation $\tilde{p}$ is the difference between the actual speed difference $p$ and recalibration $x_r$). The perceived perturbation $\tilde{p}$ serves as an input to the mapping mechanism (dark blue box). **(3. Mapping)** The mapping mechanism produces adjustment $x_m$ that can vary in magnitude to appropriately account for the perceived perturbation $\tilde{p}$ (dark blue box, $x_m$ scales with $\tilde{p}$ and matches its magnitude). **(4. Δ Motor output)** The overall adjustment to Δ motor output is computed by adding the mapping adjustment $x_m$ and recalibration adjustment $x_r$. The corner in the Δ motor output versus perturbation profile arises because mapping is computed based on the perceived perturbation $\tilde{p}$ (not the actual perturbation $p$) and is only learnt for positive $\tilde{p}$ (the experienced direction). When the perturbation is perceived to be opposite than adaptation, even if it is not, mapping is zero and the Δ motor output is constant, reflecting recalibration adjustments only (blue box, when $\tilde{p} < 0$ and $p \geq 0$ the mapping adjustment $x_m$ is zero and $u = x_r$).

**Box 1 - Figure 1. Standard paradigm and measures. (A)** Treadmill belt speeds for the standard split-belt paradigm. **(B)** Schematic time course of standard motor measures of walking adaptation: step length asymmetry – a measure of error (solid purple), Δ motor output – a measure of compensatory spatial and temporal asymmetries (dotted blue), and perturbation – the effect of the speed asymmetry on the walking pattern (dashed red). In baseline, the belts are tied, and perturbation, Δ motor output, and step length asymmetry are all ~0. In adaptation, the right leg is faster than the left such that the perturbation is positive. The Δ motor output is still ~0 in early adaptation, causing step length asymmetry errors (negative purple line). By late adaptation the Δ motor output is adapted to match the perturbation, and step length asymmetry returns to ~0.

1897 Changes to Δ motor output persist in tied-belts post-adaptation, but the perturbation is ~0, causing
1898 step length asymmetry aftereffects (positive purple line).

1899

1900 **Box 2 - Figure 1. Relevant results from Leech et al., 2018. (A)** Treadmill belt speeds and step
1901 length asymmetry time course, similar to that described in Box 1 – Figure 1. Vertical dashed gray
1902 lines indicate iterations of the speed match task, where participants adjust the speed of the right
1903 belt with a keypad to match it to the left. **(B)** Time courses of the belt speeds (top; orange = right,
1904 black = left) and step length asymmetry (bottom) in selected iterations of the speed match task.
1905 Left, "before adapt": last baseline task. Right, "after adapt": first post-adaptation task. Dotted
1906 horizontal lines depict the right speed (orange, top) and step length asymmetry magnitude (purple,
1907 bottom) at adaptation plateau (average over the last 30 strides). **(C)** Belt speed difference (top)
1908 and step length asymmetry (bottom) magnitudes at the end of the tasks shown in B. All curves
1909 show group mean ± SE, and all data is collected in the Leech at al. study (Leech et al., 2018a).

1910

1911

**A**

baseline | adaptation | Ramp Down | speed match

right

speeds

— left    - - - right

**B**

recalibration only

motor measures

recalibration

forgetting

aftereffects

forgetting in break

forgetting in task

symmetry

**C**

recalibration + mapping

motor measures

mapping

recalibration

no aftereffects    aftereffects

stride

mapping scaled down

symmetry

—— step length asym    ····· Δ motor output    - - - perturbation

**A** **Ramp Down data**

**B** **recalibration + mapping model**

motor measures

stride

right - left speed [m/s]

— Δmotor output
— perturbation

**C** **recalibration only models**

dual state

optimal control

memory of errors

right - left [m/s]

ΔBIC

recal+map better

**A** baseline | adaptation | catch (10s) | post-adaptation

1 | 1 | 1 | 1 [minutes] | 0.5 x2 | 1.5 | 3.5 | 7.5

□ tied belts | ▨ split belts | ■ speed match (30 sec)

**B** Ascend | Descend

speed [m/s]

1.5 | 1 | 0.5

30 sec | 30 sec | PSE

reference

— left | - - right

**C** group | adaptation | speed match

| group | adaptation | speed match |
|---|---|---|
| Short Ascend | 1.5m/s 0.5m/s | ↑ |
| Medium Ascend | | ↑ |
| Long Ascend | | ↑ |
| Short Descend | | ↓ |
| Medium Descend | | ↓ |
| Small Abrupt | 0.8m/s 0.4m/s | ↓ |
| Small Gradual | | ↓ |

— left | - - right

0 3 15 30 [minutes] | ↑ascend ↓descend

A

perturbation compensation [%]

Short Ascend | Medium Ascend | Long Ascend | Short Descend | Medium Descend | Small Abrupt | Small Gradual

■ motor total    □ perceptual    *[N=10 per group]*

B

Descend ← → Ascend

SLA

step length asymmetry

Δ speed [m/s]

right faster | left faster