1 Resilience metrics are robust across data qualities but sensitive to

2 community size.

4 Abstract

1. Modern biodiversity monitoring is generating increasingly multidimensional representations of wildlife populations and ecosystems. It is therefore appealing for conservation and environmental governance to combine that information into single measure of ecosystem or population health.

2. The Jacobian matrix is a common characteristic used to identify the sensitivity of simulated/mathematical systems to perturbation (a.k.a. its resilience) and predict its near future dynamics. Jacobians have therefore been suggested as a theoretically grounded measure of ecosystem resilience. Whilst historically it has been challenging to estimate the Jacobian from empirical data, recent work has proposed a suite of metrics capable of reconstructing it for real-world community using multi-species time series data.

3. Here we assess the robustness of five resilience metrics influenced by varying time series lengths and data qualities based on that seen in real-world wildlife time series. We generate data using multispecies Lotka–Volterra equations and simulate stressed and unstressed communities of varying species number. These data were then corrupted through the introduction of sampling error (to mimic varying search efforts) and truncating time series (to match the typical time series lengths reported in global biodiversity datasets such as the Living Planet Index and BIOTIME).

4. The robustness of all resilience metrics improved with time series length, whilst the amount of sampling error had little effect on their performance. However, community

25   size (number of species) dramatically altered metric capability, with larger

26   communities decreasing the reliability of resilience metric trends.

27  5. Overall, resilience metrics behave predictably across realistic data corruptions.

28   Generic resilience estimation is therefore possible from abundance time series alone,

29   and we suggest that, given the increasing availability of multivariate community data,

30   focussing on Jacobian estimates for resilience is a promising avenue of research.

31   However, we also show it is prudent to apply ecological knowledge when selecting

32   which species to contribute.

34 <u>Introduction</u>

35 Biodiversity loss is a major concern for the maintenance of ecosystem stability and

36 functioning (Cardinale et al., 2012; Oliver et al., 2015). Despite this, species extinctions are

37 accelerating (Ceballos et al., 2015) with resulting change in ecosystem services, exploitable

38 resources, and habitat cover being sufficiently impactful (Traill et al., 2011; Tylianakis et al.,

39 2008) that biodiversity policy targets have become commonplace (United Nations, 2020).

40 The conservation and political decisions behind the selection of these targets is dependent on

41 sufficiently high quality data and reliable indicators of biodiversity change (Dale & Beyeler,

42 2001). Such biodiversity indicators are ultimately used to quantify both the scale of impacts

43 and monitor mitigation actions – i.e. have the desired outcomes been achieved (Donatti et al.,

44 2020) – in an "indicator-policy cycle" (Nicholson et al., 2012) to ensure appropriate policy

45 decisions during a period of competing priorities for decision makers.

47 The Living Planet Index (LPI - Loh et al., 2005) has emerged alongside BIOTIME (Dornelas

48 et al., 2018) PREDICTS (Hudson et al., 2017) and the IUCN Red list (IUCN, 2022) as the

49 focal datasets and descriptive instruments for global biodiversity trends. They have

influenced international (United Nations, 2020, 1992), national (Environment and Climate

Change Canada, 2019), and regional (Brotons et al., 2020) environment policy, as well as

scientific discourse (Dornelas et al., 2019; Gonzalez et al., 2016; Johnson et al., 2022; Leung

et al., 2020; Loreau et al., 2022; McRae et al., 2017). It is therefore undeniable the

importance of these datasets for current and future biodiversity monitoring especially given

the relative paucity of long-term and reliable counts of species globally (Hughes et al., 2017;

McRae et al., 2017). Consequently, the available time series have driven a primary focus on

abundance changes by biodiversity indicators in absolute or relative terms (i.e. species

richness or diversity indices), despite the acknowledged sampling errors, geographical biases,

and many short time present (Gonzalez et al., 2016; Leung et al., 2020). These concerns

generate uncertain estimates of global biodiversity change (Johnson et al., 2022; Wauchope et

al., 2019) which hampers the indicator-policy cycle.

Such a focus on abundance also potentially overlooks other signals expressed by stressed

populations (Cerini et al., 2023) which are likely mechanistically important for abundance

decline and can provide earlier and more robust indications of biodiversity loss.

Unfortunately, for many practitioners, abundance data is all that is available, leading to

alternative suggestions to consider the resilience of the system (Capdevila et al., 2022;

Holling, 1973) rather than focussing on raw abundance trends. This is achieved by

reconciling the interacting nodes/species into a singular measure of population/ecosystem

health (Dakos et al., 2015; Ushio et al., 2018) – an indicator of system resilience dependent

on the response of the system rather than solely its species diversity. Here we define

resilience following Hodgson et al. (2015) as the 'capacity of a system to persist and maintain

its state and functions in the face of exogenous disturbance/stress'. Resilience therefore

consists of two major components: the ability to resist and recover from perturbations (Pimm,

1984). However, measuring ecosystem resilience (and its associated stability) from empirical

data is challenging (Capdevila et al., 2022; Donohue et al., 2013, 2016) with an ever

expanding diversity of possible metrics and indices proposed (Kéfi et al., 2019). Kéfi et al.

(2019) specifically highlight that resilience indicators are rarely considered in parallel, across

neither unidimensional (i.e. populations) nor multidimensional (i.e. communities) scales.

Empirical resilience research should therefore attempt to reconcile such differences in scale

and dimensionality. In addition, Kéfi et al. (2019) also suggest that any resilience metric must

be rooted in theory and be consistent across empirical and theoretical studies. For example,

the dominant eigenvalue of the Jacobian is common in theoretical studies due its accurate

representation of future system dynamics (Dakos, 2018; Strogatz, 2015; Yang et al., 2019),

but has rarely been reported in real-world studies (Kéfi et al., 2019) due to the challenge of

defining empirical Jacobians.

Computational advancements in the last decade have provided a novel toolbox for estimating

Jacobians and species interaction matrices from observational data (Ahmad et al., 2016;

Sugihara et al., 2012; Williamson & Lenton, 2015). These 'resilience metrics/indicators' are

grounded in dynamical system theory (Strogatz, 2015) and have the potential to quantify

resilience in the absence of true equilibrium (Lyapunov, 1992). This is obviously attractive

for biodiversity monitoring as loss of resilience/stability is indicative of oncoming decline

(Dakos et al., 2015) even if the system is away from equilibrium (which is likely the case for

ecosystems experiencing multiple stressors (Hastings et al., 2018)). However, it is unclear

how resilience metrics behave across data qualities and ecosystem community structure to

warrant their immediate usage in real-world data. Similar ambiguities have influenced the

practicality of related population collapse indicators (a.k.a. Early Warning Signals – EWSs)

with good performance reported in idealised simulated studies (Clements et al., 2015; Dakos,

2018) but failures in observational time series (Burthe et al., 2016; Zhang, 2020). There is

consequently a need to test the behaviour of the resilience metric toolbox in controlled

scenarios matching the data qualities present in biodiversity datasets such as the LPI, and

ensure their behaviour is sufficiently reliable prior to their potential usage in the indicator-

policy cycle.

In this work, we use simulated multispecies ecological communities to test the influence of

time series length and data corruption upon five resilience metrics. These resilience metrics

included a multivariate index of variability (Brock & Carpenter, 2006), Fisher information

(Ahmad et al., 2016; Karunanithi et al., 2008), and three Jacobian matrix estimators

(Grziwotz et al., 2023; Ushio et al., 2018; Williamson & Lenton, 2015). We specifically

estimated the trend of each metric in 5, 15 and 25 species communities across different scales

(populations vs communities) and show that while metric reliability generally increases with

data quality, metrics displayed weak but correct trends even in low data qualities. However,

the choice of species contributing to the metric dramatically influenced reliability leading to

recommendations that users must guide their species selection with ecological knowledge.

Methods

Simulation framework

Community dynamics were simulated using an established multi-species generalised Lotka-

Volterra competitive model (Hofbauer & Sigmund, 1998), containing a control parameter that

can drive one or more species to extinction. Specifically, this model simulates the appearance

of an invasive species (Equation 1 - (Dakos, 2018)) in an established community and is

written as:

$$\frac{\mathrm{d}N_i}{\mathrm{d}t} = r_i N_i(t) \left[1 - \frac{\sum_{j=1}^{n} \alpha_{ij} N_j(t)}{f_i(t)} + \mu_i \right] + \varepsilon_i(t)$$

$$f_i(t) = K_i(1 + \eta_i C)$$

Equation 1

where i and j are the focal and interacting species identities respectively. N_i is the resulting

population density of species i, r_i is i's intrinsic growth rate, α_{ij} is the interaction coefficient

between species i and j (or the per capita effect of species j on i), f_i is the species-specific

harvest rate function with C the control parameter, and ε is a white noise process with mean 0

and variance 0.1. Symbols K_i, μ_i, and η_i represent the initial carrying capacity, immigration

term, and species-specific response to the control parameter C for species i respectively.

The elements of the interaction matrix α_{ij} were parameterised followed Yang et al (2019). To

mimic the asymmetric species interactions of empirical food-webs, if a predator predates a

single prey species, the negative per-capita impact on the prey was set to -0.5. Conversely, in

the circumstances where multiple species are predated, one prey was randomly allocated a

strong -0.4 impact and the remaining prey set to -0.1 divided by the number of possible prey

species. The resulting positive impact on the predator was assumed dependent on its

conversion efficiency. In the absence of omnivory, that conversion efficiency was set to 0.2

while omnivorous predators were assumed to have a lower efficiency of 0.02. These

conversion efficiencies were then multiplied by the negative per-capita impact its prey

experiences to generate the positive benefit to the predator species. Interspecific competition

was assumed to be relatively strong between basal species, with interactions drawn from a

uniform distribution between 0 and -0.5, while intraspecific competition was assumed to be

stronger than any interspecific competition for simplicity (Chesson 2000). Consequently, the

diagonal of the matrix was set to -1 for basal species and -0.1 for higher trophic levels.

Interspecific competition between non-basal species was considered indirect due to their

inherent exploitation of a shared resource pool (Petchey *et al.* 2008; Yang *et al.* 2019).

A fixed trophic structure was imposed on simulated communities consisting of four trophic

levels in a chain/ladder food-web motif. This matrix was however flexible to allow varying

community sizes, with communities of 5, 15 and 25 species explored in this work. Species

were allocated to trophic levels following a 'pyramid' of species richness (Turney & Buddle,

2016) with lower trophic levels being more specious than higher trophic levels (Figure 1a).

The remaining parameters were populated following Dakos (2018). Growth rate r_i, carrying

capacity K_i, and sensitivity to the control parameter η_i were sampled from the uniform

distributions $0.9 – 1.1$, $5 – 15$, and $0 – 1$ respectively. The immigration rate μ_i was set as 0.01

for all species.

Figure 1. a) Exemplary trophic structure and intraspecific interactions for a five species community. Numbers represent species, solid paths are species interactions and arrows are intraspecific effects. b) A simulated community across stressed and unstressed conditions. The estimated inflection point is indicated by a vertical dashed line. c) Hypothesised behaviour of resilience metrics through time (trend) across time series length and search effort. All metrics (excluding Fisher information) are expected to increase through time in increasingly stressed communities and remain constant in unstressed. Increasing time series length and search effort is anticipated to improve the reliability and strength of relationship. Fisher information is expected to display negative trends. A dashed zero trend line is provided for reference.

Stable communities were then identified by estimating the community densities at

equilibrium (N^*) before calculating the Jacobian matrix J using the equation $J = \alpha_{ij}N^*$ (May

1974) and its eigenvalues. When the dominant real eigenvalue of J is positive or close to

zero, the system is locally unstable and consequently we rejected any community with a

dominant eigenvalue greater than -0.05. We then randomly selected 30 stable communities

with unique interaction matrices.

Each community was stochastically simulated using the Euler-Maruyama method at dt = 0.1,

for 100 timesteps with a control parameter value of 0. In the following period between t100

and t200, the control parameter value was incrementally increased and then held at the

maximum value of 5.0 (Dakos, 2018) from t200 to t300. For simplicity, only one species was

targeted as an invasive species as such species typically appear individually (Doherty et al.,

2016), though multiple can successively invade (O'Loughlin & Green, 2017). The species in

the third trophic level with the highest intrinsic growth rate r was selected for invasion as

invasive species success is primarily dependent on its dispersal traits (Carboni et al., 2016;

Jeschke & Strayer, 2006) with mesopredators the most pervasive invaders (Doherty et al.,

2016). Qualitatively, changing this target species does not alter the overall dynamics of

system other than the time of transition, which is mitigated here by the truncating of each

time series prior to the point of fastest community change (see *Data post-processing*). Each

community was simulated 25 times repeated across stressed and unstressed regimes resulting

in 1500 timeseries per community size (2 stress regimes, 30 communities, 25 stochastic

runs). Simulations were run using the '*DifferentialEquations*' package (Rackauckas & Nie

2017) in the Julia language (Bezanson *et al.* 2017).

Data post-processing

Each simulated time series was then altered to mimic the imperfect sampling and short time

series found in the LPI. First, we identified the time point of fastest change in the

community's first principal component using the bisection extremum surface estimator

method (Christopoulos, 2016) provided by the *inflection* R package (v.1.3.6). This ensures

that the community is stressed but no species has been yet driven to collapse (Figure 1b), and

allows us to identify practical resilience metric behaviour over timescales appropriate for

intervention (Davidson et al., 2018; Hastings, 2016). Consequently, each time series was

truncated to 10, 25, 40, 55 and 70 timepoints prior to the estimated inflection point, matching

the range of time series lengths contributing to the LPI. Each truncated time series was then

subjected to simulated searches ranging from 10 to 100% effort in 10% increments. To

achieve this, we converted the simulated continuous species biomass data to counts, to

represent natural animal populations, by multiplying the simulated data by 100 and rounding

to the nearest integer. We then drew 'observed' species counts from a binomial distribution

with probability equal to the search effort (Clements et al., 2015), which mimics both

imperfect spatial sampling and detection. Resultantly, each initial simulation was reanalysed

for each time series length and search effort combination. All post processing, metric

calculation and analysis was performed in the R language v.4.3.0 (R Core Team, 2022).

Resilience metrics

Five resilience metrics were calculated for each post-processed simulation using the R

package EWSmethods (v.1.2.0) (O'Brien et al., 2023). These metrics included Fisher

information (Karunanithi et al., 2008), a multivariate index of variability (Brock & Carpenter,

2006), two S-map estimated Jacobian indices (Grziwotz et al., 2023; Ushio et al., 2018) and a

Jacobian index estimated from multivariate autoregressive models (Williamson & Lenton,

2015). All metrics were calculated in a rolling window 50% the total time series length

following the typical practice for generic resilience indicators (Dakos et al., 2012; Lenton et

al., 2012).

Fisher information (FI) quantifies indeterminacy or the quantity of information observed data

contain on an underlying but unmeasured parameter (Ahmad et al., 2016). In practice FI is

calculated by comparing changes in the probability of detecting different states relative to

small inherent fluctuations in the time series. Typically, these fluctuations are defined by the

variance of each time series in a reference period (Karunanithi et al., 2008). FI is therefore

calculated in a rolling window along the time series, with the variability of each window

compared to that of the previous. If the variability is greater than that of the reference

variability, then the windows are binned into separate states. Here we defined this reference

variability as the standard deviation of each species across the entire time series (Karunanithi

et al., 2008). FI is consequently expected to decrease along with stability.

The multivariate index of variability (mvi) is the square root of the species covariance

matrix's dominant eigenvalue and represents a multivariate equivalent to the univariate

"variance" early warning signal (Brock & Carpenter, 2006). It is therefore expected to

increase as resilience/stability decreases.

The two S-map Jacobian indices are derived from a state space reconstructed via empirical

dynamic modelling (Park et al., 2021; Sugihara et al., 2012). In brief, both indices estimate

the Jacobian of the community by comparing the time delayed relationships between each

representative time series and predicting out from the reconstructed state-space (Ushio et al.,

2018). The coefficients of the local linear model performing this out-prediction therefore

represent the estimated α_{ij} of the community. All species data were scaled to mean zero and

unit variance prior to metric estimation, following Chang et al (2017), to ensure all species

are of equivalent magnitude to maximise the accuracy of the S-map estimated interaction

strengths.

The multivariate form of this Jacobian index (multiJI) compares all pair wise interactions

between species in the community and constructs the state space by delay embedding up to E,

where E is the number of species. The metric is therefore sensitive to the choice and number

of variables (i.e species here) contributing to the S-map calculation (Medeiros & Saavedra,

2023; Ushio et al., 2018), and limits the minimum number of time points to E. In this study,

as all species are contributing to system stability, all species are appropriate for inclusion.

However, if a species count is constant for the duration of a window, then S-map

reconstruction is impossible and therefore we excluded any species which were unchanging

for the duration of the time series. This is possible in our simulations when a species either

goes extinct prior to the shortened time series assessed, or low search efforts did not

encounter species with small populations. Similarly, as time series length was often less than

E in the specious communities (e.g. 10 timepoints but $E = 15$), for these communities we

randomly selected species to contribute to multiJI to test this additional processing error on

the capability of multiJI to accurately characterise resilience change. multiJI is expected to

increase as resilience decreases with a value greater than 1 indicating instability, whereas a

value below 1 indicates stability.

The univariate form of Jacobian index is calculated for each species independently by delay

embedding the time series against lagged versions of itself (Grziwotz et al., 2023). The

resulting α_{ij}/local linear model coefficients are therefore all 0, excluding the lower-off

diagonal (all -1), whereas the first row elements become the coefficients for the focal species.

Consequently, the system's behaviour can be approximated in the absence of complete

information and escapes concerns regarding the inclusion of variables/species (Grziwotz et

al., 2023; Medeiros et al., 2022; Ushio et al., 2018). Here we set the embedding dimension E

to 1 (1 "year") and the time lag τ to 10% the length of the time series following the coded

examples of Grziwotz *et al*. (2023). To allow comparability with the multivariate resilience

metrics, we then averaged uniJI estimates across all species (mean_uniJI) and extracted the

maximum index value observed (max_uniJI), with both metrics also expected to increase in

parallel with stressor value similar to mvi and multiJI.

Finally, the multivariate autoregressive Jacobian index (multiAR) fits autoregressive models

between multiple time series and uses the generated autocorrelation matrix's eigenvalues to

estimate the system's Jacobian dominant eigenvalue (Williamson & Lenton, 2015). The

eigenvalues of the two matrices' real components are related by the equation:

$$\lambda_j = \frac{1}{\Delta t} \log |a_j|$$

where λ_j is the Jacobian eigenvalues, a_j is the eigenvalues of the autocorrelation matrix and Δt

is the time step of the data generating process. As Δt is typically unknown for empirical data,

we have assumed it is equal to 1. multiAR is therefore conceptually similar to multiJI but

exploits lag-1 autoregressive models (Hasselmann, 1988) rather than empirical dynamical

modelling (Sugihara et al., 2012). Due to this similarity, species selection and data processing

was identical to multiJI, with the same expected behaviour; an increase in value with stress.

Resilience metric trends

All five metrics are expected to express directional change when the system is linearly

forced/stressed but remain approximately constant when unstressed (Figure 1c). We therefore

assessed the linear trend of each metric in a Bayesian mixed effects model as an alternative to

the Kendall tau correlation comparisons typically applied to such indicator comparisons

(Baruah et al., 2022; Clements et al., 2015; Dutta et al., 2018). Comparing resilience metric

ability in this way allows us to extract representative global trends while accounting for

autocorrelation and shared features between simulations (Johnson et al., 2022). Each metric's

trend across time series lengths and search efforts was consequently analysed independently

using Laplace approximation in the INLA framework (v.23.04.24) (Rue et al., 2009) due to

the quantity of data. Interactions were modelled between time (numeric, scaled between 0

and 1 to allow comparability across truncated time series lengths), whether the simulation

represents a stressed or unstressed model (factor, two levels), time series length (numeric,

divided by 10 to be of equal magnitude as search effort) and search effort (numeric) as fixed

effects. Random intercepts were also included for community identifier (factor, 30 levels)

along with an autocorrelation function to account for serial dependence within a simulation,

and random slopes for each community as each is expected to display the same trend.

Separate models were fit for each community size (5, 15, or 25 species) due to computational

constraints. Please note that mvi is modelled on the log scale due to sudden non-linearities

known to occur in this metric's trend (Brock & Carpenter, 2006). The resulting model

structure was:

$$y_{ij} \sim Normal(\mu_{ij})$$

$$\mu_{ij} = \mathrm{X}\beta + \mathrm{Z}\gamma_{ijk}$$

$$\gamma_{ijk} = \rho\epsilon_{ijk,t-1} + b_{0j} + b_{1j}time_{ijk} + \epsilon_{ikt}$$

where y_{ij} is the predicted metric value for the ith data point in the jth community and kth

simulation with mean μ_{ij}. β represents the coefficients for time, stress, time series length,

search effort and their interactions, b_{0j} the random intercepts, b_{1j} the random slope

coefficients, ρ the correlation between temporally related residuals (ε), and t the time point of

i. X and Z are design matrices for fixed and random effect variables respectively.

We set the weakly informative priors following Rue et al. (2009)'s suggestions:

$$\beta \sim Normal(0,1)$$

$$b_{0j} \sim Normal\left(0, \sigma_j\right)$$

$$b_{1j} \sim Normal\left(0, \sigma_j\right)$$

$$\sigma_j \sim lognormal(0,1)$$

Jacobian threshold transgression

As the Jacobian indices can also signal unstable periods when greater then 1 (for the S-map

derived methods) or 0 (for the autoregressive method), we also modelled the capability of

these metrics to correctly classify stressed simulations using a binomial mixed effects model.

As above, we included time series length, search effort and model stress as fixed effects, with

random intercepts for community identifier, but also included community size (numeric) as a

fixed effect with the response variable whether the threshold of 1/0 was crossed at any point

in the time series, rather than the raw metric value. The model structure was therefore:

$$y_{ij} \sim Binomial\left(\mu_{ij}\right)$$

$$\mu_{ij} = X\beta + Z\gamma_{ij}$$

$$\gamma_{ij} = b_{0j} + \epsilon_{ij}$$

with priors maintained as above.

<u>Results</u>

Metric trends

All resilience metrics display their anticipated behaviour through time and are influenced by

both the length of time series and quality of search effort (Figures 2-4). In stressed 5 species

communities (Figure 2, Tables S1-S5), increasing time series length improves the strength of

trend estimate in all metrics for a given search effort level. However, increasing search effort

alters trend differentially between metrics. For example, multiJI's and multiAR's trend is

weakened with improved search efforts while the remainder of metrics all strengthen.

Absolute strength of trend was generally weak in each metric, other than FI, owing to the

influence of stochasticity in each simulation altering the timing of metric change between

simulations and non-linearities in metric trend (Figures S1-S3).

Similarly, the conservative hierarchical and autoregressive modelling approach balloons

uncertainty as identified in other work modelling global trends from multiple local trends

(Johnson et al., 2022). In the unstressed communities, the S-map Jacobian based metrics

behave as expected across time series and search efforts (approximately 0 trend).

Figure 2. Posterior estimated slopes/trends of each resilience metric (rows) through time, across truncated time series lengths (columns) and search efforts in 5 species communities. Dark coloured estimates represent trends in unstressed communities, while light coloured estimates are those in stressed communities. The reported values are the posterior density median values (circles), with 80% (thickest bars) and 95% (thinnest bars) credible intervals.

In the 15 species communities (Figure 3, Tables S6-S10), all metrics other than multiJI and

multiAR behaved as anticipated across stressed vs unstressed communities, time series

lengths and search efforts: each metric's trend was approximately zero in the unstressed

communities but strengthened with time series length and search effort in the stressed. Both

multiJI and multiAR were unreliable in shorter, unstressed communities though became more

reliable as time series length increased. multiJI behaviour in stressed communities was stable

across time series length and search efforts and was primarily compromised by unexpectedly

positive trends in unstressed communities with shorter time series.

Figure 3. Posterior estimated slopes/trends of each resilience metric (rows) through time, across truncated time series lengths (columns) and search efforts in 15 species communities. Dark coloured estimates represent trends in unstressed communities, while light coloured estimates are those in stressed communities. The reported values are the posterior density median values (circles), with 80% (thickest bars) and 95% (thinnest bars) credible intervals.

Finally, all metrics perform less reliably in 25 species communities (Figure 4, Tables S11-

S15) relative to the smaller communities. multiJI and multiAR are unable to distinguish

stressed from unstressed communities regardless of time series length or search effort while

mean_uniJI only distinguishes stressed from unstressed in the longest and best searched

communities. FI and mvi both perform as anticipated across increasing time series lengths

and search efforts but with weaker trends than those observed in the smaller communities.

Figure 4. Posterior estimated slopes/trends of each resilience metric (rows) through time, across truncated time series lengths (columns) and search efforts in 25 species communities. Dark coloured estimates represent trends in unstressed communities, while light coloured estimates are those in stressed communities. The reported values are the posterior density median values (circles), with 80% (thickest bars) and 95% (thinnest bars) credible intervals.

Threshold transgression

For the Jacobian estimating metrics multiJI, max_uniJI and multiAR, the size of the

community was the best predictor of reported instability probability (Figure 5, Tables S16-

S18) – i.e. displaying a metric value greater than its instability threshold. For multiJI and

max_uniJI that threshold was 1 and 0 for multiAR. In multiJI and multiAR, for a constant

time series length and search effort, the 25 species community always displayed a higher

probability of reported instability than 15 or 5 (Figure 5a,c) while the opposite relationship

was observed for max_uniJI (Figure 5b); as community size decreases, the more likely

instability was reported.

Figure 5. Posterior estimated probabilities of the resilience metrics multiJI, max_uniJI and multiAR transgressing its threshold a. Estimates have been delineated by truncated time series lengths (columns) and search efforts (rows) in the three different community sizes. Dark coloured estimates represent trends in unstressed communities, while light coloured estimates are those in stressed communities. The reported values are the posterior density median values (circles), with 80% (thickest bars) and 95% (thinnest bars) credible intervals back transformed from log odds to probabilities.

Search effort did not influence the probability of instability report in stressed communities

but increased that probability in unstressed communities. Time series length generally

decreased probability of reported instability but synergised with search effort so that high

search effort resulted in equivalent probabilities across time series length. Conversely, for

unstressed communities, low search effort increased the probability of reported instability.

If the three metrics are compared, multiJI and multiAR displayed identical behaviour in

longer time series, implying both estimate similar Jacobians. Overall, multiJI displayed the

highest true positive rate (highest probabilities in stressed communities) whereas max_uniJI

displayed the lowest false positive rate (lowest probabilities in unstressed communities).

max_uniJI generally balanced the ratio better than multiJI and multiAR but is more sensitive

to low search efforts. The metric was also most robust at the highest data qualities (long time

series and high search efforts) across community sizes, but more conservative than the other

metrics at low data qualities.

<u>Discussion</u>

Quantifying population and community stability/resilience loss has emerged as a crucial

target of ecological research (Kéfi et al., 2019), mapping directly on to the needs of

practitioners and policy makers (Donohue et al., 2016; Nicholson et al., 2012). Despite the

complexity of ecological systems, political decisions often require simplifications to allow

efficient decision making and action (Donatti et al., 2020; Nicholson et al., 2012). Resilience

metrics represent a bridge between the need for indicators of stability loss and simple user

interpretation. We show here that resilience metrics can perform as anticipated in simulated

data analogous to real world biodiversity datasets though the choice of metric and

contributing time series requires caution.

Influence of data quality and community size

Our simulations suggest that each resilience metric is robust to low search efforts, but that

time series length has a significant impact on their ability to reconstruct the Jacobian matrix.

This is anticipated behaviour (Figure 1) though the multivariate Jacobian metrics (multiJI and

multiAR) unexpectedly decreased in reliability as search effort increased. It is unclear why

these metrics would behave in this way, though it is possible that, as multiJI and multiAR

displayed the highest estimated autocorrelation across metrics (autocorrelation $\approx$ 1; Tables

S1, S5, S6, S10, S11 and S15), the error introduced by low search effort acted as a low pass

filter which weakened the influence of autocorrelation and allowed the true interactions

between species to be identified. Similarly, our assumption that search effort linearly

influences metric ability may also be violated by the strong non-linearities we observe in the

raw indicator trends (Figures S1-S3).

The increasing trend in the metrics with time series length follows the typical behaviour of

biodiversity indicators (Clements et al., 2015; Gonzalez et al., 2016; Lenton et al., 2012),

with longer time series displaying more reliable estimates of the true trend (Wauchope et al.,

2019). This effect is compounded by the caveats of Sugihara et al. (2012) that a minimum of

30 time points are required for reliable empirical dynamic modelling estimates. This is

obviously unfeasible for many yearly abundance accounts reported in the LPI and other

datasets, and while we show that correct estimates can be made with these short time series,

their accuracy is weaker than in time series longer than 30 time points.

It is encouraging, however, that unstressed communities consistently showed no trend in any

resilience metric (Figures 2-4) and rarely signalled instability for metrics with a threshold

(Figure 5). This implies that if a trend is observed, then resilience/stability is decreasing prior

to abundance decline, and intervention is likely required. Note that no mechanism or driver of

stability loss is provided by resilience metrics and only phenomenological signals are

returned. These indicators consequently require system specific knowledge for successful

intervention, but this is a recurring challenge for current abundance-based biodiversity

indicators such as the LPI (Ledger et al., 2023).

Overall, the dominant determinant of resilience metric performance was the number of

species in the simulated community. The multivariate metrics estimating the communities'

Jacobian (multiJI and multiAR) were particularly influenced, with weakening performance as

community size increased. Ushio et al. (2018) acknowledge the potential for this degradation

and suggest it is dependent on the choice of species included in the Jacobian estimation. We

show here that, even when all species contributing to the system are known and measured,

including all species compromises the indices' ability. We therefore support Ushio et al.

(2018) that contributing species must be carefully chosen either based upon their trophic and

functional importance to prevent the masking of true system resilience. The univariate

Jacobian metric (mean_uniJI) also surprisingly outperformed the multivariate (multiJI and

multiAR), though it has been suggested that representative variables can provide sufficient

dynamics without masking the signal with other uninformative time series (Grziwotz et al.,

2023). This does however partially conflict with Donoghue et al. (2013, 2016) and Kéfi et al.

(2019) who clearly highlight that multidimensional measurements of resilience are required,

with a focus away from indicator species. To clarify, we agree that biodiversity and resilience

indicators need to be multidimensional, but caution against naïve species selection.

Appropriate species may be selected based upon their dominance in the ecosystem in terms

of abundance (Ushio et al., 2018), or inter-species interaction strengths (Medeiros et al.,

2022; Medeiros & Saavedra, 2023; White et al., 2020).

 A shift towards multivariate data

A general restriction of the LPI specifically is its primarily single species focus (Ledger et al.,

2023; Loh et al., 2005). BIOTIME (Dornelas et al., 2018) and PREDICTS (Hudson et al.,

2017) have greater assemblage contributions, though the latter is almost exclusively single

snapshots in time due to the project's focus on spatial questions. That said, it is clear that

biodiversity monitoring requires multidimensional representation going forward (Donohue et

al., 2013; Kéfi et al., 2019; Medeiros et al., 2022; Medeiros & Saavedra, 2023; White et al.,

2020) to provide the reliable estimates required for effective governance. Such data will

become more readily available with the advent of autonomous and remote ecological

monitoring (Besson et al., 2022), but there will be a need for tools capable of synthesising

that data in to an interpretable form for decision makers. We suggest resilience metrics can

contribute to this synthesis in extension of traditional approaches such as species diversity or

richness (Hudson et al., 2017; Turney & Buddle, 2016) due to their development specifically

focussing on species interactions. We go further, to suggest that multivariate resilience

metrics also have potential to characterise the 'trait' state-space for focal populations such a

target fisheries or forestry. Using trait measurements across plots or through time, resilience

of a population/assemblage in terms of its traits is possible which allows ecosystem managers

to ask questions regarding 'timelines of collapse' (Cerini et al., 2023) or population intactness

(Hudson et al., 2017) for the individual/population/community of management interest.

However, if multivariate data is not available, this study supports Grziwotz et al (2023)'s

findings that univariate Jacobian estimates can also be reliable indicators of system dynamics

in the absence of complete monitoring.

Comparison to critical transition indicators

A similar collection of indicators to the resilience metrics discussed here have received

extensive research interest in relation to predicting certain forms of regime shift and

population collapse (Baruah et al., 2022; Clements et al., 2015; Dakos, 2018; Dakos et al.,

2012). These 'Early Warning Signals' (EWSs) can be considered a sub-class of resilience

indicator (Dakos et al., 2015) as they quantify resilience when a system has two alternative

stable states and the capability to transition/regime shift between them. With current debates

over the commonality of alternative stable states in ecology (Dakos et al., 2015; Davidson et

al., 2023) and inconsistent ability in real world data (Burthe et al., 2016; Zhang, 2020), the

benefit of resilience metrics over EWSs is their ability to quantify stability regardless of the

presence of multiple stable states (Lyapunov, 1992; Medeiros et al., 2022; Sugihara et al.,

2012). With the complex intrinsic and extrinsic interactions driving ecosystem dynamics

possibly masking true multiple stability (Hillebrand et al., 2023) or ecosystems themselves

experiencing long transients rather than regime shifts (Hastings et al., 2018), the resilience

metrics discussed here are conceptually capable of accurate Jacobian estimation under both

heuristics. Such capability therefore expedites the general suggestions to consider non-

equilibrium resilience and stability change in ecological research (Donohue et al., 2013,

2016; Kéfi et al., 2019).

Trends vs thresholds

A key challenge for designing indicators appropriate for policy is that governance primarily

focusses on thresholds (Donohue et al., 2016; United Nations, 2020) as they allow

unambiguous interpretation and enforcement. Conversely, trends require sufficient

subjectivity that the vague language associated with trend reporting results in broad objective

language (United Nations, 2020). Many governments therefore use indicators with set target

thresholds for biodiversity rather than considering a continuous gradient of 'state' which is

suggested to be more appropriate for modern coupled socio-ecological systems (Hillebrand et

al., 2023).

This requirement for a threshold is also explored by the LPI. The debates surrounding the LPI

and its interpretation (Dornelas et al., 2019; Leung et al., 2020) ultimately revolve around

reconciling the variability of local trend directions (winner and loser populations) with the

short time series typically available (Gonzalez et al., 2016). Indeed, controlling for co-

dependencies between separate time series such as temporal, spatial and phylogenetic

autocorrelation increases the uncertainty of trend estimates (Johnson et al., 2022). This

indicator/monitoring uncertainty erodes legislators' ability to make confident governance,

interacts with other forms of uncertainty (e.g. political strategies, lobbying effort, pre-existing

laws (Dewulf & Biesbroek, 2018)) and leads to broad objectives. The Jacobian estimating

metrics we explored minimise this incongruency between trends and thresholds by

facilitating both forms of interpretation – an increase in trend is indicative of decreasing

resilience, while exceeding of the threshold value represents detection of instability.

Meanwhile, FI and mvi do not express thresholds but their superior ability relative to

Jacobian indices in this study implies trends are more reliable monitoring instruments for

ecosystem stability than instability thresholds.

Indeed, instability is not necessarily atypical of natural systems. Lake fish communities can

display consistent periods of instability though the year (Ushio et al., 2018), Canadian boreal

forests are temporarily unstable following historic fires (Héon et al., 2014), and experimental

intertidal communities recover (i.e. display transient instability) following single pulse stress

(White et al., 2020). Resilience metric thresholds are therefore only informative if instability

is prolonged.

Conclusion

Our study demonstrates resilience metrics can accurately quantify resilience of simulated

multispecies communities manipulated into realistic data quality. Contributing species/time

series is the key consideration for these metrics' functionality and requires ecological

knowledge input from the user. Resilience is strongly suggested to be under characterised

during global biodiversity monitoring (Capdevila et al., 2022; Kéfi et al., 2019) and the

resilience metrics tested here have the potential to contribute generic inference from

abundance data alone. We consequently anticipate Jacobian estimation techniques to increase

in popularity but pre-empt their use in biodiversity datasets by cautioning towards the longest

time series possible and ecologically important species.

References

Ahmad, N., Derrible, S., Eason, T., & Cabezas, H. (2016). Using Fisher Information to Track

    Stability in Multivariate Systems. *Royal Society Open Science*, *3*(11), 160582.

    https://doi.org/10.1098/rsos.160582

Baruah, G., Ozgul, A., & Clements, C. F. (2022). Community Structure Determines the

    Predictability of Population Collapse. *Journal of Animal Ecology*, *n/a*(n/a).

    https://doi.org/10.1111/1365-2656.13769

Besson, M., Alison, J., Bjerge, K., Gorochowski, T. E., Høye, T. T., Jucker, T., Mann, H. M.

    R., & Clements, C. F. (2022). Towards the fully automated monitoring of ecological

    communities. *Ecology Letters*, *25*(12), 2753–2775. https://doi.org/10.1111/ele.14123

Brock, W., & Carpenter, S. R. (2006). Variance as a Leading Indicator of Regime Shift in

    Ecosystem Services. *Ecology and Society*, *11*, 9.

Brotons, L., Pou, N., Sainz de la Maza, P., & Pont, S. (2020). *Estat de la Natura a Catalunya*

2020. Departament de Territori i Sostenibilitat, Generalitat de Catalunya.

Burthe, S. J., Henrys, P. A., Mackay, E. B., Spears, B. M., Campbell, R., Carvalho, L.,

Dudley, B., Gunn, I. D. M., Johns, D. G., Maberly, S. C., May, L., Newell, M. A.,

Wanless, S., Winfield, I. J., Thackeray, S. J., & Daunt, F. (2016). Do early warning

indicators consistently predict nonlinear change in long-term ecological data? *Journal*

of Applied Ecology, *53*(3), 666–676. https://doi.org/10.1111/1365-2664.12519

Capdevila, P., Noviello, N., McRae, L., Freeman, R., & Clements, C. F. (2022). Global

patterns of resilience decline in vertebrate populations. *Ecology Letters*, *25*(1), 240–

251. https://doi.org/10.1111/ele.13927

Carboni, M., Münkemüller, T., Lavergne, S., Choler, P., Borgy, B., Violle, C., Essl, F.,

Roquet, C., Munoz, F., DivGrass Consortium, & Thuiller, W. (2016). What it takes to

invade grassland ecosystems: Traits, introduction history and filtering processes.

Ecology Letters, *19*(3), 219–229. https://doi.org/10.1111/ele.12556

Cardinale, B. J., Duffy, J. E., Gonzalez, A., Hooper, D. U., Perrings, C., Venail, P., Narwani,

A., Mace, G. M., Tilman, D., Wardle, D. A., Kinzig, A. P., Daily, G. C., Loreau, M.,

Grace, J. B., Larigauderie, A., Srivastava, D. S., & Naeem, S. (2012). Biodiversity

loss and its impact on humanity. *Nature*, *486*(7401), 59–67.

https://doi.org/10.1038/nature11148

Ceballos, G., Ehrlich, P. R., Barnosky, A. D., García, A., Pringle, R. M., & Palmer, T. M.

(2015). Accelerated modern human–induced species losses: Entering the sixth mass

extinction. *Science Advances*, *1*(5), e1400253. https://doi.org/10.1126/sciadv.1400253

Cerini, F., Childs, D. Z., & Clements, C. F. (2023). A predictive timeline of wildlife

population collapse. *Nature Ecology & Evolution*, *7*(3), 320–331.

https://doi.org/10.1038/s41559-023-01985-2

Chang, C.-W., Ushio, M., & Hsieh, C. (2017). Empirical Dynamic Modeling for Beginners.

    Ecological Research, *32*(6), 785–796. https://doi.org/10.1007/s11284-017-1469-9

Christopoulos, D. (2016). On the Efficient Identification of an Inflection Point.

    INTERNATIONAL JOURNAL OF MATHEMATICS AND SCIENTIFIC COMPUTING

    (ISSN: 2231-5330), *6*.

Clements, C. F., Drake, J. M., Griffiths, J. I., & Ozgul, A. (2015). Factors Influencing the

    Detectability of Early Warning Signals of Population Collapse. *The American*

    Naturalist, *186*(1), 50–58. https://doi.org/10.1086/681573

Dakos, V. (2018). Identifying Best-Indicator Species for Abrupt Transitions in Multispecies

    Communities. *Ecological Indicators*, *94*, 494–502.

    https://doi.org/10.1016/j.ecolind.2017.10.024

Dakos, V., Carpenter, S. R., Brock, W. A., Ellison, A. M., Guttal, V., Ives, A. R., Kéfi, S.,

    Livina, V., Seekell, D. A., van Nes, E. H., & Scheffer, M. (2012). Methods for

    Detecting Early Warnings of Critical Transitions in Time Series Illustrated Using

    Simulated Ecological Data. *PLOS ONE*, *7*(7), e41010.

    https://doi.org/10.1371/journal.pone.0041010

Dakos, V., Carpenter, S. R., van Nes, E. H., & Scheffer, M. (2015). Resilience indicators:

    Prospects and limitations for early warnings of regime shifts. *Philosophical*

    Transactions of the Royal Society B: Biological Sciences, *370*(1659), 20130263.

    https://doi.org/10.1098/rstb.2013.0263

Dale, V. H., & Beyeler, S. C. (2001). Challenges in the development and use of ecological

    indicators. *Ecological Indicators*, *1*(1), 3–10. https://doi.org/10.1016/S1470-

    160X(01)00003-6

Davidson, T. A., Bennion, H., Reid, M., Sayer, C. D., & Whitmore, T. J. (2018). Towards

    better integration of ecology in palaeoecology: From proxies to indicators, from

594   inference to understanding. *Journal of Paleolimnology*, *60*(2), 109–116.

595   https://doi.org/10.1007/s10933-018-0032-1

596 Davidson, T. A., Sayer, C. D., Jeppesen, E., Søndergaard, M., Lauridsen, T. L., Johansson, L.

597   S., Baker, A., & Graeber, D. (2023). Bimodality and alternative equilibria do not help

598   explain long-term patterns in shallow lake chlorophyll-a. *Nature Communications*,

599   *14*(1), 398. https://doi.org/10.1038/s41467-023-36043-9

600 Dewulf, A., & Biesbroek, R. (2018). Nine lives of uncertainty in decision-making: Strategies

601   for dealing with uncertainty in environmental governance. *Policy and Society*, *37*(4),

602   441–458. https://doi.org/10.1080/14494035.2018.1504484

603 Doherty, T. S., Glen, A. S., Nimmo, D. G., Ritchie, E. G., & Dickman, C. R. (2016). Invasive

604   predators and global biodiversity loss. *Proceedings of the National Academy of

605   Sciences*, *113*(40), 11261–11265. https://doi.org/10.1073/pnas.1602480113

606 Donatti, C. I., Harvey, C. A., Hole, D., Panfil, S. N., & Schurman, H. (2020). Indicators to

607   measure the climate change adaptation outcomes of ecosystem-based adaptation.

608   *Climatic Change*, *158*(3), 413–433. https://doi.org/10.1007/s10584-019-02565-9

609 Donohue, I., Hillebrand, H., Montoya, J. M., Petchey, O. L., Pimm, S. L., Fowler, M. S.,

610   Healy, K., Jackson, A. L., Lurgi, M., McClean, D., O'Connor, N. E., O'Gorman, E. J.,

611   & Yang, Q. (2016). Navigating the complexity of ecological stability. *Ecology Letters*,

612   *19*(9), 1172–1185. https://doi.org/10.1111/ele.12648

613 Donohue, I., Petchey, O. L., Montoya, J. M., Jackson, A. L., McNally, L., Viana, M., Healy,

614   K., Lurgi, M., O'Connor, N. E., & Emmerson, M. C. (2013). On the dimensionality of

615   ecological stability. *Ecology Letters*, *16*(4), 421–429.

616   https://doi.org/10.1111/ele.12086

617 Dornelas, M., Antão, L. H., Moyes, F., Bates, A. E., Magurran, A. E., Adam, D.,

618   Akhmetzhanova, A. A., Appeltans, W., Arcos, J. M., Arnold, H., Ayyappan, N.,

Badihi, G., Baird, A. H., Barbosa, M., Barreto, T. E., Bässler, C., Bellgrove, A.,

Belmaker, J., Benedetti-Cecchi, L., … Zettler, M. L. (2018). BioTIME: A database of

biodiversity time series for the Anthropocene. *Global Ecology and Biogeography*,

27(7), 760–786. https://doi.org/10.1111/geb.12729

Dornelas, M., Gotelli, N. J., Shimadzu, H., Moyes, F., Magurran, A. E., & McGill, B. J.

(2019). A balance of winners and losers in the Anthropocene. *Ecology Letters*, *22*(5),

847–854. https://doi.org/10.1111/ele.13242

Dutta, P. S., Sharma, Y., & Abbott, K. C. (2018). Robustness of Early Warning Signals for

Catastrophic and Non-Catastrophic Transitions. *Oikos*, *127*(9), 1251–1263.

https://doi.org/10.1111/oik.05172

Environment and Climate Change Canada. (2019). *Canadian Environmental Sustainability*

Indicators: Canadian species index.

Gonzalez, A., Cardinale, B. J., Allington, G. R. H., Byrnes, J., Arthur Endsley, K., Brown, D.

G., Hooper, D. U., Isbell, F., O'Connor, M. I., & Loreau, M. (2016). Estimating local

biodiversity change: A critique of papers claiming no net loss of local diversity.

Ecology, *97*(8), 1949–1960. https://doi.org/10.1890/15-1759.1

Grziwotz, F., Chang, C.-W., Dakos, V., van Nes, E. H., Schwarzländer, M., Kamps, O.,

Heßler, M., Tokuda, I. T., Telschow, A., & Hsieh, C. (2023). Anticipating the

Occurrence and Type of Critical Transitions. *Science Advances*, *9*(1), eabq4558.

https://doi.org/10.1126/sciadv.abq4558

Hasselmann, K. (1988). PIPs and POPs: The reduction of complex dynamical systems using

principal interaction and oscillation patterns. *Journal of Geophysical Research:*

Atmospheres, *93*(D9), 11015–11021. https://doi.org/10.1029/JD093iD09p11015

Hastings, A. (2016). Timescales and the management of ecological systems. *Proceedings of*

the National Academy of Sciences, *113*(51), 14568.

https://doi.org/10.1073/pnas.1604974113

Hastings, A., Abbott, K. C., Cuddington, K., Francis, T., Gellner, G., Lai, Y.-C., Morozov, A.,

Petrovskii, S., Scranton, K., & Zeeman, M. L. (2018). Transient phenomena in

ecology. *Science*, *361*(6406), eaat6412. https://doi.org/10.1126/science.aat6412

Héon, J., Arseneault, D., & Parisien, M.-A. (2014). Resistance of the boreal forest to high

burn rates. *Proceedings of the National Academy of Sciences*, *111*(38), 13888–13893.

https://doi.org/10.1073/pnas.1409316111

Hillebrand, H., Kuczynski, L., Kunze, C., Rillo, M. C., & Dajka, J.-C. (2023). Thresholds and

tipping points are tempting but not necessarily suitable concepts to address

anthropogenic biodiversity change—An intervention. *Marine Biodiversity*, *53*(3), 43.

https://doi.org/10.1007/s12526-023-01342-3

Hodgson, D., McDonald, J. L., & Hosken, D. J. (2015). What do you mean, 'resilient'?

Trends in Ecology & Evolution, *30*(9), 503–506.

https://doi.org/10.1016/j.tree.2015.06.010

Hofbauer, J., & Sigmund, K. (1998). *Evolutionary Games and Population Dynamics*.

Cambridge University Press. https://doi.org/10.1017/CBO9781139173179

Holling, C. S. (1973). Resilience and stability of ecological systems. *Annual Review of*

Ecology and Systematics, *4*(1), 1–23.

https://doi.org/10.1146/annurev.es.04.110173.000245

Hudson, L. N., Newbold, T., Contu, S., Hill, S. L. L., Lysenko, I., De Palma, A., Phillips, H.

R. P., Alhusseini, T. I., Bedford, F. E., Bennett, D. J., Booth, H., Burton, V. J., Chng,

C. W. T., Choimes, A., Correia, D. L. P., Day, J., Echeverría-Londoño, S., Emerson, S.

R., Gao, D., … Purvis, A. (2017). The database of the PREDICTS (Projecting

Responses of Ecological Diversity In Changing Terrestrial Systems) project. *Ecology*

and Evolution, *7*(1), 145–188. https://doi.org/10.1002/ece3.2579

Hughes, B. B., Beas-Luna, R., Barner, A. K., Brewitt, K., Brumbaugh, D. R., Cerny-

Chipman, E. B., Close, S. L., Coblentz, K. E., de Nesnera, K. L., Drobnitch, S. T.,

Figurski, J. D., Focht, B., Friedman, M., Freiwald, J., Heady, K. K., Heady, W. N.,

Hettinger, A., Johnson, A., Karr, K. A., … Carr, M. H. (2017). Long-Term Studies

Contribute Disproportionately to Ecology and Policy. *BioScience*, *67*(3), 271–281.

https://doi.org/10.1093/biosci/biw185

IUCN. (2022). *The IUCN Red List of Threatened Species* (Version 2022-2) [dataset].

https://www.iucnredlist.org

Jeschke, J. M., & Strayer, D. L. (2006). Determinants of vertebrate invasion success in

Europe and North America. *Global Change Biology*, *12*(9), 1608–1619.

https://doi.org/10.1111/j.1365-2486.2006.01213.x

Johnson, T. F., Beckerman, A. P., Childs, D. Z., Griffiths, C. A., Capdevila, P., Clements, C.

F., Besson, M., Gregory, R. D., Delmas, E., Thomas, G., Evans, K., Webb, T., &

Freckleton, R. (2022). Overconfidence undermines global wildlife abundance trends.

BioRxiv, 2022.11.02.514877. https://doi.org/10.1101/2022.11.02.514877

Karunanithi, A. T., Cabezas, H., Frieden, B. R., & Pawlowski, C. W. (2008). Detection and

Assessment of Ecosystem Regime Shifts from Fisher Information. *Ecology and*

Society, *13*(1). http://www.jstor.org/stable/26267924

Kéfi, S., Domínguez-García, V., Donohue, I., Fontaine, C., Thébault, E., & Dakos, V. (2019).

Advancing our understanding of ecological stability. *Ecology Letters*, *22*(9), 1349–

1356. https://doi.org/10.1111/ele.13340

Ledger, S. E. H., Loh, J., Almond, R., Böhm, M., Clements, C. F., Currie, J., Deinet, S.,

Galewski, T., Grooten, M., Jenkins, M., Marconi, V., Painter, B., Scott-Gatty, K.,

692   Young, L., Hoffmann, M., Freeman, R., & McRae, L. (2023). Past, present, and future

693    of the Living Planet Index. *Npj Biodiversity*, *2*(1), 12. https://doi.org/10.1038/s44185-

694    023-00017-3

695 Lenton, T. M., Livina, V. N., Dakos, V., van Nes, E. H., & Scheffer, M. (2012). Early

696    Warning of Climate Tipping Points from Critical Slowing down: Comparing Methods

697    to Improve Robustness. *Philosophical Transactions. Series A, Mathematical,*

698    *Physical, and Engineering Sciences*, *370*(1962), 1185–1204.

699    https://doi.org/10.1098/rsta.2011.0304

700 Leung, B., Hargreaves, A. L., Greenberg, D. A., McGill, B., Dornelas, M., & Freeman, R.

701    (2020). Clustered versus catastrophic global vertebrate declines. *Nature*, *588*(7837),

702    267–271. https://doi.org/10.1038/s41586-020-2920-6

703 Loh, J., Green, R. E., Ricketts, T., Lamoreux, J., Jenkins, M., Kapos, V., & Randers, J.

704    (2005). The Living Planet Index: Using species population time series to track trends

705    in biodiversity. *Philosophical Transactions of the Royal Society B: Biological*

706    *Sciences*, *360*(1454), 289–295. https://doi.org/10.1098/rstb.2004.1584

707 Loreau, M., Cardinale, B. J., Isbell, F., Newbold, T., O'Connor, M. I., & de Mazancourt, C.

708    (2022). Do not downplay biodiversity loss. *Nature*, *601*(7894), E27–E28.

709    https://doi.org/10.1038/s41586-021-04179-7

710 Lyapunov, A. M. (1992). The general problem of the stability of motion. *International*

711    *Journal of Control*, *55*(3), 531–534. https://doi.org/10.1080/00207179208934253

712 McRae, L., Deinet, S., & Freeman, R. (2017). The Diversity-Weighted Living Planet Index:

713    Controlling for Taxonomic Bias in a Global Biodiversity Indicator. *PLOS ONE*, *12*(1),

714    e0169156. https://doi.org/10.1371/journal.pone.0169156

Medeiros, L. P., Allesina, S., Dakos, V., Sugihara, G., & Saavedra, S. (2022). Ranking

Species Based on Sensitivity to Perturbations under Non-Equilibrium Community

Dynamics. *Ecology Letters*, *n/a*(n/a). https://doi.org/10.1111/ele.14131

Medeiros, L. P., & Saavedra, S. (2023). Understanding the state-dependent impact of species

correlated responses on community sensitivity to perturbations. *Ecology*, *n/a*(n/a),

e4115. https://doi.org/10.1002/ecy.4115

Nicholson, E., Collen, B., Barausse, A., Blanchard, J. L., Costelloe, B. T., Sullivan, K. M. E.,

Underwood, F. M., Burn, R. W., Fritz, S., Jones, J. P. G., McRae, L., Possingham, H.

P., & Milner-Gulland, E. J. (2012). Making Robust Policy Decisions Using Global

Biodiversity Indicators. *PLOS ONE*, *7*(7), e41128.

https://doi.org/10.1371/journal.pone.0041128

O'Brien, D. A., Deb, S., Sidheekh, S., Krishnan, N. C., Sharathi Dutta, P., & Clements, C. F.

(2023). EWSmethods: An R package to forecast tipping points at the community level

using early warning signals, resilience measures, and machine learning models.

Ecography, *n/a*(n/a), e06674. https://doi.org/10.1111/ecog.06674

Oliver, T. H., Isaac, N. J. B., August, T. A., Woodcock, B. A., Roy, D. B., & Bullock, J. M.

(2015). Declining resilience of ecosystem functions under biodiversity loss. *Nature*

Communications, *6*(1), 10122. https://doi.org/10.1038/ncomms10122

O'Loughlin, L. S., & Green, P. T. (2017). Secondary invasion: When invasion success is

contingent on other invaders altering the properties of recipient ecosystems. *Ecology*

and Evolution, *7*(19), 7628–7637. https://doi.org/10.1002/ece3.3315

Park, J., Smith, C., Sugihara, G., & Deyle, E. (2021). *REDM: Empirical Dynamic Modeling*

('EDM') [Computer software]. https://CRAN.R-project.org/package=rEDM

Pimm, S. L. (1984). The complexity and stability of ecosystems. *Nature*, *307*(5949), 321–

326. https://doi.org/10.1038/307321a0

R Core Team. (2022). *R: A Language and Environment for Statistical Computing* [Computer

software]. R Foundation for statistical computing.

Rue, H., Martino, S., & Chopin, N. (2009). Approximate Bayesian inference for latent

Gaussian models by using integrated nested Laplace approximations. *Journal of the*

Royal Statistical Society: Series B (Statistical Methodology), *71*(2), 319–392.

https://doi.org/10.1111/j.1467-9868.2008.00700.x

Strogatz, S. H. (2015). *Nonlinear dynamics and chaos: With applications to physics, biology,*

chemistry, and engineering (2nd ed.). CRC Press.

https://doi.org/10.1201/9780429492563

Sugihara, G., May, R., Ye, H., Hsieh, C., Deyle, E., Fogarty, M., & Munch, S. (2012).

Detecting Causality in Complex Ecosystems. *Science*, *338*(6106), 496 LP – 500.

https://doi.org/10.1126/science.1227079

Traill, L. W., Perhans, K., Lovelock, C. E., Prohaska, A., McFallan, S., Rhodes, J. R., &

Wilson, K. A. (2011). Managing for change: Wetland transitions under sea-level rise

and outcomes for threatened species. *Diversity and Distributions*, *17*(6), 1225–1233.

https://doi.org/10.1111/j.1472-4642.2011.00807.x

Turney, S., & Buddle, C. M. (2016). Pyramids of species richness: The determinants and

distribution of species diversity across trophic levels. *Oikos*, *125*(9), 1224–1232.

https://doi.org/10.1111/oik.03404

Tylianakis, J. M., Didham, R. K., Bascompte, J., & Wardle, D. A. (2008). Global change and

species interactions in terrestrial ecosystems. *Ecology Letters*, *11*(12), 1351–1363.

https://doi.org/10.1111/j.1461-0248.2008.01250.x

United Nations. (2020). *Aichi Biodiversity Targets*. https://www.cbd.int/sp/targets/

United Nations. (1992, June 5). *Convention on biological diversity*.

Ushio, M., Hsieh, C., Masuda, R., Deyle, E. R., Ye, H., Chang, C.-W., Sugihara, G., &

Kondoh, M. (2018). Fluctuating Interaction Network and Time-Varying Stability of a

Natural Fish Community. *Nature*, *554*(7692), 360–363.

https://doi.org/10.1038/nature25504

Wauchope, H. S., Amano, T., Sutherland, W. J., & Johnston, A. (2019). When can we trust

population trends? A method for quantifying the effects of sampling interval and

duration. *Methods in Ecology and Evolution*, *10*(12), 2067–2078.

https://doi.org/10.1111/2041-210X.13302

White, L., O'Connor, N. E., Yang, Q., Emmerson, M. C., & Donohue, I. (2020). Individual

species provide multifaceted contributions to the stability of ecosystems. *Nature*

Ecology & Evolution, *4*(12), 1594–1601. https://doi.org/10.1038/s41559-020-01315-

w

Williamson, M. S., & Lenton, T. M. (2015). Detection of bifurcations in noisy coupled

systems from multiple time series. *Chaos: An Interdisciplinary Journal of Nonlinear*

Science, *25*(3), 036407. https://doi.org/10.1063/1.4908603

Yang, Q., Fowler, M. S., Jackson, A. L., & Donohue, I. (2019). The Predictability of

Ecological Stability in a Noisy World. *Nature Ecology & Evolution*, *3*(2), 251–259.

https://doi.org/10.1038/s41559-018-0794-x

Zhang, F. (2020). Early warning signals of population productivity regime shifts in global

fisheries. *Ecological Indicators*, *115*, 106371.

https://doi.org/10.1016/j.ecolind.2020.106371

785