1 **MANUSCRIPT TITLE**

2 **Affinity-tagged SMAD1 and SMAD5 mouse lines reveal transcriptional reprogramming**
3 **mechanisms during early pregnancy**
4
5 **AUTHORS**
6

7 Zian Liao[a,b,c,d], Suni Tang[a,d], Kaori Nozawa[a,d], Keisuke Shimada[e], Masahito Ikawa[e], Diana Monsivais*[a,d],
8 and Martin M. Matzuk*[a,b,c,d]
9
10 [a]Department of Pathology & Immunology, Baylor College of Medicine, Houston, TX, 77030, USA
11 [b]Department of Molecular and Human Genetics, Baylor College of Medicine, Houston, TX, 77030, USA
12 [c]Graduate Program of Genetics and Genomics, Baylor College of Medicine, Houston, TX, 77030, USA
13 [d]Center for Drug Discovery, Baylor College of Medicine, Houston, TX, 77030, USA
14 [e]Research Institute for Microbial Diseases, Osaka University, Osaka, 565-0871, Japan
15
16 *To whom correspondence should be addressed to: Martin M. Matzuk, MD, PhD Tel: +1 713-798-6451;
17 Fax: +1 713-798-5838; Email: mmatzuk@bcm.edu; and Diana Monsivais, PhD Tel: +1 713-798-1279;
18 Fax: +1 713-798-5838; Email: diana.monsivais@bcm.edu.

21 **KEY WORDS:** uterus, decidualization, SMAD proteins, bone morphogenetic proteins, progesterone,

22 endometrium

ABSTRACT

Endometrial decidualization, a prerequisite for successful pregnancies, relies on transcriptional reprogramming driven by progesterone receptor (PR) and bone morphogenetic protein (BMP)-SMAD1/SMAD5 signaling pathways. Despite their critical roles in early pregnancy, how these pathways intersect in reprogramming the endometrium into a receptive state remains unclear. To define how SMAD1 and/or SMAD5 integrate BMP signaling in the uterus during early pregnancy, we generated two novel transgenic mouse lines with affinity tags inserted into the endogenous SMAD1 and SMAD5 loci (*Smad1*$^{HA/HA}$ and *Smad5*$^{PA/PA}$). By profiling the genome-wide distribution of SMAD1, SMAD5, and PR in the mouse uterus, we demonstrated the unique and shared roles of SMAD1 and SMAD5 during the window of implantation. We also showed the presence of a conserved SMAD1, SMAD5, and PR genomic binding signature in the uterus during early pregnancy. To functionally characterize the translational aspects of our findings, we demonstrated that SMAD1/5 knockdown in human endometrial stromal cells suppressed expressions of canonical decidual markers (*IGFBP1, PRL, FOXO1)* and PR-responsive genes (*RORB*, *KLF15)*. Here, our studies provide novel tools to study BMP signaling pathways and highlight the fundamental roles of SMAD1/5 in mediating both BMP signaling pathways and the transcriptional response to progesterone (P4) during early pregnancy.

INTRODUCTION

Infertility is an emerging health issue that affects approximately 15% of couples[1]. One in five women aged 15 to 49 years old with no prior births suffers from infertility in the United States[2]. One important factor affecting fertility is failed embryo implantation or subsequent post-implantation loss due to endometrial defects. This is evident from the high number of failed pregnancies, with as many as 15% of pregnancies resulting in early pregnancy losses[3]. Understanding the molecular mechanism of how the maternal endometrium becomes suitable for embryo implantation and eventual decidualization, will be the key to eradicating global concerns related to infertility and early pregnancy losses.

The transforming growth factor β (TGFβ) family plays diverse roles in development, physiology, and pathophysiology[4,5], and in particular, signaling pathways downstream of the bone morphogenetic protein (BMP) subfamily are essential for decidual formation[6,7]. There are more than 30 TGFβ family ligands, and these ligands signal through complexes of transmembrane type 1 Activin-Like Kinases (ALK) receptors (ALK1, ALK2, ALK3, ALK6) and transmembrane type 2 receptors (BMPR2, ACVR2A, ACVR2B) and then phosphorylate downstream SMAD1 and SMAD5 proteins. Phosphorylated SMAD1/5 form heteromeric complexes with SMAD4 and translocate to the nucleus to induce specific transcriptional programs. Our laboratory and others have used genetically engineered mouse models with deletions of ligands, receptors, and downstream effectors of BMP signaling pathways to establish that BMP signaling pathways are major regulators of early pregnancy[6-12].

A successful pregnancy begins with reciprocal crosstalk between the maternal endometrium and the new blastocyst during the peri-implantation window. Effective implantation requires precise synchronization between the development of the blastocyst and the transformation of the maternal endometrium into a functional decidua. Endometrial stromal fibroblasts undergo the decidualization process in which they differentiate into unique secretory decidual cells that offer a supportive and immune-privileged microenvironment required for embryo implantation and placental development. Decidualizing stromal cells can react to individual embryos in a way that either supports the implantation and subsequent embryonic development or exerts early rejection[13,14]. Aberrant decidualization processes are observed in patients with recurrent pregnancy loss (RPL), displaying a disordered pro-inflammatory response, decreased induction of decidual marker genes, and abnormal responses to embryonic human chorionic gonadotropin[13,15,16]. In addition to affecting early pregnancy outcomes, defective decidualization is also involved in the maternal etiology of preeclampsia, causing abnormal placental phenotype[17,18]. The process of decidualization is tightly regulated by hormone signaling pathways (estrogen, E2, and progesterone, P4), as well as by BMP signaling pathways. Our recent studies found that endometrial *Smad1* deletion had no significant effect on fertility, *Smad5* conditional deletion resulted in subfertility, while double *Smad1/5* conditional deletion led to infertility due to implantation and decidualization defects[9]. The uteri of mice with double conditional *Smad1/5* deletion also displayed decreased response

to P4 during the window of implantation, suggesting synergy between the two pathways. However, the

mechanistic genomic actions of SMAD1 and/or SMAD5 in the uterus have not been explored, partly

because there are no specific antibodies that distinguish phospho-SMAD1 versus phospho-SMAD5.

In this study, we define how SMAD1/5 instructs the decidualization process using genomic approaches in

newly generated transgenic mouse lines. We inspect the potential crosstalk between P4 and BMP

signaling pathways mediated by SMAD1/5. Together, our study demonstrates that SMAD1 and SMAD5

exhibit shared and unique genomic binding features and further reveals that SMAD1/5 contributes to the

P4 response through transcriptional reprogramming during decidualization.

MATERIALS AND METHODS

Generation of Knock-in Mouse Lines

Smad5[PA/PA] knock-in (KI) mice were generated using a similar approach as previously described[19].

Briefly, single-guide RNA (sg-RNA) was designed to target the regions close to the start codon (Figure

1A,B) and the sgRNA sequence was inserted into the pX459 V2.0 plasmid (#62988, Addgene). The

reference plasmids containing PA tag sequence were constructed in pBluescript II SK (+) vector (Agilent,

Palo Alto, CA, USA). One µg of guide RNA inserted vector and 1.0 µg of reference plasmid were co-

transfected into EGRG01 embryonic stem (ES) cells. Twelve ES clones out of 48 had the expected

knock-in allele. ES cell clones that possessed the proper KI allele were injected into ICR embryos and

chimeric blastocysts were transferred into pseudopregnant females. Chimeric male mice were mated

with B6D2F1 female mice to obtain the PA-tagged SMAD5 KI heterozygous mice. Homozygous

Smad5[PA/PA] mice were maintained in the C57BL/6 J × 129S5/SvEvBrd mixed genetic background. To

generate *Smad1*[HA/HA] mice, Cas9 protein (Thermo Fisher, A36497), sg-RNA, and a repair oligo of

homology-directed repair (HDR) containing HA-tag and linker sequences were electroporated into

zygotes harvested from *in vitro* fertilization using B6D2F1 male and female mice. An ECM830

electroporation system (BTX, Holliston, MA) was used for electroporation. Subsequently, embryos were

cultured overnight to the 2-cell stage and then transferred to the oviducts of pseudopregnant CD-1 mice

(Center for Comparative Medicine, Baylor College of Medicine). Pups were further screened for

successful heterozygous or homozygous knock-in alleles by PCR using primers spanning across the HA

tag. Sequences of sgRNA, the single-stranded repair oligo for HDR and primer used for genotype are

listed in Supplementary Table 1.

Animal ethics compliance and tissue collection

All mice were housed under standard conditions of a 12-hour light/dark cycle in a vivarium with controlled

ambient temperature (70˚F ± 2˚F and 20-70% relative humidity). All mouse handling and experimental

procedures were performed under protocols approved by the Institutional Animal Care and Use

Committee of Baylor College of Medicine. All experiments were performed with female mice aged
between 7 to 12 weeks with a C57BL/6 J × 129S5/SvEvBrd mixed genetic background. All mice were
euthanized using isoflurane induction followed by cervical dislocation, and tissues were snap-frozen in
liquid nitrogen.

Cleavage Under Targets and Release Using Nuclease (CUT&RUN) Approach

Nuclei from uterine tissues were purified following a previously published protocol[20]. The experiments
were performed using pooled biological replicates from two mice that were processed as technical
replicates throughout the CUT&RUN procedure and analysis. In short, uteri were harvested from
pregnant mice at 4.5 days post coitus and washed with cold swelling buffer (10 mM Tris-HCl pH 7.5,
2 mM $MgCl_2$, 3 mM $CaCl_2$, 1X Protease Inhibitor Cocktail (PIC, Roche, 11836170001)) immediately after
collection. Then tissue was cut into small pieces (~2-3mm) using scissors, while submerged in cold
swelling buffer. Nuclear extract was prepared by dounce homogenization in cold swelling buffer (using a
size 7 dounce) and filtered using the cell strainer (100 μm, BD Biosciences). Lysate was centrifuged at
400 g for 10 min, then resuspended in lysis buffer (swelling buffer with 10% glycerol and 1% CA-630, 1X
PIC) using end-cut or wide-bore tips and incubated on ice for 5 min. Nuclei were washed twice with lysis
buffer and resuspended in lysis buffer. Next, CUT&RUN procedure largely follows a previous protocol[21].
Briefly, around 500,000 nuclei were used per reaction. 10 μl of concanavalin-coated beads (Bangs Labs,
BP531) were washed twice in Bead Activation Buffer (20 mM HEPES, pH 7.9, 10 mM KCl, 1 mM $CaCl_2$,
1 mM $MnCl_2$) for each reaction. Then, beads were added to nuclei resuspension and incubated for 10
mins at room temperature. After incubation, bead-nuclei complexes were resuspended in 100 μl
Antibody Buffer (20 mM HEPES, pH 7.5, 150 mM NaCl, 0.5 mM Spermidine, 1X PIC, 0.01% digitonin,
and 2mM EDTA) per reaction. One μg of IgG antibody (Sigma, I5006), HA antibody (EpiCypher, 13-
2010), PA antibody (Fuji Film, NZ-1) and PR antibody (Cell Signaling, D8Q2J) were added to each group
respectively. After overnight incubation at 4 °C, bead-nuclei complexes were washed twice with 200 μl
cold Dig-Wash buffer (20 mM HEPES pH=7.5, 150 mM NaCl, 0.5 mM Spermidine, 1X PIC, 0.01%
digitonin) and resuspended in 50 μl cold Dig-Wash buffer with 1 μl pAG-MNase (EpiCypher, 15-1016)
per reaction. After incubation at room temperature for 10 min, bead-nuclei complexes were washed twice
with 200 μl cold Dig-Wash buffer and resuspended in 50 μl cold Dig-Wash buffer, then 1 μl 100 mM
$CaCl_2$ was added to each reaction. The mixture was incubated at 4 °C for 2 hours and the reaction was
stopped by adding 50 μl Stop Buffer (340mM NaCl, 20 mM EDTA, 4 mM EGTA, 0.05% Digitonin, 100
ug/mL RNase A, 50 mg/mL glycogen, 0.5 ng E. coli DNA Spike-in (EpiCypher, 18-1401)) and incubate at
37 °C for 10 min. The supernatant was collected and subjected to DNA purification with phenol-
chloroform and ethanol precipitation. Sequencing libraries were prepared using NEBNext Ultra II DNA
Library Prep Kit (New England BioLabs, E7645) following manufacture's protocol. Paired-end 150 bp
sequencing was performed on a NEXTSeq550 (Illumina) platform.

Bioinformatic Analysis for CUT&RUN data and re-analysis of published single-cell RNA seq data

For CUT&Run data, raw data were de-multiplexed by bcl2fastq v2.20 with fastqc for quality control. Clean reads were mapped to reference genome mm10 by Bowtie2, with parameters of --end-to-end --very-sensitive --no-mixed --no-discordant --phred33 -I 10 -X 700. For Spike-in mapping, reads were mapped to E. coli genome U00096.3. Duplicated reads were removed, and only uniquely mapped reads were kept. Spike-in normalization was achieved through multiply primary genome coverage by scale factor (100000 / fragments mapped to E. coli genome). CUT&RUN peaks were called by SECAR[22] with the parameters of -norm -stringent -output. Track visualization was done by bedGraphToBigWig[23], bigwig files were imported to Integrative Genomics Viewer for visualization. For peak annotation, common peaks were identified with 'mergePeaks' function in HOMER v4.11[24] and then genomic annotation was added by ChIPseeker [25]. Motif analysis was conducted through HOMER v4.11 with parameter set as findMotifsGenome.pl mm10 -size 200 -mask[24]. Peak heatmaps were plotted using deepTools 2.4.2 with clustering options set to k-means[26]. For imputing the cell fractions using published mouse uterine bulk and single-cell RNA seq data, single-cell RNA-seq with cell type reference from 5.5 dpc mouse uterus was derived from GSE226417, bulk RNA-seq of the 3.5 dpc mouse uterus was derived from GSE152675. Signature matrix and cell fraction profiles were conducted through CIBERSORTx[27]. Briefly, reference gene expression signature matrix was generated by "Create Signature Matrix" module with default settings. Next, cell fractions were calculated using "Impute Cell fractions" module in the absolute mode, which scales each annotated cellular fractions to an absolute value that reflects its absolute proportion in the bulk sample. For human single-cell RNA seq data, raw data was obtained from EMBL-EBI under accession No. E-MTAB-10287. Cells with low coverage (less than 500 genes detected) were filtered, then gene counts were normalized for each cell by converting counts to quantiles and obtaining the corresponding values from a normal distribution. Then normalized cell vectors are concatenated along the gene panel. Plot visualization was conducted through CELLXGENE platform.[28]

Western Blot Analysis of Immunoprecipitation (IP-WB)

Tissues were pulverized in liquid nitrogen and then lysed using NETN buffer (20 mM Tris-HCl, pH 8.0, 150 mM NaCl, 0.5 mM EDTA, 10% Glycerol, and 0.5% NP-40). Protein concentration was determined by BCA Protein Assay Kit (Thermo Fisher, 23225). 1.5 mg of total protein lysate was used for IP. IP was performed by adding HA antibody (Cell Signaling, C29F4) or PA antibody (Fuji Film, NZ-1) to the lysate and incubate for 1 hour at 4 °C. Subsequently, protein G magnetic beads (Thermo Fisher, 88847) were added for an additional 1 hour at 4 °C. Then, the beads were washed five times with NETN buffer, and denatured in sample buffer (Thermo Fisher, NP0007) for further analysis by Western blot. For western blot procedures, briefly, denatured protein lysates were run on the 4 to 12%, Bis-Tris protein gels (Thermo Fisher, NP0321BOX) followed by electrophoretic transfer to nitrocellulose membrane. The

membrane then went through blocking by 5% milk in Tris-buffered saline with Tween20 (TBST), followed

by incubation overnight at 4 °C in the primary antibodies anti-HA (Cell Signaling, C29F4), anti-PA (Fuji

Film, NZ-1), anti-SMAD1 (Life technologies, 385400) and anti-SMAD5 (ProteinTech, 12167-1-AP) at

1:1,000 dilution. The next day, membranes were washed three times with TBST, then incubated with

horseradish peroxidase–conjugated secondary antibody for 1 h at room temperature, then washed three

times with TBST, developed and imaged on iBright Imaging System (FL1500).

Primary endometrial stromal cells isolation/RNAi/decidualization

Studies using human specimens were conducted as indicated in a protocol approved by the Institutional

Review Board at Baylor College of Medicine, H-51900. Human endometrial stromal cells were collected

from healthy volunteers' menstrual effluent as previously reported[29-31]. (N=3) In brief, samples were

collected by participants in a DIVA cup during the 4-8 hours on the first night of menses and stored in

DMEM/F12 with 10% FBS, antibiotic/antimycotic and 100µg/ml Primocin in a cold insulated pack until

processing in the laboratory on the day of collection. The effluent was digested with 5 mg/ml collagenase

and 0.2 mg/ml DNase I for 20 min at 37 °C, then the cell pellet was collected by centrifuging at 2,500 rpm

for 5 min at room temperature. Next, red blood cell lysis was performed by resuspending the cell pellet in

20 ml of 0.2% NaCl for 20 seconds and neutralized with 20 ml of 1.6% NaCl. Then the solution was then

centrifuged at 2,500 rpm for 5 min. Five ml complete medium (DMEM/F12 supplement with 10% FBS, 1X

Antibiotic-Antimycotic + 100 µg/ml Primocin) was used to resuspend the pellet and the solution was

passed through 100 µm and 20 µm cell strainer sequentially. The flowthrough containing the stromal

cells was centrifuged for 5 min at 2,500 rpm and the pellet was resuspended in 10 ml complete medium

and plated in a 10 cm dish. siRNA knockdown was performed using Lipofectamine RNAiMAX following

the manufacturer's protocol. In brief, 0.2 million stromal cells were plated in a 12-well plate one day

before transfection. On the day of transfection, 2 µl siRNA (20 µM, Dharmacon, D-001810-10, L-012723-

00-0005, L-015791-00-0005) and 3 µl Lipofectamine RNAiMAX were diluted in 50 µl Opti-MEM

respectively and then mixed to incubate at room temperature for 15 min. Then, the complex was added

dropwise onto the cells. 24 h after transfection, medium was changed to DMEM/F12 supplement with 2%

charcoal stripped FBS. Decidualization was induced by the addition of 35 nM estradiol (Sigma, E1024), 1

µM medroxyprogesterone (Sigma, 1378001), and 0.05 mM cyclic adenosine monophosphate (Axxora,

JBS-NU-1502L) for 4 days with media changes every 48 hours.

RNA Extraction and RT-qPCR

For mRNA extraction from stromal cells, cells were lysed with TRIzol and processed using the DirectZol

kit (Zymo, R2051) following manufacturer's procedures. Approximately 100 ng of mRNA was reverse

transcribed into cDNA using iScript cDNA Supermix (Bio-Rad, 1708890) and amplified using specific

primers listed in Supplementary Table 1. Primers were amplified using 2X SYBR Green Reagent (Life

Technologies, 4364346) using a BioRad CFX384 Touch Real Time PCR Detection System. Data

analysis was performed by calculating ΔΔCT value towards the geometric mean of *GAPDH, ACTB* and

RPL13A and then normalized to siCTL. P-value was determined by One-Way ANOVA test with post hoc

Tukey test using GraphPad Prism. * $P \leq 0.05$, ** $P \leq 0.01$, *** $P \leq 0.001$, **** $P \leq 0.0001$.

RESULTS

Generation of Mouse Models with Global HA-Tagged SMAD1 and PA-Tagged SMAD5 Proteins

Activation of BMP signaling pathways has been established as one of the hallmarks of the

decidualization process[32,33]. Canonically, SMAD1/5 are regarded as downstream effectors of BMP2

signaling pathways to regulate decidual-specific gene expressions[6,34]. However, our recent findings

demonstrated that SMAD1/5 can also affect the sensitivity of the endometrium towards E2 and P4

stimulation[9]. Since we observed phenotypical differences between uterine-specific single SMAD1 and

single SMAD5 deletion mice, it is beneficial to delineate the role of SMAD1 and SMAD5 in mediating P4

responses during early pregnancy. We used CRISPR technology to generate genetically engineered

knock-in mice with an HA-tagged *Smad1* allele (herein called *Smad1*[HA/HA]) and PA-tagged *Smad5* allele

(herein called *Smad5*[PA/PA]) as shown in Figure 1A, B. The HA tag and the PA tag were inserted into the

N-terminus of the SMAD1 and SMAD5 proteins, respectively. Sanger sequencing was used to confirm

genomic insertion (Figure 1A, B). To validate the global detection of tagged proteins, we performed

immunoprecipitation followed by western blot analysis on different tissues from *Smad1*[HA/HA] and

Smad5[PA/PA] mice. We confirmed the HA and PA antibodies can readily detect HA-tagged SMAD1 and

PA-tagged SMAD5 proteins at the predicted size (Figure 1C, D). We also demonstrated the molecular

size and expression pattern of HA antibody detected SMAD1 protein was comparable to the SMAD1

antibody detected SMAD1 protein across different tissue types. Similarly, PA antibody showed

comparable signal intensity to the SMAD5 antibody in detecting SMAD5 protein across different tissue

types (Figure 1C, D). Thus, we successfully generated viable mouse models with global HA-tagged

SMAD1 and PA-tagged SMAD5 proteins.

SMAD1 and SMAD5 Exhibit Shared and Unique Genomic Binding Sites During Decidualization

The BMP signaling pathway regulates multiple key events during early pregnancy[5], mediated through

receptor-regulated SMAD proteins, including SMAD1 and SMAD5. As transducers of the BMP signaling

pathway, phosphorylated SMAD1 and SMAD5 form homomeric complexes and then couple with SMAD4

to assemble hetero-oligomeric complexes in the nucleus to execute transcription programs. Our previous

studies revealed that conditional ablation of SMAD1 and SMAD5 in the uterus decreased P4 response

during the peri-implantation period, suggesting that the transcriptional activities of PR depend on

BMP/SMAD1/5 signaling[9]. Furthermore, previous genome-wide PR binding studies show that SMAD1

and SMAD4 binding motifs are enriched in PR binding sites in the uterus[35].

To determine the shared and unique transcriptional regulomes of SMAD1 and SMAD5 contributing to the

diverse effects of BMP and P4 signaling pathways during decidualization, we first utilized Cleavage

Under Targets & Release Using Nuclease (CUT&RUN)[21] coupled with next generation sequencing to

profile genomic loci bound by SMAD1, SMAD5 and PR from mouse uterine tissues. We performed

CUT&RUN on the uterine tissues collected at 4.5 days post coitus (dpc), the time when the fertilized

embryo reaches the uterus physically and initiates the decidualization program[36] (Figure 2A). After

aligning CUT&RUN reads to the mm10 mouse genome, we called peaks using Sparse Enrichment

Analysis for CUT&RUN (SEACR)[22]. To identify high confidence peaks, background noise was

normalized to IgG and the stringent criteria for peak calling in SEACR was used. After merging common

peaks from two biological replicates, we identified 118,778 peaks for SMAD1 and 166,025 total peaks for

SMAD5. We visualized the enrichment of SMAD1 and SMAD5 peaks to the overall aligned chromatin

regions with clustering for preferential enrichment, as shown in Figure 2B. Peaks in cluster 1 exhibit a

shared enrichment for both SMAD1 and SMAD5, whereas clusters 2 and 3 demonstrate preferential

enrichment for SMAD5 and SMAD1, respectively. We found that 7.55% of SMAD1 peaks and 9.53% of

SMAD5 peaks were located within the ± 3 kb of the promoter regions (Figure 2C, D). This corresponded

to 10,368 genes that were directly bound by SMAD1 at the promoter regions (± 3 kb), whereas 18,270

genes were directly bound by SMAD5 at the promoter regions (± 3 kb). Among these, 4,933 genes were

found in common between SMAD1 (47.5%) and SMAD5 (27.0%), while 2,744 and 7,427 genes were

found to be uniquely bound by SMAD1 and SMAD5, respectively, providing evidence for the shared and

unique functions of SMAD1 and SMAD5 at the transcriptional level (Supplement Figure 1A). Hence,

interpreting how the binding events correlate to biological activity requires comparisons with gene

expression profiling in a tissue-specific manner.

Identification of Direct Target Genes of SMAD1 and SMAD5 During Early Pregnancy

To pinpoint the direct target genes of SMAD1 and SMAD5, we integrated transcriptomic data from

previously published[9] SMAD1/5 double conditional knockout mice using progesterone receptor cre

(SMAD1/5 cKO) (GSE152675) with SMAD1 and SMAD5 genomic data from this paper. We cross-

compared the differentially expressed genes in the transcriptomic data to the SMAD1 and SMAD5 bound

genes, respectively. Among the 805 significantly up-regulated genes, we identified 449 genes that were

both significantly up-regulated upon SMAD1/5 depletion and were directly bound by SMAD1 and

SMAD5, whereas 187 of the up-regulated genes were bound by SMAD5 only and 30 were bound only by

SMAD1 (Figure 3A). Among the 683 significantly down-regulated genes, we identified 523 genes that

were both significantly down-regulated upon SMAD1/5 depletion and were directly bound by SMAD1 and

SMAD5, whereas 83 of the down-regulated genes were bound by SMAD5 only and 13 were bound by

SMAD1 only (Figure 3B, Supplementary Table 2).

Next, we utilized Binding and Expression Target Analysis (BETA) algorithm[37] to perform motif enrichment

analysis of the direct target genes to identify putative co-factors working together with SMAD1 and

SMAD5 in controlling gene expression (Figure 3C,D, Supplement Files1,2). "Up-targets" represent genes

that were up-regulated in the SMAD1/5 cKO mouse uteri and showed either a SMAD1 or a SMAD5

binding site in the genomic profiling data. Similarly, "down-targets" represent genes that were down-

regulated in the SMAD1/5 cKO mouse uteri and displayed either a SMAD1 or a SMAD5 binding site.

Thus, motifs enriched in the "up-targets" indicate potential repressive SMAD1/5 co-factors while motifs

enriched in the "down-targets" indicate potential SMAD1/5 co-activators. Among the "up-targets" of

SMAD1, MYB Proto-Oncogene (*Myb*)/MYB Proto-Oncogene Like 1(*Mybl1*) motif was the most highly

enriched with a P-value of 1.85E-02. *Myb* and *Mybl1* transcription factors belong to MYB gene family,

which has been well-defined in controlling cell survival, proliferation and differentiation in cancer[38]. In

addition, they have also been reported to be E2 induced in human uterine leiomyoma samples[39].

Homeobox containing 1 (*Hmbox1*) and Krüppel-like factor (*Klf*) family members (*Klf4/Klf1/Klf12*) were

also identified as potential repressive co-factors of SMAD1 with P-values of 2.85E-02 and 3.75E-02

respectively (Figure 3C). Of note, *KLF4* has been reported to inhibit the binding activity of estrogen

receptor α (ERα) to estrogen response elements in promoter regions[40]. Among the "up-targets" of

SMAD5, EBF Transcription Factor 1 (*Ebf1*) motif was the most enriched with a P-Value of 1.57E-02.

Interestingly, *Ebf1* can directly repress the transcription of Forkhead box protein O1 (*Foxo1*)[41]. It is also

recognized as downstream effector of steroid hormone receptors in the mouse uterus[42]. Additionally,

motifs from transcription factors *Zfp128* and *Otx1* were also significantly enriched in the up-regulated

genes bound by SMAD5 (Figure 3D). Taken together, our enrichment analysis provided robust evidence

for identifying novel co-factors of SMAD1/5, and such co-regulating mechanisms are in line with the

unopposed E2 response observed in the SMAD1/5 cKO mice[9]. Furthermore, odd-skipped–related genes

(*Osr1* and *Osr2*) were identified as potential co-activators for SMAD1. *Osr2* has been reported to be

highly expressed in the human endometrium[43], and it was also abundantly detected at the protein level in

the human decidual tissues[44]. Decreased OSR2 level was observed in the patients with recurrent

spontaneous abortion and knockdown of *OSR2* impairs the decidualization process in the human

endometrial stromal cells[44]. Moreover, OSR1 has been reported to suppress BMP4 expression, which in

turn reduced the Wnt/β-catenin signaling pathways during lung development in xenopus[45]. Apart from

Osr family, motifs in the Homeobox genes (HOX) were found to be enriched in the "down-targets" from

both SMAD1 and SMAD5 datasets. Specifically, *Hoxa11/Hoxd12/Hoxc10* were predicted to be co-

activators for SMAD1 while *Hoxd10* was indicated to be closely interacting with SMAD5. Indeed, HOX

genes are critical for endometrial development in normal and disease conditions and are essential during
the establishment of pregnancy[46-49].

With direct target genes of SMAD1 and SMAD5 identified, we then analyzed the Gene Ontology
enrichment for the SMAD1/5 shared up-targets and down-targets, respectively. We found that "up-
targets" genes exhibit enrichment for regulation of cell-cell adhesion, cell junction organization and
desmosomes organization (Figure 3E, Supplementary Table 3). Moreover, among the "down-targets"
genes, we found the enrichment for blood vessel / vasculature development and extracellular matrix
organization categories (Figure 3F, Supplementary Table 4). Indeed, during early pregnancy, the
stimulation from corpus-luteum derived P4 enabled the endometrium to be transformed to a receptive
state, which allows subsequent embryo attachment and develop through the epithelium into the stromal
sections[32]. During this process, apportioned direct cell-cell contacts are ensured by tight and adherent
junctions and such interactions are key in facilitating implantation and embryo invasion. In accordance
with our findings, desmosomes and adherens junctions were extensively described to decline in the early
pregnancy period, which facilitates the invasion of trophoblast through the epithelial layer[50-53]. In addition,
the stromal compartment of the endometrium also undergoes profound vascular remodeling. Precise
regulations of angiogenesis are required to establish an extensive vascular network, which is essential to
ensure blood supply and successful embryonic development[54,55]. Collectively, our findings present
evidence that emphasizes the shared roles of SMAD1 and SMAD5 in facilitating endometrial transitions
during early pregnancy.

Direct Target Genes of SMAD1 and SMAD5 Maintain the Homeostasis of Uterine Function

To discover novel direct target genes of SMAD1/5, we visualized key genes of interest from the up-
targets and down-targets. As shown in Figure 4A, data from RNA-seq represents the decrease of several
"down-targets" in the SMAD1/5 cKO mouse uteri, including Retinoic Acid-Related Orphan Receptor B
(*Rorb*), Follistatin (*Fst*), Lymphoid Enhancer Binding Factor 1 (*Lef1*), and Insulin Like Growth Factor 1
(*Igf1*). Integrative Genomics Viewer (IGV) track view shows the exemplary SMAD1/5 binding activities
near the promoter regions of *Rorb* and *Fst* Figure 4B, demonstrating that these genes are bona fide
direct target genes of SMAD1/5. *Rorb* belongs to the nuclear receptor families in the retinoic acid (RA)
signaling pathways[56] and is considered as a marker for mesenchymal progenitor cells in the stroma
compartment of the endometrium[57]. In murine models, deficient RA signaling through the perturbation of
RA receptor in the uterus leads to implantation and decidualization failure[58]. *Fst* binds several TGFβ
family ligands and thereby inhibits TGFβ family signaling extracellularly[59]. Under physiological conditions,
Fst is up-regulated in the decidua during early pregnancy. Conditional deletion of *Fst* in the mouse uterus
results in severe subfertility with a phenotype of non-receptive epithelium and poorly-differentiated
stroma[60]. Notably, RA signaling deficiency also decreases *Fst* levels in the uterus and systematic

administration of FST can fully rescue the deficient-decidualization phenotype but not the non-receptive

phenotype observed in the RA receptor mutant mice[58]. Our results suggest a direct relationship between

BMP and RA signaling pathway, accomplished by SMAD1/5 at the transcriptional level, likely

establishing a positive signaling feedback loop. Apart from being a crucial transcriptional activator,

SMAD1/5 also plays a role in repressing key gene expression pathways. Shown in Figure 4C, upon the

deletion of SMAD1/5 in the mouse uteri, several E2-responsive genes were significantly up-regulated,

including Fibroblast Growth Factor Receptor 2 (*Fgfr2*), Matrix Metallopeptidase 7 (*Mmp7*) and Wnt

Family Member 7B (*Wnt7b*). In addition, *Inhbb*, a downstream target of *Fst*[60], is also a target gene of

SMAD1/5 that resulted in transcriptional repression. SMAD1/5 binding on the *Fgfr2* and *Mmp7* genes are

exemplified in an IGV track view in Figure 4D. *Fgfr2* and its ligands regulate epithelial cell proliferation

and differentiation. Components of the Fibroblast Growth Factor (*Fgf*) signaling pathway are cyclically

expressed in the uterus and act as paracrine and/or autocrine mediators of epithelial-stromal

interactions[61,62]. During early pregnancy in mice, P4 inhibits expression of *Fgf2* in the stromal cells, which

is critical to counteract the E2-driven epithelial proliferation[61]. Similar observations are reported in gilts,

where the expression of *Fgfr2* decreased alongside with increased parity of the sows[63]. It is also

noteworthy that loss of function of *Fgfr2* in the mouse uterus leads to luminal epithelial stratification and

peri-implantation pregnancy loss[62]. Moreover, *Mmp7* and *Wnt7b* are up-regulated upon E2 stimulation

and participate in the re-epithelialization of the endometrium and implantation process, respectively[64-66].

In accordance with the phenotype of hyperproliferative endometrial epithelium during early pregnancy

observed SMAD1/5 cKO mice, we demonstrated that the suppression of key E2-responsive genes, such

as *Fgfr2* and *Mmp7,* by SMAD1/5 maintains the precise balance between E2 and P4.

To explore the major cell types regulated by SMAD1/5, first, we used CIBERSORTx[27] to analyze and

depict changes in the cell populations upon SMAD1/5 depletion in the mouse uterus during early

pregnancy. By imputing the bulk uterine gene expression profiles to previously published mouse uterine

single-cell datasets[67] using CIBERSORTx[27], we were able to compare changes across both samples and

cell types upon the SMAD1/5 perturbation in the mouse uterus. We highlight the proportional increase in

the epithelial cells, as well as the decrease in the decidual stromal cells and smooth muscle cells in mice

lacking uterine SMAD1/5 during the peri-implantation phase (Supplement Figure 1B). Such cell

populational changes are in line with the phenotypical observations of decidualization failure and

excessive proliferation in the epithelial compartment. In addition, to explore the expression patterns of

SMAD1/5 direct targets in human, we profiled the expression levels of the key "up-targets" and "down-

targets" in the different cell types of the human endometrium. Using previously published single-cell RNA

seq data of human endometrium[68], we visualized the expression patterns of suppressive targets and

activating targets of SMAD1/5 (Figure 4E). Apart from the major epithelial and stromal compartments,

SMAD1/5 target genes are also widely expressed in the immune cell populations. Such observations

reinforced the importance of the BMP signaling pathways in establishing an immune-privileged

environment at the maternal-fetal interface[69].

SMAD1 and SMAD5 Co-regulate PR Target Genes

SMAD1/5 cKO mice were infertile due to endometrial defects and displayed decreased P4 response

during the peri-implantation period[9]. Hence, we hypothesized that SMAD1 and SMAD5 act as co-

regulators of P4-responsive genes during the window of implantation and are required for endometrial

receptivity and decidualization. By determining the genomic co-occupancy of SMAD1, SMAD5 and PR,

we aimed to clarify the transcriptional interplay between the BMP and P4 signaling pathways. To this

end, we performed additional PR CUT&RUN experiments on the uteri of mice collected at 4.5 dpc and

identified 134,737 peaks showing PR binding activities (Figure 5A). Based on the k-means clustering

results of the peaks, we demonstrated clusters with shared occupancy between SMAD1/5 and PR

(cluster 1), preferential deposition in the SMAD1 (cluster 2), SMAD5 (cluster 4) and PR (clusters 3,5),

respectively. Interestingly, between clusters 3 and 5, although the primary enrichment is for PR, overall

the signal intensities for SMAD5 are higher in cluster 5. Together with previous analysis on genes

uniquely or commonly bound by SMAD1/5 (Supplement Figure 1A), we speculate such observation can

be attributed to a subset of the genes that are potentially co-regulated by SMAD5 and PR. From the

gene perspective, we identified 7,393 genes that were directly bound by PR at the promoter regions (± 3

kb), among which, 2596 genes were also concurrently bound by both SMAD1 and SMAD5 at the

promoter regions (± 3 kb) (Supplement Figure 1C).

Next, we performed KEGG pathway enrichment for the genes co-bound by SMAD1, SMAD5 and PR. As

expected, pathways critical for decidualization such as relaxin signaling pathways and WNT signaling

cascade were identified in the enrichment results (Figure 5B, Supplementary Table 5). We visualized

exemplary genes co-regulated by SMAD1, SMAD5 and PR and presented in the normalized IGV track

view (Figure 5C). We demonstrated SMAD1, SMAD5 and PR showed co-occupancy at the loci of the

SRY-Box Transcription Factor 17 (*Sox17*)*,* Inhibitor of DNA binding 2 (*Id2*)*,* Forkhead box protein O1

(*Foxo1*)*,* Insulin-like growth factor 1 (*Igf1*), Transforming growth factor beta receptor 2 (*Tgfbr2*) and

RUNX family transcription factor 1 (*Runx1*) (Figure 5C). *Sox17* has been reported as one of the direct

target genes of PR[35] and is essential for uterine functions during implantation and early pregnancy[70,71].

More recent studies also showed the importance of *Sox17* in regulating uterine epithelial–stromal

crosstalk and its indispensable role in female fertility[72]. We provided evidence that *Sox17* is also directly

regulated by SMAD1/5 complexes. Our results indicated that *Id2*, considered as canonical direct

transcriptional targets of BMP-SMAD signaling[73,74] is also regulated by PR. We also confirmed that

known P4-responsive genes such as *Tgfbr2*[75] and *Runx1*[76], as well as decidual markers such as *Foxo1*[77]

and *Igf1*[78], were co-regulated by SMAD1, SMAD5 and PR (Figure 5C).

To identify additional transcription factors that are associated with the regulatory interplay between SMAD1/5 and PR during decidualization, we performed unbiased motif analysis on the shared CUT&RUN peaks between SMAD1/5 and PR. We reported the top 10 transcription factors harboring the enriched motifs, including NANOG, Homeobox A protein family (HOXA11 and HOXA9), NK6 homeobox 1(NKX6.1), TGFB induced factor homeobox 2 (TGIF2), FOS, RUNX family transcription factor 2 (RUNX2), Androgen receptor (AR), SOX17 and Lymphoid enhancer-binding factor 1 (LEF1) (Figure 5D, Supplementary Table 6). Many of these putative interactors have been reported to interact with the SMAD proteins in other biological processes. For example, NANOG interacts with SMAD1 during mesoderm differentiation[79]. HOXA9 forms heterodimers with SMAD4, leading to BMP-driven initiation of transcription from the mouse *Opn* promoter *in vitro*[80,81]. Transcription factor AP-1 family (FOS) and RUNX2, as well as β-catenin/Lef1 complex, increase the effectiveness and specificity of DNA binding activities of SMAD1/5 in response to BMP ligand stimuli[82-84]. To further evaluate the key roles of SMAD1/5 as major uterine transcription regulators, we cross-compared the genomic binding sites of SMAD1/5 with known key transcription factors, namely aforementioned SOX17 (Supplement Figure 1F, Supplementary Table 7), as well as NR2F2 (Supplement Figure 1G, Supplementary Table 8), an essential regulator of hormonal response[85], using our CUT&RUN data sets and published mouse uterine SOX17 and NR2F2 ChIP-seq data sets (GSE118328, GSE232583). Among the annotated genes, 5402 genes are shared between SMAD1/5 and SOX17, and 1922 genes are shared between SMAD1/5 and NR2F2. Such observations indicate a potential co-regulatory mechanism between SMAD1/5 and other key uterine transcription factors in maintaining appropriate uterine functions. Overall, our analyses demonstrate that the transcriptional activity of SMAD1, SMAD5, and PR coordinate the expression of key genes required for endometrial receptivity and decidualization.

Decidualization of Human Endometrial Stromal Cells Requires SMAD1/SMAD5

We next sought to functionally characterize the role of SMAD1/5 during decidualization in human endometrial stromal cells. To do so, we examined the effect of SMAD1/5 perturbations on the decidualization of primary human endometrial stromal cells (EnSCs). EnSCs were transfected with short interfering RNAs (siRNAs) targeting each gene (*SMAD1* and *SMAD5)* and subjected to *in vitro* decidualization by treatment with E2-cAMP-and MPA (EPC) for 4.5 days (Figure 6A). We hypothesized that the combined SMAD1/5 knockdown would impair the decidualization process significantly compared to cells treated with non-targeting siRNAs. Our results demonstrated that SMAD1/5 knockdown affected decidualization and led to significantly decreased expression of the canonical decidual markers, *PRL and IGFBP1* in EnSCs (Figure 6B). The PR co-regulator, *FOXO1*[86]*,* also exhibited a significant decreasing trend in the siSMAD1/5 group. We also examined the expression level of the RA pathway regulator gene, *RORB*, and of the SMAD4-PR target gene, *KLF15*[12], following SMAD1/5 perturbation. We observed a

significant decrease in both *RORB* and *KLF15* expression upon SMAD1/5 knockdown during *in vitro*

decidualization treatment (Figure 6C). Taken together, our findings indicate SMAD1/5 can modulate PR

activity during decidualization and that this transcriptional cooperation is required for the *in vitro*

decidualization of primary human endometrial stromal cells.

DISCUSSION

SMAD proteins are canonical transcription factors that are activated in response to TGFβ family signaling

and mediate the biological effects of these pathophysiologically critical ligands[83]. While SMAD2 and

SMAD3 are downstream of TGFβs, activins, and multiple other family ligands, SMAD1 and SMAD5

preferentially transduce BMP signaling pathways and are regarded as pivotal activators for many

physiological processes, including bone development, cardiac conduction system development, and

embryonic pattern specification[87-89]. Importantly, SMAD1 and SMAD5 are implicated in diverse female

reproductive physiology and pathophysiology processes[5,9,90-92].

Due to high structural similarity, SMAD1/5 have been suggested to be redundant from the studies in

ovarian biology and chondrogenesis[91,93]. However, other studies clearly demonstrated that SMAD1/5

have different roles in governing hematopoiesis and uterine functions[9,94]. The DNA binding activities of

SMAD1 and SMAD5 have not been readily distinguished from each other due to anti-phospho antibody

limitations. To robustly define the roles of SMAD1/5 in regulating transcriptional programs *in vivo*, we

produced two genetically engineered mouse models with global knock-in of an HA tag and a PA tag in

the *Smad1* and *Smad5* loci, respectively. We showed that SMAD1 and SMAD5 not only have shared

transcriptional activities but also have unique roles in uterine physiology. In agreement with previous

studies showing that SMAD1/5 function is partially redundant[91,93], we confirmed that SMAD1/5 share a

total of 972 direct target genes in the uterus. Furthermore, we demonstrated that 43 genes were uniquely

regulated by SMAD1 whereas 270 genes are specifically regulated by SMAD5 only. Our motif analysis

also revealed distinct potential co-factors between SMAD1 and SMAD5, providing evidence at the

molecular level to mechanistically delineating the distinct roles of SMAD1 and SMAD5 in directing

cellular processes in the uterus.

Apart from directly regulating target gene expression, our data demonstrate that SMAD1/5 present as

dense genomic occupancies. To date, only a limited amount of transcription factors have been

investigated using the CUT&RUN-seq technique from the tissue samples due to antibody compatibility

issue. We recognize that the binding sites and gene number identified here are quite high; however, the

high density of binding events was also observed in the ENCODE[95] chromatin immunoprecipitation

followed by sequencing (ChIP-seq) data for SMAD1 and SMAD5 in the human K562 cells, detecting an

average of 63,563 peaks for SMAD1 and 109,682 peaks for SMAD5. (Data accessed through GSE95876

and GSE127365 from Gene Expression Omnibus). Multiple aspects can contribute to the observation of

dense SMAD1/5 genome occupancies. First, transcription factors (TFs) tend to dwell or "search and

bind" throughout the genome[96]. Such events may not yield actual biological effects but rather are due to

differences in motif binding affinities[97]. Second, apart from robust binding activities, TFs may not initiate

transcription programs owing to the lack of co-factors or favorable conditions to exert their functions[98].

Additionally, TF binding sites and target genes are unlikely to have a one-to-one relationship. TFs could

be positioned from the proximal promoter regions to hundreds of kilobases afar to modulate gene

expression. In the meantime, the same binding site could regulate multiple genes by interacting with

different promoters in different subpopulations of cells. Lastly, TFs usually direct target gene expression

in a cell-type specific manner[99]. Our genomic profiling samples were collected from whole uterus at the

time of 4.5 dpc, containing a great range of cell populations, including but not limited to the epithelium

(luminal and glandular), stroma (progenitors and differentiated cells), myometrium, endothelium, and

immune cell populations. The data is therefore expected to depict the dynamic and complex activities of

SMAD1/5 in the entire uterus. Together, the stringent filtering and normalization criteria, comparable

peak number to the published dataset and IGV track view visualization collectively validate our

CUT&RUN experiments and uncover the enriched regions as robust SMAD1/5 binding events. Our

studies also examined the role of SMAD1/5 in mediating progesterone responses at the genomic and

transcription levels. Similarly, our analysis was based on data sets generated from the whole mouse

uterus, which contains multiple compartments of the uterine structures, including but not limited to

epithelium and stroma. Published studies have shown that nuclear SMAD1/5 localize to the stroma and

epithelium during the decidualization process at 4.5 dpc, during the window of implantation[9]. Conditional

deletion of SMAD1/5 exclusively in the uterine epithelium using lactoferrin-icre (*Ltf*-icre) results in severe

subfertility due to impaired implantation and decidual development[100]. Conditional deletion of SMAD1/5/4

exclusively in the cells from mesenchymal lineage (including uterine stroma) using anti-Mullerian

hormone type 2 receptor cre (*Amhr2-cre*) results in infertility with defective decidualization[91,92]. Given the

essential roles of SMAD1/5 in both stroma and epithelium identified by previous studies, we believe that

the transcriptional co-regulatory roles of SMAD1/5 and PR reported here using the whole uterus

validates a relationship between SMAD1/5 and PR in both the stromal and epithelial compartments.

However, it does not rule out potential co-regulatory roles of SMAD1/5 and PR in the myometrium,

immune cells, and/or endothelium, given that whole uterus was used. The specific transcriptional

evaluations of SMAD1/5 in the stroma versus the epithelium would require future validations using

single-cell sequencing and/or spatial transcriptomic analysis.

Although our studies herein confirm that SMAD1 and SMAD5 proteins have distinct transcriptional

regulatory activities, our previous studies demonstrated that while SMAD5 can functionally replace

SMAD1, SMAD1 cannot replace SMAD5 in the uterus[9]. How this epistatic relationship is established in a

tissue-specific manner still needs to be determined by further biochemical investigations. In addition,

further studies are needed to uncover whether SMAD1 and SMAD5 response differently upon ligand

stimulation in the uterus, and if so, how the preference is achieved. Our study provides versatile *in vivo*

genetic tools for these questions and can advance the toolbox for the field studying BMP signaling

pathways. Because our mouse models are global knock-in mice, they will not only serve as a powerful

tool for studying BMP signaling pathways in the reproductive system but will also promote the study of

BMP signaling in other organs and tissues.

BMP signaling pathways are involved in a plethora of cellular processes and appropriate functioning of

the BMP pathway depends on the precise crosstalk with other signaling pathways. Coordinated

communication with other pathways can yield synergistic effects and lead to a complex regulatory

network of biological processes. To be specific, SMAD1/5 mediates the crosstalk with the WNT/β-catenin

pathway. WNT signaling inhibits glycogen synthase kinase 3β (GSK3β) activity and prevents SMAD1

from degradation which governs the embryonic pattern formation[101]. Also, SMAD1/5 can physically

interact with T-cell factor (TCF) or lymphoid enhancer factor (LEF) transcription factors to form

transcriptional complexes to activate the transcription of many WNT-and BMP-responsive genes[102]. In

addition, SMAD1 and SMAD5 can directly associate with Notch intracellular domain and enhance known

Notch target gene expression by binding to their regulatory DNA sequences[103]. Intriguingly, in prostate

cells, SMAD1 physically interacts with the androgen receptor (AR) and halts the androgen-stimulated

prostate cell growth[104]. Moreover, we provide first-hand evidence showing that BMP signaling pathways

converge with RA signaling pathways through the regulation of RORB by SMAD1/5. Further studies will

grant a more detailed mechanism of the positive feedback loop between BMP and RA signaling.

Our previous studies suggest that the mouse endometrium presents decreased P4 responsiveness

following the conditional deletion of SMAD1/5 in the uterus[9]. In accordance with the phenotypical

observation, we offer compelling support in our current study that SMAD1/5 work collectively with PR to

regulate their target genes and that SMAD1/5 mediate the crosstalk between BMP and P4 signaling

pathways during decidualization, a key process to ensure a successful pregnancy, and ultimately direct

the biological transformations of the uterus during early pregnancy. We provide genomic evidence that

SMAD1/5 are co-bound at around 35% of PR target genes in the mouse uterus during decidualization.

Correspondingly, in a previously published study where they performed PR ChIP-seq in the mouse

uterus after P4 stimulation, the SMAD1 motif was the 5[th] most significantly enriched sequence motifs

identified[35]. In parallel, we also identified nuclear receptor motifs (i.e., PR sequence motifs) enriched in

the SMAD1/5 binding sites (Supplement Figure 1D,E). From pathway enrichment analysis, we

demonstrate that genes with SMAD1/5 and PR bound at the promoter regions are enriched for key

pathways in directing the decidualization process, such as WNT and relaxin signaling pathways. Future

studies can benefit from analyzing binding events beyond the promoter regions. Profiling the PR genome occupancy in the SMAD1/5 deficient mice would provide an interesting perspective to reevaluate the major regulatory roles of SMAD1/5 in mediating uterine transcriptomes. In this study, we determined the overlapped transcriptional control between SMAD1/5 and PR at the gene level, and functionally validated the regulatory effect at the transcript level in a human stromal cell decidualization model. While we observe a subset of peak representations that do not overlap at the base pair level in the promoter regions, future functional screenings at the promoter level, such as luciferase reporter assays to assess transcriptional co-activation by SMAD1/5 and PR, will advance this study.

SMADs are known to recruit co-repressors (*i.e.*, Ski[105]) or co-activators (*i.e.*, p300[106]) to inhibit or activate target gene transcription, less is known about their cell-specific co-factors that confer the precise spatial-temporal control over binding activities to target genes. Our study highlights the potential co-factors by integrating both genomic and transcriptomic data to delineate signaling crosstalks that are responsible for maintaining tissue homeostasis, especially in the female reproductive tract.

Since mice only undergo decidualization upon embryo implantation whilst human stromal cells undergo cyclic decidualization in each menstrual cycle in response to rising levels of progesterone[36], asynchronous gene responses may occur in comparison between mouse models and human cells. However, cellular transformation during decidualization is conserved between mice and humans[32], which makes findings in the mouse models a valuable and transferable resource to be evaluated in human tissues. Accordingly, our functional validation studies were performed using human endometrial stromal cells induced to decidualize *in vitro* for four days, which models the early phases of decidualization. Additional transcriptomic studies of the SMAD1/5 perturbations in human endometrial stromal cells will be of great resource in understanding the entire SMAD1/5 regulomes in humans.

In summary, our findings and those of others indicate that SMAD1 and SMAD5 not only are signal transducers for BMP signaling pathways, but also engage extensively in the crosstalk with PR signaling pathways. While P4 responses are critical for early pregnancy establishment, abnormal P4 responses are implicated in diseases such as endometriosis and endometrial cancers[107-110]. Hence, our results which show that BMP and P4 signaling pathways synergize within the endometrium; these key pathways can shed light on the endometrial contribution to conditions that impact reproductive health in women, including early pregnancy loss, endometriosis, and endometrial cancer. Furthermore, we anticipate that the SMAD1/5 knock-in tagged transgenic mouse models developed herein will be useful for studying BMP/SMAD1/5 signaling pathways in other reproductive and non-reproductive tract tissues in the body.

DATA AVAILABILITY

Sequencing data and analyses are deposited in the Gene Expression Omnibus under accession number GSE237975. (Reviewer token: gludeooirhczxcj)

SUPPLEMENTARY DATA

Supplementary Table 1: DNA sequences used in the study.

Supplementary Table 2: Direct target genes of SMAD1/5.

Supplementary Table 3: Results of Gene Ontology enrichment analysis of shared direct target genes of SMAD1/5 from the up-targets.

Supplementary Table 4: Results of Gene Ontology enrichment analysis of shared direct target genes of SMAD1/5 from the down-targets.

Supplementary Table 5: Results of KEGG pathway enrichment analysis for shared genes bound by SMAD1, SMAD5, and PR.

Supplementary Table 6: Results of motif analysis results for shared peaks between SMAD1, SMAD5 and PR.

Supplementary Table 7: Gene overlapping results of SMAD1/5 and SOX17.

Supplementary Table 8: Gene overlapping results of SMAD1/5 and NR2F2.

Supplementary File 1: Complete list of motif enrichment analysis from the up-targets and down-targets for SMAD1.

Supplementary File 2: Complete list of motif enrichment analysis from the up-targets and down-targets for SMAD5.

Supplementary Figures: Supplementary Figures 1-3

AUTHOR CONTRIBUTIONS

Zian Liao: Conceptualization, Formal analysis, Methodology, Validation, Writing—original draft. Suni Tang: Data curation, Investigation, Methodology, Visualization, Writing—review & editing. Kaori Nozawa: Data curation, Investigation, Methodology, Writing—review & editing. Keisuke Shimada: Data curation, Methodology, Writing—review & editing. Masahito Ikawa: Project administration, Resources. Diana Monsivais: Conceptualization, Investigation, Supervision, Funding acquisition, Writing – review & editing. Martin M. Matzuk: Conceptualization, Supervision, Funding acquisition, Writing – review & editing.

ACKNOWLEDGEMENTS

FUNDING

This research is supported by Eunice Kennedy Shriver National Institute of Child Health and Human

Development grants HD105800 (D.M.), HD096057 (D.M.), HD032067 (M.M.M.), and HD110038

(M.M.M.). DM is supported by a Next Gen Pregnancy Award from the Burroughs Wellcome Fund

(NGP10125).

CONFLICT OF INTEREST

No conflict of interest is declared by the authors.

REFERENCE
653   BIBLIOGRAPHY AND REFERENCES
654

1   Boivin, J., Bunting, L., Collins, J. A. & Nygren, K. G. International estimates of infertility
prevalence and treatment-seeking: potential need and demand for infertility medical care. *Hum
Reprod* **22**, 1506-1512, doi:10.1093/humrep/dem046 (2007).
2   Martinez, G. M., Daniels, K. & Febo-Vazquez, I. Fertility of Men and Women Aged 15-44 in the
United States: National Survey of Family Growth, 2011-2015. *Natl Health Stat Report*, 1-17
(2018).
3   Wang, X. *et al.* Conception, early pregnancy loss, and time to clinical pregnancy: a population-
based prospective study. *Fertil Steril* **79**, 577-584, doi:10.1016/s0015-0282(02)04694-0 (2003).
4   Chang, H., Brown, C. W. & Matzuk, M. M. Genetic analysis of the mammalian transforming
growth factor-beta superfamily. *Endocr Rev* **23**, 787-823, doi:10.1210/er.2002-0003 (2002).
5   Monsivais, D., Matzuk, M. M. & Pangas, S. A. The TGF-beta Family in the Reproductive Tract.
Cold Spring Harb Perspect Biol **9**, doi:10.1101/cshperspect.a022251 (2017).
6   Lee, K. Y. *et al.* Bmp2 is critical for the murine uterine decidual response. *Mol Cell Biol* **27**, 5468-
5478, doi:10.1128/MCB.00342-07 (2007).
7   Monsivais, D. *et al.* BMP7 Induces Uterine Receptivity and Blastocyst Attachment. *Endocrinology*
158, 979-992, doi:10.1210/en.2016-1629 (2017).
8   Nagashima, T. *et al.* BMPR2 is required for postimplantation uterine function and pregnancy
maintenance. *J Clin Invest* **123**, 2539-2550, doi:10.1172/JCI65710 (2013).
9   Monsivais, D. *et al.* Endometrial receptivity and implantation require uterine BMP signaling
through an ACVR2A-SMAD1/SMAD5 axis. *Nat Commun* **12**, 3386, doi:10.1038/s41467-021-
23571-5 (2021).
10   Matzuk, M. M., Kumar, T. R. & Bradley, A. Different phenotypes for mice deficient in either
activins or activin receptor type II. *Nature* **374**, 356-360, doi:10.1038/374356a0 (1995).
11   Clementi, C. *et al.* Activin-like kinase 2 functions in peri-implantation uterine signaling in mice and
humans. *PLoS Genet* **9**, e1003863, doi:10.1371/journal.pgen.1003863 (2013).
12   Monsivais, D. *et al.* Uterine ALK3 is essential during the window of implantation. *Proc Natl Acad
Sci U S A* **113**, E387-395, doi:10.1073/pnas.1523758113 (2016).
13   Teklenburg, G. *et al.* Natural selection of human embryos: decidualizing endometrial stromal cells
serve as sensors of embryo quality upon implantation. *PLoS One* **5**, e10258,
doi:10.1371/journal.pone.0010258 (2010).

14   Brosens, J. J. *et al.* Uterine selection of human embryos at implantation. *Sci Rep* **4**, 3894,
doi:10.1038/srep03894 (2014).
15   Weimar, C. H. *et al.* Endometrial stromal cells of women with recurrent miscarriage fail to
discriminate between high- and low-quality human embryos. *PLoS One* **7**, e41424,
doi:10.1371/journal.pone.0041424 (2012).
16   Salker, M. S. *et al.* Deregulation of the serum- and glucocorticoid-inducible kinase SGK1 in the
endometrium causes reproductive failure. *Nat Med* **17**, 1509-1513, doi:10.1038/nm.2498 (2011).
17   Garrido-Gomez, T. *et al.* Defective decidualization during and after severe preeclampsia reveals
a possible maternal contribution to the etiology. *Proc Natl Acad Sci U S A* **114**, E8468-E8477,
doi:10.1073/pnas.1706546114 (2017).
18   Garrido-Gomez, T. *et al.* Preeclampsia: a defect in decidualization is associated with deficiency of
Annexin A2. *Am J Obstet Gynecol* **222**, 376 e371-376 e317, doi:10.1016/j.ajog.2019.11.1250
(2020).
19   Shimada, K. *et al.* ARMC12 regulates spatiotemporal mitochondrial dynamics during
spermiogenesis and is required for male fertility. *Proc Natl Acad Sci U S A* **118**,
doi:10.1073/pnas.2018355118 (2021).
20   Fang, B. *et al.* Circadian enhancers coordinate multiple phases of rhythmic gene transcription in
vivo. *Cell* **159**, 1140-1152, doi:10.1016/j.cell.2014.10.022 (2014).
21   Skene, P. J. & Henikoff, S. An efficient targeted nuclease strategy for high-resolution mapping of
DNA binding sites. *Elife* **6**, doi:10.7554/eLife.21856 (2017).
22   Meers, M. P., Tenenbaum, D. & Henikoff, S. Peak calling by Sparse Enrichment Analysis for
CUT&RUN chromatin profiling. *Epigenetics Chromatin* **12**, 42, doi:10.1186/s13072-019-0287-4
(2019).
23   Kent, W. J., Zweig, A. S., Barber, G., Hinrichs, A. S. & Karolchik, D. BigWig and BigBed: enabling
browsing of large distributed datasets. *Bioinformatics* **26**, 2204-2207,
doi:10.1093/bioinformatics/btq351 (2010).
24   Heinz, S. *et al.* Simple combinations of lineage-determining transcription factors prime cis-
regulatory elements required for macrophage and B cell identities. *Mol Cell* **38**, 576-589,
doi:10.1016/j.molcel.2010.05.004 (2010).
25   Yu, G., Wang, L. G. & He, Q. Y. ChIPseeker: an R/Bioconductor package for ChIP peak
annotation, comparison and visualization. *Bioinformatics* **31**, 2382-2383,
doi:10.1093/bioinformatics/btv145 (2015).
26   Ramirez, F. *et al.* deepTools2: a next generation web server for deep-sequencing data analysis.
Nucleic Acids Res **44**, W160-165, doi:10.1093/nar/gkw257 (2016).
27   Newman, A. M. *et al.* Determining cell type abundance and expression from bulk tissues with
digital cytometry. *Nat Biotechnol* **37**, 773-782, doi:10.1038/s41587-019-0114-2 (2019).
28   Initiative, C. Z.   (2023).
29   Warren, L. A. *et al.* Analysis of menstrual effluent: diagnostic potential for endometriosis. *Mol Med*
24, 1, doi:10.1186/s10020-018-0009-6 (2018).
30   Nayyar, A. *et al.* Menstrual Effluent Provides a Novel Diagnostic Window on the Pathogenesis of
Endometriosis. *Front Reprod Health* **2**, 3, doi:10.3389/frph.2020.00003 (2020).
31   Martinez-Aguilar, R. *et al.* Menstrual blood-derived stromal cells modulate functional properties of
mouse and human macrophages. *Sci Rep* **10**, 21389, doi:10.1038/s41598-020-78423-x (2020).
32   Gellersen, B. & Brosens, J. J. Cyclic decidualization of the human endometrium in reproductive
health and failure. *Endocr Rev* **35**, 851-905, doi:10.1210/er.2014-1045 (2014).
33   Magro-Lopez, E. & Munoz-Fernandez, M. A. The Role of BMP Signaling in Female Reproductive
System Development and Function. *Int J Mol Sci* **22**, doi:10.3390/ijms222111927 (2021).
34   Li, Q. *et al.* Bone morphogenetic protein 2 functions via a conserved signaling pathway involving
Wnt4 to regulate uterine decidualization in the mouse and the human. *J Biol Chem* **282**, 31725-
31732, doi:10.1074/jbc.M704723200 (2007).
35   Rubel, C. A. *et al.* Research resource: Genome-wide profiling of progesterone receptor binding in
the mouse uterus. *Mol Endocrinol* **26**, 1428-1442, doi:10.1210/me.2011-1355 (2012).

36   Ramathal, C. Y., Bagchi, I. C., Taylor, R. N. & Bagchi, M. K. Endometrial decidualization: of mice
and men. *Semin Reprod Med* **28**, 17-26, doi:10.1055/s-0029-1242989 (2010).
37   Wang, S. *et al.* Target analysis by integration of transcriptome and ChIP-seq data with BETA. *Nat*
Protoc **8**, 2502-2515, doi:10.1038/nprot.2013.150 (2013).
38   Ciciro, Y. & Sala, A. MYB oncoproteins: emerging players and potential therapeutic targets in
human cancer. *Oncogenesis* **10**, 19, doi:10.1038/s41389-021-00309-y (2021).
39   Swartz, C. D., Afshari, C. A., Yu, L., Hall, K. E. & Dixon, D. Estrogen-induced changes in IGF-I,
Myb family and MAP kinase pathway genes in human uterine leiomyoma and normal uterine
smooth muscle cell lines. *Mol Hum Reprod* **11**, 441-450, doi:10.1093/molehr/gah174 (2005).
40   Akaogi, K. *et al.* KLF4 suppresses estrogen-dependent breast cancer growth by inhibiting the
transcriptional activity of ERalpha. *Oncogene* **28**, 2894-2902, doi:10.1038/onc.2009.151 (2009).
41   Timblin, G. A. & Schlissel, M. S. Ebf1 and c-Myb repress rag transcription downstream of Stat5
during early B cell development. *J Immunol* **191**, 4676-4687, doi:10.4049/jimmunol.1301675
(2013).
42   Pan, H., Zhu, L., Deng, Y. & Pollard, J. W. Microarray analysis of uterine epithelial gene
expression during the implantation window in the mouse. *Endocrinology* **147**, 4904-4916,
doi:10.1210/en.2006-0140 (2006).
43   Fagerberg, L. *et al.* Analysis of the human tissue-specific expression by genome-wide integration
of transcriptomics and antibody-based proteomics. *Mol Cell Proteomics* **13**, 397-406,
doi:10.1074/mcp.M113.035600 (2014).
44   Ma, W. *et al.* MAX deficiency impairs human endometrial decidualization through down-regulating
OSR2 in women with recurrent spontaneous abortion. *Cell Tissue Res* **388**, 453-469,
doi:10.1007/s00441-022-03579-z (2022).
45   Rankin, S. A., Gallas, A. L., Neto, A., Gomez-Skarmeta, J. L. & Zorn, A. M. Suppression of Bmp4
signaling by the zinc-finger repressors Osr1 and Osr2 is required for Wnt/beta-catenin-mediated
lung specification in Xenopus. *Development* **139**, 3010-3020, doi:10.1242/dev.078220 (2012).
46   Du, H. & Taylor, H. S. The Role of Hox Genes in Female Reproductive Tract Development, Adult
Function, and Fertility. *Cold Spring Harb Perspect Med* **6**, a023002,
doi:10.1101/cshperspect.a023002 (2015).
47   Cakmak, H. & Taylor, H. S. Molecular mechanisms of treatment resistance in endometriosis: the
role of progesterone-hox gene interactions. *Semin Reprod Med* **28**, 69-74, doi:10.1055/s-0029-
1242996 (2010).
48   Taylor, H. S., Bagot, C., Kardana, A., Olive, D. & Arici, A. HOX gene expression is altered in the
endometrium of women with endometriosis. *Hum Reprod* **14**, 1328-1331,
doi:10.1093/humrep/14.5.1328 (1999).
49   He, B., Ni, Z. L., Kong, S. B., Lu, J. H. & Wang, H. B. Homeobox genes for embryo implantation:
From mouse to human. *Animal Model Exp Med* **1**, 14-22, doi:10.1002/ame2.12002 (2018).
50   Illingworth, I. M. *et al.* Desmosomes are reduced in the mouse uterine luminal epithelium during
the preimplantation period of pregnancy: a mechanism for facilitation of implantation. *Biol Reprod*
63, 1764-1773, doi:10.1095/biolreprod63.6.1764 (2000).
51   Paria, B. C., Zhao, X., Das, S. K., Dey, S. K. & Yoshinaga, K. Zonula occludens-1 and E-cadherin
are coordinately expressed in the mouse uterus with the initiation of implantation and
decidualization. *Dev Biol* **208**, 488-501, doi:10.1006/dbio.1999.9206 (1999).
52   Potter, S. W., Gaza, G. & Morris, J. E. Estradiol induces E-cadherin degradation in mouse uterine
epithelium during the estrous cycle and early pregnancy. *J Cell Physiol* **169**, 1-14,
doi:10.1002/(SICI)1097-4652(199610)169:1<1::AID-JCP1>3.0.CO;2-S (1996).
53   Grund, S. & Grummer, R. Direct Cell(-)Cell Interactions in the Endometrium and in Endometrial
Pathophysiology. *Int J Mol Sci* **19**, doi:10.3390/ijms19082227 (2018).
54   Schatz, F., Guzeloglu-Kayisli, O., Arlier, S., Kayisli, U. A. & Lockwood, C. J. The role of decidual
cells in uterine hemostasis, menstruation, inflammation, adverse pregnancy outcomes and
abnormal uterine bleeding. *Hum Reprod Update* **22**, 497-515, doi:10.1093/humupd/dmw004
(2016).

55 Evans, J. *et al.* Fertile ground: human endometrial programming and lessons in health and
disease. *Nat Rev Endocrinol* **12**, 654-667, doi:10.1038/nrendo.2016.116 (2016).
56 Stehlin-Gaon, C. *et al.* All-trans retinoic acid is a ligand for the orphan nuclear receptor ROR beta.
Nat Struct Biol **10**, 820-825, doi:10.1038/nsb979 (2003).
57 Spitzer, T. L. *et al.* Perivascular human endometrial mesenchymal stem cells express pathways
relevant to self-renewal, lineage specification, and functional phenotype. *Biol Reprod* **86**, 58,
doi:10.1095/biolreprod.111.095885 (2012).
58 Yin, Y., Haller, M. E., Chadchan, S. B., Kommagani, R. & Ma, L. Signaling through retinoic acid
receptors is essential for mammalian uterine receptivity and decidualization. *JCI Insight* **6**,
doi:10.1172/jci.insight.150254 (2021).
59 Chang, C. Agonists and Antagonists of TGF-beta Family Ligands. *Cold Spring Harb Perspect Biol*
8, doi:10.1101/cshperspect.a021923 (2016).
60 Fullerton, P. T., Jr., Monsivais, D., Kommagani, R. & Matzuk, M. M. Follistatin is critical for mouse
uterine receptivity and decidualization. *Proc Natl Acad Sci U S A* **114**, E4772-E4781,
doi:10.1073/pnas.1620903114 (2017).
61 Li, Q. *et al.* The antiproliferative action of progesterone in uterine epithelium is mediated by
Hand2. *Science* **331**, 912-916, doi:10.1126/science.1197454 (2011).
62 Filant, J., DeMayo, F. J., Pru, J. K., Lydon, J. P. & Spencer, T. E. Fibroblast growth factor
receptor two (FGFR2) regulates uterine epithelial integrity and fertility in mice. *Biol Reprod* **90**, 7,
doi:10.1095/biolreprod.113.114496 (2014).
63 Lim, W., Bae, H., Bazer, F. W. & Song, G. Stimulatory effects of fibroblast growth factor 2 on
proliferation and migration of uterine luminal epithelial cells during early pregnancy. *Biol Reprod*
96, 185-198, doi:10.1095/biolreprod.116.142331 (2017).
64 Russo, L. A. *et al.* Regulated expression of matrix metalloproteinases, inflammatory mediators,
and endometrial matrix remodeling by 17beta-estradiol in the immature rat uterus. *Reprod Biol*
Endocrinol **7**, 124, doi:10.1186/1477-7827-7-124 (2009).
65 Tenvergert, E., Gillespie, M. W., Kingma, J. & Klasen, H. Abortion attitudes, 1984-1987-1988:
effects of item order and dimensionality. *Percept Mot Skills* **74**, 627-642,
doi:10.2466/pms.1992.74.2.627 (1992).
66 Hayashi, K. *et al.* Wnt genes in the mouse uterus: potential regulation of implantation. *Biol*
Reprod **80**, 989-1000, doi:10.1095/biolreprod.108.075416 (2009).
67 Yang, M. *et al.* Spatiotemporal insight into early pregnancy governed by immune-featured stromal
cells. *Cell* **186**, 4271-4288 e4224, doi:10.1016/j.cell.2023.08.020 (2023).
68 Garcia-Alonso, L. *et al.* Mapping the temporal and spatial dynamics of the human endometrium in
vivo and in vitro. *Nat Genet* **53**, 1698-1711, doi:10.1038/s41588-021-00972-2 (2021).
69 PrabhuDas, M. *et al.* Immune mechanisms at the maternal-fetal interface: perspectives and
challenges. *Nat Immunol* **16**, 328-334, doi:10.1038/ni.3131 (2015).
70 Guimaraes-Young, A., Neff, T., Dupuy, A. J. & Goodheart, M. J. Conditional deletion of Sox17
reveals complex effects on uterine adenogenesis and function. *Dev Biol* **414**, 219-227,
doi:10.1016/j.ydbio.2016.04.010 (2016).
71 Hirate, Y. *et al.* Mouse Sox17 haploinsufficiency leads to female subfertility due to impaired
implantation. *Sci Rep* **6**, 24171, doi:10.1038/srep24171 (2016).
72 Wang, X. *et al.* SOX17 regulates uterine epithelial-stromal cross-talk acting via a distal enhancer
upstream of Ihh. *Nat Commun* **9**, 4421, doi:10.1038/s41467-018-06652-w (2018).
73 Hollnagel, A., Oehlmann, V., Heymer, J., Ruther, U. & Nordheim, A. Id genes are direct targets of
bone morphogenetic protein induction in embryonic stem cells. *J Biol Chem* **274**, 19838-19845,
doi:10.1074/jbc.274.28.19838 (1999).
74 Miyazono, K. & Miyazawa, K. Id: a target of BMP signaling. *Sci STKE* **2002**, pe40,
doi:10.1126/stke.2002.151.pe40 (2002).
75 Holloran, S. M. *et al.* Reciprocal fine-tuning of progesterone and prolactin-regulated gene
expression in breast cancer cells. *Mol Cell Endocrinol* **511**, 110859,
doi:10.1016/j.mce.2020.110859 (2020).

76 Dinh, D. T. *et al.* Progesterone receptor mediates ovulatory transcription through RUNX
transcription factor interactions and chromatin remodelling. *Nucleic Acids Res*,
doi:10.1093/nar/gkad271 (2023).
77 Vasquez, Y. M. *et al.* FOXO1 regulates uterine epithelial integrity and progesterone receptor
expression critical for embryo implantation. *PLoS Genet* **14**, e1007787,
doi:10.1371/journal.pgen.1007787 (2018).
78 Shi, J. W. *et al.* An IGF1-expressing endometrial stromal cell population is associated with human
decidualization. *BMC Biol* **20**, 276, doi:10.1186/s12915-022-01483-0 (2022).
79 Suzuki, A. *et al.* Nanog binds to Smad1 and blocks bone morphogenetic protein-induced
differentiation of embryonic stem cells. *Proc Natl Acad Sci U S A* **103**, 10294-10299,
doi:10.1073/pnas.0506945103 (2006).
80 Shi, X., Bai, S., Li, L. & Cao, X. Hoxa-9 represses transforming growth factor-beta-induced
osteopontin gene transcription. *J Biol Chem* **276**, 850-855, doi:10.1074/jbc.M005955200 (2001).
81 Shi, X., Yang, X., Chen, D., Chang, Z. & Cao, X. Smad1 interacts with homeobox DNA-binding
proteins in bone morphogenetic protein signaling. *J Biol Chem* **274**, 13711-13717,
doi:10.1074/jbc.274.19.13711 (1999).
82 Feng, X. H. & Derynck, R. Specificity and versatility in tgf-beta signaling through Smads. *Annu*
Rev Cell Dev Biol **21**, 659-693, doi:10.1146/annurev.cellbio.21.022404.142018 (2005).
83 Massague, J., Seoane, J. & Wotton, D. Smad transcription factors. *Genes Dev* **19**, 2783-2810,
doi:10.1101/gad.1350705 (2005).
84 Derynck, R. & Budi, E. H. Specificity, versatility, and control of TGF-beta family signaling. *Sci*
Signal **12**, doi:10.1126/scisignal.aav5183 (2019).
85 Lee, D. K. *et al.* Suppression of ERalpha activity by COUP-TFII is essential for successful
implantation and decidualization. *Mol Endocrinol* **24**, 930-940, doi:10.1210/me.2009-0531 (2010).
86 Vasquez, Y. M. *et al.* FOXO1 is required for binding of PR on IRF4, novel transcriptional regulator
of endometrial stromal decidualization. *Mol Endocrinol* **29**, 421-433, doi:10.1210/me.2014-1292
(2015).
87 Wu, M., Chen, G. & Li, Y. P. TGF-beta and BMP signaling in osteoblast, skeletal development,
and bone formation, homeostasis and disease. *Bone Res* **4**, 16009, doi:10.1038/boneres.2016.9
(2016).
88 van Weerd, J. H. & Christoffels, V. M. The formation and function of the cardiac conduction
system. *Development* **143**, 197-210, doi:10.1242/dev.124883 (2016).
89 Whitman, M. Smads and early developmental signaling by the TGFbeta superfamily. *Genes Dev*
12, 2445-2462, doi:10.1101/gad.12.16.2445 (1998).
90 Middlebrook, B. S., Eldin, K., Li, X., Shivasankaran, S. & Pangas, S. A. Smad1-Smad5 ovarian
conditional knockout mice develop a disease profile similar to the juvenile form of human
granulosa cell tumors. *Endocrinology* **150**, 5208-5217, doi:10.1210/en.2009-0644 (2009).
91 Pangas, S. A. *et al.* Conditional deletion of Smad1 and Smad5 in somatic cells of male and
female gonads leads to metastatic tumor development in mice. *Mol Cell Biol* **28**, 248-257,
doi:10.1128/MCB.01404-07 (2008).
92 Rodriguez, A. *et al.* SMAD Signaling Is Required for Structural Integrity of the Female
Reproductive Tract and Uterine Function During Early Pregnancy in Mice. *Biol Reprod* **95**, 44,
doi:10.1095/biolreprod.116.139477 (2016).
93 Retting, K. N., Song, B., Yoon, B. S. & Lyons, K. M. BMP canonical Smad signaling through
Smad1 and Smad5 is required for endochondral bone formation. *Development* **136**, 1093-1104,
doi:10.1242/dev.029926 (2009).
94 McReynolds, L. J., Gupta, S., Figueroa, M. E., Mullins, M. C. & Evans, T. Smad1 and Smad5
differentially regulate embryonic hematopoiesis. *Blood* **110**, 3881-3890, doi:10.1182/blood-2007-
04-085753 (2007).
95 Consortium, E. P. An integrated encyclopedia of DNA elements in the human genome. *Nature*
489, 57-74, doi:10.1038/nature11247 (2012).
96 Chen, J. *et al.* Single-molecule dynamics of enhanceosome assembly in embryonic stem cells.
Cell **156**, 1274-1285, doi:10.1016/j.cell.2014.01.062 (2014).

97 Swinstead, E. E. *et al.* Steroid Receptors Reprogram FoxA1 Occupancy through Dynamic
Chromatin Transitions. *Cell* **165**, 593-605, doi:10.1016/j.cell.2016.02.067 (2016).
98 Chen, C. H. *et al.* Determinants of transcription factor regulatory range. *Nat Commun* **11**, 2472,
doi:10.1038/s41467-020-16106-x (2020).
99 Arvey, A., Agius, P., Noble, W. S. & Leslie, C. Sequence and chromatin determinants of cell-type-
specific transcription factor binding. *Genome Res* **22**, 1723-1734, doi:10.1101/gr.127712.111
(2012).
100 Tang, S., Cope, D. I., Vasquez, Y. M. & Monsivais, D. BMP/SMAD1/5 Signaling in the
Endometrial Epithelium Is Essential for Receptivity and Early Pregnancy. *Endocrinology* **163**,
doi:10.1210/endocr/bqac043 (2022).
101 Fuentealba, L. C. *et al.* Integrating patterning signals: Wnt/GSK3 regulates the duration of the
BMP/Smad1 signal. *Cell* **131**, 980-993, doi:10.1016/j.cell.2007.09.027 (2007).
102 Labbe, E., Letamendia, A. & Attisano, L. Association of Smads with lymphoid enhancer binding
factor 1/T cell-specific factor mediates cooperative signaling by the transforming growth factor-
beta and wnt pathways. *Proc Natl Acad Sci U S A* **97**, 8358-8363, doi:10.1073/pnas.150152697
(2000).
103 Zavadil, J., Cermak, L., Soto-Nieves, N. & Bottinger, E. P. Integration of TGF-beta/Smad and
Jagged1/Notch signalling in epithelial-to-mesenchymal transition. *EMBO J* **23**, 1155-1165,
doi:10.1038/sj.emboj.7600069 (2004).
104 Qiu, T., Grizzle, W. E., Oelschlager, D. K., Shen, X. & Cao, X. Control of prostate cell growth:
BMP antagonizes androgen mitogenic activity with incorporation of MAPK signals in Smad1.
EMBO J **26**, 346-357, doi:10.1038/sj.emboj.7601499 (2007).
105 Luo, K. *et al.* The Ski oncoprotein interacts with the Smad proteins to repress TGFbeta signaling.
Genes Dev **13**, 2196-2206, doi:10.1101/gad.13.17.2196 (1999).
106 Pouponnot, C., Jayaraman, L. & Massague, J. Physical and functional interaction of SMADs and
p300/CBP. *J Biol Chem* **273**, 22865-22868, doi:10.1074/jbc.273.36.22865 (1998).
107 Brosens, J. J. & Gellersen, B. Death or survival--progesterone-dependent cell fate decisions in
the human endometrial stroma. *J Mol Endocrinol* **36**, 389-398, doi:10.1677/jme.1.02060 (2006).
108 Yilmaz, B. D. & Bulun, S. E. Endometriosis and nuclear receptors. *Hum Reprod Update* **25**, 473-
485, doi:10.1093/humupd/dmz005 (2019).
109 Janzen, D. M. *et al.* Progesterone receptor signaling in the microenvironment of endometrial
cancer influences its response to hormonal therapy. *Cancer Res* **73**, 4697-4710,
doi:10.1158/0008-5472.CAN-13-0930 (2013).
110 MacLean, J. A., 2nd & Hayashi, K. Progesterone Actions and Resistance in Gynecological
Disorders. *Cells* **11**, doi:10.3390/cells11040647 (2022).
929

Figure 1

Figure 1: Mouse models with global HA tagged SMAD1 and PA tagged SMAD5 proteins. A-B)

Schematic approaches for generating Smad1HA/HA and Smad5PA/PA knock-in mouse lines. Sanger

sequencing of the genotyping results are included as validation of knock-in sequence. Black and blue

boxes indicate untranslated and coding regions, respectively. **C-D)** Immunoblot (IB) analysis of the

immunoprecipitation (IP) of HA tagged SMAD1 and PA tagged SMAD5 proteins from different tissues of

the tagged mouse lines. Wild type (WT) mice were used as negative controls. Antibodies used for IB and

IP are as labeled. Targeted bands of SMAD1 and SMAD5 are indicated by red arrows.

Figure 2

Figure 2: Genomic profiling of SMAD1 and SMAD5 binding sites during decidualization *in vivo*. A)
Diagram outlining experimental approaches for tissue collection, processing, and CUT&RUN. B)
Heatmaps and summary plots showing the enrichment of SMAD1 and SMAD5 binding peaks from one
exemplary replicate. Clustering was conducted using k-means algorithm. The colors in the summary
plots correspond to clusters labeled in the heatmap below. **C-D**) Feature distribution of the annotated
peaks for the SMAD1 (C) binding sites and SMAD5 (D) binding sites.

Figure 3

Figure 3: **SMAD1 and SMAD5 show unique direct target genes during early pregnancy. A-B**) Venn

diagrams showing the shared and unique direct up-target genes (A) and down-target genes (B) of

SMAD1, SMAD5 Numbers indicate genes numbers. **C-D**) Motif enrichment analysis from the up-targets

and down-targets for SMAD1 (C) and SMAD5 (D). **E-F**) Dot plot showing Gene Ontology enrichment

950   analysis of shared direct target genes of SMAD1/5 from the up-targets (E) and the down-targets (F),

951   respectively. Dot size represents the gene ratio in the enriched categories compared to background

952   genes, dot colors reflect P-value.

Figure 4

Figure 4: Direct target genes of SMAD1/5 mediate uterine homeostasis. **A**) Bar graph of normalized

Fragments Per Kilobase of transcript per Million mapped reads (FPKM) of downregulated transcripts in

the Control and SMAD1/5 cKO groups as indicated by the label. Histograms represent average +/- SEM

of experiments uteri from Control mice (N=3) and SMAD1/5 cKO mice (N=4). Analyzed by an unpaired t-

test. **B**) Integrative Genomics Viewer (IGV) track view of SMAD1, SMAD5 binding activities. Gene loci

are as indicated in the figure, genomic coordinates are annotated in mm10. **C**) Bar graph of FPKM of up

regulated transcripts in the Control and SMAD1/5 cKO groups as indicated by the label. **D**) IGV track

view of SMAD1, SMAD5 binding activities. Gene loci are as indicated in the figure, genomic coordinates

are annotated in mm10. **E**) Dot plot showing the gene expression pattern of the key SMAD1/5 direct

target genes in different cell types from published human endometrium single-cell RNA-seq dataset.

Figure 5

Figure 5: SMAD1 and SMAD5 co-regulate PR target genes. A) Heatmaps and summary plots

showing the enrichment comparison between SMAD1, SMAD5 and PR binding peaks from one

exemplary replicate. Clustering was conducted using k-means algorithm. The colors in the summary

968   plots correspond to clusters labeled in the heatmap below. **B**) Dot plot showing KEGG pathway

969   enrichment analysis for shared genes bound by SMAD1, SMAD5, and PR. **C**) IGV track view of SMAD1,

970   SMAD5 and PR binding activities. Gene loci are as indicated in the figure, genomic coordinates are

971   annotated in mm10. **D**) Table of motif analysis results for shared peaks between SMAD1, SMAD5 and

972   PR, with P-value and motif annotation specified for each motif.

Figure 6

Figure 6: SMAD1 and SMAD5 are required for PR responses during decidualization of human endometrial stromal cells. A) Schematic approach and timeline outlining in vitro decidualization for endometrial stromal cells (EnSCs). **B-C**) RT-qPCR results showing mRNA levels of *PRL*, *IGFBP1*, *FOXO1*, *RORB* and *KLF15* after SMAD1/5 perturbation using siRNAs. Data are normalized to siCTL-Veh

979   for visualization. Bar graphs represent average +/- SEM of experiments on cells from three different

980   individuals with technical triplicates. Analyzed by a One-Way ANOVA test with post hoc Tukey test.

Supplement Figure 1:

Supplement Figure 1: Gene numbers with SMAD1/5 promoter binding activities and motif

analysis of SMAD1/5 peaks. A) Venn diagrams showing the shared and unique genes bound by

SMAD1 or SMAD5 in the +/- 3kb region of the promoter regions. **B**) Cell type compositions in the

deconvoluted RNA-seq from control mice (Control 1-3) and SMAD1/5 PR-Cre mice (KO1-4). Numbers

plotted in the table are absolute values that reflects the absolute proportion of each cell type in the given

sample. A larger number indicates a higher proportion of that cell type in the mixture. **C**) Venn diagrams

showing the shared and unique genes bound by SMAD1, SMAD5 or PR in the +/- 3kb of the promoter

regions. **D-E**) Table of motif analysis results for unique peaks for SMAD1(D) and SMAD5 (E), with P-

value and motif annotation specified for each motif. **F)** Venn diagrams showing the shared and unique

genes bound by SMAD1, SMAD5 or SOX17 in the +/- 25kb regions near transcription start site. **G)** Venn

diagrams showing the shared and unique genes bound by SMAD1, SMAD5 or NR2F2 in the +/- 25kb

regions near transcription start site.

Supplement Figure 2:

995 **Supplement Figure 2: Knockdown effect validation of SMAD1/5 perturbation. A-B**) RT-qPCR

996 results showing mRNA levels of SMAD1(A) and SMAD5 (B) after siRNA treatments in both Veh and EPC

997 conditions. Data are normalized to siCTL for visualization. Bar graphs represent average +/- SEM of

998 experiments on cells from three different individuals with technical triplicates. Analyzed by a One-Way

999 ANOVA test with post hoc Tukey test.

1000

Supplement Figure 3:

A

B

1001

1002   **Supplement Figure 3: Genotype of the knock-in mouse lines**. **A**) Schematic design of the genotype

1003   primers for Smad1HA/HA and Smad5PA/PA mouse lines. **B**) Exemplary gel electrophoresis of PCR

1004   products derived from homozygous knock-in mice, heterozygous mice, and WT mice using genotyping

1005   primers.

Uncropped Western Blots for Figure 1