The genomic landscape of metallic color variation in ground beetles

Yi-Ming Weng[1,6]*, David H. Kavanaugh[2], Bryan Rubio-Perez[3], Jad Salman[3], Mikhail A.

Kats[3,4,5], Sean D. Schoville[1]

1. University of Wisconsin-Madison, Department of Entomology, Madison, WI, USA

2. California Academy of Sciences, Department of Entomology, San Francisco, CA, USA

3. University of Wisconsin-Madison, Department of Electrical and Computer Engineering,

Madison, WI, USA

4. University of Wisconsin-Madison, Department of Physics, Madison, WI, USA

5. University of Wisconsin-Madison, Department of Material Science and Engineering, Madison,

WI, USA.

6. Okinawa Institute of Science and Technology Graduate University, Okinawa, Japan

* Corresponding author: Yi-Ming Weng

Email:

weng76111@gmail.com

ORCID identifier:

Yi-Ming Weng: 0000-0002-8243-5061

David H. Kavanaugh: 0000-0001-8777-9771

Mikhail A. Kats: 0000-0003-4897-4720

Sean D. Schoville: 0000-0001-7364-434X

Abstract

The metallic color variation of beetles is a spectacular feature that has inspired diverse human cultures. However, little is known about the genetic basis of this trait or its ecological importance. In this study, we characterize the geographical distribution, optical mechanism, genetic basis, and ecological and evolutionary importance of metallic color variation in the *Nebria ingens* complex, an alpine ground beetle in the Sierra Nevada, California. We find that elytral color varies continuously across two allopatric species (from black *N. ingens* to green *N. riversi*), with hybrid populations showing intermediate coloration, and we demonstrate that the metallic color is generated from multilayer reflectors in the epicuticle of the elytra. By applying association mapping in natural populations (wild-GWAS) using high-density genotype variants, we identify five promising candidate genes covarying with metallic variation, with known roles in cuticle formation and pigmentation pathways. This finding, together with a significant correlation between color variation and water availability, suggests that metallic variation evolves as a local adaptation to environmental variation in the *N. ingens* complex.

Keywords: metallic color, Wild-GWAS, phenotypic evolution, functional ecology, *Nebria*, insects

Introduction

The impression created by bright metallic coloration in beetle wings is so stunning that diverse human cultures have used them in art and jewelry ('beetlewing art') for thousands of years (Ratcliffe, 2006; Rivers et al., 2003). This remarkable phenotypic trait is all the more impressive when its tremendous variation among beetles (order Coleoptera), a biodiverse group including more than 500,000 species, is recognized to have repeatedly evolved (Seago et al., 2009). The optical mechanism of metallic color in insects is well known, generated by the internal structure of the exoskeleton (Andersen, 2009). In Coleoptera, the most prevalent internal structure is a multilayered reflector in the epicuticle or exocuticle, where the layers with alternating refractive index reflect light in a specific range of wavelengths (Barry et al., 2020; Seago et al., 2009). For species exhibiting multilayered reflectors, variations may involve differences in saturation (chroma), hue, or both. The thickness of the alternating layers primarily governs the hue, with thicker layers producing longer peak wavelengths and thinner layers resulting in shorter ones (Kurachi et al., 2002). Meanwhile, saturation can be influenced by the number of layers or the overall tanning level of the cuticle. As a result of the discovery of the optical mechanism of metallic colors in insects, these microstructures have become an important resource for bioinspired materials (Biró et al., 2010; Biró & Vigneron, 2011; Colusso et al., 2017).

Although metallic coloration is known to be polymorphic in numerous beetle species (Kozlov et al., 2022; Strickland et al., 2019), little is known about the genetic basis of this phenotypic trait. Compared to beetles, the genetic basis of wing coloration in butterflies has been more extensively studied and could provide hypotheses for testing the roles of similar genes in regulating multilayer reflectors in beetles. For example, transcription factors such as *optix*, *Antennapedia*, *aristaless1*, *bric-a-brac* (*bab*), and *spalt* are involved in various pigmentation pathways and in regulating the photonic microstructures on butterfly scales (Ficarrotta et al., 2022; Livraghi et al., 2025; Mazo-Vargas et al., 2022; Prakash et al., 2022; Reed et al., 2020; Thayer et al., 2020; Zhang et al., 2017). These transcription factors may also regulate the synthesis, transport, or deposition of melanized layers in the beetle cuticle. On the other hand, genes involved in the tyrosine-mediated tanning pathway such as *ebony*, *tan*, *arylalkylamine N-acetyltransferase* (*aaNAT*), *laccase2*, *yellow*, *pale* (*tyrosine hydroxylase*), *Dopa decarboxylase* (*DDC*), along with potential transcription factors such as *pannier* (*pnr*) and *POU domain motif 3*

(*pdm3*), are responsible for the production and distribution of melanin pigments and could

therefore contribute to patterning the multilayer reflector (Arakane et al., 2016a; Niimi & Ando,

2021).

Since studies of diverse beetle species have shown that metallic color variation is typically

additively determined by a small number of genetic loci (Favila et al., 2000; Fujiyama &

Arimoto, 1988; Gotoh & Lavine, 2014; Mossakowski & Paarmann, 2014a; Vasconcellos-Neto,

1988), and only infrequently dependent on genotype-environment interactions (Davis et al.,

2008), it should be possible to identify the genes underlying the metallic color variation by

association approaches (Wellenreuther & Hansson, 2016).

In this study, we aim to investigate the genetic basis of metallic color from the multilayer

reflector in the cuticle of alpine ground beetles, the *Nebria ingens* species complex, by using

whole genome sequence data and association mapping in natural populations (wild-GWAS). The

N. ingens species complex is endemic to the Sierra Nevada in California, where it is found in the

high elevation nival zone that is covered by snow throughout most of the year. Two species with

divergent color in their elytra (appearing metallic green or black to human perception), *N. ingens*

Horn and *N. riversi* Van Dyke, are geographically isolated, but form a broad admixture zone of

intermediately colored individuals (Schoville et al., 2012; Weng et al., 2021) (**Figure 1**). The

morphologically variable and genetically admixed populations make this species complex a great

study system to apply wild-GWAS approaches (Buerkle & Lexer, 2008; Lindtke et al., 2013).

Replicate admixed populations disrupt linkage disequilibrium across the genome, increasing

power to isolate causative loci underlying the divergent phenotypes of the two species.

Furthermore, the nearly continuous color variation improves the statistical power of association

analyses. Leveraging this system, we identify candidate for the genetic basis of metallic color in

beetles which could be tested in the future expression regulation or gene editing approaches.

Materials and Methods

Animal, genotyping, and color variation

Adult beetles of the *N. ingens* complex (n=383) were collected from 27 populations,

encompassing the entire distribution range of the species complex (**Figure 1**). Whole-genome

resequencing data from 374 beetles were generated in our previous study using the GATK

variant calling pipeline (Van der Auwera & O'Connor, 2020; Weng et al., 2024). After removing

samples lacking color phenotypic data and loci with minor allele frequency less than 0.05 using

vcftools v0.1.16 (Danecek et al., 2011), we performed GWAS using a genotype matrix with 355

beetles and 526,717 SNP+indel variants. To quantify optical variation, we measured the

reflectance spectra from the elytral surface of 368 beetles using a hyperspectral camera (Specim

IQ). The average spectrum was measured from each pixel in the brightest elytral region (8-600

pixels, median=190 pixels) to represent the individual reflectance spectrum. The mean

reflectance spectrum was then used to calculate the CIELAB color for each elytron (Robertson,

1977) using the CIE standard illuminant D65 with a custom MATLAB script. Chroma was

calculated as the distance from the neutral gray point (0,0) in the a*b* plane to represent the

color saturation of the elytra.

Cuticular structure of the metallic hue

The elytra of three beetles, representing *N. ingens*, *N. riversi*, and a genetically admixed hybrid

lineage, were used to examine the cuticular microstructure with transmission electron

microscopy (TEM). We used a FEI Tecnai T12 microscope in the School of Medicine and Public

Health at UW-Madison. The samples were stored in -20°C with 95% ethanol before use. The

images with clear structure were selected to measure the thickness of the epicuticle. We took 13-

30 vertical transects to repeat measurements of thickness for each specimen using ImageJ

(Abràmoff et al., 2004). Counts and thickness of layers were used to predict the reflectance

spectra of each specimen using a multilayer reflectance calculator provided by Filmetrics[®]

Corporation, with the refractive indices of electron dense and electron lucent being 1.7 and 1.4,

respectively (**Figure 2**). These indices are consistent across different beetle taxa (Bernard &

Miller, 1968; Kurachi et al., 2002). The calculated reflectance spectrum was subsequently used

to compare with the observed spectra from the same individual beetle, except for the green

elytrum for which we compared the calculated spectrum to other bright green beetles from the

same population, to verify the source of the metallic color (**Figure 2**).

Genome-wide association mapping

To identify SNP and indel markers associated with the color phenotype, we performed GWAS

using latent factor mixed models (LFMM) and partial redundancy analysis (pRDA) implemented

using R package lfmm and vegan, respectively (Caye et al., 2019; Oksanen et al., 2025). In

LFMM analysis, four latent factors were specified to account for population structure, which

includes a pattern of isolation by distance across seven ancestral population clusters (Weng,

Kavanaugh, et al., 2021). The regularized least squares estimate of causal variants were obtained

using both ridge and lasso regression. To reduce the false-discovery rate (FDR), we first

calibrated the p values with genomic inflation factors (GIF), then adjusted the calibrated p values

using the Benjamini-Hochberg multiple-testing procedure. Variants with the adjusted p value

less than 0.01 in ridge or lasso regression were considered candidates of LFMM. In pRDA

analysis, genotype data was first transformed into plink format (.raw) using vcftools v0.1.16

(Danecek et al., 2011). To adjust the linear effects of chroma on genotype while accounting for

the variation from the population structure, the first four PC axes calculated with SNPRelate

were included as conditioning variables (Zheng & Zheng, 2013). Variants falling outside of four

standard deviations of first RDA axis loading are considered candidate variants for chroma. The

intersection of candidate variants from the two analyses were used to find the intersecting genes

from the published genome using bedtools v2.28 (Quinlan, 2014; Weng, Francoeur, et al., 2021).

Gene intersection was defined by variants falling into the genetic region and their 2 kilobase

upstream (5') flanking region, as regulatory elements could be located thousands of base pairs

from the 5' flanking region. Finally, we assessed to annotated function of the candidate genes by

referencing their best blast hit against NCBI nr database and InterPro function using

InterProScan (Blum et al., 2025; Camacho et al., 2009; Jones et al., 2014), and compared

candidate genes with our previously published genes with signal of selective sweeps from

OmegaPlus to identify overlapping genes potentially involved in both color variation and

evolutionary adaptation (Alachiotis et al., 2012; Weng et al., 2024).

Testing for environmental selection on color variation

Because color variation is a key diagnostic character separating two species in the *N. ingens*

complex, it may be associated with adaptation to local environments. We investigated the

correlation between color variation, water loss rate of beetles, and the snow water equivalent

(SWE) in the habitats of the sampled populations. To quantify the water loss rate, 87 beetles

were placed individually in a tube in a sealed container, which contained lithium chloride (LiCl)

as a desiccant to maintain a relative humidity between 20 and 30 percent. The temperature was

controlled at 5°C during the experiment, which is similar to the natural foraging temperature in

the field. The water loss rate was calculated by weighing beetles before and after 24 hours of

desiccation. Habitat SWE data were extracted from published Sierra Nevada snow analysis

results using the R package raster v3.6-3, with further methodological details provided in our

previous study (Margulis et al., 2016; Weng et al., 2024). Spearman's rank correlation coefficient

(ρ) was used to assess the correlation since the data were non-normal in distribution.

Results

Continuous variation and the mechanism controlling metallic color

We examined 368 beetles from 27 populations across the distribution of the species complex. In

general, the reflectance spectra of green beetle elytra have a peak of 500 to 600 nm wavelength,

while beetles with black elytra have a lower reflectance amplitude and loss of the characteristic

reflectance peak resulting in the green color (**Figure 2**). By converting the reflectance spectra to

the CIELAB color space (L*a*b*) using the CIE standard illuminant D65, we found that many,

but not all beetles identified as *N. riversi* reflect a green hue, while most beetles identified as *N.*

ingens or as hybrids reflect a dark green to black hue (**Figure 1**). The color variation is

continuously distributed and thus likely to be a quantitative trait encoded by multiple genes.

From the elytral cross section, TEM images of green, dark green, and black elytra

demonstrate variation in the inner epicuticle (cuticulin layer), which is most pronounced in the

green elytra (**Figure 2**). There are 1-3 electron dense layers 50-100 nm thick (darker layers in the

images due to reduced permeability for the electron beam), alternating with similar-width

electron lucent layers, serving as a multilayered reflector (**Figure S1**). In the dark green elytra,

only one electron dense layer is found, and no clear multilayer reflector was observed in the

black elytra throughout the examined regions. Calculated spectra based on the thickness of the

multilayer reflectors generally matches the visual color and observed spectra, with more

alternating layers producing a stronger amplitude of light at a peak wavelength between 500 and

600 nm, roughly corresponding to the color green (**Figure 2**).

Color associated genes and their roles in color determination

211   Among the 526,717 scanned variants, 892 (0.17%) passed the adjusted *p*-value threshold

at 0.01 in ridge or lasso regression models and 573 (0.11%) fall outside of 4th standard deviation

of first RDA axis loading distribution, where 195 (0.037%) variants are found shared in both

candidate lists and considered candidate variants. These candidate variants intersect 127 genes

including 65 genes with functional annotations (**Table S1**) and five with known functions in

different pathways of insect cuticle formation and pigmentation (**Figure 3**). These include

yellow-e and *pdm3* in the tyrosine-mediated tanning pathway, *aristaless* in the ommochrome

pathway, *chico* (encoding insulin receptor substrate (IRS) protein) in the Insulin/IGF signaling

pathways through PI3K/Akt, and a gene encoding Pyrokinin-1 receptors-like protein (*PK1*-like),

a type of G-protein coupled receptor in the pyrokinin signaling pathway (Matsumoto et al., 1990;

Shakhmantsir et al., 2014). All five genes have intersecting variants within intragenic regions,

and the variants in *pdm3* and *chico* are in the coding region. Finally, the candidate genes *chico*,

PK1-like, and small G protein signaling modulator 3 (*GPSM3*) homolog were previously

identified in genome-wide scans for selective sweeps.

Correlation between color variation, water loss rate, and SWE

To test the physiological and ecological function of metallic color, we performed pairwise

correlation analysis using chroma, water loss rate, and seasonal water availability in the nival

zone (represented by measurements of snow water equivalent in September, SWE) (**Figure 4**).

The results showed a significant positive correlation between beetle color and SWE (ρ=0.45,

95% CI=0.35 to 0.53; *p*= 2.2e-16), with higher color chroma occurring in more snow-rich

habitats, and a significant positive correlation between water loss rate and SWE (ρ=0.26, 95%

CI=0.06 to 0.44; *p*=0.014). We also found a positive trend between color variation and water loss

rate but that was not significant (ρ=0.21, 95% CI=-0.06 to 0.45; *p*=0.12).

Discussion

Despite a long-standing interest in the metallic color of insects, distinct knowledge gaps in our

understanding of color variation remain. Beyond the lack of knowledge for the most biodiverse

animal group, the beetles, gaps include the nebulous genetic basis of metallic color variation. By
applying wild-GWAS to an alpine ground beetle species complex with admixed populations, we
found promising candidate genes that are statistically associated with the metallic variation
generated from multilayer reflectors in the epicuticle and functionally involved in cuticular
structure and melanization. These include genes from several known pathways involved in
butterfly color patterns.

Genetic basis of the metallic color in the N. ingens complex
Color variation can be categorized by optical mechanisms which vary across insect lineages. For
instance, in Coleoptera, iridescent color generated by three-dimensional photonic structures has
been found in weevils and longhorn beetles, whereas multilayer reflectors are widespread in
distantly related lineages of jewel beetles (Buprestidae), ground beetles (Carabidae), leaf beetles
(Chrysomelidae), and scarab beetles (Scarabaeidae) (Seago et al., 2009, 2019). Although
structural layering may be common to the ancestors of beetles, structural coloration has been
repeatedly gained, lost and modified (Seago et al., 2019). Yet very little is known about how
genetically complex this phenotypic variation is across beetles.
In this study, we found that a multilayer reflector is responsible for the metallic color
variation of the *N. ingens* complex. This is directly supported by the presence of the layers with
alternating refractive index in the epicuticle, with their estimated reflectance spectra matching
the elytral colors and the observed spectra (**Figure 2**). Such resemblance of the calculated and
observed reflectance spectra has been used to determine the optical mechanism of metallic color
in many other insects (Bermúdez-Ureña et al., 2020; Kurachi et al., 2002; J. Lee et al., 2020;
Luna et al., 2013; Noyes et al., 2007; Parker et al., 1998; Pasteels et al., 2016). In the *N. ingens*
complex, the amplitude of the peak between 500 and 600 nm in the reflectance spectrum is
directly correlated with the number of alternating layers in the epicuticle, representing a form of
saturation variation characterized by changes in amplitude at specific wavelengths caused by
differing numbers of layers with consistent thickness. It is structurally distinct from phenotypes
where the reflectance peak shifts across wavelengths due to differences in layer thickness and
therefore should be regarded as a separate category of color variation. In nature, this type of
saturation variation seen in the *N. ingens* complex appears to be less common but has been

reported in different beetle families (Gotoh & Lavine, 2014; Mossakowski & Paarmann, 2014b;
Stanbrook et al., 2021).

Knowing that the metallic color variation in the *N. ingens* complex is regulated by multilayer
reflectors, we conducted a wild-GWAS to identify genetic variants associated with this color
variation. We discovered 65 genes containing variants that significantly correlate with the
chroma of the elytra, including five genes with known roles in cuticle pigmentation processes
(**Figure 3**). Among these five genes, *yellow-e* and *pdm3* in the tyrosine-mediated tanning
pathway are most likely to contribute to the color variation in the *N. ingens* complex, as the
multilayer reflectors are primarily composed of alternating layers with different degree of
melanin deposition (Hamza & Zahran, 2023; Onelli et al., 2017; Parker et al., 1998). Yellow
protein (dopachrome conversion enzyme) from the *yellow* gene family advances melanin
biosynthesis, a vital biochemical pathway altering the insect cuticle profile by converting
tyrosine into pigments (e.g., melanin and dopamine-melanin) (Arakane et al., 2016b; Noh et al.,
2016). This gene has been identified as a key mediator of the melanin biosynthesis pathway
across multiple insect species (Gompel et al., 2005; Li & Christensen, 2011; Noh et al., 2015;
Zhang, Martin, et al., 2017). On the other hand, the transcription factor *pdm3* is known for its
role in regulating neural development, but it has also been identified as a repressor of abdominal
pigmentation in *Drosophila melanogaster*, potentially through co-expression with other melanin
biosynthesis transcription factors such as *Abdominal-B* and *doublesex* (David et al., 2025;
Hughes et al., 2023; Kim & Cho, 2020; Rogers et al., 2014; Yassin et al., 2016). However, the
function of *pdm3* orthologs in beetle (Coleoptera) species remains unexplored, highlighting the
need for further studies that investigate its role in beetle cuticle pigmentation. In contrast, *chico*
and *PK1-like* genes are indirectly involved in cuticle pigmentation through the Insulin/IGF and
pyrokinin signaling pathways, respectively (Matsumoto et al., 1990; Shakhmantsir et al., 2014).
Although their involvement is indirect, signals of positive selection from the selective sweeps
have been previously identified in these genes in the *N. ingens* complex, underscoring the
importance of further investigating their roles in the pigmentation of beetle elytra (Weng et al.,
2024). Finally, *aristaless*, a transcription factor known to influence appendage formation and
scale color in butterflies, has been identified as a candidate gene (Bayala et al., 2023). However,
aristaless is involved in ommochrome pigment production through the ommochrome pathway.

Since melanin is the dominant pigment in insect cuticles, we consider *aristaless* the least likely candidate among the five. In summary, we identified five genes linked to beetle elytra color, with *yellow-e* and *pdm3* most likely influencing melanin-based pigmentation. *Chico* and *PK1*-like show indirect roles and signs of selection, while *aristaless*, involved in a different pigment pathway, is the least likely candidate.

The evolutionary and ecological role of metallic color in the <u>N. ingens</u> complex

The ecological function of metallic color has been considered an important characteristic in biotic interactions, both within and between species (Clusella-Trullas & Nielsen, 2020). In beetles, sexual selection, mimicry, camouflage, or aposematism are widely discussed functions for metallic color (Kjernsmo et al., 2022; Lee et al., 2018; Vulinec, 1997; Xu et al., 2010). But just how relevant biotic interactions are for the diversity of color patterns in beetles remains unclear (Ospina-Rozo et al., 2024; Schultz & Hadley, 1987). Other studies have shown that metallic color may instead result from adaptation to abiotic environmental factors (Davis et al., 2008; Keinath et al., 2020). In particular, pigmentation from the melanization pathway in the epidermal and serosal cuticle is thought to be associated with desiccation resistance (Bai et al., 2022; Farnesi et al., 2017; Wang et al., 2021). In the subgenus *Catonebria*, which includes the *N. ingens* complex, metallic color variation has been gained and lost many times with a general trend that metallic species occur in cooler and wetter regions compared to related species with darker, less or non-metallic cuticle. For example, among species of the *Nebria gebleri* species complex (Kavanaugh et al., 2021), those from more northern and coastal regions (northern Cascade Range and northern Rocky Mountains; namely, *N. gebleri* Dejean and *N. cascadensis* Kavanaugh) are more vividly metallic than those from the Sierra Nevada and other southern ranges in California (i.e., *N. albimontis* Kavanaugh, *N. rathvoni* LeConte, and *N. siskiyouensis* Kavanaugh). The same general pattern is seen among members of the *meanyi* species complex as well as in other closely related *Nebria* subgenera. To investigate this phenomenon, we explored the possibility that metallic variation might be associated with water availability in *N. ingens* habitats, with the expectation that the black individuals could have higher fitness in drier and warmer habitats. Evidence from other beetle species has shown a relationship between the *yellow* gene, cuticle melanization, and water retention ability (Noh et al., 2015). This leads to the

hypothesis that the loss of metallic color, as a byproduct of melanin accumulation, is the result of

adaptation to arid environments. Our results show correlations between chroma and habitat SWE,

as well as between water loss rate and habitat SWE, indicating that less melanized beetles are

more commonly found in habitats with lower snow covers (**Figure 4A & 4B**). Moreover,

individuals inhabiting these drier environments tend to exhibit better water retention ability.

However, the correlation between chroma and water loss rate is moderate and not statistically

significant (**Figure 4C**), suggesting that additional factors could be contributing to variation in

water retention ability.

Study limitations and future directions

While our wild-GWAS approach successfully identified genetic variants associated with beetle

elytra color variation, several limitations may have influenced the strength of the association

signals. Despite efforts to minimize noise, challenges in accurately quantifying color, such as

measurement errors due to varying specimen conditions, could have introduced variability into

the data. Additionally, nearly half of the candidate genes predicted from the reference genome

lack functional annotation, which means that important genes may have been overlooked in our

analysis. Nonetheless, we identified five promising candidate genes that warrant further

functional investigation to better understand their roles in producing the metallic coloration in

beetles.

Acknowledgments

We thank Filmetrics® for granting permission to use the reflectance calculator. We thank the

National Parks of the United States, including Yosemite, Sequoia and Kings Canyon National

Parks, for providing permission to conduct research in the parks. This research was financially

supported by the Valentine Eastern Sierra Reserve (UC Santa Barbara Natural Reserve System)

through VESR Graduate Student Funding and the Ministry of Education of the Republic of

China (Taiwan) through the Government Scholarship to Study Abroad to Y.M.W. We thank the

PhD committee of Y.M.W., consisting of Daniel Young, Carol Lee, Prashant Sharma, and Jun

Zhu, for providing suggestions regarding this study. The authors have no conflicts of interests to

declare.

Author Contributions: YMW, DHK, and SDS developed the original idea; YMW and SDS

designed the experiments. The quantification of color variation was conducted by YMW, BRP,

and JS and other analyses were performed by YMW. YMW and SDS wrote the first draft of the

manuscript and DHK, BRP, JS, and MAK provided comments to improve the manuscript.

References

References

Abràmoff, M. D., Magalhães, P. J., & Ram, S. J. (2004). Image processing with ImageJ. *Biophotonics*
International, *11*(7), 36–42.

Alachiotis, N., Stamatakis, A., & Pavlidis, P. (2012). OmegaPlus: A scalable tool for rapid detection of
selective sweeps in whole-genome datasets. *Bioinformatics*, *28*(17), 2274–2275.
https://doi.org/10.1093/bioinformatics/bts419

Andersen, S. O. (2009). Exoskeleton. In *Encyclopedia of insects* (pp. 339–342). Elsevier.

Arakane, Y., Noh, M. Y., Asano, T., & Kramer, K. J. (2016a). Tyrosine Metabolism for Insect Cuticle
Pigmentation and Sclerotization. In E. Cohen & B. Moussian (Eds.), *Extracellular Composite Matrices in*
Arthropods (pp. 165–220). Springer International Publishing. https://doi.org/10.1007/978-3-319-40740-
1_6

Arakane, Y., Noh, M. Y., Asano, T., & Kramer, K. J. (2016b). Tyrosine Metabolism for Insect Cuticle
Pigmentation and Sclerotization. In E. Cohen & B. Moussian (Eds.), *Extracellular Composite Matrices in*
Arthropods (pp. 165–220). Springer International Publishing. https://doi.org/10.1007/978-3-319-40740-
1_6

Bai, T.-T., Pei, X.-J., Liu, T.-X., Fan, Y.-L., & Zhang, S.-Z. (2022). Melanin synthesis genes BgTH and BgDdc
affect body color and cuticle permeability in Blattella germanica. *Insect Science*, *29*(6), 1552–1568.
https://doi.org/10.1111/1744-7917.13024

Barry, M. A., Berthier, V., Wilts, B. D., Cambourieux, M.-C., Bennet, P., Pollès, R., Teytaud, O., Centeno,
E., Biais, N., & Moreau, A. (2020). Evolutionary algorithms converge towards evolved biological photonic
structures. *Scientific Reports*, *10*(1), 12024. https://doi.org/10.1038/s41598-020-68719-3

Bayala, E. X., VanKuren, N., Massardo, D., & Kronforst, M. R. (2023). Aristaless1 has a dual role in
appendage formation and wing color specification during butterfly development. *BMC Biology*, *21*(1),
100. https://doi.org/10.1186/s12915-023-01601-6

Bermúdez-Ureña, E., Kilchoer, C., Lord, N. P., Steiner, U., & Wilts, B. D. (2020). Structural Diversity with
Varying Disorder Enables the Multicolored Display in the Longhorn Beetle *Sulawesiella rafaelae*.
iScience, *23*(7), 101339. https://doi.org/10.1016/j.isci.2020.101339

Bernard, G. D., & Miller, W. H. (1968). Interference filters in the corneas of Diptera. *Investigative*
Ophthalmology, *7*(4), 416–434.

Biró, L. P., Kertész, K., Horváth, E., Márk, G. I., Molnár, G., Vértesy, Z., Tsai, J.-F., Kun, A., Bálint, Z., &
Vigneron, J. P. (2010). Bioinspired artificial photonic nanoarchitecture using the elytron of the beetle
Trigonophorus rothschildi varians as a 'blueprint.' *Journal of the Royal Society Interface*, *7*(47), 887–894.

Biró, L. P., & Vigneron, J. P. (2011). Photonic nanoarchitectures in butterflies and beetles: Valuable
sources for bioinspiration. *Laser & Photonics Reviews*, *5*(1), 27–51.
https://doi.org/10.1002/lpor.200900018

Blum, M., Andreeva, A., Florentino, L. C., Chuguransky, S. R., Grego, T., Hobbs, E., Pinto, B. L., Orr, A.,
Paysan-Lafosse, T., Ponamareva, I., Salazar, G. A., Bordin, N., Bork, P., Bridge, A., Colwell, L., Gough, J.,

Haft, D. H., Letunic, I., Llinares-López, F., … Bateman, A. (2025). InterPro: The protein sequence
classification resource in 2025. *Nucleic Acids Research*, *53*(D1), D444–D456.
https://doi.org/10.1093/nar/gkae1082

Buerkle, C. A., & Lexer, C. (2008). Admixture as the basis for genetic mapping. *Trends in Ecology &*
Evolution, *23*(12), 686–694. https://doi.org/10.1016/j.tree.2008.07.008

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., & Madden, T. L. (2009).
BLAST+: Architecture and applications. *BMC Bioinformatics*, *10*(1), 421. https://doi.org/10.1186/1471-
2105-10-421

Caye, K., Jumentier, B., Lepeule, J., & François, O. (2019). LFMM 2: Fast and Accurate Inference of Gene-
Environment Associations in Genome-Wide Studies. *Molecular Biology and Evolution*, *36*(4), 852–860.
https://doi.org/10.1093/molbev/msz008

Clusella-Trullas, S., & Nielsen, M. (2020). The evolution of insect body coloration under changing
climates. *Current Opinion in Insect Science*, *41*, 25–32. https://doi.org/10.1016/j.cois.2020.05.007

Colusso, E., Perotto, G., Wang, Y., Sturaro, M., Omenetto, F., & Martucci, A. (2017). Bioinspired stimuli-
responsive multilayer film made of silk–titanate nanocomposites. *Journal of Materials Chemistry C*,
5(16), 3924–3931. https://doi.org/10.1039/C7TC00149E

Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., Handsaker, R. E., Lunter, G.,
Marth, G. T., Sherry, S. T., McVean, G., & Durbin, R. (2011). The variant call format and VCFtools.
Bioinformatics, *27*(15), 2156–2158. https://doi.org/10.1093/bioinformatics/btr330

David, J. R., Denis, B., Francelle, P., Lemaire, A., Das, A., Mohanty, S., Gibert, P., & Yassin, A. (2025).
Phenotypic plasticity as a function of genetic polymorphism: Thermal dominance reversal in Drosophila
species with contrasting melanism (p. 2025.05.30.657080). bioRxiv.
https://doi.org/10.1101/2025.05.30.657080

Davis, A. L. v., Brink, D. J., Scholtz, C. H., Prinsloo, L. C., & Deschodt, C. M. (2008). Functional implications
of temperature-correlated colour polymorphism in an iridescent, scarabaeine dung beetle. *Ecological*
Entomology, *33*(6), 771–779. https://doi.org/10.1111/j.1365-2311.2008.01033.x

Farnesi, L. C., Vargas, H. C. M., Valle, D., & Rezende, G. L. (2017). Darker eggs of mosquitoes resist more
to dry conditions: Melanin enhances serosal cuticle contribution in egg resistance to desiccation in
Aedes, Anopheles and Culex vectors. *PLOS Neglected Tropical Diseases*, *11*(10), e0006063.
https://doi.org/10.1371/journal.pntd.0006063

Favila, M. E., Ruiz-Lizarraga, G., & Nolasco, J. (2000). Inheritance of a Red Cuticular Color Mutation in the
Scarab Beetle CANTHON CYANELLUS CYANELLUS LeConte (Coleoptera: Scarabaeidae). *The Coleopterists*
Bulletin, *54*(4), 541–545. https://doi.org/10.1649/0010-
065X(2000)054%255B0541:IOARCC%255D2.0.CO;2

Ficarrotta, V., Hanly, J. J., Loh, L. S., Francescutti, C. M., Ren, A., Tunström, K., Wheat, C. W., Porter, A. H.,
Counterman, B. A., & Martin, A. (2022). A genetic switch for male UV iridescence in an incipient species
pair of sulphur butterflies. *Proceedings of the National Academy of Sciences*, *119*(3), e2109255118.
https://doi.org/10.1073/pnas.2109255118

Fujiyama, S., & Arimoto, K. (1988). Genetics of the two colour forms of Chrysolina aurichalcea
(Mannerheim) (Coleoptera: Chrysomelidae) and their gene frequencies in two mountainous areas of
central Honshu, Japan. In P. Jolivet, E. Petitpierre, & T. H. Hsiao (Eds.), *Biology of Chrysomelidae* (pp.
205–215). Springer Netherlands. https://doi.org/10.1007/978-94-009-3105-3_12

Gompel, N., Prud'homme, B., Wittkopp, P. J., Kassner, V. A., & Carroll, S. B. (2005). Chance caught on the
wing: Cis-regulatory evolution and the origin of pigment patterns in Drosophila. *Nature*, *433*(7025), 481–
487. https://doi.org/10.1038/nature03235

Gotoh, H., & Lavine, L. C. (2014). Genetic Control of Color Polymorphism in the Stag Beetle Cyclommatus
metallifer Boisduval (Coleoptera: Lucanidae). *The Coleopterists Bulletin*, *68*(2), 209–213.

Hamza, A., & Zahran, N. (2023). Melanin pigmentation variations in the larval cuticle of almond moth,
Ephestia cautella caused by gamma radiation. *Scientific Reports*, *13*, 21012.
https://doi.org/10.1038/s41598-023-48154-w

Hughes, J. T., Williams, M. E., Rebeiz, M., & Williams, T. M. (2023). Widespread cis- and trans-regulatory
evolution underlies the origin, diversification, and loss of a sexually dimorphic fruit fly pigmentation
trait. *Journal of Experimental Zoology Part B: Molecular and Developmental Evolution*, *340*(2), 143–161.
https://doi.org/10.1002/jez.b.23068

Jones, P., Binns, D., Chang, H.-Y., Fraser, M., Li, W., McAnulla, C., McWilliam, H., Maslen, J., Mitchell, A.,
Nuka, G., Pesseat, S., Quinn, A. F., Sangrador-Vegas, A., Scheremetjew, M., Yong, S.-Y., Lopez, R., &
Hunter, S. (2014). InterProScan 5: Genome-scale protein function classification. *Bioinformatics*, *30*(9),
1236–1240. https://doi.org/10.1093/bioinformatics/btu031

Kavanaugh, D. H., Maddison, D. R., Simison, W. B., Schoville, S. D., Schmidt, J., Faille, A., Moore, W.,
Pflug, J. M., Archambeault, S. L., Hoang, T., & Chen, J.-Y. (2021). Phylogeny of the supertribe Nebriitae
(Coleoptera, Carabidae) based on analyses of DNA sequence data. *ZooKeys*, *1044*, 41–152.
https://doi.org/10.3897/zookeys.1044.62245

Keinath, S., Frisch, J., Müller, J., Mayer, F., & Rödel, M.-O. (2020). Spatio-Temporal Color Differences
Between Urban and Rural Populations of a Ground Beetle During the Last 100 Years. *Frontiers in Ecology*
and Evolution, *7*. https://www.frontiersin.org/articles/10.3389/fevo.2019.00525

Kim, Y., & Cho, K.-O. (2020). POU domain motif3 (Pdm3) induces wingless (wg) transcription and is
essential for development of larval neuromuscular junctions in Drosophila. *Scientific Reports*, *10*(1), 517.
https://doi.org/10.1038/s41598-020-57425-9

Kjernsmo, K., Lim, A. M., Middleton, R., Hall, J. R., Costello, L. M., Whitney, H. M., Scott-Samuel, N. E., &
Cuthill, I. C. (2022). Beetle iridescence induces an avoidance response in naïve avian predators. *Animal*
Behaviour, *188*, 45–50. https://doi.org/10.1016/j.anbehav.2022.04.005

Kozlov, M. V., Oudendijk, Z., Forsman, A., Lanta, V., Barclay, M. V. L., Gusarov, V. I., Gustafsson, B.,
Huang, Z.-Z., Kruglova, O. Y., Marusik, Y. M., Mikhailov, Y. E., Mutanen, M., Schneider, A., Sekerka, L.,
Sergeev, M. E., Zverev, V., & Zvereva, E. L. (2022). Climate shapes the spatiotemporal variation in color
morph diversity and composition across the distribution range of Chrysomela lapponica leaf beetle.
Insect Science, *29*(3), 942–955. https://doi.org/10.1111/1744-7917.12966

Kurachi, M., Takaku, Y., Komiya, Y., & Hariyama, T. (2002). The origin of extensive colour polymorphism in Plateumaris sericea (Chrysomelidae, Coleoptera). *Naturwissenschaften*, *89*(7), 295–298. https://doi.org/10.1007/s00114-002-0332-0

Lee, C. Y., Yo, S. P., Clark, R. W., Hsu, J. Y., Liao, C. P., Tseng, H. Y., & Huang, W. S. (2018). The role of different visual characters of weevils signalling aposematism to sympatric lizard predators. *Journal of Zoology*, *306*(1), 36–47. https://doi.org/10.1111/jzo.12567

Lee, J., Terlier, T., Jang, Y. J., Lee, K.-B., & Lee, Y. (2020). Structural colors and physical properties of elytra in the jewel beetle, Chrysochroa fulgidissima, using surface analytical techniques. *Surface and Interface Analysis*, *52*(10), 656–664. https://doi.org/10.1002/sia.6807

Li, J., & Christensen, B. M. (2011). Biological Function of Insect Yellow Gene Family. In T. Liu & L. Kang (Eds.), *Recent Advances in Entomological Research: From Molecular Biology to Pest Management* (pp. 121–131). Springer. https://doi.org/10.1007/978-3-642-17815-3_7

Lindtke, D., González-Martínez, S. C., Macaya-Sanz, D., & Lexer, C. (2013). Admixture mapping of quantitative traits in Populus hybrid zones: Power and limitations. *Heredity*, *111*(6), 474–485. https://doi.org/10.1038/hdy.2013.69

Livraghi, L., Hanly, J. J., Loh, L. S., Henry, A., Keck, C., Shirey, V. M., Tsai, C.-C., Yu, N., Belleghem, S. M. V., Roberts, W. M., Boggs, C. L., & Martin, A. (2025). Genetic basis of an adaptive polymorphism controlling butterfly silver iridescence. *Current Biology*, *35*(9), 2154-2163.e7. https://doi.org/10.1016/j.cub.2025.03.028

Luna, A., Macías, D., Skigin, D., Inchaussandague, M., Schinca, D., Gigli, M., & Vial, A. (2013). Characterization of the iridescence-causing multilayer structure of the Ceroglossus suturalis beetle using bio-inspired optimization strategies. *Optics Express*, *21*(16), 19189–19201. https://doi.org/10.1364/OE.21.019189

Margulis, S. A., Cortés, G., Girotto, M., & Durand, M. (2016). *A Landsat-Era Sierra Nevada Snow Reanalysis (1985–2015)*. https://doi.org/10.1175/JHM-D-15-0177.1

Matsumoto, S., Kitamura, A., Nagasawa, H., Kataoka, H., Orikasa, C., Mitsui, T., & Suzuki, A. (1990). Functional diversity of a neurohormone produced by the suboesophageal ganglion: Molecular identity of melanization and reddish colouration hormone and pheromone biosynthesis activating neuropeptide. *Journal of Insect Physiology*, *36*(6), 427–432. https://doi.org/10.1016/0022-1910(90)90060-S

Mazo-Vargas, A., Langmüller, A. M., Wilder, A., van der Burg, K. R. L., Lewis, J. J., Messer, P. W., Zhang, L., Martin, A., & Reed, R. D. (2022). Deep cis-regulatory homology of the butterfly wing pattern ground plan. *Science*, *378*(6617), 304–308. https://doi.org/10.1126/science.abi9407

Mossakowski, D., & Paarmann, W. (2014a). Elytral surface structure in Poecilus lepidus (Coleoptera: Carabidae): What about the nature of its inheritance? *EJE*, *111*(5), 601–607. https://doi.org/10.14411/eje.2014.087

Mossakowski, D., & Paarmann, W. (2014b). Elytral surface structure in Poecilus lepidus (Coleoptera: Carabidae): What about the nature of its inheritance? *EJE*, *111*(5), 601–607. https://doi.org/10.14411/eje.2014.087

Niimi, T., & Ando, T. (2021). *Evo-devo* of wing colour patterns in beetles. *Current Opinion in Genetics &*
Development, *69*, 97–102. https://doi.org/10.1016/j.gde.2021.02.007

Noh, M. Y., Kramer, K. J., Muthukrishnan, S., Beeman, R. W., Kanost, M. R., & Arakane, Y. (2015). Loss of
function of the *yellow-e* gene causes dehydration-induced mortality of adult *Tribolium castaneum*.
Developmental Biology, *399*(2), 315–324. https://doi.org/10.1016/j.ydbio.2015.01.009

Noh, M. Y., Muthukrishnan, S., Kramer, K. J., & Arakane, Y. (2016). Cuticle formation and pigmentation in
beetles. *Current Opinion in Insect Science*, *17*, 1–9. https://doi.org/10.1016/j.cois.2016.05.004

Noyes, J. A., Vukusic, P., & Hooper, I. R. (2007). Experimental method for reliably establishing the
refractive index of buprestid beetle exocuticle. *Optics Express*, *15*(7), 4351–4358.
https://doi.org/10.1364/OE.15.004351

Oksanen, J., Simpson, G. L., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., O'Hara, R. B., Solymos,
P., Stevens, M. H. H., Szoecs, E., Wagner, H., Barbour, M., Bedward, M., Bolker, B., Borcard, D., Borman,
T., Carvalho, G., Chirico, M., Caceres, M. D., … Weedon, J. (2025). *vegan: Community Ecology Package*.
https://vegandevs.github.io/vegan/

Onelli, O. D., Kamp, T. van de, Skepper, J. N., Powell, J., Rolo, T. dos S., Baumbach, T., & Vignolini, S.
(2017). Development of structural colour in leaf beetles. *Scientific Reports*, *7*(1), 1373.
https://doi.org/10.1038/s41598-017-01496-8

Ospina-Rozo, L., Medina, I., Hugall, A., Rankin, K. J., Roberts, N. W., Roberts, A., Mitchell, A., Reid, C. A.
M., Moussalli, A., & Stuart-Fox, D. (2024). Polarization and reflectance are linked to climate, size and
mechanistic constraints in a group of scarab beetles. *Scientific Reports*, *14*(1), 29349.
https://doi.org/10.1038/s41598-024-80325-1

Parker, A. R., Mckenzie, D. R., & Large, M. C. J. (1998). Multilayer Reflectors in Animals Using Green and
Gold Beetles as Contrasting Examples. *Journal of Experimental Biology*, *201*(9), 1307–1313.
https://doi.org/10.1242/jeb.201.9.1307

Pasteels, J. M., Deparis, O., Mouchet, S. R., Windsor, D. M., & Billen, J. (2016). Structural and physical
evidence for an endocuticular gold reflector in the tortoise beetle, *Charidotella ambita*. *Arthropod*
Structure & Development, *45*(6), 509–518. https://doi.org/10.1016/j.asd.2016.10.008

Prakash, A., Finet, C., Banerjee, T. D., Saranathan, V., & Monteiro, A. (2022). *Antennapedia* and *optix*
regulate metallic silver wing scale development and cell shape in *Bicyclus anynana* butterflies. *Cell*
Reports, *40*(1), 111052. https://doi.org/10.1016/j.celrep.2022.111052

Quinlan, A. R. (2014). BEDTools: The Swiss-Army Tool for Genome Feature Analysis. *Current Protocols in*
Bioinformatics, *47*(1), 11.12.1-11.12.34. https://doi.org/10.1002/0471250953.bi1112s47

Ratcliffe, B. C. (2006). Scarab Beetles in Human Culture. *Coleopterists Society Monographs. Patricia*
Vaurie Series, *5*, 85–101.

Reed, R. D., Selegue, J. E., Zhang, L., & Brunetti, C. R. (2020). Transcription factors underlying wing
margin color patterns and pupal cuticle markings in butterflies. *EvoDevo*, *11*(1), 10.
https://doi.org/10.1186/s13227-020-00155-w

Rivers, V. Z., Motte-Florac, É., & Thomas, J. (2003). Emeralds on wing: Jewel beetles in textiles and
adornment. *Insects" in Oral Literature and Traditions*.

Robertson, A. R. (1977). The CIE 1976 Color-Difference Formulae. *Color Research & Application*, *2*(1), 7–
11. https://doi.org/10.1002/j.1520-6378.1977.tb00104.x

Rogers, W. A., Grover, S., Stringer, S. J., Parks, J., Rebeiz, M., & Williams, T. M. (2014). A survey of the
trans-regulatory landscape for *Drosophila melanogaster* abdominal pigmentation. *Developmental*
Biology, *385*(2), 417–432. https://doi.org/10.1016/j.ydbio.2013.11.013

Schoville, S. D., Roderick, G. K., & Kavanaugh, D. H. (2012). Testing the 'Pleistocene species pump' in
alpine habitats: Lineage diversification of flightless ground beetles (Coleoptera: Carabidae: Nebria) in
relation to altitudinal zonation. *Biological Journal of the Linnean Society*, *107*(1), 95–111.
https://doi.org/10.1111/j.1095-8312.2012.01911.x

Schultz, T. D., & Hadley, N. F. (1987). Structural Colors of Tiger Beetles and Their Role in Heat Transfer
through the Integument. *Physiological Zoology*, *60*(6), 737–745.
https://doi.org/10.1086/physzool.60.6.30159990

Seago, A. E., Brady, P., Vigneron, J.-P., & Schultz, T. D. (2009). Gold bugs and beyond: A review of
iridescence and structural colour mechanisms in beetles (Coleoptera). *Journal of The Royal Society*
Interface, *6*(suppl_2), S165–S184. https://doi.org/10.1098/rsif.2008.0354.focus

Seago, A. E., Oberprieler, R., & Saranathan, V. K. (2019). Evolution of Insect Iridescence: Origins of Three-
Dimensional Photonic Crystals in Weevils (Coleoptera: Curculionoidea). *Integrative and Comparative*
Biology, *59*(6), 1664–1672. https://doi.org/10.1093/icb/icz040

Shakhmantsir, I., Massad, N. L., & Kennell, J. A. (2014). Regulation of cuticle pigmentation in drosophila
by the nutrient sensing insulin and TOR signaling pathways. *Developmental Dynamics: An Official*
Publication of the American Association of Anatomists, *243*(3), 393–401.
https://doi.org/10.1002/dvdy.24080

Stanbrook, R. A., Harris, W. E., Wheater, C. P., & Jones, M. (2021). Evidence of phenotypic plasticity
along an altitudinal gradient in the dung beetle Onthophagus proteus. *PeerJ*, *9*, e10798.
https://doi.org/10.7717/peerj.10798

Strickland, L. R., Arias, C. F., Rodriguez, V., Johnston, J. S., McMillan, W. O., & Windsor, D. (2019).
Inheritance, distribution and genetic differentiation of a color polymorphism in Panamanian populations
of the tortoise beetle, Chelymorpha alternans (Coleoptera: Chrysomelidae). *Heredity*, *122*(5), 558–569.
https://doi.org/10.1038/s41437-018-0149-z

Thayer, R. C., Allen, F. I., & Patel, N. H. (2020). Structural color in Junonia butterflies evolves by tuning
scale lamina thickness. *eLife*, *9*, e52187. https://doi.org/10.7554/eLife.52187

Van der Auwera, G. A., & O'Connor, B. D. (2020). *Genomics in the cloud: Using Docker, GATK, and WDL in*
Terra. O'Reilly Media.

Vasconcellos-Neto, J. (1988). Genetics of Chelymorpha cribraria, Cassidinae: Colour patterns and their
ecological meanings. In P. Jolivet, E. Petitpierre, & T. H. Hsiao (Eds.), *Biology of Chrysomelidae* (pp. 217–
232). Springer Netherlands. https://doi.org/10.1007/978-94-009-3105-3_13

Vulinec, K. (1997). Iridescent dung beetles: A different angle. *Florida Entomologist*, 132–141.

Wang, Y., Ferveur, J.-F., & Moussian, B. (2021). Eco-genetics of desiccation resistance in Drosophila.
Biological Reviews, *96*(4), 1421–1440. https://doi.org/10.1111/brv.12709

Wellenreuther, M., & Hansson, B. (2016). Detecting Polygenic Evolution: Problems, Pitfalls, and
Promises. *Trends in Genetics*, *32*(3), 155–164. https://doi.org/10.1016/j.tig.2015.12.004

Weng, Y.-M., Francoeur, C. B., Currie, C. R., Kavanaugh, D. H., & Schoville, S. D. (2021). A high-quality
carabid genome assembly provides insights into beetle genome evolution and cold adaptation.
Molecular Ecology Resources, *21*(6), 2145–2165. https://doi.org/10.1111/1755-0998.13409

Weng, Y.-M., Kavanaugh, D. H., & Schoville, S. D. (2021). Drainage basins serve as multiple glacial refugia
for alpine habitats in the Sierra Nevada Mountains, California. *Molecular Ecology*, *30*(3), 826–843.
https://doi.org/10.1111/mec.15762

Weng, Y.-M., Kavanaugh, D. H., & Schoville, S. D. (2024). Evidence for Admixture and Rapid Evolution
During Glacial Climate Change in an Alpine Specialist. *Molecular Biology and Evolution*, *41*(7), msae130.
https://doi.org/10.1093/molbev/msae130

Xu, M., Seago, A. E., Sutherland, T. D., & Weisman, S. (2010). Dual structural color mechanisms in a
scarab beetle. *Journal of Morphology*, *271*(11), 1300–1305. https://doi.org/10.1002/jmor.10870

Yassin, A., Delaney, E. K., Reddiex, A. J., Seher, T. D., Bastide, H., Appleton, N. C., Lack, J. B., David, J. R.,
Chenoweth, S. F., Pool, J. E., & Kopp, A. (2016). The pdm3 Locus Is a Hotspot for Recurrent Evolution of
Female-Limited Color Dimorphism in Drosophila. *Current Biology: CB*, *26*(18), 2412–2422.
https://doi.org/10.1016/j.cub.2016.07.016

Zhang, L., Martin, A., Perry, M. W., van der Burg, K. R. L., Matsuoka, Y., Monteiro, A., & Reed, R. D.
(2017). Genetic Basis of Melanin Pigmentation in Butterfly Wings. *Genetics*, *205*(4), 1537–1550.
https://doi.org/10.1534/genetics.116.196451

Zhang, L., Mazo-Vargas, A., & Reed, R. D. (2017). Single master regulatory gene coordinates the
evolution and development of butterfly color and iridescence. *Proceedings of the National Academy of*
Sciences, *114*(40), 10707–10712. https://doi.org/10.1073/pnas.1709058114

Zheng, X., & Zheng, M. X. (2013). Package 'SNPRelate.' *A Package for Parallel Computing Toolset for*
Relatedness and Principal Component Analysis of SNP Data, *1093*.

Figures

Figure 1. Color variation of the elytra in the *Nebria ingens* species complex. (A) Quantification

of color using the a* (green-red axis) and b* (blue-yellow axis) color space, where chroma is

defined as the distance of each point from the achromatic center (0,0). (B) Map showing the

geographic distribution of color variation across sampling sites: the northernmost six populations

correspond to *N. riversi*, the southernmost eleven populations to *N. ingens*, and the intermediate

populations represent admixed groups.

Figure 2. Comparisons of modeled spectra calculated from measurements of the multilayer reflectors in the epicuticle of individuals with observed reflectance spectra in different elytral colors (A-C: green, D-F: dark green, G-I: black). The TEM images demonstrate the pattern of multilayer reflectors in the epicuticle. In each spectrum plot, the thin lines represent the means of modeled spectra from different observation points (see Supplemental methods) and the grey area shows the 95% confidence interval from measurements of different vertical transects, and the measured spectra are denoted with colored thick lines.

Figure 3. The scatter plot with mean BH-adjusted p-value of LFMM (x axis) and RDA absolute loading (y axis). Each dot represents a variant where the close dots are the outlier (candidate) variants defined by LFMM BH-adjusted p-value of ridge or lasso less than 0.01, and RDA loading falling outside of 4th standard deviation.

Figure 4. The correlations among elytral color (chroma), water loss rate, and mean September snow water equivalent (SWE) in habitat of the *N. ingens* complex. From left to right: (A) SWE vs. chroma; (B) SWE vs. water loss rate, and (C) water loss rate vs. chroma. The correlations were calculated using Spearman's rank correlation coefficient, and the grey areas indicate the 95% confidence interval of the estimate.