Individual Behavioral Variability Across Time and Contexts in *Dendrobates tinctorius*

Poison Frogs

Katharina M Soto[1,2], Faith O Hardin[1], Harmen P Alleyne[1], Eva K Fischer[1,3*]

[1] Department of Evolution, Ecology, and Behavior; School of Integrative Biology, University of

Illinois Urbana-Champaign, Urbana IL 61801, USA

[2] Program of Evolution, Ecology, and Conservation, School of Integrative Biology; University of

Illinois Urbana-Champaign, Urbana IL 61801, USA

[3] Carl R. Woese Institute for Genomic Biology, University of Illinois Urbana-Champaign, Urbana

IL 61801, USA

*Corresponding author:

Eva K Fischer

Department of Evolution, Ecology, and Behavior

School of Integrative Biology

University of Illinois Urbana-Champaign

505 S Goodwin Ave

Urbana, IL 61801

efischer@illinois.edu

Abstract

Consistent individual differences in behavior ("animal personality") have consequences for

individual fitness, evolutionary trajectories, and species' persistence. Such differences have

been documented across a wide range of animals, though amphibians are generally

underrepresented in this research area. The aim of our study was to examine consistent

individual differences in Dyeing poison frogs, *Dendrobates tinctorius*. We evaluated repeatability

in behaviors including activity, exploration, and boldness to assess consistency of behaviors

across different temporal, experimental, and environmental contexts. We found repeatability in

activity and exploration across time and contexts. In contrast, we observed context-specific

behavior for our metrics of boldness, with consistent individual differences only for some

measures. Further, while activity and exploration displayed consistent correlations across

contexts, relationships between activity and boldness were context dependent. Our findings

document the presence of consistent individual differences in behavior in *D. tinctorius* poison

frogs, and also reveal context-dependent behavioral differences, highlighting the complex

relationship between consistent individual differences and context-specific responses in animal

behavior.

Statement of Significance

The concept of animal personality centers on the existence of consistent individual differences

in behavior. However, behavioral responses can depend on context, and consistent individual

differences in one context do not guarantee consistent differences in another. To address this

question, we assessed activity, exploration, and boldness in captive-bred poison frogs

(*Dendrobates tinctorius*) across time and environmental contexts. Our comprehensive approach

revealed consistent individual differences in some behaviors and context-specificity in others.

While activity and exploration were generally repeatable and correlated with one another,

46   boldness was not. Especially in view of the emphasis on measures of boldness in the animal

47   personality literature, our findings emphasize the importance of reiterative and holistic

48   approaches in the study of animal behavior.

50   **Keywords**: behavioral syndromes, animal personality, repeatability, individual variation,

51   Dendrobatid

Introduction

There is growing evidence that consistent individual differences in behavior exist across taxa, and that these differences can have important consequences for individual fitness, adaptation, and species' persistence (Bell et al. 2009). For example, bold and aggressive individuals may be superior at territory defense but inferior at parental care (Duckworth 2006). Such trade-offs can lead to the establishment of alternative behavioral strategies within populations and potentially shape evolutionary trajectories (Smith and Blumstein 2008; Dochtermann and Dingemanse 2013). Moreover, certain personality types may fare better in the face of increasing human-induced global change, for example if they are better able to adapt to increasingly urban habitats (Pamela et al. 2015; Murgui and Hedblom 2017) or more likely to invade novel environments (Brodin et al. 2013; Carvalho et al. 2013). Increased recognition of the complex consequences of consistent individual behavioral differences have led to an increase in the study of animal personalities across taxa. Animal personality is increasingly invoked to explain behavioral differences in natural and captive populations, and to speculate on how these differences may influence survival and adaptation. However, many studies do so without confirming consistent individual differences across multiple contexts and over time. Assuming – rather than testing – that individual behavioral variation is stable across time and contexts can introduce bias, leading to adaptive inferences based on isolated incidents or context-specific behavior (Bell et al. 2009). Moreover, design and interpretation of behavioral assays that facilitate cross-species behavioral comparisons remain a challenge (Uher 2008; O'Connell and Hofmann 2011; Odom et al. 2021).

Animal personality is broadly defined as consistent individual differences in behavior (Réale et al. 2010; Carere and Maestripieri 2013). These differences are typically characterized by quantifying behavioral repeatability, defined as the proportion of variance explained by within-individual variation compared to between-individual variation (Bell et al. 2009; Bergmüller 2010; Garamszegi and Herczeg 2012). Animal personality encompasses an array of behavioral

traits, often with a particular focus on boldness:shyness, exploration:avoidance, and activity

(Réale et al. 2007; Garamszegi and Herczeg 2012; Michelangeli et al. 2019). Further, these

traits can become "packaged together" into behavioral syndromes (Sih et al. 2004). For

instance, studies spanning diverse taxa such as common voles (*Microtus arvali*) (Herde and

Eccard 2013), convict cichlids (*Amatitlania nigrofasciata*) (Mazué et al. 2015), sea turtles

(*Chelonia mydas)* (Kudo et al. 2021), and great tits (*Parus major*) (Cole and Quinn 2011)

document positive correlations between boldness, activity, and exploration.

While the presence of animal personalities and behavioral syndromes have been

documented in animals from primates to beetles, there is bias in taxonomic representation.

While there is extensive research in non-human animals including birds (e.g., Réale et al. 2007;

Carere and Locurto 2011) and fish (e.g., Bell 2005; Smith and Blumstein 2010; Mazué et al.

2015), amphibians have received comparatively less attention. Indeed, recent cross-taxon

reviews mention one or no studies on amphibians (Moiron et al. 2019; Cabrera et al. 2021).

These biases likely arise in part from historical assumptions that amphibians were "largely

instinctive bound machines controlled by external stimuli" (Maier and Schneirla 1964), yet it is

increasingly clear that these creatures make complex, context dependent behavioral decisions

(Burghardt 2013; Kelleher et al. 2018). Studying individual behavioral variation in amphibians is

critical for advancing our understanding of how these animals navigate risks and opportunities,

such as the trade-off between mating advertisement and predator avoidance (Stamps 2007; Sih

and Del Giudice 2012). Importantly, this knowledge has broader implications for understanding

evolutionary trajectories, ecological interactions, and conservation needs, which are of particular

importance given the alarming global decline of amphibian populations (Kelleher et al. 2018).

Among anurans (frogs and toads), poison frogs (family Dendrobatidae) offer compelling

study subjects for exploring individual variation and correlations in behavior. Poison frogs are

known for their bright coloration, toxicity, and behavior. These species' vivid hues are an honest

signal of their toxicity and have coevolved with a diurnal lifestyle (Summers 2003; Maan and

Cummings 2012). Poison frogs exhibit an array of interesting behaviors, including anti-predator

defenses (Breed 2008), aggressive territoriality (Crothers and Cummings 2015), parental care

(Roland and O'Connell 2015; Fischer et al. 2019), spatial learning and navigation (Liu et al.

2016; Pašukonis et al. 2022), and behavioral plasticity (Ringler et al. 2017; Peignier et al. 2023).

Taken together, these characteristics have garnered poison frogs increasing attention in the

realm of animal personality and behavioral syndromes. Existing studies in poison frogs have

examined consistent individual differences in activity (Sonnleitner et al. 2022), antipredator

behavior (Peignier et al. 2023), and aggression (Chaloupka et al. 2022). Nonetheless, a

knowledge gap remains regarding the extent of consistent individual differences across time

and context in poison frogs specifically, and amphibians more generally.

Dyeing poison frogs, *Dendrobates tinctorius*, have long been popular in zoos, aquaria,

and the pet trade, and have recently received increasing research attention in the wild (Rojas

and Pašukonis 2019; Fouilloux et al. 2021; Pašukonis et al. 2022) and the lab (Fischer et al.

2019, Fischer and O'Connell 2020; Sonnleitner et al. 2022). *D. tinctorius* boast a variety of

bright color morphs without sexual dimorphism in coloration. In the wild, males and females

maintain home ranges of a similar size, have similar daily movement patterns, and exhibit

aggression toward same- and opposite-sex individuals (Rojas and Endler 2013; Pašukonis et al.

2022). Unusually among frogs, males of the species do not produce conspicuous advertisement

calls, though they do produce low amplitude courtship vocalization (Rojas and Pašukonis 2019).

Like other Dendrobatids, *D. tinctorius* provide extended parental care that includes defense,

hydration, and cleaning of terrestrially deposited embryos, and transportation of tadpoles

piggyback to pools of water upon hatching (Stynoski et al. 2015; Rojas and Pašukonis 2019). In

D. tinctorius, tadpole transport is typically performed by males, however, females will

occasionally take over (Rojas and Pašukonis 2019; Fischer and O'Connell 2020). Alongside

their fascinating traits, existing work on *D. tinctorius* in both the field and the lab, and the ability

to ethically acquire animals make them an excellent study system.

We investigated personality and behavioral syndromes across contexts and timeframes

in *D. tinctorius*. Through a set of three experiments, we repeatedly assessed activity,

exploration, and boldness – in the same and in different assays – to test whether commonly

used personality metrics are robust to differences in assay design and timing (Fig. 1). Following

the methodologies of Kelleher et al. (2018), our experiments included an open field test (OFT), a

threat stimulus test (TST), a novel environment test (NET), and an activity in a familiar

environment test (AFET) (Fig. 1). All experiments were built upon the open field test, a widely

utilized method for assessing activity levels, exploratory behavior, and boldness across taxa

(Bell et al. 2009; Carere and Locurto 2011). Importantly, individuals were assayed multiple times

in each experiment (i.e., experienced multiple 'rounds' of each 'trial type' - terminology we use

throughout the manuscript). This design allowed us to assess repeatability of commonly used

behavioral metrics within and across contexts, and in response to multiple stimuli.

We predicted consistent individual differences in activity, exploration, and boldness

across multiple rounds and the three distinct experiments. Further, we anticipated a behavioral

syndrome, in which frogs with higher activity levels would also display increased boldness and

exploration. Finally, we expected frogs to perceive the threat stimulus and the novel

environment as stressful and exhibit context-dependent behavioral responses. Our overarching

goal was to provide a comprehensive, nuanced understanding of how consistent individual

differences in behavior manifest across time and context, and to reflect on the implications of

these patterns for extrapolation to ecological and evolutionary outcomes.

Methods

Experimental overview

In Experiment 1, we conducted an open field test (OFT) in an artificial environment to

establish baseline activity, exploration, boldness, and the relationships among them (Fig. 1).

This test reflects the standard, basic design used for open field tests. In Experiment 2, we

transitioned to an open field test (OFT) in a naturalistic environment, additionally assessing

behavior before, during, and after exposure to a threat stimulus test (TST) (Fig. 1). In

Experiment 3, we maintained the naturalistic environment, but extended the duration of each

trial including the baseline open field test (OFT) and an activity in a familiar environment test

(AFET), and additionally performed an extended duration novel environment test (NET) in an

artificial environment (Fig. 1).

D. tinctorius were housed in our animal facility at the University of Illinois Urbana-

Champaign. Terraria contained soil, moss, leaf litter, live plants, coconut shelters, and a small

pond. For housing and all experiments, we maintained terraria and test arenas at 21°–23° C

and 80%–90% humidity on a 12L:12D light cycle. We fed frogs *Drosophila hydei* or *Drosophila*

melanogaster dusted with vitamins three times weekly.

All behavioral experiments were conducted between March and April 2022. We used a

total of 48 captive bred frogs (N= 20 adult males, N= 12 adult females, and N= 17 juveniles of

unknown sex). All frogs were assayed in Experiments 1 and 2. In Experiment 3, we used a

subset of 19 frogs (N= 6 adult males, N= 8 adult females, and N= 5 juveniles; see additional

details below). For Experiments 1 and 2, we ran frogs sequentially in groups of five per week,

excluding the last group which had only four frogs. We followed the same protocol in a second

round of trials with the same groups of frogs. In essence, each individual underwent two

identical rounds, with a 5-week interval in between. For Experiment 3, we ran frogs in groups of

nine per week through a series of sequential trials that lasted a combined 25 hours (details

below). These trials were repeated three times with a 3-week interval between each round.

Group size for all experiments was chosen to make all behavioral testing feasible within a given

daily time window and to balance individuals of different color morphs across groups. Group

sizes and time intervals in Experiment 3 were modified to allow for the longer testing time

course while still assaying the same frogs in all experiments within the same timeframe.

Although frogs varied in age, all individuals were sexually inexperienced. Previous studies have

demonstrated that fed or digesting amphibians exhibit decreased levels of activity compared to

fasted individuals (Andrade et al. 2005). To ensure that the frogs were not actively digesting

during the experiments, we fasted frogs for 24 hours before behavioral trials.

Experiment 1: Artificial open field test

In Experiment 1, we investigated exploration, activity, and boldness in an open field test

(OFT) in a novel, artificial environment (Fig. 1). The open field arena was a circular enclosure

with a diameter of 108 cm. To maintain humidity and moisture, we lined the arena with white

bench paper soaked in reversed-osmosis (RO) water. For each observation, we placed a single

frog at the center of the arena as a standardized starting point and recorded behavior for 15

minutes using an overhead camera (DVC, HDV-604S). Individuals were immediately returned to

their home terraria following trials. The 15-minute testing time was chosen based on previous

tests in amphibians and other species. We quantified exploration as the proportion of areas a

frog visited, activity as the number of times a frog crossed the line between areas, and boldness

as time spent in the center of area (Fig. 2) (additional details below).

Experiment 2: Naturalistic open field and threat stimulus test

In Experiment 2, we quantified activity and exploration in a naturalistic open field. The

behavioral arena was a 20 x 30 x 20 cm (L x W x H) plexiglass tank that mimicked the frogs'

home terraria, including soil substrate, moss, leaves, and a live plant positioned in one corner.

The arena was completely covered from the outside with white contact paper, excluding a 24 x

17 cm (W x H) exposed area opposite the plant for the presentation of the threat stimulus. As in

Experiment 1, we placed a single frog at the center of the arena at the beginning of each

observation and used an overhead camera to record behavior. We conducted three trials to

investigate behavior (1) pre-stimulus open field test (OFT) (15 minutes after introduction to

arena), (2) during threat stimulus test (TST) (3 minutes of looming stimulus), and (3) post-

stimulus open field test (for 15 minutes after the stimulus) (Fig.1). The 15-minute testing time

was chosen based on previous studies and to align with Experiment 1. The duration of the

looming stimulus was determined through pilot testing. Test trials on frogs not included in the

study demonstrated that frogs typically showed a reaction within 3 minutes or froze throughout

the entire stimulus presentation.

We assessed activity and exploration using the same methods as in Experiment 1 (Fig.

2). Additionally, we measured boldness by observing the frogs' responses to a looming stimulus

(Fig. 2). Previous studies using leopard frogs (*Lithobates pipiens*) have shown reliable

avoidance behavior in response to looming objects (Ingle 1990; Waldeck and Gruberg 1995).

To present the looming stimulus, we used an iPad (Apple, 6th generation). The iPad was

covered in white tape, except for the display screen, and placed in the exposed area of the

arena (as shown Fig. 1). The looming stimulus was an expanding black circle projected

repeatedly onto the iPad screen for a duration of three minutes. To prevent habituation and

predictability, random pauses ranging from 1 to 30 seconds were incorporated into the stimulus

presentation. During the pre- and post- exposure trials, the iPad displayed a white screen to

match the surrounding environment. During the looming stimulus, we recorded activity and

exploration, as well as boldness based on whether the frogs approached or displayed

avoidance behavior in response to the stimulus.

Experiment 3: Naturalistic open field, familiar environment, and novel environment tests

In Experiment 3, we increased the duration of each trial to capture more prolonged

patterns in the frogs' behavior. To conduct these longer trials concurrently with Experiments 1

and 2 while also maintaining our repeated measures design, we selected a subset of individuals

(N= 19) from the previous experiments and limited our observations to two behaviors (activity

and boldness). The behavioral arena was constructed using the same materials as Experiment

2. However, the dimensions of the arena were twice as long at 30 x 60 x 20 cm (L x W x H) and

the arena split into two halves. One side of the arena mimicked the frogs' home terraria as in

Experiment 2. The other side mimicked the novel, artificial environment of Experiment 1, with

only white, moistened paper flooring. The two sides of the arena were separated by a white

cardboard barrier, which could be removed without disturbing the frog or the soil structure of the

natural chamber.

We conducted three trials to investigate the frogs' activity and boldness: (1) an open field

test (OFT) in a novel, naturalistic environment (the first 90 minutes after introduction to the

area), (2) activity in a familiar, natural environment test (AFET) (60 minutes after a 24 hour

acclimation period in the arena), and (3) exposure to a novel, artificial environment test (NET)

(60 minutes after the removal of the barrier between the familiar and novel environment) (Fig.

1). Time points were chosen to characterize differences based on longer observations (60

minutes = 4x the open field trial time in Experiments 1 and 2) and extended habituation to the

test arena (and immediate 90-minute trial and a second trial after 24 hours). Trials were

conducted sequentially, with each frog remaining in the arena for 25 hours while the trials

progressed. We conducted two subsequent rounds of trials for each individual, with three weeks

between each round. Due to the extended duration of each trial, we focused our analyses on

activity and boldness. This decision was made to streamline data collection and because we

expected – and found – a positive correlation between activity and exploration. We quantified

activity using the same methods as in Experiment 1 and 2, number of line crosses. To assess

boldness, we measured the frogs' latency to cross into the novel environment when the barrier

was lifted (Fig. 2).

Data analysis

We describe general methods here with modifications for each experiment below. We

coded videos using Behavioral Observation Research Interactive Software (BORIS) ver. 8.13

(Friard and Gamba 2016). Experiment type was not blind because assay identity was obvious

from videos, but all analyses were blind to individual identity. Behavioral data was analyzed with

R v.4.2.4 (R Core Team 2022) in Rstudio v.4.2.2. (RStudio Team 2021). For all response

variables that were normally distributed, we used the raw data for analysis. Response variables

that did not meet expectations of normality were transformed as described below. For all

analyses, outcomes were considered significant at $p <= 0.05$ and all posthoc comparisons were

Tukey adjusted for multiple hypothesis testing.

We used the lmer function in the lmerTest package (Kuznetsova et al. 2017) for all linear

models, and always included frog ID as a random effect and round (i.e., repeated

measurements in the same assay) as a fixed effect. Additional fixed effects varied based on

experiment (see below). We tested for the effects of sex on behavior and found no significant

results. Therefore, and because not all individuals could be sexed at the time of the experiment,

we did not include sex as a factor in the analyses described below. We ran Type III ANOVAs on

linear models to generate test statistics and p-values for main effects and interactions.

Subsequently, we used the emmeans package (Lenth 2021) to run pairwise post-hoc

comparisons where appropriate.

We performed repeatability analyses using the rptR package (Stoffel et al. 2017), with

95% confidence intervals (CI) based on 1000 bootstrapping runs. Frog ID was used as a

random effect for repeatability (R) among individuals. We used Pearson correlation for normally

distributed data and Kendall correlation for non-normally distributed data to assess relationships

between behaviors. The correlation coefficients were interpreted using these criteria: ≈0.1 a

small effect, ≈0.3 a moderate effect, and >0.5 a strong effect (Garamszegi et al. 2013).

Analysis Experiment 1: Artificial open field test

We coded videos in BORIS using a transparent overlay that divided the behavioral arena

into 40 areas: 24 on the ground and 16 on the wall. From videos, we quantified exploration as

the proportion of areas a frog visited and activity as the number of times a frog crossed a line

between areas. Movement between areas was recorded when the majority of the frogs' body

entered into a new area. We measured boldness as time spent in the center four areas of the

arena (Fig. 2). We log transformed the activity and boldness data to improve normality. We ran

linear models for each behavior (activity, exploration, boldness) as described above, including

round as a fixed effect and frog ID as a random effect. We performed repeatability analysis with

Gaussian distributions for exploration and Poisson distributions for activity and boldness (Stoffel

et al. 2017). Data from 11 individuals was lost due to equipment failure, leaving a final sample

size of N=38 individuals. Analyses and results are based on these samples.

Analysis Experiment 2: Naturalistic open field threat stimulus test

We coded videos in BORIS using a transparent overlay that divided the behavioral arena

into 32 areas: 16 on the ground and 16 on the wall. We quantified and analyzed activity and

exploration as in Experiment 1, with the addition of a fixed effect of trial (pre-stimulus, during

stimulus, and post-stimulus) to test for pre- and post- stimulus behavior differences. To compare

trials of different duration, we divided each behavior by the total trial time.

For response to stimulus, which we used as a measure of boldness, we divided the

arena into three sections: (1) at stimulus (the two wall areas overlaying the iPad, the two floor

areas directly in front the iPad, and one wall area immediately to the left of the iPad), (2) near

stimulus (the two wall areas and the four floor areas surrounding the "at stimulus" area), and (3)

away from stimulus (the remaining areas) (Fig. 2). We noted the frogs' position before and after

stimulus presentation. We used a Chi-Squared test in the Tidyverse package (Wickham et al.

2019) for pairwise comparisons of responses (moved away, moved toward, or froze) (R Core

Team 2022). We fit rptR models with Gaussian distributions to test for repeatability of each

behavior across rounds and across trials (Stoffel et al. 2017).

Analysis Experiment 3: Naturalistic open field novel environment test

We coded videos in BORIS using a transparent overlay that divided the behavioral arena

into 96 areas: 16 areas on both the natural and novel arena floor, and 16 areas on the walls of

each chamber. We recorded activity as the number of times a frog crossed the line between

areas as in Experiments 1 and 2. After the divider was removed, we recorded the time it took for

a frog to move into the novel area as latency to explore, which we interpreted as a measure of

boldness (Fig. 2). We recorded activity on both sides of the chamber after the barrier was

removed. We ran linear models for activity and latency to explore, including round (i.e., repeated

measurements in the assay), trial (baseline, after acclimation, in a novel environment), and trial:

round interaction as fixed effects, and frog ID as a random effect. As in Experiment 1, behaviors

were divided by trial duration when analyzing trial effects.

To determine whether activity predicted an individual's latency to cross into a new area,

we filtered the data to only include observations taken during the novel environment test. We

then fit a similar linear mixed effect model to predict latency to enter the novel area with fixed

effects of activity, round, and their interaction, and frog ID as a random effect. We fit rptR

models with Gaussian distributions to test for repeatability of each behavior across rounds and

across trials (Stoffel et al. 2017).

Cross-experiment analysis of activity and exploration

In addition to behavioral analyses within experiments, we were interested in the behavior

of individuals across experiments. To compare exploration between Experiments 1 and 2, which

had different arena sizes, we calculated the proportion of areas explored. We did not quantify

exploration in Experiment 3, and it was therefore excluded from this analysis (Fig. 2). We

compared the ratio of areas explored in Experiment 1 to both the pre-stimulus and post-stimulus

exploration in Experiment 2 using linear mixed models, with exploration rate and experiment as

fixed effects (Bates et al. 2015). We tested repeatability using a Gaussian distribution (Stoffel et

al. 2017).

In all experiments, we measured activity by counting the number of line crosses (Fig. 2).

To compare across experiments of different duration without introducing biases based on trial

time, we divided the number of lines crossed by the total trial time to calculate a line crossing

rate. We compared the subset of individuals (n= 19) in Experiment 3 with the same individuals

from Experiments 1 and 2. Baseline activity measurements were similar across all experiments,

taken prior to either the presentation of a stressor (threat stimulus test) or an environmental

stressor (novel environment test) (Fig. 1). We also compared the post-stimulus activity in

Experiment 2 to the activity observed during the novel environment test in Experiment 3,

reasoning that these trials were similar in representing responses following an environmental

change. We analyzed behavioral differences using activity rate as the response variable,

experiment as a fixed effect, and frog ID as a random effect (Bates et al. 2015). We calculated

post hoc comparisons and repeatability as before.

Results

Experiment 1: Artificial open field test

Frogs displayed repeatability in exploration (R= 0.278, p= 0.029) (Fig. 3a) and activity

(R= 0.279, p= 0.032) (Fig. 3b) and across both rounds of an open field test separated by five

weeks. However, time in the center of the arena, which we interpreted as a measure of

boldness, was not repeatable (R= 0.044, p= 0.395). We found no significant effect of round for

activity ($F_{1,37}$= 3.390, p= 0.074), exploration ($F_{1,37}$= 1.392, p= 0.245), or boldness ($F_{1,74}$= 0.628,

p= 0.431), suggesting that frogs' behavior did not change in response to previous experience in

the arena. Activity and exploration exhibited a strong positive correlation (corr= 0.767, p<

0.001). Boldness was not correlated with either activity (corr= -0.150, p= 0.196) or exploration

(corr= -0.119, p= 0.307) (Table 1).

Experiment 2: Naturalistic open field and threat stimulus test

Activity and exploration were repeatable across pre- and post- stimulus open field tests

in both round 1 (activity: R= 0.593, p< 0.001; exploration: R= 0.421, p< 0.001) and round 2

(activity: R= 0.529, p< 0.001; exploration: R= 0.449, p< 0.001). We observed the same trend

across rounds within trial (activity: R= 0.123, p= 0.006; exploration: R= 0.177, p< 0.001),

aligning with the results from Experiment 1 (Table 1). In contrast, during the threat stimulus test,

there was no repeatability of activity (R= 0.201, p= 0.069) or exploration (R= 0.175, p= 0.145)

across rounds.

We found an effect of trial on exploration ($F_{2,230}$= 112.143, p< 0.001) (Fig. 4a) and

activity ($F_{1,234}$= 170.128, p< 0.001) (Fig. 4b), with frogs decreasing behaviors during the threat

stimulus test and then increasing post stimulus (Fig. 4). We observed no significant round effect

for activity ($F_{1,232}$= 0.570, p= 0.451) or exploration ($F_{1,234}$= 0.292, p= 0.590), nor an interaction

between round and trial ($F_{2,228}$= 0.506, p= 0.604; $F_{2,228}$= 0.216, p= 0.801), indicating that frogs

behaved consistently across multiple experiences in the arena. Activity and exploration were

highly correlated across all trials (pre-stimulus, during stimulus, and post-stimulus) (corr= 0.7–

0.8, p< 0.001).

We evaluated boldness by quantifying frogs' responses to the threat (looming) stimulus.

Frogs' responses to the looming stimulus were not repeatable across rounds (R= 0, p= 1). This

could be in part because overall among individual variation was low, as the majority of frogs

demonstrated a freezing response during stimulus presentation rather than moving away ($\chi^2_{1,96}$=

19.755, p< 0.001) or towards ($\chi^2_{1,96}$= 41.796, p< 0.001) the stimulus. No significant effect of

round was observed for boldness ($\chi^2_{2,96}$= 22.429, p= 0.401).

We found significant correlations between boldness (response to stimulus) and frogs'

activity (tau= 0.155, p= 0.002) and exploration (tau= 0.155, p= 0.002) during the pre- and post-

stimulus trials. These trends suggest that frogs displaying higher levels of activity and

exploration prior to and after the stimulus were also more inclined to approach the stimulus

during the threat stimulus trial. These correlations between boldness, activity, and exploration

contrast with the results obtained in Experiment 1 (Table 1).

Experiment 3: Naturalistic open field and novel environment test

Frogs were repeatable for activity both across trials (R= 0.321, p<0.011) and across

rounds (R= 0.651, p= 0.001). We found that frogs moved more with repeated experience in the

arena across all trials ($F_{2,143}$= 3.919, p= 0.022), with round 1 having significantly less movement

than round 2 (t_{36}=-3.423, p=0.004) and Round 3 (t_{36}= -3.784, p= 0.002), but no significant

difference between rounds 2 and 3 (t_{36}= -0.360 p=0.931) (Fig. 5a). Similarly, we found that frogs

took less time to cross the boundary into the novel area during the novel environment test as

rounds progressed ($F_{2,36}$= 13.100, p< 0.001), with frogs taking longer to cross in round 1

compared to round 2 (t_{36}= 2.837, p= 0.019) and round 3 (t_{36}= 5.108, p< 0.001), but not in round

2 compared to round 3 (t_{36}= 2.270, p= 0.073) (Fig 5b). We found significant repeatability for

latency to explore across rounds (R= 0.388, p= 0.003) suggesting all frogs responded to

repeated experience similarly.

We use latency to cross into a novel environment as an indicator of boldness,

interpreting shorter latencies as bolder. We found a significant negative correlation between

activity and latency to cross into a novel environment during the novel environment test (corr= -

0.665, p< 0.001), indicative of a positive relationship between activity and this measure of

boldness (Fig 5c). Additionally, we found that boldness was influenced by both activity ($F_{1,38}$=

28.795, p<0.001) and round ($F_{2,38}$= 6.546, p= 0.004), indicating that more active frogs moved

into the novel area more quickly, and all frogs moved into the novel area more quickly with

repeated experience.

Cross-experiment analysis of activity and exploration

Frogs exhibited consistent individual differences in exploration (R= 0.171, p= 0.005)

across Experiments 1 and 2. For activity, baseline activity during the open field tests were

repeatable between the naturalistic environments of Experiment 2 and Experiment 3 (R= 0.207,

p< 0.001), and between the artificial environment of Experiment 1 and the naturalistic

environment of Experiment 3 (R= 0.171, p= 0.041). Activity was not repeatable between the

artificial environment of Experiment 1 and the naturalistic environment of Experiment 2 (R=

0.029, p= 0.401). Activity was repeatable across stressor trials (Fig. 1) (i.e., threat stimulus test

in Experiment 2 and novel environment test in Experiment 3) (R= 0.243, p= 0.003).

Discussion

We investigated animal personality and behavioral syndromes in captive-bred poison

frogs (*Dendrobates tinctorius*) by characterizing behavioral differences across multiple

environmental contexts and timepoints. We conducted three experiments to assess activity,

exploration, and boldness in the same individuals and found repeatability in activity and

exploration across time and contexts. In contrast, we observed context-specific behavior for our

metrics of boldness, with consistent individual differences only for some measures. The stability

of individual differences in behavior can have implications for how species respond to shifting

environmental conditions (Wong and Candolin 2015; Merrick and Koprowski 2017), thereby

shaping evolutionary trajectories and ecological interactions (Shaw 2020; Stiegler et al. 2022).

Despite their declining populations, amphibians remain relatively understudied in behavioral

research. The results of our work help to fill this gap, address broader issues in the field of

"animal personality", and set the stage for future work in *D. tinctorius*.

Our study revealed widespread repeatability for activity and exploration, evident across

time, trials, and experimental contexts. These findings suggest that measures of activity and

exploration are indeed reliable, and measurements in one assay or at one timepoint are good

predictors of an individual's behavior in another context or at another time. We also recovered

the commonly described activity:exploration behavioral syndrome (Réale et al. 2007). While we

expected a positive relationship between activity and exploration (i.e., exploration is impossible

without movement), we note that there is room for variability in this relationship. For instance,

frogs might have displayed a lot of movement but only within a restricted area, leading to

heightened activity but limited exploration. The strong, significant relationships we found

(correlations > 0.76), suggest that these behaviors are indeed tightly linked and can therefore

reliably predict one another as well as some other behaviors (e.g., latency to explore a novel

area, see below).

The above outcomes align with prior research on animal personality across taxa (Réale

et al. 2007; Kelleher et al. 2018), and complement a recent investigation on another

Dendrobatid, *Allobates femoralis*, which demonstrated the efficacy of both artificial and

naturalistic setups in capturing consistent individual differences in behavior (Bégué et al. 2023).

However, a study on *D. tinctorius* froglets by Sonnleitner et al. (2022) yielded somewhat

different findings: the authors found significant repeatability at the family level but no significant

repeatability at the individual level. These results suggests that genetic factors influence

behavior, and this effect outweighs individual differences when related individuals are reared

under similar conditions. We were unable to control for relatedness in our study, but assume

genetic differences explain at least some of the behavioral variation we document. A non-

mutually exclusive explanation is that individuals in the study of Sonnleitner et al. (2022) were

much younger than those used here, and individual differences in behavior may become more

pronounced with age and experience (Bell et al. 2009). We note that our findings here and

those of Sonnleitner et al. (2022) are from captive-bred individuals. While *D. tinctorius* readily

exhibit behaviors of interest (e.g., parental care and aggression) under captive conditions,

influences of lab adaptation and genetic differences between wild and captive animals surely

also influence behavior. In any case, while consistent individual differences are being

increasingly documented across diverse taxa, contrasting results even in the same species

underscore the importance of considering various contexts and timepoints when extrapolating

from single studies to evolutionary and ecological outcomes.

While we observed a high degree of repeatability in activity and exploration across time

and contexts, the behaviors we considered as measures of boldness demonstrated contextual

dependence. Our first measure was the commonly used metric of time spent in the center of a

novel arena (Carter et al. 2012; Yuen et al. 2017; Baker et al. 2018; Rudolfová et al. 2022). We

did not observe consistent individual differences in this measure, nor was it correlated with other

behaviors. As our next measure of boldness, we assayed response to a threat stimulus in a

familiar environment, aligning with prior studies (Ingle 1990; Waldeck and Gruberg 1995; Carter

et al. 2012; Yuen et al. 2017). Once again, we did not find repeatability for boldness, although

we did observe a correlation between this measure of boldness, activity, and exploration. In our

final approach, we measured boldness as the latency to enter a novel environment, a less

common yet established method for evaluating boldness (White et al. 2013; Magnhagen et al.

2014; Theódórsson and Ólafsdóttir 2022). With this approach, we observed both repeatability

and a correlation between boldness and activity. Taken together, our results suggest that

commonly used boldness methodologies may not measure the same personality trait.

Despite the popularity of measuring boldness in animal personality studies, our findings

align with those of others who have explicitly compared outcomes across different metrics

categorized as 'boldness' and found conflicting results. For example, our approach and

outcomes are similar to that of Yuen et al. (2017) in their study on African striped mice

(*Rhabdomys pumilio*). The authors investigated whether distinct experimental methods were

indeed measuring boldness. Like our approach, they employed both artificial and natural

environments, and conducted two boldness tests, an open field, and a startle test. Yuen et al.

(2017) highlighted the distinction between these methods: the startle test evaluated risk-taking

behavior after threat exposure, potentially measuring anxiety rather than boldness, while the

open field test did not incorporate a preceding threat and proved to be a more reliable method

for gauging boldness in their species. Carter et al. (2012) found related results when examining

chacma baboons (*Papio ursinus*), concluding that threatening situations may measure anxiety,

not boldness. Moreover, the use of artificial open field tests, particularly in evaluation of time

spent in the center, may not consistently capture boldness. Indeed, a recent study on Zebrafish

(*Danio rerio*) suggests that thigmotaxis may not effectively measure boldness or anxiety-like

behavior in this species (Rajput et al. 2022).

Translating these observations to our own results, our open field test focused on

evaluating time in center, the threat stimulus test observed behavior in the presence of a threat,

whereas the novel environment test examined behavior in a new and unfamiliar environment.

While we initially considered all tests as measures of boldness, our data suggest that the

specific nature of the challenges presented in each experiment and the environmental context

led to varying types and intensities of behavioral responses, and in turn distinct patterns of

repeatability and behavioral correlations. Our findings also confirm the importance of

considering the environment when measuring behavioral correlations: we observed correlations

between boldness, activity, and exploration (i.e., behavioral syndromes) in naturalistic

environments (Experiments 2 and 3) but not in artificial settings (Experiment 1) (Table 1).

Alongside those of others, our findings highlight that the environment and assay in which

individuals are assessed can exert a profound influence on their behavior (e.g., Niemelä and

Dingemanse 2014; Bégué et al. 2023) and the necessity of evaluating individual variation

across diverse contexts (Bell et al. 2009).

Taken together, these observations emphasize the challenges of adapting standard

methodologies across closely and distantly related taxa. For example, poison frogs are diurnal

unlike most frogs, and amphibians are generally less active than mammals. While behavioral

and life-history differences between taxa are obvious and themselves of interest, they provide

fundamental challenges for the design, implementation, and interpretation of comparative

520 studies (Uher 2008; O'Connell and Hofmann 2011; Odom et al. 2021). When findings align

521 across taxa, we assume shared core principles; however, when patterns differ it is more difficult

522 to understand whether organisms differ fundamentally in, for example, relationships between

523 activity and boldness, or whether we have not "asked them the right question" with our chosen

524 assay. Fortunately, with a growing literature that includes more species and taxa, these

525 challenges are also opportunities to build our understanding through comparative data and

526 thinking.

528 *Conclusions*

529  Our findings demonstrate consistent individual differences in behavior across time and

530 context in *D. tinctorius*, while also highlighting the subtle complexities of studying animal

531 personality. While we found strong evidence for individual consistency in activity and

532 exploration, our measures of boldness were variable and context dependent. Our results

533 demonstrate the value of assessing multiple behavioral traits and the relationships among them

534 to build a comprehensive view of behavioral repeatability and variability within and across

535 species (Sih et al. 2004; Bell et al. 2009; Sih and Del Giudice 2012). Broadly, our results add to

536 the growing literature emphasizing the importance of evaluating context-dependent patterns of

537 individual behavioral variation, even for assays commonly considered as measuring the same

538 trait (Bell et al. 2009; Carter et al. 2012; Yuen et al. 2017). More specifically, our findings

539 contribute to the growing recognition of behavioral complexity in anuran amphibians, with

540 implications for amphibian research, including their role as ecological indicators amid habitat

541 degradation and climate change (Grant et al. 2019). Our findings here pave the way for future

542 research on the impact of consistent individual differences on reproductive success, survival,

543 and community dynamics.

Acknowledgments

We thank the members of the Fischer Lab for feedback on previous versions of the

manuscript and help with frog husbandry. For their insight on statistical analyses, we thank Drs.

Jen Moss and Sarah Westrick. We thank two anonymous reviewers and the associate editor for

comments that improved the manuscript. Our work would not be possible without the animal

care support of the Division of Animal Resources (DAR) at UIUC.

Funding

This work is/was supported by the USDA National Institute of Food and Agriculture Hatch

project 1026333 (ILLU-875-984 to KMS), a UIUC Graduate College Master's Fellowship (to

KMS), a UIUC Illinois Distinguished Fellowship (to FOH), and UIUC Laboratory Start-up funds

(to EKF).

Data availability

Data are included in the Supplementary information.

Author Contributions

KMS, FOH, HPA, and EKF conceived of and planned the experiments. KMS, FOH, and HPA

carried out the experiments. KMS, FOH, and EKF contributed to data analysis and interpretation

of the results. KMS wrote the manuscript with assistance from all authors.

Compliance with Ethical Standards

Disclosure of potential conflicts of interest

The authors declare no competing interests.

Ethics approval

All animal care and experimental procedures were approved by the University of Illinois Urbana-

Champaign Animal Care and Use Committee (protocol #20147) and followed all institutional and

national guidelines for the use of animals in research.

References

Andrade DV, Cruz-Neto AP, Abe AS, Wang T (2005) Specific dynamic action in ectothermic

vertebrates: a review of the determinants of postprandial metabolic response in fishes,

amphibians, and reptiles. In: Stark JM, Wang T (eds) Physiological and ecological

adaptations to feeding in vertebrates. Science Publishers, Enfield, pp 306-324

Baker MR, Goodman AC, Santo JB, Wong RY (2018) Repeatability and reliability of exploratory

behavior in proactive and reactive zebrafish, *Danio rerio*. Sci Rep 8:12114.

https://doi.org/10.1038/s41598-018-30630-3

Bates D, Mächler M, Bolker BM, Walker SC (2015) Fitting linear mixed-effects models using

lme4. J Stat Softw 67:1-48. https://doi.org/10.18637/jss.v067.i01

Bégué L, Peignier M, Ringler E (2023) The link between animal personality and habitat selection

in males of the Neotropical poison frog *Allobates femorali*s. Behaviour 160:257–274.

https://doi.org/10.1163/1568539X-bja10202

Bell AM (2005) Behavioural differences between individuals and two populations of stickleback

(*Gasterosteus aculeatus*). J Evol Biol 18:464–473. https://doi.org/10.1111/j.1420-

9101.2004.00817.x

Bell AM, Hankison SJ, Laskowski KL (2009) The repeatability of behaviour: a meta-analysis.

Anim Behav 77:771–783. https://doi.org/10.1016/j.anbehav.2008.12.022

Bergmüller R (2010) Animal personality and behavioural syndromes. In: Kappeler P (ed) Animal

Behavior: Evolution and Mechanisms. Springer, Berlin, pp 587-621

Breed M (2008) Warning colouration predicts behavioural and morphological differences in the

strawberry poison-dart frog. Thesis, Uppsala University

Brodin T, Lind MI, Wiberg MK, Johansson F (2013) Personality trait differences between

mainland and island populations in the common frog (*Rana temporaria*). Behav Ecol

Sociobiol 67:135–143. https://doi.org/10.1007/s00265-012-1433-1

Burghardt GM (2013) Environmental enrichment and cognitive complexity in reptiles and

amphibians: Concepts, review, and implications for captive populations. Appl Anim Behav

Sci 147:286–298. https://doi.org/10.1016/j.applanim.2013.04.013

Cabrera D, Nilsson JR, Griffen BD (2021) The development of animal personality across

Ontogeny: A cross-species review. Anim Behav 173:137–144.

https://doi.org/10.1016/j.anbehav.2021.01.003

Carere C, Locurto C (2011) Interaction between animal personality and animal cognition. Curr

Zool 57:491–498. https://doi.org/10.1093/czoolo/57.4.491

Carere C, Maestripieri D (2013) Animal personalities: behavior, physiology, and evolution.

University of Chicago Press, Chicago

Carter AJ, Marshall HH, Heinsohn R, Cowlishaw G (2012) How not to measure boldness: Novel

object and antipredator responses are not the same in wild baboons. Anim Behav 84:603–

609. https://doi.org/10.1016/j.anbehav.2012.06.015

Carvalho CF, Leitão AV., Funghi C, Batalha HR, Reis S, Mota PG, Lopes RJ, Cardoso GC

(2013) Personality traits are related to ecology across a biological invasion. Behav Ecol

24:1081–1091. https://doi.org/10.1093/beheco/art034

Chaloupka S, Peignier M, Stückler S, Araya-Ajoy Y, Walsh P, Ringler M, Ringler E (2022)

Repeatable territorial aggression in a neotropical poison frog. Front Ecol Evol 10:881387.

https://doi.org/10.3389/fevo.2022.881387

Cole, Ella F, Quinn, John L (2011) Personality and problem-solving performance explain

competitive ability in the wild. Proc R Soc Lond B 279:1168–1175

Crothers LR, Cummings ME (2015) A multifunctional warning signal behaves as an agonistic

status signal in a poison frog. Behav Ecol 26:560–568.

https://doi.org/10.1093/beheco/aru231

Dochtermann NA, Schwab T, Sih A (2014) The contribution of additive genetic variation to

personality variation: Heritability of personality. Proc R Soc B 282:20142201.

https://doi.org/10.1098/rspb.2014.2201

Duckworth RA (2006) Behavioral correlations across breeding contexts provide a mechanism

for a cost of aggression. Behav Ecol 17:1011–1019. https://doi.org/10.1093/beheco/arl035

Fischer EK, O'Connell LA (2020) Hormonal and neural correlates of care in active versus

observing poison frog parents. Horm Behav 120:104696.

https://doi.org/10.1016/j.yhbeh.2020.104696

Fischer EK, Roland AB, Moskowitz NA, Vidoudez C, Ranaivorazo N, Tapia EE, Trauger SA,

Vences M, Coloma LA, O'Connell LA (2019) Mechanisms of convergent egg provisioning

in poison frogs. Curr Biol 29:4145-4151.e3. https://doi.org/10.1016/j.cub.2019.10.032

Friard O, Gamba M (2016) BORIS: a free, versatile open-source event-logging software for

video/audio coding and live observations. Methods Ecol Evol 7:1325–1330.

https://doi.org/10.1111/2041-210X.12584

Fouilloux CA, Serrano Rojas SJ, Carvajal-Castro JD, Valkonen JK, Gaucher P, Fischer M,

Pašukonis A, Rojas B (2021) Pool choice in a vertical landscape: tadpole-rearing site

flexibility in phytotelm-breeding frogs. Ecol Evol 11:9021-9038.

https://doi.org/10.1002/ece3.7741

Garamszegi LZ, Herczeg G (2012) Behavioural syndromes, syndrome deviation and the within-

and between-individual components of phenotypic correlations: When reality does not meet

statistics. Behav Ecol Sociobiol 66:1651–1658. https://doi.org/10.1007/s00265-012-1439-8

Garamszegi LZ, Markó G, Herczeg G (2013) A meta-analysis of correlated behaviors with

implications for behavioral syndromes: Relationships between particular behavioral traits.

Behav Ecol 24:1068–1080. https://doi.org/10.1093/beheco/art033

Grant EHC, Muths E, Schmidt BR, Petrovan SO (2019) Amphibian conservation in the

Anthropocene. Biol Conserv 236:543–547. https://doi.org/10.1016/j.biocon.2019.03.003

Herde A, Eccard JA (2013) Consistency in boldness, activity and exploration at different stages

of life. BMC Ecol 103:61–68. https://doi.org/10.1186/1472-6785-13-49

Ingle DJ (1990) Visually elicited evasive behavior in frogs. Bioscience 40:284–291.

https://doi.org/10.2307/1311265

Kelleher SR, Silla AJ, Byrne PG (2018) Animal personality and behavioral syndromes in

amphibians: a review of the evidence, experimental approaches, and implications for

conservation. Behav Ecol Sociobiol 72:79. https://doi.org/10.1007/s00265-018-2493-7

Kudo H, Nishizawa H, Uchida K, Sato K (2021) Boldness–exploration behavioral syndrome in

wild sub-adult green sea turtles caught at Oita, Japan. Appl Anim Behav Sci 236:105216.

https://doi.org/10.1016/j.applanim.2021.105216

Kuznetsova A, Brockhoff PB, Christensen RHB (2017) lmerTest package: tests in linear mixed

effects models. J Stat Softw 82:1–26. https://doi.org/10.18637/JSS.V082.I13

Lenth R (2021) emmeans: Estimated Marginal Means, aka Least-Squares Means. R package

version 1.8.5, https://github.com/rvlenth/emmeans.

Liu Y, Day LB, Summers K, Burmeister SS (2016) Learning to learn: Advanced behavioural

flexibility in a poison frog. Anim Behav 111:167–172.

https://doi.org/10.1016/j.anbehav.2015.10.018

Maan ME, Cummings ME (2012) Poison frog colors are honest signals of toxicity, particularly for

bird predators. Am Nat 179:E1-E14. https://doi.org/10.1086/663197

Magnhagen C, Wacker S, Forsgren E, Myhre LC, Espy E, Amundsen T (2014) Context

consistency and seasonal variation in boldness of male two-spotted gobies. PLoS ONE

9:e93354. https://doi.org/10.1371/journal.pone.0093354

Maier NR, Schneirla TC (1964) Principles of animal psychology. Dover, London

Mazué GPF, Dechaume-Moncharmont FX, Godin JGJ (2015) Boldness-exploration behavioral

syndrome: Interfamily variability and repeatability of personality traits in the young of the

convict cichlid (*Amatitlania siquia*). Behav Ecol 26:900–908.

https://doi.org/10.1093/beheco/arv030

Merrick MJ, Koprowski JL (2017) Should we consider individual behavior differences in applied

wildlife conservation studies? Biol Conserv 209:34–44.

https://doi.org/10.1016/j.biocon.2017.01.021

Michelangeli M, Chapple DG, Goulet CT, Bertram MG, Wong BBM (2019) Behavioral

syndromes vary among geographically distinct populations in a reptile. Behav Ecol 30:393–

401. https://doi.org/10.1093/beheco/ary178

Moiron M, Laskowski KL, Niemelä PT. 2019. Individual differences in behaviour explain

variation in survival: A meta-analysis. Ecol Lett 23:399–408.

https://doi.org/10.1111/ele.13438

Murgui E, Hedblom M (eds) (2017) Ecology and conservation of birds in urban environments.

Springer, Cham. https://doi.org/10.1007/978-3-319-43314-1

Niemelä PT, Dingemanse NJ (2014) Artificial environments and the study of "adaptive"

personalities. Trends Ecol Evol 29:245–247. https://doi.org/10.1016/j.tree.2014.02.007

O'Connell LA, Hofmann HA (2011) Genes, hormones, and circuits: An integrative approach to

study the evolution of social behavior. Front Neuroendocrinol 32:320–335.

https://doi.org/10.1016/j.yfrne.2010.12.004

Odom KJ, Araya-Salas M, Morano JL et al (2021) Comparative bioacoustics: A roadmap for

quantifying and comparing animal sounds across diverse taxa. Biol Rev 96:1135–1159.

https://doi.org/10.1111/brv.12695

Pamela Delarue EM, Kerr SE, Lee Rymer T (2015) Habitat complexity, environmental change

and personality: A tropical perspective. Behav Process 120:101–110.

https://doi.org/10.1016/j.beproc.2015.09.006

Pašukonis A, Serrano-Rojas SJ, Fischer MT et al (2022) Contrasting parental roles shape sex

differences in poison frog space use but not navigational performance. eLife 11:e80483.

https://doi.org/10.7554/eLife.80483

Peignier M, Araya-Ajoy YG, Bégué L, Chaloupka S, Dellefont K, Walsh P, Ringler M, Ringler E

(2022) Exploring links between personality traits and their social and non-social

environments in wild poison frogs. Behav Ecol Sociobiol 76:93.

https://doi.org/10.1007/s00265-022-03202-9

Peignier M, Bégué L, Ringler M, Szabo B, Ringler E (2023) Regardless of personality, males

show similar levels of plasticity in territory defense in a Neotropical poison frog. Sci Rep

13:3435. https://doi.org/10.1038/s41598-023-30546-7

R Core Team (2022) R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria, http://www.R-project.org

Rajput N, Parikh K, Kenney JW (2022) Beyond bold versus shy: Zebrafish exploratory behavior

falls into several behavioral clusters and is influenced by strain and sex. Biol Open

11:bio05944. https://doi.org/10.1242/bio.059443

Réale D, Dingemanse NJ, Kazem AJN, Wright J (2010) Evolutionary and ecological approaches

to the study of personality. Phil Trans R Soc B 365:3937–3946.

https://doi.org/10.1098/rstb.2010.0222

Réale D, Reader SM, Sol D, McDougall PT, Dingemanse NJ (2007) Integrating animal

temperament within ecology and evolution. Biol Rev 82:291–318

Ringler E, Barbara Beck K, Weinlein S, Ludwig H, Ringler M (2017) Adopt, ignore, or kill? Male

poison frogs adjust parental decisions according to their territorial status. Sci Rep 7:43544.

https://doi.org/10.1038/srep43544

Rojas B, Endler JA (2013) Sexual dimorphism and intra-populational colour pattern variation in

the aposematic frog *Dendrobates tinctorius*. Evol Ecol 27:739–753.

https://doi.org/10.1007/s10682-013-9640-4

Rojas B, Pašukonis A (2019) From habitat use to social behavior: Natural history of a voiceless

poison frog, *Dendrobates tinctorius*. PeerJ 7:e7648. https://doi.org/10.7717/peerj.7648

Roland AB, O'Connell LA (2015) Poison frogs as a model system for studying the neurobiology

of parental care. Curr Opin Behav Sci 6:76–81.

https://doi.org/10.1016/j.cobeha.2015.10.002

RStudio Team (2021) RStudio: Integrated Development Environment for R. RStudio, PBC,

Boston, MA

Rudolfová V, Petrásek T, Antošová E, Frynta D, Landová E, Valeš K, Nekovářová T (2022)

Inter-individual differences in laboratory rats as revealed by three behavioural tasks. Sci

Rep 12:9361. https://doi.org/10.1038/s41598-022-13288-w

Shaw AK (2020) Causes and consequences of individual variation in animal movement. Mov

Ecol 8:12. https://doi.org/10.1186/s40462-020-0197-x

Sih A, Bell A, Johnson JC (2004) Behavioral syndromes: An ecological and evolutionary

overview. Trends Ecol Evol 19:372–378. https://doi.org/10.1016/j.tree.2004.04.009

Sih A, Del Giudice M (2012) Linking behavioural syndromes and cognition: A behavioural

ecology perspective. Phil Trans R Soc B 367:2762–2772.

https://doi.org/10.1098/rstb.2012.0216

Smith BR, Blumstein DT (2008) Fitness consequences of personality: A meta-analysis. Behav

Ecol 19:448–455. https://doi.org/10.1093/beheco/arm144

Smith BR, Blumstein DT (2010) Behavioral types as predictors of survival in Trinidadian guppies

(*Poecilia reticulata*). Behav Ecol 21:919–926. https://doi.org/10.1093/beheco/arq084

Sonnleitner R, Alanen E, Fouilloux C, Valkonen JK, Rojas B (2022) Genetic factors influence

behavioural repeatability in juvenile poison frogs. bioRxiv, 2022.09.19.508512.

https://doi.org/10.1101/2022.09.19.508512

Stamps, Judy A (2007) Growth-mortality tradeoffs and 'personality traits' in animals. Ecol Lett

10:355–363. https://doi.org/10.1111/j.1461-0248.2007.01034.x

Stiegler J, Lins A, Dammhahn M, Kramer-Schadt S, Ortmann S, Blaum N (2022) Personality

drives activity and space use in a mammalian herbivore. Mov Ecol 10:33.

https://doi.org/10.1186/s40462-022-00333-6

Stoffel MA, Nakagawa S, Schielzeth H (2017) rptR: repeatability estimation and variance

decomposition by generalized linear mixed-effects models. Methods Ecol Evol 8:1639–

1644. https://doi.org/10.1111/2041-210X.12797

Stynoski JL, Schulte LM, Rojas B (2015) Poison frogs. Curr Biol 25:1026–1028.

https://doi.org/10.1016/j.cub.2015.06.044

Summers K (2003) Convergent evolution of bright coloration and toxicity in frogs. P Natl Acad

Sci USA 100:12533–12534. https://doi.org/10.1073/pnas.2335928100

Theódórsson BK, Ólafsdóttir GÁ (2022) Laboratory measures of boldness correlate with

ecological niche in threespine stickleback. Ecol Evol 12:e9235.

https://doi.org/10.1002/ece3.9235

Uher J (2008) Comparative personality research: Methodological approaches. Eur J Pers

22:427–455. https://doi.org/10.1002/per.680

Waldeck RF, Gruberg ER (1995) Studies on the optic chiasm of the leopard frog. Brain Behav

Evol 46:84–94. https://doi.org/10.1016/j.anbehav.2022.10.005

White JR, Meekan MG, McCormick MI, Ferrari MCO (2013) A comparison of measures of

boldness and their relationships to survival in young fish. PLoS ONE 8:e68900.

https://doi.org/10.1371/journal.pone.0068900

Wickham H, Averick M, Bryan J et al (2019) Welcome to the Tidyverse. J Open Source Softw

4:1686. https://doi.org/10.21105/joss.01686

Wilson ADM, Godin J-GJ, Ward AJW (2010) Boldness and reproductive fitness correlates in the

eastern mosquitofish, *Gambusia holbrooki*. Ethology 116:96–104.

https://doi.org/10.1111/j.1439-0310.2009.01719.x

Wong BBM, Candolin U (2015) Behavioral responses to changing environments. Behav Ecol

26:665–673. https://doi.org/10.1093/beheco/aru183

Yuen CH, Schoepf I, Schradin C, Pillay N (2017) Boldness: are open field and startle tests

measuring the same personality trait? Anim Behav 128:143–151.

https://doi.org/10.1016/j.anbehav.2017.04.009

    Tables

789   **Table 1** Overview of statistical results for behavioral repeatability, correlations, and linear

790   models. Exploration was not quantified in Experiment 3. Significance interpreted as: R= 0.2-0.4

791   moderate repeatability, corr≥0.5 strong correlation, p< 0.05 significant. Pearson correlation

792   (corr) used for normal data, Kendall Rank correlation (tau) used for non-parametric data

Repeatability			
Behavior	**Experiment 1**	**Experiment 2**	**Experiment 3**
Activity	**R= 0.279, p= 0.032**	**R= 0.123, p= 0.006**	**R= 0.321, p= 0.011**
Exploration	**R= 0.278, p= 0.029**	**R= 0.177, p< 0.001**	NA
Boldness	R= 0.044, p= 0.395	R= 0, p= 1	**R= 0.388, p= 0.003**
Behavioral Syndromes			
Syndrome	**Experiment 1**	**Experiment 2**	**Experiment 3**
Activity:Exploration	**corr= 0.767, p< 0.001**	**corr= 0.911, p< 0.001**	NA
Activity:Boldness	corr= -0.150, p= 0.196	**tau= 0.155, p= 0.002**	**corr= -0.665, p< 0.001**
Exploration:Boldness	corr= -0.119, p= 0.307	**tau= 0.155, p= 0.002**	NA
ANOVAs			
Activity	**Experiment 1**	**Experiment 2**	**Experiment 3**
Trial	NA	$F_{1,234}$= 170.128, p< 0.001	$F_{2,143}$= 18.387, p<0.001
Round	$F_{1,37}$= 3.390, p= 0.074	$F_{1,232}$= 0.570, p= 0.451	$F_{2,143}$=3.919, p= 0.022
Trial:Round	NA	$F_{2,228}$= 0.216, p= 0.801	$F_{4,143}$ = 1.356, p= 0.252
Exploration	**Experiment 1**	**Experiment 2**	**Experiment 3**
Trial	NA	$F_{2,230}$= 112.143, p< 0.001	NA
Round	$F_{1,37}$= 1.392, p= 0.245	$F_{1,234}$= 0.292, p= 0.590	NA
Trial:Round	NA	$F_{2,228}$= 0.506, p= 0.604	NA
Boldness	**Experiment 1**	**Experiment 2**	**Experiment 3**
Round	$F_{1,74}$= 0.628, p= 0.431	$\chi^2_{2,96}$= 22.429, p= 0.401	$F_{2,36}$ = 13.100, p< 0.001

Figures

Fig 1 Overview of experimental design. All experiments were built on the framework of an open

field test and expanded to explore differences in time and context. For each experiment, trial

type (i.e., different phases within each experiment), trial duration, and number of rounds (i.e.,

repeated experiences in the same assay) are shown. OFT = open field experiment, TST =

threat stimulus test, AFET = activity in a familiar environment test, NET = novel environment

test. Colors represent corresponding trial types that were compared across experiments (see

"Cross Analysis" Methods)

Fig 2 Overview of analysis methods. For all experiments, we quantified activity as the number of lines crossed from one area into the next and exploration as the total number of areas visited (not assessed in Experiment 3). Note that crossing back and forth from one area to another increases our measure of activity but not exploration. We quantified boldness as time in center (Experiment 1), response to a looming stimulus (Experiment 2), and latency to cross into a novel environment (Experiment 3)

816 **Fig 3** Behavior in an artificial environment open field test (Experiment 1). Frogs exhibited

817 repeatability in (a) exploration (R= 0.278, p= 0.029) and (b) activity (R= 0.279, p= 0.032) across

818 two rounds separated by five weeks. Three frogs are highlighted to showcase consistent

819 individual differences in and the relationship between exploration and activity

Fig 4 Behavior in a naturalistic open field before, during, and after a threat stimulus (Experiment 2). Exploration (a) and activity (b) decrease during stimulus presentation compared to pre-stimulus and post-stimulus ($F_{2,230}$= 112.143, p< 0.001; $F_{1,234}$= 170.128, p< 0.001), and these patterns were repeatable across the two rounds of the experiment. Data are collapsed across rounds, as no significant effect of round or a trial by round interaction were observed for exploration or activity. The boxes show the median, the first and third quartiles, and whiskers as 1.5x the interquartile range. Dots show individual data points.

Fig 5 Behavior in a familiar, naturalistic environment and during the novel environment test

(Experiment 3). (a) Activity increased with repeated experience in the assays (i.e., across rounds

for the OFT, AFET, and NET; ($F_{2,143}$= 3.919, p= 0.022). (b) All frogs explored the novel area more

quickly in subsequent rounds ($F_{2,36}$ = 13.100, p< 0.001), and (c) more active frogs explored the

novel area sooner (corr = -0.655, p< 0.001). Letters in (a) and (b) indicate significant pair-wise

differences between rounds. The boxes show the median, the first and third quartiles, and

whiskers as 1.5x the interquartile range. Dots show individual data points.