Individual Behavioral Variability Across Time and Contexts in *Dendrobates tinctorius*

Poison Frogs

Katharina M Soto[1,2], Faith O Hardin[1], Harmen P Alleyne[1], Eva K Fischer[1,3]

[1] Department of Evolution, Ecology, and Behavior; School of Integrative Biology; University of

Illinois Urbana-Champaign

[2] Program of Evolution, Ecology, and Conservation; School of Integrative Biology; University of

Illinois Urbana-Champaign

[3] Carl R. Woese Institute for Genomic Biology, University of Illinois Urbana-Champaign, Urbana

IL 61801, USA

*Corresponding author:

Eva K Fischer

Department of Evolution, Ecology, and Behavior

School of Integrative Biology

University of Illinois Urbana-Champaign

505 S Goodwin Ave

Urbana, IL 61801

efischer@illinois.edu

Abstract

Consistent individual differences in behavior (aka, "animal personality") have consequences for

individual fitness, adaptive trajectories, and species' persistence. Such differences have been

documented across a wide range of animals, though amphibians are generally

underrepresented in this research area. The aim of our study was to examine consistent

individual differences in poison frogs (*Dendrobates tinctorius*). We evaluated repeatability in

behaviors including activity, exploration, and boldness to assess consistency of behaviors

across different temporal, experimental, and environmental contexts. We found repeatability in

activity and exploration across time and contexts. In contrast, we observed context-specific

behavior for our metrics of boldness, with consistent individual differences only for some

measures. Further, while activity and exploration displayed consistent correlations across

contexts, relationships between activity and boldness were context dependent. Our findings

document the presence of consistent individual differences in behavior in poison frogs,

challenging historic assumptions about the simplicity of amphibian behavior. Nonetheless, our

approach testing the same individuals across multiple time points and assays also reveals

context-dependent differences, highlighting the complex relationship between consistent

individual differences and context-specific responses in animal behavior.

Statement of Significance

The concept of animal personality centers on the existence of consistent individual differences

in behavior. However, behavioral responses can also depend on the context, and consistent

individual differences in one context do not guarantee consistent differences in another. To

address this question, we assessed activity, exploration, and boldness in captive-bred poison

frogs (*Dendrobates tinctorius*) across time and environmental contexts. Our comprehensive

approach revealed consistent individual differences in some behaviors and context-specificity in

others. While activity and exploration were generally repeatable and correlated with one

another, boldness was not. Especially in view of the emphasis on measures of boldness in the

animal personality literature, our findings emphasize the importance of reiterative and holistic

approaches in the study of animal behavior.

Keywords: behavioral syndromes, animal personality, repeatability, individual variation,

Dendrobatid

Introduction

There is growing evidence that consistent individual differences in behavior exist across taxa, and that these differences can have important consequences for individual fitness, adaptive trajectories, and species' persistence (Bell et al. 2009). For example, bold and aggressive individuals may be superior at territory defense but inferior at parental care (Duckworth 2006). Such trade-offs can have consequences for individual fitness, leading to the establishment of alternative behavioral strategies within populations and have been argued to constrain adaptive trajectories (Smith and Blumstein 2008; Dochtermann and Dingemanse 2013). Moreover, certain personality types may fare better in the face of increasing human-induced global change, for example if they are better able to adapt to increasingly urban habitats (Pamela et al. 2015; Murgui and Hedblom 2017) or more likely to invade novel environments (Brodin et al. 2013; Carvalho et al. 2013). Increased recognition of the complex consequences of consistent individual behavioral differences have led to an increase in the study of animal personalities across taxa. Animal personality is increasingly invoked to explain behavioral differences in natural and captive populations, and to speculate on how these differences may influence survival and adaptation. However, many studies do so without confirming consistent individual differences across multiple contexts and over time. Assuming – rather than testing – that individual behavioral variation is stable across contexts and time can introduce bias, leading to adaptive inferences based on isolated incidents or context-specific behavior (Bell et al. 2009).

Animal personality is broadly defined as consistent individual differences in behavior (Réale et al. 2010; Carere and Maestripieri 2013). These differences are typically characterized by quantifying behavioral repeatability, defined as the proportion of variance explained by within-individual variation compared to between-individual variation (Bell et al. 2009; Bergmüller 2010; Garamszegi and Herczeg 2012). Animal personality encompasses an array of behavioral traits, often with a particular focus on boldness:shyness, exploration:avoidance, and activity

(Réale et al. 2007; Garamszegi and Herczeg 2012; Michelangeli et al. 2019). Further, these

traits can become "packaged together" into behavioral syndromes (Sih et al. 2004). For

instance, studies spanning diverse taxa such as common voles (*Microtus arvali*) (Herde and

Eccard 2013), convict cichlids (*Amatitlania nigrofasciata*) (Mazué et al. 2015), sea turtles

(*Chelonia mydas)* (Kudo et al. 2021), and great tits (*Parus major*) (Cole and Quinn 2011)

document positive correlations between boldness, activity, and exploration.

While the presence of animal personalities and behavioral syndromes have been

documented in animals from primates to beetles, there is bias in taxonomic representation.

While there is extensive research in non-human animals including birds (e.g., Réale et al. 2007;

Carere and Locurto 2011) and fish (e.g., Bell 2005; Smith and Blumstein 2010; Mazué et al.

2015), amphibians have received comparatively less attention (but see Peignier et al. 2022;

Sonnleitner et al. 2022). These biases arise in part from historical assumptions about the

(behavioral) complexity of different animals. Amphibians were long viewed as behaviorally

simple and stereotyped (Stamps 2007; Sih and Del Giudice 2012), yet evidence is accumulating

that these creatures make complex, context dependent behavioral decisions (Kelleher et al.

2018). Studying individual behavioral variation in amphibians is critical for advancing our

understanding of how these animals navigate risks and opportunities, such as the trade-off

between mating advertisement boldness and predator avoidance (Stamps 2007; Sih and Del

Giudice 2012). Importantly, this knowledge has broader implications for understanding adaptive

strategies, ecological interactions, and conservation needs of amphibians, which is of particular

importance given the alarming global decline of amphibian populations (Kelleher et al. 2018).

Among anurans (frogs and toads), poison frogs (family Dendrobatidae) offer compelling

study subjects for exploring individual variation and correlations in behavior. Poison frogs are

known for their bright coloration, toxicity, and interesting behaviors. These species' vivid hues

are an honest signal of their toxicity, and both traits have coevolved with bold behavior and a

conspicuous, diurnal lifestyle as compared to most anurans (Summers 2003; Maan and

Cummings 2012). Poison frogs exhibit a rich array of complex behaviors, encompassing anti-

predator defenses (Breed 2008), aggressive territoriality (Crothers and Cummings 2015),

parental care (Roland and O'Connell 2015; Fischer et al. 2019), spatial learning and navigation

(Liu et al. 2016; Pašukonis et al. 2022), and behavioral plasticity (Ringler et al. 2017; Peignier et

al. 2023). Taken together, these characteristics have garnered poison frogs increasing attention

in the realm of animal personality and behavioral syndromes. Existing studies in poison frogs

have examined consistent individual differences in activity (Sonnleitner et al. 2022), antipredator

behavior (Peignier et al. 2023), and aggression (Chaloupka et al. 2022). Despite these initial

insights, there remains a substantial knowledge gap regarding the extent of consistent individual

differences across time and context in poison frogs specifically, and amphibians more generally.

We used the Dyeing Poison Frog, *Dendrobates tinctorius*, to investigate personality and

behavioral syndromes across contexts and timeframes. We repeatedly assessed activity,

exploration, and boldness – in the same and in different assays – to test whether commonly

used personality metrics are robust to differences in assay design and timing (Fig. 1). Following

the methodologies of Kelleher et al. (2018), our experiments included an open field test, a threat

stimulus test, a novel environment test, and an activity in a familiar vs novel environment test

(Fig. 1). All experiments were built upon the open field test, a widely utilized method for

assessing activity levels, exploratory behavior, and boldness across taxa (Bell et al. 2009;

Carere and Locurto 2011).

In Experiment 1, we conducted an open field test in an artificial environment to establish

baseline activity, exploration, boldness, and the relationships among them (Fig. 1). This test

reflects the standard, basic design used for open field tests. In Experiment 2, we transitioned to

an open field test in a naturalistic environment, additionally assessing behavior before, during,

and after exposure to a threat stimulus (Fig. 1). In Experiment 3, we maintained the naturalistic

environment, but extended the duration of each trial, and directly compared behavior in a

familiar, naturalistic versus a novel, artificial environment (Fig. 1). Importantly, individuals were

assayed multiple times in each experiment (i.e., experienced multiple 'rounds' of each 'trial type'

- terminology we use throughout the manuscript). This design allowed us to assess repeatability

of commonly used behavioral metrics within and across contexts, and in response to multiple

stimuli.

We predicted consistent individual differences in activity, exploration, and boldness

across multiple rounds and the three distinct experiments. Further, we anticipated a behavioral

syndrome, in which frogs with higher activity levels would also display increased boldness and

exploration. Finally, we expected frogs to perceive the threat stimulus and the novel

environment as stressful and exhibit context-dependent behavioral responses. Our overarching

goal was to provide a comprehensive, nuanced understanding of how consistent individual

differences manifest across diverse time and context, and to reflect on the implications of these

patterns for extrapolation to ecological and evolutionary outcomes.

Methods

Animals

D. tinctorius were housed in our animal facility at the University of Illinois Urbana-

Champaign. Terraria contained soil, moss, leaf litter, live plants, coconut shelters, and a small

pond. For housing and all experiments, we maintained terraria and test arenas at 21°–23° C

and 80%–90% humidity on a 12L:12D light cycle. We fed frogs *Drosophila hydei* or *Drosophila*

melanogaster dusted with vitamins three times weekly. All animal care and experimental

procedures were approved by the University of Illinois Animal Care and Use Committee (IACUC

protocol #20147).

All behavioral experiments were conducted between March and April 2022. We used a

total of 48 captive bred frogs (N= 20 adult males, N= 12 adult females, and N= 17 juveniles of

unknown sex). All frogs were assayed in Experiments 1 and 2. In Experiment 3, we used a

subset of 19 frogs (N= 6 adult males, N= 8 adult females, and N= 5 juveniles; see additional

details below). For Experiments 1 and 2, we ran frogs sequentially in groups of five per week,

excluding the last group which had only four frogs. We followed the same protocol in a second

round of trials with the same groups of frogs. In essence, each individual underwent two

identical rounds, with a 5-week interval in between. For Experiment 3, we ran frogs in groups of

9 per week through a series of sequential trials that lasted a combined 25 hours (details below).

These trials were repeated three times with three weeks between each round. Although frogs

varied in age, all individuals were sexually inexperienced. Previous studies have demonstrated

that fed or digesting amphibians exhibit decreased levels of activity compared to fasted

individuals (Andrade et al. 2005). To ensure that the frogs were not actively digesting during the

experiments, we fasted frogs for 24 hours before behavioral trials.

Experiment 1: Artificial open field test

In Experiment 1, we investigated exploration, activity, and boldness in an open field test

in a novel, artificial environment (Fig. 1). The open field arena was a circular enclosure with a

diameter of 108 mm. To maintain humidity and moisture, we lined the arena with white bench

paper soaked in reversed-osmosis (RO) water. For each observation, we placed a single frog at

the center of the arena as a standardized starting point and recorded behavior for 15 minutes

using an overhead camera (DVC, HDV-604S). Individuals were immediately returned to their

home terraria following trials. We quantified exploration as the proportion of areas a frog visited,

activity as the number of times a frog crossed the line between areas, and boldness as time

spent in the center of area (Fig. 2) (additional details below).

Experiment 2: Naturalistic open field and threat stimulus test

In Experiment 2, we quantified activity and exploration in a naturalistic open field. The

behavioral arena was a 305 x 205 x 203 mm plexiglass tank that mimicked the frogs' home

terraria, including soil substrate, moss, leaves, and a live plant positioned in one corner. The

arena was completely covered from the outside with white contact paper, excluding a 240 x

169.5 mm exposed area opposite the plant for the presentation of the threat stimulus. As in

Experiment 1, we placed a single frog at the center of the arena at the beginning of each

observation and used an overhead camera to record behavior. We conducted three trials to

investigate behavior (1) pre-stimulus open field test (15 minutes after introduction to arena), (2)

during threat stimulus test (3 minutes of looming stimulus), and (3) post-stimulus open field test

(for 15 minutes after the stimulus).

We assessed activity and exploration using the same methods as in Experiment 1 (Fig.

2). Additionally, we measured boldness by observing the frogs' responses to a looming stimulus

(Fig. 2). Previous studies using leopard frogs (*Lithobates pipiens*) have shown reliable

avoidance behavior in response to looming objects (Ingle 1990; Waldeck and Gruberg 1995).

To present the looming stimulus, we used an iPad (Apple, 6th generation). The iPad was

covered in white tape, except for the display screen, and placed in the exposed area of the

arena (as shown Fig. 1). The looming stimulus was an expanding black circle projected

repeatedly onto the iPad screen for a duration of three minutes. To prevent habituation, random

pauses ranging from 1 to 30 seconds were incorporated into the stimulus presentation. During

the pre- and post- exposure trials, the iPad displayed a white screen to match the surrounding

environment. During the looming stimulus, we recorded activity and exploration, as well as

boldness based on whether the frogs approached or displayed avoidance behavior in response

to the stimulus.

Experiment 3: Naturalistic open field and novel environment test

In Experiment 3, we increased the duration of each trial to capture more prolonged

patterns in the frogs' behavior. To conduct these longer trials concurrently with Experiments 1

and 2 while also maintaining our repeated measures design, we selected a subset of individuals

(N= 19) from the previous experiments. The behavioral arena was constructed using the same

materials as Experiment 2. However, the dimensions of the arena were twice as long (610 x 305

x 203 mm) and the arena split into two halves. One side of the arena mimicked the frogs' home

terraria as in Experiment 2. The other side mimicked the novel, artificial environment of

Experiment 1, with only white, moistened paper flooring. The two sides of the arena were

separated by a white cardboard barrier, which could be removed without disturbing the frog or

the soil structure of the natural chamber.

We conducted three trials to investigate the frogs' activity and boldness: (1) baseline

activity in the novel, naturalistic environment (the first 90 minutes after introduction to the area),

(2) activity in after familiarization to the naturalistic environment (60 minutes after a 24 hour

acclimation period in the arena), and (3) exposure to a novel, artificial environment (60 minutes

after the removal of the barrier between the familiar and novel environment) (Fig. 1). Trials were

conducted sequentially, with each frog remaining in the arena for 25 hours while the trials

progressed. We conducted two subsequent rounds of trials for each individual, with three weeks

between each round. Due to the extended duration of each trial, we focused our analyses on

activity and boldness. We quantified activity using the same methods as in Experiment 1 and 2,

number of line crosses. To assess boldness, we measured the frogs' latency to cross into the

novel environment when the barrier was lifted (Fig. 2).

Data analysis

We describe general methods here with modifications for each experiment below. We

coded videos using Behavioral Observation Research Interactive Software (BORIS) ver. 8.13

(Friard and Gamba 2016). Experiment type was not blinded because assay identity was obvious

from videos, but all analyses were blind to individual identity. Behavioral data was analyzed with

R v.4.2.4 (R Core Team, 2022) in Rstudio v.4.2.2. (RStudio Team 2021). For all response

variables that were normally distributed, we used the raw data for analysis. Response variables

235 that did not meet expectations of normality were transformed as described below. For all

236 analyses, outcomes were considered significant at p < = 0.05.

237   We used the lmer function in the lmerTest package (Kuznetsova et al. 2017) for all linear

238 models, and always included frog ID as a random effect and round (i.e., repeated

239 measurements in the same assay) as a fixed effect. Additional fixed effects varied based on

240 experiment (see below). We tested for the effects of sex on behavior and found no significant

241 results. Therefore, and because not all individuals could be sexed at the time of the experiment,

242 we did not include sex as a factor in the analyses described below. We ran Type III ANOVAs on

243 linear models to generate test statistics and p-values for main effects and interactions.

244 Subsequently, we used the emmeans package (Lenth 2021) to run pairwise post-hoc

245 comparisons where appropriate.

246   We performed repeatability analyses using the rptR package (Stoffel et al. 2017), with

247 95% confidence intervals (CI) based on 1000 bootstrapping runs. Frog ID was used as a

248 random effect for repeatability (R) among individuals. We used Pearson correlation for normally

249 distributed data and Kendall correlation for non-normally distributed data to assess relationships

250 between behaviors. The correlation coefficients were interpreted using these criteria: ≈0.1 a

251 small effect, ≈0.3 a moderate effect, and>0.5 a strong effect (Garamszegi et al. 2013).

253 *Analysis Experiment 1: Artificial open field test*

254   We coded videos in BORIS using a transparent overlay that divided the behavioral arena

255 into 40 areas: 24 on the ground and 16 on the wall. From videos, we quantified exploration as

256 the proportion of areas a frog visited and activity as the number of times a frog crossed a line

257 between areas. Movement between areas was recorded when the majority of the frogs' body

258 entered into a new area. We measured boldness as time spent in the center four areas of the

259 arena (Fig. 2). We log transformed the activity and boldness data to improve normality. We ran

260 linear models for each behavior (activity, exploration, boldness) as described above, including

round as a fixed effect and frog ID as a random effect. We performed repeatability analysis with

Gaussian distributions for exploration and Poisson distributions for activity and boldness (Stoffel

et al. 2017). Data from 11 individuals was lost due to equipment failure, leaving a final sample

size of N=38 individuals. Analyses and results are based on these samples.

Analysis Experiment 2: Naturalistic open field threat stimulus test

We coded videos in BORIS using a transparent overlay that divided the behavioral arena

into 32 areas: 16 on the ground and 16 on the wall. We quantified and analyzed activity and

exploration as in Experiment 1, with the addition of a fixed effect of trial (pre-stimulus, during

stimulus, and post-stimulus) to test for pre- and post- stimulus behavior differences. To compare

trials of different duration, we divided each behavior by the total trial time.

For response to stimulus, which we used as a measure of boldness, we divided the

arena into three sections: (1) at stimulus (the two wall areas overlaying the iPad, the two floor

areas directly in front the iPad, and one wall area immediately to the left of the iPad), (2) near

stimulus (the two wall areas and the four floor areas surrounding the "at stimulus" area), and (3)

away from stimulus (the remaining areas) (Fig. 2). We noted the frogs' position before and after

stimulus presentation. We used a Chi-Squared test in the Tidyverse package (Wickham et al.

2019) for pairwise comparisons of responses (moved away, moved toward, or froze) (R Core

Team 2022). We fit rptR models with Gaussian distributions to test for repeatability of each

behavior across rounds and across trials (Stoffel et al. 2017).

Analysis Experiment 3: Naturalistic open field novel environment test

We coded videos in BORIS using a transparent overlay that divided the behavioral arena

into 96 areas: 16 areas on both the natural and novel arena floor, and 16 areas on the walls of

each chamber. We recorded activity as the number of times a frog crossed the line between

areas as in Experiments 1 and 2. After the divider was removed, we recorded the time it took for

a frog to move into the novel area as latency to explore, which we interpreted as a measure of

boldness (Fig. 2). We recorded activity on both sides of the chamber after the barrier was

removed. We ran linear models for activity and latency to explore, including round (i.e., repeated

measurements in the assay), trial (baseline, after acclimation, in a novel environment), and trial:

round interaction as fixed effects, and frog ID as a random effect. As in Experiment 1, behaviors

were divided by trial duration when analyzing trial effects.

To determine whether activity predicted an individual's latency to cross into a new area,

we filtered the data to only include observations taken during the novel environment test. We

then fit a similar linear mixed effect model to predict latency to enter the novel area with fixed

effects of activity, round, and their interaction, and frog ID as a random effect. We fit rptR

models with Gaussian distributions to test for repeatability of each behavior across rounds and

across trials (Stoffel et al. 2017).

Cross-experiment analysis of activity and exploration

In addition to behavioral analyses within experiments, we were interested in the behavior

of individuals across experiments. To compare exploration between Experiments 1 and 2, which

had different arena sizes, we calculated the proportion of areas explored. We did not quantify

exploration in Experiment 3, and it was therefore excluded from this analysis (Fig. 2). We

compared the ratio of areas explored in Experiment 1 to both the pre-stimulus and post-stimulus

exploration in Experiment 2 using linear mixed models, with exploration rate and experiment as

fixed effects (Bates et al. 2015). We tested repeatability using a Gaussian distribution (Stoffel et

al. 2017).

In all experiments, we measured activity by counting the number of line crosses (Fig. 2).

To compare across experiments of different duration, we divided the number of lines crossed by

the total trial time to calculate a line crossing rate. We compared the subset of individuals (n=

19) in Experiment 3 with the same individuals from Experiments 1 and 2. Baseline activity

measurements were similar across all experiments, taken prior to either the presentation of a

stressor (looming stimulus test) or an environmental stressor (novel environment test) (Fig. 1).

We also compared the post-stimulus activity in Experiment 2 to the activity observed during the

novel environment test in Experiment 3, reasoning that these trials were similar in representing

responses following an environmental change. We analyzed behavioral differences using

activity rate and experiment as fixed effects and frog ID as a random effect (Bates et al. 2015).

We calculated post hoc comparisons and repeatability as before.

Results

Experiment 1: Artificial open field test

324   Frogs displayed repeatability in exploration (R= 0.278, p= 0.029) (Fig. 3a) and activity

(R= 0.279, p= 0.032) (Fig. 3b) and across both rounds separated by five weeks. However, time

in the center of the arena, which we interpret as a measure of boldness, was not repeatable (R=

0.044, p= 0.395). No significant effect of round was observed for activity ($F_{1,37}$= 3.390, p= 0.074)

or exploration ($F_{1,37}$= 1.392, p= 0.245, or boldness ($F_{1,74}$= 0.628, p= 0.431), suggesting that

frogs' behavior did not change in response to previous experience in the arena. Activity and

exploration exhibited a strong positive correlation (corr= 0.767, p< 0.001). Boldness was not

correlated with either activity (corr= -0.150, p= 0.196) or exploration (corr= -0.119, p= 0.307)

(Table 1).

Experiment 2: Naturalistic open field and threat stimulus test

335   Activity and exploration were repeatable across pre- and post- stimulus trials in both

round 1 (activity: R= 0.593, p< 0.001; exploration: R= 0.421, p< 0.001) and round 2 (activity: R=

0.529, p< 0.001; exploration: R= 0.449, p< 0.001). We observed the same trend across rounds

within trial (activity: R= 0.123, p= 0.006; exploration: R= 0.177, p< 0.001), aligning with the

results from Experiment 1 (Table 1). In contrast, during the looming stimulus presentation, there

was no repeatability of activity (R= 0.201, p= 0.069) or exploration (R= 0.175, p= 0.145) across

rounds.

We found an effect of trial on exploration ($F_{2,230}$= 112.143, p< 0.001) (Fig. 4a) and

activity ($F_{1,234}$= 170.128, p< 0.001) (Fig. 4b), with frogs decreasing behaviors during the stimulus

and then increasing post-stimulus (Fig. 4). We observed no significant round effect for activity

($F_{1,232}$= 0.570, p= 0.451) or exploration ($F_{1,234}$= 0.292, p= 0.590), nor an interaction between

round and trial ($F_{2,228}$= 0.506, p= 0.604; $F_{2,228}$= 0.216, p= 0.801), indicating that frogs behaved

consistently across multiple experiences in the arena. Activity and exploration were highly

correlated across all trials (pre-stimulus, during stimulus, and post-stimulus) (corr= 0.7–0.8, p<

0.001).

We evaluated boldness by quantifying frogs' responses to the looming stimulus. Frogs'

responses to the looming stimulus were not repeatable across rounds (R= 0, p= 1). We note this

is likely because overall variation was low, as the majority of frogs demonstrated a freezing

response during stimulus presentation rather than moving away ($\chi^2_{1,96}$= 19.755, p< 0.001) or

towards ($\chi^2_{1,96}$= 41.796, p< 0.001) the stimulus. No significant effect of round was observed for

boldness ($\chi^2_{2,96}$= 22.429, p= 0.401).

We found significant correlations between boldness (response to stimulus) and frogs'

activity (tau= 0.155, p= 0.002) and exploration (tau= 0.155, p= 0.002) during the pre- and post-

stimulus trials. These trends suggest that frogs displaying higher levels of activity and

exploration prior to and after the stimulus were also more inclined to approach the stimulus

during the threat stimulus trial. These correlations between boldness, activity, and exploration

contrast with the results obtained in Experiment 1 (Table 1).

Experiment 3: Naturalistic open field and novel environment test

Frogs were repeatable for activity, both across trials (R= 0.321, p<0.011) and across

rounds (R= 0.651, p= 0.001). We found that frogs moved more with repeated experience in the

arena ($F_{2,143}$= 5.971, p= 0.003), with Round 1 having significantly less movement than Round 2

(t_{36}=-3.423, p=0.004) and Round 3 (t_{36}= -3.784, p= 0.002), but no significant difference between

Round 2 and 3 (t_{36}= -0.360 p=0.931) (Fig. 5a). Similarly, we found that frogs took less time to

cross the boundary into the novel area as rounds progressed ($F_{2,36}$= 13.100, p< 0.001), with

frogs taking longer to cross in Round 1 compared to Round 2 (t_{36}= 2.837, p= 0.019) and Round

3 (t_{36}= 5.108, p< 0.001), and was approaching significance between Round 2 and 3 (t_{36}= 2.270,

p= 0.073). We found significant repeatability for latency to explore across rounds (R= 0.388, p=

0.003) suggesting all frogs responded to repeated experience similarly.

We found a significant negative correlation between activity and boldness (*e.g.,* latency

to cross into novel environment; corr= -0.665, p< 0.001). Additionally, we found that boldness

was influenced by both activity ($F_{1,38}$= 28.795, p<0.001) and round ($F_{2,38}$= 6.546, p= 0.004),

indicating that more active frogs moved into the novel area more quickly, and all frogs moved

into the novel area more quickly with repeated experience.

Cross-experiment analysis of activity and exploration

Frogs exhibited consistent individual differences in exploration (R= 0.171, p= 0.005)

across Experiments 1 and 2. For activity, baseline activity was repeatable between the

naturalistic environments of Experiment 2 and Experiment 3 (R= 0.207, p< 0.001), and between

the artificial environment of Experiment 1 and the naturalistic environment of Experiment 3 (R=

0.171, p= 0.041). Activity was not repeatable between the artificial environment of Experiment 1

and the naturalistic environment of Experiment 2 (R= 0.029, p= 0.401). Activity was repeatable

across stressor trials (Fig. 1) (i.e., threat stimulus in Experiment 2 and novel environment in

Experiment 3) (R= 0.243, p= 0.003) (Fig. 5b).

Discussion

We investigated animal personality and behavioral syndromes in captive-bred poison

frogs (*Dendrobates tinctorius*) by characterizing behavioral differences across multiple

environmental contexts and timepoints. We conducted three experiments to assess activity,

exploration, and boldness in the same individuals across varying conditions. We found

repeatability in activity and exploration across time and contexts. In contrast, we observed

context-specific behavior for our metrics of boldness, with consistent individual differences only

for some measures of boldness. The stability of individual differences in behavior can carry

profound implications for how species adapt to shifting environmental conditions (Wong and

Candolin 2015; Merrick and Koprowski 2017), thereby shaping the evolutionary trajectory and

ecological interactions of the species (Shaw 2020; Stiegler et al. 2022). Despite their declining

populations, amphibians remain relatively understudied in this research area. The results of our

work help to fill this gap, address broader issues in the field of "animal personality", and set the

stage for future work in *D. tinctorius*.

Our study revealed widespread repeatability for activity and exploration, evident across

time, trials, and experimental contexts. These findings suggest that measures of activity and

exploration are indeed reliable, and measurements in one assay or at one timepoint are likely to

predict an individual's behavior in another context or at another time. We also recovered the

commonly described activity:exploration behavioral syndrome (Réale et al., 2007). While we

expected a positive relationship between activity and exploration (i.e., exploration is impossible

without movement), we note that there is room for variability in this relationship. For instance,

frogs might have displayed a lot of movement but only within a restricted area, leading to

heightened activity but limited exploration. The strong, significant relationships we recover

(correlations > 0.76), suggest that these behaviors are indeed tightly linked and can therefore

predict one another as well as some other behaviors (e.g., latency to explore a novel area, see

below) under certain circumstances.

The above outcomes align with prior research on animal personality across taxa (Réale

et al. 2007; Kelleher et al. 2018), and complement a recent investigation on another

Dendrobatid, *Allobates femoralis*, which demonstrated the efficacy of both artificial and

naturalistic setups in capturing consistent individual differences in behavior (Bégué et al. 2023).

However, a study on *D. tinctorius* froglets by Sonnleitner et al. (2022) yielded somewhat

different findings: the authors found significant repeatability at the family level but no significant

repeatability at the individual level. These results suggests that genetic factors influence

behavior and this effect outweighs individual differences when related individuals are reared

under similar conditions. We were unable to control for relatedness in our study, but assume

genetic differences explain at least some of the behavioral variation we document. A non-

mutually exclusive explanation is that individuals in the study of Sonnleitner et al. (2022) were

much younger than those used here, and individual differences in behavior may become more

pronounced with age and experience (Bell et al. 2009). In any case, while consistent individual

differences are being increasingly documented across diverse taxa, contrasting results even in

the same species underscore the importance of considering various contexts and timepoints

when extrapolating from single studies to evolutionary and ecological outcomes.

While we observed a high degree of repeatability in activity and exploration across time

and contexts, the behaviors we considered as measures of boldness demonstrated contextual

dependence. Our first measure was the commonly used metric of time spent in the center of a

novel arena (Carter et al. 2012; Yuen et al. 2017; Baker et al. 2018; Rudolfová et al. 2022). We

did not observe consistent individual differences in this measure, nor was it correlated with other

behaviors. As our next measure of boldness, we assessed response to a threat stimulus in a

familiar environment, aligning with prior studies (Ingle 1990; Waldeck and Gruberg 1995; Carter

et al. 2012; Yuen et al. 2017). Once again, we did not find repeatability for boldness, although

we did observe a correlation between this measure of boldness, activity, and exploration. In our

final approach, we measured boldness as the latency to enter a novel environment, a less

443 common yet established method for evaluating boldness (White et al. 2013; Magnhagen et al.

444 2014; Theódórsson and Ólafsdóttir 2022). With this approach, we observed both repeatability

445 and a correlation between boldness and activity. Taken together, our results suggest that

446 commonly used boldness methodologies may not measure the same personality trait.

447  Despite the popularity of measuring boldness in animal personality studies, our findings

448 align with those of others who have explicitly compared outcomes across different metrics

449 categorized as 'boldness' and found conflicting results. For example, our approach and

450 outcomes are similar to that of Yuen et al. (2017) in their study on African striped mice

451 (*Rhabdomys pumilio*). The authors investigated whether distinct experimental methods were

452 indeed measuring boldness. Similar to our approach, they employed both artificial and natural

453 environments, and conducted two boldness tests, an open field, and a startle test. Yuen et al.

454 (2017) highlighted the distinction between these methods. The startle test evaluated risk-taking

455 behavior after threat exposure, potentially measuring anxiety rather than boldness, while the

456 open field test did not incorporate a preceding threat and proved to be a more precise method

457 for gauging boldness in their species. Carter et al. (2012) found related results when examining

458 chacma baboons (*Papio ursinus*), concluding that threatening situations may measure anxiety,

459 not boldness. Moreover, the use of artificial open field tests, particularly in evaluation of time

460 spent in the center, may not consistently capture boldness. Indeed, a recent study on Zebrafish

461 (*Danio rerio*) suggests that thigmotaxis may not effectively measure boldness or anxiety-like

462 behavior in this species (Rajput et al. 2022).

463  Translating these observations to our own results, our open field test focused on

464 evaluating time in center, the threat stimulus test observed behavior in the presence of a threat,

465 whereas the novel environment test examined behavior in a new and unfamiliar environment.

466 While we initially considered all tests as measures of boldness, our data suggest that the

467 specific nature of the challenges presented in each experiment and the environmental context

468 led to varying types and intensities of behavioral responses, and in turn distinct patterns of

repeatability and behavioral correlations.

Our findings also confirm the importance of considering the environment when

measuring behavioral correlations: we observed correlations between boldness, activity, and

exploration (i.e., behavioral syndromes) in naturalistic environments (Experiments 2 and 3) but

not in artificial settings (Experiment 1) (Table 1). Alongside those of others, our observations

highlight that the environment in which individuals are assessed can exert a profound influence

on their behavior (e.g., Niemelä and Dingemanse 2014; Bégué et al. 2023) and the necessity of

evaluating individual variation across diverse contexts (Bell et al. 2009).

Conclusions

Our findings demonstrate consistent individual differences in behavior across time and

context in *D. tinctorius*, while also highlighting the subtle complexities of studying animal

personality. While we found strong evidence for individual consistency in activity and

exploration, our measures of boldness were variable and context dependent. Our results

demonstrate the value of assessing multiple behavioral traits and the relationships among them

to build a comprehensive view of behavioral repeatability and variability within and across

species (Sih et al. 2004; Bell et al. 2009; Sih and Del Giudice 2012). Broadly, our results add to

the growing literature emphasizing the importance of evaluating context-dependent patterns of

individual behavioral variation, even for assay commonly considered as measuring the same

trait (Bell et al. 2009; Carter et al. 2012, Yuen et al. 2017). More specifically, our findings

contribute to the growing recognition of behavioral complexity in anuran amphibians, with

implications for amphibian research, including their role as ecological indicators amid habitat

degradation and climate change (Grant et al. 2019). Our findings here pave the way for future

research on the impact of consistent individual differences on reproductive success, survival,

and community dynamics.

Acknowledgments

497 We thank the members of the Fischer Lab for feedback on previous versions of the

498 manuscript and help with frog husbandry. For their insight on statistical analyses, we thank Drs.

499 Jen Moss and Sarah Westrick. Our work would not be possible without the animal care support

500 of the Division of Animal Resources (DAR) at UIUC.

Funding

503 This work is/was supported by the USDA National Institute of Food and Agriculture Hatch

504 project 1026333 (ILLU-875-984 to KMS), a UIUC Graduate College Master's Fellowship (to

505 KMS), a UIUC Illinois Distinguished Fellowship (to FOH), and UIUC Laboratory Start-up funds

506 (to EKF).

Code availability

509 All raw data and analysis code will be available on Dryad upon publication.

Author Contributions

512 KMS, FOH, HPA, and EKF conceived of and planned the experiments. KMS, FOH, and HPA

513 carried out the experiments. KMS, FOH, and EKF contributed to data analysis and interpretation

514 of the results. KMS wrote the manuscript with assistance from all authors.

Conflict of Interest

517 None.

References

Andrade D V, Cruz-Neto AP, Abe AS, Wang T (2005) Specific dynamic action in ectothermic

vertebrates: a review of the determinants of postprandial metabolic response in fishes,

amphibians, and reptiles

Baker MR, Goodman AC, Santo JB, Wong RY (2018) Repeatability and reliability of exploratory

behavior in proactive and reactive zebrafish, *Danio rerio*. Sci Rep.

https://doi.org/10.1038/s41598-018-30630-3

Bates D, Mächler M, Bolker BM, Walker SC (2015) Fitting linear mixed-effects models using

lme4. J Stat Softw 67. https://doi.org/10.18637/jss.v067.i01

Bégué L, Peignier M, Ringler E (2023) The link between animal personality and habitat selection

in males of the Neotropical poison frog *Allobates femorali*s. Behaviour 160:257–274.

https://doi.org/10.1163/1568539X-bja10202

Bell AM (2005) Behavioural differences between individuals and two populations of stickleback

(*Gasterosteus aculeatus*). J Evol Biol 18:464–473. https://doi.org/10.1111/j.1420-

9101.2004.00817.x

Bell AM, Hankison SJ, Laskowski KL (2009) The repeatability of behaviour: a meta-analysis.

Anim Behav 77:771–783. https://doi.org/10.1016/j.anbehav.2008.12.022

Bergmüller R (2010) Animal personality and behavioural syndromes. Springer, Berlin,

Heidelberg

Breed M (2008) Warning colouration predicts behavioural and morphological differences in the

strawberry poison-dart frog. Thesis, Uppsala University

Brodin T, Lind MI, Wiberg MK, Johansson F (2013) Personality trait differences between

mainland and island populations in the common frog (*Rana temporaria*). Behav Ecol

Sociobiol 67:135–143. https://doi.org/10.1007/s00265-012-1433-1

Carere C, Locurto C (2011) Interaction between animal personality and animal cognition. Curr

Zool 57:491–498. https://doi.org/10.1093/czoolo/57.4.491

Carere C, Maestripieri D (2013) Animal personalities: behavior, physiology, and evolution.

University of Chicago Press

Carter AJ, Marshall HH, Heinsohn R, Cowlishaw G (2012) How not to measure boldness: Novel

object and antipredator responses are not the same in wild baboons. Anim Behav 84:603–

609. https://doi.org/10.1016/j.anbehav.2012.06.015

Carvalho CF, Leitão AV., Funghi C, Batalha HR, Reis S, Mota PG, Lopes RJ, Cardoso GC

(2013) Personality traits are related to ecology across a biological invasion. Behav Ecol

24:1081–1091. https://doi.org/10.1093/beheco/art034

Chaloupka S, Peignier M, Stückler S, Araya-Ajoy Y, Walsh P, Ringler M, Ringler E (2022)

Repeatable Territorial Aggression in a Neotropical Poison Frog. Front Ecol Evol 10:1–7.

https://doi.org/10.3389/fevo.2022.881387

Cole, Ella F, Quinn, John L (2011) Personality and problem-solving performance explain

competitive ability in the wild. Proc R Soc B 279:1168–1175

Crothers LR, Cummings ME (2015) A multifunctional warning signal behaves as an agonistic

status signal in a poison frog. Behav Ecol 26:560–568.

https://doi.org/10.1093/beheco/aru231

Dochtermann NA, Schwab T, Sih A (2014) The contribution of additive genetic variation to

personality variation: Heritability of personality. Proc R Soc B 282.

https://doi.org/10.1098/rspb.2014.2201

Duckworth RA (2006) Behavioral correlations across breeding contexts provide a mechanism

for a cost of aggression. Behav Ecol 17:1011–1019. https://doi.org/10.1093/beheco/arl035

Fischer EK, Roland AB, Moskowitz NA, Vidoudez C, Ranaivorazo N, Tapia EE, Trauger SA,

Vences M, Coloma LA, O'Connell LA (2019) Mechanisms of Convergent Egg Provisioning

in Poison Frogs. Curr Biol 29:4145-4151.e3. https://doi.org/10.1016/j.cub.2019.10.032

Friard O, Gamba M (2016) BORIS: a free, versatile open-source event-logging software for

video/audio coding and live observations. Methods Ecol Evol 7:1325–1330.

https://doi.org/10.1111/2041-210X.12584

Garamszegi LZ, Herczeg G (2012) Behavioural syndromes, syndrome deviation and the within-

and between-individual components of phenotypic correlations: When reality does not meet

statistics. Behav Ecol Sociobiol 66:1651–1658. https://doi.org/10.1007/s00265-012-1439-8

Garamszegi LZ, Markó G, Herczeg G (2013) A meta-analysis of correlated behaviors with

implications for behavioral syndromes: Relationships between particular behavioral traits.

Behav Ecol 24:1068–1080. https://doi.org/10.1093/beheco/art033

Grant EHC, Muths E, Schmidt BR, Petrovan SO (2019) Amphibian conservation in the

Anthropocene. Biol Conserv 236:543–547. https://doi.org/10.1016/j.biocon.2019.03.003

Herde A, Eccard JA (2013) Consistency in boldness , activity and exploration at different stages

of life. BMC Ecol 103:61–68. https://doi.org/10.1186/1472-6785-13-49

Ingle DJ (1990) Visually Elicited Evasive Behavior in Frogs. Bioscience 40:284–291.

https://doi.org/10.2307/1311265

Kelleher SR, Silla AJ, Byrne PG (2018) Animal personality and behavioral syndromes in

amphibians: a review of the evidence, experimental approaches, and implications for

conservation. Behav Ecol Sociobiol 72:79. https://doi.org/10.1007/s00265-018-2493-7

Kudo H, Nishizawa H, Uchida K, Sato K (2021) Boldness–exploration behavioral syndrome in

wild sub-adult green sea turtles caught at Oita, Japan. Appl Anim Behav Sci 236:105216.

https://doi.org/10.1016/j.applanim.2021.105216

Kuznetsova A, Brockhoff PB, Christensen RHB (2017) lmerTest Package: Tests in Linear Mixed

Effects Models. J Stat Softw 82:1–26. https://doi.org/10.18637/JSS.V082.I13

Lenth R (2021) emmeans: Estimated Marginal Means, aka Least-Squares Means. R package

version 1.8.5

Liu Y, Day LB, Summers K, Burmeister SS (2016) Learning to learn: Advanced behavioural

flexibility in a poison frog. Anim Behav 111:167–172.

https://doi.org/10.1016/j.anbehav.2015.10.018

Maan ME, Cummings ME (2012) Poison frog colors are honest signals of toxicity, particularly for

bird predators. The Am Nat 179. https://doi.org/10.1086/663197

Magnhagen C, Wacker S, Forsgren E, Myhre LC, Espy E, Amundsen T (2014) Context

consistency and seasonal variation in boldness of male two-spotted gobies. PLoS One 9.

https://doi.org/10.1371/journal.pone.0093354

Mazué GPF, Dechaume-Moncharmont FX, Godin JGJ (2015) Boldness-exploration behavioral

syndrome: Interfamily variability and repeatability of personality traits in the young of the

convict cichlid (*Amatitlania siquia*). Behav Ecol 26:900–908.

https://doi.org/10.1093/beheco/arv030

Merrick MJ, Koprowski JL (2017) Should we consider individual behavior differences in applied

wildlife conservation studies? Biol Conserv 209:34–44.

https://doi.org/10.1016/j.biocon.2017.01.021

Michelangeli M, Chapple DG, Goulet CT, Bertram MG, Wong BBM (2019) Behavioral

syndromes vary among geographically distinct populations in a reptile. Behav Ecol 30:393–

401. https://doi.org/10.1093/beheco/ary178

Murgui E, Hedblom M (2017) Ecology and conservation of birds in urban environments. Ecology

and Conservation of Birds in Urban Environments 1–525. https://doi.org/10.1007/978-3-

319-43314-1

Niemelä PT, Dingemanse NJ (2014) Artificial environments and the study of "adaptive"

personalities. Trends Ecol Evol 29:245–247. https://doi.org/10.1016/j.tree.2014.02.007

Pamela Delarue EM, Kerr SE, Lee Rymer T (2015) Habitat complexity, environmental change

and personality: A tropical perspective. Behav Process 120:101–110.

https://doi.org/10.1016/j.beproc.2015.09.006

Pašukonis A, Serrano-Rojas SJ, Fischer MT, et al (2022) Contrasting parental roles shape sex

differences in poison frog space use but not navigational performance. Elife 11:1–31.

https://doi.org/10.7554/eLife.80483

Peignier M, Araya-Ajoy YG, Bégué L, Chaloupka S, Dellefont K, Walsh P, Ringler M, Ringler E

(2022) Exploring links between personality traits and their social and non-social

environments in wild poison frogs. Behav Ecol Sociobiol 76.

https://doi.org/10.1007/s00265-022-03202-9

Peignier M, Bégué L, Ringler M, Szabo B, Ringler E (2023) Regardless of personality, males

show similar levels of plasticity in territory defense in a Neotropical poison frog. Sci Rep

13:1–9. https://doi.org/10.1038/s41598-023-30546-7

R Core Team (2022) R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria

Rajput N, Parikh K, Kenney JW (2022) Beyond bold versus shy: Zebrafish exploratory behavior

falls into several behavioral clusters and is influenced by strain and sex. Biol Open 11.

https://doi.org/10.1242/bio.059443

Réale D, Dingemanse NJ, Kazem AJN, Wright J (2010) Evolutionary and ecological approaches

to the study of personality. Phil Trans Biol Sci 365:3937–3946.

https://doi.org/10.1098/rstb.2010.0222

Réale D, Reader SM, Sol D, McDougall PT, Dingemanse NJ (2007) Integrating animal

temperament within ecology and evolution. Biol Rev 82:291–318

Ringler E, Barbara Beck K, Weinlein S, Ludwig H, Ringler M (2017) Adopt, ignore, or kill? Male

poison frogs adjust parental decisions according to their territorial status. Sci Rep 7:1–6.

https://doi.org/10.1038/srep43544

Roland AB, O'Connell LA (2015) Poison frogs as a model system for studying the neurobiology

of parental care. Curr Opin Behav Sci 6:76–81.

https://doi.org/10.1016/j.cobeha.2015.10.002

RStudio Team (2021) RStudio: Integrated Development Environment for R. RStudio, PBC,

Boston, MA

Rudolfová V, Petrásek T, Antošová E, Frynta D, Landová E, Valeš K, Nekovářová T (2022)

Inter-individual differences in laboratory rats as revealed by three behavioural tasks. Sci

Rep 12:1–11. https://doi.org/10.1038/s41598-022-13288-w

Shaw AK (2020) Causes and consequences of individual variation in animal movement. Mov

Ecol 8:1–12. https://doi.org/10.1186/s40462-020-0197-x

Sih A, Bell A, Johnson JC (2004) Behavioral syndromes: An ecological and evolutionary

overview. Trends Ecol Evol 19:372–378. https://doi.org/10.1016/j.tree.2004.04.009

Sih A, Del Giudice M (2012) Linking behavioural syndromes and cognition: A behavioural

ecology perspective. Phil Trans Biol Sci 367:2762–2772.

https://doi.org/10.1098/rstb.2012.0216

Smith BR, Blumstein DT (2008) Fitness consequences of personality: A meta-analysis. Behav

Ecol 19:448–455. https://doi.org/10.1093/beheco/arm144

Smith BR, Blumstein DT (2010) Behavioral types as predictors of survival in Trinidadian guppies

(Poecilia reticulata). Behav Ecol 21:919–926. https://doi.org/10.1093/beheco/arq084

Sonnleitner R, Alanen E, Fouilloux C, Valkonen JK, Rojas B (2022) Genetic factors influence

behavioural repeatability in juvenile poison frogs. bioRxiv 2022.09.19.508512; doi:

https://doi.org/10.1101/2022.09.19.508512

Stamps, Judy A (2007) Growth-mortality tradeoffs and 'personality traits' in animals. Ecol Lett

10:355–363. https://doi.org/10.1111/j.1461-0248.2007.01034.x.

Stiegler J, Lins A, Dammhahn M, Kramer-Schadt S, Ortmann S, Blaum N (2022) Personality

drives activity and space use in a mammalian herbivore. Mov Ecol 10:1–12.

https://doi.org/10.1186/s40462-022-00333-6

Stoffel MA, Nakagawa S, Schielzeth H (2017) rptR: repeatability estimation and variance

decomposition by generalized linear mixed-effects models. Methods Ecol Evol 8:1639–

1644. https://doi.org/10.1111/2041-210X.12797

Summers K (2003) Convergent evolution of bright coloration and toxicity in frogs. Proc Natl

Acad Sci USA 100:12533–12534. https://doi.org/10.1073/pnas.2335928100

Theódórsson BK, Ólafsdóttir GÁ (2022) Laboratory measures of boldness correlate with

ecological niche in threespine stickleback. Ecol Evol 12:1–9.

https://doi.org/10.1002/ece3.9235

Waldeck, R F, Gruberg, ER (1995) Studies on the optic chiasm of the leopard frog. Brain Behav

Evol 46:84–94

https://doi.org/10.1016/j.anbehav.2022.10.005

White JR, Meekan MG, McCormick MI, Ferrari MCO (2013) A Comparison of Measures of

Boldness and Their Relationships to Survival in Young Fish. PLoS One 8.

https://doi.org/10.1371/journal.pone.0068900

Wickham H, Averick M, Bryan J, et al (2019) Welcome to the Tidyverse. J Open Source Softw

4:1686. https://doi.org/10.21105/joss.01686

Wilson ADM, Godin JGJ, Ward AJW (2010) Boldness and reproductive fitness correlates in the

eastern mosquitofish, *gambusia holbrooki*. Ethology 116:96–104.

https://doi.org/10.1111/j.1439-0310.2009.01719.x

Wong BBM, Candolin U (2015) Behavioral responses to changing environments. Behav Ecol

26:665–673. https://doi.org/10.1093/beheco/aru183

Yuen CH, Schoepf I, Schradin C, Pillay N (2017) Boldness: are open field and startle tests

measuring the same personality trait? Anim Behav 128:143–151.

https://doi.org/10.1016/j.anbehav.2017.04.009

694
695

Tables

Table 1. Overview of statistical results for behavioral repeatability, correlations, and linear
models. Exploration was not quantified in Experiment 3 and therefore no results are presented.
Significance interpreted as: R= 0.2-0.4 moderate repeatability, corr≥0.5 strong correlation, p<
0.05 significant. Pearson correlation (corr) used for normal data, Kendall Rank correlation (tau)
used for non-parametric data

Repeatability			
Behavior	**Experiment 1**	**Experiment 2**	**Experiment 3**
Activity	**R= 0.279, p= 0.032**	**R= 0.123, p= 0.006**	**R= 0.321, p= 0.011**
Exploration	**R= 0.278, p= 0.029**	**R= 0.177, p< 0.001**	NA
Boldness	R= 0.044, p= 0.395	R= 0, p= 1	**R= 0.388, p= 0.003**
Behavioral Syndromes			
Syndrome	**Experiment 1**	**Experiment 2**	**Experiment 3**
Activity:Exploration	**corr= 0.767, p< 0.001**	**corr= 0.911, p< 0.001**	NA
Activity:Boldness	corr= -0.150, p= 0.196	**tau= 0.155, p= 0.002**	**corr= -0.665, p< 0.001**
Exploration:Boldness	corr= -0.119, p= 0.307	**tau= 0.155, p= 0.002**	NA
ANOVAs			
Activity	**Experiment 1**	**Experiment 2**	**Experiment 3**
Trial	NA	$F_{1,234}$**= 170.128, p< 0.001**	$F_{2,143}$**= 6.312, p= 0.002**
Round	$F_{1,37}$= 3.390, p= 0.074	$F_{1,232}$= 0.570, p= 0.451	$F_{2,143}$**=5.971, p= 0.003**
Trial:Round	NA	$F_{2,228}$= 0.216, p= 0.801	$F_{4,,143}$ = 1.356, p= 0.252
Exploration	**Experiment 1**	**Experiment 2**	**Experiment 3**
Trial	NA	$F_{2,230}$**= 112.143, p< 0.001**	NA
Round	$F_{1,37}$= 1.392, p= 0.245	$F_{1,234}$= 0.292, p= 0.590	NA
Trial:Round	NA	$F_{2,228}$= 0.506, p= 0.604	NA
Boldness	**Experiment 1**	**Experiment 2**	**Experiment 3**
Round	$F_{1,74}$= 0.628, p= 0.431	$\chi^2_{2,96}$= 22.429, p= 0.401	$F_{2,36}$ **= 13.100, p< 0.001**

  Figures

Figure 1. Overview of experimental design. All experiments were built on the framework of an
open field test and expanded to explore differences in time and context. For each experiment,
trial type (i.e., different phases within each experiment), trial duration, and number of rounds
(i.e., repeated experiences in the same assay) are shown. OFT = open field experiment, TST =
threat stimulus test, AFET = activity in a familiar environment test, NET = novel environment
test. Colors represent corresponding trial types that were compared across experiments (See
"Cross Analysis" Methods)

715
Figure 2. Overview of analysis methods. For all experiments, we quantified activity as the
number of lines crossed from one area into the next and exploration as the total number of
areas visited (not assessed in Experiment 3). Note that crossing back and forth from one area to
another increases our measure of activity but not exploration. We quantified boldness as time in
center (Experiment 1), response to a looming stimulus (Experiment 2), and latency to cross into
a novel environment (Experiment 3)

723
724 **Figure 3.** Behavior in an artificial environment open field test (Experiment 1). Frogs exhibited
725 repeatability in (a) exploration (R= 0.278, p= 0.029) and (b) activity (R= 0.279, p= 0.032) across
726 two rounds separated by five weeks. Three frogs are highlighted to showcase consistent
727 individual differences in and the relationship between exploration and activity

728
Figure 4. Behavior in a naturalistic open field before, during, and after a threat stimulus
(Experiment 2). Exploration (a) and activity (b) decrease during stimulus presentation compared
to pre-stimulus and post-stimulus ($F_{2,230}= 112.143$, $p< 0.001$; $F_{1,234}= 170.128$, $p< 0.001$), and
these patterns were repeatable across the two rounds of the experiment. Data are collapsed
across rounds, as no significant effect of round or a trial by round interaction were observed for
exploration or activity

Figure 5. Behavior in a familiar, naturalistic environment versus a novel environment
(Experiment 3). (a) Activity increased with repeated experience in the assay (i.e., across rounds;
($F_{2,143}$= 5.971, p= 0.003). (b) All frogs explored the novel area more quickly in subsequent
rounds ($F_{2,36}$ = 13.100, p< 0.001), and (c) more active frogs explored the novel area sooner (corr
= -0.655, p< 0.001). Letters in (a) and (b) indicate significant pair-wise differences between
rounds
742