Title:	Regulation of the mouse ventral tegmental area by melanin-concentrating hormone
Abbreviated title:	MCH-mediated regulation of the VTA
Authors:	C Duncan Spencer, Persephone A Miller, Jesukhogie G Williams-Ikhenoba, Ralitsa G Nikolova, Melissa J Chee
Affiliation:	Department of Neuroscience, Carleton University, Ottawa, ON, K1S 5B6, Canada
Correspondence to:	Melissa Chee
	HS 5309
	1125 Colonel By Drive
	Ottawa, ON, K1S 5B6
	Canada
	melissa.chee@carleton.ca

Number of pages:	45
Number of figures:	6
Number of words	
Abstract:	234
Introduction:	646
Discussion:	1496

Conflicts of Interest: The authors have no conflicts of interest to declare.

Acknowledgements: This study is supported by Natural Sciences and Engineering Research Council of Canada (NSERC) Discovery Grant RGPIN-2017-06272 (MJC), NSERC Canadian Graduate Master's Scholarship and the Ontario QEII-GSST program (CDS), Carleton University's Internship-Carleton University Research Experience for Undergraduate Students (I-CUREUS) (PAM, JWI).

Abstract

Melanin-concentrating hormone (MCH) acts via its sole receptor MCHR1 in rodents and is an important regulator of homeostatic behaviors like feeding, sleep, and mood to impact overall energy balance. The loss of MCH signaling by MCH or MCHR1 deletion produces hyperactive mice with increased energy expenditure, and these effects are consistently associated with a hyperdopaminergic state. We recently showed that MCH suppresses dopamine release in the nucleus accumbens, which principally receives dopaminergic projections from the ventral tegmental area (VTA), but the mechanisms underlying MCH-regulated dopamine release are not clearly defined. MCHR1 expression is widespread and includes dopaminergic VTA cells. However, as the VTA is a neurochemically diverse structure, we assessed *Mchr1* gene expression at glutamatergic, GABAergic, and dopaminergic VTA cells and determined if MCH inhibited the activity of VTA cells and/or their local microcircuit. *Mchr1* expression was robust in major VTA cell types, including most dopaminergic (78%) or glutamatergic cells (52%) and some GABAergic cells (38%). Interestingly, MCH directly inhibited dopaminergic and GABAergic cells but did not regulate the activity of glutamatergic cells. Rather, MCH produced a delayed increase in excitatory input to dopamine cells and a corresponding decrease in GABAergic input to glutamatergic VTA cells. Our findings suggested that MCH may acutely suppress dopamine release while disinhibiting local glutamatergic signaling to restore dopamine levels. This indicated that the VTA is a target of MCH action, which may provide bidirectional regulation of energy balance.

Significance Statement

Role of melanin-concentrating hormone (MCH) on energy balance may converge on the dopamine system via the mesolimbic pathway, as loss of MCH or MCH receptor (MCHR1) signaling increases hyperactivity and energy expenditure associated with a hyperdopaminergic state. MCH can suppress dopamine release within the mesolimbic pathway, but its underlying mechanism is not known. We thus determined if MCH could inhibit dopamine release through direct actions within the ventral tegmental area (VTA). We found that MCH directly inhibited dopaminergic VTA cells, but MCH also disinhibited excitatory input to dopamine cells. Therefore, we showed that the VTA is a putative target site supporting dopamine-dependent actions of MCH.

Introduction

Melanin-concentrating hormone (MCH) is produced largely in the lateral hypothalamic zone to promote positive energy balance by acting through its receptor MCHR1. MCHR1 is a G-protein coupled receptor (Chambers et al., 1999; Lembo et al., 1999; Saito et al., 1999) and typically inhibits cells via $G_{i/o}$ activation (Gao & Pol, 2001). Mice overexpressing *Mch* may overeat and develop diet-induced obesity (Ludwig et al., 2001), while *Mch* knockout mice (*Mch-KO*) are lean, hyperactive, and have an increased metabolic rate (Kokkotou et al., 2005; Shimada et al., 1998). Similarly, *Mchr1* deletion also results in leanness, hyperactivity, and increased energy expenditure (Åstrand et al., 2004; Chen et al., 2002; Marsh et al., 2002). Although MCH administration stimulates feeding (Qu et al., 1996; Sears et al., 2010), transgenic models of *Mch* or *Mchr1* deletion do not similarly effect feeding (Lord et al., 2022). By contrast, hyperactivity and increased energy expenditure are consistent features following the transgenic *Mch* or *Mchr1* deletion.

A hyperdopaminergic state underlies the hyperactivity of *Mch-* or *Mchr1-KO* animals. There is an elevated amount of dopamine in the accumbens nucleus of *Mch-KO* mice (Pissios et al., 2008) and rats (Mul et al., 2011). Furthermore, treating *Mch-KO* mice with a dopamine reuptake inhibitor leads to greater accumulation of dopamine at the accumbens, and this can enhance hyperlocomotor activity (Pissios et al., 2008). Consistently, MCH can suppress dopamine release in the accumbens, and the inhibitory actions of MCH are mediated in part by *Mchr1* expression at GABAergic accumbens neurons (Chee et al., 2019).

Dopaminergic afferents in the accumbens originate from the ventral tegmental area (VTA) (Albanese & Minciacchi, 1983) and form the mesolimbic pathway implicated in the

regulation of reward and motivation. The VTA is a heterogeneous area comprising

dopaminergic, GABAergic, and glutamatergic cells (Morales & Margolis, 2017), as well as cells

that are both GABAergic and glutamatergic (Root, Mejias-Aponte, Qi, et al., 2014; Yamaguchi

et al., 2007), dopaminergic and GABAergic (Stamatakis et al., 2013; Tritsch et al., 2012), or

dopaminergic and glutamatergic (Dal Bo et al., 2004; Stuber et al., 2010; Sulzer et al., 1998;

Tecuapetla et al., 2010). Interestingly, dopaminergic VTA neurons receive inputs from local

GABA and glutamate cells (Morales & Margolis, 2017), but other brain regions, such as the

lateral hypothalamic area, projecting to the VTA may also regulate dopaminergic VTA neurons

(Nieh et al., 2016).

MCH neurons may directly innervate multiple VTA cell types, including GABAergic,

glutamatergic, and dopaminergic cells (Faget et al., 2016). Optogenetic activation of MCH

neurons can increase dopamine release in the accumbens and increase sucrose preference over a

non-nutritive sweetener (Domingos et al., 2013). However, *Mch-KO* mice do not show deficits in

nutrient-sensing because the nutritive value associated with sucrose is ascribed to glutamate

release from MCH neurons (Schneeberger et al., 2018). Nevertheless, this does not preclude

direct MCH actions on VTA neurons that can inhibit dopamine release.

The VTA comprises MCH-immunoreactive fibers (Bittencourt et al., 1992) and *Mchr1*

mRNA (Hervieu et al., 2000), but the identity of MCHR1-expressing VTA cells and

mechanism(s) of MCH action in the VTA have not been defined. We determined *Mchr1* mRNA

expression on GABAergic, glutamatergic, or dopaminergic VTA cells identified by the vesicular

GABA transporter (*Vgat*), vesicular glutamate transporter 2 (*Vglut2*), or tyrosine hydroxylase

(TH), respectively. We found that MCH directly hyperpolarized dopaminergic or GABAergic

VTA cells but not glutamatergic cells. As VTA dopamine cells receive both GABAergic and

glutamatergic afferents (Marisela Morales & Margolis, 2017), we also determined if MCH may

regulate GABAergic or glutamatergic synaptic transmission at dopaminergic VTA cells.

Interestingly, MCH did not alter GABAergic transmission at dopaminergic VTA cells, but it

produced a delayed increase in glutamatergic input. This may reflect a disinhibition of

glutamatergic tone onto dopamine cells in the VTA, as MCH can suppress GABAergic

transmission at glutamate cells. These findings supported both direct and indirect MCH action

that regulates neuronal activity or synaptic transmission at dopaminergic VTA cells.

Materials and Methods

110   All animal procedures were approved by the Animal Care Committee at Carleton

111 University. All mice were bred on a C57BL/6J background and group housed in a 12:12 hour

112 light:dark cycle with *ad libitum* access to water and standard mouse chow (Teklad Global Diets

113 2014, Envigo, Mississauga, Canada).

Quantitative real-time PCR

WT mice 6–20 weeks old (N = 16) were anesthetized with an intraperitoneal injection

(ip) of 7% chloral hydrate (700 mg/kg). The brain was rapidly removed from the skull and

samples of the striatum, hypothalamus, hippocampus, cerebellum, and VTA were microdissected

and flash-frozen in liquid nitrogen. Tissue samples were homogenized in TRIzol (Invitrogen,

Carlsbad, CA), and RNA was isolated by chloroform and precipitated by isopropanol. The RNA

pellet was washed with 70% ethanol and air-dried before resuspending in distilled water. RNA

purity was determined by a 260/280 absorbance ratio of ~2.0 using a NanoDrop Lite

Spectrophotometer (Thermo Scientific, Waltham, MA).

RNA samples were treated with DNase and reverse transcribed for cDNA synthesis with

an iScript gDNA Clear cDNA Synthesis Kit (1725035, Bio-Rad, Hercules, CA). We performed

quantitative RT-PCR using a CFX Connect Real-Time PCR Detection System thermal cycler

(1855201, Bio-Rad). Custom primers (Invitrogen) for mouse glyceraldehyde-3-phosphate

dehydrogenase (*Gapdh*) and *Mchr1* mRNA comprised the following sequences: *Gapdh* forward,

5'- ATGTGTCCGTCGTGGATCTGA-3'; *Gapdh* reverse, 5'-

ATGCCTGCTTCACCACCTTCTT-3'; *Mchr1* forward, 5'- CAATGCCAGCAACATCTCC-3';

Mchr1 reverse, 5'- ACCAAACACTGAAGGCATGA-3'.

Mchr1 mRNA expression in each brain region was calculated via double delta Ct analysis

and standardized to the expression levels of *Gapdh* of the same mouse. Relative *Mchr1* mRNA

expression in each brain region was then normalized to the level of *Mchr1* mRNA in the

hypothalamus for each sex.

Histology

Tissue preparation. Mice (12–20 weeks old) were anaesthetized with 7% chloral

hydrate (700 mg/kg, ip) and underwent transcardiac perfusion with cold (4 °C) sterile saline

(0.9% NaCl) followed by pre-chilled (4°C) phosphate buffered formalin (10%; VWR, Radnor,

PA).

Immunohistochemistry (IHC). Brain tissue from WT (N = 6) and *Mchr1-KO* mice (N

= 2) were post-fixed in 20% sucrose prepared in formalin (4 h) and then cryoprotected with 20%

sucrose prepared in phosphate buffered saline (PBS, pH 7.4) for 16 h until sectioning using a

freezing microtome (Spencer Lens Co., Buffalo, NY) into five series of 30 μm coronal sections.

MCHR1 immunoreactivity was labeled as previously described (Diniz et al., 2020). In brief,

brain tissue was treated with 0.3% hydrogen peroxide and incubated with a rabbit anti-MCHR1

antibody (1:1,000; PA5-24182, UC2738292, Invitrogen, Carlsbad, CA) for 48 h at 4 ℃ in a 3%

normal donkey serum (NDS) solution. Tissue was thoroughly rinsed, incubated with a

biotinylated donkey anti-rabbit IgG antibody (1:500; 711-065-152, Lot 101909, Jackson

ImmunoResearch Laboratories) at RT for 1 h, and treated with avidin-biotin-horseradish

peroxidase (1:1:833; PK-6100, Vector Labratories, Burlingame, CA) at RT for 30 min before

incubating with 0.5% biotinylated (EZ-Link sulfo-NHS-LC-biotin; ThermoFisher Scientific,

Waltham, MA) tyramine (Sigma-Aldrich; Sigma Chemical; St. Louis, MO, USA) for 20 min

(RT). All sections were then incubated with an Alexa Fluor 647-conjugated streptavidin (1:500;

016-600-084, Lot 135095, Jackson ImmunoResearch Laboratories) and NeuroTrace (1:50;

N21479, ThermoFisher) for 2 h at RT. Following staining, all tissue was thoroughly rinsed and

immediately mounted onto Superfrost Plus microscope glass slides (ThermoFisher) and

coverslipped with Prolong Diamond Antifade mounting media (P36961, Invitrogen).

After IHC processing for MCHR1 immunoreactivity, one additional series were

incubated with a sheep anti-TH antibody (1:1,000; GTX82570, Lot 822005648, GeneTex, Irvine,

CA) at RT overnight. Sections were then incubated with a donkey anti-sheep secondary antibody

conjugated with Alexa Fluor 594 (1:500; 715-496-150, Lot 82079, JacksonImmuno Research

Laboratories) and NeuroTrace (1:50) for 2 h at RT. Tissues were rinsed, immediately mounted,

and coverslipped as described above.

Thick tissue IHC. Brain slices following patch-clamp recordings were fixed in formalin

for at least 24 h, treated with a 2% Triton-X PBS solution for 45 min at RT to permeabilize the

tissue, and then incubated with a sheep anti-TH primary antibody (1:1,000; GeneTex) at 4 °C

for 72 h. After rinsing, the tissue was incubated with a streptavidin-conjugated Cy3 (1:500; 016-

160-084, Lot 146014, JacksonImmuno Research Laboratories) and donkey anti-sheep conjugated

with Cy5 (1:500; 713-176-147, Lot 70443, JacksonImmuno Research Laboratories) secondary

antibodies. The tissues were then rinsed and immediately mounted onto Superfrost Plus

microscope glass slides and coverslipped with Prolong Diamond Antifade mounting media).

Combined *in situ* hybridization (ISH) and IHC. Brains from WT mice (N = 6) were

post-fixed in 10% formalin overnight, cryoprotected in 20% sucrose with 0.5% sodium-azide (24

h) and sectioned using a freezing microtome into five series of 30 μm coronal sections. The

series designated for ISH were immediately mounted onto Superfrost Plus microscope slides, air-

dried at room temperature (RT; 21–23 °C; 1 h) and −20 °C (30 min), and then stored at −80 °C.

Adjacent brain series were reserved for Nissl staining, RNAscope probe labeling, and negative

control, respectively. One slide from a separate series served as the positive control. We

determined the coexpression of *Mchr1* with *Slc32a1* (*Vgat*) and *Slc17a6* (*Vglut2*) by RNAscope-

enabled ISH and with TH by subsequent IHC in the same brain tissue. ISH procedures were

completed based on manufacturer instructions for RNAscope ISH on fixed-frozen mouse brain

tissue (Multiplex Fluorescent Reagent Kit v2 User Manual; Advanced Cell Diagnostics (ACD),

Newark, CA) as previously described (Bono et al., 2022), unless indicated otherwise.

In brief, slide-mounted brain tissues were dehydrated in an ethanol gradient, treated with

hydrogen peroxide and target retrieval procedure for 5 min (99 °C), and then dehydrated and air-

dried overnight (RT). Tissues of interest were isolated by drawing a hydrophobic barrier and

treated with Protease III in a HybEZ oven (40 °C, 30 min). Slides were then incubated with a

cocktail of *Mm-Mchr1* (317491, ACD) in channel 1 (C1), *Mm-Slc32a1-C2* in C2 (319191-C2,

ACD), and *Mm-Slc17a6* in C3 (319171-C3, ACD) probes (40 °C, 2 h). Concentrated *Mm-*

Slc32a1 C2 and *Mm-Slc17a6* C3 probes were added to the diluted *Mm-Mchr1* C1 probe at a ratio

of 1:50. A volume of 5–8 drops of the probe mixture was added to each slide. *Bacilus*

dihydrodipicolinate reductase (*dapB*, 320871, ACD) and *Mm-Ppib* (313911, ACD) probes were

used as the negative and positive control, respectively. Hybridization signals were amplified by

sequential incubation at 40 °C with AMP-1 (30 min; 323101, ACD), AMP-2 (30 min; 323102,

ACD), and AMP-3 (15 min; 323103, ACD).

C1 *Mchr1* hybridization was developed (40 °C, 15 min with HRP-C1; 323104, ACD) and

tyramide signal amplification (TSA) plus Opal 690 (1:750, 40 °C, 30 min; FP1497001KT,

Akoya Biosciences, Marlborough, MA), then quenched with HRP Blocker (40 °C, 15 min;

323107, ACD). C2 *Vgat* and C3 *Vglut2* hybridization were sequentially developed with HRP-C2

(323105, ACD) with TSA plus Cyanine 3 (1:1,500; NEL44E001KT, PerkinElmer, Waltham,

MA) and HRP-C3 (323106, ACD) with TSA plus Opal 520 (1:750; FP1487001KT, Akoya

Biosciences). C2 signal was similarly quenched with HRP Blocker (40 °C, 15 min) before

proceeding to C3 development.

Following ISH of probe tissue, the tissues were treated for IHC labeling of TH

immunoreactivity based on ACD directions for combined ISH and IHC (Advanced Cell

Diagnostics, 2020) and as previously described (Mickelsen et al., 2019). Tissues were blocked

with 3% NDS solution prepared in PBS (700 μl) for 30 min (RT) and then incubated with anti-

sheep TH primary antibody (1:1,000; GTX82570, Lot 822005648, GeneTex, Irvine, CA) 2 h

(RT). Slides were thoroughly rinsed in PBS and then incubated with a donkey anti-sheep

secondary antibody conjugated with Alexa Fluor 594 (1:500; 715-496-150, Lot 82079,

JacksonImmuno Research) for 30 min (RT). Slides were rinsed in Wash Buffer (320058, ACD),

treated with DAPI Reagent (323108, ACD) for 30 s (RT), and immediately coverslipped with

ProLong Gold Antifade Mountant (Fisher Scientific). Slides were kept in a dark drying drawer at

RT overnight and then transferred to –20 °C for storage until imaging.

Nissl stain. Sections underwent Nissl staining as previously described (Bono et al., 2022;

Negishi et al., 2020).

Microscopy

Brightfield imaging. Nissl-stained tissue was acquired with a Nikon Eclipse Ti2 inverted

microscope (Nilon Instruments Inc., Mississauga, Canada) equipped with a motorized stage and

DS-Ri2 color camera (Nikon) using a Plan Apochromat 10x objective (0.45 NA). Brightfield

images were stitched at 20% overlap with NIS Elements software.

Confocal imaging. Images from Nissl-stained tissue were aligned to tiled, overview

confocal images showing the VTA in relation to other midbrain regions acquired using the 405-

nm wavelength laser with a 4x Plan Apochromat objective (0.2 NA). Stitched, high resolution

confocal frames from IHC- and/or ISH-treated tissues were captured from within the overview

image.

Confocal imaging of MCHR1 immunoreactivity underwent scans with a 10x and/or 20x

(0.75 NA) Plan Apochromat objectives using a Nikon C2 confocal system. Images were acquired

with a 405- and 647-nm wavelength laser to visualize NeuroTrace- and Alexa Fluor 647-labeled

signals. Tissue that was also stained for TH had additional images acquired with a 561-nm

wavelength laser to visualize Alexa Fluor 594-labeled signal, and fluorescence signals from the

561- and 647-nm channels were pseudocoloured magenta and green, respectively.

Confocal imaging of combined ISH- and IHC-treated tissue underwent scans with 20x

Plan Apochromat objective using a Nikon C2 confocal system and acquired with a 405-, 488-,

and 647-nm wavelength laser to visualize DAPI-, Opal 520-, and Alexa Fluor 647-labeled

signals for nuclear stains, *Vglut2* hybridization, and *Mchr1* hybridization, respectively.

Simultaneously at the same optical section, Cyanine 3- and Alexa Fluor 594-labeled signals to

visualize *Vgat* hybridization and TH were acquired by spectral imaging using a 561-nm

wavelength excitation laser in the continuous band (CB) setting to generate a lambda stack

acquired with sequential 10-nm bandwidths spanning the wavelength range of 580 nm to 640

nm. The spectral profiles were unmixed to separate Cyanine 3 and Alexa Fluor 594 signals,

which were most distinguished at 590 nm and 620 nm, respectively, and then merged with

signals acquired by the 405-, 488-, and 647-nm wavelength laser to generate one image file

displaying signals from all five fluorophores.

Tissues that served as the negative control were also acquired using the same

combination of conventional confocal and spectral unmixing for probe tissue. Background

fluorescence related to nonspecific binding from ISH treatment was determined using the

negative control tissue by adjusting the look up table (LUT) values for each of the three image

channels to yield a resultant black image. The LUT range was then applied to images acquired

from the probe tissue to subtract background fluorescence, thus any remaining signal observed

from the probe tissue was considered to be positive hybridization signals.

Confocal imaging of IHC-treated cells in thick tissue underwent scans with a 20x Plan

Apochromat objective in a Nikon C2 confocal system and acquired with a 488-, 561-, and 647-

nm wavelength laser to visualize native EGFP fluorescence, Cy3-labelled signal for biocytin-

filled cell, and Cy5-labelled signals for TH immunoreactivity, respectively. Cy3 signal was

pseudocoloured blue.

Neuroanatomical analyses

Plane-of-section analysis. Nissl or fluorescent-Nissl images were parcellated in Adobe

Illustrator CS6 (Adobe Inc., San Jose, CA) based on the regions defined by the *Allen Reference*

Atlas (*ARA*; Dong, 2008), as previously described (Bono et al., 2022; Negishi et al., 2020).

Parcellations for MCHR1-ir cells were directly applied from fluorescent-Nissl signals captured

within the same image acquisition. High magnification confocal images of ISH-treated tissues

were indirectly parcellated from brightfield images of an adjacent Nissl-stained section, which

was aligned to the low magnification, overview confocal image of the probe tissue.

Identifying MCHR1- and TH-ir signals. Our MCHR1 antibody (1:1,000; PA5-24182,

Invitrogen, Carlsbad, CA) labeled MCHR1 immunoreactivity at the primary cilium of neurons

(Diniz et al., 2020). Only cilia, identified by their long spindle-like appearance greater than 1 μm

(Bansal et al., 2021; Diniz et al., 2020), associated with a NeuroTrace-labeled somata was

counted as a MCHR1-expressing cell. Only TH staining at the somata that colocalized with

DAPI-, NeuroTrace-, or EGFP- labeled cell was considered. Only signals within the boundaries

of the VTA, as determined by our plane-of-section analysis, were included in our analyses.

Identifying hybridization signals. Signal criteria was independently-established based

on the signal-to-background level for each hybridization signal. *Mchr1* hybridization signal

appeared as punctate "dots" with clusters centered tightly around a DAPI-labelled nucleus (Bono

et al., 2022). *Vgat* hybridization appeared as dot clusters that took the shape of cell bodies and

were colocalized to a DAPI-labelled nucleus. Compared to *Vgat* hybridization, dot clusters

following *Vglut2* hybridization was less dense within the cell and may appear as punctate dots.

Mapping. A circle was placed over each positively-labeled immunoreactive or

hybridized cell, and the relative positions of the circles were transferred, as previously described

(Bono et al., 2022; Negishi et al., 2020), to coronal brain atlas templates (Dong, 2008) to

generate standardized maps of *Mchr1*, *Vglut2*, *Vgat*, and TH expression. Cell counts at each atlas

level were determined from cells mapped and counted unilaterally.

Number of cells reported for IHC and combined ISH and IHC studies were corrected for

oversampling, as previously described (Chee et al., 2013; Negishi et al., 2020; Bono et al., 2022),

using the Abercrombie formula $P = A\left(\frac{M}{L+M}\right)$ (Abercrombie, 1946): where P is the corrected

count, A is the original count, M is the mean tissue thickness (IHC, 13.6 μm; ISH+IHC, 10.9

μm) determined from five brain slices, and L is the mean cell diameter (IHC, 15.6 μm;

ISH+IHC, 16.0 μm) determined from 80 cells.

Electrophysiology

Animals. *Th-cre;L10-Egfp*, *Vgat-cre ;L10-Egfp*, and *Vglut2-cre;L10-Egfp* reporter mice

were generated by crossing *Th-cre* (Lindeberg et al., 2004), *Vgat-cre* (028862, JAX; Vong et al.,

2011), or *Vglut2-cre* (028863, JAX; Vong et al., 2011) mice, respectively to homozygous *lox-*

STOP-lox-L10-Egfp reporter mice (Krashes et al., 2014). All transgenic lines were kindly

provided by Dr. B. B. Lowell (Beth Israel Deaconess Medical Center, Boston, MA).

Acute slice preparation. Mice (4–10 weeks old) were anaesthetized with 7% chloral

hydrate (700 mg/kg, ip) and transcardially perfused with ice cold, carbogenated (95% O_2, 5%

CO_2) slice artificial cerebrospinal fluid (ACSF) solution (300 mOsm/L; pH 7.4) containing (in

mM) 92 NMDG, 2.5 KCl, 1.25 NaH_2PO_4, 30 $NaHCO_3$, 20 HEPES, 25 glucose, 2 thiourea, 5 Na-

ascorbate, 3 Na-pyruvate, 0.5 $CaCl_2 \cdot 2H_2O$, and 10 $MgSO_4$ (Ting et al., 2018). Brains were

sectioned on a vibratome (VT 1000S, Leica Biosystems, Wetzlar, Germany) in slice ACSF to

produce 250 μm-thick coronal slices that were immediately transferred to carbogenated slice

solution at 37 °C for 5 min. Slices were then incubated in a carbogenated bath ACSF (300

mOsm/L) containing (in mM) 124 NaCl, 3 KCl, 1.3 $MgSO_4$, 1.4 NaH_2PO_4, 10 D-glucose, 26

$NaHCO_3$, and 2.5 $CaCl_2$ for 5 min and then allowed to recover in bath ACSF (RT) for at least 1

hr prior to recordings.

Patch-clamp recordings. Brain slices containing the VTA, which primarily

encompassed L83–87, were placed in a slice chamber, superfused (2–2.5 ml/min) with

carbogenated bath ACSF pre-warmed (37 °C) by a temperature controller (Warner Instrument

Corporation, Hamden, CT), and held in place by a platinum ring strung with nylon fiber made in-

house.

VTA cells were visualized with infrared differential interference contrast microscopy at

40x magnification on either an Examiner.A1 microscope (Zeiss, Oberkochen, Germany)

equipped with an AxioCam camera (Zeiss) and Axiovision software (Zeiss) or with an Eclipse

FN1 microscope (Nikon) equipped with a pco.panda 4.2 camera (Excelitas PCO GmbH,

Kelheim, Germany) and NIS-Elements Imaging software (Nikon). EGFP-labeled cells were

identified using epifluorescence illumination from a halogen (HXP120, Zeiss) or mercury lamp

(C-SHG1, Nikon).

Borosilicate glass pipettes (593800; A-M Systems) were pulled on a P-1000

Flaming/Brown micropipette puller (Sutter Instruments; Novato, CA) and backfilled (7–10 MΩ)

with a potassium-based internal solution (290 mOsm/L, pH 7.24) containing (in mM) 120 K-

gluconate, 10 KCl, 10 HEPES, 1 $MgCl_2$, 1 EGTA, 4 MgATP, 0.5 NaGTP, and 10

phosphocreatine to assess membrane properties and glutamatergic events. A cesium-based

internal solution (290 mOsm/L, pH 7.24) containing (in mM) 128 CsMS, 11 KCl, 10 HEPES,

0.1 CaCl2, 1 EGTA, 4 MgATP, 0.5 NaGTP) was used to record GABAergic events. Voltage-

clamp recordings were acquired at a holding potential of $V_h = -60$ mV and -5 mV for excitatory

and inhibitory events, respectively using a MultiClamp 700B amplifier (Molecular Devices, San

Jose, CA) and digitized using a Digidata 1440A (Molecular Devices). Biocytin (0.4%; Cayman

Chemical, Ann Arbor, MI) was added to the internal solution for post-hoc thick tissue IHC in a

subset of recorded cells. Traces were acquired at 10 kHz and filtered at 3 kHz by pClamp 10.3 or

11 software (Molecular Devices), as previously described (M. Chee et al., 2010).

Drug application. Pharmacological applications of MCH (H-1482, Bachem, Torrance,

CA) was applied by bath perfusion or puff application after a baseline period of at least 5 min or

1 min, respectively, where the membrane potential fluctuated by less than 10%. Cells were held

for at least 20 min during the MCH-washout period to assess the reversibility of MCH-mediated

effects on the membrane potential. Where applicable, tetrodotoxin citrate (TTX; 500 nM; T-550,

Alomone labs, Jerusalem, Israel) or the MCHR1-antagonist TC-MCH 7c (10 μM; 4365, Tocris,

Toronto, Ontario, Canada), was bath applied and allowed to saturate the slice for at least 10 min

prior to MCH application and then maintained during the MCH application and washout period

as well.

Puff-applied MCH was delivered via a puff-pipette backfilled with 3 μM MCH (or bath ACSF)

positioned within 20–40 μm away from the patched cell held in the whole-cell configuration. A

short "puff" was applied by manually applying positive pressure via the puff-pipette, and a

solution ripple can be seen to contact the patched cell.

Experimental design and statistical analyses

qPCR studies. Comparison of *Mchr1* gene expression in wildtype mice (N = 6–7/sex)

were analyzed using a between-subject design. *Mchr1* expression in different regions were

analyzed using a two-way mixed model ANOVA with Šidák post-hoc testing to determine sex

differences.

Neuroanatomical studies. Comparisons of the total number of MCHR1-expressing cells

were analyzed using a between-subject design from (N = 3/sex) and compared via unpaired *t* test,

and differences in the distribution of MCHR1-expressing cells across each *ARA* level between

sexes was determined by the two-way mixed model ANOVA.

Comparisons of *Mchr1-*, *Vglut2-*, *Vgat-*, or TH-expressing cells were analyzed from male

and female wildtype mice (N = 6) using a between-subject design, and a one-way ANOVA was

used to identify distinct VTA cell groups. Sex differences were determined by two-way

ANOVA.

Electrophysiological studies. Acute brain slices were prepared from male and female

Th-cre;L10-Egfp (N = 27), *Vgat-cre;L10-Egfp* (N = 8), and *Vglut2-cre;L10-Egfp* (N = 8) reporter

mice. Up to three cells were recorded from three to four coronal slices, where the VTA sits

between the medial mammillary nucleus and compact part of the substantia nigra in rostral

sections and sits beside the interpeduncular nucleus in caudal sections. Recordings from EGFP-

labeled VTA cells of each reporter line was guided by atlas maps of *Mchr1* expression in each

cell type. Electrical signals recorded were processed in Clampfit 10.7 (Molecular Devices).

The resting membrane potential (RMP) was sampled every 1 s and averaged into 30-s

bins in bath-applied treatments and sampled every 100 ms and averaged into 1-s bins in puff-

applied treatments. Cells deemed to be MCH-responsive exhibited a change in RMP lasting at

least 2 min that was greater than 2-times the standard deviation of the mean RMP at baseline.

MCH-mediated change in RMP (Δ RMP) was sampled within 7 min after MCH application or

within 5 s of puff application. Comparisons of mean RMP before and after MCH application was

determined via a paired *t* test and reported with effect size (η^2). Comparisons of MCH-mediated

Δ RMP over time in the presence or absence of TTX was determined using within-subject design

and repeated measures two-way ANOVA with Šidák post-hoc testing.

Excitatory (EPSC) and inhibitory postsynaptic current (IPSC) events were detected using

MiniAnalysis (Synaptosoft, Decatur, GA). Event detection threshold was set to 5-times the root

mean square noise level for each recording. The frequency of EPSCs and IPSCs were indirectly

determined by the interevent interval between events using MiniAnalysis (Synaptosoft, Decatur,

GA) and averaged into 30-s bins. The baseline event frequency or amplitude was determined as

the mean event frequency averaged over 5 min immediately prior to MCH bath application.

Comparisons of mean event frequency or amplitude before and after MCH application was

determined via a paired *t* test and reported with effect size (η^2). Comparisons of MCH-mediated

effects on postsynaptic current events over time in the presence or absence of TTX or TC-MCH

7c was determined using within-subject design and repeated measures two-way ANOVA with

Šidák post-hoc testing.

Graphs. Only responders were included in our graphical representations, unless a null

response was referenced. All data graphs were prepared using Prism 9 (GraphPad, San Diego,

CA). Results were considered statistically significant at $p < 0.05$. All data are reported as group

mean ± SEM. Figures were assembled in Adobe Illustrator 2020 (Adobe Inc.). Numeral in

parentheses in graphs represent the number of animals (N) or cells (n).

Results

Moderate *Mchr1* mRNA and MCHR1 protein expression in the mouse VTA

Mchr1 mRNA and protein have been reported in the rat VTA (Hervieu et al., 2000; Saito

et al., 2001), and we first determined *Mchr1* mRNA and/or protein expression in the VTA of

male and female wildtype mice. While the level of *Mchr1* mRNA varied across brain regions

$(F(4, 44) = 81.6, p < 0.0001$, ANOVA), it was similar between male and female mice $(F(1, 11) =$

$0.6, p = 0.449$, ANOVA), and there was no interaction between brain region and sex $(F(4, 44) =$

$0.8, p = 0.561$, ANOVA). As expected, *Mchr1* expression was highest in the striatum, which

included the nucleus accumbens and caudate putamen, and barely detectable in the cerebellum

(**Figure 1A**). Importantly, *Mchr1* expression in the VTA was comparable to the hypothalamus

(male: $t(44) = 1.5, p = 1.000$; female: $t(44) = 1.0, p = 1.000$) and hippocampus (male: $t(44) = 1.3,$

$p = 1.000$; female: $t(44) = 0.6, p = 1.000$; **Figure 1A**).

The VTA of wildtype male and female mice also comprised MCHR1-immunoreactive

(MCHR1-ir) signals, which appeared long and spindle-like and were consistent with MCHR1

expression observed on the primary cilium of neurons (**Figure 1B**). No MCHR1-ir labeling was

detected in the VTA of *Mchr1-KO* mice (**Figure 1C**). Male and female wildtype mice had a

similar distribution pattern of MCHR1-expressing cells throughout the VTA ($F_{(1, 4)} = 0.02$, p =

0.891, ANOVA; **Figure 1D**). Additionally, there was no difference in the total number of

MCHR1-expressing cells in the VTA of male (159 ± 48, n = 3) and female (171 ± 29, n = 3)

mice ($t_{(4)} = 0.2$, p = 0.835, $\eta^2 = 0.01$, unpaired *t* test).

Neurochemical diversity of VTA cells

The VTA comprises multiple cell types, including dopaminergic, GABAergic, or

glutamatergic cells that have distinct distribution patterns (M. Morales & Root, 2014; Marisela

Morales & Margolis, 2017). To identify the neurochemical identities of VTA cell types, we

determined if *Mchr1* hybridization colocalized with *Vglut2* or *Vgat* hybridization in

glutamatergic or GABAergic cells, respectively, or with TH immunoreactivity in dopaminergic

cells within the same tissue. As there were no sex differences in *Mchr1* mRNA (**Figure 1A**) or

protein expression (**Figure 1D**), we herein combined our neuroanatomical analyses from male

and female mice.

The overall number of VTA cells labeled by *Vglut2* mRNA, *Vgat* mRNA, and/or TH

immunoreactivity decreased posteriorly (**Figure 2A**, gray line). We compared the distribution of

Vglut2 mRNA, *Vgat* mRNA, and TH-ir cells within the VTA to determine their relative

prevalence rostrocaudally, and we distinguished at least five subtypes of VTA cells ($F_{(4, 50)} =$

13.4, p = 0.0001, ANOVA). *Vglut2* cells were most prominent anteriorly then gradually

diminished towards the posterior VTA (**Figure 2A**), and they were usually medially distributed

within the VTA (**Figure 2B**). *Vgat* cells comprised ~30% of VTA cells counted through the

anterior and middle sections of the VTA (L80–87) but made up the majority of cells in the

posterior VTA (L88–90) (**Figure 2A**). TH-ir cells were most numerous in the middle of the VTA

and comprised ~40% of cells counted (**Figure 2A**). Both *Vgat* and TH cells tended to be laterally

distributed, though some *Vgat* cells were spread medially (**Figure 2B**).

In addition, we also found that VTA cells may express two or more neurotransmitter

markers. The most common were those expressing both *Vglut2* and *Vgat* that comprised 5–10%

of VTA cells and up to 16% of cells towards the middle of the VTA (L86–87; **Figure 2A**). Cells

expressing TH and *Vglut2* emerged towards the posterior VTA (L85–89) but comprised <6% at

its peak (**Figure 2A**) so were not mapped. Similarly, just 0.25% of cells exhibited TH and *Vgat*,

and cells exhibiting *Vglut2*, *Vgat*, and TH were even rarer (<0.05%; data not shown).

Neurochemical and spatial distribution of *Mchr1* mRNA expression in the VTA

Mchr1 mRNA expression was widely distributed throughout the VTA (**Figure 3A**) and

across all major VTA cell types (**Figure 3B–E**), but there was no sex difference in the

heterogeneity of *Mchr1* VTA cells (F(1, 24) = 0.01, p = 0.924, ANOVA; **Figure 3-1A**).

Interestingly, about one in four (25%) *Mchr1* cells (*Mchr1*-only) do not coexpress TH, *Vglut2*, or

Vgat and are yet to be defined neurochemically (**Figure 3F**). These *Mchr1*-only cells can be seen

throughout the VTA (**Figure 3G**) and tend to be less prominent in the VTA of female mice, as

sex may interact with *Vgat* expression in *Mchr1* cells (F(1, 8) = 4.8, p = 0.059, ANOVA; **Figure

3-1B**). Consistent with the rostrocaudal composition of VTA cells, *Mchr1* expression was

predominantly in *Vglut2* cells rostrally, TH cells in the middle sections of the VTA, and in *Vgat*

cells caudally (**Figure 3G**).

About half (52%) of *Vglut2* VTA cells expressed *Mchr1* mRNA (**Figure 4-1**), which was

evenly distributed throughout *Vglut2* cells in the medial and anterior VTA but only sparsely

distributed posteriorly (**Figure 4A**). By contrast, few *Vgat* cells in the anterior VTA expressed

Mchr1 mRNA (**Figure 4-2**), but in the middle and posterior VTA sections where *Mchr1*-

expressing *Vgat* cells were more common, they tend to be laterally distributed or have no

discernible pattern, respectively (**Figure 4B**). Interestingly, the majority of TH-ir VTA neurons

(78%) expressed *Mchr1* mRNA (**Figure 4-3**), and while *Mchr1*-expressing TH cells were found

sparingly in the anterior VTA, they were abundant in middle VTA sections, especially near the

ventral and lateral borders of the VTA (**Figure 4C**). In addition, although dual *Vglut2* and *Vgat*

cells comprised just one-tenth of all VTA cells, nearly 60% of them expressed *Mchr1* mRNA

(**Figure 4-4**) and were most likely found in the dorsomedial aspects of the VTA (**Figure 4D**).

Taken together, these findings indicated the abundance of *Mchr1* mRNA throughout the VTA in

glutamatergic, GABAergic, and/or dopaminergic VTA cells.

MCH inhibited *Vgat* and *Th* VTA cells but not *Vglut2* VTA cells

Based on the spatial distribution of *Mchr1* expression in *Vglut2*, *Vgat*, and TH VTA cells

(**Figure 4**), we performed whole-cell patch-clamp recordings in EGFP-labeled *Vglut2*$^{\text{EGFP}}$,

Vgat$^{\text{EGFP}}$, and *Th*$^{\text{EGFP}}$ VTA cells from *Vglut2-cre;L10-Egfp* (**Figure 5A*i***), *Vgat-cre;L10-Egfp*

(**Figure 5B*i***), and *Th-cre;L10-Egfp* reporter mice (**Figure 5C*i***), respectively. MCH (3 μM) bath

application had no effect on the resting membrane potential (baseline: -59.6 ± 3.2 mV;

MCH: -59.7 ± 3.8 mV, n = 6 cells, t(5) = 0.20, p = 0.849, $\eta^2 = 0.01$, paired t test) of *Vglut2*$^{\text{EGFP}}$

VTA cells (n = 0/6 cells, N = 3; **Figure 5A*ii***). By contrast, MCH hyperpolarized *Vgat*$^{\text{EGFP}}$

(baseline: -47.5 ± 2.3 mV; MCH: -52.9 ± 2.7 mV, n = 5/14 cells, N = 5, t(4) = 2.80, p = 0.049,

$\eta^2 = 0.66$, paired t test; **Figure 5B*ii***) and *Th*$^{\text{EGFP}}$ cells (baseline: -53.5 ± 3.6 mV; MCH:

-57.8 ± 2.8 mV, n = 6/16 cells, N = 12, t(5) = 3.15, p = 0.025, $\eta^2 = 0.66$, paired t test; **Figure**

5C*ii*) within 10 min and persisted for at least 25 min before gradually returning to baseline.

While the proportion of MCH-sensitive $Vgat^{EGFP}$ VTA cells could be predicted by the

availability of *Mchr1* expression at *Vgat* cells (**Figure 4-2**), the proportion of MCH-sensitive

Th^{EGFP} cells was lower than predicted by the prevalence of *Mchr1* expression at TH-ir cells

(**Figure 4-3**). We first confirmed that over 80% (n = 13/16 cells, N = 7) of Th^{EGFP} cells in patch-

clamp recordings expressed TH immunoreactivity (**Figure 6A**). We then determined the

proportion of TH-ir cells that expressed MCHR1 protein (**Figure 6B**). Interestingly, only about

one-third of TH-ir cells within the span of the VTA targeted for patch-clamp recordings (L84–

87) expressed MCHR1 (**Figure 6C**), which is comparable with a lower (38%) proportion of

MCH-sensitive Th^{EGFP} cells surveyed by patch-clamp recordings (**Figure 5C*ii***).

We then assessed if MCH-mediated inhibition of $Vgat^{EGFP}$ and Th^{EGFP} VTA cells was

activity-dependent by pretreatment with the voltage-gated sodium channel blocker tetrodotoxin

(TTX). Since the effect of bath-applied MCH was so long-lasting, we first established if MCH-

sensitive VTA cells could be identified by a short puff application of MCH (**Figure 7A*i***) lasting

<5 seconds to allow rapid recovery of the resting membrane potential (**Figure 7A*ii***). A single

puff of MCH produced an acute membrane hyperpolarization (baseline: −55.5 ± 4.2 mV; Δ

RMP: −6.7 ± 1.8 mV, n = 4) that peaked within 10 seconds and returned to baseline within one

minute. A second puff of MCH applied 3–5 minutes later at the same cell elicited a similar

hyperpolarization (baseline: −55.9 ± 4.3 mV; Δ RMP: −4.3 ± 1.7 mV, n = 4) and time course

($F_{(1, 6)}$ = 3.3, p = 0.120, ANOVA; **Figure 7A*iii***). Importantly, puff application of ACSF

produced visibly different outcomes than the first MCH puff ($F_{(1, 5)}$ = 6.8, p = 0.047, ANOVA)

or second MCH puff ($F_{(1, 5)}$ = 30.7, p = 0.003, ANOVA), as it did not effect changes in

membrane potential (baseline: −56.3 ± 2.7 mV; Δ RMP: −0.2 ± 0.6 mV, n = 3; **Figure 7A*iii***).

This thus showed that a localized MCH puff could identify MCH-sensitive VTA cells by

enabling short-lived but reproducible membrane hyperpolarizations that were not significantly

desensitized by repeated application. We then puff-applied MCH to identify MCH-sensitive

$Vgat^{EGFP}$ and Th^{EGFP} VTA cells and compared the effect of MCH before and after TTX (500 nM)

treatment. MCH-mediated hyperpolarization at $Vgat^{EGFP}$ VTA cells (baseline: -56.3 ± 1.7 mV; Δ

RMP: -1.8 ± 1.2 mV, n = 4) persisted in the presence of TTX (baseline: -55.2 ± 1.3 mV; Δ

RMP: -1.1 ± 0.4 mV, n = 4) over time ($F_{(1, 6)} = 0.002$, p = 0.965, ANOVA; **Figure 7B**).

Similarly, Th^{EGFP} VTA cells remained sensitive to the peak effect of MCH (baseline: $-57.1 \pm$

3.6 mV; Δ RMP: -5.4 ± 1.6 mV, n = 5) in the presence of TTX (baseline: -50.1 ± 3.4 mV; Δ

RMP: -4.3 ± 1.0 mV, n = 5) and over time ($F_{(1, 8)} = 0.3$, p = 0.623, ANOVA; **Figure 7C**).

These findings indicated that MCH directly hyperpolarized $Vgat^{EGFP}$ and Th^{EGFP} VTA cells.

MCH increased glutamatergic input to *Th* VTA cells

Dopaminergic output from the VTA can be regulated by local glutamatergic and

GABAergic circuitry (Bariselli et al., 2016; Morales & Margolis, 2017; Wang et al., 2015), thus

we assessed if MCH regulated the GABAergic and/or glutamatergic input to Th^{EGFP} VTA cells.

Bath application of MCH did not alter the frequency (baseline: 0.9 ± 0.1 Hz; MCH: 0.9 ± 0.1 Hz,

n = 9, $t_{(8)} = 0.33$, p = 0.750, $\eta^2 = 0.01$, paired t test; **Figure 8A**, **B**) or amplitude (baseline: 13.1

± 0.8 pA; MCH: 13.4 ± 0.8 pA, n = 9, $t_{(8)} = 0.60$, p = 0.563, $\eta^2 = 0.04$, paired t test; **Figure 8C**)

of sIPSC events arriving at Th^{EGFP} VTA cells.

Interestingly, MCH elicited a delayed 3.1 ± 0.9-fold increase in baseline sEPSC

frequency (1.3 ± 0.4 Hz, n = 8) at Th^{EGFP} VTA cells that gradually peaked about 25 minutes

following MCH application (**Figure 8D**, **8E*i***). To determine if this delayed sEPSC frequency

increase was MCH-dependent, we tracked the change in sEPSC frequency in the absence of

MCH (ACSF only) or in the presence of a MCHR1 antagonist, TC-MCH 7c. Following MCH

treatment, sEPSC frequency increased, from 1.3 ± 0.4 Hz to 2.6 ± 0.6 (n = 8, t(7) = 3.17, p =

0.016, $\eta^2 = 0.59$, paired t test), which was not seen when bathed in ACSF only (baseline:

1.0 ± 0.4 Hz; MCH: 0.9 ± 0.5 Hz, n = 4, t(3) = 0.34, p = 0.760, $\eta^2 = 0.03$, paired t test). The

MCH-mediated increase in sEPSC frequency was also not seen when MCH was applied in the

presence of TC-MCH 7c (baseline: 0.9 ± 0.4 Hz; MCH in TC-MCH 7c: 0.8 ± 0.3 Hz, n = 7, t(6)

= 0.29, p = 0.780, $\eta^2 = 0.01$, paired t test). There was a main effect of drug over time (F(2, 16) =

4.6, p = 0.027, ANOVA; **Figure 8E*i***), as the time course of fold change in sEPSC frequency

with MCH treatment was significantly different than in ACSF (t(16) = 2.6, p = 0.036, Sidak

post-test; **Figure 8E*i***) or when MCH was applied in the presence of TC-MCH 7c (t(15) = 2.4, p

= 0.058, Sidak post-test; **Figure 8E*i***). This difference was largely attributed to a delayed increase

in sEPSC frequency that was prominent >20 min after drug treatment. Overall, the fold change in

sEPSC frequency in MCH (2.8 ± 0.6, n = 8) was significantly greater than in ACSF only

(0.8 ± 0.2, n = 4; t(16) = 2.93, p = 0.029, ANOVA) or when applied at Th^{EGFP} VTA cells

pretreated with TC-MCH 7c (0.8 ± 0.1, n = 7; t(16) = 3.33, p = 0.013, ANOVA; **Figure 8E*ii***).

There was no main effect of MCH on sEPSC amplitudes over time (F(2, 16) = 2.3, p = 0.130;

ANOVA; **Figure 8F*i***) or when comparing their fold change within each condition (F(2, 16) =

0.18, p = 0.176; one-way ANOVA; **Figure 8F*ii***).

To determine the role of upstream targets that facilitate excitatory input to Th^{EGFP} VTA

cells, we isolated presynaptic excitatory input (mEPSC) with TTX pretreatment (**Figure 9A**).

MCH-mediated glutamate release at Th^{EGFP} VTA cells was largely abolished in the presence of

TTX (**Figure 9B**). However, a slight 1.5 ± 0.4-fold (baseline: 1.2 ± 0.5 Hz; MCH: 1.5 ± 0.6 Hz,

n = 6; t(5) = 2.27, p = 0.072, $\eta^2 = 0.51$, paired t test) increase in mEPSC frequency can be

detected at some Th^{EGFP} VTA (**Figure 9B**), thus it is possible that MCH mediated glutamate

release directly at glutamatergic terminals. There was no change in mEPSC amplitude following

MCH treatment (baseline: 14.9 ± 1.9 pA; MCH: 14.9 ± 1.7 pA, n = 6; t(5) = 0.05, p = 0.961, η^2 =

0.001, paired t test) or over time (**Figure 9C**). We thus considered that MCH may recruit

upstream targets that indirectly regulate excitatory input at Th^{EGFP} VTA cells.

As MCHR1 is a known inhibitory $G_{i/o}$-protein coupled receptor (Hawes et al., 2000), we

determined if the MCH-mediated increase in glutamatergic tone at Th^{EGFP} VTA cells may be

ascribed to GABAergic disinhibition at glutamatergic nerve terminals (**Figure 9D**). We recorded

from $Vglut2^{EGFP}$ VTA cells and found that MCH produced a prolonged 0.4 ± 0.1-fold decrease in

sIPSC frequency (baseline: 1.7 ± 0.4 Hz; MCH: 1.0 ± 0.3 Hz, n = 5; t(4) = 5.08, p = 0.007, η^2 =

0.87, paired t test; **Figure 9E**) but not amplitude of sIPSC events (baseline: 14.4 ± 1.3 pA; MCH:

19.3 ± 5.0 pA, n = 5; t(4) = 0.95, p = 0.395, η^2 = 0.19, paired t test; **Figure 9F**). Overall, these

recordings suggested that, in addition to directly inhibiting Th VTA cells, MCH may also engage

a local disynaptic circuit from $Vgat$ and $Vglut2$ VTA cells to regulate VTA output (**Figure 9G**).

Discussion

MCH can directly and indirectly reglate Th VTA cells. MCHR1-expressing cells were

widespread through the VTA, including at glutamatergic $Vglut2$, GABAergic $Vgat$, and

dopaminergic TH cells, thus providing a neuroanatomical basis for MCH action in the VTA.

Electrophysiological recordings showed that MCH hyperpolarized and inhibited Th cells and

$Vgat$ cells. Interestingly, MCH did not directly regulate $Vglut2$ cells but disinhibited $Vglut2$

afferents leading to increased excitatory transmission at *Th* cells. Therefore, functional MCHR1

activation in the VTA may provide bidirectional regulation of dopamine output from the VTA.

The majority, nearly 80%, of TH-ir VTA cells expressed *Mchr1* mRNA, and MCH

directly hyperpolarized 38% of *Th* VTA cells in an activity-independent manner, which is

consistent with MCHR1 protein expression at TH-ir VTA cells. Furthermore, it is possible that

native EGFP fluorescence may not reflect active TH cells, as Cre expression in TH-negative

VTA cells may comprise up to half of all EGFPTh cells (Lammel et al., 2015). However, when

patching within the VTA, we found over 80% of patched EGFPTh cells expressed TH, indicating

more robust EGFPTh expression in TH cells than previously reported.

MCHR1 signaling has known inhibitory actions (Chambers et al., 1999; Hawes et al.,

2000; Lembo et al., 1999; Saito et al., 1999), including within the mesolimbic system connecting

the VTA and ventral striatum. In the accumbens, MCH hyperpolarizes medium spiny neurons

(Georgescu et al., 2005) and suppresses dopamine release (Chee et al., 2019). Furthermore, loss

of MCH signaling increases accumbens dopamine accumulation leading to hyperactivity and

increased energy expenditure (M. J. Chee et al., 2019; Mul et al., 2011; Pissios et al., 2008). As

the main source of dopaminergic afferents to the accumbens arises from the VTA, it stands to

reason that MCH may regulate dopamine release by direct inhibition of VTA neurons.

Interestingly, MCH action within the VTA caused a delayed increase in excitatory input

to *Th* VTA cells. MCHR1 activation engages $G_{i/o}$- or G_q-protein systems (Pissios et al., 2003;

Sears et al., 2010) to elicit inhibitory or excitatory actions, respectively. We considered that

MCH may act presynaptically to stimulate glutamatergic afferents, but the MCH-mediated

increase in glutamatergic transmission was abolished in the presence of TTX. This suggested that

the stimulatory effect of MCH on glutamatergic afferents was indirect, activity-dependent, and

polysynaptic. We thus examined if MCH may indirectly recruit glutamatergic afferents detected

at *Th* VTA cells by suppressing GABAergic input onto *Vglut2* VTA cells or axons. We found

that MCH directly inhibited GABAergic VTA cells and suppressed GABAergic input to *Vglut2*

VTA cells. In effect, MCH can suppress GABAergic activity to disinhibit glutamatergic

transmission at *Th* VTA cells, and we posit that this GABA-dependent microcircuit within the

VTA may serve to restore the output of *Th* VTA cells that were acutely inhibited by MCH

(**Figure 9G**).

VTA *Vglut2* cells are known to innervate (Dobi et al., 2010) and regulate dopaminergic

VTA cells (Bariselli et al., 2016; Wang et al., 2015). MCH had no effect on the membrane

potential of *Vglut2* VTA cells. *Vglut2* cells were more abundant in the anterior VTA (Morales &

Root, 2014; Root et al., 2018; Yamaguchi et al., 2007), thus it is possible that the null effect of

MCH at *Vglut2* cells is ascribed to differences in electrophysiological effects in the anterior and

posterior VTA (Guan et al., 2012). It may thus be insightful to track the rostrocaudal distribution

of subsequent patch-clamp recordings. By contrast, MCH increased glutamatergic tone at *Th*

VTA cells. *Vglut2* VTA cells form local synapses with dopaminergic and non-dopaminergic

VTA neurons (Dobi et al., 2010) and are pivotal regulators of local VTA circuitry and output.

Glutamatergic VTA cells can drive dopamine release and promote conditioned place preference

(Wang et al., 2015) or drug-induced conditioned place preference (Mead & Stephens, 1999;

Slusher et al., 2001; Tzschentke & Schmidt, 1998) that can be enhanced by food restriction

(Stuber et al., 2002) to assess drug reward (Hiroi & White, 1991). Interestingly, *Mchr1* deletion

can block the effect of food restriction on amphetamine-induced conditioned place preference

(Geuzaine et al., 2014). However, as *Vglut2* cells within the VTA were not MCH-sensitive, it

would be important to consider that MCH may indirectly modulate dopamine output at

glutamatergic afferents arising from outside the VTA, such as from the prefrontal cortex (Beier

et al., 2015) or lateral hypothalamus (Geisler & Wise, 2008; Watabe-Uchida et al., 2012).

Furthermore, since we did not detect postsynaptic MCH effects at the soma of *Vglut2* cells, we

predicted that MCHR1 distribution within *Vglut2* cells would accumulate at presynaptic

terminals to regulate glutamate release at projection sites (**Figure 9G**, *asterisk*).

Prominent *Vgat* expression in the posterior VTA may partly form the tail of the VTA

(tVTA) or rostromedial tegmental nucleus (RMTg; Jhou et al., 2009). The GABAergic cell

cluster in the posterior VTA also reflected the highest proportion of *Vgat* VTA cells that

expressed *Mchr1*. This supports the hypothesis that the posterior VTA is anatomically and

functionally unique (Lahti et al., 2016; Smith et al., 2019) and suggests that this subregion is a

site of MCH action. Sex may interact with *Vgat* expression at *Mchr1* cells in the VTA, as there is

a larger portion of *Mchr1* cells expressing *Vgat* in the VTA of female mice. Therefore, MCH

may contribute to sex-specific regulation of VTA output, potentially via *Vgat* cells in females or

by yet characterized *Mchr1*-only cells in males. The tVTA/RMTg has been implicated in

responsiveness to alcohol as exposure to alcohol stimulates tVTA/RMTg neurons, and rats will

self-administer ethanol to the posterior but not anterior VTA (Guan et al., 2012). Interestingly,

Mchr1 deletion suppresses alcohol-induced conditioned place preference (Karlsson et al., 2016),

and MCHR1 blockade decreases self-administration of alcohol (Cippitelli et al., 2010). The

posterior VTA supplies dense GABAergic projections to other VTA neurons (Barrot et al., 2012;

Jhou et al., 2009; Lecca et al., 2012) and provides strong inhibitory tone to dopaminergic VTA

neurons (Barrot et al., 2012; Bouarab et al., 2019; Guan et al., 2012). While MCH did not

regulate GABAergic synaptic transmission at *Th* VTA cells, it inhibited GABAergic

transmission at glutamatergic cells. Furthermore, MCH directly inhibited *Vgat* cells, thus, taken

together, MCH action at *Vgat* cells in the posterior VTA may weaken inhibition of *Th* VTA cells

to indirectly facilitate dopaminergic output.

Consistent with findings in rats (Hervieu et al., 2000; Saito et al., 2001), the relative

expression of *Mchr1* mRNA in the VTA was comparable to that in the hypothalamus and

hippocampus. *Mchr1* expression was visible in all major subtypes of VTA cells, including cells

that coexpressed two or more chemical messengers. *Vgat*/*Vglut2* cells were the most common

subpopulation coexpressing a combination of TH, *Vgat*, or *Vglut2* and is consistent with previous

reports of GABA and glutamate corelease from VTA cells (Root et al., 2018; Root, Mejias-

Aponte, Zhang, et al., 2014). VTA neurons may also be both glutamatergic and dopaminergic

(Morales & Root, 2014; Morales & Margolis, 2017; Yamaguchi et al., 2011), but unlike the rat

VTA, the incidence of colocalization between TH and *Vglut2* was low. In rats, most

dopaminergic and glutamatergic VTA cells were at or very close to the midline (Yamaguchi et

al., 2011) and includes midline regions encompassing the interfascicular nucleus raphe, rostral

linear nucleus raphe, and central linear nucleus raphe as subregions of the VTA (Morales &

Root, 2014; Morales & Margolis, 2017; Yamaguchi et al., 2011). However, our analyses were

based on the neuroanatomical boundaries of the VTA as defined by the *Allen Reference Atlas*

(Dong, 2008) and so did not include midline outside the VTA where dual TH/*Vglut2* cells may

be most abundant. Finally, although VTA cells may co-release dopamine and GABA (Stamatakis

et al., 2013; Tritsch et al., 2012), coexpression of TH and *Vgat* did not form a substantial

population within the VTA. GABA release does not require vesicular transport via vGAT

because the vesicular monoamine transporter 2 (vMAT2) can fully support GABA release from

dopaminergic cells (Tritsch et al., 2012), thus it is not surprising that TH/*Vgat* coexpression was

low.

In conclusion, our findings suggest that MCHR1 expression at multiple VTA cell types

can modulate the local VTA microcircuit to provide bidirectional regulation of dopamine output

from the VTA. MCH directly inhibited dopaminergic VTA cells, which may acutely suppress

dopamine release. However, MCH also inhibited GABAergic VTA cells, which may disinhibit

glutamatergic input to dopamine cells. This GABA-mediated disinhibition has a delayed time

course and may act to restore dopamine levels acutely lowered by the direct MCH-mediated

inhibition of dopamine cells. Paradoxically, this may be in contrast to prior findings that *Mchr1*

deletion from *Vgat* neurons results in a hyperdopaminergic state (Chee et al., 2019), but this and

previous findings may not be mutually exclusive, as MCH can have site-specific effects at

different nodes of the mesolimbic dopamine pathway. It would be of interest to test our proposed

model (**Figure 9G**) in future studies and determine if *Mchr1* deletion from dopaminergic cells

would also elicit a hyperdopaminergic state, or whether disrupting GABAergic transmission

would abolish the facilitation of glutamatergic input to dopamine cells.

References

Advanced Cell Diagnostics. (2020). RNAscope ® Multiplex Fluorescent v2 Assay combined

with Immunofluorescence - Integrated Co-Detection Workflow (ICW) Combinations.

Advanced Cell Diagnostics Inc., 1–5.

Albanese, A., & Minciacchi, D. (1983). Organization of the ascending projections from the

ventral tegmental area: A multiple fluorescent retrograde tracer study in the rat. *Journal of*

Comparative Neurology, *216*(4), 406–420. https://doi.org/10.1002/cne.902160406

Åstrand, A., Bohlooly, M., Larsdotter, S., Mahlapuu, M., Andersén, H., Tornell, J., Ohlsson, C.,

Snaith, M., & Morgan, D. G. A. (2004). Mice lacking melanin-concentrating hormone

receptor 1 demonstrate increased heart rate associated with altered autonomic activity.

American Journal of Physiology - Regulatory Integrative and Comparative Physiology,

287(4 56-4), 749–758. https://doi.org/10.1152/ajpregu.00134.2004

Bansal, R., Engle, S. E., Kamba, T. K., Brewer, K. M., Lewis, W. R., & Berbari, N. F. (2021).

Artificial intelligence approaches to assessing primary cilia. *Journal of Visualized*

Experiments, *2021*(171), 1–21. https://doi.org/10.3791/62521

Bariselli, S., Glangetas, C., Tzanoulinou, S., & Bellone, C. (2016). Ventral tegmental area

subcircuits process rewarding and aversive experiences. *Journal of Neurochemistry*, *139*(6),

1071–1080. https://doi.org/10.1111/jnc.13779

Barrot, M., Sesack, S. R., Georges, F., Pistis, M., Hong, S., & Jhou, T. C. (2012). Braking

dopamine systems: A new GABA master structure for mesolimbic and nigrostriatal

functions. *Journal of Neuroscience*, *32*(41), 14094–14101.

https://doi.org/10.1523/JNEUROSCI.3370-12.2012

Beier, K. T., Steinberg, E. E., DeLoach, K. E., Xie, S., Miyamichi0, K., Schwarz, L., Gao, X. J.,

Kremer, E. J., Malenka, R. C., & Luo, L. (2015). Circuit Architecture of VTA Dopamine

Neurons Revealed by Systematic Input–Output Mapping. *Cell*, *162*(3), 622–634.

https://doi.org/10.1016/j.cell.2015.07.015.Circuit

Bittencourt, J. C., Presse, F., Arias, C., Peto, C., Vaughan, J., Nahon, J. ☐L, Vale, W., &

Sawchenko, P. E. (1992). The melanin☐concentrating hormone system of the rat brain: An

immuno☐ and hybridization histochemical characterization. *Journal of Comparative*

Neurology, *319*(2), 218–245. https://doi.org/10.1002/cne.903190204

Bono, B. S., Koziel Ly, N. K., Miller, P. A., Williams-Ikhenoba, J., Dumiaty, Y., & Chee, M. J.

(2022). Spatial distribution of beta-klotho mRNA in the mouse hypothalamus, hippocampal

region, subiculum, and amygdala. *Journal of Comparative Neurology*, *530*(10), 1634–1657.

https://doi.org/10.1002/cne.25306

Bouarab, C., Thompson, B., & Polter, A. M. (2019). VTA GABA Neurons at the Interface of

Stress and Reward. *Frontiers in Neural Circuits*, *13*(December), 1–12.

https://doi.org/10.3389/fncir.2019.00078

Chambers, J., Ames, R. S., Bergsma, D., Muir, A., Fitzgerald, L. R., Hervieu, G., Dytko, G. M.,

Foley, J. J., Martin, J., Liu, W. S., Park, J., Ellis, C., Ganguly, S., Konchar, S., Cluderay, J.,

Leslie, R., Wilson, S., & Sarau, H. M. (1999). Melanin-concentrating hormone is the

cognate ligand for the orphan G- protein-coupled receptor SLC-1. *Nature*, *400*(6741), 261–

265. https://doi.org/10.1038/22313

Chee, M. J., Hebert, A. J., Briançon, N., Flaherty, S. E., Pissios, P., & Maratos-Flier, E. (2019).

Conditional deletion of melanin-concentrating hormone receptor 1 from GABAergic

neurons increases locomotor activity. *Molecular Metabolism*, *29*(August), 114–123. https://doi.org/10.1016/j.molmet.2019.08.018

Chee, M., Myers, M. G., Price, C. J., & Colmers, W. F. (2010). Neuropeptide Y suppresses anorexigenic output from the ventromedial nucleus of the hypothalamus. *Journal of Neuroscience*, *30*(9), 3380–3390. https://doi.org/10.1523/JNEUROSCI.4031-09.2010

Chen, Y., Hu, C., Hsu, C. K., Zhang, Q., Bi, C., Asnicar, M., Hsiung, H. M., Fox, N., Slieker, L. J., Yang, D. D., Heiman, M. L., & Shi, Y. (2002). Targeted disruption of the melanin-concentrating hormone receptor-1 results in hyperphagia and resistance to diet-induced obesity. *Endocrinology*, *143*(7), 2469–2477. https://doi.org/10.1210/endo.143.7.8903

Cippitelli, A., Karlsson, C., Shaw, J. L., Thorsell, A., Gehlert, D. R., & Heilig, M. (2010). Suppression of alcohol self-administration and reinstatement of alcohol seeking by melanin-concentrating hormone receptor 1 (MCH1-R) antagonism in Wistar rats. *Psychopharmacology*, *211*(4), 367–375. https://doi.org/10.1007/s00213-010-1891-y

Dal Bo, G., St.-Gelais, F., Danik, M., Williams, S., Cotton, M., & Trudeau, L. E. (2004). Dopamine neurons in culture express VGLUT2 explaining their capacity to release glutamate at synapses in addition to dopamine. *Journal of Neurochemistry*, *88*(6), 1398–1405. https://doi.org/10.1046/j.1471-4159.2003.02277.x

Diniz, G. B., Battagello, D. S., Klein, M. O., Bono, B. S. M., Ferreira, J. G. P., Motta-Teixeira, L. C., Duarte, J. C. G., Presse, F., Nahon, J. L., Adamantidis, A., Chee, M. J., Sita, L. V., & Bittencourt, J. C. (2020). Ciliary melanin-concentrating hormone receptor 1 (MCHR1) is widely distributed in the murine CNS in a sex-independent manner. *Journal of Neuroscience Research*, *98*(10), 2045–2071. https://doi.org/10.1002/jnr.24651

Dobi, A., Margolis, E. B., Wang, H. L., Harvey, B. K., & Morales, M. (2010). Glutamatergic and

nonglutamatergic neurons of the ventral tegmental area establish local synaptic contacts

with dopaminergic and nondopaminergic neurons. *Journal of Neuroscience*, *30*(1), 218–

229. https://doi.org/10.1523/JNEUROSCI.3884-09.2010

Domingos, A. I., Sordillo, A., Dietrich, M. O., Liu, Z. W., Tellez, L. A., Vaynshteyn, J., Ferreira,

J. G., Ekstrand, M. I., Horvath, T. L., de Araujo, I. E., & Friedman, J. M. (2013).

Hypothalamic melanin concentrating hormone neurons communicate the nutrient value of

sugar. *ELife*, *2013*(2), 1–15. https://doi.org/10.7554/eLife.01462

Dong, H. (2008). *The Allen reference atlas: A digital color brain atlas of the C57BL/6J male*

mouse. John Wiley & Sons.

Faget, L., Osakada, F., Duan, J., Ressler, R., Johnson, A. B., Proudfoot, J. A., Yoo, J. H.,

Callaway, E. M., & Hnasko, T. S. (2016). Afferent Inputs to Neurotransmitter-Defined Cell

Types in the Ventral Tegmental Area. *Cell Reports*, *15*(12), 2796–2808.

https://doi.org/10.1016/j.celrep.2016.05.057

Gao, X.-B., & Pol, A. N. van den. (2001). Melanin concentrating hormone depresses synaptic

activity of glutamate and GABA neurons from rat lateral hypothalamus. *Journal of*

Physiology, *533*(1), 237–252.

Geisler, S., & Wise, R. A. (2008). Functional implications of glutamatergic projections to the

ventral tegmental area. *Reviews in the Neurosciences*, *19*(4–5), 227–244.

https://doi.org/10.1515/REVNEURO.2008.19.4-5.227

Georgescu, D., Sears, R. M., Hommel, J. D., Barrot, M., Bolaños, C. A., Marsh, D. J., Bednarek,

M. A., Bibb, J. A., Maratos-Flier, E., Nestler, E. J., & DiLeone, R. J. (2005). The

hypothalamic neuropeptide melanin-concentrating hormone acts in the nucleus accumbens

to modulate feeding behavior and forced-swim performance. *Journal of Neuroscience*,

25(11), 2933–2940. https://doi.org/10.1523/JNEUROSCI.1714-04.2005

Geuzaine, A., Tyhon, A., Grisar, T., Brabant, C., Lakaye, B., & Tirelli, E. (2014). Amphetamine

reward in food restricted mice lacking the melanin-concentrating hormone receptor-1.

Behavioural Brain Research, *262*, 14–20. https://doi.org/10.1016/j.bbr.2013.12.052

Guan, Y., Xiao, C., Krnjević, K., Xie, G., Zuo, W., & Ye, J.-H. (2012). GABAergic Actions

Mediate Opposite Ethanol Effects on Dopaminergic Neurons in the Anterior and Posterior

Ventral Tegmental Area. *Journal of Pharmacology and Experimental Therapeutics*, *341*(1),

33–42. https://doi.org/10.1124/jpet.111.187963

Hawes, B. E., Erin, K. I. L., Green, B., O'Neill, K. I. M., Fried, S., & Graziano, M. P. (2000).

The melanin-concentrating hormone receptor couples to multiple G proteins to activate

diverse intracellular signaling pathways. *Endocrinology*, *141*(12), 4524–4532.

https://doi.org/10.1210/endo.141.12.7833

Hervieu, G. J., Cluderay, J. E., Harrison, D., Meakin, J., Maycox, P., Nasir, S., & Leslie, R. A.

(2000). The distribution of the mRNA and protein products of the melanin-concentrating

hormone (MCH) receptor gene, slc-1, in the central nervous system of the ratt. *European*

Journal of Neuroscience, *12*(4), 1194–1216. https://doi.org/10.1046/j.1460-

9568.2000.00008.x

Hiroi, N., & White, N. M. (1991). The amphetamine conditioned place preference: differential

involvement of dopamine receptor subtypes and two dopaminergic terminal areas. *Brain*

Research, *552*(1), 141–152. https://doi.org/10.1016/0006-8993(91)90672-I

Jhou, T., Fields, H., Baxter, M., & Saper, C. (2009). The rostromedial tegmental nucleus

(RMTg), a major GABAergic afferent to midbrain dopamine neurons, selectively encodes

aversive stimuli and promotes behavioral inhibition. *Neuron*, *61*(5), 786–800.

https://doi.org/10.1016/j.neuron.2009.02.001.The

Karlsson, C., Rehman, F., Damdazic, R., Atkins, A. L., Schank, J. R., Gehlert, D. R., Steensland,

P., Thorsell, A., & Heilig, M. (2016). The melanin-concentrating hormone-1 receptor

modulates alcohol-induced reward and DARPP-32 phosphorylation. *Psychopharmacology*,

233(12), 2355–2363. https://doi.org/10.1007/s00213-016-4285-y

Kokkotou, E., Jeon, J. Y., Wang, X., Marino, F. E., Carlson, M., Trombly, D. J., & Maratos-

Flier, E. (2005). Mice with MCH ablation resist diet-induced obesity through strain-specific

mechanisms. *American Journal of Physiology - Regulatory Integrative and Comparative*

Physiology, *289*(1 58-1), 117–124. https://doi.org/10.1152/ajpregu.00861.2004

Krashes, M. J., Shah, B. P., Madara, J. C., Olson, D. P., Strochlic, D. E., Garfield, A. S., Vong,

L., Pei, H., Watabe-Uchida, M., Uchida, N., Liberles, S. D., & Lowell, B. B. (2014). An

excitatory paraventricular nucleus to AgRP neuron circuit that drives hunger. *Nature*,

507(7491), 238–242. https://doi.org/10.1038/nature12956.A

Lahti, L., Haugas, M., Tikker, L., Airavaara, M., Voutilainen, M. H., Anttila, J., Kumar, S.,

Inkinen, C., Salminen, M., & Partanen, J. (2016). Differentiation and molecular

heterogeneity of inhibitory and excitatory neurons associated with midbrain dopaminergic

nuclei. *Development (Cambridge)*, *143*(3), 516–529. https://doi.org/10.1242/dev.129957

Lammel, S., Steinberg, E. E., Földy, C., Wall, N. R., Beier, K., Luo, L., & Malenka, R. C.

(2015). Diversity of Transgenic Mouse Models for Selective Targeting of Midbrain

Dopamine Neurons. *Neuron*, *85*(2), 429–438.

https://doi.org/10.1053/j.gastro.2016.08.014.CagY

Lecca, S., Melis, M., Luchicchi, A., Muntoni, A. L., & Pistis, M. (2012). Inhibitory inputs from rostromedial tegmental neurons regulate spontaneous activity of midbrain dopamine cells and their responses to drugs of abuse. *Neuropsychopharmacology*, *37*(5), 1164–1176. https://doi.org/10.1038/npp.2011.302

Lembo, P. M. C., Grazzini, E., Cao, J., Hubatsch, D. A., Pelletier, M., Hoffert, C., St-Onge, S., Pou, C., Labrecque, J., Groblewski, T., O'Donnell, D., Payza, K., Ahmad, S., & Walker, P. (1999). The receptor for the orexigenic peptide melanin-concentrating hormone is a G-protein-coupled receptor. *Nature Cell Biology*, *1*(5), 267–271. https://doi.org/10.1038/12978

Lindeberg, J., Usoskin, D., Bengtsson, H., Gustafsson, A., Kylberg, A., Söderström, S., & Ebendal, T. (2004). Transgenic expression of Cre recombinase from the tyrosine hydroxylase locus. *Genesis (United States)*, *40*(2), 67–73. https://doi.org/10.1002/gene.20065

Lord, M. N., Subramanian, K., Kanoski, S. E., Noble, E. E., Angeles, L., Section, E. B., California, S., & Angeles, L. (2022). Melanin-concentrating hormone and food intake control: sites of action, peptide interactions, and appetition. *Peptides*, *137*, 1–23. https://doi.org/10.1016/j.peptides.2020.170476.Melanin-concentrating

Ludwig, D. S., Tritos, N. A., Mastaitis, J. W., Kulkarni, R., Kokkotou, E., Elmquist, J., Lowell, B., Flier, J. S., & Maratos-Flier, E. (2001). Melanin-concentrating hormone overexpression in transgenic mice leads to obesity and insulin resistance. *Journal of Clinical Investigation*,

107(3), 379–386. https://doi.org/10.1172/JCI10660

Marsh, D. J., Weingarth, D. T., Novi, D. E., Chen, H. Y., Trumbauer, M. E., Chen, A. S., Guan,

X. M., Jiang, M. M., Feng, Y., Camacho, R. E., Shen, Z., Frazier, E. G., Yu, H., Metzger, J.

M., Kuca, S. J., Shearman, L. P., Gopal-Truter, S., MacNeil, D. J., Strack, A. M., … Qian,

S. (2002). Melanin-concentrating hormone 1 receptor-deficient mice are lean, hyperactive,

and hyperphagic and have altered metabolism. *Proceedings of the National Academy of*

Sciences of the United States of America, *99*(5), 3240–3245.

https://doi.org/10.1073/pnas.052706899

Mead, A. N., & Stephens, D. N. (1999). CNQX but not NBQX prevents expression of

amphetamine-induced place preference conditioning: a role for the glycine site of the

NMDA receptor, but not AMPA receptors. *The Journal of Pharmacology and Experimental*

Therapeutics, *290*(1), 9–15.

Mickelsen, L. E., Bolisetty, M., Chimileski, B. R., Fujita, A., Beltrami, E. J., Costanzo, J. T.,

Naparstek, J. R., Robson, P., & Jackson, A. C. (2019). Single-cell transcriptomic analysis of

the lateral hypothalamic area reveals molecularly distinct populations of inhibitory and

excitatory neurons. *Nature Neuroscience*, *22*(4), 642–656. https://doi.org/10.1038/s41593-

019-0349-8

Morales, M., & Root, D. H. (2014). Glutamate neurons within the midbrain dopamine regions.

Neuroscience, *282*, 60–68. https://doi.org/10.1016/j.neuroscience.2014.05.032

Morales, Marisela, & Margolis, E. B. (2017). Ventral tegmental area: Cellular heterogeneity,

connectivity and behaviour. *Nature Reviews Neuroscience*, *18*(2), 73–85.

https://doi.org/10.1038/nrn.2016.165

Mul, J. D., la Fleur, S. E., Toonen, P. W., Afrasiab-Middelman, A., Binnekade, R., Schetters, D.,

Verheij, M. M. M., Sears, R. M., Homberg, J. R., Schoffelmeer, A. N. M., Adan, R. A. H.,

DiLeone, R. J., de Vries, T. J., & Cuppen, E. (2011). Chronic loss of melanin-concentrating

hormone affects motivational aspects of feeding in the rat. *PLoS ONE*, *6*(5).

https://doi.org/10.1371/journal.pone.0019600

Negishi, K., Payant, M. A., Schumacker, K. S., Wittmann, G., Butler, R. M., Lechan, R. M.,

Steinbusch, H. W. M., Khan, A. M., & Chee, M. J. (2020). Distributions of hypothalamic

neuron populations coexpressing tyrosine hydroxylase and the vesicular GABA transporter

in the mouse. *Journal of Comparative Neurology*, *528*(11), 1833–1855.

https://doi.org/10.1002/cne.24857

Nieh, E. H., Vander Weele, C. M., Matthews, G. A., Presbrey, K. N., Wichmann, R., Leppla, C.

A., Izadmehr, E. M., & Tye, K. M. (2016). Inhibitory Input from the Lateral Hypothalamus

to the Ventral Tegmental Area Disinhibits Dopamine Neurons and Promotes Behavioral

Activation. *Neuron*, *90*(6), 1286–1298. https://doi.org/10.1016/j.neuron.2016.04.035

Pissios, P., Frank, L., Kennedy, A. R., Porter, D. R., Marino, F. E., Liu, F. F., Pothos, E. N., &

Maratos-Flier, E. (2008). Dysregulation of the Mesolimbic Dopamine System and Reward

in MCH-/- Mice. *Biological Psychiatry*, *64*(3), 184–191.

https://doi.org/10.1016/j.biopsych.2007.12.011

Pissios, P., Trombly, D. J., Tzameli, I., & Maratos-Flier, E. (2003). Melanin-concentrating

hormone receptor 1 activates extracellular signal-regulated kinase and synergizes with Gs-

coupled pathways. *Endocrinology*, *144*(8), 3514–3523. https://doi.org/10.1210/en.2002-

0004

Qu, D., Ludwig, D. S., Gammeltoftt, S., Piper, M., Pelleymountert, M. A., Cullent, M. J.,

Mathes, W. F., Przypek, J., Kanarek, R., & Ii, E. M. (1996). A role for melanin-

concentrating hormone in the central regulation of feeding behaviour. *Nature*, *380*(March),

243–247.

Root, D. H., Mejias-Aponte, C. A., Qi, J., & Morales, M. (2014). Role of glutamatergic

projections from ventral tegmental area to lateral Habenula in aversive conditioning.

Journal of Neuroscience, *34*(42), 13906–13910. https://doi.org/10.1523/JNEUROSCI.2029-

14.2014

Root, D. H., Mejias-Aponte, C. A., Zhang, S., Wang, H. L., Hoffman, A. F., Lupica, C. R., &

Morales, M. (2014). Single rodent mesohabenular axons release glutamate and GABA.

Nature Neuroscience, *17*(11), 1543–1551. https://doi.org/10.1038/nn.3823

Root, D. H., Zhang, S., Barker, D. J., Miranda-Barrientos, J., Liu, B., Wang, H. L., & Morales,

M. (2018). Selective Brain Distribution and Distinctive Synaptic Architecture of Dual

Glutamatergic-GABAergic Neurons. *Cell Reports*, *23*(12), 3465–3479.

https://doi.org/10.1016/j.celrep.2018.05.063

Saito, Y., Cheng, M., Leslie, F. M., & Civelli, O. (2001). Expression of the melanin-

concentrating hormone (MCH) receptor mRNA in the rat brain. *Journal of Comparative*

Neurology, *435*(1), 26–40. https://doi.org/10.1002/cne.1191

Saito, Y., Nothacker, H., Wang, Z., Lin, S. H. S., Leslie, F., & Civelli, O. (1999). Molecular

characterization of the melanin-concentrating- hormone receptor. *Nature*, *400*(July), 265–

269.

Schneeberger, M., Tan, K., Nectow, A. R., Parolari, L., Caglar, C., Azevedo, E., Li, Z.,

Domingos, A., & Friedman, J. M. (2018). Functional analysis reveals differential effects of

glutamate and MCH neuropeptide in MCH neurons. *Molecular Metabolism*, *13*(May), 83–

89. https://doi.org/10.1016/j.molmet.2018.05.001

Sears, R. M., Liu, R. J., Narayanan, N. S., Sharf, R., Yeckel, M. F., Laubach, M., Aghajanian, G.

K., & DiLeone, R. J. (2010). Regulation of nucleus accumbens activity by the hypothalamic

neuropeptide melanin-concentrating hormone. *Journal of Neuroscience*, *30*(24), 8263–

8273. https://doi.org/10.1523/JNEUROSCI.5858-09.2010

Shimada, M., Tritos, N. A., Lowell, B. B., Flier, J. S., & Maratos-Flier, E. (1998). Mice lacking

melanin- concentrating hormone are hypophagic and lean. *Nature*, *396*(December), 3–7.

Slusher, B. S., Thomas, A., Paul, M., Schad, C. A., & Ashby, C. R. (2001). Expression and

acquisition of the conditioned place preference response to cocaine in rats is blocked by

selective inhibitors of the enzyme N-Acetylated-α-Linked-Acidic Dipeptidase

(NAALADase). *Synapse*, *41*(1), 22–28. https://doi.org/10.1002/syn.1056

Smith, R. J., Vento, P. J., Chao, Y. S., Good, C. H., Thomas, C., & Ground, A. P. (2019). Gene

expression and neurochemical characterization of the rostromedial tegmental nucleus

(RMTg) in rats and mice. *Brain Struct Funct*, *224*(1), 219–238.

https://doi.org/10.1007/s00429-018-1761-7.Gene

Stamatakis, A. M., Jennings, J. H., Ung, R. L., Blair, G. A., Weinberg, J., Neve, R. L., Boyce, F.,

Mattis, J., Deisseroth, K., & Stuber, G. D. (2013). A unique population of ventral tegmental

area neurons inhibits the lateral habenula to promote reward. *Neuron*, *80*(4), 1–23.

https://doi.org/10.1016/j.neuron.2013.08.023.A

Stuber, G. D., Evans, S. B., Higgins, M. S., Pu, Y., & Figlewicz, D. P. (2002). Food restriction

modulates amphetamine-conditioned place preference and nucleus accumbens dopamine

release in the rat. *Synapse*, *46*(2), 83–90. https://doi.org/10.1002/syn.10120

Stuber, G. D., Hnasko, T. S., Britt, J. P., Edwards, R. H., & Bonci, A. (2010). Dopaminergic

terminals in the nucleus accumbens but not the dorsal striatum corelease glutamate. *Journal*

of Neuroscience, *30*(24), 8229–8233. https://doi.org/10.1523/JNEUROSCI.1754-10.2010

Sulzer, D., Joyce, M. P., Lin, L., Geldwert, D., Haber, S. N., Hattori, T., & Rayport, S. (1998).

Dopamine neurons make glutamatergic synapses in vitro. *Journal of Neuroscience*, *18*(12),

4588–4602. https://doi.org/10.1523/jneurosci.18-12-04588.1998

Tecuapetla, F., Patel, J. C., Xenias, H., English, D., Tadros, I., Shah, F., Berlin, J., Deisseroth,

K., Rice, M. E., Tepper, J. M., & Koos, T. (2010). Glutamatergic signaling by mesolimbic

dopamine neurons in the nucleus accumbens. *Journal of Neuroscience*, *30*(20), 7105–7110.

https://doi.org/10.1523/JNEUROSCI.0265-10.2010

Ting, J. T., Lee, B. R., Chong, P., Soler-Llavina, G., Cobbs, C., Koch, C., Zeng, H., & Lein, E.

(2018). Preparation of Acute Brain Slices Using an Optimized N-Methyl-D-glucamine

Protective Recovery Method. *Journal of Visualized Experiments□: JoVE*, *132*, 1–13.

https://doi.org/10.3791/53825

Tritsch, N. X., Ding, J. B., & Sabatini, B. L. (2012). Dopaminergic neurons inhibit striatal output

via non-canonical release of GABA. *Nature*, *490*, 262–266.

https://doi.org/doi:10.1038/nature11466

Tzschentke, T. M., & Schmidt, W. J. (1998). Blockade of morphine- and amphetamine-induced

conditioned place preference in the rat by riluzole. *Neuroscience Letters*, *242*, 114–116.

papers2://publication/uuid/3FF1E7DE-1CAD-497E-A55B-478F8D972A24

930 Vong, L., Ye, C., Yang, Z., Choi, B., Jr., S. C., & Lowell, B. B. (2011). Leptin Action on

931   GABAergic Neurons Prevents Obesity and Reduces Inhibitory Tone to POMC Neurons.

932   *Neuron*, *71*(1), 142–154. https://doi.org/10.1016/j.neuron.2011.05.028.Leptin

933 Wang, H. L., Qi, J., Zhang, S., Wang, H., & Morales, M. (2015). Rewarding effects of optical

934   stimulation of ventral tegmental area glutamatergic neurons. *Journal of Neuroscience*,

935   *35*(48), 15948–15954. https://doi.org/10.1523/JNEUROSCI.3428-15.2015

936 Watabe-Uchida, M., Zhu, L., Ogawa, S. K., Vamanrao, A., & Uchida, N. (2012). Whole-Brain

937   Mapping of Direct Inputs to Midbrain Dopamine Neurons. *Neuron*, *74*(5), 858–873.

938   https://doi.org/10.1016/j.neuron.2012.03.017

939 Yamaguchi, T., Sheen, W., & Morales, M. (2007). Glutamatergic neurons are present in the rat

940   ventral tegmental area. *Eur J Neurosci*, *25*(1), 106–118. https://doi.org/doi:10.1111/j.1460-

941   9568.2006.05263.x.

942 Yamaguchi, T., Wang, H. L., Li, X., Ng, T. H., & Morales, M. (2011). Mesocorticolimbic

943   glutamatergic pathway. *Journal of Neuroscience*, *31*(23), 8476–8490.

944   https://doi.org/10.1523/JNEUROSCI.1598-11.2011

Legends

Figure 1. *Mchr1* mRNA and MCHR1 protein expression in wildtype mouse VTA. qPCR analysis of *Mchr1* mRNA from the hypothalamus (HT), striatum (STR), hippocampus (HC), ventral tegmental area (VTA), and cerebellum (CB) of male and female wildtype mice (**A**). Confocal photomicrographs of MCHR1-immunoreactive signals on Nissl-stained (NeuroTrace) soma in wildtype (**B**) and *Mchr1-KO* (**C**) VTA at low (***i***) and high magnification (***ii***, from dash-outlined region in ***i***). Representative examples of the presence (white arrowhead) or absence (open arrowhead) of MCHR1 immunoreactivity at the primary cilium of VTA cells. Number of MCHR1-immunoreactive cells throughout the VTA of male and female wildtype mice (**D**). VTA levels were assigned with reference to the *Allen Reference Atlas* (*ARA*; Dong, 2008). Two-way ANOVA with Bonferroni post-test: ****, $p < 0.0001$ (STR vs all regions; **A**). Scale bars: ***i***, 100 µm; ***ii***, 40 µm. IF, interfascicular nucleus raphe; IPN, interpeduncular nucleus; VTA, ventral tegmental area.

Figure 2. Neurochemical characterization of VTA cells. Number (right *y*-axis, gray line) and proportion of each cell type (left *y*-axis, stacked colored bars) at each *Allen Reference Atlas* level (Dong, 2008) rostrocaudally from VTA slices expressing *Vglut2* mRNA hybridization, *Vgat* mRNA hybridization, and tyrosine hydroxylase (TH) immunoreactivity (N = 6 mice; **A**). Representative coronal brain maps of VTA cells that are *Vglut2*-, *Vgat*- and/or TH-positive (**B**). Each panel includes the nomenclature, atlas level and Bregma value (β) assigned to that tissue in accordance with the *Allen Reference Atlas* (Dong, 2008).

Figure 3. *Mchr1* expression on diverse VTA cell types. Representative low (**A**) and high magnification confocal photomicrographs (**B**–**E**), from corresponding dash-outlined region in **A**)

of *Vglut2* mRNA, *Vgat* mRNA, and/or TH immunoreactivity (***i***) that coexpress (white

arrowhead) or lack (open arrowhead) *Mchr1* mRNA hybridization in VTA cells (***ii***). Relative

proportion of *Mchr1*-expressing VTA cells overall (**F**) and at each level, as determined with

reference to the *Allen Reference Atlas* (Dong, 2008), of the VTA (N = 6; **G**). Scale bars: **A**, 250

µm; **B**–**E**, 30 µm. IF, interfascicular nucleus raphe; IPN, interpeduncular nucleus; ml, medial

lemniscus; RN red nucleus; SNc, substantia nigra, compact part; SNr, substantia nigra, reticular

part; VTA, ventral tegmental area.

Figure 4. Spatial distribution of *Mchr1* mRNA in heterogeneous VTA cells. *Mchr1* mRNA

hybridization expression in *Vglut2* (**A**; **Figure 4-1**), *Vgat* (**B**; **Figure 4-2**)*,* TH (**C**; **Figure 4-3**),

or dual *Vglut2* and *Vgat* VTA cells (**D**; **Figure 4-4**) mapped to a representative anterior (***i***),

middle (***ii***), and posterior (***iii***) VTA section using *Allen Reference Atlas* (Dong, 2008) templates

at the atlas level or Bregma value (β) indicated (*top*).

Figure 5. MCH hyperpolarized GABAergic and dopaminergic VTA cells. Representative

epifluorescence image of coronal VTA brain slices (***i***) where cells expressing native EGFP

fluorescence (inset in ***i***) from *Vglut2-cre;L10-Egfp* (**A**), *Vgat-cre;L10-Egfp* (**B**), and *Th-cre;L10-*

Egfp reporter mice (**C**) guided whole-cell patch-clamp recordings to determine the change in

resting membrane potential (Δ RMP) over time following the bath application of 3 µM MCH (***ii***)

and proportion of MCH-responsive cells (donut plot in ***ii***). Scale bars: ***i***, 250 µm; *inset* in ***i***, 20

µm. ml, medial lemniscus; SNc, substantia nigra, compact part; SNr, substantia nigra, reticular

part; VTA, ventral tegmental area.

Figure 6. Selectivity of TH and MCHR1 coexpression in *Th*^EGFP VTA cells. Merged confocal

photomicrograph (***i***) of native EGFP fluorescence (***ii***) and TH immunoreactivity (***iii***) in biocytin-

filled (*iv*) cells from a subset of patch-clamp recordings from the VTA of *Th-cre;L10-Egfp*

reporter mice (**A**). Merged confocal photomicrograph at low (*i*) and high magnification (*ii*, from

dash-outlined region in *i*) of TH- (*iii*) and MCHR1-immunoreactive signals (*iv*) on Nissl-stained

(NeuroTrace) soma in wildtype VTA (**B**). Representative examples of the presence (white

arrowhead) or absence (open arrowhead) of MCHR1 immunoreactivity at the primary cilium of

VTA TH cells or non-TH-labeled cells (*). Number and proportion (donut plot) of MCHR1-

positive (red) or -negative (pink) TH cells at each VTA level in accordance with the *Allen*

Reference Atlas (*ARA*; Dong, 2008) of male and female wildtype mice (N = 4; **C**). Scale bars: **A**,

30 μm; **B***i*, 200 μm; **B***ii*–*iv*, 25 μm. IF, interfascicular nucleus raphe; IPN, interpeduncular

nucleus; VTA, ventral tegmental area.

Figure 7. Rapid detection of direct MCH-mediated hyperpolarization at GABAergic and

dopaminergic VTA cells. MCH-sensitive VTA cells identified by a localized puff of MCH

applied within 20 μm of the patched cell (**A***i*). Representative sample trace of the membrane

potential change following a puff of MCH (red shaded region) lasting <5 s (**A***ii*). Change in

resting membrane potential (Δ RMP) over time following the first (black circles) and second

localized puff (white circles) of 3 μM MCH or of bath ACSF (gray circles) onto a VTA cell

(**A***iii*). RMP change over time following localized puff of 3 μM MCH in the presence (red

outlined circles) or absence of 500 nM TTX (black circles) at *Vgat*$^{\text{EGFP}}$ (**B**) and *Th*$^{\text{EGFP}}$ VTA

cells (**C**). Scale bars: **A***i*, 25 μm; **A***ii*, 3 mV, 10 s.

Figure 8. **Delayed MCH-mediated increase of excitatory input to *Th*$^{\text{EGFP}}$ VTA cells.**

Representative sample trace of spontaneous inhibitory postsynaptic current (sIPSC) events at

baseline, following 3 μM MCH application, and MCH washout (*left*) from patch-clamp

recordings at *Th*$^{\text{EGFP}}$ cells (*right* schematic; **A**). Time course of the relative fold change in sIPSC

frequency (**B**) and amplitude (**C**) relative to baseline values following bath application of 3 μM

MCH. Representative sample trace of spontaneous excitatory postsynaptic current (sEPSC)

events in ACSF, 3 μM MCH, and in MCH following pretreatment with 10 μM TC-MCH 7c (*left*)

from patch-clamp recordings at Th^{EGFP} cells (*right* schematic; **D**). Time course of the relative (*i*)

and peak (*ii*) fold change in sEPSC frequency (**E**) and amplitude (**F**) relative to baseline values

following bath application of 3 μM MCH (black), bath ACSF only (gray), or MCH after

pretreatment with 10 μM TC-MCH 7c (blue) for at least 10 minutes. One-way ANOVA: *, $p <$

0.05 or two-way ANOVA: †, $p < 0.05$ with Šidák multiple comparisons posttest: ^, $p < 0.05$.

Scale bars: 10 pA, 2.5 s (**A**, **D**).

Figure 9. MCH-mediated disinhibition of local excitatory input to *Th* VTA cells.

Representative sample trace of miniature excitatory postsynaptic current (mEPSC) events in the

presence of 500 nM TTX at baseline, following 3 μM MCH application, and MCH washout (*left*)

from patch-clamp recordings at Th^{EGFP} cells (*right* schematic; **A**). Time course of the relative

fold change in mEPSC frequency (**B**) and amplitude (**C**) relative to baseline values following

bath application of 3 μM MCH in slices pretreated with 500 nM TTX. Representative sample

trace of spontaneous inhibitory postsynaptic current (sIPSC) events at baseline, following 3 μM

MCH application, and MCH washout (*left*) from patch-clamp recordings at $Vglut2^{EGFP}$ cells

(*right* schematic; **D**). Time course of the relative fold change in sIPSC frequency (**E**) and

amplitude (**F**) relative to baseline values following bath application of 3 μM MCH. Model of

bidirectional, MCH-mediated regulation of dopamine release from the VTA to nucleus

accumbens (NAc) via the mesolimbic pathway (**G**). Schematic showing inferred MCHR1

distribution within the VTA to elicit MCH-mediated direct inhibition of Th^{EGFP} VTA cells (*i*) and

delayed, indirect GABA-mediated disinhibition of glutamatergic input to Th^{EGFP} VTA cells (*ii*).

Putative MCHR1 expression (*****) along efferent projections of VTA *Vglut2*^{EGFP} cells, which were

not MCH-responsive. Scale bars: 5 pA, 2.5 s (**A**); 10 pA, 2.5 s (**D**).

Extended Data

Figure 3-1. Relative distribution of *Mchr1*-expressing VTA cells in male and female mice

(**A**). Sex may interact with *Vgat* expression to shift the distribution of *Mchr1* mRNA (**B**).

Figure 4-1. Distribution of *Mchr1*-expressing *Vglut2* VTA cells. Number of *Mchr1*-positive

(dark green) or -negative (light green) *Vglut2* cells relative to all VTA cells counted (gray) at

each level in accordance with the *Allen Reference Atlas* (*ARA*; Dong, 2008) (N = 6; **A**). Presence

(dark green circles) or absence (light green circles) of *Mchr1* mRNA hybridization in *Vglut2*

cells throughout the anteroposterior extent of the VTA (**B**–**L**) mapped to *ARA*. Each panel

includes the nomenclature, atlas level and Bregma value (β) assigned to that tissue in accordance

with the *ARA*.

Figure 4-2. Distribution of *Mchr1*-expressing *Vgat* VTA cells. Number of *Mchr1*-positive

(dark yellow) or -negative (light yellow) *Vgat* cells relative to all VTA cells counted (gray) at

each level in accordance with the *Allen Reference Atlas* (*ARA*; Dong, 2008) (N = 6; **A**). Presence

(dark yellow circles) or absence (light yellow circles) of *Mchr1* mRNA hybridization in *Vgat*

cells throughout the anteroposterior extent of the VTA (**B**–**L**) mapped to *ARA*. Each panel

includes the nomenclature, atlas level and Bregma value (β) assigned to that tissue in accordance

with the *ARA*.

Figure 4-3. Distribution of *Mchr1*-expressing TH VTA cells. Number of *Mchr1*-positive (red)

or -negative (pink) TH-immunoreactive cells relative to all VTA cells counted (gray) at each

level in accordance with the *Allen Reference Atlas* (*ARA*; Dong, 2008) (N = 6; **A**). Presence (red

circles) or absence (pink circles) of *Mchr1* mRNA hybridization in TH cells throughout the

anteroposterior extent of the VTA (**B**–**L**) mapped to *ARA*. Each panel includes the nomenclature,

atlas level and Bregma value (β) assigned to that tissue in accordance with the *ARA*.

Figure 4-4. Distribution of *Mchr1* mRNA expression *Vgat*- and *Vglut2*-positive VTA cells.

Number of *Mchr1*-positive (dark blue) or -negative (light blue) cells expressing both *Vglut2* and

Vgat mRNA relative to all VTA cells counted (gray) at each level in accordance with the *Allen*

Reference Atlas (*ARA*; Dong, 2008) (N = 6; **A**). Presence (dark blue circles) or absence (light

blue circles) of *Mchr1* mRNA hybridization in dual-labeled *Vglut2* and *Vgat* cells throughout the

anteroposterior extent of the VTA (**B**–**L**) mapped to *ARA*. Each panel includes the nomenclature,

atlas level and Bregma value (β) assigned to that tissue in accordance with the *ARA*.

Figure 1

Figure 2

F

Mchr1 VTA cells (%)

- 2 — *Mchr1*/*Vglut2*/TH
- 9 — *Mchr1*/*Vglut2*/*Vgat*
- 25 — *Mchr1*/TH
- 18 — *Mchr1*/*Vgat*
- 21 — *Mchr1*/*Vglut2*
- 25 — *Mchr1* only

G

Mchr1 VTA cells (%)

Allen Reference Atlas level

- *Mchr1*/*Vglut2*/TH
- *Mchr1*/*Vglut2*/*Vgat*
- *Mchr1*/TH
- *Mchr1*/*Vgat*
- *Mchr1*/*Vglut2*
- *Mchr1* only

Figure 7