Title:	Regulation of the mouse ventral tegmental area by melanin-concentrating hormone
Abbreviated title:	MCH-mediated regulation of the VTA
Authors:	C Duncan Spencer, Persephone A Miller, Jesukhogie G Williams-Ikhenoba, Ralitsa G Nikolova, Melissa J Chee
Affiliation:	Department of Neuroscience, Carleton University, Ottawa, ON, K1S 5B6, Canada
Correspondence to:	Melissa Chee
	HS 5309
	1125 Colonel By Drive
	Ottawa, ON, K1S 5B6
	Canada
	melissa.chee@carleton.ca

Number of pages:	45
Number of figures:	6
Number of words	
Abstract:	234
Introduction:	646
Discussion:	1499

Conflicts of Interest: The authors have no conflicts of interest to declare.

Acknowledgements: This study is supported by Natural Sciences and Engineering Research Council of Canada (NSERC) Discovery Grant RGPIN-2017-06272 (MJC), NSERC Canadian Graduate Master's Scholarship and the Ontario QEII-GSST program (CDS), Carleton University's Internship-Carleton University Research Experience for Undergraduate Students (I-CUREUS) (JWI).

Abstract

Melanin-concentrating hormone (MCH) acts via its sole receptor MCHR1 in rodents and is an important regulator of homeostatic behaviors like feeding, sleep, and mood to impact overall energy balance. The loss of MCH signaling by MCH or MCHR1 deletion produces hyperactive mice with increased energy expenditure, and these effects are consistently associated with a hyperdopaminergic state. We recently showed that MCH suppresses dopamine release in the nucleus accumbens, which principally receives dopaminergic projections from the ventral tegmental area (VTA), but the mechanisms underlying MCH-regulated dopamine release are not clearly defined. MCHR1 expression is widespread and includes dopaminergic VTA cells. However, as the VTA is a neurochemically diverse structure, we assessed *Mchr1* gene expression at glutamatergic, GABAergic, and dopaminergic VTA cells and determined if MCH inhibited the activity of VTA cells and/or their local microcircuit. *Mchr1* expression was robust in major VTA cell types, including most dopaminergic (78%) or glutamatergic cells (52%) and some GABAergic cells (38%). Interestingly, MCH directly inhibited dopaminergic and GABAergic cells but did not regulate the activity of glutamatergic cells. Rather, MCH produced a delayed increase in excitatory input to dopamine cells and a corresponding decrease in GABAergic input to glutamatergic VTA cells. Our findings suggested that MCH may acutely suppress dopamine release while disinhibiting local glutamatergic signaling to restore dopamine levels. This indicated that the VTA is a target of MCH action, which may provide bidirectional regulation of energy balance.

Significance Statement

Role of melanin-concentrating hormone (MCH) on energy balance may converge on the dopamine system via the mesolimbic pathway, as loss of MCH or MCH receptor (MCHR1) signaling increases hyperactivity and energy expenditure associated with a hyperdopaminergic state. MCH can suppress dopamine release within the mesolimbic pathway, but its underlying mechanism is not known. We thus determined if MCH could inhibit dopamine release through direct actions within the ventral tegmental area (VTA). We found that MCH directly inhibited dopaminergic VTA cells, but MCH also disinhibited excitatory input to dopamine cells. Therefore, we showed that the VTA is a putative target site supporting dopamine-dependent actions of MCH.

Introduction

Melanin-concentrating hormone (MCH) is produced largely in the lateral hypothalamic zone to promote positive energy balance by acting through its receptor MCHR1. MCHR1 is a G-protein coupled receptor (Chambers et al., 1999; Lembo et al., 1999; Saito et al., 1999) and typically inhibits cells via $G_{i/o}$ activation (Gao & Pol, 2001). Mice overexpressing *Mch* may overeat and develop diet-induced obesity (Ludwig et al., 2001), while *Mch* knockout mice (*Mch-KO*) are lean, hyperactive, and have an increased metabolic rate (Kokkotou et al., 2005; Shimada et al., 1998). Similarly, *Mchr1* deletion also results in leanness, hyperactivity, and increased energy expenditure (Åstrand et al., 2004; Chen et al., 2002; Marsh et al., 2002). Although MCH administration stimulates feeding (Qu et al., 1996; Sears et al., 2010), transgenic models of *Mch* or *Mchr1* deletion do not similarly effect feeding (Lord et al., 2022). By contrast, hyperactivity and increased energy expenditure are consistent features following the transgenic *Mch* or *Mchr1* deletion.

A hyperdopaminergic state underlies the hyperactivity of *Mch-* or *Mchr1-KO* animals. There is an elevated amount of dopamine in the accumbens nucleus of *Mch-KO* mice (Pissios et al., 2008) and rats (Mul et al., 2011). Furthermore, treating *Mch-KO* mice with a dopamine reuptake inhibitor leads to greater accumulation of dopamine at the accumbens, and this can enhance hyperlocomotor activity (Pissios et al., 2008). Consistently, MCH can suppress dopamine release in the accumbens, and the inhibitory actions of MCH are mediated in part by *Mchr1* expression at GABAergic accumbens neurons (Chee et al., 2019).

Dopaminergic afferents in the accumbens originate from the ventral tegmental area (VTA) (Albanese & Minciacchi, 1983) and form the mesolimbic pathway implicated in the

regulation of reward and motivation. The VTA is a heterogeneous area comprising

dopaminergic, GABAergic, and glutamatergic cells (Morales & Margolis, 2017), as well as cells

that are both GABAergic and glutamatergic (Root, Mejias-Aponte, Qi, et al., 2014; Yamaguchi

et al., 2007), dopaminergic and GABAergic (Stamatakis et al., 2013; Tritsch et al., 2012), or

dopaminergic and glutamatergic (Dal Bo et al., 2004; Stuber et al., 2010; Sulzer et al., 1998;

Tecuapetla et al., 2010). Interestingly, dopaminergic VTA neurons receive inputs from local

GABA and glutamate cells (Morales & Margolis, 2017), but other brain regions, such as the

lateral hypothalamic area, projecting to the VTA may also regulate dopaminergic VTA neurons

(Nieh et al., 2016).

MCH neurons may directly innervate multiple VTA cell types, including GABAergic,

glutamatergic, and dopaminergic cells (Faget et al., 2016). Optogenetic activation of MCH

neurons can increase dopamine release in the accumbens and increase sucrose preference over a

non-nutritive sweetener (Domingos et al., 2013). However, *Mch-KO* mice do not show deficits in

nutrient-sensing because the nutritive value associated with sucrose is ascribed to glutamate

release from MCH neurons (Schneeberger et al., 2018). Nevertheless, this does not preclude

direct MCH actions on VTA neurons that can inhibit dopamine release.

The VTA comprises MCH-immunoreactive fibers (Bittencourt et al., 1992) and *Mchr1*

mRNA (Hervieu et al., 2000), but the identity of MCHR1-expressing VTA cells and

mechanism(s) of MCH action in the VTA have not been defined. We determined *Mchr1* mRNA

expression on GABAergic, glutamatergic, or dopaminergic VTA cells identified by the vesicular

GABA transporter (*Vgat*), vesicular glutamate transporter 2 (*Vglut2*), or tyrosine hydroxylase

(TH), respectively. We found that MCH directly hyperpolarized dopaminergic or GABAergic

VTA cells but not glutamatergic cells. As VTA dopamine cells receive both GABAergic and

glutamatergic afferents (Marisela Morales & Margolis, 2017), we also determined if MCH may

regulate GABAergic or glutamatergic synaptic transmission at dopaminergic VTA cells.

Interestingly, MCH did not alter GABAergic transmission at dopaminergic VTA cells, but it

produced a delayed increase in glutamatergic input. This may reflect a disinhibition of

glutamatergic tone onto dopamine cells in the VTA, as MCH can suppress GABAergic

transmission at glutamate cells. These findings supported both direct and indirect MCH action

that regulates neuronal activity or synaptic transmission at dopaminergic VTA cells.

Materials and Methods

110 All animal procedures were approved by the Animal Care Committee at Carleton

111 University. All mice were group housed in a 12:12 hour light:dark cycle with *ad libitum* access

112 to water and standard mouse chow (Teklad Global Diets 2014, Envigo, Mississauga, Canada).

Quantitative real-time PCR

WT mice 6–20 weeks old (N = 16) were anesthetized with an intraperitoneal injection

(ip) of 7% chloral hydrate (700 mg/kg). The brain was rapidly removed from the skull and

samples of the striatum, hypothalamus, hippocampus, cerebellum, and VTA were microdissected

and flash-frozen in liquid nitrogen. Tissue samples were homogenized in TRIzol (Invitrogen,

Carlsbad, CA), and RNA was isolated by chloroform and precipitated by isopropanol. The RNA

pellet was washed with 70% ethanol and air-dried before resuspending in distilled water. RNA

purity was determined by a 260/280 absorbance ratio of ~2.0 using a NanoDrop Lite

Spectrophotometer (Thermo Scientific, Waltham, MA).

RNA samples were treated with DNase and reverse transcribed for cDNA synthesis with

an iScript gDNA Clear cDNA Synthesis Kit (1725035, Bio-Rad, Hercules, CA). We performed

quantitative RT-PCR using a CFX Connect Real-Time PCR Detection System thermal cycler

(1855201, Bio-Rad). Custom primers (Invitrogen) for mouse glyceraldehyde-3-phosphate

dehydrogenase (*Gapdh*) and *Mchr1* mRNA comprised the following sequences: *Gapdh* forward,

5'- ATGTGTCCGTCGTGGATCTGA-3'; *Gapdh* reverse, 5'-

ATGCCTGCTTCACCACCTTCTT-3'; *Mchr1* forward, 5'- CAATGCCAGCAACATCTCC-3';

Mchr1 reverse, 5'- ACCAAACACTGAAGGCATGA-3'.

Mchr1 mRNA expression in each brain region was calculated via double delta Ct analysis

and standardized to the expression levels of *Gapdh* of the same mouse. Relative *Mchr1* mRNA

expression in each brain region was then normalized to the level of *Mchr1* mRNA in the

hypothalamus for each sex.

Histology

Tissue preparation. Mice (12–20 weeks old) were anaesthetized with 7% chloral

hydrate (700 mg/kg, ip) and underwent transcardiac perfusion with cold (4 °C) sterile saline

(0.9% NaCl) followed by pre-chilled (4°C) phosphate buffered formalin (10%; VWR, Radnor,

PA).

Immunohistochemistry (IHC). Brain tissue from WT (N = 6) and *Mchr1-KO* mice (N

= 2) were post-fixed in 20% sucrose prepared in formalin (4 h) and then cryoprotected with 20%

sucrose prepared in phosphate buffered saline (PBS, pH 7.4) for 16 h until sectioning using a

freezing microtome (Spencer Lens Co., Buffalo, NY) into five series of 30 μm coronal sections.

MCHR1 immunoreactivity was labeled as previously described (Diniz et al., 2020). In brief,

brain tissue was treated with 0.3% hydrogen peroxide and incubated with a rabbit anti-MCHR1

antibody (1:1,000; PA5-24182, Invitrogen, Carlsbad, CA) for 48 h at 4 ºC in a 3% normal

donkey serum (NDS) solution. Tissue was thoroughly rinsed, incubated with a biotinylated

donkey anti-rabbit IgG antibody (1:500; 711-065-152, Lot 101909, Jackson ImmunoResearch

Laboratories) at RT for 1 h, and treated with avidin-biotin-horseradish peroxidase (1:1:833; PK-

6100, Vector Labratories, Burlingame, CA) at RT for 30 min before incubating with 0.5%

biotinylated (EZ-Link sulfo-NHS-LC-biotin; ThermoFisher Scientific, Waltham, MA) tyramine

(Sigma-Aldrich; Sigma Chemical; St. Louis, MO, USA) for 20 min (RT). All sections were then

incubated with an Alexa Fluor 647-conjugated streptavidin (1:500; 016-600-084, Lot 135095,

Jackson ImmunoResearch Laboratories) and NeuroTrace (1:50; N21479, ThermoFisher) for 20

min (RT). Following staining, all tissue was thoroughly rinsed and immediately mounted onto

Superfrost Plus microscope glass slides (ThermoFisher) and coverslipped with Prolong Diamond

Antifade mounting media (P36961, Invitrogen).

Combined *in situ* hybridization (ISH) and IHC. Brains from WT mice (N = 6) were

post-fixed in 10% formalin overnight, cryoprotected in 20% sucrose with 0.5% sodium-azide (24

h), and sectioned using a freezing microtome into five series of 30 μm coronal sections. The

series designated for ISH were immediately mounted onto Superfrost Plus microscope slides, air-

dried at room temperature (RT; 21–23 °C; 1 h) and −20 °C (30 min), and then stored at −80 °C.

Adjacent brain series were reserved for Nissl staining, RNAscope probe labeling, and negative

control, respectively. One slide from a separate series served as the positive control. We

determined the coexpression of *Mchr1* with *Slc32a1* (*Vgat*) and *Slc17a6* (*Vglut2*) by RNAscope-

enabled ISH and with TH by subsequent IHC in the same brain tissue. ISH procedures were

completed based on manufacturer instructions for RNAscope ISH on fixed-frozen mouse brain

tissue (Multiplex Fluorescent Reagent Kit v2 User Manual; Advanced Cell Diagnostics (ACD),

Newark, CA) as previously described (Bono et al., 2022), unless indicated otherwise.

In brief, slide-mounted brain tissues were dehydrated in an ethanol gradient, treated with

hydrogen peroxide and target retrieval procedure for 5 min (99 °C), and then dehydrated and air-

dried overnight (RT). Tissues of interest were isolated by drawing a hydrophobic barrier and

treated with Protease III in a HybEZ oven (40 °C, 30 min). Slides were then incubated with a

cocktail of *Mm-Mchr1* (317491, ACD) in channel 1 (C1), *Mm-Slc32a1-C2* in C2 (319191-C2,

ACD), and *Mm-Slc17a6* in C3 (319171-C3, ACD) probes (40 °C, 2 h). Concentrated *Mm-*

Slc32a1 C2 and *Mm-Slc17a6* C3 probes were added to the diluted *Mm-Mchr1* C1 probe at a ratio

of 1:50. A volume of 5–8 drops of the probe mixture was added to each slide. *Bacilus*

dihydrodipicolinate reductase (*dapB*, 320871, ACD) and *Mm-Ppib* (313911, ACD) probes were

used as the negative and positive control, respectively. Hybridization signals were amplified by

sequential incubation at 40 °C with AMP-1 (30 min; 323101, ACD), AMP-2 (30 min; 323102,

ACD), and AMP-3 (15 min; 323103, ACD).

C1 *Mchr1* hybridization was developed (40 °C, 15 min with HRP-C1; 323104, ACD) and

tyramide signal amplification (TSA) plus Opal 690 (1:750, 40 °C, 30 min; FP1497001KT,

Akoya Biosciences, Marlborough, MA), then quenched with HRP Blocker (40 °C, 15 min;

323107, ACD). C2 *Vgat* and C3 *Vglut2* hybridization were sequentially developed with HRP-C2

(323105, ACD) with TSA plus Cyanine 3 (1:1,500; NEL44E001KT, PerkinElmer, Waltham,

MA) and HRP-C3 (323106, ACD) with TSA plus Opal 520 (1:750; FP1487001KT, Akoya

Biosciences). C2 signal was similarly quenched with HRP Blocker (40 °C, 15 min) before

proceeding to C3 development.

Following ISH of probe tissue, the tissues were treated for IHC labeling of TH

immunoreactivity based on ACD directions for combined ISH and IHC (Advanced Cell

Diagnostics, 2020) and as previously described (Mickelsen et al., 2019). Tissues were blocked

with 3% NDS solution prepared in PBS (700 μl) for 30 min (RT) and then incubated with anti-

sheep TH primary antibody (1:1,000; GTX82570, Lot 822005648, GeneTex, Irvine, CA) 2 h

(RT). Slides were thoroughly rinsed in PBS and then incubated with a donkey anti-sheep

secondary antibody conjugated with Alexa Fluor 594 (1:500; 715-496-150, Lot 82079,

JacksonImmuno Research) for 30 min (RT). Slides were rinsed in Wash Buffer (320058, ACD),

treated with DAPI Reagent (323108, ACD) for 30 s (RT), and immediately coverslipped with

ProLong Gold Antifade Mountant (Fisher Scientific). Slides were kept in a dark drying drawer at

RT overnight and then transferred to −20 °C for storage until imaging.

Nissl stain. Sections underwent Nissl staining as previously described (Bono et al., 2022;

Negishi et al., 2020).

Microscopy

Brightfield imaging. Nissl-stained tissue was acquired with a Nikon Eclipse Ti2 inverted

microscope (Nilon Instruments Inc., Mississauga, Canada) equipped with a motorized stage and

DS-Ri2 color camera (Nikon) using a Plan Apochromat 10x objective (0.45 NA). Brightfield

images were stitched at 20% overlap with NIS Elements software.

Confocal imaging. Images from Nissl-stained tissue were aligned to tiled, overview

confocal images showing the VTA in relation to other midbrain regions acquired using the 405-

nm wavelength laser with a 4x Plan Apochromat objective (0.2 NA). Stitched, high resolution

confocal frames from IHC- and/or ISH-treated tissues were captured from within the overview

image.

Confocal imaging of MCHR1 immunoreactivity underwent scans with a 10x and/or 20x

(0.75 NA) Plan Apochromat objectives using a Nikon C2 confocal system. Images were acquired

with a 405- and 647-nm wavelength laser to visualize NeuroTrace- and Alexa Fluor 647-labeled

signals.

Confocal imaging of combined ISH- and IHC-treated tissue underwent scans with 20x

Plan Apochromat objective using a Nikon C2 confocal system and acquired with a 405-, 488-,

and 647-nm wavelength laser to visualize DAPI-, Opal 520-, and Alexa Fluor 647-labeled

signals for nuclear stains, *Vglut2* hybridization, and *Mchr1* hybridization, respectively.

Simultaneously at the same optical section, Cyanine 3- and Alexa Fluor 594-labeled signals to

visualize *Vgat* hybridization and TH were acquired by spectral imaging using a 561-nm

wavelength excitation laser in the continuous band (CB) setting to generate a lambda stack

acquired with sequential 10-nm bandwidths spanning the wavelength range of 580 nm to 640

nm. The spectral profiles were unmixed to separate Cyanine 3 and Alexa Fluor 594 signals,

which were most distinguished at 590 nm and 620 nm, respectively, and then merged with

signals acquired by the 405-, 488-, and 647-nm wavelength laser to generate one image file

displaying signals from all five fluorophores.

Tissues that served as the negative control were also acquired using the same

combination of conventional confocal and spectral unmixing for probe tissue. Background

fluorescence related to nonspecific binding from ISH treatment was determined using the

negative control tissue by adjusting the look up table (LUT) values for each of the three image

channels to yield a resultant black image. The LUT range was then applied to images acquired

from the probe tissue to subtract background fluorescence, thus any remaining signal observed

from the probe tissue was considered to be positive hybridization signals.

Neuroanatomical analyses

Plane-of-section analysis. Nissl or fluorescent-Nissl images were parcellated in Adobe

Illustrator CS6 (Adobe Inc., San Jose, CA) based on the regions defined by the *Allen Reference*

Atlas (*ARA*; Dong, 2008), as previously described (Bono et al., 2022; Negishi et al., 2020).

Parcellations for MCHR1-ir cells were directly applied from fluorescent-Nissl signals captured

within the same image acquisition. High magnification confocal images of ISH-treated tissues

were indirectly parcellated from brightfield images of an adjacent Nissl-stained section, which

was aligned to the low magnification, overview confocal image of the probe tissue.

Identifying MCHR1-ir signals. Our MCHR1 antibody (1:1,000; PA5-24182, Invitrogen,

Carlsbad, CA) labeled MCHR1 immunoreactivity at the primary cilium of neurons (Diniz et al.,

2020). Only cilia associated with a NeuroTrace-labeled somata was counted as a MCHR1-

expressing cell. Similarly, only TH-ir signals that colocalized to a DAPI-labeled cell was

considered. Only signals within the boundaries of the VTA, as determined by our plane-of-

section analysis, were included in our analyses.

Identifying hybridization signals. Signal criteria was independently-established based

on the signal-to-background level for each hybridization signal. *Mchr1* hybridization signal

appeared as punctate "dots" with clusters centered tightly around a DAPI-labelled nucleus (Bono

et al., 2022). *Vgat* hybridization appeared as dot clusters that took the shape of cell bodies and

were colocalized to a DAPI-labelled nucleus. Compared to *Vgat* hybridization, dot clusters

following *Vglut2* hybridization was less dense within the cell and may appear as punctate dots.

Mapping. A circle was placed over each positively-labeled immunoreactive or

hybridized cell, and the relative positions of the circles were transferred, as previously described

(Bono et al., 2022; Negishi et al., 2020), to coronal brain atlas templates (Dong, 2008) to

generate standardized maps of *Mchr1*, *Vglut2*, *Vgat*, and TH expression. Cell counts at each atlas

level were determined from cells mapped and counted unilaterally.

Electrophysiology

Animals. *Th-cre;L10-Egfp*, *Vgat-cre ;L10-Egfp*, and *Vglut2-cre;L10-Egfp* reporter mice

were generated by crossing *Th-cre* (Lindeberg et al., 2004), *Vgat-cre* (028862, JAX; Vong et al.,

2011), or *Vglut2-cre* (028863, JAX; Vong et al., 2011) mice, respectively to homozygous *lox-*

STOP-lox-L10-Egfp reporter mice (Krashes et al., 2014). All transgenic lines were kindly

provided by Dr. B. B. Lowell (Beth Israel Deaconess Medical Center, Boston, MA).

Acute slice preparation. Mice (4–10 weeks old) were anaesthetized with 7% chloral

hydrate (700 mg/kg, ip) and transcardially perfused with ice cold, carbogenated (95% O_2, 5%

CO_2) slice artificial cerebrospinal fluid (ACSF) solution (300 mOsm/L; pH 7.4) containing (in

mM) 92 NMDG, 2.5 KCl, 1.25 NaH_2PO_4, 30 $NaHCO_3$, 20 HEPES, 25 glucose, 2 thiourea, 5 Na-

ascorbate, 3 Na-pyruvate, 0.5 $CaCl_2 \cdot 2H_2O$, and 10 $MgSO_4$ (Ting et al., 2018). Brains were

sectioned on a vibratome (VT 1000S, Leica Biosystems, Wetzlar, Germany) in slice ACSF to

produce 250 μm-thick slices that were immediately transferred to carbogenated slice solution at

37 °C for 5 min. Slices were then incubated in a carbogenated bath ACSF (300 mOsm/L)

containing (in mM) 124 NaCl, 3 KCl, 1.3 $MgSO_4$, 1.4 NaH_2PO_4, 10 D-glucose, 26 $NaHCO_3$, and

2.5 CaCl$_2$ for 5 min and then allowed to recover in bath ACSF (RT) for at least 1 hr prior to

recordings.

Patch-clamp recordings. Brain slices containing the VTA were placed in a slice

chamber, superfused (2–2.5 ml/min) with carbogenated bath ACSF pre-warmed (37 ^{o}C) by a

temperature controller (Warner Instrument Corporation, Hamden, CT), and held in place by a

platinum ring strung with nylon fiber made in-house.

VTA cells were visualized with infrared differential interference contrast microscopy at

40x magnification on either an Examiner.A1 microscope (Zeiss, Oberkochen, Germany)

equipped with an AxioCam camera (Zeiss) and Axiovision software (Zeiss) or with an Eclipse

FN1 microscope (Nikon) equipped with a pco.panda 4.2 camera (Excelitas PCO GmbH,

Kelheim, Germany) and NIS-Elements Imaging software (Nikon). EGFP-labeled cells were

identified using epifluorescence illumination from a halogen (HXP120, Zeiss) or mercury lamp

(C-SHG1, Nikon).

Borosilicate glass pipettes (593800; A-M Systems) were pulled on a P-1000

Flaming/Brown micropipette puller (Sutter Instruments; Novato, CA) and backfilled (7–10 MΩ)

with a potassium-based internal solution (290 mOsm/L, pH 7.24) containing (in mM) 120 K-

gluconate, 10 KCl, 10 HEPES, 1 MgCl$_2$, 1 EGTA, 4 MgATP, 0.5 NaGTP, and 10

phosphocreatine to assess membrane properties and glutamatergic events. A cesium-based

internal solution (290 mOsm/L, pH 7.24) containing (in mM) 128 CsMS, 11 KCl, 10 HEPES,

0.1 CaCl2, 1 EGTA, 4 MgATP, 0.5 NaGTP) was used to record GABAergic events. Voltage-

clamp recordings were acquired at a holding potential of $V_h = -60$ mV and -5 mV for excitatory

and inhibitory events, respectively using a MultiClamp 700B amplifier (Molecular Devices, San

Jose, CA) and digitized using a Digidata 1440A (Molecular Devices). Traces were acquired and

filtered at 1 kHz by pClamp 10.3 or 11 software (Molecular Devices).

Drug application. Pharmacological applications of MCH (H-1482, Bachem, Torrance,

CA) was applied by bath perfusion or puff application after a baseline period of at least 5 min or

1 min, respectively, where the membrane potential fluctuated by less than 10%. Cells were held

for at least 20 min during the MCH-washout period to assess the reversibility of MCH-mediated

effects on the membrane potential. Where applicable, tetrodotoxin citrate (TTX; T-550, Alomone

labs, Jerusalem, Israel) was bath applied and allowed to saturate the slice for at least 10 min prior

to MCH application and then maintained during the MCH-washout period as well.

Puff-applied MCH was delivered via a puff-pipette backfilled with 3 μM MCH (or bath ACSF)

positioned within 20–40 μm away from the patched cell held in the whole-cell configuration. A

short "puff" was applied by manually applying positive pressure via the puff-pipette, and a

solution ripple can be seen to contact the patched cell.

Experimental design and statistical analyses

qPCR studies. Comparison of *Mchr1* gene expression in wildtype mice (N = 6–7/sex)

were analyzed using a between-subject design. *Mchr1* expression in different regions were

analyzed using a two-way mixed model ANOVA with Šidák post-hoc testing to determine sex

differences.

Neuroanatomical studies. Comparisons of the total number of MCHR1-expressing cells

were analyzed using a between-subject design from (N = 3/sex) and compared via unpaired *t* test,

and differences in the distribution of MCHR1-expressing cells across each *ARA* level between

sexes was determined by the two-way mixed model ANOVA.

Comparisons of *Mchr1*-, *Vglut2*-, *Vgat*-, or TH-expressing cells were analyzed from male

and female wildtype mice (N = 6) using a between-subject design, and a one-way ANOVA was

used to identify distinct VTA cell groups. Sex differences were determined by two-way

ANOVA.

Electrophysiological studies. Acute brain slices were prepared from male and female

Th-cre;L10-Egfp (N = 23), *Vgat-cre;L10-Egfp* (N = 8), and *Vglut2-cre;L10-Egfp* (N = 6) reporter

mice. Up to three cells were recorded from three to four slices, where the VTA sits between the

medial mammillary nucleus and compact part of the substantia nigra in rostral sections and sits

beside the interpeduncular nucleus in caudal sections. Electrical signals recorded were processed

in Clampfit 10.7 (Molecular Devices).

The resting membrane potential (RMP) was sampled every 1 s and averaged into 15-s

bins in bath-applied treatments and sampled every 100 ms and averaged into 1-s bins in puff-

applied treatments. MCH-mediated change in RMP (Δ RMP) was sampled within 7 min after

MCH application or within 5 s of puff application. Comparisons of MCH-mediated Δ RMP over

time in the presence or absence of TTX was determined using within-subject design and repeated

measures two-way ANOVA with Šidák post-hoc testing, and comparisons of mean Δ RMP were

determined by paired *t* test.

The frequency of spontaneous excitatory postsynaptic current (sEPSC) events were

indirectly determined by the interevent interval between sEPSCs using MiniAnalysis

(Synaptosoft, Decatur, GA) and averaged into 15-s bins. The baseline sEPSC frequency was

determined as the mean sEPSC frequency averaged over 5 min immediately prior to MCH bath

application.

Graphs. All data graphs were prepared using Prism 9 (GraphPad, San Diego, CA).

Results were considered statistically significant at $p < 0.05$. All data are reported as group mean

$\pm$ SEM. Figures were assembled in Adobe Illustrator 2020 (Adobe Inc.). Numeral in parentheses

in graphs represent the number of animals (N) or cells (n).

Results

Moderate *Mchr1* mRNA and MCHR1 protein expression in the mouse VTA

Mchr1 mRNA and protein have been reported in the rat VTA (Hervieu et al., 2000; Saito

et al., 2001), and we first determined *Mchr1* mRNA and/or protein expression in the VTA of

male and female wildtype mice. While the level of *Mchr1* mRNA varied across brain regions

($F_{(4, 44)} = 81.6$, $p < 0.0001$, ANOVA), it was similar between male and female mice ($F_{(1, 11)} =$

0.6, $p = 0.449$, ANOVA), and there was no interaction between brain region and sex ($F_{(4, 44)} =$

0.8, $p = 0.561$, ANOVA). As expected, *Mchr1* expression was highest in the striatum, which

included the nucleus accumbens and caudate putamen, and barely detectable in the cerebellum

(**Figure 1A**). Importantly, *Mchr1* expression in the VTA was comparable to the hypothalamus

(male: $t_{(44)} = 1.5$, $p = 1.000$; female: $t_{(44)} = 1.0$, $p = 1.000$) and hippocampus (male: $t_{(44)} = 1.3$,

$p = 1.000$; female: $t_{(44)} = 0.6$, $p = 1.000$; **Figure 1A**).

The VTA of wildtype male and female mice also comprised MCHR1-immunoreactive

(MCHR1-ir) signals, which had a long and slender appearance and were consistent with MCHR1

expression observed on the primary cilium of neurons (**Figure 1B**). No MCHR1-ir labeling was

detected in the VTA of *Mchr1-KO* mice (**Figure 1C**). Male and female wildtype mice had a

similar distribution pattern of MCHR1-expressing cells throughout the VTA ($F_{(1, 4)} = 0.02$, $p =$

0.891, ANOVA; **Figure 1D**). Additionally, there was no difference in the total number of

MCHR1-expressing cells in the VTA of male (159 ± 48, n = 3) and female (171 ± 29, n = 3)

mice (t(4) = 0.2, p = 0.835, unpaired *t* test).

Neurochemical diversity of VTA cells

The VTA comprises multiple cell types, including dopaminergic, GABAergic, or

glutamatergic cells that have distinct distribution patterns (M. Morales & Root, 2014; Marisela

Morales & Margolis, 2017). To identify the neurochemical identities of VTA cell types, we

determined if *Mchr1* hybridization colocalized with *Vglut2* or *Vgat* hybridization in

glutamatergic or GABAergic cells, respectively, or with TH immunoreactivity in dopaminergic

cells within the same tissue. As there were no sex differences in *Mchr1* mRNA (**Figure 1A**) or

protein expression (**Figure 1C**), we herein combined our neuroanatomical analyses from male

and female mice.

The overall number of VTA cells labeled by *Vglut2* mRNA, *Vgat* mRNA, and/or TH

immunoreactivity decreased posteriorly (**Figure 2A**, gray line). We compared the distribution of

Vglut2 mRNA, *Vgat* mRNA, and TH-ir, cells within the VTA to determine their relative

prevalence rostrocaudally, and we could distinguish at least five subtypes of VTA cells (F(4, 50)

= 13.4, p = 0.0001, ANOVA). *Vglut2* cells were most prominent anteriorly then gradually

diminished towards the posterior VTA (**Figure 2A**), and they were usually medially distributed

within the VTA (**Figure 2B**). *Vgat* cells comprised ~30% of VTA cells counted through the

anterior and middle sections of the VTA (L80–87) but made up the majority of cells in the

posterior VTA (L88–90) (**Figure 2A**). TH-ir cells were most numerous in the middle of the VTA

and comprised ~40% of cells counted (**Figure 2A**). Both *Vgat* and TH cells tended to be laterally

distributed, though some *Vgat* cells were spread medially (**Figure 2B**).

In addition, we also found that VTA cells may express two or more neurotransmitter

markers. The most common were those expressing both *Vglut2* and *Vgat* that comprised 5–10%

of VTA cells and up to 16% of cells towards the middle of the VTA (L86–87; **Figure 2A**). Cells

expressing TH and *Vglut2* emerged towards the posterior VTA (L85–89) but comprised <6% at

its peak (**Figure 2A**) so were not mapped. Similarly, just 0.25% of cells exhibited TH and *Vgat*,

and cells exhibiting *Vglut2*, *Vgat*, and TH were even rarer (<0.05%; data not shown).

Neurochemical and spatial distribution of *Mchr1* mRNA expression in the VTA

Mchr1 mRNA expression was widely distributed throughout the VTA (**Figure 3A**) and

across all major VTA cell types (**Figure 3B**–**E**), but there was no sex difference in the

heterogeneity of *Mchr1* VTA cells (F(1, 24) = 0.01, p = 0.924, ANOVA; **Figure 3-1A**).

Interestingly, about one in four *Mchr1* cells (25%) do not coexpress TH, *Vglut2*, or *Vgat* and are

yet to be defined neurochemically (**Figure 3F**). These *Mchr1*-only cells can be seen throughout

the VTA (**Figure 3G**) and tend to be less prominent in the VTA of female mice, as sex may

interact with *Vgat* expression in *Mchr1* cells (F(1, 8) = 4.8, p = 0.059, ANOVA; **Figure 3-1B**).

Consistent with the rostrocaudal composition of VTA cells, *Mchr1* expression was

predominantly in *Vglut2* cells rostrally, TH cells in the middle sections of the VTA, and in *Vgat*

cells caudally (**Figure 3G**).

About half (52%) of *Vglut2* VTA cells expressed *Mchr1* mRNA (**Figure 4-1**), which was

evenly distributed throughout *Vglut2* cells in the medial and anterior VTA but only sparsely

distributed posteriorly (**Figure 4A**). By contrast, few *Vgat* cells in the anterior VTA expressed

Mchr1 mRNA (**Figure 4-2**), but in the middle and posterior VTA sections where *Mchr1*-

expressing *Vgat* cells were more common, they tend to be laterally distributed or have no

discernible pattern, respectively (**Figure 4B**). Interestingly, the majority of TH-ir VTA neurons

(78%) expressed *Mchr1* mRNA (**Figure 4-3**), and while *Mchr1*-expressing TH cells were found

sparingly in the anterior VTA, they were abundant in middle VTA sections, especially near the

ventral and lateral borders of the VTA (**Figure 4C**). In addition, although dual *Vglut2* and *Vgat*

cells comprised just one-tenth of all VTA cells, nearly 60% of them expressed *Mchr1* mRNA

(**Figure 4-4**) and were most likely found in the dorsomedial aspects of the VTA (**Figure 4D**).

Taken together, these findings indicated the abundance of *Mchr1* mRNA throughout the VTA in

glutamatergic, GABAergic, and/or dopaminergic VTA cells.

MCH inhibited *Vgat* and *Th* VTA cells but not *Vglut2* VTA cells

Based on the spatial distribution of *Mchr1* expression in *Vglut2*, *Vgat*, and TH VTA cells

(**Figure 4),** we performed whole-cell patch-clamp recordings in EGFP-labeled (**Figure 5A**)

$Vglut2^{EGFP}$, $Vgat^{EGFP}$, and Th^{EGFP} VTA cells from *Vglut2-cre;L10-Egfp*, *Vgat-cre;L10-Egfp*, and

Th-cre;L10-Egfp reporter mice, respectively. MCH (3 μM) bath application had no effect on the

resting membrane potential of $Vglut2^{EGFP}$ VTA cells (–0.3 ± 0.9 mV, n = 0/6 cells, N = 3;

Figure 5B). By contrast, MCH hyperpolarized $Vgat^{EGFP}$ (–4.4 ± 1.3 mV, n = 5/14 cells, N = 5;

Figure 5C) and Th^{EGFP} cells (–4.8 ± 1.3 mV, n = 6/16 cells, N = 12; **Figure 5D**) within 10 min

and persisted for at least 25 min before gradually returning to baseline.

We then assessed if MCH-mediated inhibition of $Vgat^{EGFP}$ and Th^{EGFP} VTA cells was

activity-dependent by pretreatment with the voltage-gated sodium channel blocker tetrodotoxin

(TTX). Since the effect of bath-applied MCH was so long-lasting, we first established if MCH-

sensitive VTA cells could be identified by a short puff application of MCH lasting < 5 seconds to

allow rapid recovery of the resting membrane potential. A single puff of MCH produced an acute

membrane hyperpolarization that peaked within 10 seconds and returned to baseline within one

minute (−6.7 ± 1.8 mV, n = 4; **Figure 5E**). A second puff of MCH applied 3–5 minutes later at

the same cell elicited a similar hyperpolarization timecourse (−4.3 ± 1.7 mV, n = 4; $F(1, 6) = 3.3$,

$p = 0.120$, ANOVA; **Figure 5E*iii***). Importantly, puff application of ACSF produced visibly

different outcomes than the first MCH puff ($F(1, 5) = 6.8$, $p = 0.047$, ANOVA) or second MCH

puff ($F(1, 5) = 30.7$, $p = 0.003$, ANOVA), as it did not effect changes in membrane potential

(−0.2 ± 0.6 mV, n = 3; **Figure 5E*iii***). This thus showed that a localized MCH puff could identify

MCH-sensitive VTA cells by enabling short-lived but reproducible membrane

hyperpolarizations that were not significantly desensitized by repeated application. We then puff-

applied MCH to identify MCH-sensitive *Vgat*[EGFP] and *Th*[EGFP] VTA cells and compared the

effect of MCH before and after TTX (500 nM) treatment. MCH-mediated hyperpolarization at

Vgat[EGFP] VTA cells (−1.8 ± 1.2 mV, n = 4) persisted in the presence of TTX (−1.1 ± 0.4 mV, n

= 4; $F(1, 6) = 0.002$, $p = 0.965$, ANOVA; **Figure 5F**). Similarly, *Th*[EGFP] VTA cells remained

sensitive to MCH (−5.4 ± 1.6 mV, n = 5) in the presence of TTX (−4.3 ± 1.0 mV, n = 5; $F(1, 8) =$

0.3, $p = 0.623$, ANOVA; **Figure 5G**). These findings indicated that MCH directly

hyperpolarized *Vgat*[EGFP] and *Th*[EGFP] VTA cells.

MCH increased glutamatergic input to *Th* VTA cells

Dopaminergic output from the VTA can be regulated by local glutamatergic and

GABAergic circuitry (Bariselli et al., 2016; Morales & Margolis, 2017; Wang et al., 2015), thus

we assessed if MCH regulated the GABAergic and/or glutamatergic input to *Th*[EGFP] VTA cells.

Bath application of MCH did not alter the frequency of sIPSC events arriving at *Th*[EGFP] VTA

cells (**Figure 6A**). Interestingly, MCH elicited a delayed increase in sEPSC frequency

$(1.4 \pm 0.3$ Hz, n = 8) at Th^{EGFP} VTA cells that gradually peaked about 25 minutes following

MCH application (**Figure 6B**), and this delayed increase was MCH-dependent, as control

recordings where ACSF was bath applied in the absence of MCH did not alter sEPSC frequency

$(F(1, 10) = 11.3, p = 0.007$, ANOVA; **Figure 6B**).

To determine the role of upstream targets that facilitate excitatory input to Th^{EGFP} VTA

cells, we isolated presynaptic excitatory input (mEPSC) with TTX pretreatment. MCH applied in

the presence of TTX did not alter mEPSC frequency at Th^{EGFP} VTA cells (**Figure 6C**), thus

suggesting that MCH may recruit upstream targets that indirectly regulate excitatory input at

Th^{EGFP} VTA cells. As MCHR1 is a known inhibitory $G_{i/o}$-protein coupled receptor (Hawes et al.,

2000), we determined the MCH-mediated increase in glutamatergic tone at Th^{EGFP} VTA cells

may be ascribed to GABAergic disinhibition at glutamatergic nerve terminals. We recorded from

$Vglut2^{EGFP}$ VTA cells and found that MCH decreased the frequency of sIPSC events $(-0.7 \pm$

0.3 Hz; **Figure 6D**). Overall, these recordings suggested that, in addition to directly inhibiting Th

VTA cells, MCH may also engage a local disynaptic circuit from $Vgat$ and $Vglut2$ VTA cells to

regulate VTA output (**Figure 6E**).

Discussion

MCH can act within the VTA to provide direct and indirect regulation of Th VTA cells.

MCHR1-expressing cells were widespread throughout the VTA, including at glutamatergic

$Vglut2$, GABAergic $Vgat$, and dopaminergic TH cells, thus providing a neuroanatomical basis

for MCH action in the VTA. Accordingly, electrophysiological recordings showed that MCH

hyperpolarized and inhibited *Th* cells and *Vgat* cells. Interestingly, MCH did not have a direct

effect on *Vglut2* cells, but it disinhibited *Vglut2* afferents and increased excitatory synaptic

transmission at *Th* cells. Therefore, functional MCHR1 activation in the VTA may provide

bidirectional regulation of dopamine output from the VTA.

The majority, nearly 80%, of TH-ir VTA cells expressed *Mchr1* mRNA, and MCH

directly hyperpolarized 38% of *Th* VTA cells in an activity-independent manner. The lower rate

of MCH-responsive *Th-cre* cells may reflect Cre expression in TH-negative VTA cells that may

comprise up to half of all EGFPTh cells (Lammel et al., 2015). Furthermore, although MCHR1

has been reported in TH VTA cells (Diniz et al., 2020), the level of *Mchr1* mRNA required for

protein translation is not known and may be insufficient for MCHR1 expression. MCHR1

signaling has known inhibitory actions (Chambers et al., 1999; Hawes et al., 2000; Lembo et al.,

1999; Saito et al., 1999), including within the mesolimbic system that connects the VTA to the

ventral striatum. In the accumbens, MCH can hyperpolarize medium spiny neurons (Georgescu

et al., 2005) and suppress dopamine release (Chee et al., 2019). Furthermore, loss of MCH

signaling can increase dopamine accumulation in the accumbens leading to hyperactivity and

increased energy expenditure (Chee et al., 2019; Mul et al., 2011; Pissios et al., 2008). As the

main source of dopaminergic afferents to the accumbens arises from the VTA, it stands to reason

that MCH may regulate dopamine release by direct inhibition of VTA neurons.

Interestingly, MCH action within the VTA caused a delayed increase in excitatory input

to *Th* VTA cells. MCHR1 activation can engage $G_{i/o}$- or G_q-protein systems (Pissios et al., 2003;

Sears et al., 2010) to elicit inhibitory or excitatory actions, respectively. Therefore, we

considered that MCH may act presynaptically to stimulate glutamatergic afferents, but the MCH-

mediated increase in glutamatergic transmission was abolished in the presence of TTX. This

suggested that the stimulatory effect of MCH on glutamatergic afferents was indirect, activity-

dependent, and polysynaptic. We thus examined if MCH may indirectly recruit glutamatergic

afferents detected at *Th* VTA cells by suppressing GABAergic input onto *Vglut2* VTA cells or

axons. We found that MCH directly inhibited GABAergic VTA cells and suppressed

GABAergic input to *Vglut2* VTA cells. In effect, MCH can suppress GABAergic activity to

disinhibit glutamatergic transmission at *Th* VTA cells, and we posit that this GABA-dependent

microcircuit within the VTA may serve to restore the output of *Th* VTA cells that were acutely

inhibited by MCH (**Figure 6E**).

VTA *Vglut2* cells are known to innervate (Dobi et al., 2010) and regulate the activity of

dopaminergic VTA cells (Bariselli et al., 2016; Wang et al., 2015). Consistent with previous

reports (Morales & Root, 2014; Root et al., 2018; Yamaguchi et al., 2007), *Vglut2* cells,

including *Mchr1*-expressing *Vglut2* cells were more abundant in the anterior VTA. However,

unlike MCH-mediated inhibition at *Th* cells, MCH had no effect on the membrane potential of

Vglut2 VTA cells, but rather, MCH increased glutamatergic tone at *Th* VTA cells. *Vglut2* VTA

cells form local synapses with dopaminergic and non-dopaminergic VTA neurons (Dobi et al.,

2010) and are pivotal regulators of local VTA circuitry and output. Glutamatergic VTA cells can

drive dopamine release and promote conditioned place preference (Wang et al., 2015) or drug-

induced conditioned place preference (Mead & Stephens, 1999; Slusher et al., 2001; Tzschentke

& Schmidt, 1998) that can be enhanced by food restriction (Stuber et al., 2002) to assess drug

reward (Hiroi & White, 1991). Interestingly, *Mchr1* deletion can block the effect of food

restriction on amphetamine-induced conditioned place preference (Geuzaine et al., 2014).

However, as *Vglut2* cells within the VTA were not MCH-sensitive, it would be important to

consider that MCH may indirectly modulate dopamine output at glutamatergic afferents arising

from outside the VTA, such as from the prefrontal cortex or lateral hypothalamus (Geisler &

Wise, 2008). Furthermore, since we did not detect postsynaptic MCH effects at the soma of

Vglut2 cells, we predicted that MCHR1 distribution within *Vglut2* cells would accumulate at

presynaptic terminals to regulate glutamate release at projection sites (**Figure 6E**, *asterisk*).

523   Prominent *Vgat* expression in the posterior VTA may partly form the tail of the VTA

(tVTA) or rostromedial tegmental nucleus (RMTg; Jhou et al., 2009). The GABAergic cell

cluster in the posterior VTA also reflected the highest proportion of *Vgat* VTA cells that

expressed *Mchr1*. This supports the hypothesis that the posterior VTA is anatomically and

functionally unique (Lahti et al., 2016; Smith et al., 2019) and suggests that this subregion is a

site of MCH action. Sex may interact with *Vgat* expression at *Mchr1* cells in the VTA, as there is

a larger portion of *Mchr1* cells expressing *Vgat* in the VTA of female mice. This suggested that

MCH may contribute to sex-specific regulation of VTA output, potentially via *Vgat* cells in

females or by yet characterized *Mchr1*-only cells in males. The tVTA/RMTg has been implicated

in responsiveness to alcohol as exposure to alcohol stimulates tVTA/RMTg neurons, and rats

will self-administer ethanol to the posterior but not anterior VTA (Guan et al., 2012).

Interestingly, *Mchr1* deletion suppresses alcohol-induced conditioned place preference (Karlsson

et al., 2016), and MCHR1 blockade decreases self-administration of alcohol (Cippitelli et al.,

2010). The posterior VTA supplies dense GABAergic projections to other VTA neurons (Barrot

et al., 2012; Jhou et al., 2009; Lecca et al., 2012) and provides strong inhibitory tone to

dopaminergic VTA neurons (Barrot et al., 2012; Bouarab et al., 2019; Guan et al., 2012). While

MCH did not regulate GABAergic synaptic transmission at *Th* VTA cells, it inhibited

GABAergic transmission at glutamatergic cells. Furthermore, MCH directly inhibited *Vgat* cells,

thus, taken together, MCH action at *Vgat* cells in the posterior VTA may weaken inhibition of *Th*

VTA cells to indirectly facilitate dopaminergic output.

Consistent with findings in rats (Hervieu et al., 2000; Saito et al., 2001), the relative

expression of *Mchr1* mRNA in the VTA was comparable to that in the hypothalamus and

hippocampus. *Mchr1* expression was visible in all major subtypes of VTA cells, including cells

that coexpressed two or more chemical messengers. The *Vgat*/*Vglut2* cells were the most

common subpopulation coexpressing a combination of TH, *Vgat*, or *Vglut2*. The availability of

Vgat/*Vglut2* cells throughout the VTA is consistent with findings demonstrating there is a

population of VTA neurons that corelease GABA and glutamate (Root, Mejias-Aponte, Zhang,

et al., 2014) and express both *Vgat* and *Vglut2* (Root et al., 2018). VTA neurons may also be

both glutamatergic and dopaminergic (Morales & Root, 2014; Morales & Margolis, 2017;

Yamaguchi et al., 2011), but unlike the rat VTA, the incidence of colocalization between TH and

Vglut2 was low. In rats, most dopaminergic and glutamatergic VTA cells were at or very close to

the midline (Yamaguchi et al., 2011) and includes the midline regions between the left and right

VTA hemispheres to encompass the interfascicular nucleus raphe, rostral linear nucleus raphe,

and central linear nucleus raphe as subregions of the VTA (Morales & Root, 2014; Morales &

Margolis, 2017; Yamaguchi et al., 2011). However, our analyses were based on the

neuroanatomical boundaries of the VTA as defined by the *Allen Reference Atlas* (Dong, 2008)

and so did not include the midline regions where dual TH/*Vglut2* cells may be most abundant.

Finally, although VTA cells may co-release dopamine and GABA (Stamatakis et al., 2013;

Tritsch et al., 2012), coexpression of TH and *Vgat* did not form a substantial population within

the VTA. GABA release does not require vesicular transport via vGAT because the vesicular

monoamine transporter 2 (vMAT2) can fully support GABA release from dopaminergic cells

(Tritsch et al., 2012), thus it is not surprising that TH/*Vgat* coexpression was low.

In conclusion, our findings suggest that MCHR1 expression at multiple VTA cell types

can modulate the local VTA microcircuit to provide bidirectional regulation of dopamine output

from the VTA. MCH directly inhibited dopaminergic VTA cells, which may acutely suppress

dopamine release. However, MCH also inhibited GABAergic VTA cells, which may disinhibit

glutamatergic input to dopamine cells. This GABA-mediated disinhibition has a delayed

timecourse and may act to restore dopamine levels acutely lowered by the direct MCH-mediated

inhibition of dopamine cells. Paradoxically, this may be in contrast to prior findings that *Mchr1*

deletion from *Vgat* neurons results in a hyperdopaminergic state (Chee et al., 2019), but this and

previous findings may not be mutually exclusive, as MCH can have site-specific effects at

different nodes of the mesolimbic dopamine pathway. It would be of interest to test our proposed

model (**Figure 6E**) in future studies and determine if *Mchr1* deletion from dopaminergic cells

would also elicit a hyperdopaminergic state, or whether disrupting GABAergic transmission

would abolish the facilitation of glutamatergic input to dopamine cells.

References

Advanced Cell Diagnostics. (2020). RNAscope ® Multiplex Fluorescent v2 Assay combined

with Immunofluorescence - Integrated Co-Detection Workflow (ICW) Combinations.

Advanced Cell Diagnostics Inc., 1–5.

Albanese, A., & Minciacchi, D. (1983). Organization of the ascending projections from the

ventral tegmental area: A multiple fluorescent retrograde tracer study in the rat. *Journal of*

Comparative Neurology, *216*(4), 406–420. https://doi.org/10.1002/cne.902160406

Åstrand, A., Bohlooly, M., Larsdotter, S., Mahlapuu, M., Andersén, H., Tornell, J., Ohlsson, C.,

Snaith, M., & Morgan, D. G. A. (2004). Mice lacking melanin-concentrating hormone

receptor 1 demonstrate increased heart rate associated with altered autonomic activity.

American Journal of Physiology - Regulatory Integrative and Comparative Physiology,

287(4 56-4), 749–758. https://doi.org/10.1152/ajpregu.00134.2004

Bariselli, S., Glangetas, C., Tzanoulinou, S., & Bellone, C. (2016). Ventral tegmental area

subcircuits process rewarding and aversive experiences. *Journal of Neurochemistry*, *139*(6),

1071–1080. https://doi.org/10.1111/jnc.13779

Barrot, M., Sesack, S. R., Georges, F., Pistis, M., Hong, S., & Jhou, T. C. (2012). Braking

dopamine systems: A new GABA master structure for mesolimbic and nigrostriatal

functions. *Journal of Neuroscience*, *32*(41), 14094–14101.

https://doi.org/10.1523/JNEUROSCI.3370-12.2012

Bittencourt, J. C., Presse, F., Arias, C., Peto, C., Vaughan, J., Nahon, J. □L, Vale, W., &

Sawchenko, P. E. (1992). The melanin□concentrating hormone system of the rat brain: An

immuno□ and hybridization histochemical characterization. *Journal of Comparative*

Neurology, *319*(2), 218–245. https://doi.org/10.1002/cne.903190204

Bono, B. S., Koziel Ly, N. K., Miller, P. A., Williams-Ikhenoba, J., Dumiaty, Y., & Chee, M. J. (2022). Spatial distribution of beta-klotho mRNA in the mouse hypothalamus, hippocampal region, subiculum, and amygdala. *Journal of Comparative Neurology*, *530*(10), 1634–1657. https://doi.org/10.1002/cne.25306

Bouarab, C., Thompson, B., & Polter, A. M. (2019). VTA GABA Neurons at the Interface of Stress and Reward. *Frontiers in Neural Circuits*, *13*(December), 1–12. https://doi.org/10.3389/fncir.2019.00078

Chambers, J., Ames, R. S., Bergsma, D., Muir, A., Fitzgerald, L. R., Hervieu, G., Dytko, G. M., Foley, J. J., Martin, J., Liu, W. S., Park, J., Ellis, C., Ganguly, S., Konchar, S., Cluderay, J., Leslie, R., Wilson, S., & Sarau, H. M. (1999). Melanin-concentrating hormone is the cognate ligand for the orphan G- protein-coupled receptor SLC-1. *Nature*, *400*(6741), 261–265. https://doi.org/10.1038/22313

Chee, M. J., Hebert, A. J., Briançon, N., Flaherty, S. E., Pissios, P., & Maratos-Flier, E. (2019). Conditional deletion of melanin-concentrating hormone receptor 1 from GABAergic neurons increases locomotor activity. *Molecular Metabolism*, *29*(August), 114–123. https://doi.org/10.1016/j.molmet.2019.08.018

Chen, Y., Hu, C., Hsu, C. K., Zhang, Q., Bi, C., Asnicar, M., Hsiung, H. M., Fox, N., Slieker, L. J., Yang, D. D., Heiman, M. L., & Shi, Y. (2002). Targeted disruption of the melanin-concentrating hormone receptor-1 results in hyperphagia and resistance to diet-induced obesity. *Endocrinology*, *143*(7), 2469–2477. https://doi.org/10.1210/endo.143.7.8903

Cippitelli, A., Karlsson, C., Shaw, J. L., Thorsell, A., Gehlert, D. R., & Heilig, M. (2010).

Suppression of alcohol self-administration and reinstatement of alcohol seeking by melanin-

concentrating hormone receptor 1 (MCH1-R) antagonism in Wistar rats.

Psychopharmacology, *211*(4), 367–375. https://doi.org/10.1007/s00213-010-1891-y

Dal Bo, G., St.-Gelais, F., Danik, M., Williams, S., Cotton, M., & Trudeau, L. E. (2004).

Dopamine neurons in culture express VGLUT2 explaining their capacity to release

glutamate at synapses in addition to dopamine. *Journal of Neurochemistry*, *88*(6), 1398–

1405. https://doi.org/10.1046/j.1471-4159.2003.02277.x

Diniz, G. B., Battagello, D. S., Klein, M. O., Bono, B. S. M., Ferreira, J. G. P., Motta-Teixeira,

L. C., Duarte, J. C. G., Presse, F., Nahon, J. L., Adamantidis, A., Chee, M. J., Sita, L. V., &

Bittencourt, J. C. (2020). Ciliary melanin-concentrating hormone receptor 1 (MCHR1) is

widely distributed in the murine CNS in a sex-independent manner. *Journal of*

Neuroscience Research, *98*(10), 2045–2071. https://doi.org/10.1002/jnr.24651

Dobi, A., Margolis, E. B., Wang, H. L., Harvey, B. K., & Morales, M. (2010). Glutamatergic and

nonglutamatergic neurons of the ventral tegmental area establish local synaptic contacts

with dopaminergic and nondopaminergic neurons. *Journal of Neuroscience*, *30*(1), 218–

229. https://doi.org/10.1523/JNEUROSCI.3884-09.2010

Domingos, A. I., Sordillo, A., Dietrich, M. O., Liu, Z. W., Tellez, L. A., Vaynshteyn, J., Ferreira,

J. G., Ekstrand, M. I., Horvath, T. L., de Araujo, I. E., & Friedman, J. M. (2013).

Hypothalamic melanin concentrating hormone neurons communicate the nutrient value of

sugar. *ELife*, *2013*(2), 1–15. https://doi.org/10.7554/eLife.01462

Dong, H. (2008). *The Allen reference atlas: A digital color brain atlas of the C57BL/6J male*

mouse. John Wiley & Sons.

Faget, L., Osakada, F., Duan, J., Ressler, R., Johnson, A. B., Proudfoot, J. A., Yoo, J. H.,

Callaway, E. M., & Hnasko, T. S. (2016). Afferent Inputs to Neurotransmitter-Defined Cell

Types in the Ventral Tegmental Area. *Cell Reports*, *15*(12), 2796–2808.

https://doi.org/10.1016/j.celrep.2016.05.057

Gao, X.-B., & Pol, A. N. van den. (2001). Melanin concentrating hormone depresses synaptic

activity of glutamate and GABA neurons from rat lateral hypothalamus. *Journal of*

Physiology, *533*(1), 237–252.

Geisler, S., & Wise, R. A. (2008). Functional implications of glutamatergic projections to the

ventral tegmental area. *Reviews in the Neurosciences*, *19*(4–5), 227–244.

https://doi.org/10.1515/REVNEURO.2008.19.4-5.227

Georgescu, D., Sears, R. M., Hommel, J. D., Barrot, M., Bolaños, C. A., Marsh, D. J., Bednarek,

M. A., Bibb, J. A., Maratos-Flier, E., Nestler, E. J., & DiLeone, R. J. (2005). The

hypothalamic neuropeptide melanin-concentrating hormone acts in the nucleus accumbens

to modulate feeding behavior and forced-swim performance. *Journal of Neuroscience*,

25(11), 2933–2940. https://doi.org/10.1523/JNEUROSCI.1714-04.2005

Geuzaine, A., Tyhon, A., Grisar, T., Brabant, C., Lakaye, B., & Tirelli, E. (2014). Amphetamine

reward in food restricted mice lacking the melanin-concentrating hormone receptor-1.

Behavioural Brain Research, *262*, 14–20. https://doi.org/10.1016/j.bbr.2013.12.052

Guan, Y., Xiao, C., Krnjević, K., Xie, G., Zuo, W., & Ye, J.-H. (2012). GABAergic Actions

Mediate Opposite Ethanol Effects on Dopaminergic Neurons in the Anterior and Posterior

Ventral Tegmental Area. *Journal of Pharmacology and Experimental Therapeutics*, *341*(1),

33–42. https://doi.org/10.1124/jpet.111.187963

666 Hawes, B. E., Erin, K. I. L., Green, B., O'Neill, K. I. M., Fried, S., & Graziano, M. P. (2000).

667  The melanin-concentrating hormone receptor couples to multiple G proteins to activate

668  diverse intracellular signaling pathways. *Endocrinology*, *141*(12), 4524–4532.

669  https://doi.org/10.1210/endo.141.12.7833

670 Hervieu, G. J., Cluderay, J. E., Harrison, D., Meakin, J., Maycox, P., Nasir, S., & Leslie, R. A.

671  (2000). The distribution of the mRNA and protein products of the melanin-concentrating

672  hormone (MCH) receptor gene, slc-1, in the central nervous system of the ratt. *European*

673  *Journal of Neuroscience*, *12*(4), 1194–1216. https://doi.org/10.1046/j.1460-

674  9568.2000.00008.x

675 Hiroi, N., & White, N. M. (1991). The amphetamine conditioned place preference: differential

676  involvement of dopamine receptor subtypes and two dopaminergic terminal areas. *Brain*

677  *Research*, *552*(1), 141–152. https://doi.org/10.1016/0006-8993(91)90672-I

678 Jhou, T., Fields, H., Baxter, M., & Saper, C. (2009). The rostromedial tegmental nucleus

679  (RMTg), a major GABAergic afferent to midbrain dopamine neurons, selectively encodes

680  aversive stimuli and promotes behavioral inhibition. *Neuron*, *61*(5), 786–800.

681  https://doi.org/10.1016/j.neuron.2009.02.001.The

682 Karlsson, C., Rehman, F., Damdazic, R., Atkins, A. L., Schank, J. R., Gehlert, D. R., Steensland,

683  P., Thorsell, A., & Heilig, M. (2016). The melanin-concentrating hormone-1 receptor

684  modulates alcohol-induced reward and DARPP-32 phosphorylation. *Psychopharmacology*,

685  *233*(12), 2355–2363. https://doi.org/10.1007/s00213-016-4285-y

686 Kokkotou, E., Jeon, J. Y., Wang, X., Marino, F. E., Carlson, M., Trombly, D. J., & Maratos-

687  Flier, E. (2005). Mice with MCH ablation resist diet-induced obesity through strain-specific

mechanisms. *American Journal of Physiology - Regulatory Integrative and Comparative*

Physiology, *289*(1 58-1), 117–124. https://doi.org/10.1152/ajpregu.00861.2004

Krashes, M. J., Shah, B. P., Madara, J. C., Olson, D. P., Strochlic, D. E., Garfield, A. S., Vong,

L., Pei, H., Watabe-Uchida, M., Uchida, N., Liberles, S. D., & Lowell, B. B. (2014). An

excitatory paraventricular nucleus to AgRP neuron circuit that drives hunger. *Nature*,

507(7491), 238–242. https://doi.org/10.1038/nature12956.A

Lahti, L., Haugas, M., Tikker, L., Airavaara, M., Voutilainen, M. H., Anttila, J., Kumar, S.,

Inkinen, C., Salminen, M., & Partanen, J. (2016). Differentiation and molecular

heterogeneity of inhibitory and excitatory neurons associated with midbrain dopaminergic

nuclei. *Development (Cambridge)*, *143*(3), 516–529. https://doi.org/10.1242/dev.129957

Lammel, S., Steinberg, E. E., Földy, C., Wall, N. R., Beier, K., Luo, L., & Malenka, R. C.

(2015). Diversity of Transgenic Mouse Models for Selective Targeting of Midbrain

Dopamine Neurons. *Neuron*, *85*(2), 429–438.

https://doi.org/10.1053/j.gastro.2016.08.014.CagY

Lecca, S., Melis, M., Luchicchi, A., Muntoni, A. L., & Pistis, M. (2012). Inhibitory inputs from

rostromedial tegmental neurons regulate spontaneous activity of midbrain dopamine cells

and their responses to drugs of abuse. *Neuropsychopharmacology*, *37*(5), 1164–1176.

https://doi.org/10.1038/npp.2011.302

Lembo, P. M. C., Grazzini, E., Cao, J., Hubatsch, D. A., Pelletier, M., Hoffert, C., St-Onge, S.,

Pou, C., Labrecque, J., Groblewski, T., O'Donnell, D., Payza, K., Ahmad, S., & Walker, P.

(1999). The receptor for the orexigenic peptide melanin-concentrating hormone is a G-

protein-coupled receptor. *Nature Cell Biology*, *1*(5), 267–271.

https://doi.org/10.1038/12978

Lindeberg, J., Usoskin, D., Bengtsson, H., Gustafsson, A., Kylberg, A., Söderström, S., &

Ebendal, T. (2004). Transgenic expression of Cre recombinase from the tyrosine

hydroxylase locus. *Genesis (United States)*, *40*(2), 67–73.

https://doi.org/10.1002/gene.20065

Lord, M. N., Subramanian, K., Kanoski, S. E., Noble, E. E., Angeles, L., Section, E. B.,

California, S., & Angeles, L. (2022). Melanin-concentrating hormone and food intake

control: sites of action, peptide interactions, and appetition. *Peptides*, *137*, 1–23.

https://doi.org/10.1016/j.peptides.2020.170476.Melanin-concentrating

Ludwig, D. S., Tritos, N. A., Mastaitis, J. W., Kulkarni, R., Kokkotou, E., Elmquist, J., Lowell,

B., Flier, J. S., & Maratos-Flier, E. (2001). Melanin-concentrating hormone overexpression

in transgenic mice leads to obesity and insulin resistance. *Journal of Clinical Investigation*,

107(3), 379–386. https://doi.org/10.1172/JCI10660

Marsh, D. J., Weingarth, D. T., Novi, D. E., Chen, H. Y., Trumbauer, M. E., Chen, A. S., Guan,

X. M., Jiang, M. M., Feng, Y., Camacho, R. E., Shen, Z., Frazier, E. G., Yu, H., Metzger, J.

M., Kuca, S. J., Shearman, L. P., Gopal-Truter, S., MacNeil, D. J., Strack, A. M., … Qian,

S. (2002). Melanin-concentrating hormone 1 receptor-deficient mice are lean, hyperactive,

and hyperphagic and have altered metabolism. *Proceedings of the National Academy of*

Sciences of the United States of America, *99*(5), 3240–3245.

https://doi.org/10.1073/pnas.052706899

Mead, A. N., & Stephens, D. N. (1999). CNQX but not NBQX prevents expression of

amphetamine-induced place preference conditioning: a role for the glycine site of the

NMDA receptor, but not AMPA receptors. *The Journal of Pharmacology and Experimental*

Therapeutics, *290*(1), 9–15.

Mickelsen, L. E., Bolisetty, M., Chimileski, B. R., Fujita, A., Beltrami, E. J., Costanzo, J. T.,

Naparstek, J. R., Robson, P., & Jackson, A. C. (2019). Single-cell transcriptomic analysis of

the lateral hypothalamic area reveals molecularly distinct populations of inhibitory and

excitatory neurons. *Nature Neuroscience*, *22*(4), 642–656. https://doi.org/10.1038/s41593-

019-0349-8

Morales, M., & Root, D. H. (2014). Glutamate neurons within the midbrain dopamine regions.

Neuroscience, *282*, 60–68. https://doi.org/10.1016/j.neuroscience.2014.05.032

Morales, Marisela, & Margolis, E. B. (2017). Ventral tegmental area: Cellular heterogeneity,

connectivity and behaviour. *Nature Reviews Neuroscience*, *18*(2), 73–85.

https://doi.org/10.1038/nrn.2016.165

Mul, J. D., la Fleur, S. E., Toonen, P. W., Afrasiab-Middelman, A., Binnekade, R., Schetters, D.,

Verheij, M. M. M., Sears, R. M., Homberg, J. R., Schoffelmeer, A. N. M., Adan, R. A. H.,

DiLeone, R. J., de Vries, T. J., & Cuppen, E. (2011). Chronic loss of melanin-concentrating

hormone affects motivational aspects of feeding in the rat. *PLoS ONE*, *6*(5).

https://doi.org/10.1371/journal.pone.0019600

Negishi, K., Payant, M. A., Schumacker, K. S., Wittmann, G., Butler, R. M., Lechan, R. M.,

Steinbusch, H. W. M., Khan, A. M., & Chee, M. J. (2020). Distributions of hypothalamic

neuron populations coexpressing tyrosine hydroxylase and the vesicular GABA transporter

in the mouse. *Journal of Comparative Neurology*, *528*(11), 1833–1855.

https://doi.org/10.1002/cne.24857

Nieh, E. H., Vander Weele, C. M., Matthews, G. A., Presbrey, K. N., Wichmann, R., Leppla, C.

A., Izadmehr, E. M., & Tye, K. M. (2016). Inhibitory Input from the Lateral Hypothalamus

to the Ventral Tegmental Area Disinhibits Dopamine Neurons and Promotes Behavioral

Activation. *Neuron*, *90*(6), 1286–1298. https://doi.org/10.1016/j.neuron.2016.04.035

Pissios, P., Frank, L., Kennedy, A. R., Porter, D. R., Marino, F. E., Liu, F. F., Pothos, E. N., &

Maratos-Flier, E. (2008). Dysregulation of the Mesolimbic Dopamine System and Reward

in MCH-/- Mice. *Biological Psychiatry*, *64*(3), 184–191.

https://doi.org/10.1016/j.biopsych.2007.12.011

Pissios, P., Trombly, D. J., Tzameli, I., & Maratos-Flier, E. (2003). Melanin-concentrating

hormone receptor 1 activates extracellular signal-regulated kinase and synergizes with Gs-

coupled pathways. *Endocrinology*, *144*(8), 3514–3523. https://doi.org/10.1210/en.2002-

0004

Qu, D., Ludwig, D. S., Gammeltoftt, S., Piper, M., Pelleymountert, M. A., Cullent, M. J.,

Mathes, W. F., Przypek, J., Kanarek, R., & Ii, E. M. (1996). A role for melanin-

concentrating hormone in the central regulation of feeding behaviour. *Nature*, *380*(March),

243–247.

Root, D. H., Mejias-Aponte, C. A., Qi, J., & Morales, M. (2014). Role of glutamatergic

projections from ventral tegmental area to lateral Habenula in aversive conditioning.

Journal of Neuroscience, *34*(42), 13906–13910. https://doi.org/10.1523/JNEUROSCI.2029-

14.2014

Root, D. H., Mejias-Aponte, C. A., Zhang, S., Wang, H. L., Hoffman, A. F., Lupica, C. R., &

Morales, M. (2014). Single rodent mesohabenular axons release glutamate and GABA.

Nature Neuroscience, *17*(11), 1543–1551. https://doi.org/10.1038/nn.3823

Root, D. H., Zhang, S., Barker, D. J., Miranda-Barrientos, J., Liu, B., Wang, H. L., & Morales,

M. (2018). Selective Brain Distribution and Distinctive Synaptic Architecture of Dual

Glutamatergic-GABAergic Neurons. *Cell Reports*, *23*(12), 3465–3479.

https://doi.org/10.1016/j.celrep.2018.05.063

Saito, Y., Cheng, M., Leslie, F. M., & Civelli, O. (2001). Expression of the melanin-

concentrating hormone (MCH) receptor mRNA in the rat brain. *Journal of Comparative*

Neurology, *435*(1), 26–40. https://doi.org/10.1002/cne.1191

Saito, Y., Nothacker, H., Wang, Z., Lin, S. H. S., Leslie, F., & Civelli, O. (1999). Molecular

characterization of the melanin-concentrating- hormone receptor. *Nature*, *400*(July), 265–

269.

Schneeberger, M., Tan, K., Nectow, A. R., Parolari, L., Caglar, C., Azevedo, E., Li, Z.,

Domingos, A., & Friedman, J. M. (2018). Functional analysis reveals differential effects of

glutamate and MCH neuropeptide in MCH neurons. *Molecular Metabolism*, *13*(May), 83–

89. https://doi.org/10.1016/j.molmet.2018.05.001

Sears, R. M., Liu, R. J., Narayanan, N. S., Sharf, R., Yeckel, M. F., Laubach, M., Aghajanian, G.

K., & DiLeone, R. J. (2010). Regulation of nucleus accumbens activity by the hypothalamic

neuropeptide melanin-concentrating hormone. *Journal of Neuroscience*, *30*(24), 8263–

8273. https://doi.org/10.1523/JNEUROSCI.5858-09.2010

Shimada, M., Tritos, N. A., Lowell, B. B., Flier, J. S., & Maratos-Flier, E. (1998). Mice lacking

melanin- concentrating hormone are hypophagic and lean. *Nature*, *396*(December), 3–7.

Slusher, B. S., Thomas, A., Paul, M., Schad, C. A., & Ashby, C. R. (2001). Expression and

acquisition of the conditioned place preference response to cocaine in rats is blocked by

selective inhibitors of the enzyme N-Acetylated-α-Linked-Acidic Dipeptidase

(NAALADase). *Synapse*, *41*(1), 22–28. https://doi.org/10.1002/syn.1056

Smith, R. J., Vento, P. J., Chao, Y. S., Good, C. H., Thomas, C., & Ground, A. P. (2019). Gene

expression and neurochemical characterization of the rostromedial tegmental nucleus

(RMTg) in rats and mice. *Brain Struct Funct*, *224*(1), 219–238.

https://doi.org/10.1007/s00429-018-1761-7.Gene

Stamatakis, A. M., Jennings, J. H., Ung, R. L., Blair, G. A., Weinberg, J., Neve, R. L., Boyce, F.,

Mattis, J., Deisseroth, K., & Stuber, G. D. (2013). A unique population of ventral tegmental

area neurons inhibits the lateral habenula to promote reward. *Neuron*, *80*(4), 1–23.

https://doi.org/10.1016/j.neuron.2013.08.023.A

Stuber, G. D., Evans, S. B., Higgins, M. S., Pu, Y., & Figlewicz, D. P. (2002). Food restriction

modulates amphetamine-conditioned place preference and nucleus accumbens dopamine

release in the rat. *Synapse*, *46*(2), 83–90. https://doi.org/10.1002/syn.10120

Stuber, G. D., Hnasko, T. S., Britt, J. P., Edwards, R. H., & Bonci, A. (2010). Dopaminergic

terminals in the nucleus accumbens but not the dorsal striatum corelease glutamate. *Journal*

of Neuroscience, *30*(24), 8229–8233. https://doi.org/10.1523/JNEUROSCI.1754-10.2010

Sulzer, D., Joyce, M. P., Lin, L., Geldwert, D., Haber, S. N., Hattori, T., & Rayport, S. (1998).

Dopamine neurons make glutamatergic synapses in vitro. *Journal of Neuroscience*, *18*(12),

4588–4602. https://doi.org/10.1523/jneurosci.18-12-04588.1998

Tecuapetla, F., Patel, J. C., Xenias, H., English, D., Tadros, I., Shah, F., Berlin, J., Deisseroth,

K., Rice, M. E., Tepper, J. M., & Koos, T. (2010). Glutamatergic signaling by mesolimbic dopamine neurons in the nucleus accumbens. *Journal of Neuroscience*, *30*(20), 7105–7110. https://doi.org/10.1523/JNEUROSCI.0265-10.2010

Ting, J. T., Lee, B. R., Chong, P., Soler-Llavina, G., Cobbs, C., Koch, C., Zeng, H., & Lein, E. (2018). Preparation of Acute Brain Slices Using an Optimized N-Methyl-D-glucamine Protective Recovery Method. *Journal of Visualized Experiments : JoVE*, *132*, 1–13. https://doi.org/10.3791/53825

Tritsch, N. X., Ding, J. B., & Sabatini, B. L. (2012). Dopaminergic neurons inhibit striatal output via non-canonical release of GABA. *Nature*, *490*, 262–266. https://doi.org/doi:10.1038/nature11466

Tzschentke, T. M., & Schmidt, W. J. (1998). Blockade of morphine- and amphetamine-induced conditioned place preference in the rat by riluzole. *Neuroscience Letters*, *242*, 114–116. papers2://publication/uuid/3FF1E7DE-1CAD-497E-A55B-478F8D972A24

Vong, L., Ye, C., Yang, Z., Choi, B., Jr., S. C., & Lowell, B. B. (2011). Leptin Action on GABAergic Neurons Prevents Obesity and Reduces Inhibitory Tone to POMC Neurons. *Neuron*, *71*(1), 142–154. https://doi.org/10.1016/j.neuron.2011.05.028.Leptin

Wang, H. L., Qi, J., Zhang, S., Wang, H., & Morales, M. (2015). Rewarding effects of optical stimulation of ventral tegmental area glutamatergic neurons. *Journal of Neuroscience*, *35*(48), 15948–15954. https://doi.org/10.1523/JNEUROSCI.3428-15.2015

Yamaguchi, T., Sheen, W., & Morales, M. (2007). Glutamatergic neurons are present in the rat ventral tegmental area. *Eur J Neurosci*, *25*(1), 106–118. https://doi.org/doi:10.1111/j.1460-9568.2006.05263.x.

Yamaguchi, T., Wang, H. L., Li, X., Ng, T. H., & Morales, M. (2011). Mesocorticolimbic

glutamatergic pathway. *Journal of Neuroscience*, *31*(23), 8476–8490.

https://doi.org/10.1523/JNEUROSCI.1598-11.2011

Legends

Figure 1. *Mchr1* mRNA and MCHR1 protein expression in wildtype mouse VTA. qPCR

analysis of *Mchr1* mRNA from the hypothalamus (HT), striatum (STR), hippocampus (HC),

ventral tegmental area (VTA), and cerebellum (CB) of male and female wildtype mice (**A**).

Confocal photomicrographs of MCHR1-immunoreactive signals on Nissl-stained (NeuroTrace)

soma in wildtype (**B**) and *Mchr1-KO* (**C**) VTA at low (***i***) and high magnification (***ii***, from dash-

outlined region in ***i***). Representative examples of the presence (white arrowhead) or absence

(open arrowhead) of MCHR1 immunoreactivity at the primary cilium of VTA cells. Number of

MCHR1-immunoreactive cells throughout the VTA of male and female wildtype mice (**D**). VTA

levels were assigned with reference to the *Allen Reference Atlas* (*ARA*; Dong, 2008). Two-way

ANOVA with Bonferroni post-test: ****, $p < 0.0001$ (STR vs all regions; **A**). Scale bars: ***i***, 100

µm; ***ii***, 40 µm. IF, interfascicular nucleus raphe; IPN, interpeduncular nucleus; VTA, ventral

tegmental area.

Figure 2. Neurochemical characterization of VTA cells. Number (right *y*-axis, gray line) and

proportion of each cell type (left *y*-axis, stacked colored bars) at each *Allen Reference Atlas* level

(Dong, 2008) rostrocaudally from VTA slices expressing *Vglut2* mRNA hybridization, *Vgat*

mRNA hybridization, and tyrosine hydroxylase (TH) immunoreactivity (N = 6 mice; **A**).

Representative coronal brain maps of VTA cells that are *Vglut2-*, *Vgat-* and/or TH-positive (**B**).

Each panel includes the nomenclature, atlas level and Bregma value (β) assigned to that tissue in

accordance with the *Allen Reference Atlas* (Dong, 2008).

Figure 3. *Mchr1* expression on diverse VTA cell types. Representative low (**A**) and high

magnification confocal photomicrographs (**B**–**E**), from corresponding dash-outlined region in **A**)

of *Vglut2* mRNA, *Vgat* mRNA, and/or TH immunoreactivity (*i*) that coexpress (white

arrowhead) or lack (open arrowhead) *Mchr1* mRNA hybridization in VTA cells (*ii*). Relative

proportion of *Mchr1*-expressing VTA cells overall (**F**) and at each level, as determined with

reference to the *Allen Reference Atlas* (Dong, 2008), of the VTA (N = 6; **G**). Scale bars: **A***,* 250

µm; **B**–**E**, 30 µm. IF, interfascicular nucleus raphe; IPN, interpeduncular nucleus; ml, medial

lemniscus; RN red nucleus; SNc, substantia nigra, compact part; SNr, substantia nigra, reticular

part; VTA, ventral tegmental area.

Figure 4. Spatial distribution of *Mchr1* mRNA in heterogeneous VTA cells. *Mchr1* mRNA

hybridization expression in *Vglut2* (**A**; **Figure 4-1**), *Vgat* (**B**; **Figure 4-2***),* TH (**C**; **Figure 4-3**),

or dual *Vglut2* and *Vgat* VTA cells (**D**; **Figure 4-4**) mapped to a representative anterior (*i*),

middle (*ii*), and posterior (*iii*) VTA section using *Allen Reference Atlas* (Dong, 2008) templates

at the atlas level or Bregma value (β) indicated (*top*).

Figure 5. MCH directly hyperpolarized GABAergic and dopaminergic VTA cells.

Representative epifluorescence image of native EGFP fluorescence from VTA cells (*inset*) in

Vglut2-cre;L10-Egfp (*Vglut2*^{EGFP}), *Vgat-cre;L10-Egfp* (*Vgat*^{EGFP}), and *Th-cre;L10-Egfp* (*Th*^{EGFP})

reporter mice guided whole-cell patch-clamp recordings within the VTA (**A**). Change in resting

membrane potential (Δ RMP) over time and the proportion of MCH-responsive cells (donut plot)

at *Vglut2*^{EGFP} (**B**), *Vgat* ^{EGFP} (**C**), and *Th*^{EGFP} VTA cells (**D**) following the bath application of 3

µM MCH. MCH-sensitive VTA cells identified by a localized puff of MCH applied within 20

µm of the patched cell (**E***i*). Representative sample trace of the membrane potential change

following a puff of MCH lasting <5 s (**E***ii*). RMP change over time following the first (black

circles) and second (white circles) localized puff of 3 µM MCH or of bath ACSF (gray circles)

onto a VTA cell (**E***iii*). RMP change over time following localized puff of 3 µM MCH in the

presence (red outlined circles) or absence of 500 nM TTX (black circles) at $Vgat^{\text{EGFP}}$ (**F**) and

Th^{EGFP} VTA cells (**G**). Scale bars: **A**, 250 μm; **A** *inset*, 20 μm; **E***i*, 25 μm; **E***ii*, 3 mV, 10 s. SNc,

substantia nigra, compact part; SNr, substantia nigra, reticular part; VTA, ventral tegmental area.

Figure 6. **MCH-mediated disinhibition of local excitatory input to *Th* VTA cells.** Change in

spontaneous inhibitory postsynaptic current (Δ sIPSC) event frequency over time following bath

application of 3 μM MCH at EGFP-labeled Th^{EGFP} VTA cells from *Th-cre;L10-Egfp* mice (**A**).

Change in spontaneous excitatory postsynaptic current (Δ sEPSC) event frequency at Th^{EGFP}

VTA cells following 3 μM MCH (black circles) or bath ACSF (gray circles) application (**B**).

Change in miniature excitatory postsynaptic current (Δ mEPSC) event frequency at Th^{EGFP} VTA

cells following MCH application in the presence of 500 nM TTX (**C**). Change in sIPSC

frequency at $Vglut2^{\text{EGFP}}$ VTA cells following MCH application (**D**). Model of bidirectional,

MCH-mediated regulation of dopamine release from the VTA to nucleus accumbens (NAc) via

the mesolimbic pathway (**E**). Schematic showing inferred MCHR1 distribution within the VTA

to elicit MCH-mediated direct inhibition of Th^{EGFP} VTA cells (***i***) and delayed, indirect GABA-

mediated disinhibition of glutamatergic input to Th^{EGFP} VTA cells (***ii***). Putative MCHR1

expression (*****) along efferent projections of VTA $Vglut2^{\text{EGFP}}$ cells, which were not MCH-

responsive.

Extended Data

Figure 3-1. Relative distribution of *Mchr1*-expressing VTA cells in male and female mice

(**A**). Sex may interact with *Vgat* expression to shift the distribution of *Mchr1* mRNA (**B**).

Figure 4-1. Distribution of *Mchr1*-expressing *Vglut2* VTA cells. Number of *Mchr1*-positive

(dark green) or -negative (light green) *Vglut2* cells relative to all VTA cells counted (gray) at

each level in accordance with the *Allen Reference Atlas* (*ARA*; Dong, 2008) (N = 6; **A**). Presence

(dark green circles) or absence (light green circles) of *Mchr1* mRNA hybridization in *Vglut2*

cells throughout the anteroposterior extent of the VTA (**B**–**L**) mapped to *ARA*. Each panel

includes the nomenclature, atlas level and Bregma value (β) assigned to that tissue in accordance

with the *ARA*.

Figure 4-2. Distribution of *Mchr1*-expressing *Vgat* VTA cells. Number of *Mchr1*-positive

(dark yellow) or -negative (light yellow) *Vgat* cells relative to all VTA cells counted (gray) at

each level in accordance with the *Allen Reference Atlas* (*ARA*; Dong, 2008) (N = 6; **A**). Presence

(dark yellow circles) or absence (light yellow circles) of *Mchr1* mRNA hybridization in *Vgat*

cells throughout the anteroposterior extent of the VTA (**B**–**L**) mapped to *ARA*. Each panel

includes the nomenclature, atlas level and Bregma value (β) assigned to that tissue in accordance

with the *ARA*.

Figure 4-3. Distribution of *Mchr1*-expressing TH VTA cells. Number of *Mchr1*-positive (red)

or -negative (pink) TH-immunoreactive cells relative to all VTA cells counted (gray) at each

level in accordance with the *Allen Reference Atlas* (*ARA*; Dong, 2008) (N = 6; **A**). Presence (red

circles) or absence (pink circles) of *Mchr1* mRNA hybridization in TH cells throughout the

anteroposterior extent of the VTA (**B**–**L**) mapped to *ARA*. Each panel includes the nomenclature,

atlas level and Bregma value (β) assigned to that tissue in accordance with the *ARA*.

Figure 4-4. Distribution of *Mchr1* mRNA expression *Vgat*- and *Vglut2*-positive VTA cells.

Number of *Mchr1*-positive (dark blue) or -negative (light blue) cells expressing both *Vglut2* and

Vgat mRNA relative to all VTA cells counted (gray) at each level in accordance with the *Allen*

Reference Atlas (*ARA*; Dong, 2008) (N = 6; **A**). Presence (dark blue circles) or absence (light

blue circles) of *Mchr1* mRNA hybridization in dual-labeled *Vglut2* and *Vgat* cells throughout the

anteroposterior extent of the VTA (**B**–**L**) mapped to *ARA*. Each panel includes the nomenclature,

atlas level and Bregma value (β) assigned to that tissue in accordance with the *ARA*.

A

Relative *Mchr1* expression

Brain region: HT, STR, HC, VTA, CB

Male (7)
Female (6)

**** ****

B **WT**

i NeuroTrace MCHR1

VTA
IF
IPN

ii NeuroTrace MCHR1

C *Mchr1-KO*

i NeuroTrace MCHR1

VTA
IF
IPN

ii NeuroTrace MCHR1

D

Number of MCHR1 cells

Allen Reference Atlas level: 81, 82, 83, 84, 85, 86, 87, 88, 89, 90

Male (3)
Female (3)

Figure 2

Figure 3

Figure 6