Virulence and transmission biology of the widespread, ecologically important pathogen

of zooplankton, *Spirobacillus cienkowskii*

Authors

Nina Wale[a,b,c,d], Claire B. Freimark[d,e], Justin Ramirez[d], Marcin K. Dziuba[d] Ahmad Y. Kafri[d],

Rebecca Bilich[d,f] , and Meghan A. Duffy[d]

[a]Department of Microbiology & Molecular Genetics, Michigan State University, East Lansing,

Michigan, USA

[b]Department of Integrative Biology, Michigan State University, East Lansing, Michigan, USA,

[c]Program in Ecology, Evolution and Behavior, Michigan State University, East Lansing,

Michigan, USA

[d] Department of Ecology & Evolutionary Biology, University of Michigan, Ann Arbor, Michigan,

USA

[e] Present address: Department of Ecology & Evolutionary Biology, University of California Irvine,

Irvine, California, USA

[f] Present address: School of Medicine, Wayne State University, Detroit, Michigan, USA

Abstract

Spirobacillus cienkowskii (*Spirobacillus,* hereafter) is a widely distributed bacterial pathogen of

zooplankton (*Daphnia* sp.) that has significant impacts on zooplankton population dynamics.

Despite its ecological importance, little is known about *Spirobacillus'* virulence and transmission

biology. These knowledge gaps stymie our understanding of *Daphnia-Spirobacillus* interactions

and the ecology of the newly described family *Silvanigrellaceae,* of which *Spirobacillus* is the

longest studied member. Here, we present our work to fill these gaps, experimentally. We found

that *Spirobacillus* is among the most virulent of *Daphnia* pathogens, killing its host within a week

of exposure and reducing host fecundity. Our experiments further suggest that *Spirobacillus*

requires host death to transmit horizontally i.e., it is an "obligate killer"*,* since it did not transmit

horizontally among hosts unless the host died or was destroyed. In a series of further

experiments characterizing the dynamics of horizontal transmission, we found that *Spirobacillus'*

per capita rate of transmission is low, does not change with host density or pathogen inoculum

size, and decreases rapidly with time spent in the environment. Lastly, we complemented our

experimental work with a field survey of *Spirobacillus'* prevalence within ephippia, seed-like

cases containing *Daphnia* offspring. We found *Spirobacillus* DNA in almost 40% of ephippia,

suggesting that *Spirobacillus* is vertically transmitted and/or can invade ephippia and use them

to endure environmental stress, as their hosts do. Our work fills critical gaps in our

understanding of this ecologically important pathogen's biology.

Importance

Spirobacillus cienkowskii is a bacterial pathogen of zooplankton, first described in the 19^th

Century and recently placed in a new family of bacteria, the *Silvanigrellaceae*. *Spirobacillus*

causes large epidemics in lake zooplankton populations and increases the probability that

zooplankton will be eaten by predators. However, little is known about how *Spirobacillus*

transmits among hosts, to what extent it reduces host survival and reproduction (i.e., how

virulent it is) and what role virulence plays in *Spirobacillus'* life cycle. Here, we experimentally

quantified *Spirobacillus*' virulence and showed that *Spirobacillus* must kill its host to transmit

horizontally. We also found evidence that *Spirobacillus* may transmit vertically via *Daphnia*'s

seed-like egg sacks. Our work will help scientists to (i) understand *Spirobacillus* epidemics, (ii)

use *Spirobacillus* as a model pathogen for the study of host-parasite interactions and (iii) better

understand the unusual group of bacteria to which *Spirobacillus* belongs.

Introduction

Interactions among pathogens and *Daphnia* can have significant impacts on *Daphnia* population

dynamics, evolution and trophic interactions, and serve as important models of host-pathogen

interactions (1–8). One of the oldest *Daphnia* pathogens known to science is *Spirobacillus*

cienkowskii, a gram-negative bacterium that belongs to the newly characterized family,

Silvanigrellaceae (9). First found in *Daphnia magna* in Ukraine in the 1880s (10), *Spirobacillus*

cienkowskii (hereafter *Spirobacillus*) has since been found in a wide range of *Cladocera* at

locations across the globe (11–13). In *Daphnia* populations in Michigan, where *Spirobacillus'*

dynamics have been most intensively studied, *Spirobacillus* causes large epidemics, depleting

the numbers of hosts and increasing their vulnerability to predation (4, 14, 15). *Spirobacillus'*

relatives in the *Silvanigrellaceae* may also be host-associated, and perhaps even pathogenic,

as evidenced by their presence on frog skin and in the tissues of a sick person (16, 17). By

better understanding the biology of *Spirobacillus* we thus stand to gain a deeper understanding

of the eco-epidemiological dynamics of zooplankton and the biology of a newly described group

of bacteria, the *Silvanigrellaceae*.

A first step toward this goal is to fill critical gaps in our understanding of *Spirobacillus*' virulence

and transmission biology, which will together drive its evolutionary and epidemiological

dynamics (18). These gaps exist because *Spirobacillus*' has been difficult to establish in the

laboratory; nevertheless, field studies, limited laboratory investigations and genomic analyses

have yielded some insights and hypotheses. For example, the observation that *Spirobacillus*-

infected*,* asexually reproducing *Daphnia* rarely bear offspring and always die from infection (11,

19), gave rise to the hypothesis that *Spirobacillus* is an "obligate-killer" i.e., a pathogen that

requires host death to transmit horizontally. This hypothesis remains untested, however.

Similarly, very little is known about the dynamics of *Spirobacillus'* horizontal transmission or the

ecological factors that impact it. An experimental investigation demonstrated that *Spirobacillus*

transmits via water (19), presumably by motile free-living stages and/or spore-like stages that

have been observed via microscopy and for which there is genomic evidence (10, 20). The

infectivity of free-living stages, and the rate at which their infectivity decays, are critical drivers of

the epidemiological dynamics and virulence of environmentally transmitted pathogens (21–23)

and can be impacted by a range of abiotic and biotic factors (22, 24, 25). Thus, understanding

the ecology of horizontal transmission is essential if we are to explain *Spirobacillus'* epidemic

dynamics. Lastly, to better understand how *Spirobacillus* persists between the seasons when

epidemics occur, further investigation is required to establish if *Spirobacillus* is capable of

vertical, as well as horizontal, transmission, as prior observations suggest (11).

In 2016, we overcame a major obstacle to filling these knowledge gaps by establishing

Spirobacillus in *in vivo* culture (19) and thereafter performed a series of experiments to answer

fundamental questions about *Spirobacillus'* infection biology, which we present here. Our

experiments demonstrate that *Spirobacillus* is a highly virulent, obligate killer that dramatically

reduces host fecundity. Consistent with previous laboratory observations, but surprising in the

light of the large epidemics that have been observed in the field, we show that the

transmissibility of *Spirobacillus* is consistently very low, does not vary with host density or

parasite inoculum size (i.e., the amount of bacterium to which hosts were exposed, hereafter

"dose") and declines rapidly after host death. Lastly, consistent with the hypotheses that

Spirobacillus is capable of vertical transmission or invades ephippia while in a free-living state in

the water column, we show that a high frequency of field-collected ephippia harbor

Spirobacillus. Our work advances our understanding of the basic biology of this little studied

pathogen and leads to multiple, testable hypotheses as to how *Spirobacillus* can persist in

Daphnia populations, despite its apparently low horizontal transmission rate.

Materials & Methods

Individual-scale experiments

To investigate the virulence and transmission dynamics of *Spirobacillus*, we conducted several

"individual-scale" experiments, wherein individual *Daphnia* (hereafter "recipients') were exposed

to either infected hosts (hereafter "donors") or their homogenized remains (Table 1). Briefly, in

experiment 1 we investigated whether *Spirobacillus* is an obligate killer by exposing hosts to

either live infected animals, the homogenate of an infected animal or water in which infected

animals had died. In experiments 2 and 3, we respectively investigated whether pathogen dose

Exp.	Question(s)	Independent variable (treatments)	Pathogen dose (# host's worth material)	Duration exposed	Duration monitored (block)
1	Is *Spirobacillus* an obligate killer? What is the impact of infection on host fitness?	Status of donor (alive, dead, crushed, unexposed)	1	6hr	21d
2	Is *Spirobacillus'* transmission rate dose-dependent?	# donors per beaker (2, 3, unexposed)	0, 2, 3	continuous	11d *(a)* 14d *(b)*
3	Does *Spirobacillus'* transmission rate change with host density?	Density of hosts per beaker (1, 2)	1	continuous	10d *(a, b)* 12d *(c, d)*
4	How does *Spirobacillus'* infectiousness change following host destruction?	Time since donor death (0, 1, 2, 4, 6d)	0.5	24hr	14d

Table 1: Focal questions addressed by individual-scale experiments. Exp, experiment; d, days; hr, hours. "continuous" exposure means that animals remained in infectious slurry for the experiment's duration.

and host density, two variables that commonly alter infectious disease transmission (26, 27),

impact the per capita rate of *Spirobacillus* transmission. Lastly, since *Spirobacillus* is a water-

borne pathogen whose transmission dynamics are likely to depend on its survival in the

environment, in experiment 4 we investigated how the infectiousness of *Spirobacillus* varies with

increasing time spent incubating in the environment, in the absence of hosts. Since the

availability of donors was limited by the number of donors in our *in vivo* cultures on the day of

experimental setup, the sample size of each individual experiment was limited and some

experiments were necessarily performed in several blocks. Each of these individual-scale

experiments shared a similar format and schedule, as follows.

Hosts & parasites

"Recipient" hosts were 5-6 day old *Daphnia dentifera* of the L6D9 clone, which was originally

collected from Dogwood Lake (Greene-Sullivan State Forest, Indiana, USA). Infected, 'donor'

hosts were harvested from *in vivo* cultures that we maintained in the Duffy Lab at the University

of Michigan from February 2016 until the spring of 2020, when the COVID-19 pandemic caused

their loss. These cultures originated from infected animals also isolated from Dogwood Lake.

We identified donor hosts by their opacity and red coloration, which signifies the terminal stage

of infection (19).

Experimental exposure & monitoring

To expose recipients to *Spirobacillus,* we either added donors or their homogenized remains to

the water in which recipients were housed (Table 1). Specifically, on the first day of the

experiment, denoted day 0, we placed recipients in 50mL beakers filled with 25mL filtered lake

water. In experiment 1, we exposed recipients to live donors, the remains of hosts that had died

the previous day, or a "slurry" made of homogenized hosts. To make the slurry, we

homogenized donor hosts in MilliQ water using a motorized pestle and brought the solution up

to a known volume. We then mixed the "donor-slurry" thoroughly and added a fraction of the

slurry to each beaker so that each beaker received ~ one donor host's worth of putatively

infectious material. In addition, we included an "unexposed" treatment that was treated the

same way except that we did not add live donors, dead infected hosts or crushed hosts to the

beakers.

In experiments 2-4, we exposed recipients via donor slurry only. We introduced a

volume of slurry equivalent to between 0.5 to 3 donor host's worth of infectious material

(depending on the experiment, cf. Table 1). We varied parasite inoculum in this way because

methods to quantify the amount of *Spirobacillus* in a sample via e.g., microscopy or CFU have

not been developed. In experiment 2, we also included an "unexposed" treatment that did not

receive any donor slurry.

In experiment 4, we sought to understand how environmental conditions – independent

of hosts – alter the infectiousness of *Spirobacillus* over time. Accordingly, we added donor-slurry

to each beaker before the recipient was added. We then placed the water-slurry solution in an

incubator at 22°C, until a predetermined interval (0, 1, 2, 4 or 6 days) had elapsed, at which

point we added a recipient.

In experiments 2 & 3, we exposed recipients for the duration of the experiment. In experiments

1 & 4, we exposed recipients for 6 and 24 hours respectively, after which time they were

removed, rinsed in filtered lake water and placed in clean filtered lake water. We chose an

exposure period of 6-hours in Experiment 1 based on preliminary data on mean survival time of

symptomatic, *Spirobacillus*-infected hosts (Fig. S1). These data suggested that greater than

50% of alive donors would remain alive during the exposure period. Nevertheless, 48 of 79

donors in the alive treatment died during the exposure period; none of the recipients became

infected, however.

We monitored recipients daily for the above-described symptoms of infection (19). Although our

data (cf. Results) suggests that hosts that appear asymptomatic after a week or more are not

infected, we cannot exclude the possibility that "uninfected" hosts became infected but

subsequently cleared the infection. As such, our experiments are best interpreted as evidence

of the effects of experimental treatment on the probability that symptomatic *Spirobacillus*-

infection is observed.

In experiments 1, 2 and 4, we allowed infection to proceed until host death; we hence use these

experiments to quantify the impact of infection on survival. In experiment 3, recipients were

preserved for subsequent analyses (not presented here) as soon as they became symptomatic.

To quantify the impact of infection on fecundity, we removed and counted the number of

offspring produced by each host in experiment 1, three times a week, and in block b of

experiment 2 (daily). In all other experiments, we removed offspring from beakers daily but did

not quantify their number.

In experiment 1, we preserved recipients that died from an unknown cause or were alive

on the last day of the experiment in 1.5mL tubes filled with 180μL TE buffer and froze them at -

20°C. To assess whether asymptomatic infection occurred, we assessed a subset of the stored,

asymptomatic hosts for infection by PCR, as previously described (19).

Husbandry

All experiments were conducted under a 16-8 hour light-dark cycle at 22°C. Animals were fed

250μl of a 1 million cells/ml solution of *Ankistrodesmus falcatus* daily (yielding 10,000 cells/mL

food), with the exception of experiment 4 wherein animals were fed 500μl (20,000 cells/mL) per

day.

Mesocosm-scale experiments

We supplemented our individual-level experiments with investigations of the dynamics of

Spirobacillus infection in mesocosms, akin to those used to maintain infection in the laboratory.

All experiments were maintained under a 16-8 hour light-dark cycle at 22°C and fed 20ml of

algal solution (1 million cells/mL *A. falcatus*), three times a week.

To complement experiment 4, and to assess the extent to which mesocosms contain infectious

material as time elapses, we performed an experiment of the following design. We added 80

recipients, denoted the `primary cohort', to 1L beakers containing 800ml filtered lake water. We

then added 4 donors to this mesocosm. After a predefined interval (1, 3, or 5 days), we removed

the primary cohort to a secondary beaker containing filtered lake water and added a cohort of

80 `secondary' recipients to the original mesocosm. We monitored whether any infected animals

appeared in the primary and secondary cohorts for 15 days following their initial exposure to the

water in the original beaker. Accordingly, we used the 'primary' cohort to indicate whether the

focal mesocosm was *ever* infectious, and the secondary cohort to indicate whether any

infectious material remained in the beaker after the time interval had lapsed. We ran 3 replicates

of each of the 1 and 3-day treatments and 4 of the 5-day treatment.

To quantify the dynamics of infections in *in vivo* cultures, we monitored the prevalence of

infection (as indicated by symptomatic animals) and total host density in 1L mesocosms. On day

0, we filled each 1L mesocosm with 800ml filtered lake water. We added 75 recipients to each

mesocosm. To estimate the density of animals and prevalence of infection in mesocosms, we

took 20 x 20ml samples on each sampling day using a handmade scoop. We demonstrated in

preliminary experiments that the estimate of infection prevalence stabilizes at or before 20

samples were taken. We temporarily retained each sample, counted the number of symptomatic

and asymptomatic animals and then stirred the mesocosm before taking the next sample. After

all samples were taken, we replaced the water and animals back into mesocosm.

Quantification of *Spirobacillus* in field-collected ephippia

To assess whether *Spirobacillus* DNA was present inside ephippia, we collected ephippia from

the field and quantified *Spirobacillus* via ddPCR.

In May and November 2019, we collected sediment from Pickerel Lake (Washtenaw County,

Michigan, USA), wherein *Spirobacillus* infected animals have been observed over several years

(28). To separate the ephippia from the sediment, we placed small portions of the sediment (~4

grams) on a 243μm plankton net filter and rinsed it with deionized water (diH$_2$O) to remove

small particles and liquid reside. We transferred the material retained on the filter to a watch

glass containing diH$_2$O, inspected it under a dissecting microscope and isolated ephippia of

Daphnia dentifera following Rogalski et al. (29). We then surface-sterilized the ephippia by

placing them in 70% ethanol for 2 mins and finally placed each ephippium in a separate 1.5 ml

centrifuge tube filled with PBS buffer. We homogenized the ephippium using a sterile motor-

powered pestle and stored the tubes at -20℃.

Next, we extracted whole genomic DNA from each ephippium using the DNeasy Blood and

Tissue kit (Qiagen®), initiated by adding Proteinase K into the thawed solution and then

following the manufacturer's protocol. Extracted DNA was stored at -20℃ prior to analysis by

digital PCR, as follows. Each reaction contained 20µl of eluted DNA, 10µl of mastermix

(QIAcuity™ Probe PCR Kit), 0.72 µl of forward primer Spiro70F (5'-

AACTGTAGCTAAGACCGCGT-3') and 0.72 µl of reverse primer Spiro210R (5'-

CCAGCTATCCATCTTCGCCT-3'), 0.36 µl of dual-quenched probe SpiroP (5'-

ACCGTAAGGCCAGCAGCCATAAGATGA-3') labeled with fluorescent Dye ROX at the 5' end

and Iowa Black® RQ dark quencher at the 3' end, 0.5 µl of restriction enzyme EcoRI and 13.5 µl

of RNase-free water. We used a high volume of DNA because we expected that the ephippia

would contain a low concentration of bacterial DNA. The PCR was performed in a 24-well

Nanoplate 26K (Qiagen®), using a QIAcuity One Digital PCR System (Qiagen®). PCR

conditions were 2 min at 95℃, followed by 40 cycles of 15 sec at 95℃ followed by 1 min at 55℃.

We first performed post-PCR imaging using an exposure time of 300ms and, when it was

necessary to increase the separation of signal from noise, reimaged the plates using either

200ms or 100ms. The results were analyzed with QIAcuity Software Suite 2.0.20. Finally, we

converted the output – number of gene copies in the reaction mix – to the total copy

number/ephippium. Note that, since we do not know the number of copies of the 16SrRNA

copies in the *Spirobacillus* genome, copy number should not be interpreted as an absolute

estimate of the number of cells contained within each ephippium.

We designed the aforementioned primers and probe using NCBI Primer-BLAST (30) to target

the *Spirobacillus* 16S rRNA gene. We verified the assay's specificity by using it to assay DNA of

three other members of the family *Silvanigrellaceae, Silvanigrella aquatica*, *Silvanigrella*

paludirubra and *Fluviispira multicolorata.* The assay did not amplify DNA from these species.

Statistical analysis

We performed statistical analysis using R (version 4.2.1). To test for an association between

treatment and the probability of infection, we ran Chi-square tests or (when expected values<5)

Fisher's Exact Tests, following (31). To verify the result of these analyses and to obtain

estimates of per capita infection rates (and confidence intervals around said estimates) for

presentation in figures, we ran binomial general linearized models (GLMS) and (where

necessary) controlled for block. In all case, the two approaches yielded the same results *vis-a-*

vis the significance of treatment. We removed asymptomatic animals that died on or before day

8 – the last day on which we ever observed symptoms – from the analysis of transmission rates,

reasoning that these animals may not have had sufficient time to display symptoms.

To quantify the impact of infection on fecundity, we used Poisson GLMs. We corrected for

overdispersion, which was found in all cases following (32) i.e., using a quasi-Poisson model.

Animals whose fecundity was not recorded throughout their lifetime were removed from this

analysis.

To analyze the effect of treatment on time to death and symptoms, we employed survival

analyses using the *surv* package (33). These analyses allow us to include all the individual-level

time series data, including data relating to animals that died before day 9 or which were not

tracked throughout the entirety of a time-period. We fit Kaplan-Meier survival curves to time-

series data *vis-à-vis* host death and the appearance of symptoms and used them to derive

estimates of the median (+/- 95% CI) time-to -death and -symptoms, respectively. To ask

whether *Spirobacillus* had a significant impact on lifespan, we used log rank tests to compare

the survival curves. In addition, in Figure 1b we plot the results of (i) a "competing risks"

analysis, which we used to quantify the probability that an animal either becomes symptomatic

or dies without showing symptoms (dashed line), and (ii) a "multi-state" analysis, which we used

to quantify the probability that an animal died following the appearance of symptoms (black line)

(34).

Results

<u>Virulence</u>

Symptomatic *Spirobacillus* infection ("infection", hereafter) had a 100% fatality rate in our

individual-level experiments (Fig. 1a). Symptoms developed rapidly – on average, by day 6 –

and were followed by death within a day, on average (Fig. 1b; Fig. S1, restricted mean survival

time of a symptomatic animal 9.6 hours). As a result, *Spirobacillus* infection had a significant

impact on host lifespan (χ^2 = 290, df = 1, p<0.001). In experiment 1, the median survival time of

infected animals fell from over 21 days (i.e., longer than the duration of the experiment) in

infected, but exposed, animals to to 7 days (95% CI 7-8) in uninfected animals.

Spirobacillus infection significantly reduced the fecundity of infected animals, as compared to

uninfected treatment-mates (Fig. 2a, *infection status* δAIC= 187, df=1). Over the course of

experiment 1, for example, infected animals had ten times fewer offspring than uninfected

animals in the same treatment. The effect of *Spirobacillus* on host fecundity does not seem to

stem only from its effect on host lifespan. During the first week of experiments 1 and 2**,**

recipients that later became symptomatic had fewer offspring than those that did not become

infected (Fig. 2b). Thus, the negative effects of *Spirobacillus* infection on host fitness are

apparent even before symptoms are detectable.

<u>Transmission among asexually reproducing *Daphnia*</u>

Spirobacillus required host death or destruction to transmit (Fig. 3). In experiment 1,

symptomatic infection was only observed in animals exposed to material from donors that had

been destroyed by homogenization or had died naturally (Fig. 3). To assess whether

asymptomatic infections may have arisen, 46 asymptomatic animals (including 20 that were

exposed to alive donors) were assessed for infection via PCR. None were positive (data not

shown). These results suggest that *Spirobacillus* is an obligate killer and that asymptomatic

infection does not occur.

Across all individual-scale experiments, the mean per capita probability of infection did not

exceed 50% in any treatment (Figs. 3-5). The per capita probability of infection was not

impacted by pathogen dose (Fig. 4a, p= 0.7) or the density of animals in the beaker (Figure 4b,

p = 0.2).

The infectiousness of *Spirobacillus* declined with time spent in water in the absence of hosts

(Fig. 5*, time since host destruction* χ^2 = 31, df =1, p<0.001), in experiment 4 were we made

observations at the individual scale. *Spirobacillus* that was not incubated in the environment for

any period generated infected hosts at a rate of 20**%**. Our data suggests that, after just 3 days

of incubation, the probability that a host became infected fell to <1% (Fig. 5). We observed a

similar decline in infectivity in mesocosm-scale experiments (where, unlike in experiment 4, we

cannot disentangle the effect of hosts and the environment on *Spirobacillus'* infectivity). None of

3 mesocosms that generated infections in the "primary cohort" were infectious 5 days after

inoculation and only 1 (of 3) generated any infections 3 days after inoculation.

We did not observe classical epidemic dynamics in mesocosms (Fig. 6). With the exception of 1

mesocosm, the prevalence of symptomatic hosts cycled up and down at a level below or

equivalent to the initial prevalence. In no mesocosm did the prevalence of symptomatic hosts

ever grow for a period longer than the incubation period of 7 days (which, given that

Spirobacillus is an obligate killer, is equivalent to the time-to-death of symptomatic hosts as

estimated above). Furthermore, the density of symptomatic hosts did not lag behind that of

asymptomatic hosts, as is expected (21); indeed, the dynamics of these two cohorts bear scant

relationship to one another.

Presence within ephippia

We found *Spirobacillus* DNA in 39% of the 41 *Daphnia dentifera* ephippium assayed. Positive

ephippia contained 83 copies of the 16srRNA gene, on average (range 4 to 860) (Figure 7).

Discussion

Despite *Spirobacillus cienkowksii's* long history of study and widespread distribution,

fundamental aspects of its biology have remained uncharacterized. Our experiments reveal that

Spirobacillus is among the most virulent of *Daphnia* pathogens yet described (33–35), killing its

host almost twice as rapidly as *Metschnikowia bicuspidata* (35), the pathogen of *Daphnia*

dentifera with the next most rapid measured impact on host death. In addition to killing hosts

rapidly, *Spirobacillus* reduces host fitness by reducing host fecundity. *Spirobacillus'* impact on

host reproduction is equivalent to that of another common parasite of *Daphnia*, the castrator

Pasteuria ramosa, but *Spirobacillus'* effect likely comes about via a different mechanism.

Pasteuria sterilizes hosts weeks before it kills them, presumably via chemical manipulation of

the host (36). By contrast, *Spirobacillus* primarily impacts offspring production by reducing host

lifespan, though it also reduces fecundity prior to their early death. Given that we have

observed field- and laboratory- infected *Daphnia* birth on the day they die (data not shown), we

think it unlikely that *Spirobacillus* directly interferes with host reproductive functions but rather

reduces host reproduction by inducing e.g., nutritional stress.

We demonstrated that *Spirobacillus'* effects on mortality are intimately connected with its

capacity to transmit horizontally. Specifically, in experiment 1, we showed that *Spirobacillus*

requires the death of its host to transmit horizontally consistent with the hypothesis that it is an

obligate killer. Given that mechanical destruction of the host permitted infection, host death may

not need to be induced by the pathogen, however. Rather, destruction of hosts by "sloppy"

predators may facilitate transmission (37, 38). Though our experiments were not explicitly

designed to explore the timescale of transmission following host death, the "alive" treatment of

experiment 1 reveals something about it. Several of the "alive" donors died within the 6-hour

exposure period, yet none of the recipients became infected. This implies that either there is a

lag between host death and the production of viable transmission stages (unlikely given that

crushing hosts yielded infections) or, more likely, that it takes some time for motile pathogens to

escape and find a new host.

In these laboratory conditions, the per capita rate at which susceptible hosts become infected

(or the force of *Spirobacillus* infection (FOI)) was low and was not influenced by two ecological

factors that are expected to change FOI. FOI is a product of a transmission constant, the

density of infected hosts or pathogen in the environment, and susceptible host density (27). Yet

Spirobacillus' FOI did not increase with pathogen dose in experiment 2, nor between experiment

3 (where recipients were exposed to 1 host worth of parasite) and experiment 2 (where

recipients were exposed to 2-3 host's worth of parasite). This behavior contrasts markedly with

that of *Pasteuria ramosa*, for example. In *D. magna* housed in similar experimental conditions,

only a small fraction (~0.001%) of the *Pasteuria* spores contained in a fully infected host is

sufficient to infect ~50% of new hosts, and a dose of ~10% of the spores in a fully infected host

all but guarantees infection ((39–41)(42). Our data are not sufficient to conclude that there is no

relationship between pathogen dose and *Spirobacillus'* FOI, however. Dose-response

relationships are rarely linear, and, instead, tend to be decelerating (42, 43). One hypothesis for

our observations, therefore, is that we were working in the decelerating region of the dose

response curve in experiments 2 and 3 and that had we explored a larger range of smaller

doses, we would have observed a relationship between inoculum size and FOI. Similarly, there

may well be a relationship between host density and FOI, that went unobserved because of

experiment 3's limited sample size and/or range of treatments. Indeed, that new infections arose

in our mesocosms – which contained a higher density (~2.5x) of susceptible hosts but were

initiated with a much smaller inoculum than our individual-scale experiments – implies that

density-dependence may play a role in *Spirobacillus'* transmission dynamics. Ironically, we were

unable to explore a wide range of density and dose treatments or boost the power of our

experiments via greater replication because the poor transmission of *Spirobacillus* in *in vivo*

cultures limited the availability of donors.

Spirobacillus' limited environmental persistence, as demonstrated in experiment 4, combined

with its low force of infection, likely explain why we did not observe epidemics in *in vivo* cultures.

The extent to which environmentally transmitted pathogens can persist in the environment is a

critical determinant of their fitness and epidemiological dynamics (22, 25, 44–46). For obligate

killers like *Spirobacillus*, whose survival in the environment (at least in our experiments) does

not exceed the timescale of host population dynamics, the ability to invade a host population

increases with transmission rate and virulence, and declines as the decay rate of infectious

stages increases (47). So, although *Spirobacillus* is highly virulent, its low FOI and rapid rate of

decay in the environment may provide a "break" on its ability to cause epidemics in the

laboratory. Instead, we observe peaks and troughs in the prevalence of infection, with a

periodicity of ~6 days (at least at the beginning of infections, when small differences among

individuals have not accumulated to dampen the cycles). We hypothesize that these dynamics

are a result of the strong, relatively invariant periodicity of the time to symptoms (6-7 days) and

time to death (7-8 days), combined with *Spirobacillus'* limited environmental persistence.

Why, then, does *Spirobacillus* generate epidemics in the field (and not in the lab)? We have two

hypotheses. Our first assumes that *Spirobacillus* is an obligate parasite and attributes our

findings to a discrepancy between the conditions in the lab and those under which transmission

occurs in nature. For example, in a study of *Spirobacillus* in the water-column, Thomas et al.

(48) found the highest density of *Spirobacillus* at the surface-water interface, a region

characterized by low temperatures, oxygen and light (49, 50). If this region is the location of

transmission in nature, our experiments poorly represent them. Alternatively, it is possible that

our recipient hosts were less susceptible than the hosts that become infected in nature e.g.,

because of their age or genotype or nutritional status. Given that we have observed infection

across host ages in the field and have generated infections at equivalent rates in other hosts

clones (unpublished data), we think that differences in the abiotic conditions of the lab vs. the

field are more likely to explain our inability to recover epidemic dynamics in the lab, than

differences in their biotic conditions.

Our second, more heterodox, hypothesis is that *Spirobacillus* is a facultative pathogen of

Daphnia, whose epidemic dynamics are not entirely a function of replication within asexually

reproducing *Daphnia*. If our experiments do reflect *Spirobacillus* behavior in nature, it is unlikely

that it could make a living as an obligate pathogen because of its poor horizontal transmission

rate. Instead, it is possible that *Spirobacillus* primarily replicates on particles (including ephippia)

in the water column and/or sediment and employs the same traits that allow for particle-finding, -

attachment and -exploitation to parasitize hosts, when the opportunity arises. In accordance

with this hypothesis, *Spirobacillus* and the other *Silvanigrellaceae* exhibit a suite of traits,

including pleiomorphism, small size and the genomic capacity to make flagella and perform

chemotaxis (20), that are characteristic of copiotrophic aquatic microbes capable of facultative

predation or parasitism, e.g., the *Vibrios* and *Bdellovibrios* (51, 52)*.* Under this model, either (i)

Spirobacillus outbreaks in *Daphnia* result from "blooms" in the particle-associated population

that spillover into the *Daphnia* population, or (ii) there is an ecological cue that stimulates a

switch from a free-living or particle associated lifestyle to a *Daphnia*-associated lifestyle.

Comparative genomic analyses of the *Spirobacillus* genome with that of predatory and/or

parasitic species, and investigations of the distribution and abundance of *Spirobacillus* in

different particulate and non-particulate fractions of the water column, could shed further light on

this hypothesis.

Our discovery that *Spirobacillus* is harbored within ephippia is consistent with the idea that

Spirobacillus may not entirely rely on asexually reproducing hosts for its fitness and could help

us to explain a range of observations made in the past and in this study. We found *Spirobacillus*

DNA in almost 40% of the ephippia assayed. This observation, combined with a previous

observation that ephippia inside infected *Daphnia* contain bacteria (11), suggests that

Spirobacillus may be vertically transmitted from sexually reproducing females to ephippia. (We

note that vertical transmission would include bacteria that are contained within the ephippium

but not the embryo itself.) That said, these data do not exclude the possibility that ephippia

become colonized by *Spirobacillus* after they had been released from *Daphnia* and, given that

Spirobacillus' prevalence rarely exceeds 10% even during epidemics, its comparatively high

prevalence in ephippia (39%) is difficult to explain with respect to vertical transmission alone.

Irrespective of the mechanism by which *Spirobacillus* enters the ephippia, the ability to exploit

ephippia could explain *Spirobacillus'* inter-epidemic persistence and hypervirulence. Rodrigues

et al. observed that *Spirobacillus*-infected hosts hatch directly from ephippia (11). This

observation, coupled within our findings, suggests that *Spirobacillus* might use the hardy

ephippia to persist through the winter, before emerging from them and seeding the next year's

epidemics. Given the high energetic costs of sporulation (53)*,* invading a hardy structure rather

than making one could have significant advantages. The capacity to survive within ephippia

might also allow *Spirobacillus* to offset or avoid paying the costs of virulence, if it is an obligate

pathogen, per the "curse of the pharaoh" hypothesis (23).

Our work reveals much about the life history of a bacterium that has a long and significant

history in bacteriology and sets a foundation for the use of *Spirobacillus* as a model of host-

parasite interactions at the individual host and evolutionary timescale. Carotenoid production in

Spirobacillus was first described in 1959 at the dawn of carotenoid chemistry (12, 54), and

bacterial carotenoids have since received significant attention in biotechnology and medicine,

wherein carotenoids are considered virulence factors (55). Here, we found that *Spirobacillus*

infection is virulent, reliably symptomatic and proceeds from initiation to death within a week.

These features, coupled with our prior description of natural variation in the color of infected

hosts (and thence, presumably, carotenoid production) (19), could make *Spirobacillus* a

tractable, *in vivo* model with which to study how bacterial secondary metabolites alter infection

dynamics and virulence. Future work to identify conditions under which infection can be

guaranteed will further boost its tractability in this context. Indeed, future work aimed at

elucidating the mode(s) of *Spirobacillus* transmission could help us to make new insights about

the evolutionary dynamics of virulence, as well as carotenoid's role in it. The dynamics of

virulence evolution in pathogens capable of both horizontal and vertical (i.e., "mixed-mode")

transmission and/or environmental replication are poorly understood as compared to pathogens

with strictly horizontal and/or direct transmission (56); virulence evolution in obligate killers is

also relatively poorly studied. Our work suggests that mixed-mode transmission and/or

environmental replication is possible in *Spirobacillus*. Better understanding these aspects of

Spirobacillus' biology could help us to build and test ideas about how obligate killers evolve

when their fitness does not rely solely on horizontal transmission. In conclusion, by establishing

Spirobacillus in the laboratory for the first time, we established critical facets of *Spirobacillus'*

infection biology that can be harnessed by biomedical biologists and ecologists alike.

Acknowledgements

We thank Libby Davenport (U. Michigan) for help designing the dPCR assay and Martin Hahn

(U. Innsbruck) for the samples of *Silvanigrellaceae* that we used to validate said assay.

Bibliography

1. Duncan AB, Little TJ. 2007. Parasite-driven genetic change in a natural population of
Daphnia. Evolution 61:796–803.

2. Duffy MA, Ochs JH, Penczykowski RM, Civitello DJ, Klausmeier CA, Hall SR. 2012.
Ecological context influences epidemic size and parasite-driven evolution. Science 335:1636–
1638.

3. Duffy MA, Brassil CE, Hall SR, Tessier AJ, Cáceres CE, Conner JK. 2008. Parasite-mediated
disruptive selection in a natural Daphnia population. BMC evolutionary biology 8:80.

4. Duffy MA, Hall SR, Tessier AJ, Huebner M. 2005. Selective predators and their parasitized
prey: Are epidemics in zooplankton under top-down control? Limnol Oceanogr 50:412–420.

5. Johnson PTJ, Stanton DE, Preu ER, Forshay KJ, Carpenter SR. 2006. Dining on disease: how
interactions between infection and environment affect predation risk. Ecology 87:1973–1980.

6. Wale N, Fuller RC, Johnsen S, Turrill ML, Duffy MeghanA. 2021. The visual ecology of
selective predation: Are unhealthy hosts less stealthy hosts? Ecol Evol
https://doi.org/10.1002/ece3.8464.

7. Ebert D. 2008. Host–parasite coevolution: Insights from the *Daphnia*–parasite model system.
Curr Opin Microbiol 11:290–301.

8. Ebert D. 2022. Daphnia as a versatile model system in ecology and evolution. Evodevo 13:16.

9. Hahn MW, Pitt A. 2017. Silvanigrella, p. 1–9. *In* Bergey's Manual of Systematics of Archaea
and Bacteria.

10. Metchnikoff E. 1889. Contributions a l'etude du pleomorphisme des bacteriens. Annales de
l'Institut Pasteur 3:61–68.

11. Rodrigues JLM, Duffy MA, Tessier AJ, Ebert D, Mouton L, Schmidt TM. 2008.
Phylogenetic characterization and prevalence of Spirobacillus cienkowskii, a red-pigmented,
spiral-shaped bacterial pathogen of freshwater Daphnia species. Applied and Environmental
Microbiology 74:1575–1582.

12. Green J. 1959. Carotenoid pigment in Spirobacillus cienkowskii Metchnikoff, a pathogen of
Cladocera. Nature 183:56–57.

13. Mioduchowska M, Czyż MJ, Gołdyn B, Kilikowska A, Namiotko T, Pinceel T, Łaciak M,
Sell J. 2018. Detection of bacterial endosymbionts in freshwater crustaceans: the applicability of
non-degenerate primers to amplify the bacterial 16S rRNA gene. Peerj 6:e6039.

14. Hall SR, Duffy MA, Cáceres CE. 2005. Selective predation and productivity jointly drive
complex behavior in host-parasite systems. The American Naturalist 165:70–81.

15. Duffy MA, Hall SR. 2008. Selective Predation and Rapid Evolution Can Jointly Dampen
Effects of Virulent Parasites on Daphnia Populations. Am Nat 171:499–510.

16. Peng N, Xu W, Luo H-M, Ouyang P-W, Jiang B, Zhang Z-H, Cao Y-D, Tan L-M, Qu P-H,
Xie L-Y. 2021. Pigmentibacter ruber gen. nov., sp. nov., a novel bacterium of the family
Silvanigrellaceae isolated from human blood. Antonie Van Leeuwenhoek 1–9.

17. Walke JB, Becker MH, Hughey MC, Swartwout MC, Jensen RV, Belden LK. 2015. Most of
the Dominant Members of Amphibian Skin Bacterial Communities Can Be Readily Cultured.
Appl Environ Microb 81:6589–6600.

18. Anderson RM, May RM. 1979. Population biology of infectious diseases: Part I. Nature
280:361–367.

19. Wale N, Turrill ML, Duffy MA. 2019. A colorful killer: Daphnia infected with the bacterium
Spirobacillus cienkowskii exhibit unexpected color variation. Ecology 100.

20. Bresciani L, Lemos LN, Wale N, Lin JY, Strauss AT, Duffy MA, Rodrigues JLM. 2018.
Draft genome sequence of "Candidatus Spirobacillus cienkowskii," a pathogen of freshwater
Daphnia species, reconstructed from hemolymph metagenomic reads. Microbiology Resource
Announcements 7.

21. Cortez MH, Weitz JS. 2013. Distinguishing between Indirect and Direct Modes of
Transmission Using Epidemiological Time Series. Am Nat 181:E43–E52.

22. Turner WC, Kamath PL, Heerden H van, Huang Y-H, Barandongo ZR, Bruce SA, Kausrud
K. 2021. The roles of environmental variation and parasite survival in virulence–transmission
relationships. Roy Soc Open Sci 8:210088.

23. Gandon S. 1998. The curse of the pharoah hypothesis. Proc Royal Soc Lond Ser B Biological
Sci 265:1545–1552.

24. Sánchez KF, Zhong B, Agudelo JA, Duffy MA. 2023. Infectivity of the parasite
Metschnikowia bicuspidata is decreased by time spent as a transmission spore, but exposure to
phycotoxins in the water column has no effect. Freshwater Biol 68:1020–1030.

25. Rumschlag SL, Roth SA, McMahon TA, Rohr JR, Civitello DJ. 2022. Variability in
environmental persistence but not per capita transmission rates of the amphibian chytrid fungus
leads to differences in host infection prevalence. J Anim Ecol 91:170–181.

26. McCallum H, Fenton A, Hudson PJ, Lee B, Levick B, Norman R, Perkins SE, Viney M,
Wilson AJ, Lello J. 2017. Breaking beta: deconstructing the parasite transmission function.
Philosophical Transactions Royal Soc B Biological Sci 372:20160084.

27. Bjørnstad ON. 2023. Epidemics, Models and Data Using R. R https://doi.org/10.1007/978-3-
031-12056-5.

28. Gowler CD, Rogalski MA, Shaw CL, Hunsberger KK, Duffy MA. 2021. Density, parasitism,
and sexual reproduction are strongly correlated in lake Daphnia populations. Ecol Evol
11:10446–10456.

29. Rogalski MA, Leavitt PR, Skelly DK. 2017. Daphniid zooplankton assemblage shifts in
response to eutrophication and metal contamination during the Anthropocene. Proc Royal Soc B
Biological Sci 284:20170865.

30. Ye J, Coulouris G, Zaretskaya I, Cutcutache I, Rozen S, Madden TL. 2012. Primer-BLAST:
A tool to design target-specific primers for polymerase chain reaction. Bmc Bioinformatics
13:134.

31. Field A, Miles J, Field Z. 2012. Discovering Statistics Using R. SAGE Publications Ltd.

32. Zuur AF, Ieno EN, Walker NJ, Saveliev AA, Smith GM. 2009. Mixed Effects Modelsand
Extensions in Ecology with R.

33. Therneau TM. 2023. A Package for Survival Analysis in R. https://CRAN.R-
project.org/package=survival.

34. Therneau T, Crowson C, Atkinson E. Multi-state models and competing risks. https://cran.r-
project.org/web/packages/survival/vignettes/compete.pdf. Retrieved 2022.

35. Clay PA, Dhir K, Rudolf VHW, Duffy MA. 2019. Within-Host Priority Effects
Systematically Alter Pathogen Coexistence. Am Nat 193:187–199.

36. Clerc M, Ebert D, Hall MD. 2015. Expression of parasite genetic variation changes over the
course of infection: implications of within-host dynamics for the evolution of virulence.
Proceedings of the Royal Society of London, Series B: Biological Sciences 282:20142820–
20142820.

37. Duffy MA, Cáceres CE, Hall SR. 2019. Healthy herds or predator spreaders? Insights from
the plankton into how predators suppress and spread disease, p. 458–479. *In* Wilson, K, Fenton,
A, Tompkins, & D (eds.), Wildlife Disease Ecology: Linking Theory to Data and Application.
Cambridge University Press.

38. Cáceres CE, Knight CJ, Hall SR. 2009. Predator–spreaders: Predation can enhance parasite
success in a planktonic host–parasite system. Ecology 90:2850–2858.

39. Ben-Ami F, Regoes RR, Ebert D. 2008. A quantitative test of the relationship between
parasite dose and infection probability across different hostparasite combinations. Proc Royal
Soc B Biological Sci 275:853–859.

40. Ben-Ami F, Ebert D, Regoes RR. 2010. Pathogen Dose Infectivity Curves as a Method to
Analyze the Distribution   of Host Susceptibility: A Quantitative Assessment of Maternal
Effects   after Food Stress and Pathogen Exposure. Am Nat 175:106–115.

41. Regoes RR, HOTTINGER JW, SYGNARSKI L, EBERT D. 2003. The infection rate of
Daphnia magna by Pasteuria ramosa conforms with the mass-action principle. Epidemiology
Amp Infect 131:957–966.

42. Lunn TJ, Restif O, Peel AJ, Munster VJ, Wit E de, Sokolow S, Doremalen N van, Hudson P,
McCallum H. 2019. Dose–response and transmission: the nexus between reservoir hosts,
environment and recipient hosts. Philosophical Transactions Royal Soc B Biological Sci
374:20190016.

43. Clay PA, Cortez MH, Duffy MA. 2021. Dose relationships can exacerbate, mute, or reverse
the impact of heterospecific host density on infection prevalence. Ecology 102:e03422.

44. Overholt EP, Duffy MA, Meeks MP, Leach TH, Williamson CE. 2020. Light exposure
decreases infectivity of the *Daphnia* parasite *Pasteuria ramosa*. J Plankton Res 42:41–44.

45. Li S, Eisenberg JNS, Spicknall IH, Koopman JS. 2009. Dynamics and Control of Infections
Transmitted From Person to Person Through the Environment. Am J Epidemiol 170:257–265.

46. Ogbunugafor CB, Miller-Dickson MD, Meszaros VA, Gomez LM, Murillo AL, Scarpino
SV. 2020. Variation in microparasite free-living survival and indirect transmission can modulate
the intensity of emerging outbreaks. Sci Rep-uk 10:20786.

47. Tate AT. 2016. The Interaction of Immune Priming with Different Modes of Disease
Transmission. Front Microbiol 7:1102.

48. Thomas S, Bertram C, Rensburg K van, Cáceres C, Duffy M. 2011. Spatiotemporal
dynamics of free-living stages of a bacterial parasite of zooplankton. Aquat Microb Ecol 63:265–
272.

49. Wetzel RG. 2001. Light in Inland Waters, p. 49–69. *In* Third. Elsevier.

50. Morris RL, Schmidt TM. 2013. Shallow breathing: bacterial life at low O2. Nat Rev
Microbiol 11:205–212.

51. Sockett RE. 2009. Predatory Lifestyle of Bdellovibrio bacteriovorus. Microbiology+ 63:523–
539.

52. Lutz C, Erken M, Noorian P, Sun S, McDougald D. 2013. Environmental reservoirs and
mechanisms of persistence of Vibrio cholerae. Front Microbiol 4:375.

53. O'Hara MB, Hageman JH. 1990. Energy and calcium ion dependence of proteolysis during
sporulation of Bacillus subtilis cells. J Bacteriol 172:4161–4170.

54. Yabuzaki J. 2017. Carotenoids Database: structures, chemical fingerprints and distribution
among organisms. Database 2017:bax004.

55. Liu GY, Nizet V. 2009. Color me bad: microbial pigments as virulence factors. Trends
Microbiol 17:406–413.

56. Pandey A, Mideo N, Platt TG. 2022. Virulence Evolution of Pathogens That Can Grow in
Reservoir Environments. Am Nat 199:141–158.

Fig 1. *Spirobacillus* kills its host shortly after symptoms appear, dramatically reducing host lifespan. a) Mean probability of survival (+/- 95% confidence interval, shaded area) following exposure to *Spirobacillus* of infected (red) or uninfected (black) hosts in experiments 1-4. Estimates derived from a Kaplan-Meier survival curve. Crosses indicate when animals were removed from the experiments for a reason other than death e.g., the experiment ended or, in experiment 3, when infected animals were preserved before death. b) Estimated probability that a host was symptomatic (dashed line) and dead after displaying symptoms (solid line) following exposure to *Spirobacillus*, across experiments 1, 2 and 4. The estimated probabilities take such small values because most of the animals exposed in these experiments did not become infected (cf. Fig. 3, 4a, 5)

Fig 2. *Spirobacillus* **infection reduces the fecundity of infected** *Daphnia,* **even during the short period that they are alive**. a) Cumulative no. of offspring produced per host over the course of experiments 1 and 2. b) Total offspring produced in the first week post exposure (p.e.), by hosts that went on to be infected (red) or remained uninfected (white). Note that animals were exposed while juveniles, hence the limited reproduction between days 0-2. In both plots, each individual point represents data from an individual host.

Fig. 3 *Spirobacillus* requires the demise of its host to transmit horizontally. Mean infection rates (+/- standard error) in each treatment of experiment 3, as estimated from a generalized linear model.

Fig. 4 The per capita transmission rate of *Spirobacillus* is not altered by inoculum size or recipient host density. Mean infection rates (+/- standard error) in a) experiment 2 and b) experiment 3, as calculated from a generalized linear model specifying treatment as the sole predictor variable. Smaller shaded points indicate the proportion infected in different experimental blocks. Inclusion of block in the statistical model does not change the significance of treatment in either case. The 100% infection rates in blocks a and d of a) and b), respectively are likely an artefact of small sample size (n = 2 in each case). The pathogen dose in panel a is given in terms of the number of donor hosts' worth of material.

Fig. 5 The infectivity of *Spirobacillus* declines rapidly with time. Mean proportion of animals infected (red line), +/- standard error (grey shading). Quantities estimated from a generalized linear model with time since donor death as the sole predictor variable. Points indicate the proportion infected in the indicated experimental block. Neither the statistical significance of time since donor death nor the trend indicated changes with the inclusion of block in the model.

Fig. 6 Dynamics of *Spirobacillus* infection in laboratory mesocosms. The prevalence of symptomatic hosts (blue), and the density of asymptomatic host (black) and symptomatic hosts (red), in 800ml mesocosms over 28 days. Asterisks indicate the initial value of parameters in day 0, when the mesocosm were established. Error bars around prevalence estimated from a binomial distribution.

Fig. 7. *Spirobacillus* is present in field-collected *Daphnia dentifera* ephippia (resting eggs). Quantity of *Spirobacillus* rRNA gene copies within individual ephippia collected from a Michigan lake where *Spirobacillus* epidemics have previously been observed.