Poly ADP-ribosylation regulates *Arc* expression and promotes adaptive stress-coping

Eliyahu Dahan[1], Tze'ela Taub[1], Arthur Vovk [1], Jade Manier [1], Raphael Avneri[1], Elad Lax[1,*]

[1]Department of Molecular Biology, Ariel University, Ariel, Israel

[*]Corresponding author: eladl@ariel.ac.il

Abstract

Rapid adaptation to stressful events is essential for survival and requires acute stress response and stress-coping strategy. Yet, the molecular mechanisms that govern this coping strategy have to be fully discovered. Poly ADP-ribosylation (PARylation) is a post-translational protein modification that plays a vital role in histone modification following neuronal activity and is required for long-term memory formation. Previous studies showed that inhibition of Parp1, the primary enzyme that catalyzes Poly ADP-ribose polymers (PAR), impaired long-term memory formation. However, the target genes of Parp1-induced PARylation following acute stress are unknown. Here, we showed that the forced swim test, a well-established acute stress paradigm, induced elevated cortical PARylation and increased the expression of activity-dependent genes in mice. Systemic pharmacological Parp1 inhibition with ABT888 (Veliparib), impaired stress-coping in a second forced swim test done 24 hours later as measured by diminished immobility response. This effect was associated with reduced expression of Arc, but not other activity-dependent genes, 1 hour after Parp1 inhibition, suggesting Arc PARylation regulates its expression and successful stress-coping behavior.

1. Introduction

Long-term memory formation of stressful events is an adaptive mechanism enabling the optimization of coping strategies in the future. However, under abnormal circumstances, highly stressful events and persistent reactivation of said memory can lead to undesired outcomes such as avoidance behavior and post-traumatic stress disorder (Ressler et al., 2022). The formation of long-term memories requires *de-novo* protein synthesis preceded by time-dependent

transcriptional alteration (Alberini, 2008; Bisaz, Travaglia, & Alberini, 2014; Dudai & Eisenberg, 2004; Eric R. Kandel, 2001).

Poly ADP-ribosylation (PARylation) is a post-translational protein modification mostly known for its role in DNA repair. PARylation also promotes gene expression, presumably through increasing chromatin accessibility by evicting the linker histone H1 (Azad et al., 2018; Fontán-Lozano et al., 2010). The enzyme Poly (ADP-ribose) polymerase 1 (Parp1) is the most abundant member of the PARP family in mammals (Homburg et al., 2000; Ray Chaudhuri & Nussenzweig, 2017). Parp1 catalyzes the polymerization of ADP-ribose, sourced in nicotinamide adenine dinucleotide (NAD+), to yield linear or branched structured poly ADP-ribose polymers (PAR).

Previous findings by us and others indicate that Parp1 is necessary for forming and retrieving long-term memories of feeding, stress, and cocaine-seeking behaviors (Goldberg, Visochek, Giladi, Gozes, & Cohen-Armon, 2009; Hernández et al., 2009; E. Lax et al., 2017; Scobie et al., 2014). Poly ADP-ribosylation (PARylation) is a rapid and transient posttranslational modification that occurs within 10-15 minutes after depolarization and neuronal activity (Cohen-Armon et al., 2004; Homburg et al., 2000; Visochek et al., 2005). The PARylation of histone H1 regulates chromatin de-condensation and enables memory consolidation. However, it is still not clear which downstream genes are upregulated following histone PARylation. Stressful events that cause neuronal activity trigger the expression of immediate-early and activity-dependent genes required for further neuronal activation, memory formation, and neuroplasticity (Flavell & Greenberg, 2008; McEwen et al., 2015). We hypothesize that a forced swim test (FST), a highly stressful event, will initiate a fast histone PARylation alongside the mRNA expression of immediate-early and activity-dependent genes. We further speculated that systemic pharmacological inhibition of Parp1 before the FST will impair memory formation. This Parp inhibition will also reduce the mRNA expression of some known neuronal activity-related, immediate-early, and activity-dependent genes, thus allowing us to identify the mRNA expression of which of these genes are PARylation-dependent.

2. Methods

2.1 Animals

In the current work, we used 8-9 weeks old C57BL/6J mice (Envigo, Israel). We included equal numbers of both male and female mice for all the experiments. Mice had ad libitum access to standard chow and drinking water and were kept at constant light-dark cycles for 12 hours each, at 22℃.

All animal experiments were approved by the Institutional Animal Care and Use Committee of Ariel University, Israel (approval No. IL-225-07-21). Institutional guidelines were followed for the proper and humane use of animals in research.

2.2 Force Swim Test (FST)

The forced-swim test (FST) was used to induce acute stress. Mice were gently put in a glass cylinder (height: 30 cm, diameter: 10 cm) filled with water (22-33℃) with no option to escape for 6 minutes. Next, the mice were removed from the cylinder and dried in a warm, dry cage for several moments. The mice were euthanized by cervical dislocation at the following time points after FST: 15 min, 60 min, and 24h. Naïve mice not introduced to the FST were used as a control group. Cortical tissues were immediately removed, snapped frozen in liquid nitrogen, and then stored at -80℃ for later use.

2.3 Systemic ABT-888 administration

ABT-888 (Veliparib®) is a selective PARP1 and PARP2 inhibitor with higher efficacy towards PARP1 (Thorsell et al., 2017), which crosses the brain-blood-barrier (Donawho et al., 2007). We dissolved ABT-888 hydrochloride in saline and injected 15 mg/kg of ABT888 intraperitoneally at 0.1 ml volume. The injection took place 50 min before the FST based on pharmacokinetic experiments (Donawho et al., 2007) and as we reported before (Lax et al., 2017). Equal volumes of saline were injected to control mice. For a subset of mice, twenty-four hours after the first FST, a second FST was done to assess changes in the duration of immobility as a measure of the adaptive stress response as described before (Gutièrrez-Mecinas et al., 2011; Reul, 2014; Saunderson et al.,

2016; West, 1990). The experiment was video-recorded, and immobility time was manually quantified by a trained experimenter blind to the experimental conditions.

2.4 Protein extraction and Western Blot

Protein extraction and immunoblotting protocols were performed as we described before (Lax et al., 2018). Briefly, frozen samples were lysed using RIPA buffer, loaded into 7% SDS-PAGE gel, and transferred to a nitrocellulose membrane. Membranes were first blocked with 5% non-fat milk in TBST for 1 hour and then probed with antibodies against Parp1 and PAR. Beta-actin was used as a reference protein for normalization. On completing the FST, brains were immediately removed to measure Parp1 levels and Parp1 activation (i.e., PARylation) by western blot. Detailed descriptions of the primary and secondary antibodies are detailed in supplemental Table 1. Anti-Parp1, anti-PAR, and anti-beta-actin primary antibodies were diluted 1:1000. The secondary goat anti-mouse antibody was diluted 1: 5000. To detect protein levels, membranes were exposed to an ECL solution, and images were collected using a GelDoC device (Bio-Rad). Image quantification was done with ImageJ software.

2.5 RNA extraction, reverse transcription, and quantitative PCR (q-PCR)

We extracted RNA using an RNeasy mini kit (Qiagen) following the manufacturer's protocol. Next, we used GoScript Reverse Transcription Mix, Oligo (dT) kit (Promega, #A2791) to reverse transcribe the RNA into cDNA following the manufacturer protocol.

The qPCR procedure was performed using the Hy-syber power mix (hylabs, Israel) and included primers for the genes listed below in supplemental Table 2. RNA expression levels were normalized to beta-actin expression levels, and values were calculated using the $\Delta\Delta$CT method.

2.6 Statistics

Data are expressed as mean ±SEM unless otherwise stated. The data were analyzed by one-way ANOVA followed by Tukey's post-hoc tests. For ABT888 effect on immobility in the FST tests, two-way repeated measures ANOVA with time and treatment as the main factors followed by pair-wise Bonferroni-corrected t-tests. Comparisons between two groups in the qPCR experiments

were made with the ΔΔCT method and one-sample two-tailed t-tests. P-values smaller than 0.05 were considered statistically significant. Possible outliers were visible in the data; therefore, we applied the Iglewicz and Hoaglin's test for outliers under the strict criterion of z-score ≥|3.5|, which excluded no more than a single data point from each experimental group.

3. Results

3.1 FST triggers elevated cortical PARylation

Mice were exposed to the FST for 6 minutes and were sacrificed 15 min, 1 hour, or 24 hours later (Fig. 1A). PARylation was rapidly elevated 15 minutes after the FST and then returned to basal levels at 1-hour post-FST and remained at those level 24 hours later (Fig 1B; left). This elevated PARylation was followed by a slower increase in Parp1 levels, the primary enzyme that catalyzes PAR, 1-hour post-FST. Parp1 levels were reduced 24 hours later (Fig 1B; right). These findings suggest that FST triggers PARylation, in line with previous studies that showed neuronal- and neurotrophic- activity-induced PARylation (Azad et al., 2018; Homburg et al., 2000; Visochek et al., 2005).

3.2 FST induces upregulation of cortical immediate-early genes

Since FST causes extensive neuronal activation in the cortex (Connor, Kelly, & Leonard, 1997; Slattery, Neumann, & Cryan, 2011), we examined the mRNA expression levels of several immediate-early genes known to be transcribed upon neuronal activation. The mRNA expression levels were measured at the same time points as the PARylation levels (15 minutes, 1 hour, or 24 hours post-FST). We found a significant increase in Arc, c-Fos, Egr1, and Egr3 mRNA levels 15 minutes to 1 hour after the FST. The mRNA levels returned to baseline levels 24 hours later, and in the case of Egr3, mRNA levels returned to baseline levels already after 1 hour (Fig 2). Egr2, Bdnf, Nrn1, and Nr4a1 did not show a significant increase in mRNA levels, although, in all these genes, a trend toward increased mRNA expression was observed (Fig 2).

3.3 Parp inhibition impairs stress coping behavior in the FST

Next, we explored whether elevated PARylation induces the commonly seen shift in stress coping strategy from swimming as an active strategy into floating (i.e., immobility) as a passive strategy (Commons, Cholanians, Babb, & Ehlinger, 2017). For this aim, we injected the mice with the Parp inhibitor ABT888 (15 mg/kg IP, total volume of 0.1ml) or saline as a vehicle 50 minutes before the first FST. Twenty-four hours later, we re-exposed the mice to a second FST and recorded the immobility time during these two FST sessions (Fig 4A). We found that ABT888-injected mice showed no significant increase in immobility time, while control mice did show this shift in behavior (Fig 4B). This finding suggests that PARylation promotes the observed shift in stress coping strategy, likely through gene-expression regulation of immediate-early genes.

3.4 Parp inhibition reduces Arc expression after FST

Based on our findings, we speculated that the FST-induced increase in mRNA expression of at least some of the immediate-early genes is due to chromatin PARylation and, hence, relaxation that enables transcription. To identify which of the immediate-early genes were regulated through PARylation, we repeated the injection of ABT888 with the same settings, introduced the mice to the FST, and one hour later, we measured mRNA levels of the immediate-early genes that were significantly elevated in the previous experiment (i.e., Arc, C-Fos, Egr1; Fig 4A). We found that Arc (but not c-fos and Egr1) mRNA expression was significantly reduced following FST due to Parp inhibition (Fig 4B). This implies that Arc is a downstream target of FST-induced PARylation.

4. Discussion

Neuronal and behavioral responses to stressful events are required for proper adaptation to stress. However, the underlying molecular mechanisms that govern this response are yet to be fully explored. Here, we showed that PARylation, a necessity for long-term memory formation (Cohen-Armon et al., 2004), is induced shortly after exposure to FST and promotes adaptive stress coping.

PARylation happens shortly after neuronal activation or a neurotrophic stimulus (Homburg et al., 2000; Visochek et al., 2005; Wang et al., 2012). Parp1 catalyzes PAR chain formation on itself (i.e., auto-PARylation) and nearby histones, thus likely increasing chromatin accessibility (Azad et al., 2018; Kraus & Lis, 2003). This, in turn, enables elevated gene expression of immediate-early genes. An earlier *in-vitro* study found that high-frequency stimulation of cultured rat cerebral neurons induced PARylation-mediated expression of Arc, c-Fos, and Egr1 (Visochek et al., 2016). However, whether PARylation-mediated expression of these and other immediate-early genes occurs *in-vivo* in response to stressful stimuli was unknown. We hypothesized that (a) PARylation would be induced upon FST as part of the expected cortical neuronal response and stress adaptation; (b) that transcription of neuronal activity-dependent immediate-early genes that promotes neuronal plasticity and long-term memory formation are likely targets of this FST-induced PARylation (Cohen-Armon et al., 2004; Goldberg et al., 2009; Hernández et al., 2009; Visochek et al., 2016); and (c) that Parp inhibition will reduce the expression of these genes upon FST, resulting in reduced immobility in the second FST due to impaired adaptation response to the stress induced by the first FST.

As expected, we found that FST rapidly induced elevated cortical PARylation 15 minutes after FST, accompanied by a slower increase in Parp1, the primary enzyme that catalyzes PARylation. Further, systemic Parp1 inhibition diminished adaptive stress response in a second FST test. Gene-expression analysis of genes that were elevated after FST revealed that Parp1 inhibition reduced the expression of Arc, a known regulator of synaptic plasticity and protein-synthesis-dependent long-term potentiation (Korb & Finkbeiner, 2011; Messaoudi et al., 2007). These findings partly recapitulate previous *ex-vitro* work demonstrating a PARylation-dependent increase in Arc, C-Fos, and Egr1 expression following high-frequency electrical stimulation of rat cortical neurons (Visochek et al., 2016). We did not detect a PARylation-dependent increase in C-fos or Egr1; this partial discrepancy between the two studies may be due to the different experimental settings and species used (rats vs. mice).

Intra-cerebral and systemic PARP inhibitor applications were done before to manipulate reconsolidation and extinction of fear conditioning successfully (Elharrar, Dikshtein, Meninger-Mordechay, Lichtenstein, & Yadid, 2021; Goldberg et al., 2009; Inaba, Tsukagoshi, & Kida, 2015). However, the downstream effects of this inhibition were unknown. Our findings expand the role of PARylation in modulating behaviors to include also stress coping. Further, we filled the gap in knowledge between the behavioral effects of PARP inhibition and the observed PARylation-dependent long-term potentiation and increased expression of some immediate-early genes after *ex-vivo* stimulation.

Our experiments were focused on several potential target genes; further research is needed to explore the transcriptome-wide effects of FST-induced PARylation and PARP inhibition. In addition, we examined the impact of FST and PARP inhibition in the cerebral cortex to closely follow the tissue selection done in previous reports (Fontán-Lozano et al., 2010; Goldberg et al., 2009; Visochek et al., 2005). Future studies are required to decipher the effects of FST on PARylation and gene expression in different subregions of the cortex and other relevant brain regions, such as the hippocampus and the amygdala. We recognize these limitations and present our findings as a first step towards understanding the PARylation-dependent mechanisms of activity-dependent gene expression regulation required for proper stress-coping behavior.

Declaration of Competing Interests

We have no conflicts of interest to declare.

Acknowledgments

This work was funded by an internal grant from Ariel University. The Porsolt forced swim test icon by DBCLS https://togotv.dbcls.jp/en/pics.html is licensed under CC-BY 4.0 Unported https://creativecommons.org/licenses/by/4.0/.

References

Alberini, C. M. (2008). The role of protein synthesis during the labile phases of memory: Revisiting the skepticism. *Neurobiology of Learning and Memory*, *89*(3), 234–246. https://doi.org/10.1016/j.nlm.2007.08.007

Azad, G. K., Ito, K., Sailaja, B. S., Biran, A., Nissim-Rafinia, M., Yamada, Y., … Meshorer, E. (2018). PARP1-dependent eviction of the linker histone H1 mediates immediate early gene expression during neuronal activation. *Journal of Cell Biology*, *217*(2), 473–481. https://doi.org/10.1083/jcb.201703141

Bisaz, R., Travaglia, A., & Alberini, C. M. (2014). The neurobiological bases of memory formation: From physiological conditions to psychopathology. *Psychopathology*, *47*(6), 347–356. https://doi.org/10.1159/000363702

Cohen-Armon, M., Visochek, L., Katzoff, A., Levitan, D., Susswein, A. J., Klein, R., … Schwartz, J. H. (2004). Long-term memory requires polyADP-ribosylation. *Science*, *304*(5678), 1820–1822. https://doi.org/10.1126/science.1096775

Commons, K. G., Cholanians, A. B., Babb, J. A., & Ehlinger, D. G. (2017). The Rodent Forced Swim Test Measures Stress-Coping Strategy, Not Depression-like Behavior. *ACS Chemical Neuroscience*, *8*(5), 955–960. https://doi.org/10.1021/acschemneuro.7b00042

Connor, T. J., Kelly, J. P., & Leonard, B. E. (1997). Forced swim test-induced neurochemical, endocrine, and immune changes in the rat. *Pharmacology Biochemistry and Behavior*, *58*(4), 961–967. https://doi.org/10.1016/S0091-3057(97)00028-2

Donawho, C. K., Luo, Y., Luo, Y., Penning, T. D., Bauch, J. L., Bouska, J. J., … Frost, D. J. (2007). ABT-888, an orally active poly(ADP-ribose) polymerase inhibitor that potentiates DNA-damaging agents in preclinical tumor models. *Clinical Cancer Research*, *13*(9), 2728–2737. https://doi.org/10.1158/1078-0432.CCR-06-3039

Dudai, Y., & Eisenberg, M. (2004). Rites of passage of the engram: Reconsolidation and the lingering consolidation hypothesis. *Neuron*, *44*(1), 93–100. https://doi.org/10.1016/j.neuron.2004.09.003

Elharrar, E., Dikshtein, Y., Meninger-Mordechay, S., Lichtenstein, Y., & Yadid, G. (2021). Modulation of parp-1 activity in a broad time window attenuates memorizing fear. *International Journal of Molecular Sciences*, *22*(12). https://doi.org/10.3390/ijms22126170

Eric R. Kandel. (2001). The Molecular Biology of Memory Storage: A Dialogue Between Genes and Synapses. *Science*, *294*(5544), 1030–1038. https://doi.org/10.1126/science.1067020

Flavell, S. W., & Greenberg, M. E. (2008). Expression and Plasticity of the Nervous System. *Annu Rev Neurosci*, 563–590. https://doi.org/10.1146/annurev.neuro.31.060407.125631.Signaling

Fontán-Lozano, Á., Suárez-Pereira, I., Horrillo, A., Del-Pozo-Martín, Y., Hmadcha, A., & Carrión, Á. M. (2010). Histone H1 poly[ADP]-ribosylation regulates the chromatin alterations

required for learning consolidation. *Journal of Neuroscience*, *30*(40), 13305–13313. https://doi.org/10.1523/JNEUROSCI.3010-10.2010

Goldberg, S., Visochek, L., Giladi, E., Gozes, I., & Cohen-Armon, M. (2009). PolyADP-ribosylation is required for long-term memory formation in mammals. *Journal of Neurochemistry*, *111*(1), 72–79. https://doi.org/10.1111/j.1471-4159.2009.06296.x

Gutièrrez-Mecinas, M., Trollope, A. F., Collins, A., Morfett, H., Hesketh, S. A., Kersanté, F., & Reul, J. M. H. M. (2011). Long-lasting behavioral responses to stress involve a direct interaction of glucocorticoid receptors with ERK1/2-MSK1-Elk-1 signaling. *Proceedings of the National Academy of Sciences of the United States of America*, *108*(33), 13806–13811. https://doi.org/10.1073/pnas.1104383108

Hernández, A. I., Wolk, J., Hu, J. Y., Liu, J., Kurosu, T., Schwartz, J. H., & Schacher, S. (2009). Poly-(ADP-ribose) polymerase-1 is necessary for long-term facilitation in Aplysia. *Journal of Neuroscience*, *29*(30), 9553–9562. https://doi.org/10.1523/JNEUROSCI.1512-09.2009

Homburg, S., Visochek, L., Moran, N., Dantzer, F., Priel, E., Asculai, E., … Cohen-Armon, M. (2000). A fast signal-induced activation of poly(ADP-ribose) polymerase: A novel downstream target of phospholipase C. *Journal of Cell Biology*, *150*(2), 293–307. https://doi.org/10.1083/jcb.150.2.293

Inaba, H., Tsukagoshi, A., & Kida, S. (2015). PARP-1 activity is required for the reconsolidation and extinction of contextual fear memory. *Molecular Brain*, *8*(1), 21–25. https://doi.org/10.1186/s13041-015-0153-7

Korb, E., & Finkbeiner, S. (2011). Arc in synaptic plasticity: From gene to behavior. *Trends in Neurosciences*, *34*(11), 591–598. https://doi.org/10.1016/j.tins.2011.08.007

Kraus, W. L., & Lis, J. T. (2003). PARP goes transcription. *Cell*, *113*(6), 677–683. https://doi.org/10.1016/S0092-8674(03)00433-1

Lax, E., Friedman, A., Massart, R., Barnea, R., Abraham, L., Cheishvili, D., … Yadid, G. (2017). PARP-1 is required for retrieval of cocaine-associated memory by binding to the promoter of a novel gene encoding a putative transposase inhibitor. *Molecular Psychiatry*, *22*(4), 570–579. https://doi.org/10.1038/mp.2016.119

Lax, Elad, DoCarmo, S., Enuka, Y., Sapozhnikov, D., Welikovitch, L., Mahmood, N., … Szyf, M. (2018). Methyl-CpG binding domain 2 (Mbd2) is an Epigenetic Regulator of Autism-Risk Genes and Cognition. *Methyl-CpG Binding Domain 2 (Mbd2) Deficiency Causes Cognitive, Social and Emotional Deficits and down-Regulation of Ortholog Autism Risk Genes*, *2*(July), 247197. https://doi.org/10.1038/s41398-023-02561-9

McEwen, B. S., Bowles, N. P., Gray, J. D., Hill, M. N., Hunter, R. G., Karatsoreos, I. N., & Nasca, C. (2015). Mechanisms of stress in the brain. *Nature Neuroscience*, *18*(10), 1353–1363. https://doi.org/10.1038/nn.4086

Messaoudi, E., Kanhema, T., Soulé, J., Tiron, A., Dagyte, G., Da Silva, B., & Bramham, C. R. (2007). Sustained Arc/Arg3.1 synthesis controls long-term potentiation consolidation

through regulation of local actin polymerization in the dentate gyrus in vivo. *Journal of Neuroscience*, *27*(39), 10445–10455. https://doi.org/10.1523/JNEUROSCI.2883-07.2007

Ray Chaudhuri, A., & Nussenzweig, A. (2017). The multifaceted roles of PARP1 in DNA repair and chromatin remodelling. *Nature Reviews Molecular Cell Biology*, *18*(10), 610–621. https://doi.org/10.1038/nrm.2017.53

Ressler, K. J., Berretta, S., Bolshakov, V. Y., Rosso, I. M., Meloni, E. G., Rauch, S. L., & Carlezon, W. A. (2022). Post-traumatic stress disorder: clinical and translational neuroscience from cells to circuits. *Nature Reviews Neurology*, *18*(5), 273–288. https://doi.org/10.1038/s41582-022-00635-8

Reul, J. M. H. M. (2014). Making memories of stressful events: A journey along epigenetic, gene transcription, and signaling pathways. *Frontiers in Psychiatry*, *5*(JAN), 1–11. https://doi.org/10.3389/fpsyt.2014.00005

Saunderson, E. A., Spiers, H., Mifsud, K. R., Gutierrez-Mecinas, M., Trollope, A. F., Shaikh, A., … Reul, J. M. H. M. (2016). Stress-induced gene expression and behavior are controlled by DNA methylation and methyl donor availability in the dentate gyrus. *Proceedings of the National Academy of Sciences of the United States of America*, *113*(17), 4830–4835. https://doi.org/10.1073/pnas.1524857113

Scobie, K. N., Damez-Werno, D., Sun, H. S., Shao, N. Y., Gancarz, A., Panganiban, C. H., … Nestler, E. J. (2014). Essential role of poly(ADP-ribosyl)ation in cocaine action. *Proceedings of the National Academy of Sciences of the United States of America*, *111*(5), 2005–2010. https://doi.org/10.1073/pnas.1319703111

Slattery, D. A., Neumann, I. D., & Cryan, J. F. (2011). Transient inactivation of the infralimbic cortex induces antidepressant-like effects in the rat. *Journal of Psychopharmacology*, *25*(10), 1295–1303. https://doi.org/10.1177/0269881110368873

Thorsell, A. G., Ekblad, T., Karlberg, T., Löw, M., Pinto, A. F., Trésaugues, L., … Schüler, H. (2017). Structural Basis for Potency and Promiscuity in Poly(ADP-ribose) Polymerase (PARP) and Tankyrase Inhibitors. *Journal of Medicinal Chemistry*, *60*(4), 1262–1271. https://doi.org/10.1021/acs.jmedchem.6b00990

Visochek, L., Grigoryan, G., Kalal, A., Milshtein-Parush, H., Gazit, N., Slutsky, I., … Cohen-Armon, M. (2016). A PARP1-ERK2 synergism is required for the induction of LTP. *Scientific Reports*, *6*(April), 1–16. https://doi.org/10.1038/srep24950

Visochek, Leonid, Steingart, R. A., Vulih-Shultzman, I., Klein, R., Priel, E., Gozes, I., & Cohen-Armon, M. (2005). PolyADP-ribosylation is involved in neurotrophic activity. *Journal of Neuroscience*, *25*(32), 7420–7428. https://doi.org/10.1523/JNEUROSCI.0333-05.2005

Wang, S. H., Liao, X. M., Liu, D., Hu, J., Yin, Y. Y., Wang, J. Z., & Zhu, L. Q. (2012). NGF promotes long-term memory formation by activating poly(ADP-ribose) polymerase-1. *Neuropharmacology*, *63*(6), 1085–1092. https://doi.org/10.1016/j.neuropharm.2012.06.050

West, A. P. (1990). Neurobehavioral studies of forced swimming: The role of learning and memory in the forced swim test. *Progress in Neuropsychopharmacology and Biological Psychiatry*, *14*(6), 863–877. https://doi.org/10.1016/0278-5846(90)90073-P

Figure 1: (**A**) Experimental timeline: Force Swimming Test and euthanization of the experimental groups; (**B**) Expression levels of cortical PAR and PARP1 at the time points described in (A). One-way ANOVAs for PAR levels: $F_{(3,18)} = 5.243$, $p=0.0089$; one-way ANOVA for Parp1 levels: $F_{(3,20)} = 6.748$, $p=0.0025$, followed by Tukey's post-hoc test. n=5-6/group. *$p<0.05$, **$p<0.01$.

Figure 2: Expression levels of immediate-early and activity-dependent genes at several time points following FST. One-way ANOVAs for Arc: $F_{(3, 20)} = 5.991$, $p=0.0044$; Bdnf: $F_{(3, 20)} = 0.5675$, $p=0.6428$; Nr4a1: $F_{(3, 20)} = 0.7207$, $p=0.5513$; Nrn1: $F_{(3, 20)} = 1.741$, $p=0.1909$; C-Fos: $F_{(3, 20)} = 12.96$, $p<0.0001$; Egr1: $F_{(3, 20)} = 7.051$, $p=0.0020$; Egr2: $F_{(3, 20)} = 2.578$, $p=0.0823$; Egr3: $F_{(3, 20)} = 3.400$, $p=0.0378$. Followed by Tukey's post-hoc test. n=6/group. *$p<0.05$, **$p<0.01$, ***$p<0.001$.

Figure 3: (**A**) Experimental timeline: ABT-888 / saline injection, followed by two FSTs, 24h apart. (**B**) Immobility time (in seconds) for each of the FSTs. Two-way repeated-measures ANOVA with a main effect of time $F_{(1,17)}=21.09$, $p=0.0003$, no main effect of treatment $F_{(1,17)}=3.81$, $p=0.0675$ and no interaction $F_{(1,17)}=1.01$, $p=0.3288$, followed by pair-wise Bonferroni-corrected t-tests. n=10 for the saline group and n=9 for the ABT888 group. *$p<0.02$; n.s: no significant.

Figure 4: (**A**) Experimental timeline: ABT-888 / saline injection, followed by FST and mice euthanization 1-h after the end of the FST. (**B**) The relative expression level of the genes elevated 1 h after FST: Arc, C-fos, and Erg-1. T-test, **$p<0.01$, n=12 for the saline group and n=15 for the ABT888 group.

A

Experimental time-line

0 min.
ABT-888 / Saline injection

24 h.
2ⁿᵈ FST

50 min.
1ˢᵗ FST

Force Swim Test (FST)

B

A

Experimental timeline

0 min.

116 min.
(60 min.)

ABT-888 / Saline injection

Euthanizing mice

50 min.
Start FST

B