Deep learning models of RNA base-pairing structures make accurate zero-shot predictions of

base-base interactions of RNA complexes

Mei Lang[1], Thomas Litfin[2], Ke Chen[1], Jian Zhan[1,3 a], and Yaoqi Zhou[1,2a]

[1]Institute of Systems and Physical Biology, Shenzhen Bay Laboratory, Shenzhen, 518107, China

[2]Institute for Glycomics, Griffith University, Parklands Dr, Southport, QLD 4222, Australia

[3]Ribopeutic Inc, Guangzhou International Bio Island, Guangdong, 510320, China

[a]For correspondence, please email to Dr. Yaoqi Zhou (zhouyq@szbl.ac.cn) or Dr. Jian Zhan

(zhanjian@szbl.ac.cn).

ABSTRACT

The intricate network of RNA-RNA interactions, crucial for orchestrating essential cellular

processes like transcriptional and translational regulation, has been unveiling through high-

throughput techniques and computational predictions. With the emergence of deep learning

methodologies, the question arises: how do these cutting-edge techniques for base-pairing

prediction compare to traditional free-energy-based approaches, particularly when applied to

the challenging domain of interaction prediction via chain concatenation? In this study, we

employ base pairs derived from three-dimensional RNA complex structures as the gold standard

benchmark to assess the performance of 23 different methods, including recently developed

deep learning models. Our results demonstrate that the deep-learning-based methods, SPOT-

RNA can be generalized to make accurate zero-shot predictions of RNA-RNA interactions not only

between previously unseen RNA structures but also between RNAs without monomeric

structures. The finding underscores the potential of deep learning as a robust tool for advancing

our understanding of these complex molecular interactions.

Introduction

Recent advancements in high-throughput techniques have unveiled a complex network of RNA-

RNA interactions (RRIs) critical for governing transcriptional and translational processes. These

interactions are pivotal in the biogenesis of various RNA molecules, including mRNAs, rRNA, tRNA,

microRNAs, and circRNAs[1-3]. Large-scale detection of RRIs has been achieved through innovative

approaches that combine cross-linking techniques with high-throughput sequencing. Notably,

techniques such as PARIS[4], SPLASH[5], LIGR-seq[6], and COMRADES[7] have employed psoralen or its

derivatives, as well as formaldehyde in the case of RIC-Seq[8], for cross-linking. While these

methods hold promise, they are not without limitations stemming from probe biases and ligation

efficiencies[1-3]. For example, KARR-seq, PARIS, and RIC-seq can only achieve about 55% true

positive rate at 20% false positive rate for human 18S rRNA[5]. Furthermore, many of these high-

throughput techniques have yet to achieve the single nucleotide resolution.

Attaining the nucleotide-level resolution in RNA structures has historically relied on traditional

structure-determination methods such as X-ray crystallography, Nuclear Magnetic Resonance

(NMR), and Cryo-electron microscopy. Yet, compared to proteins, determining RNA structures

presents formidable challenges due to the unique physiochemical properties of nucleotides and

the inherent fragility of RNA structures[9]. This is reflected from the fact that only a meagre 3% of

structures in the Protein Data Bank contain RNAs, with even fewer dedicated to RNA-RNA

complexes (681 as of March 16, 2023, before redundancy removal)[10]. This stark contrast becomes

even more pronounced when considering the extensive collection of more than 31 million

noncoding RNA sequences catalogued in the RNAcentral database [11]. Given the cost and

challenges associated with experimental approaches, there is an imperative need for

development of complementary computational prediction techniques.

The existing methods for predicting RNA-RNA interactions (RRIs) can be broadly classified into

alignment-based, free-energy-based, and homology modeling approaches[12,13]. Alignment-based

techniques, such as GUUGle[14] and RIsearch[15], focus on inter-RNA base pairs while overlooking

potential intra-RNA interactions. Free-energy-based methods can be categorized into those

considering only intermolecular interactions for expediency (such as RNAhybrid[16], RNAduplex[17],

RNAplex-c[18], and DuplexFold[19]), those factoring in intramolecular interactions based on solvent

accessibility (such as RNAup[20], IntaRNA[21], RNAplex-a[22], and AccessFold[23]), and those

accommodating both intra- and inter-molecular base pairs through sequence concatenation

(such as PAIRFOLD[24], RNAcofold[25] and biFold[19]) or without restrictions (such as RactIP[26]).

Homology-based techniques, exemplified by TargetRNA2[27], CopraRNA[28], RNAaliduplex[17] and

PETcofold[29], utilize evolutionary information to infer binding.

Presently, 'de novo' RRI prediction methods predominantly rely on free-energy-based approaches,

limited by their approximate energy or scoring functions, akin to the challenges faced in RNA

secondary structure prediction[30]. Recent advancements have seen the emergence of deep

learning-based methods, starting with SPOT-RNA[31], which achieved the first end-to-end

prediction of intra-RNA base pairs. Subsequent developments include mxfold2[32], UFold[33], and

2dRNA[34]. To further enhance prediction accuracy, SPOT-RNA2[35] was developed to integrate

evolutionary profiles and mutational coupling data generated by RNAcmap[36].

In this study, we conducted a comprehensive benchmark of various methods for predicting inter-

RNA interactions. Our evaluation encompassed traditional energy-based techniques and newly

developed deep learning models based on simple sequence concatenation. To ensure a rigorous

assessment, we employed base pairs derived from experimentally-determined RNA-RNA complex

structures and eliminated monomer structures employed for training of SPOT-RNA and SPOT-

RNA2 through a strict structural similarity cutoff (TM-score[37] <0.3), in addition to sequence

similarity cut off by CD-HIT. This challenging set of the complex of unseen structures revealed

significant improvements of SPOT-RNA's performance over the other 22 methods evaluated,

underscoring the transferability of deep learning from intra-RNA to inter-RNA interaction

prediction. The performance by SPOT-RNA is robust against different types of RNA complexes.

Further tests on training-family-excluded test sets confirm the robustness of SPOT-RNA in

heterodimeric complexes of RNAs, in particular.

Results

Method Comparison on Inter-RNA Base Pair Prediction

In this study, we compared 23 different RRI predictors on a benchmark set of 64 RNA-RNA pairs

after excluding all monomer structures remotely similar to the RNA structures employed in the

training and validation sets for SPOT-RNA and SPOT-RNA2. We evaluated their performance in

untrained inter-RNA base pair prediction through precision/recall curves and F1 score

distributions, as shown in Figure 1. The performance metrics, including overall F1-score,

Mathews' correlation coefficient (MCC) values, and the median F1-score and standard deviation

of individual RNA pairs, are also summarized in Table 1. Predictors with probabilistic outputs are

represented by precision-recall (PR) curves, while others are represented as single points. For all

methods with chain concatenation for predicting RNA-RNA interactions, a low case "c" is

appended to the method name. They are RNAfoldc, UFoldc, MXfold2c, EternaFoldc, SPOT-RNAc,

and SPOT-RNA2c. No linker was employed because adding a 3-nucleotide link did not lead to

performance improvement (See Methods).

As shown in Figure 1A, SPOT-RNA2c is the most accurate predictor at low sensitivity (<0.2).

However, the overall PR curve given by SPOT-RNAc has the best performance. We can also

measure the performance by the overall F1-Score for all RNA pairs and the median F1-Score for

individual RNA pairs. The thresholds for determining F1-scores of SPOT-RNAc and SPOT-RNA2c in

the test set were set according to the thresholds for producing the highest F1-scores in the

validation dataset for SPOT-RNAc and SPOT-RNA2c, respectively. Table 1 confirmed the result

from the PR curve that SPOT-RNAc achieved the best overall performance with an overall F1-

Score of 0.583, outperforming SPOT-RNA2c (the overall F1-Score of 0.561) and EternaFold (F1-

Score of 0.512). SPOT-RNAc improves over SPOT-RNA2c by more than 4% and outperforms other

methods by over 14%, a pattern similarly observed in MCC values (Table 1).

Figure 1B presents the distribution of F1 scores for individual RNA pairs, including median, 25th,

and 75th percentiles. SPOT-RNAc continues to achieve the best performance with the highest

median F1 score of 0.582, outperforming the next best SPOT-RNA2c (median F1-score of 0.533)

with a 10% improvement. The improvement of SPOT-RNAc over all methods are statistically

significant except EternaFold (Table 1).

Figure 1: Performance comparison of 23 methods for inter-RNA base-pair prediction on the 64

complexes of RNA structures unseen by SPOT-RNA and SPOT-RNA2. All structures in the test set

have the structural similarity score TM-Score<0.3 compared to the monomeric structures used in

training and validating SPOT-RNA and SPOT-RNA2 methods. (A) Precision-recall curves (for those

methods with probabilistic outputs) or points given by 23 methods (B) Distribution of F1 scores

for inter-RNA base pair prediction for individual RNA pairs by the same 23 methods. Each boxplot

shows the median, 25th, and 75th percentiles, with outliers represented by "•". SPOT-RNAc

exhibits the best performance for both overall and individual measures of F1-scores.

Table 1: Performance Comparison of 23 Predictors of Inter-RNA Base Pairs on 64 Complexes of

RNA Structures Unseen by SPOT-RNA and SPOT-RNA2. All single-chain structures in the test set

have the structural similarity score TM-Score<0.3 compared to the monomeric structures used in

training and validating SPOT-RNA and SPOT-RNA2 methods.

Methods	PPV/ Precision	Sensitivity/Recall	Overall F1-Score	MCC	Individual F1-Score Median ± Std	P-value
GUUGle	0.157	0.105	0.126	0.124	0.000 ± 0.241	7.90e^{-15}
RIsearch	0.176	0.252	0.207	0.205	0.000 ± 0.319	2.03e^{-12}
RNAplex-cA*	0.463	0.148	0.224	0.259	0.000 ± 0.296	1.56e^{-11}
RNAaliduplex*	0.469	0.148	0.225	0.261	0.000 ± 0.296	1.56e^{-11}
DuplexFold	0.181	0.455	0.259	0.281	0.200 ± 0.344	6.12e^{-08}
RNAplex-c	0.189	0.442	0.265	0.283	0.216 ± 0.343	1.10e^{-08}
RNAduplex	0.194	0.491	0.278	0.303	0.229 ± 0.329	8.49e^{-09}
RNAplex-a	0.359	0.285	0.318	0.316	0.000 ± 0.331	1.43e^{-09}
RNAup	0.285	0.365	0.320	0.318	0.000 ± 0.342	4.13e^{-12}
AccessFold	0.317	0.507	0.390	0.397	0.349 ± 0.315	6.05e^{-06}
PETcoFold*	0.434	0.365	0.397	0.395	0.376 ± 0.352	9.24e^{-05}
RNAfoldc	0.384	0.582	0.463	0.469	0.446 ± 0.318	0.007
NUPACK	0.446	0.483	0.464	0.461	0.474 ± 0.355	0.001
MXfold2c*	0.391	0.603	0.474	0.482	0.448 ± 0.320	0.004
UFoldc	0.422	0.545	0.476	0.476	0.455 ± 0.302	0.002
bifold	0.412	0.566	0.477	0.480	0.454 ± 0.325	0.014
IntaRNA2.0	0.536	0.436	0.481	0.481	0.405 ± 0.377	9.75e^{-05}
PairFold	0.427	0.562	0.485	0.486	0.500 ± 0.339	0.007
RNAmultifold	0.423	0.582	0.490	0.493	0.494 ± 0.325	0.015
RNAcoFold	0.424	0.582	0.491	0.494	0.494 ± 0.325	0.015
EternaFoldc	0.473	**0.618**	0.512	0.519	0.498 ± 0.314	0.088
SPOT-RNA2c*	0.583	0.540	0.561	0.559	0.533 ± 0.331	0.012
SPOT-RNAc	**0.620**	0.551	**0.583**	**0.583**	**0.582** ± 0.326	

Note: The overall F1-score is harmonic mean of precision and recall for all RNA pairs. PPV denotes

positive predictive value. MCC denotes Matthews' correlation coefficient. The star * denotes the

use of evolution information. Methods with an ending of "c" indicate the use of chain

concatenation for RNA-RNA interaction prediction. Median F1 means the median F1 value of

single RNA. The P-value of a given method was computed by against the result from SPOT-RNAc.

Figure 2: Performance Comparison of Intra-RNA Base Pair Prediction by 12 methods (A) Precision-

recall curves and points illustrate the performance rankings. (B) A distribution of F1-scores for

intra-RNA base pair prediction. SPOT-RNA and SPOT-RNAc represent intramolecular base pair

prediction as single and concatenated chains, respectively. Evolution-based SPOT-RNA2c (or

SPOT-RNA2) outperforms SPOT-RNAc (or SPOT-RNA) for intra-RNA base pairs.

It's important to assess how these methods perform on intra-molecular interactions, although

not all RRI methods offer predictions for such interactions. We remove these RNAs without intra-

base-pairing structures. This leads to 77 RNA chains. Figure 2A compares the PR curves or PR

points given by 12 methods. Interestingly, PR curves indicate that SPOT-RNA2c now has the best

performance. The overall performance according to overall F1-scores given by SPOT-RNA2c (Table

2) is the highest, surpassing the next best methods (SPOT-RNA2 without chain concatenation) by

a 6% improvement in F1-score, and the third best (RNAmultifold or RNAcoFold) by 12%, with

SPOT-RNAc ranking as the fifth best. This trend is consistent with MCC values.

Table 2: Performance Comparison of 12 Predictors for Predicting Intra-RNA Base Pairs on 77

unseen RNA structures that have intra secondary structure in 64 RNA-RNA Complex Structures.

Methods	PPV/ Precision	Sensitivity /Recall	Overall F1-Score	MCC	Individual F1-Score Median ± Std	P-value
PETcoFold*	0.338	0.112	0.168	0.194	0.000 ± 0.217	$1.62e^{-15}$
UFoldc	0.471	0.574	0.517	0.519	0.452 ± 0.278	$5.33e^{-07}$
SPOT-RNA	0.584	0.486	0.531	0.531	0.476 ± 0.271	$1.51e^{-06}$
MXfold2c	0.538	0.536	0.537	0.536	0.444 ± 0.291	0.0001
RNAfoldc	0.528	0.578	0.552	0.551	0.494 ± 0.287	$1.06e^{-05}$
PairFold	0.581	0.525	0.552	0.551	0.483 ± 0.308	$8.15e^{-05}$
SPOT-RNAc	0.653	0.493	0.562	0.567	0.481 ± 0.291	0.0005
EternaFoldc	0.567	0.581	0.574	0.573	0.511 ± 0.294	0.001
RNAcoFold	0.587	0.566	0.576	0.576	0.467 ± 0.305	0.001
RNAmultifold	0.587	0.566	0.576	0.576	0.467 ± 0.305	0.001
SPOT-RNA2*	0.589	0.629	0.608	0.607	0.489 ± 0.310	0.007
SPOT-RNA2c*	**0.626**	**0.665**	**0.645**	**0.644**	**0.560 ± 0.294**	-

Note: The overall F1-score is harmonic mean of precision and recall for all RNA pairs. PPV denotes

positive predictive value. MCC denotes Matthew's correlation coefficient. The star * indicates the

use of evolution information. Std means standard deviation. Median F1 means the median F1

value of individual RNA chains. The P-value of a given method is calculated by against the result

of SPOT-RNA2c.

Figure 2B further examines the distribution of F1-scores for individual RNAs. SPOT-RNA2c

continues to lead with the highest median F1-score at 0.560, while EternaFold follows closely

with the second-best median F1-score of 0.511. The differences in F1-score distributions between

SPOT-RNA2c and other methods are all statistically significant, with a p-value of 0.001 when

comparing SPOT-RNA2c to the second best EternaFold (Table 2).

Intuitively, a better intra-RNA base-pairing prediction should lead to a better inter-RNA base-

pairing prediction. However, although SPOT-RNA2c has the best performance for intra-RNA

interaction prediction, it is SPOT-RNAc with the best performance for inter-RNA interaction

prediction. If we remove these RNA RRI pairs of which both chains do not have intra-base-pairing

structures, this leads to 53 RRI pairs. Figure 3A compares intermolecular F1-scores for individual

RNA pairs from SPOT-RNAc with the average intramolecular F1-scores. No correlation was found.

Similar uncorrelated intra- and inter-RNA F1-scores are observed for SPOT-RNA2c (Figure 3B).

This suggests that the evolution information contained in SPOT-RNA2c did contain the co-

evolution information for predicting intra-RNA, but not inter-RNA interactions.

Figure 3 No Correlation between Inter-RNA F1-scores and Intra-RNA F1-scores. (A) Inter-RNA F1-

scores versus the average Intra-RNA F1-scores of SPOT-RNAc for 53 RNA complex structures with

intra-RNA base pairs for both chains. (B) Inter-RNA F1-scores versus Intra-RNA F1-scores of SPOT-

RNA2c for 53 RNA complex structures.

We also assessed the impact of sequence concatenation on the intra-RNA base-pair prediction

(i.e., SPOT-RNA versus SPOT-RNAc, SPOT-RNA2 versus SPOT-RNA2c). Table 2 shows that SPOT-

RNAc/SPOT-RNA2c are better than SPOT-RNA/SPOT-RNA2 based on either overall F1-score or the

median of individual F1-scores. In both cases, the difference is statistically significant with a p-

value of 0.0007 between SPOT-RNA and SPOT-RNAc and a p-value of 0.007 between SPOT-RNA2

and SPOT-RNA2c. This indicates that knowing the binding partner improves the intra-RNA base

pair prediction.

We observed that some inter-RNA (and intra-RNA) interactions were predicted with F1 score of 0

as shown in Figures 3A and 3B. To understand the reason behind these poor predictions, we

examined F1-scores given by SPOT-RNAc as a function of sequence length (L1+L2) in Figure 4A.

No obvious correlation was found. However, when we plotted F1-scores against the number of

true inter-RNA base pairs divided by the square root of (L1*L2), a clear and strong correlation

emerged with a Pearson's correlation coefficient (PCC) of 0.464 (Figure 4B). Thus, poor

predictions, including those with F1-scores of 0, can be attributed to the scarcity of inter-RNA

contacts relative to the sequence lengths. This observation holds true for intra-RNA base pair

prediction as well: intra-RNA interactions with F1-scores of 0 also involve very few intra-RNA base

pairs (<5).

Figure 4: Relationship Between Inter-RNA Base Pairing Prediction and Sequence/Interaction

Characteristics. (A) The inter-RNA F1 scores for individual RNA-RNA complexes from SPOT-RNAc

plotted against the sum of sequence lengths (L1+L2). (B) The inter-RNA F1 scores for individual

RNA-RNA complexes from SPOT-RNAc plotted against the normalized number of inter-RNA base

pairs (the number of true inter-RNA base pairs divided by the $\sqrt{L1 \times L2}$). The performance does

not correlate with sequence length but is related to the normalized number of inter-RNA base

pairs.

203 One question is whether the method performance depends on RNA types. We classified 64

204 complexes into rRNA-containing complexes (only 8), snRNA/snoRNA/mRNA-containing complexes

205 (40), and RNA complexes lacking intra-RNA interactions (16). SPOT-RNAc is consistently the best

206 in each category as shown Supplementary Table S2 for the overall F1-score and MCC values. It is

207 of note that even for RNA complexes lacking intra-RNA interactions (i.e. these extreme cases of

208 unstructured RNAs are not employed for training in neither SPOT-RNA, nor SPOT-RNA2), SPOT-

209 RNAc continues to provide the best overall F1-score and MCC values, although its performance

210 for averaging over individual complexes is statistically similar to the those of some energy-based

211 techniques. The consistency of SPOT-RNAc performance among different RNA types

212 demonstrates that deep learning, even with a limited number of 3D structures for training, can

213 yield generalizable models for inter-RNA base pair prediction.

215 Despite the attempt to remove structural homologs from training by structural alignment, some

216 RNA families in training sets (rRNA and spliceosome complexes) were still included in the above

217 64-complex test sets because RNAs in the same family do not necessarily have an identical

218 structure. We further examined the effect of excluding-training families in intra-base pair

219 prediction by establishing new datasets. They were obtained by removing training/validation

220 families from 64 test complexes and locating additional new RNA complexes (up to 2024/02/28)

221 (See methods). These two datasets are mixed and separated into 15 hetero dimeric complexes

222 and 13 homo-dimeric complexes. For 15 hetero-dimer complexes, although the difference

223 between SPOT-RNAc and some methods are statistically insignificant, SPOT-RNAc remains the

224 best performance among all the methods compared for the overall performance (Table 3),

whereas SPOT-RNA2 remains the best for intra-RNA base-pair prediction (Supplementary Table

S3) with an overall F1-score of 0.589, followed by RNAmultifold (0.586) and RNAcoFold (0.586). If

ranked by median F1-score, SPOT-RNA2c remains the best for intra-RNA interaction predition.

Most did poorly with a median F1-score of 0 except SPOT-RNA2c, due to low number of intra-

RNA contacts for this set of the data. All also did very poorly with median F1-score of 0 for inter-

RNA base-pair prediction in the 13-homodimer set (Supplementary Table S4), whereas SPOT-

RNA2 is the best performance with an overall F1-score of 0.750, followed by SPOT-RNA (0.587)

and RNA-RNA2c (0.529) for intra-RNA base-pair prediction (Supplementary Table S5).

Table 3 Performance comparison of all predictors on 15 hetero-dimer complexes of inter-RNA

base pairs that are not in RNA families containing in TR0/VL0 (bpRNA) and TR1/VL1 for training

SPOT-RNA and SPOT-RNA2.

Methods	PPV/ Precision	Sensitivity/Recall	Overall F1-Score	MCC	Individual F1-Score Median ± Std	P-value
GUUGle	0.433	0.318	0.367	0.367	0.000 ± 0.355	$9.06e^{-05}$
RNAaliduplex*	0.647	0.331	0.438	0.460	0.381 ± 0.359	0.0005
RNAplex-cA*	0.647	0.331	0.438	0.460	0.381 ± 0.359	0.0005
RNAup	0.464	0.764	0.577	0.592	0.615 ± 0.227	$3.35e^{-06}$
RIsearch	0.472	0.744	0.578	0.589	0.781 ± 0.347	0.0399
RNAduplex	0.469	0.807	0.593	0.611	0.846 ± 0.301	0.1529
RNAplex-a	0.676	0.561	0.613	0.613	0.652 ± 0.271	0.0014
PETcoFold*	0.684	0.561	0.616	0.617	0.638 ± 0.306	0.0092
DuplexFold	0.559	0.780	0.651	0.657	0.846 ± 0.313	0.1867
RNAplex-c	0.572	0.780	0.660	0.665	0.846 ± 0.313	0.1867

PairFold	0.634	0.807	0.710	0.713	0.846 ± 0.283	0.2091
UFold	0.659	0.793	0.720	0.721	0.833 ± 0.274	0.1352
bifold	0.684	0.793	0.734	0.734	0.846 ± 0.278	0.2157
RNAfoldc	0.672	0.820	0.739	0.740	0.833 ± 0.273	0.1101
NUPACK	0.698	0.797	0.744	0.744	0.846 ± 0.271	0.3273
EternaFoldc	0.684	0.816	0.744	0.745	0.833 ± 0.269	0.0099
RNAmultifold	0.702	0.797	0.746	0.746	0.846 ± 0.276	0.1303
RNAcoFold	0.702	0.797	0.746	0.746	0.846 ± 0.276	0.1303
SPOT-RNA2c*	0.721	0.787	0.753	0.751	0.821 ± 0.269	0.0713
IntaRNA2.0	0.713	0.800	0.754	0.753	**0.846** ± 0.272	0.3950
MXfold2	0.684	**0.839**	0.754	0.756	0.833 ± 0.210	0.5833
AccessFold	0.696	0.826	0.755	0.756	0.833 ± 0.221	0.4844
SPOT-RNAc	**0.745**	0.823	**0.782**	**0.781**	0.833 ± 0.247	-

Note: The overall F1-score is harmonic mean of precision and recall for all RNA pairs. PPV denotes

positive predictive value. MCC denotes Matthews' correlation coefficient. The star * denotes the

use of evolution information. Methods with an ending of "c" indicate the use of chain

concatenation for RNA-RNA interaction prediction. Median F1 means the median F1 value of F1

values for single RNAs. The P-value of a given method was calculated by against the result from

SPOT-RNAc

We selected one example to illustrate SPOT-RNA's performance in RRI prediction. Figure 4A

displays predicted and actual base-pairing maps for the subunits L2a rRNA complexed with L3b

rRNA from Chlamydomonas reinhardtii mitoribosome (PDB ID 7PKT, chain ID 2 and chain ID 3).

Figure 5A displays the intra- and inter-base-pairing maps of these two RNAs with Figure 5B for

inter-base-pairing maps only. The F1 scores for intra-RNA base pairs are 0.32 for L2a rRNA and

0.30 for L3b rRNA, while the F1-score for inter-RNA base pairs is 0.571. Correctly predicted base

pairs are highlighted with red dots in Figures 5B and 5C, and their 3D locations were shown as

red-colored bases in Figure 5C. For this complex structure, precision for inter-RNA base pairs is

0.435, and sensitivity is relatively high (0.769).

Figure 5: Accurate Prediction of Key Inter-RNA Base-pairing Contacts in RNA Complexes by SPOT-

RNAc. Complex structure of large subunits of the Chlamydomonas reinhardtii mitoribosome

(L2aRNA and L3bRNA in PDB ID 7PKT) with true distance-contact map and predicted intra and

inter-RNA base pairs (A), inter-RNA base pairs only (B), and 3-D structure (C). Predicted intra-base

pairs and inter-base pairs are denoted by black and red dots, respectively, in the base-pairing

maps. In the 3-D structure (C), correctly predicted inter-RNA base pairs are highlighted in red.

Discussion

This study represents a comprehensive benchmark for assessing more than 20 methods in

predicting RNA-RNA interactions. Previous benchmarks were constrained to interactions involving

small RNAs without intra-RNA base pairs. For instance, Lai and Meyer [13] compared 14 RRI

methods based on experimentally confirmed interactions in fungal snoRNA-rRNA and bacterial

sRNA-mRNA pairs. Umu and Gardner[12] examined 15 RRI methods using a dataset focused on

short linear base-pair matching. Antonov et al.[38] compared 13 RRI methods on mammalian

lncRNAs with experimentally proven hybridizations. In contrast, our study presents the first

comprehensive benchmark of 23 RRI prediction methods using known RRI interactions derived

from 3D structures at the base pair level. Notably, most of these complexes (53/64) include RNAs

with 3D structures and intra-RNA base pairs. Additionally, this study marks the first inclusion of

deep-learning based methods for comparisons, utilizing chain concatenation.

In contrast to proteins, the available non-redundant data for RNA structures is limited. This raises

concerns about the generalizability of deep learning models trained on such limited data.

Previous studies by Szikszai et al [39] and Qiu [40] highlighted the challenges of deep-learning models

when applied to unseen families not present in the training and validation sets. SPOT-RNA and

SPOT-RNA2 were initially trained with a set of bpRNA dataset after removing all sequences in

bpRNA with known 3D structures by CD-HIT (80% sequence identity cut-off), followed by transfer

learning with 3D-structure-derived base pairs. To evaluate the adaptability of SPOT-RNA and

SPOT-RNA2 beyond their training and validation data, we placed them to the test of untrained

inter-RNA interactions. To avoid possible "over-trained" intra-RNA interactions affecting

prediction of inter-RNA interactions, we excluded all single-chain structures in the complex test

set with the structural similarity score (TM-score)≥0.3, compared to those in the training and

validation sets of SPOT-RNA. Interestingly, the performance of intra-RNA base-pair prediction

does not correlate with the performance of inter-RNA base-pair prediction (Figure 3A). Lacking

correlation suggests that the performance for predicting inter-RNA interactions is independent of

the performance for predicting intra-RNA interactions and the dataset for inter-RNA interactions

can be employed as an independent test set for the methods trained for predicting intra-RNA

interactions. Despite the above precaution, training/validation families were not entirely

excluded because RNAs in the same family may not have an identical structure. Thus, we further

established two family-excluded test sets (15 hetero-dimeric RNAs and 13 homo-dimeric RNAs).

SPOT-RNA and SPOT-RNA2 remain best performance for inter and intra-RNA base-pair prediction,

respectively, except for inter-base-pair prediction for homodimeric RNAs, for which no methods

did well. The generalization capability of SPOT-RNA is further illustrated with its robust

performance across different RNA types, even for RNAs with few intra-RNA structures

(Supplementary Table S2).

SPOT-RNA2, which incorporates evolution and co-evolution information, outperforms SPOT-RNA

for intra-RNA base pairs, aligning with previous findings (Table 2) [35]. However, SPOT-RNA2c

underperforms SPOT-RNAc for inter-RNA base pairs. Notably, SPOT-RNA2 exhibits a positive

correlation with the number of effective homologous sequences for intra-RNA base-pair

prediction (PCC=0.5, p-value=2×10^{-6}, Supplementary Figure 1C), but this correlation nearly

diminishes for SPOT-RNA2c for intra-RNA base pairs (PCC=0.27, p-value=0.05) and turned

negative for inter-RNA base pair prediction (PCC=-0.3, p-value=0.02). This suggests that using

linked sequences in SPOT-RNA2c for homology search may have provided harmful information

for inferring inter-RNA interactions. In the future, it may be necessary to utilize sequences from

the same species for homology searches, as co-evolution information can only be detected

through inter-species comparisons via Multiple Sequence Alignment (MSA) pairing as has been

done for proteins[41]. However, the failure of SPOT-RNA2 in predicting inter-RNA interactions

between homodimers indicates that the coevolution signals between RNAs may be too weak to

capture.

For predicting RNA-RNA interactions, we concatenated two chains (A and B) as a single chain. To

eliminate artificial sequence-order dependence, we predicted results for both AB and BA chains

and then calculated the average. Interestingly, we found that in some cases, one sequence order

(e.g., BA) outperformed the other (i.e., AB). Upon closer examination, we discovered for some

RNA pairs that the order with a shorter separation in sequence positions for contacting base pairs

tended to perform better. This observation is intuitive, as longer-range interactions are inherently

more challenging to predict. However, there were some outliers deserving further studies.

Methods

321   Benchmark datasets

We retrieved all RNA structures from the Protein Data Bank (PDB) in March 2023[10] and

specifically selected structures featuring two RNA chains with a minimum of 5 inter-RNA base

pairs. The identification of base pairs was carried out using DSSR[42]. For the base-pair prediction in

complex structures, interacting base pairs inside the complex are positive samples, and unpaired

bases between them are naturally classified as negative samples. To eliminate redundancy, we

applied CD-HIT-EST[43], removing binding pairs with over 80% sequence identity between either

chain. This initial step resulted in 155 unique RNA-RNA interaction (RRI) pairs.

To ensure stringency, we further filtered out RRI pairs that exhibited single-chain structural

similarities with any RNAs in the SPOT-RNA training set, defined by TM-score ≥0.3 using RNA-

align[44] with the length of the query sequence for normalization. This rigorous process yielded a

final benchmark set comprising 64 unique RRI pairs. The PDB IDs of the benchmark set can be

found in Supplementary Table S6. To further build a dataset excluding RNA families employed in

training/validating SPOT-RNA, we identified 14 complexes out of 64 test complexes whose RNAs

do not belong to any RFAM families with RNAs contained in training/validation TR0/VL0 (bpRNA)

and TR1/VL1 (RNAs with 3D structures) sets of SPOT-RNA.

To expand this dataset, we re-downloaded all RNA complexes on 2024/02/28 from the protein

data bank and processed by DSSR. Potential complexes were extracted from asymmetric units

and biological assemblies and then filtered such that only structures with 6 or more

intermolecular and 3 or more intramolecular base pairs (canonical and non-canonical) per

subunit were retained in the dataset. Any complex with a subunit of length >400 nucleotides

(almost exclusively rRNA and snRNA) or containing only subunits with length less than 25

nucleotides were removed from the dataset. Remaining RNA were filtered against the SPOT-RNA

training/validation sets (TR0/VL0/TR1/VL1) using blastn (-task blastn-short) with an e-value cutoff

of 0.001 and using RNA-align with a cutoff of 0.45. Identical duplicates were removed by

selecting structures with the largest number of base-pairing interactions. Two RNA (7r6q - rRNA

and 6ltp - crRNA) were also removed after manual inspection. Finally, we found that except for

three of the new complexes, all other complexes are homooligomers and among the original

complexes, only two of them are homooligomeric. Therefore, we integrated these two datasets,

resulting in a homo-dimer data set containing 13 complexes and a hetero-dimer data set

containing 15 complexes.

Performance evaluation

We evaluated performance using common metrics: Recall (sensitivity), Precision, and the F1

score. PPV/Precision is TP/(TP+FP), Sensitivity/Recall is TP/(TP+FN), and the F1 score is

2(Recall*Precision)/(Recall + Precision). Here, TP, FP, and FN are true positive, false positive and

false negative, respectively. We also calculated Matthew's correlation coefficient (MCC) to

provide a balanced measure as below:

360
$$MCC = \frac{TP \times TN - FP \times FN}{\sqrt{(TP + FP)(TP + FN)(TN + FP)(TN + FN)}},$$

where TN denotes true negatives. MCC considers true negatives (TN) and measures the

correlation between expected and observed classes. It ranges from 0 (no correlation) to 1

(highest correlation).

Methods for interaction prediction

We summarized the methods compared in Table 4, with detailed settings for each method

provided in the Supplementary method description in the Supplementary Material. As per Lai

and Meyer[13], we categorized the algorithms into four types: 1) 'Interaction only' methods predict

intermolecular hybridization, ignoring RNA secondary structures; 2) 'Accessibility' methods

consider RNA secondary structures using a partition function for unpaired probability; 3)

'Concatenation' algorithms treat two input sequences as a single chain and predict a joint

secondary structure; and 4) 'Complex joint' methods also predict joint secondary structures but

without concatenating the input sequences.

We used concatenation for comparing recent deep-learning methods with traditional free-

energy-based methods. When dealing with concatenated chains, we made predictions for both

sequence orders (AB and BA) and reported the average if probabilities were predicted. For those

methods providing a two-state prediction, we considered the union of base pairs predicted for

both sequence orders as a positive prediction (i.e. a positive prediction from either AB or BA

concatenation will be considered as positive).

We experimented with and without a three-nucleotide linker (AAA, UUU, CCC, or GGG) and

found that direct concatenation without any linker yielded slightly better results, although not

statistically significant compared to AAA/CCC linkers (Supplementary Table 1). Therefore, we

report results based on direct concatenation without any linkers.

Table 4 RRI interaction tools employed in this study for comparison, listed according to the year

of publication along with their categories, the use of evolution information (MSA), the algorithm,

and the capability of predicting intra-RNA base pairs.

Methods	Ref	Year	Category[a]	MSA[b]	Algorithms[c]	Intra-RNA base pairs[d]
RNAfoldc[e]	[45]	2004	concatenation	No	MFE	Yes
PairFold	[24]	2005	concatenation	No	MFE	Yes
RNAup	[20]	2006	accessibility	No	MFE	No

GUUGle	14	2006	interaction only	No	MFE	No
RNAcoFold	25	2006	concatenation	No	partition + MFE	Yes
NUPACK	46	2007	concatenation	No	MFE	No
RNAplex-c	18	2008	interaction only	No	MFE	No
RNAplex-a	18	2008	accessibility	No	MFE	No
RNAplex-cA	18	2008	interaction only	Yes	MFE	No
bifold	19	2010	complex joint	No	MFE	No
DuplexFold	47	2010	interaction only	No	MFE	No
PETcoFold	29	2011	complex joint	Yes	MFE	Yes
RNAduplex	17	2011	interaction only	No	MFE	No
RNAaliduplex	17	2011	interaction only	Yes	MFE	No
RNAmultifold	17	2011	concatenation	No	MFE	Yes
RIsearch	15	2012	interaction only	No	MFE	No
AccessFold	23	2016	concatenation	No	MFE	No
IntaRNA2.0	48	2017	accessibility	No	MFE + partition	No
SPOT-RNAc[e]	31	2019	concatenation	No	DL	Yes
SPOT-RNA2c[e]	35	2021	concatenation	Yes	DL	Yes
MXfold2c[e]	32	2021	concatenation	No	DL	Yes
UFoldc[e]	33	2022	concatenation	No	DL	Yes
EternaFoldc[e]	49	2022	concatenation	No	MFE	Yes

[a] Category: the broad category of the method (see text for more details).

[b]MSA—indicate whether it takes multiple sequence alignment as input.

[c]Algorithm: MFE: minimum free energy, Partition; partition function, DL: deep learning.

[d]Intra-RNA Base Pairs —whether the output of a software also contains base pairing information

for intramolecular interactions.

[e]c – indicates concatenation of two chains. For example, SPOT-RNAc and SPOT-RNA2c denotes

SPOT-RNA and SPOT-RNA2 with sequence concatenation, respectively, to distinguish from the

methods dedicated to individual chains (SPOT-RNA and SPOT-RNA2, respectively).

Homology search

For these methods that do not predicted RRI by employing a linked chain such as RNAplex-cA,

PETcoFold, RNAaliduplex and SPOT-RNA2, we searched the homologs using the single chain by

RNAcmap3[50] which is based on MARS database (a comprehensive database by including the

RNAcentral, MG-RAST, GWH, MGnify and NCBI's nucleotide databases). For SPOT-RNA2c using a

link chain, we searched the homologs with the linked chain by RNAcmap3. The median number

of homologous sequences found is 1467 for single chains and 940 for linked chains.

Data and code availability

All data and codes are available at https://github.com/meilanglang/RNA-RNA-Interaction

Acknowledgement

We express our gratitude to Dr. Jaswinder Singh for valuable discussions and insights. This

research received support from the National Natural Science Foundation of China (Grant #

22350710182), the Shenzhen Science and Technology Program (Grant No.

KQTD20170330155106581) and benefited from access to the supercomputing resources at

Shenzhen Bay Laboratory.

Conflict of Interest.

All authors declare no financial interest. Zhan and Zhou are the CEO and the chair of the scientific

advisor board for Ribopeutic, respectively.

Reference

1.   Dai, X., Zhang, S. & Zaleta-Rivera, K. RNA: interactions drive functionalities. *Mol Biol Rep* 47,

1413–1434 (2020).

2.   Singh, S., Shyamal, S. & Panda, A. C. Detecting RNA–RNA interactome. *WIREs RNA* 13, e1715

(2022).

3.   Gong, J., Ju, Y., Shao, D. & Zhang, Q. C. Advances and challenges towards the study of RNA-

RNA interactions in a transcriptome-wide scale. *Quant Biol* 6, 239–252 (2018).

4.   Lu, Z. *et al.* RNA Duplex Map in Living Cells Reveals Higher-Order Transcriptome Structure.

Cell 165, 1267–1279 (2016).

5.   Aw, J. G. A. *et al.* In Vivo Mapping of Eukaryotic RNA Interactomes Reveals Principles of

Higher-Order Organization and Regulation. *Mol Cell* 62, 603–617 (2016).

6.   Sharma, E., Sterne-Weiler, T., O'Hanlon, D. & Blencowe, B. J. Global Mapping of Human RNA-

RNA Interactions. *Mol Cell* 62, 618–626 (2016).

7.   Ziv, O. *et al.* COMRADES determines in vivo RNA structures and interactions. *Nat Methods* 15,

785–788 (2018).

8.   Cai, Z. *et al.* RIC-seq for global in situ profiling of RNA–RNA spatial interactions. *Nature* 582,

432–437 (2020).

9.   Xu, B. *et al.* Recent advances in RNA structurome. *Sci China Life Sci* 65, 1285–1324 (2022).

10.   Burley, S. K. *et al.* RCSB Protein Data Bank: powerful new tools for exploring 3D structures of

biological macromolecules for basic and applied research and education in fundamental biology,

biomedicine, biotechnology, bioengineering and energy sciences. *Nucleic Acids Res* 49, D437–

D451 (2021).

11.   RNAcentral Consortium. RNAcentral 2021: secondary structure integration, improved

sequence search and new member databases. *Nucleic Acids Res* 49, D212–D220 (2021).

12.   Umu, S. U. & Gardner, P. P. A comprehensive benchmark of RNA–RNA interaction prediction

tools for all domains of life. *Bioinformatics* 33, 988–996 (2017).

13.   Lai, D. & Meyer, I. M. A comprehensive comparison of general RNA-RNA interaction

prediction methods. *Nucleic Acids Res* 44, e61 (2016).

14.   Gerlach, W. & Giegerich, R. GUUGle: a utility for fast exact matching under RNA

complementary rules including G-U base pairing. *Bioinformatics* 22, 762–764 (2006).

15.   Wenzel, A., Akbasli, E. & Gorodkin, J. RIsearch: fast RNA-RNA interaction search using a

simplified nearest-neighbor energy model. *Bioinformatics* 28, 2738–2746 (2012).

16.   Krüger, J. & Rehmsmeier, M. RNAhybrid: microRNA target prediction easy, fast and flexible.

Nucleic Acids Res 34, W451-454 (2006).

17.   Lorenz, R. *et al.* ViennaRNA Package 2.0. *Algorithms Mol Biol* 6, 26 (2011).

18.   Tafer, H. & Hofacker, I. L. RNAplex: a fast tool for RNA-RNA interaction search. *Bioinformatics*

24, 2657–2663 (2008).

19.   Reuter, J. S. & Mathews, D. H. RNAstructure: software for RNA secondary structure

prediction and analysis. *BMC Bioinformatics* 11, 129 (2010).

20.   Mückstein, U. *et al.* Thermodynamics of RNA-RNA binding. *Bioinformatics* 22, 1177–1182

(2006).

21.   Busch, A., Richter, A. S. & Backofen, R. IntaRNA: efficient prediction of bacterial sRNA targets

incorporating target site accessibility and seed regions. *Bioinformatics* 24, 2849–2856 (2008).

22. Tafer, H., Amman, F., Eggenhofer, F., Stadler, P. F. & Hofacker, I. L. Fast accessibility-based

prediction of RNA-RNA interactions. *Bioinformatics* 27, 1934–1940 (2011).

23. DiChiacchio, L., Sloma, M. F. & Mathews, D. H. AccessFold: predicting RNA-RNA interactions

with consideration for competing self-structure. *Bioinformatics* 32, 1033–1039 (2016).

24. Andronescu, M., Zhang, Z. C. & Condon, A. Secondary structure prediction of interacting

RNA molecules. *J Mol Biol* 345, 987–1001 (2005).

25. Bernhart, S. H. *et al.* Partition function and base pairing probabilities of RNA heterodimers.

Algorithms Mol Biol 1, 3 (2006).

26. Kato, Y. *et al.* RactIP: fast and accurate prediction of RNA-RNA interaction using integer

programming. *Bioinformatics* 26, i460-466 (2010).

27. Kery, M. B., Feldman, M., Livny, J. & Tjaden, B. TargetRNA2: identifying targets of small

regulatory RNAs in bacteria. *Nucleic Acids Res* 42, W124-129 (2014).

28. Wright, P. R. *et al.* Comparative genomics boosts target prediction for bacterial small RNAs.

Proc Natl Acad Sci U S A 110, E3487-3496 (2013).

29. Seemann, S. E., Richter, A. S., Gesell, T., Backofen, R. & Gorodkin, J. PETcofold: predicting

conserved interactions and structures of two multiple alignments of RNA sequences.

Bioinformatics 27, 211–219 (2011).

30. Zhao, Y., Wang, J., Zeng, C. & Xiao, Y. Evaluation of RNA secondary structure prediction for

both base-pairing and topology. *Biophys Rep* 4, 123–132 (2018).

31. Singh, J., Hanson, J., Paliwal, K. & Zhou, Y. RNA secondary structure prediction using an

ensemble of two-dimensional deep neural networks and transfer learning. *Nat Commun* 10, 5407

(2019).

32. Sato, K., Akiyama, M. & Sakakibara, Y. RNA secondary structure prediction using deep

learning with thermodynamic integration. *Nat Commun* 12, 941 (2021).

33. Fu, L. *et al.* UFold: fast and accurate RNA secondary structure prediction with deep learning.

Nucleic Acids Res 50, e14 (2022).

34. Mao, K., Wang, J. & Xiao, Y. Length-Dependent Deep Learning Model for RNA Secondary

Structure Prediction. *Molecules* 27, 1030 (2022).

35. Singh, J. *et al.* Improved RNA Secondary Structure and Tertiary Base-pairing Prediction Using

Evolutionary Profile, Mutational Coupling and Two-dimensional Transfer Learning. *Bioinformatics*

btab165 (2021) doi:10.1093/bioinformatics/btab165.

36. Zhang, T. *et al.* RNAcmap: A Fully Automatic Pipeline for Predicting Contact Maps of RNAs by

Evolutionary Coupling Analysis. *Bioinformatics* btab391 (2021)

doi:10.1093/bioinformatics/btab391.

37. Gong, S., Zhang, C. & Zhang, Y. RNA-align: quick and accurate alignment of RNA 3D

structures based on size-independent TM-scoreRNA. *Bioinformatics* 35, 4459–4461 (2019).

38. Antonov, I. V., Mazurov, E., Borodovsky, M. & Medvedeva, Y. A. Prediction of lncRNAs and

their interactions with nucleic acids: benchmarking bioinformatics tools. *Briefings in*

Bioinformatics 20, 551–564 (2019).

39. Szikszai, M., Wise, M., Datta, A., Ward, M. & Mathews, D. H. Deep learning models for RNA

secondary structure prediction (probably) do not generalize across families. *Bioinformatics* 38,

3892–3899 (2022).

40. Qiu, X. Sequence similarity governs generalizability of de novo deep learning models for RNA

secondary structure prediction. *PLOS Computational Biology* 19, e1011047 (2023).

41. Improved prediction of protein-protein interactions using AlphaFold2 | Nature

Communications. https://www.nature.com/articles/s41467-022-28865-w.

42. Lu, X.-J., Bussemaker, H. J. & Olson, W. K. DSSR: an integrated software tool for dissecting the

spatial structure of RNA. *Nucleic Acids Res* 43, e142 (2015).

43. Li, W. & Godzik, A. Cd-hit: a fast program for clustering and comparing large sets of protein

or nucleotide sequences. *Bioinformatics* 22, 1658–1659 (2006).

44. Zheng, J., Xie, J., Hong, X. & Liu, S. RMalign: an RNA structural alignment tool based on a

novel scoring function RMscore. *BMC Genomics* 20, 276 (2019).

45. Mathews, D. H. *et al.* Incorporating chemical modification constraints into a dynamic

programming algorithm for prediction of RNA secondary structure. *Proc Natl Acad Sci U S A* 101,

7287–7292 (2004).

46. Dirks, R. M., Bois, J. S., Schaeffer, J. M., Winfree, E. & Pierce, N. A. Thermodynamic Analysis

of Interacting Nucleic Acid Strands. *SIAM Rev.* 49, 65–88 (2007).

47. Reuter, J. S. & Mathews, D. H. RNAstructure: software for RNA secondary structure

prediction and analysis. *BMC Bioinformatics* 11, 129 (2010).

48. Mann, M., Wright, P. R. & Backofen, R. IntaRNA 2.0: enhanced and customizable prediction

of RNA–RNA interactions. *Nucleic Acids Research* 45, W435–W439 (2017).

49. Wayment-Steele, H. K. *et al.* RNA secondary structure packages evaluated and improved by

high-throughput experiments. *Nat Methods* 19, 1234–1242 (2022).

50. Chen, K., Litfin, T., Singh, J., Zhan, J. & Zhou, Y. MARS and RNAcmap3: The Master Database

of All Possible RNA Sequences and Its Integration with RNAcmap for RNA Homology Search.

Genomics, Proteomics, & Bioinformatics, in press (2024).

525