Reduced discrimination between signals of danger and safety but not overgeneralization is linked to exposure to childhood adversity in healthy adults

Maren Klingelhöfer-Jens[1], Katharina Hutterer[2], Miriam A. Schiele[3], Elisabeth J. Leehr[4], Dirk Schümann[1], Karoline Rosenkranz[1], Joscha Böhnlein[4], Jonathan Repple[4,5], Jürgen Deckert[2], Katharina Domschke[3], Udo Dannlowski[4], Ulrike Lueken[6,7], Andreas Reif[5,8], Marcel Romanos[9], Peter Zwanzger[10,11], Paul Pauli[12], Matthias Gamer[12], & Tina B. Lonsdorf[13]

[1] Institute for Systems Neuroscience, University Medical Center Hamburg-Eppendorf, Hamburg, Germany

[2] Department of Psychiatry, Psychosomatics and Psychotherapy, Center of Mental Health, University Hospital of Würzburg, University of Würzburg, Würzburg, Germany

[3] Department of Psychiatry and Psychotherapy, Medical Center - University of Freiburg, Faculty of Medicine, University of Freiburg, Freiburg, Germany

[4] Institute for Translational Psychiatry, University of Münster, Münster, Germany

[5] Department of Psychiatry, Psychosomatic Medicine and Psychotherapy, University Hospital Frankfurt – Goethe University, Frankfurt am Main, Germany

[6] Department of Psychology, Humboldt-Universität zu Berlin, Berlin, Germany

[7] German Center of Mental Health (DZPG), partner site Berlin-Potsdam, Berlin, Germany

[8] Fraunhofer Institute for Translational Medicine and Pharmacology ITMP, Frankfurt am Main, Germany

[9] Department of Child and Adolescent Psychiatry, Psychosomatics and Psychotherapy, Center of Mental Health, University Hospital of Würzburg, University of Würzburg, Würzburg, Germany

[10] Kbo Inn Salzach Hospital Clinical Center for Psychiatry, Wasserburg am Inn, Germany

[11] Department of Psychiatry, Ludwig-Maximilian-University Munich, Munich, Germany

[12] Department of Psychology and Center of Mental Health, Julius Maximilians University of Würzburg, Würzburg, Germany

[13] Department of Psychology, Biological Psychology and Cognitive Neuroscience, University of Bielefeld, Bielefeld, Germany

The authors made the following contributions. Maren Klingelhöfer-Jens: Conceptualization, Software, Validation, Formal analysis, Writing - Original Draft, Writing - Review & Editing, Visualization; Katharina Hutterer: Investigation, Ressources, Data Curation, Writing - Review & Editing; Miriam A. Schiele: Investigation, Data Curation, Writing - Review & Editing, Supervision, Project Administration; Elisabeth J. Leehr: Investigation, Writing - Review & Editing, Project Administration; Dirk Schümann: Investigation, Data Curation; Karoline Rosenkranz: Investigation, Data Curation; Joscha Böhnlein: Investigation, Writing - Review & Editing, Project Administration; Jonathan Repple: Investigation, Writing - Review & Editing, Project Administration; Jürgen Deckert: Ressources, Writing - Review & Editing, Supervision, Project Administration, Funding Acquisition; Katharina Domschke: Ressources, Writing - Review & Editing, Supervision, Project Administration, Funding Acquisition; Udo Dannlowski: Ressources, Writing - Review & Editing, Supervision, Project Administration, Funding Acquisition; Ulrike Lueken: Ressources, Writing - Review & Editing, Supervision, Project Administration, Funding Acquisition; Andreas Reif: Ressources, Writing - Review & Editing, Supervision, Project Administration, Funding Acquisition; Marcel Romanos: Ressources, Writing - Review & Editing, Supervision, Project Administration, Funding Acquisition; Peter Zwanzger: Ressources, Writing - Review & Editing, Supervision, Project Administration, Funding Acquisition; Paul Pauli: Ressources, Writing - Review & Editing, Supervision, Project Administration, Funding Acquisition; Matthias Gamer: Ressources, Writing - Review & Editing, Supervision, Project Administration, Funding Acquisition; Tina B. Lonsdorf: Conceptualization, Methodology, Ressources, Writing - Original Draft, Writing - Review & Editing, Supervision, Project Administration, Funding Acquisition.

Correspondence concerning this article should be addressed to Maren Klingelhöfer-Jens, University Medical Center Hamburg-Eppendorf, Martinistrasse 52, Bldg. W34, 20246 Hamburg, Germany. E-mail: m.klingelhoefer-jens@uke.de

Abstract

Childhood adversity is a strong predictor of developing psychopathological conditions. Multiple theories on the mechanisms underlying this association have been suggested which, however, differ in the operationalization of 'exposure'. Altered (threat) learning mechanisms represent central mechanisms by which environmental inputs shape emotional and cognitive processes and ultimately behavior. 1402 healthy participants underwent a fear conditioning paradigm (acquisition training, generalization), while acquiring skin conductance responses (SCRs) and ratings (arousal, valence, and contingency). Childhood adversity was operationalized as (1) dichotomization, and following (2) the specificity model, (3) the cumulative risk model, and (4) the dimensional model. Individuals exposed to childhood adversity showed blunted physiological reactivity in SCRs, but not ratings, and reduced CS+/CS- discrimination during both phases, mainly driven by attenuated CS + responding. The latter was evident across different operationalizations of 'exposure' following the different theories. None of the theories tested showed clear explanatory superiority. Notably, a remarkably different pattern of increased responding to the CS- is reported in the literature for anxiety patients, suggesting that individuals exposed to childhood adversity may represent a specific sub-sample. We highlight that theories linking childhood adversity to (vulnerability to) psychopathology need refinement.

Keywords: Childhood Adversity, Childhood Maltreatment, Early Adversity, Adverse Childhood Experiences, Fear Conditioning, Generalization, Skin Conductance, SCR, Fear Ratings

Reduced discrimination between signals of danger and safety but not overgeneralization is linked to exposure to childhood adversity in healthy adults

Introduction

Exposure to childhood adversity - particularly in early life - is a strong predictor for developing somatic, psychological, and psychopathological conditions (Anda et al., 2006; Baldwin, Coleman, Francis, & Danese, 2024; Danese, Pariante, Caspi, Taylor, & Poulton, 2007; Danese & Widom, 2023; Felitti, 2002; Gilbert et al., 2009; Green et al., 2010; Heim & Nemeroff, 2002; Hosseini-Kamkar et al., 2023; Klauke, Deckert, Reif, Pauli, & Domschke, 2010; McLaughlin et al., 2012; Moffitt et al., 2007; Samaey et al., 2024; Teicher, Gordon, & Nemeroff, 2022) and is hence associated with substantial individual suffering as well as societal costs (Hughes et al., 2021). Exposure to childhood adversity is rather common, with nearly two-thirds of individuals experiencing one or more traumatic events prior to their 18th birthday (McLaughlin et al., 2013). While not all trauma-exposed individuals develop psychopathological conditions, there is some evidence of a dose-response relationship (Danese et al., 2009; Smith & Pollak, 2021; Young et al., 2019). As this potential relationship is not yet fully clear, understanding the mechanisms by which childhood adversity becomes biologically embedded and contributes to the pathogenesis of stress-related somatic and mental disorders is central to the development of targeted intervention and prevention programmes. Learning is a core mechanism through which environmental inputs shape emotional and cognitive processes and ultimately behavior. Thus, learning mechanisms are key candidates potentially underlying the biological embedding of exposure to childhood adversity and their impact on development and risk for psychopathology (McLaughlin & Sheridan, 2016).

Fear conditioning is a prime translational paradigm for testing potentially altered (threat) learning mechanisms following exposure to childhood adversity under laboratory conditions. The fear conditioning paradigm typically consists of different experimental phases (Lonsdorf et al., 2017). During fear acquisition training, a neutral cue is paired with an aversive event such as an electrotactile stimulation or a loud aversive human scream (unconditioned stimulus, US). Through these pairings, an association between both stimuli is formed and the previously neutral cue becomes a conditioned stimulus (CS+) that elicits conditioned responses. In human differential conditioning experiments, typically a second neutral cue is never paired with the US and serves as a control or safety stimulus (i.e., CS-). During a subsequent fear extinction training phase, both the CS+ and the CS- are presented without the US which leads to a gradual waning of conditioned responding. A fear generalization phase includes additional stimuli (i.e., generalization stimuli; GSs) that are perceptually or conceptually similar to the CS+ and CS- (e.g., generated through merging perceptual properties of the CS+ and CS-) which allows for the investigation of what degree to which conditioned responding generalizes to similar cues.

Fear acquisition as well as extinction are considered as experimental models of the development and exposure-based treatment of anxiety- and stress-related disorders. Fear generalization is in principle adaptive in ensuring survival ('better safe than sorry'), but broad overgeneralization can become burdensome for patients. Accordingly, maintaining the ability to distinguish between signals of danger (i.e., CS+) and safety (i.e., CS-) under aversive circumstances is crucial, as it is assumed to be beneficial for healthy functioning (Hölzel et al., 2016) and predicts resilience to life stress (Craske et al., 2012), while reduced discrimination between the CS+ and the CS- has been linked to pathological anxiety (Duits et al., 2015; Lissek et al., 2005): Meta-analyses suggest that patients suffering from anxiety- and stress-related disorders show enhanced responding to the safe CS- during fear acquisition (Duits et al., 2015).

During extinction, patients exhibit stronger defensive responses to the CS+ and a trend toward increased discrimination between the CS+ and CS- compared to controls, which may indicate delayed and/or reduced extinction (Duits et al., 2015). Furthermore, meta-analytic evidence also suggests stronger generalization to cues similar to the CS+ in patients and more linear generalization gradients (Cooper, van Dis, et al., 2022; Dymond, Dunsmoor, Vervliet, Roche, & Hermans, 2015; Fraunfelter, Gerdes, & Alpers, 2022). Hence, aberrant fear acquisition, extinction, and generalization processes may provide clear and potentially modifiable targets for intervention and prevention programs for stress-related psychopathology (McLaughlin & Sheridan, 2016).

In sharp contrast to these threat learning patterns observed in patient samples, a recent review provided converging evidence that exposure to childhood adversity is linked to reduced CS discrimination, driven by blunted responding to the CS+ during experimental phases characterized through the presence of threat (i.e., acquisition training and generalization, Ruge et al., 2024). Of note, this pattern was observed in mixed samples (healthy, at risk, patients), in pediatric samples, and in adults exposed to childhood adversity as children. The latter suggests that recency of exposure or developmental timing may not play a major role, even though there is some evidence pointing towards accelerated pubertal and neural (connectivity) development in exposed children (Machlin, Miller, Snyder, McLaughlin, & Sheridan, 2019; Silvers et al., 2016). There is, however, no evidence pointing towards differences in extinction learning or generalization gradients between individuals exposed and unexposed to childhood adversity (for a review, see Ruge et al., 2024).

Ruge et al. (2024) also highlighted operationalization as a key challenge in the field hampering the interpretation of findings across studies and consequently cumulative knowledge

generation. Operationalization of exposure to childhood adversity, and hence the translation of theoretical accounts of the role of childhood adversity into statistical tests, is an ongoing discussion in the field (McLaughlin, Sheridan, Humphreys, Belsky, & Ellis, 2021; Pollak & Smith, 2021; Smith & Pollak, 2021). Historically, childhood adversity has been conceptualized rather broadly considering different adversity types lumped into a single category. This follows from the (implicit) assumption that any exposure to an adverse event will have similar and additive effects on the individual and its (neuro-biological) development (Smith & Pollak, 2021). Accordingly, childhood adversity has often been considered as a cumulative measure ('cumulative risk approach'; McLaughlin et al., 2021; Smith & Pollak, 2021). An alternative approach Sheridan & McLaughlin (2014) posits that different types of adverse events have a distinct impact on individuals and their (neuro-biological) development through distinct mechanisms ('specificity approach'; McLaughlin et al., 2021; Smith & Pollak, 2021). Currently, distinguishing between threat and deprivation exposure represents the prevailing approach (McLaughlin, DeCross, Jovanovic, & Tottenham, 2019), which has been formalized in the (two-)dimensional model of adversity and psychopathology (DMAP; Machlin et al., 2019; McLaughlin & Sheridan, 2016; McLaughlin et al., 2021; McLaughlin, Sheridan, & Lambert, 2014; Sheridan & McLaughlin, 2014, 2016). To this end, exposure to threat-related childhood adversity has been suggested to be specifically linked to altered emotional and fear learning (Sheridan & McLaughlin, 2014).

Yet, there is converging evidence from different fields of research suggesting that the effects of exposure to childhood adversity are cumulative, non-specific and rather unlikely to be tied to specific types of adverse events (Danese et al., 2009; Smith & Pollak, 2021; Young et al., 2019) - with few exceptions (Colich, Rosen, Williams, & McLaughlin, 2020; McLaughlin, Weissman, & Bitrán, 2019). This is also supported by the recent review of Ruge et al. (2024).

However, the different theoretical accounts have not yet been directly compared in a single fear conditioning study.

Here, we aim to fill this gap in an extraordinarily large sample of healthy adults (N = 1402) recruited in the context of a multi-centric study conducted at the Universities of Münster, Würzburg, and Hamburg, Germany. Participants underwent a differential fear conditioning paradigm including a fear acquisition and generalization phase using female faces as CSs and GSs and a female scream as US, while we measured skin conductance responses (SCRs) and different ratings types (i.e., arousal, valence, and US contingency). For SCRs and fear ratings, we calculated three different outcomes: CS discrimination (i.e., the difference between CS+ and CS-responses), the linear deviation score (LDS) as an index of the linearity of the generalization gradient (Kaczkurkin et al., 2017), and the general reactivity which was defined as the reactivity across all phases (for more details, see Materials and methods section). We also performed manipulation checks to verify whether the experimental manipulations had the intended effect.

We operationalize childhood adversity exposure through different approaches: Our main analyses employ the approach adopted by most publications in the field (see Ruge et al., 2024 for a review) - dichotomization of the sample into exposed vs. unexposed individuals based on published cut-offs for the Childhood Trauma Questionnaire [CTQ; Bernstein et al. (2003); Wingenfeld et al. (2010)]. Individuals were classified as exposed to childhood adversity if at least one CTQ subscale met the published cut-off (Bernstein & Fink, 1998; Häuser, Schmutzer, & Glaesmer, 2011) for at least moderate exposure (i.e., emotional abuse $\geq$ 13, physical abuse $\geq$ 10, sexual abuse $\geq$ 8, emotional neglect $\geq$ 15, physical neglect $\geq$ 10).

In addition, we provide exploratory analyses that attempt to translate dominant (verbal) theoretical accounts (McLaughlin et al., 2021; Pollak & Smith, 2021) on the impact of exposure to childhood adversity into statistical tests. At the same time, we acknowledge that such a translation is not unambiguous and these exploratory analyses should be considered as showcasing a set of plausible solutions. With this, we aim to facilitate comparability, replicability, and cumulative knowledge generation in the field as well as providing a solid base for hypothesis generation (Ruge et al., 2024) and refinement of theoretical accounts. More precisely, we attempted to exploratively translate a) the cumulative risk approach, which is based on the assumed key role of cumulative childhood adversity exposure, b) the specificity model, which considers specific types of exposure (in the present study: abuse and neglect), and c) the dimensional model, which also considers specific exposure types but controls for the effects of one another, into statistical tests applied to our dataset. Furthermore, we compiled challenges that arise in the practical implementation of these verbal theories into statistical models (for more details, see Materials and methods section, Table 2).

Based on the recently reviewed literature (Ruge et al., 2024), we expected less discrimination between signals of danger (CS+) and safety (CS-) in exposed individuals as compared to those unexposed to childhood adversity - primarily due to reduced CS+ responses - during both the fear acquisition and the generalization phase. Based on the literature (Ruge et al., 2024), we did not expect group differences in generalization gradients.

Methods and materials

Participants

In total, 1678 healthy participants (age_M = 25.26 years, age_{SD} = 5.58 years, female = 60.10%, male = 39.30%) were recruited in a multi-centric study at the Universities of Münster, Würzburg, and Hamburg, Germany (SFB TRR58). Data from parts of the Würzburg sample have been reported previously (Herzog et al., 2021; Imholze et al., 2023; Schiele, Reinhard, et al., 2016; Schiele, Ziegler, et al., 2016; Stegmann et al., 2019). These previous reports, also those focusing on experimental fear conditioning (Schiele, Reinhard, et al., 2016; Stegmann et al., 2019), addressed, however, research questions different from the ones investigated here (see also Supplementary Material for details). The study was approved by the local ethics committees of the three Universities (Münster: 2016-131-b-S, Ethics Committee Westfalen-Lippe; Würzburg: Votum 07/08, Ethics Committee of the Medical Faculty of the University of Würzburg; Hamburg: PV2755, Ethics Committee of the General Medical Council Hamburg) and was conducted in agreement with the Declaration of Helsinki Current and/or lifetime diagnosis of DSM-IV mental Axis-I disorders, as assessed by the German version of the Mini International Psychiatric Interview (Sheehan et al., 1998), led to exclusion from the study (see Supplementary Material for additional exclusion criteria). All participants provided written informed consent and received 50 € as compensation.

A reduced number of 1402 participants (age_M = 25.38 years, age_{SD} = 5.76 years, female = 60.30%, male = 39.70%) were included in the statistical analyses because 276 participants were excluded due to missing data (CTQ: n = 21, ratings: n = 78, SCRs: n = 182), for technical reasons, and due to deviating from the study protocol. Five participants had missing CTQ and missing SCR data. Thus, the sum of exclusions in specific outcome measures does not add up to

the total number of exclusions. We did not exclude physiological SCR non-responders or non-learners, as this procedure has been shown to induce bias through predominantly excluding specific subpopulations (e.g., high trait anxiety), which may be particularly prevalent in individuals exposed to childhood adversity (Lonsdorf et al., 2019). See Table 1 and Supplementary Material for additional sample information including trait anxiety and depression scores (see Supplementary Figure 6 and 7), zero-order correlations (Pearson's correlation coefficient) between trait anxiety, depression scores, and childhood adversity (see Supplementary Figure 1) as well as information on socioeconomic status (see Supplementary Figure 2).

Table 1: Descriptive information on the subsamples being exposed or unexposed to childhood adversity

Variable	Exposed	Unexposed	Statistics
N	203 (14%)	1199 (86%)	$X^2(1) = 707.57, p < .001$
Female/Male	124 (61%) / 79 (39%)	721 (61%) / 478 (39%)	$X^2(1) = 0.03, p = .858$
Age (M/SD)	26.80 (6.99)	25.14 (5.50)	$t(246.1) = -3.21, p < .001, d = 0.29$
STAI-T sum (M/SD)	38.73 (9.52)	34.04 (7.83)	$t(250.4) = -6.65, p < .001, d = 0.58$
ADS-K sum (M/SD)	8.71 (6.31)	6.69 (5.70)	$t(261) = -4.28, p < .001, d = 0.35$

Note. STAI-T = State-Trait Anxiety Inventory, Trait scale (Spielberger, 1983), ADS-K = Allgemeine Depressionsskala - Kurzform (short version of the Center for Epidemiological Studies-Depression Scale, CES-D; Hautzinger and Bailer, 1993). Individuals were classified as exposed to childhood adversity if at least one subscale met the published cut-off (Bernstein & Fink, 1998; Häuser et al., 2011) for at least moderate exposure (i.e., emotional abuse ≥ 13, physical abuse ≥ 10, sexual abuse ≥ 8, emotional neglect ≥ 15, physical neglect ≥ 10).

Procedure

Fear conditioning and generalization paradigm. Participants underwent a fear conditioning and generalization paradigm which was adapted from Lau et al. (2008) and described previously in detail (Herzog et al., 2021; Schiele, Reinhard, et al., 2016; Stegmann et al., 2019). Details are also provided in brief in the Supplementary Material (see also Supplementary Figure 3).

Ratings

At the end of each experimental phase (habituation, acquisition training, and generalization) as well as after half of the total acquisition and generalization trials, participants provided ratings of the faces with regards to valence, arousal (9-point Likert-scales; from 1 = very unpleasant/very calm to 9 = very pleasant/very arousing) and US contingencies (11-point Likert-scale; from 0 to 100% in 10% increments). As the US did not occur during the habituation phase, contingency ratings were not provided after this phase. For reasons of comparability, valence ratings were inverted.

Physiological data recordings and processing

Skin conductance was recorded continuously using Brainproducts V-Amp-16 and Vision Recorder software (Brainproducts, Gilching, Germany) at a sampling rate of 1000 Hz from the non-dominant hand (thenar and hypothenar eminences) using two Ag/AgCl electrodes. Data were analyzed offline using BrainVision Analyzer 2 software (Brainproducts, Gilching, Germany). The signal was filtered offline with a high cut-off filter of 1 Hz and a notch filter of 50 Hz. Amplitudes of SCRs were quantified by using the Trough-to-peak (TTP) approach. According to published guidelines (Boucsein et al., 2012), the response onset was defined as between 900–

4000 ms after stimulus onset and the peak between 2000–6000 ms after stimulus onset. A minimum response criterion of 0.02 μS was applied, with lower índividual responses scored as zero (i.e., non-responses). Note that previous work using this sample (Schiele, Reinhard, et al., 2016; Stegmann et al., 2019) had used square-root transformations but we decided to employ a log-transformation and range-correction (i.e., dividing each SCR by the maximum SCR per participant). We used log-transformation and range-correction for SCR data because these transformations are standard practice in our laboratory and we strive for methodological consistency across different projects (e.g., Ehlers, Nold, Kuhn, Klingelhöfer-Jens, & Lonsdorf, 2020; Kuhn, Mertens, & Lonsdorf, 2016; Scharfenort, Menz, & Lonsdorf, 2016; Sjouwerman & Lonsdorf, 2020; Sjouwerman, Niehaus, & Lonsdorf, 2015). Additionally, log-transformed and range-corrected data are generally assumed to approximate a normal distribution more closely and exhibit lower error variance, which leads to larger effect sizes (Lykken, 1972; Lykken & Venables, 1971; Sjouwerman, Illius, Kuhn, & Lonsdorf, 2022). Additionally, on a descriptive level, this combination of transformations appear to offer greater reliability compared to using raw data alone (Klingelhöfer-Jens, Ehlers, Kuhn, Keyaniyan, & Lonsdorf, 2022).

Psychometric assessment

Participants completed a computerized battery of questionnaires (for a full list, see Stegmann et al., 2019) prior to the experiment including a questionnaire with general questions asking, for example, about the socioeconomic status (SES), the German versions of the trait version of the State-Trait Anxiety Inventory (STAI-T, Spielberger, 1983), the CTQ-SF (Bernstein et al., 2003; Wingenfeld et al., 2010) and the short version of the Center for Epidemiological Studies-Depression Scale (CES-D, in Germany: Allgemeine Depressionsskala - Kurzform, ADS-K, Hautzinger & Bailer, 1993) The CTQ contains 28 items for the retrospective assessment of

childhood adversity across five subscales (emotional, physical, and sexual abuse, as well as

emotional and physical neglect; for internal consistency, see Supplementary Material), and a

control scale. The STAI-T consists of 20 items addressing trait anxiety (Laux & Spielberger,

1981; Spielberger, 1983), and the ADS-K includes 15 items assessing depressiveness during the

past 7 days.

Operationalization of 'exposure'.   We implemented different approaches to

operationalize exposure to childhood adversity in the main analyses and exploratory analyses (see

Table 2). In the main analyses, we followed the approach most commonly employed in the field

of research on childhood adversity and threat learning - using the moderate exposure cut-off of

the CTQ (for a recent review see Ruge et al. (2024)). In addition, the heterogeneous

operationalizations of classifying individuals into exposed and unexposed to childhood adversity

in the literature (Koppold, Kastrinogiannis, Kuhn, & Lonsdorf, 2023; Ruge et al., 2024) hampers

comparison across studies and hence cumulative knowledge generation. Therefore, we also

provide exploratory analyses (see below) in which we employ different operationalizations of

childhood adversity exposure.

Table 2: *Operationalization of childhood adversity in different theoretical approaches and challenges of their statistical translation*

Approach name and reference	Operationalization of childhood adversity	Challenges in translating theory into a statistical model
Main analyses		
Moderate exposure based on CTQ (exposed vs. unexposed)	**Short description**: At least one subscale met the published cut-off for at least moderate exposure (Bernstein & Fink,1998; Häuser et al., 2011). The moderate cut-off was chosen, as it was recently identified as the most commonly used in the literature (for a review see Ruge et al., 2023). **Procedure:** Dichotomization of the sample into exposed vs. unexposed individuals based on published cut-offs: emotional abuse >= 13, physical abuse >= 10, sexual abuse >= 8, emotional neglect >= 15, physical neglect >= 10. Such cut-off of moderate exposure was employed in previous work by our team (Koppold et al., 2023) and in the literature (Ruge et al., 2023) **Statistical test:** See Materials and methods: Statistical analyses	• Not based on an existing theory but on what is commonly used in the literature (Ruge et al., 2023) • Different cut-offs published (for a discussion, see Ruge et al., 2023) • (Statistical) Challenges linked to dichotomization of an inherently continuous variable
Exploratory analyses		
Cumulative risk model	**Short description:** Based on the assumed key role	• Problem with CTQ sum score: it assigns the same 'value' to all

| (Evans et al., 2013; McEwen 2003) | of cumulative exposure (exposure intensity and frequency)

Procedure a): Classification into four severity groups (no, low, moderate, severe exposure) based on cut-offs published by Bernstein & Fink (1998)

Statistical test a): Comparison of conditioned responding of the four severity groups by using one-way ANOVAs

Procedure b): Number of subscales exceeding an at least moderate cut-off based on Bernstein & Fink (1998) and Häuser et al. (2011)

Statistical test b): Number of sub-scales exceeding an at least moderate cut-off as a predictor and conditioned responding as the criterion in simple linear regression models | CM types (see also 'General operationalizational challenges' below)
- Number of subscales exceeding cut-off: calculate ANOVA or regression?
- Cumulative risk scores are based on the implicit assumption that different types of adverse events affect the same mechanisms and are of equal impact |
| **Specificity model**

(McMahon et al., 2003; Pollak et al., 2000; Pollak & Tolley-Schell, 2004) | **Short description:** Consideration of specific exposure types (abuse vs. neglect)

Procedure: Summing up the CTQ subscales of emotional abuse, physical abuse, and sexual abuse yielding a composite score for exposure to 'abuse' and summing up the subscales of emotional neglect and physical neglect to yield a composite score for 'neglect' (or threat vs. deprivation as done by Sheridan et al., 2017)

Statistical test: The abuse and neglect composite | - What qualifies as a specific exposure type? (i.e., subscales or composite scales for neglect vs. abuse?)
- Which exposure subcategories are 'too specific' or 'too broad'? (A heterogeneous category may obscure potentially relevant discrete associations)
- Include only participants who experienced only one specific type but not any other types despite this being rather artificial due to high co-occurrences of different exposure types and requiring extremely large samples? Which cut-off should be used then to define exposure? We decided to include all participants in the |

	scores are tested for associations with conditioned responding in separate regression models. In our sample n = 52 and n = 96 individuals were exposed to abuse only and neglect only, respectively, while n = 55 reported to have experienced both abuse and neglect. We included all participants in all analyses as done previously (Sheridan et al., 2017)	analyses as done in previous studies (Sheridan et al., 2017) • Lack of specificity of exposure subtypes (e.g., sexual abuse also has an emotional component)
Dimensional model (McLaughlin et al. 2016; McLaughlin et al., 2021)	**Short description:** Consideration of specific exposure types (i.e., abuse and neglect) that are assumed to co-occur and be controlled for the effect of one another (as opposed to the specificity model) **Procedure:** See specificity model **Statistical test:** Abuse and neglect scores are tested for associations with conditioned responding in a single linear regression model in which the influence of the other exposure type is controlled for	• Ongoing debate on the multicollinearity of multiple types of childhood adversity in one model (McLaughlin et al., 2021; Pollak & Smith, 2021)
General operationalizational challenges	• Non-comparability of dimensional and categorical approaches: CTQ sum score assumes an equal contribution of all items which contradicts different thresholds for being considered as exposed for different subscales (e.g., lower cut-off for sexual abuse as compared to emotional neglect) • Associations in a full sample may differ from associations in the group of exposed individuals only which is a challenge for the interpretation of data • Multiple cut-offs published (Bernstein et al. 1997; Bernstein & Fink, 1998)	

	• Specific challenges relating to abuse and neglect: They
	    o often co-occur
	    o are not the only relevant dimensions (e.g., unpredictability, loss)
	    o are not strongly supported as distinct dimensions in the literature (Carozza et al., 2022; Smith & Pollak, 2021)
	• Heterogeneity in the assessment of childhood adversity across studies - both with respect to the assessment tools (e.g., questionnaires, interviews) as well as with respect to the operationalization of adversity (i.e., definition)
	• Different response formats (yes/no vs. specification of duration and frequency) and the number of trauma types/events included in assessment tools impact on prevalence rates and potentially also associations between the number of adverse experiences and symptom severity (e.g., Contractor et al., 2018)
	• Distinction between stressful events and trauma is often unclear (Richter-Levin & Sandi, 2021)

Statistical analyses

Manipulation checks were performed to test for successful fear acquisition and generalization (for more details, see Supplementary Material). Following previous studies (Imholze et al., 2023; Stegmann et al., 2019), we calculated three different outcomes for each participant for SCRs and ratings: CS discrimination (for acquisition training and the generalization phase), the linear deviation score (LDS; only for the generalization phase) as an index of the linearity of the generalization gradient (Kaczkurkin et al., 2017), and the general reactivity (across all phases including habituation, acquisition training and the generalization phase). CS discrimination was calculated by separately averaging responses to CS+ and CS- across trials (except the first acquisition trial) and subtracting averaged CS- responses from averaged CS+ responses. The first acquisition trial was excluded as no learning could possibly have taken place due to the delay conditioning paradigm. The LDS was calculated by subtracting the mean responses to all GSs from the mean responses to both CSs during the generalization phase. To calculate the general reactivity in SCRs and ratings, trials were averaged across all stimuli (CSs and GSs) and phases (i.e., habituation, acquisition training and generalization phase). Note that raw SCRs were used for analyses of general physiological reactivity.

CS discrimination during acquisition training and the generalization phase, LDS, and general reactivity were compared between participants who were exposed and unexposed to childhood adversity by using two-tailed independent-samples t-tests. For CS discrimination in SCRs, a two-way mixed ANOVA was conducted to examine the effect of childhood adversity exposure on responses to the CS+ and CS- by including CS type and childhood adversity exposure as independent variables. As the interaction between CS type and childhood adversity exposure was statistically significant, post hoc two-tailed paired t-tests were used to compare

SCRs between CS+ and CS- within each group and independent-samples t-tests to contrast responses to each CS between exposed and unexposed participants.

Exploratory analyses. Additionally, the different ways of classifying individuals as exposed or unexposed to childhood adversity in the literature (Koppold et al., 2023; for discussion see Ruge et al., 2024) hinder comparison across studies and hence cumulative knowledge generation. Therefore, we also conducted exploratory analyses using different approaches to operationalize exposure to childhood adversity (see Table 2 for details). Note that no correction for alpha inflation was applied in these analyses, given their exploratory nature. To compare the explanatory strengths of the included theories, all effect sizes from the exploratory tests were converted to the absolute value of Cohen's d as the direction is not relevant in this context. When their value fell outside the confidence intervals of the effect sizes of the main analysis (LeBel, McCarthy, Earp, Elson, & Vanpaemel, 2018), this was inferred as meaningful differences in explanatory strengths.

Analyses of trait anxiety and depression symptoms. To further characterize our sample, we compared individuals being unexposed to exposed to childhood adversity on trait anxiety and depression scores by using Welch tests due to unequal variances.

On the request of a reviewer, we additionally investigated the association of childhood adversity as operationalized by the different models used in our explanatory analyses (i.e., cumulative risk, specificity, and dimensional model) and trait anxiety as well as depression scores (see Supplementary Figure 7). By using STAI-T and ADS-K scores as independent variables, we calculated a) a comparison of conditioned responding of the four severity groups (i.e., no, low, moderate, severe exposure to childhood adversity) using one-way ANVOAs and the association with the number of sub-scales exceeding an at least moderate cut-off in simple linear regression

models for the implementation of the cumulative risk model, and b) the association with the CTQ

abuse and neglect composite scores in separate linear regression models for the implementation

of the specificity/dimensional models. On request of the reviewer, we also calculated the Pearson

correlation between trait anxiety (i.e., STAI-T scores), depression scores (i.e., ADS-K scores),

and conditioned responding in SCRs (see Supplementary Table 8).

In statistical procedures where the assumption of homogeneity of variance was not met,

Welch's tests, robust trimmed means ANOVAs (Mair & Wilcox, 2020), and regressions with

robust standard errors using the HC3 estimator (Hayes & Cai, 2007) were calculated instead of t-

tests, ANOVAs and regressions, respectively. Note that for robust mixed ANOVAs, the WRS2

package in R (Mair & Wilcox, 2020) does not provide an effect size. In the main analyses, post

hoc t-test or Welch's tests were corrected for multiple comparisons by using the Holm correction.

As post hoc tests for robust ANOVAs, Yuen independent samples t-test for trimmed means were

calculated including the explanatory measure of effect size (values of 0.10, 0.30, and 0.50

represent small, medium, and large effect sizes, respectively; Mair and Wilcox, 2020). Even

though such rules of thumb have to be interpreted with caution, we provide these benchmarks

here as this effect size might be somewhat unknown.

Following previous calls for a stronger focus on measurement reliability (Cooper,

Dunsmoor, Koval, Pino, & Steinman, 2022; Klingelhöfer-Jens et al., 2022), we also provide

information on split-half reliability for SCRs as well as Cronbach's alpha for the CTQ in the

Supplementary Material. For all statistical analyses described above, the a priori significance

level was set to $\alpha = 0.05$. For data analysis and visualizations as well as for the creation of the

manuscript, we used R (Version 4.1.3; R Core Team, 2022b) and the R-packages *apa* (Aust &

Barth, 2020; Version 0.3.3; Gromer, 2020), *car* (Version 3.0.10; Fox & Weisberg, 2019; Fox,

Weisberg, & Price, 2020), *carData* (Version 3.0.4; Fox et al., 2020), *chisq.posthoc.test* (Version

0.1.2; Ebbert, 2019), *cocor* (Version 1.1.3; Diedenhofen & Musch, 2015), *data.table* (Version

1.13.4; Dowle & Srinivasan, 2020), *DescTools* (Version 0.99.42; Andri et mult. al., 2021), *dplyr*

(Version 1.1.4; Wickham, François, Henry, & Müller, 2022), *effectsize* (Version 0.8.8; Ben-

Shachar, Lüdecke, & Makowski, 2020), *effsize* (Version 0.8.1; Torchiano, 2020), *ez* (Version

4.4.0; Lawrence, 2016), *flextable* (Version 0.9.6; Gohel, 2021), *forcats* (Version 0.5.0; Wickham,

2020), *foreign* (Version 0.8.82; R Core Team, 2022a), *ftExtra* (Version 0.6.4; Yasumoto, 2023),

GGally (Version 2.1.2; Schloerke et al., 2021), *ggExtra* (Version 0.10.0; Attali & Baker, 2022),

gghalves (Version 0.1.1; Tiedemann, 2020), *ggpattern* (Version 1.0.1; FC, Davis, & ggplot2

authors, 2022), *ggplot2* (Version 3.5.1; Wickham, 2016), *ggpubr* (Version 0.4.0; Kassambara,

2020), *ggsankey* (Version 0.0.99999; Sjoberg, 2023), *ggsignif* (Version 0.6.3; Constantin & Patil,

2021), *gridExtra* (Version 2.3; Auguie, 2017), *haven* (Version 2.3.1; Wickham & Miller, 2020),

here (Version 1.0.1; Müller, 2020), *kableExtra* (Version 1.3.1; Zhu, 2020), *knitr* (Version 1.37;

Xie, 2015), *lm.beta* (Version 1.5.1; Behrendt, 2014), *lme4* (Version 1.1.26; Bates, Mächler,

Bolker, & Walker, 2015), *lmerTest* (Version 3.1.3; Kuznetsova, Brockhoff, & Christensen,

2017), *lmtest* (Version 0.9.38; Zeileis & Hothorn, 2002), *MatchIt* (Version 4.4.0; Ho, Imai, King,

& Stuart, 2011), *Matrix* (Version 1.4.0; Bates & Maechler, 2021), *officedown* (Version 0.2.4;

Gohel & Ross, 2022), *papaja* (Version 0.1.2; Aust & Barth, 2020), *patchwork* (Version 1.2.0;

Pedersen, 2020), *performance* (Version 0.12.0; Lüdecke, Ben-Shachar, Patil, Waggoner, &

Makowski, 2021), *psych* (Version 2.0.9; Revelle, 2020), *purrr* (Version 1.0.2; Henry &

Wickham, 2020), *readr* (Version 2.1.4; Wickham & Hester, 2020), *reshape2* (Version 1.4.4;

Wickham, 2007), *rstatix* (Version 0.7.0; Kassambara, 2021), *sandwich* (Zeileis, 2004, 2006;

Version 3.0.1; Zeileis, Köll, & Graham, 2020), *sjPlot* (Version 2.8.16; Lüdecke, n.d.), *stringr*

(Version 1.5.1; Wickham, 2019), *tibble* (Version 3.2.1; Müller & Wickham, 2021), *tidyr* (Version

1.3.1; Wickham & Girlich, 2022), *tidyverse* (Version 1.3.0; Wickham et al., 2019), *tinylabels*

(Version 0.2.3; Barth, 2022), *WRS2* (Version 1.1.4; Mair & Wilcox, 2020b), and *zoo* (Version

1.8.8; Zeileis & Grothendieck, 2005).

Results

Exposed and unexposed participants were equally distributed across data recording sites

$(\chi^2(3) = 3.72, p = .293)$.

Main Effect of Task

In brief, and as reported previously (Herzog et al., 2021; Schiele, Reinhard et. al., 2016),

the fear acquisition was successful in SCRs as well as ratings in the full sample (all p's < 0.001;

see Supplementary Material for details). During fear generalization, the expected generalization

gradient was observed with a gradual increase in SCRs and ratings with increasing similarity to

the CS+ (all p's < 0.01 except for the comparisons of SCRs to CS- vs. GS4 as well as GS1

vs. GS2 which were non-significant; see Supplementary Material).

Association between different outcomes and exposure to childhood adversity

During both the acquisition training and generalization phase, CS discrimination in SCRs

was significantly lower in individuals exposed to childhood adversity as compared to unexposed

individuals (see Table 3 and Figure 1; for trial-by-trial responses, see Supplementary Figure 4).

Post hoc analyses (i.e., ANOVAs) revealed that childhood adversity exposure significantly

interacted with stimulus type (acquisition training: $F(1, 1400) = 5.42, p = .020, \eta_p^2 < .01$;

generalization test: $F(1, 1051) = 5.37, p = 0.021$): SCRs to the CS+ during both acquisition

training and the generalization phase were significantly lower in exposed as compared to

unexposed individuals (acquisition training: $t(1400) = 2.54$, $p = .011$, $d = 0.14$; generalization

test: $t(194.1) = 3.51$, $p = 0.001$, *explanatory measure of effect size* = 0.179; see Figure 2) but not

for the CS- (acquisition training: $t(1400) = 0.75$, $p = .452$, $d = 0.04$; generalization test: $t(178.9) =$

1.63, $p = 0.104$, *explanatory measure of effect size* = 0.09). For ratings, no significant effects of

exposure to childhood adversity were observed in CS discrimination (see Table 3).

No significant effect of exposure to childhood adversity in either SCRs or ratings was

observed for generalization gradients (see Table 3 and Figure 3). It is, however, also evident from

the generalization gradients that both groups differ specifically in reactivity to the CS+ (see

above and Figure 3).

In addition, general physiological reactivity in SCRs (i.e., raw amplitudes) was

significantly lower in participants exposed to childhood adversity compared to unexposed

participants (see Table 3 and Figure 4) while there were no differences between both groups in

general rating response levels (see Table 3).

At the request of a reviewer, we repeated our main analyses by using linear mixed models

including age, sex, school degree (i.e., to approximate socioeconomic status), and exposure to

childhood adversity as fixed effects as well as site as random effect. These analyses yielded

comparable results demonstrating a significant effect of childhood adversity on CS discrimination

during acquisition training and the generalization phase as well as on general reactivity, but not

on the generalization gradients in SCRs (see Supplementary Table 2 A). Consistent with the

results of the main analyses reported in our manuscript, we did not observe any significant effects

of childhood adversity on the different types of ratings when using mixed models (see

Supplementary Table 2 B-D). Some of the mixed model analyses showed significantly lower CS

discrimination during acquisition training and generalization, and lower general reactivity in males compared to females (see Supplementary Table 2 for details).

Table 3: Results of t-tests comparing CS discrimination, the linear deviation score (i.e., strength of generalization), and general reactivity between exposed and unexposed participants

Outcome	Phase	Measure	t	df	p	Cohen's d	LL (95% CI)	UL (95% CI)
CS discrimination	ACQ	SCR	**2.33**	**1,400**	**0.020**	**-0.18**	**-0.33**	**-0.03**
		Arousal ratings	-1.52	1,400	0.128	0.12	-0.03	0.26
		Valence ratings	0.20	1,400	0.845	-0.01	-0.16	0.13
		Contigency ratings	0.70	1,400	0.484	-0.05	-0.20	0.10
	GEN	SCR	**2.34**	**1,400**	**0.020**	**-0.18**	**-0.33**	**-0.03**
		Arousal ratings	-0.28	1,400	0.777	0.02	-0.13	0.17
		Valence ratings	0.06	1,400	0.953	0.00	-0.15	0.14
		Contigency ratings	0.58	1,400	0.560	-0.04	-0.19	0.10
LDS	GEN	SCR	1.41	295	0.158	-0.10	-0.25	0.05
		Arousal ratings	-0.62	1,400	0.538	0.05	-0.10	0.20
		Valence ratings	0.30	1,400	0.765	-0.02	-0.17	0.13
		Contigency ratings	-0.95	1,400	0.344	0.07	-0.08	0.22
General reactivity	ALL	SCR	**2.06**	**1,400**	**0.040**	**-0.16**	**-0.31**	**-0.01**
		Arousal ratings	-0.10	1,400	0.920	0.01	-0.14	0.16
		Valence ratings	0.83	1,400	0.408	-0.06	-0.21	0.09
		Contigency ratings	-0.97	250	0.334	0.07	-0.09	0.24

Note. ACQ = acquisition training, GEN = generalization phase, LDS = linear deviation score. Bold numbers indicate significant results ($p < 0.05$).

Figure 1: Illustration of CS discrimination in SCRs during acquisition training (A-B) and generalization phase (C-D) for individuals unexposed (gray) and exposed (pink) to childhood adversity. Barplots (A and C) with error bars represent means and standard errors of the means (SEMs) including $n_{unexposed} = 1199$ and $n_{exposed} = 203$, respectively. The statistical parameters presented in A) and C) are derived from two-tailed independent-samples t-tests. The a priori significance level was set to $\alpha = 0.05$. Distributions of the data are illustrated in the raincloud plots (B and D). Points next to the densities represent the CS discrimination of each participant averaged across phases. Boxes of boxplots represent the interquartile range (IQR) crossed by the median as a bold line, ends of whiskers represent the minimum/maximum value in the data within the range of 25th/75th percentiles ± 1.5 IQR. For trial-by-trial SCRs across all phases, see Supplementary Figure 4. log = log-transformed, rc = range-corrected.

Figure 2: Illustration of SCRs during acquisition training (A-B) and the generalization phase (C-D) for individuals unexposed (gray) and exposed (pink) to childhood adversity separated by stimulus types (CS+ dark shades, CS-: light shades). Barplots (A and C) with error bars represent means and SEMs including $n_{unexposed} = 1199$ and $n_{exposed} = 203$, respectively. The presented statistical parameters are derived from a two-tailed independent-samples t-test (A) and a Yuen

independent-samples t-test for trimmed means (C). The a priori significance level was set to α = 0.05. Distributions of the data are illustrated in the raincloud plots (B and D). Points next to the densities represent the SCRs of each participant as a function of stimulus type averaged across phases. Boxes of boxplots represent the IQR crossed by the median as a bold line, ends of whiskers represent the minimum/maximum value in the data within the range of 25th/75th percentiles ±1.5 IQR. CS = conditioned stimulus, log = log-transformed, rc = range-corrected.

Figure 3: Illustration of SCRs to the different stimulus types during the generalization phase (i.e., generalization gradients) for individuals unexposed (gray) and exposed (pink) to childhood adversity. Ribbons represent SEMs including $n_{unexposed}$ = 1199 and $n_{exposed}$ = 203, respectively. CS = conditioned stimulus, GS = generalization stimuli with gradual perceptual similarity to the CS+ and CS-, respectively. log = log-transformed, rc = range-corrected. * $p < .05$.

Figure 4: Illustration of general reactivity in SCRs across all experimental phases for individuals

unexposed (gray) and exposed (pink) to childhood adversity. Barplots (A) with error bars

represent means and SEMs including $n_{unexposed} = 1199$ and $n_{exposed} = 203$, respectively. The

statistical parameters presented in A) are derived from a two-tailed independent-samples t-test.

The a priori significance level was set to $\alpha = 0.05$. Distributions of the data are illustrated in the

raincloud plots (B). Points next to the densities represent the general reactivity of each participant

averaged across all phases. Boxes of boxplots represent the interquartile range IQR crossed by

the median as a bold line, ends of whiskers represent the minimum/maximum value in the data

within the range of 25th/75th percentiles ±1.5 IQR.

Exploratory analyses

The cumulative risk model operationalized through the different cut-offs for no, low,

moderate and severe exposure (Bernstein & Fink, 1998) did not yield any significant results for

any outcome measure and experimental phase (see Supplementary Table 3). However, on a

descriptive level (see Figure 5), it seems that indeed exposure to an at least moderate cut-off level

may induce behavioral and physiological changes (see main analysis, Bernstein & Fink, 1998).

This might suggest that the cut-off for exposure commonly applied in the literature (see Ruge et

al., 2024) may indeed represent a reasonable approach.

Cumulative risk operationalized as the number of CTQ subscales exceeding the moderate

cut-off (Bernstein & Fink, 1998), however, revealed that a higher number of subscales exceeding

the cut-off predicted lower CS discrimination in SCRs ($F(1, 1400) = 6.86$, $p = 0.009$, $R^2 = 0.005$)

and contingency ratings ($F(1, 1400) = 4.08$, $p = 0.044$, $R^2 = 0.003$) during acquisition training

(see Figure 6 and Supplementary Table 4 for an exemplary illustration of SCRs during

acquisition training). This was driven by significantly lower SCRs to the CS+ ($F(1, 1400) = 5.42$,

$p = 0.02$, $R^2 = 0.004$) while for contingency ratings no significant post hoc tests were identified

(all $p > 0.05$). For an illustration, of how the different adversity types (i.e., subscales) are

distributed among the different numbers of subscales, see Supplementary Figure 5.

The operationalization of childhood adversity in the context of the specificity model tests

the association between exposure to abuse and neglect experiences on conditioned responding

statistically independently, while the dimensional model controls for each other's impact (see

Table 2 for details and Figure 7 for an exemplary illustration of SCRs during acquisition

training). Despite these conceptual and operationalizational differences, results are converging.

More precisely, no significant effect of exposure to abuse was observed on CS discrimination, the

strength of generalization (i.e., LDS), or general reactivity in any of the outcome measures and in

any experimental phase (see Supplementary Table 5 and 7). In contrast, a significant negative

association between exposure to neglect and CS discrimination in SCRs was observed during

acquisition training (specificity model: $F(1, 1400) = 6.4$, $p = 0.012$, $R^2 = 0.005$; dimensional

model: $F(3, 1398) = 2.91$, $p = 0.234$, $R^2 = 0.006$), which is contrary to the predictions of the dimensional model, that posits a specific role for abuse but not neglect (Machlin et al., 2019; McLaughlin et al., 2021). Post hoc tests revealed that in both models, effects were driven by significantly lower SCRs to the CS+ (specificity model: $F(1, 1400) = 6.13$, $p = 0.013$, $R^2 = 0.004$, dimensional model: $\beta = -0.004$, $t(1398) = -1.97$, $p = 0.049$, $r = -0.07$). Within the dimensional model framework, the issue of multicollinearity among predictors (i.e., different childhood adversity types) is frequently discussed (McLaughlin et al., 2021; Smith & Pollak, 2021). If we apply the rule of thumb of a variance inflation factor (VIF) > 10, which is often used in the literature to indicate concerning multicollinearity (e.g., Hair, Anderson, Tatham, & Black, 1995; Mason, Gunst, & Hess, 1989; Neter, Wasserman, & Kutner, 1989), we can assume that multicollinearity was not a concern in our study (abuse: VIF = 8.64; neglect: VIF = 7.93). However, some authors state that VIFs should not exceed a value of 5 (e.g., Akinwande, Dikko, and Samson (2015)), while others suggest that these rules of thumb are rather arbitrary (O'brien, 2007).

Furthermore, the statistical analyses of the specificity model additionally revealed that greater exposure to neglect significantly predicted a generally lower SCR reactivity ($F(1, 1400) = 4.3$, $p = 0.038$, $R^2 = 0.003$) as well as lower a CS discrimination in contingency ratings during both acquisition training ($F(1, 1400) = 5.58$, $p = 0.018$, $R^2 = 0.004$) and the generalization test ($F(1, 1400) = 6.33$, $p = 0.012$, $R^2 = 0.005$; see Supplementary Table 6). These were driven by significantly higher CS- responding in contingency ratings (acquisition training: $F(1, 1400) = 4.62$, $p = 0.032$, $R^2 = 0.003$; generalization test: $F(1, 1400) = 8.38$, $p = 0.004$, $R^2 = 0.006$) in individuals exposed to neglect.

To explore the explanatory power of different theories, we exemplarily compared the absolute values of Cohen's d of all exploratory analyses including CS discrimination in SCRs during acquisition training with the absolute values of the Cohen's d confidence intervals of our main analyses. We chose CS discrimination during fear acquisition training for this test, because the most convergent results across theories were observed during this experimental phase. None of the effect sizes from the exploratory analyses (cumulative risk, severity groups: d = 0.14; cumulative risk, number of subscales exceeding an at least moderate cut-off: d = 0.20; specificity model, abuse: d = 0.10; specificity model, neglect: d = 0.19; dimensional model: d = 0.18) fell outside the confidence intervals of our main results (i.e., an at least moderate childhood adversity exposure: [0.03; 0.33]). Hence, we found no evidence of differential explanatory strengths among theories.

Figure 5: Means and standard errors of the mean of CS discrimination in SCRs during acquisition training (A) and the generalization phase (B), Linear deviation score (LDS) (C), and general reactivity in SCRs (D) for the four CTQ severity groups respectively. The dashed line indicates the moderate CTQ cut-off frequently used in the literature and hence also employed in our main analyses: On a descriptive level, CS discrimination in SCRs during acquisition training and generalization test as well as the strength of generalization (i.e., LDS) and the general reactivity are lower in all groups exposed to childhood adversity at an at least moderate level as compared to those with no or low exposure - which corresponds to the main analyses (see above). log = log-transformed, rc = range-corrected.

A

B

C

Figure 6: Scatterplots with marginal densities illustrating the associations between the number of CTQ subscales exceeding a moderate or higher cut-off (Häuser et al., 2011) and CS discrimination in SCRs (A) as well as SCRs to the CS+ (B) and CS- (C) during acquisition training. log = log-transformed, rc = range-corrected.

Figure 7: Scatterplots with marginal densities illustrating the associations between CTQ composite scores of abuse (left panel) and neglect (right panel) and CS discrimination in SCRs (A and B) as well as SCRs to the CS+ (C and D) and CS- (E and F) during acquisition training. Note that the different ranges of CTQ composite scores result from summing up two and three subscales for the neglect and abuse composite scores, respectively (see also Table 2 for more details). log = log-transformed, rc = range-corrected, R squ. = R squared.

Analyses of trait anxiety and depression symptoms

As expected, participants exposed to childhood adversity reported significantly higher trait anxiety and depression levels than unexposed participants (all p's < 0.001; see Table 1 and Supplementary Figure 6). This pattern remained unchanged when childhood adversity was operationalized differently - following the cumulative risk approach, the specificity, and dimensional model (see methods). These additional analyses all indicated a significant positive relationship between exposure to childhood adversity and trait anxiety as well as depression scores irrespective of the specific operationalization of 'exposure' (see Supplementary Figure 7).

CS discrimination during acquisition training and the generalization phase, generalization gradients, and general reactivity in SCRs were unrelated to trait anxiety and depression scores in this sample with the exception of a significant association between depression scores and CS discrimination during fear acquisition training (see Supplementary Table 8). More precisely, a very small but significant negative correlation was observed indicating that high levels of depression were associated with reduced levels of CS discrimination (r = -0.057, p = 0.033). The correlation between trait anxiety levels and CS discrimination during fear acquisition training was not statistically significant but on a descriptive level, high trait anxiety scores were also

linked to lower CS discrimination scores (r = -0.05, p = 0.06) although we highlight that this should not be overinterpreted in light of the large sample. However, both correlations (i.e., CS-discrimination during fear acquisition training and trait anxiety as well as depression, respectively) did not statistically differ from each other (z = 0.303, p = 0.762, Dunn & Clark, 1969). Interestingly, and consistent with our results showing that the relationship between childhood adversity and CS discrimination was mainly driven by significantly lower CS+ responses in exposed individuals, trait anxiety and depression scores were significantly associated with SCRs to the CS+, but not to the CS- during acquisition training (see Supplementary Table 8).

Discussion

The objective of this study was to examine the relationship between the exposure to childhood adversity and conditioned responding using a large community sample of healthy participants. This relationship might represent a potential mechanistic route linking experience-dependent plasticity in the nervous system and behavior related to risk of and resilience to psychopathology. In additional exploratory analyses, we examined these associations through different approaches by translating key theories in the literature into statistical models. In line with the conclusion of a recent systematic literature review (Ruge et al., 2024), individuals exposed to (an at least moderate level of) childhood adversity exhibited reduced CS discrimination in SCRs during both acquisition training and the generalization phase compared to those classified as unexposed (i.e., no or low exposure). Generalization gradients themselves were, however, comparable between exposed and unexposed individuals.

The systematic literature search by Ruge et al. (2024) revealed that the pattern of decreased CS discrimination, driven primarily by reduced CS+ responding, was observed despite substantial heterogeneity in childhood adversity assessment and operationalization, and despite differences in the experimental paradigms. Although both individuals without mental disorders exposed to childhood adversity and patients suffering from anxiety- and stress-related disorders (e.g., Duits et al., 2015) show reduced CS discrimination, it is striking that the response pattern of individuals exposed to childhood adversity (i.e., reduced responding to the CS+) is remarkably different from what is typically observed in patients (i.e., enhanced responding to the CS-). It should be noted, however, that childhood adversity exposure status was not considered in this meta-analysis (Duits et al., 2015). As exposure to childhood adversity represents a particularly strong risk factor for the development of later psychopathology, these seemingly contrary findings warrant an explanation. In this context, it is important to note that all individuals included in the present study were mentally healthy - at least up to the assessment. Hence, it may be an obvious explanation that reduced CS discrimination driven by decreased CS+ responding may represent a resilience rather than a risk factor because individuals exposed to childhood adversity in our sample are mentally healthy despite being exposed to a strong risk factor.

In fact, there is substantial heterogeneity in individual trajectories and profiles in the aftermath of such exposures in humans and rodents (Russo, Murrough, Han, Charney, & Nestler, 2012). While some individuals remain resilient despite exposure, others develop psychopathological conditions (Galea, Nandi, & Vlahov, 2005). Consequently, the sample of the present study may represent a specific subsample of exposed individuals who are developing along a resilient trajectory. Thus, it can be speculated that reduced physiological reactivity to a signal of threat (e.g., CS+) may protect the individual from overwhelming physiological and/or emotional responses to potentially recurrent threats (for a discussion, see Ruge et al. (2024)).

Similar concepts have been proposed as 'emotional numbing' in post-traumatic stress disorders (for a review, see e.g., Litz & Gray, 2002).

While this seems a plausible theoretical explanation, decreased CS discrimination driven by reduced CS+ responding is observed rather consistently and most importantly irrespective of whether the investigated samples were healthy, at risk, or included patients (Ruge et al., 2024). Thus, this response pattern which was also observed in our study might be a specific characteristic of childhood adversity exposure distinct from the response pattern generally observed in patients suffering from anxiety- and stress-related disorders (i.e., increased responding to the CS-) - even though individuals exposed to childhood adversity in this sample indeed showed significantly higher anxiety and depression despite being free of any categorical diagnoses (see Supplementary Material) which was also previously reported by e.g., Spinhoven et al. (2011) and Kuhn et al. (2016). Interestingly, in our study, trait anxiety and depression scores were mostly unrelated to SCRs, defined by CS discrimination and generalization gradients based on SCRs as well as general SCR reactivity, with the exception of a significant - albeit minute - relationship between CS discrimination during acquisition training and depression scores (see above). Although reported associations in the literature are heterogeneous (Lonsdorf et al., 2017), we may speculate that they may be mediated by childhood adversity. We conducted additional mediation analyses (data not shown) which, however, did not support this hypothesis. As the potential links between reduced CS discrimination in individuals exposed to childhood adversity and the developmental trajectories of psychopathological symptoms are still not fully understood, future work should investigate these further in - ideally - prospective studies.

In addition to reduced CS discrimination in SCRs, a generally blunted electrodermal responding was observed, which may, however, be mainly driven by substantially reduced CS+

responses. Yet, it is noteworthy that reduced skin conductance in children exposed to childhood adversity was also observed during other tasks such as attention regulation during interpersonal conflict (Pollak, Vardi, Putzer Bechner, & Curtin, 2005), or passively viewing slides with emotional or cognitive content (Carrey, Butter, Persinger, & Bialik, 1995), whereas other studies did not find such an association (Ben-Amitay, Kimchi, Wolmer, & Toren, 2016). Moreover, in various threat-related studies, also enhanced responding or no significant differences were observed across outcome measures (Estrada, Richards, Gee, & Baskin-Sommers, 2020; Huskey, Taylor, & Friedman, 2022; Jovanovic et al., 2009, 2022; Kreutzer & Gorka, 2021; Lis et al., 2020; Pole et al., 2007; Rowland et al., 2022; Thome et al., 2018; D. A. Young et al., 2018). While generally blunted responding might be particularly related to decreased CS+ responding in the present study, differences in general reactivity need to be taken into account for data analyses and interpretation - in particular as the exclusion of physiological non-responders or so-called 'non-learners' (i.e., individuals not showing a minimum discrimination score between SCRs to the CS+ and CS-) has been common in the field until recently (for a critical discussion, see Lonsdorf et al., 2019). Future work should also investigate reactivity to the unconditioned stimulus, which was not implemented here and may shed light on potential differences in general reactivity unaffected by associative learning processes (see e.g., Harnett et al., 2019; Machlin et al., 2019).

Contrary to the association between CS discrimination as well as general (electrodermal) reactivity and exposure to childhood adversity, no such relationship was found for generalization gradients. In a subsample of this study, it was previously observed that fear generalization phenotypes explained less variance as compared to CS discrimination and general reactivity (Stegmann et al., 2019), and CS discrimination as well as general reactivity but not fear generalization predicted increases in anxiety and depression scores during the COVID-19

pandemic (Imholze et al., 2023). The lack of associations with fear generalization measures (i.e., LDS) may be specific to the paradigm and sample used in these studies, but it may also be an interesting lead for future work to disentangle the relationship between CS discrimination, general reactivity, and generalization gradients, as they have been suggested to be interrelated (Imholze et al., 2023; Stegmann et al., 2019).

In sum, the current results converge with the literature in identifying reduced CS discrimination and decreased CS+ responding as key characteristics in individuals exposed to childhood adversity. As highlighted recently (see Koppold et al., 2023; Ruge et al., 2024), the various operationalizations of childhood adversity as well as general trauma (Karstoft & Armour, 2023) represent a challenge for integrating the current results into the existing literature. Hence, future studies should focus on in-depth phenotyping (Ruge et al., 2024), an elaborate classification of adversity subtypes (Pollak & Smith, 2021), and methodological considerations (Ruge et al., 2024). Besides optimizing the operationalization of childhood adversity, there are also initiatives to advance the field in data processing and analysis, such as developing methods to address multicollinearity in childhood adversity data (Brieant, Sisk, Keding, Cohodes, & Gee, 2024).

Several proposed (verbal) theories describe the association between (specific) childhood adversity types and behavioral as well as physiological consequences differently and there is currently a heated debate rather than consensus on this issue (McLaughlin et al., 2021; Pollak & Smith, 2021; Smith & Pollak, 2021). In the field, most often a dichotomization in exposed vs. unexposed individuals is used (for a review, see Ruge et al., 2024). We adopted this typical approach of an at least moderate exposure cut-off from the literature for our main analyses, despite the well-known statistical disadvantages of artificially dichotomizing variables that are

(presumably) dimensional in nature (Cohen, 1983). It is noteworthy, however, that this cut-off appears to map rather well onto the psychophysiological response patterns observed here (see Figure 5). More precisely, our exploratory results of applying different exposure cut-offs (low, moderate, severe, no exposure) seem to indicate that indeed a moderate exposure level is 'required' for the manifestation of physiological differences, suggesting that childhood adversity exposure may not have a linear or cumulative effect.

Of note, comparing individuals exposed vs. unexposed to an at least moderate level of childhood adversity is not derived from any of the existing theories, but rather from practices in the literature (see Ruge et al., 2024). For this reason, we aimed at an exploratory translation of key (verbal) theories into statistical models (see Table 2). Several important topical and methodological take-home messages can be drawn from this endeavor: First, the translation of these verbal theories into precise statistical tests proved to be a rather challenging task paved by operationalizational ambiguity. We have collected some key challenges in Table 2 and conclude that current verbal theories are, at least to a certain degree, ill-defined, as our attempt has disclosed a multiverse of different, equally plausible ways to test them - even though we provide only a limited number of exemplary tests. Second, despite these challenges, the results of most tests converged in identifying an effect of childhood adversity on reduced CS discrimination in SCRs during acquisition training, which is reassuring when aiming to integrate results based on different operationalizations. Third, none of the theories appears to be explanatorily superior. Fourth, our results are not in line with predictions of the dimensional model (Machlin et al., 2019; McLaughlin et al., 2021) which posits a specific association between exposure to threat- but not deprivation-related childhood adversity and fear conditioning performance. If anything, our results point in the opposite direction.

Taken together, neither considering childhood adversity as a broad category, nor different subtypes have consistently shown to strongly map onto biological mechanisms (for an in-depth discussion, see Smith & Pollak, 2021). Hence, even though it is currently the dominant view in the field, that considering the potentially distinct effects of dissociable adversity types holds promise to provide mechanistic insights into how childhood adversity becomes biologically embedded (Berens, Jensen, & Nelson, 2017; Kuhlman, Chiang, Horn, & Bower, 2017; Smith & Pollak, 2021), we emphasize the urgent need for additional exploration, refinement, and testing of current theories. This is particularly important in light of diverging evidence pointing towards different conclusions.

Some limitations of this work are worth noting: First, despite our observation of significant associations between exposure to childhood adversity and fear conditioning performance in a large sample, it should be noted that effect sizes were small. Second, we cannot provide a comparison of potential group differences in unconditioned responding to the US. This is, however, important as this comparison may explain group differences in conditioned responding - a mechanism that remains unexplored to date (Ruge et al., 2024). Third, the use of the CTQ, which is the most commonly used questionnaire in the field (see Ruge et al., 2024), comes with a number of disadvantages. Most prominently, the CTQ focuses exclusively on the presence or absence of exposure without consideration of individual and exposure characteristics that have been shown to be of crucial relevance (Danese & Widom, 2023; see Smith & Pollak, 2021), such as controllability, burdening, exposure severity, duration, and developmental timing. These characteristics are embedded in the framework of the topological approach (Smith & Pollak, 2021), another important model linking childhood adversity exposure to negative outcomes, which, however, was not evaluated in the present work. Testing this model requires an extremely large dataset including in-depth phenotyping, which was not available here, but may be

an important avenue for future work. Fourth, across all theories, significant effects of childhood adversity have been shown primarily on physiological reactivity (i.e., SCR). Whether these findings are specific to SCRs or might generalize to other physiological outcome measures such as fear-potentiated startle, heart rate, or local changes in neural activation, remains an open question for future studies.

In sum, when ultimately aiming to understand the impact of exposure to adversity on the development of psychopathological symptoms (Anda et al., 2006; Felitti, 2002; Gilbert et al., 2009; Green et al., 2010; Heim & Nemeroff, 2001; McLaughlin et al., 2012; Moffitt et al., 2007; Teicher et al., 2022), it is crucial to understand the biological mechanisms through which exposure to adversity 'gets under the skin'. To achieve this, emotional-associative learning can serve as a prime translational model for fear and anxiety disorders: One plausible mechanism is the ability to distinguish threat from safety, which is key to an individual's ability to dynamically adapt to changing environmental demands (Craske et al., 2012; Vervliet, Craske, & Hermans, 2013) - an ability that appears to be impaired in individuals with a history of childhood adversity. This mechanism is of particular relevance to the development of stress- and anxiety-related psychopathology, as the identification of risk but also resilience factors following exposure to childhood adversity is essential for the development of effective intervention and prevention programs.

Data and Code Availability Statement

The data will be made available to editors and reviewers only, as publicly sharing of individual-level data was not included in the informed consent forms. Instead, the forms specified that the data would be published anonymously as a collective dataset. At the time the study was planned, data sharing was not a common practice. Therefore, participants were not asked to consent to individual-level data sharing and were assured that their data would be used exclusively for the purposes specified in the consent forms. This restriction also applies to de-identified and processed versions of the individual-level data. R Markdown files that include the code for all analyses and generate this manuscript are openly available at Zenodo (https://doi.org/ https://doi.org/10.5281/zenodo.14851004; Klingelhöfer-Jens et al., 2023)

Acknowledgements

The authors thank Julia Ruge for critical reviewing. KD, MAS and PZ are members of the Anxiety Disorders Research Network (ADRN) of the European College of Neuropsychopharmacology (ECNP).

Funding

This work was supported by the German Research Foundation (DFG) – project number 44541416 – TRR 58 'Fear, Anxiety, Anxiety Disorders', subproject Z02 to JD, KD, UD, TBL, UL, AR, MR, PP; subproject B01 to PP, subproject B07 to TBL, subproject C02 to KD and JD, subproject C10 to MG.

Conflict of Interest

The authors declare no competing financial interests.

References

Akinwande, M. O., Dikko, H. G., & Samson, A. (2015). Variance Inflation Factor: As a

Condition for the Inclusion of Suppressor Variable(s) in Regression Analysis. *Open

Journal of Statistics*, *05*(07), 754–767. https://doi.org/10.4236/ojs.2015.57075

Anda, R. F., Felitti, V. J., Bremner, J. D., Walker, J. D., Whitfield, Ch., Perry, B. D., … Giles, W.

H. (2006). The enduring effects of abuse and related adverse experiences in childhood: A

convergence of evidence from neurobiology and epidemiology. *European Archives of

Psychiatry and Clinical Neuroscience*, *256*(3), 174–186. https://doi.org/10.1007/s00406-

005-0624-4

Andri et mult. al., S. (2021). *DescTools: Tools for descriptive statistics*. Retrieved from

https://cran.r-project.org/package=DescTools

Attali, D., & Baker, C. (2022). *ggExtra: Add marginal histograms to 'ggplot2', and more

'ggplot2' enhancements*. Retrieved from https://CRAN.R-project.org/package=ggExtra

Auguie, B. (2017). *gridExtra: Miscellaneous functions for "grid" graphics*. Retrieved from

https://CRAN.R-project.org/package=gridExtra

Aust, F., & Barth, M. (2020). *papaja: Prepare reproducible APA journal articles with R

Markdown*. Retrieved from https://github.com/crsh/papaja

Baldwin, J. R., Coleman, O., Francis, E. R., & Danese, A. (2024). Prospective and Retrospective

Measures of Child Maltreatment and Their Association With Psychopathology: A

Systematic Review and Meta-Analysis. *JAMA Psychiatry*.

https://doi.org/10.1001/jamapsychiatry.2024.0818

Barth, M. (2022). *tinylabels: Lightweight variable labels*. Retrieved from https://cran.r-project.org/package=tinylabels

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, *67*(1), 1–48. https://doi.org/10.18637/jss.v067.i01

Bates, D., & Maechler, M. (2021). *Matrix: Sparse and dense matrix classes and methods*. Retrieved from https://CRAN.R-project.org/package=Matrix

Behrendt, S. (2014). *Lm.beta: Add standardized regression coefficients to lm-objects*. Retrieved from https://CRAN.R-project.org/package=lm.beta

Ben-Amitay, G., Kimchi, N., Wolmer, L., & Toren, P. (2016). Psychophysiological Reactivity in Child Sexual Abuse. *Journal of Child Sexual Abuse*, *25*(2), 185–200. https://doi.org/10.1080/10538712.2016.1124309

Ben-Shachar, M. S., Lüdecke, D., & Makowski, D. (2020). effectsize: Estimation of effect size indices and standardized parameters. *Journal of Open Source Software*, *5*(56), 2815. https://doi.org/10.21105/joss.02815

Berens, A. E., Jensen, S. K. G., & Nelson, C. A. (2017). Biological embedding of childhood adversity: From physiological mechanisms to clinical implications. *BMC Medicine*, *15*(1), 135. https://doi.org/10.1186/s12916-017-0895-4

Bernstein, D. P., Ahluvalia, T., Pogge, D., & Handelsman, L. (1997). Validity of the Childhood Trauma Questionnaire in an Adolescent Psychiatric Population. *Journal of the American*

Academy of Child & Adolescent Psychiatry, *36*(3), 340–348.

https://doi.org/10.1097/00004583-199703000-00012

Bernstein, D. P., & Fink, L. (1998). *Childhood Trauma Questionnaire: A retrospective self-report manual.* San Antonio, TX: The Psychological Corporation.

Bernstein, D. P., Stein, J. A., Newcomb, M. D., Walker, E., Pogge, D., Ahluvalia, T., … Zule, W. (2003). Development and validation of a brief screening version of the Childhood Trauma Questionnaire. *Child Abuse & Neglect*, *27*(2), 169–190. https://doi.org/10.1016/S0145-2134(02)00541-0

Boucsein, W., Fowles, D. C., Grimnes, S., Ben-Shakhar, G., roth, W. T., Dawson, M. E., … Society for Psychophysiological Research Ad Hoc Committee on Electrodermal Measures. (2012). Publication recommendations for electrodermal measurements. *Psychophysiology*, *49*(8), 1017–1034. https://doi.org/10.1111/j.1469-8986.2012.01384.x

Brieant, A., Sisk, L. M., Keding, T. J., Cohodes, E. M., & Gee, D. G. (2024). Leveraging multivariate approaches to advance the science of early-life adversity. *Child Abuse & Neglect*, 106754. https://doi.org/10.1016/j.chiabu.2024.106754

Carozza, S., Holmes, J., & Astle, D. E. (2022). Testing Deprivation and Threat: A Preregistered Network Analysis of the Dimensions of Early Adversity. *Psychological Science*, *33*(10), 1753–1766. https://doi.org/10.1177/09567976221101045

Carrey, N. J., Butter, H. J., Persinger, M. A., & Bialik, R. J. (1995). Physiological and cognitive correlates of child abuse. *Journal of the American Academy of Child and Adolescent Psychiatry*, *34*(8), 1067–1075. https://doi.org/10.1097/00004583-199508000-00017

Cohen, J. (1983). The Cost of Dichotomization. *Applied Psychological Measurement*, *7*(3), 249–
253. https://doi.org/10.1177/014662168300700301

Colich, N. L., Rosen, M. L., Williams, E. S., & McLaughlin, K. A. (2020). Biological aging in
childhood and adolescence following experiences of threat and deprivation: A systematic
review and meta-analysis. *Psychological Bulletin*, *146*(9), 721–764.
https://doi.org/10.1037/bul0000270

Constantin, A.-E., & Patil, I. (2021). ggsignif: R package for displaying significance brackets for
'ggplot2'. *PsyArxiv*. https://doi.org/10.31234/osf.io/7awm6

Contractor, A. A., Brown, L. A., & Weiss, N. H. (2018). Relation between lifespan polytrauma
typologies and post-trauma mental health. *Comprehensive Psychiatry*, *80*, 202–213.
https://doi.org/10.1016/j.comppsych.2017.10.005

Cooper, S. E., Dunsmoor, J. E., Koval, K. A., Pino, E. R., & Steinman, S. A. (2022). TEST–
RETEST reliability of human threat conditioning and generalization across a 1-TO-2-WEEK
interval. *Psychophysiology*. https://doi.org/10.1111/psyp.14242

Cooper, S. E., van Dis, E. A. M., Hagenaars, M. A., Krypotos, A.-M., Nemeroff, C. B., Lissek,
S., … Dunsmoor, J. E. (2022). A meta-analysis of conditioned fear generalization in
anxiety-related disorders. *Neuropsychopharmacology*. https://doi.org/10.1038/s41386-
022-01332-2

Craske, M. G., Wolitzky-Taylor, K. B., Mineka, S., Zinbarg, R., Waters, A. M., Vrshek-
Schallhorn, S., … Ornitz, E. (2012). Elevated responding to safe conditions as a specific
risk factor for anxiety versus depressive disorders: Evidence from a longitudinal

investigation. *Journal of Abnormal Psychology*, *121*(2), 315–324.

https://doi.org/10.1037/a0025738

Danese, A., Moffitt, T. E., Harrington, H., Milne, B. J., Polanczyk, G., Pariante, C. M., … Caspi,

A. (2009). Adverse childhood experiences and adult risk factors for age-related disease:

Depression, inflammation, and clustering of metabolic risk markers. *Archives of*

Pediatrics & Adolescent Medicine, *163*(12), 1135–1143.

https://doi.org/10.1001/archpediatrics.2009.214

Danese, A., Pariante, C. M., Caspi, A., Taylor, A., & Poulton, R. (2007). Childhood maltreatment

predicts adult inflammation in a life-course study. *Proceedings of the National Academy*

of Sciences, *104*(4), 1319–1324. https://doi.org/10.1073/pnas.0610362104

Danese, A., & Widom, C. S. (2023). Associations Between Objective and Subjective Experiences

of Childhood Maltreatment and the Course of Emotional Disorders in Adulthood. *JAMA*

Psychiatry, e232140. https://doi.org/10.1001/jamapsychiatry.2023.2140

Diedenhofen, B., & Musch, J. (2015). Cocor: A comprehensive solution for the statistical

comparison of correlations. *PLoS ONE*, *10*(4), e0121945.

https://doi.org/10.1371/journal.pone.0121945

Dowle, M., & Srinivasan, A. (2020). *Data.table: Extension of 'data.frame'*. Retrieved from

https://CRAN.R-project.org/package=data.table

Duits, P., Cath, D. C., Lissek, S., Hox, J. J., Hamm, A. O., Engelhard, I. M., … Baas, J. M. P.

(2015). UPDATED META-ANALYSIS OF CLASSICAL FEAR CONDITIONING IN

THE ANXIETY DISORDERS: Review: Updated Meta-Analysis of Fear Conditioning in

Anxiety Disorders. *Depression and Anxiety*, *32*(4), 239–253.

https://doi.org/10.1002/da.22353

Dunn, O. J., & Clark, V. (1969). Correlation Coefficients Measured on the Same Individuals.

Journal of the American Statistical Association, *64*(325), 366–377.

https://doi.org/10.1080/01621459.1969.10500981

Dymond, S., Dunsmoor, J. E., Vervliet, B., Roche, B., & Hermans, D. (2015). Fear

Generalization in Humans: Systematic Review and Implications for Anxiety Disorder

Research. *Behavior Therapy*, *46*(5), 561–582. https://doi.org/10.1016/j.beth.2014.10.001

Ebbert, D. (2019). *Chisq.posthoc.test: A post hoc analysis for pearson's chi-squared test for

count data*. Retrieved from https://CRAN.R-project.org/package=chisq.posthoc.test

Ehlers, M. R., Nold, J., Kuhn, M., Klingelhöfer-Jens, M., & Lonsdorf, T. B. (2020). Revisiting

potential associations between brain morphology, fear acquisition and extinction through

new data and a literature review. *Scientific Reports*, *10*(1), 19894.

https://doi.org/10.1038/s41598-020-76683-1

Estrada, S., Richards, C., Gee, D. G., & Baskin-Sommers, A. (2020). Exposure to violence and

nonassociative learning capability confer risk for violent behavior. *Journal of Abnormal

Psychology*, *129*(7), 748–759. https://doi.org/10.1037/abn0000579

Evans, G. W., Li, D., & Whipple, S. S. (2013). Cumulative risk and child development.

Psychological Bulletin, *139*(6), 1342–1396. https://doi.org/10.1037/a0031808

FC, M., Davis, T. L., & ggplot2 authors. (2022). *Ggpattern: 'ggplot2' pattern geoms*. Retrieved

from https://CRAN.R-project.org/package=ggpattern

Felitti, V. J. (2002). The relationship of adverse childhood experiences to adult health: Turning

gold into lead/ Belastungen in der Kindheit und Gesundheit im Erwachsenenalter: die

Verwandlung von Gold in Blei. *Zeitschrift für Psychosomatische Medizin und

Psychotherapie*, *48*(4), 359–369. https://doi.org/10.13109/zptm.2002.48.4.359

Fox, J., & Weisberg, S. (2019). *An R companion to applied regression* (Third). Thousand Oaks

CA: Sage. Retrieved from https://socialsciences.mcmaster.ca/jfox/Books/Companion/

Fox, J., Weisberg, S., & Price, B. (2020). *carData: Companion to applied regression data sets*.

Retrieved from https://CRAN.R-project.org/package=carData

Fraunfelter, L., Gerdes, A. B. M., & Alpers, G. W. (2022). Fear one, fear them all: A systematic

review and meta-analysis of fear generalization in pathological anxiety. *Neuroscience and

Biobehavioral Reviews*, *139*, 104707. https://doi.org/10.1016/j.neubiorev.2022.104707

Fredrikson, M., Annas, P., Georgiades, A., Hursti, T., & Tersman, Z. (1993). Internal consistency

and temporal stability of classically conditioned skin conductance responses. *Biological

Psychology*, *35*(2), 153–163. https://doi.org/10.1016/0301-0511(93)90011-V

Galea, S., Nandi, A., & Vlahov, D. (2005). The Epidemiology of Post-Traumatic Stress Disorder

after Disasters. *Epidemiologic Reviews*, *27*(1), 78–91.

https://doi.org/10.1093/epirev/mxi003

Gilbert, R., Widom, C. S., Browne, K., Fergusson, D., Webb, E., & Janson, S. (2009). Burden

and consequences of child maltreatment in high-income countries. *The Lancet*, *373*(9657),

68–81. https://doi.org/10.1016/S0140-6736(08)61706-7

Gohel, D. (2021). *Flextable: Functions for tabular reporting*. Retrieved from https://CRAN.R-project.org/package=flextable

Gohel, D., & Ross, N. (2022). *Officedown: Enhanced 'r markdown' format for 'word' and 'PowerPoint'*. Retrieved from https://CRAN.R-project.org/package=officedown

Green, J. G., McLaughlin, K. A., Berglund, P. A., Gruber, M. J., Sampson, N. A., Zaslavsky, A. M., & Kessler, R. C. (2010). Childhood Adversities and Adult Psychiatric Disorders in the National Comorbidity Survey Replication I: Associations With First Onset of *DSM-IV* Disorders. *Archives of General Psychiatry*, *67*(2), 113. https://doi.org/10.1001/archgenpsychiatry.2009.186

Gromer, D. (2020). *Apa: Format outputs of statistical tests according to APA guidelines*. Retrieved from https://CRAN.R-project.org/package=apa

Hair, J. F., Anderson, R. E., Tatham, R. L., & Black, W. C. (1995). *Multivariate Data Analysis* (3rd ed.). New York: Macmillan.

Harnett, N. G., Wheelock, M. D., Wood, K. H., Goodman, A. M., Mrug, S., Elliott, M. N., … Knight, D. C. (2019). Negative life experiences contribute to racial differences in the neural response to threat. *NeuroImage*, *202*, 116086. https://doi.org/10.1016/j.neuroimage.2019.116086

Häuser, W., Schmutzer, G., & Glaesmer, H. (2011). Maltreatment in childhood and adolescence. *Deutsches Ärzteblatt International*, *108*(17), 287–294.

Hautzinger, M., & Bailer, M. (1993). *Allgemeine Depressions-Skala (ADS) [German version of the Center for Epidemiological Studies Depression Scale (CES-D)]*. Weinheim: Belz.

Hayes, A. F., & Cai, L. (2007). Using heteroskedasticity-consistent standard error estimators in

OLS regression: An introduction and software implementation. *Behavior Research

Methods*, *39*(4), 709–722. https://doi.org/10.3758/bf03192961

Heim, C., & Nemeroff, C. B. (2001). The role of childhood trauma in the neurobiology of mood

and anxiety disorders: Preclinical and clinical studies. *Biological Psychiatry*, *49*(12),

1023–1039. https://doi.org/10.1016/S0006-3223(01)01157-X

Heim, C., & Nemeroff, C. B. (2002). Neurobiology of early life stress: Clinical studies. *Seminars

in Clinical Neuropsychiatry*, *7*(2), 147–159. https://doi.org/10.1053/scnp.2002.33127

Henry, L., & Wickham, H. (2020). *Purrr: Functional programming tools*. Retrieved from

https://CRAN.R-project.org/package=purrr

Herzog, K., Andreatta, M., Schneider, K., Schiele, M. A., Domschke, K., Romanos, M., … Pauli,

P. (2021). Reducing Generalization of Conditioned Fear: Beneficial Impact of Fear

Relevance and Feedback in Discrimination Training. *Frontiers in Psychology*, *12*, 665711.

https://doi.org/10.3389/fpsyg.2021.665711

Ho, D. E., Imai, K., King, G., & Stuart, E. A. (2011). MatchIt: Nonparametric preprocessing for

parametric causal inference. *Journal of Statistical Software*, *42*(8), 1–28.

https://doi.org/10.18637/jss.v042.i08

Hölzel, B. K., Brunsch, V., Gard, T., Greve, D. N., Koch, K., Sorg, C., … Milad, M. R. (2016).

Mindfulness-Based Stress Reduction, Fear Conditioning, and The Uncinate Fasciculus: A

Pilot Study. *Frontiers in Behavioral Neuroscience*, *10*.

https://doi.org/10.3389/fnbeh.2016.00124

Hosseini-Kamkar, N., Varvani Farahani, M., Nikolic, M., Stewart, K., Goldsmith, S.,

Soltaninejad, M., … Leyton, M. (2023). Adverse Life Experiences and Brain Function: A

Meta-Analysis of Functional Magnetic Resonance Imaging Findings. *JAMA Network*

Open, *6*(11), e2340018. https://doi.org/10.1001/jamanetworkopen.2023.40018

Hughes, K., Ford, K., Bellis, M. A., Glendinning, F., Harrison, E., & Passmore, J. (2021). Health

and financial costs of adverse childhood experiences in 28 European countries: A

systematic review and meta-analysis. *The Lancet Public Health*, *6*(11), e848–e857.

https://doi.org/10.1016/S2468-2667(21)00232-2

Huskey, A., Taylor, D. J., & Friedman, B. H. (2022). "Generalized unsafety" as fear inhibition to

safety signals in adults with and without childhood trauma. *Developmental Psychobiology*,

64(4). https://doi.org/10.1002/dev.22242

Imholze, C., Hutterer, K., Gall, D., Dannlowski, U., Domschke, K., Leehr, E. J., … Gamer, M.

(2023). Prediction of Changes in Negative Affect During the COVID-19 Pandemic by

Experimental Fear Conditioning and Generalization Measures: A Longitudinal Study.

Zeitschrift für Psychologie, *231*(2), 137–148. https://doi.org/10.1027/2151-2604/a000523

Infantolino, Z. P., Luking, K. R., Sauder, C. L., Curtin, J. J., & Hajcak, G. (2018). Robust is not

necessarily reliable: From within-subjects fMRI contrasts to between-subjects

comparisons. *NeuroImage*, *173*, 146–152.

https://doi.org/10.1016/j.neuroimage.2018.02.024

Jovanovic, T., Blanding, N. Q., Norrholm, S. D., Duncan, E., Bradley, B., & Ressler, K. J.

(2009). Childhood abuse is associated with increased startle reactivity in adulthood.

Depression and Anxiety, *26*(11), 1018–1026. https://doi.org/10.1002/da.20599

Jovanovic, T., Wiltshire, C. N., Reda, M. H., France, J., Wanna, C. P., Minton, S. T., … Stenson, A. F. (2022). Uncertain in the face of change: Lack of contingency shift awareness during extinction is associated with higher fear-potentiated startle and PTSD symptoms in children. *International Journal of Psychophysiology*, *178*, 90–98. https://doi.org/10.1016/j.ijpsycho.2022.06.008

Kaczkurkin, A. N., Burton, P. C., Chazin, S. M., Manbeck, A. B., Espensen-Sturges, T., Cooper, S. E., … Lissek, S. (2017). Neural Substrates of Overgeneralized Conditioned Fear in PTSD. *American Journal of Psychiatry*, *174*(2), 125–134. https://doi.org/10.1176/appi.ajp.2016.15121549

Karstoft, K.-I., & Armour, C. (2023). What we talk about when we talk about trauma: Content overlap and heterogeneity in the assessment of trauma exposure. *Journal of Traumatic Stress*, *36*(1), 71–82. https://doi.org/10.1002/jts.22880

Kassambara, A. (2020). *Ggpubr: 'ggplot2' based publication ready plots*. Retrieved from https://CRAN.R-project.org/package=ggpubr

Kassambara, A. (2021). *Rstatix: Pipe-friendly framework for basic statistical tests*. Retrieved from https://CRAN.R-project.org/package=rstatix

Klauke, B., Deckert, J., Reif, A., Pauli, P., & Domschke, K. (2010). Life events in panic disorder-an update on "candidate stressors". *Depression and Anxiety*, *27*(8), 716–730. https://doi.org/10.1002/da.20667

Klingelhöfer-Jens, M., Ehlers, M. R., Kuhn, M., Keyaniyan, V., & Lonsdorf, T. B. (2022). Robust group- but limited individual-level (longitudinal) reliability and insights into cross-

phases response prediction of conditioned fear. *eLife*, *11*, e78717.

https://doi.org/10.7554/eLife.78717

Koppold, A., Kastrinogiannis, A., Kuhn, M., & Lonsdorf, T. B. (2023). Watching with Argus

eyes: Characterization of emotional and physiological responding in adults exposed to

childhood maltreatment and/or recent adversity. *Psychophysiology*, *60*(7), e14253.

https://doi.org/10.1111/psyp.14253

Kreutzer, K. A., & Gorka, S. M. (2021). Impact of Trauma Type on Startle Reactivity to

Predictable and Unpredictable Threats. *Journal of Nervous & Mental Disease*, *209*(12),

899–904. https://doi.org/10.1097/NMD.0000000000001394

Kuhlman, K. R., Chiang, J. J., Horn, S., & Bower, J. E. (2017). Developmental

psychoneuroendocrine and psychoneuroimmune pathways from childhood adversity to

disease. *Neuroscience and Biobehavioral Reviews*, *80*, 166–184.

https://doi.org/10.1016/j.neubiorev.2017.05.020

Kuhn, M., Mertens, G., & Lonsdorf, T. B. (2016). State anxiety modulates the return of fear.

International Journal of Psychophysiology, *110*, 194–199.

https://doi.org/10.1016/j.ijpsycho.2016.08.001

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest package: Tests in

linear mixed effects models. *Journal of Statistical Software*, *82*(13), 1–26.

https://doi.org/10.18637/jss.v082.i13

Lau, J. Y. F., Lissek, S., Nelson, E. E., Lee, Y., Roberson-Nay, R., Poeth, K., … Pine, D. S.

(2008). Fear conditioning in adolescents with anxiety disorders: Results from a novel

experimental paradigm. *Journal of the American Academy of Child and Adolescent Psychiatry*, *47*(1), 94–102. https://doi.org/10.1097/chi.0b01e31815a5f01

Laux, L., & Spielberger, C. D. (1981). *Das State-Trait-Angstinventar: STAI*. Weinheim: Beltz.

Lawrence, M. A. (2016). *Ez: Easy analysis and visualization of factorial experiments*. Retrieved from https://CRAN.R-project.org/package=ez

LeBel, E. P., McCarthy, R. J., Earp, B. D., Elson, M., & Vanpaemel, W. (2018). A Unified Framework to Quantify the Credibility of Scientific Findings. *Advances in Methods and Practices in Psychological Science*, *1*(3), 389–402. https://doi.org/10.1177/2515245918787489

Lis, S., Thome, J., Kleindienst, N., Mueller-Engelmann, M., Steil, R., Priebe, K., … Bohus, M. (2020). Generalization of fear in post-traumatic stress disorder. *Psychophysiology*, *57*(1), e13422. https://doi.org/10.1111/psyp.13422

Lissek, S., Powers, A. S., McClure, E. B., Phelps, E. A., Woldehawariat, G., Grillon, C., & Pine, D. S. (2005). Classical fear conditioning in the anxiety disorders: A meta-analysis. *Behaviour Research and Therapy*, *43*(11), 1391–1424. https://doi.org/10.1016/j.brat.2004.10.007

Litz, B. T., & Gray, M. J. (2002). Emotional numbing in posttraumatic stress disorder: Current and future research directions. *The Australian and New Zealand Journal of Psychiatry*, *36*(2), 198–204. https://doi.org/10.1046/j.1440-1614.2002.01002.x

Lonsdorf, T. B., Klingelhöfer-Jens, M., Andreatta, M., Beckers, T., Chalkia, A., Gerlicher, A., …

Merz, C. J. (2019). Navigating the garden of forking paths for data exclusions in fear

conditioning research. *eLife*, *8*, e52465. https://doi.org/10.7554/eLife.52465

Lonsdorf, T. B., Menz, M. M., Andreatta, M., Fullana, M. A., Golkar, A., Haaker, J., … Merz, C.

J. (2017). Don't fear 'fear conditioning': Methodological considerations for the design and

analysis of studies on human fear acquisition, extinction, and return of fear. *Neuroscience

and Biobehavioral Reviews*, *77*, 247–285. https://doi.org/10.1016/j.neubiorev.2017.02.026

Lüdecke, D. (n.d.). *sjPlot: Data visualization for statistics in social science*. Retrieved from

https://CRAN.R-project.org/package=sjPlot

Lüdecke, D., Ben-Shachar, M. S., Patil, I., Waggoner, P., & Makowski, D. (2021). performance:

An R package for assessment, comparison and testing of statistical models. *Journal of

Open Source Software*, *6*(60), 3139. https://doi.org/10.21105/joss.03139

Lykken, D. T. (1972). Range correction applied to heart rate and to GSR data. *Psychophysiology*,

9(3), 373–379. https://doi.org/10.1111/j.1469-8986.1972.tb03222.x

Lykken, D. T., & Venables, P. H. (1971). Direct measurement of skin conductance: A proposal

for standardization. *Psychophysiology*, *8*(5), 656–672. https://doi.org/10.1111/j.1469-

8986.1971.tb00501.x

Lynam, D. R., Hoyle, R. H., & Newman, J. P. (2006). The perils of partialling: Cautionary tales

from aggression and psychopathy. *Assessment*, *13*(3), 328–341.

https://doi.org/10.1177/1073191106290562

Machlin, L., Miller, A. B., Snyder, J., McLaughlin, K. A., & Sheridan, M. A. (2019). Differential

Associations of Deprivation and Threat With Cognitive Control and Fear Conditioning in

Early Childhood. *Frontiers in Behavioral Neuroscience*, *13*, 80.

https://doi.org/10.3389/fnbeh.2019.00080

Mair, P., & Wilcox, R. (2020). Robust statistical methods in R using the WRS2 package.

Behavior Research Methods, *52*(2), 464–488. https://doi.org/10.3758/s13428-019-01246-

w

Mason, R. L., Gunst, R. F., & Hess, J. L. (1989). *Statistical Design and Analysis of Experiments:

Applications to Engineering and Science*. New York: Wiley.

McEwen, B. S. (2003). Mood disorders and allostatic load. *Biological Psychiatry*, *54*(3), 200–

207. https://doi.org/10.1016/S0006-3223(03)00177-X

McLaughlin, K. A., DeCross, S. N., Jovanovic, T., & Tottenham, N. (2019). Mechanisms linking

childhood adversity with psychopathology: Learning as an intervention target. *Behaviour

Research and Therapy*, *118*, 101–109. https://doi.org/10.1016/j.brat.2019.04.008

McLaughlin, K. A., Greif Green, J., Gruber, M. J., Sampson, N. A., Zaslavsky, A. M., & Kessler,

R. C. (2012). Childhood Adversities and First Onset of Psychiatric Disorders in a National

Sample of US Adolescents. *Archives of General Psychiatry*, *69*(11), 1151.

https://doi.org/10.1001/archgenpsychiatry.2011.2277

McLaughlin, K. A., Koenen, K. C., Hill, E. D., Petukhova, M., Sampson, N. A., Zaslavsky, A.

M., & Kessler, R. C. (2013). Trauma exposure and posttraumatic stress disorder in a

national sample of adolescents. *Journal of the American Academy of Child and Adolescent

Psychiatry*, *52*(8), 815–830.e14. https://doi.org/10.1016/j.jaac.2013.05.011

McLaughlin, K. A., & Sheridan, M. A. (2016). Beyond Cumulative Risk: A Dimensional

Approach to Childhood Adversity. *Current Directions in Psychological Science*, *25*(4),

239–245. https://doi.org/10.1177/0963721416655883

McLaughlin, K. A., Sheridan, M. A., Humphreys, K. L., Belsky, J., & Ellis, B. J. (2021). The

Value of Dimensional Models of Early Experience: Thinking Clearly About Concepts and

Categories. *Perspectives on Psychological Science*, *16*(6), 1463–1472.

https://doi.org/10.1177/1745691621992346

McLaughlin, K. A., Sheridan, M. A., & Lambert, H. K. (2014). Childhood adversity and neural

development: Deprivation and threat as distinct dimensions of early experience.

Neuroscience & Biobehavioral Reviews, *47*, 578–591.

https://doi.org/10.1016/j.neubiorev.2014.10.012

McLaughlin, K. A., & Sheridan, M. A. (2016). Beyond Cumulative Risk: A Dimensional

Approach to Childhood Adversity. *Current Directions in Psychological Science*, *25*(4),

239–245. https://doi.org/10.1177/0963721416655883

McLaughlin, K. A., Sheridan, M. A., Gold, A. L., Duys, A., Lambert, H. K., Peverill, M.,

Heleniak, C., Shechner, T., Wojcieszak, Z., & Pine, D. S. (2016). Maltreatment Exposure,

Brain Structure, and Fear Conditioning in Children and Adolescents.

Neuropsychopharmacology: Official Publication of the American College of

Neuropsychopharmacology, *41*(8), 1956–1964. https://doi.org/10.1038/npp.2015.365

McLaughlin, K. A., Weissman, D., & Bitrán, D. (2019). Childhood Adversity and Neural

Development: A Systematic Review. *Annual Review of Developmental Psychology*, *1*,

277–312. https://doi.org/10.1146/annurev-devpsych-121318-084950

McMahon, S. D., Grant, K. E., Compas, B. E., Thurm, A. E., & Ey, S. (2003). Stress and
psychopathology in children and adolescents: Is there evidence of specificity?: Specificity
and stress. *Journal of Child Psychology and Psychiatry*, *44*(1), 107–133.
https://doi.org/10.1111/1469-7610.00105

Moffitt, T. E., Caspi, A., Harrington, H., Milne, B. J., Melchior, M., Goldberg, D., & Poulton, R.
(2007). Generalized anxiety disorder and depression: Childhood risk factors in a birth
cohort followed to age 32. *Psychological Medicine*, *37*(03), 441.
https://doi.org/10.1017/S0033291706009640

Moriarity, D. P., & Alloy, L. B. (2021). Back to Basics: The Importance of Measurement
Properties in Biological Psychiatry. *Neuroscience & Biobehavioral Reviews*, *123*, 72–82.
https://doi.org/10.1016/j.neubiorev.2021.01.008

Müller, K. (2020). *Here: A simpler way to find your files*. Retrieved from https://CRAN.R-
project.org/package=here

Müller, K., & Wickham, H. (2021). *Tibble: Simple data frames*. Retrieved from https://CRAN.R-
project.org/package=tibble

Neter, J., Wasserman, W., & Kutner, M. H. (1989). *Applied Linear Regression Models*.
Homewood, IL: Irwin.

O'brien, R. M. (2007). A Caution Regarding Rules of Thumb for Variance Inflation Factors.
Quality & Quantity, *41*(5), 673–690. https://doi.org/10.1007/s11135-006-9018-6

Pedersen, T. L. (2020). *Patchwork: The composer of plots*. Retrieved from https://CRAN.R-
project.org/package=patchwork

Pole, N., Neylan, T. C., Otte, C., Metzler, T. J., Best, S. R., Henn-Haase, C., & Marmar, C. R. (2007). Associations between childhood trauma and emotion-modulated psychophysiological responses to startling sounds: A study of police cadets. *Journal of Abnormal Psychology*, *116*(2), 352–361. https://doi.org/10.1037/0021-843X.116.2.352

Pollak, S. D., Cicchetti, D., Hornung, K., & Reed, A. (2000). Recognizing emotion in faces: Developmental effects of child abuse and neglect. *Developmental Psychology*, *36*(5), 679–688. https://doi.org/10.1037/0012-1649.36.5.679

Pollak, S. D., & Tolley-Schell, S. (2004). Attention, Emotion, and the Development of Psychopathology. In M. I. Posner (Hrsg.), *Cognitive neuroscience of attention* (S. 357–368). The Guilford Press.

Pollak, S. D., & Smith, K. E. (2021). Thinking Clearly About Biology and Childhood Adversity: Next Steps for Continued Progress. *Perspectives on Psychological Science*, *16*(6), 1473–1477. https://doi.org/10.1177/17456916211031539

Pollak, S. D., Vardi, S., Putzer Bechner, A. M., & Curtin, J. J. (2005). Physically abused children's regulation of attention in response to hostility. *Child Development*, *76*(5), 968–977. https://doi.org/10.1111/j.1467-8624.2005.00890.x

R Core Team. (2022a). *Foreign: Read data stored by 'minitab', 's', 'SAS', 'SPSS', 'stata', 'systat', 'weka', 'dBase', ...* Retrieved from https://CRAN.R-project.org/package=foreign

R Core Team. (2022b). *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing. Retrieved from https://www.R-project.org/

Revelle, W. (2020). *Psych: Procedures for psychological, psychometric, and personality research*. Evanston, Illinois: Northwestern University. Retrieved from https://CRAN.R-project.org/package=psych

Richter-Levin, G., & Sandi, C. (2021). Title: "Labels Matter: Is it stress or is it Trauma?". *Translational Psychiatry*, *11*(1), 385. https://doi.org/10.1038/s41398-021-01514-4

Rowland, G. E., Mekawi, Y., Michopoulos, V., Powers, A., Fani, N., Bradley, B., … Stevens, J. S. (2022). Distinctive impacts of sexual trauma versus non-sexual trauma on PTSD profiles in highly trauma-exposed, Black women. *Journal of Affective Disorders*, *317*, 329–338. https://doi.org/10.1016/j.jad.2022.08.099

Ruge, J., Ehlers, M. R., Kastrinogiannis, A., Klingelhöfer-Jens, M., Koppold, A., Abend, R., & Lonsdorf, T. B. (2024). How adverse childhood experiences get under the skin: A systematic review, integration and methodological discussion on threat and reward learning mechanisms. *eLife*, *13*, e92700. https://doi.org/10.7554/eLife.92700

Russo, S. J., Murrough, J. W., Han, M.-H., Charney, D. S., & Nestler, E. J. (2012). Neurobiology of resilience. *Nature Neuroscience*, *15*(11), 1475–1484. https://doi.org/10.1038/nn.3234

Samaey, C., Lecei, A., Achterhof, R., Hagemann, N., Hermans, K. S. F. M., Hiekkaranta, A. P., … Van Winkel, R. (2024). Anticipating future threats: Evidence for association, but not interaction, of childhood adversity and identity development with threat anticipation in adolescence. *Early Intervention in Psychiatry*, eip.13511. https://doi.org/10.1111/eip.13511

Scharfenort, R., Menz, M., & Lonsdorf, T. B. (2016). Adversity-induced relapse of fear: Neural mechanisms and implications for relapse prevention from a study on experimentally

induced return-of-fear following fear conditioning and extinction. *Translational Psychiatry*, *6*, e858. https://doi.org/10.1038/tp.2016.126

Schiele, M. A., Herzog, K., Kollert, L., Schartner, C., Leehr, E. J., Böhnlein, J., … Domschke, K. (2020). Extending the vulnerability–stress model of mental disorders: Three-dimensional *NPSR1* environment coping interaction study in anxiety. *The British Journal of Psychiatry*, *217*(5), 645–650. https://doi.org/10.1192/bjp.2020.73

Schiele, M. A., Reinhard, J., Reif, A., Domschke, K., Romanos, M., Deckert, J., & Pauli, P. (2016). Developmental aspects of fear: Comparing the acquisition and generalization of conditioned fear in children and adults. *Developmental Psychobiology*, *58*(4), 471–481. https://doi.org/10.1002/dev.21393

Schiele, M. A., Ziegler, C., Holitschke, K., Schartner, C., Schmidt, B., Weber, H., … Domschke, K. (2016). Influence of 5-HTT variation, childhood trauma and self-efficacy on anxiety traits: A gene-environment-coping interaction study. *Journal of Neural Transmission (Vienna, Austria: 1996)*, *123*(8), 895–904. https://doi.org/10.1007/s00702-016-1564-z

Schloerke, B., Cook, D., Larmarange, J., Briatte, F., Marbach, M., Thoen, E., … Crowley, J. (2021). *GGally: Extension to 'ggplot2'*. Retrieved from https://CRAN.R-project.org/package=GGally

Sheehan, D. V., Lecrubier, Y., Sheehan, K. H., Amorim, P., Janavs, J., Weiller, E., … Dunbar, G. C. (1998). The Mini-International Neuropsychiatric Interview (M.I.N.I.): The development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *The Journal of Clinical Psychiatry*, *59 Suppl 20*, 22-33;quiz 34-57.

Sheridan, M. A., & McLaughlin, K. A. (2014). Dimensions of early experience and neural development: Deprivation and threat. *Trends in Cognitive Sciences*, *18*(11), 580–585. https://doi.org/10.1016/j.tics.2014.09.001

Sheridan, M. A., & McLaughlin, K. A. (2016). Neurobiological models of the impact of adversity on education. *Current Opinion in Behavioral Sciences*, *10*, 108–113. https://doi.org/10.1016/j.cobeha.2016.05.013

Sheridan, M. A., Peverill, M., Finn, A. S., & McLaughlin, K. A. (2017). Dimensions of childhood adversity have distinct associations with neural systems underlying executive functioning. *Development and Psychopathology*, *29*(5), 1777–1794. https://doi.org/10.1017/S0954579417001390

Silvers, J. A., Lumian, D. S., Gabard-Durnam, L., Gee, D. G., Goff, B., Fareri, D. S., … Tottenham, N. (2016). Previous Institutionalization Is Followed by Broader Amygdala-Hippocampal-PFC Network Connectivity during Aversive Learning in Human Development. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *36*(24), 6420–6430. https://doi.org/10.1523/JNEUROSCI.0038-16.2016

Sjoberg, D. (2023). *Ggsankey: Sankey, alluvial and sankey bump plots*.

Sjouwerman, R., Illius, S., Kuhn, M., & Lonsdorf, T. B. (2022). A data multiverse analysis investigating non-model based SCR quantification approaches. *Psychophysiology*. https://doi.org/10.1111/psyp.14130

Sjouwerman, R., & Lonsdorf, T. B. (2020). Experimental boundary conditions of reinstatement-induced return of fear in humans: Is reinstatement in humans what we think it is? *Psychophysiology*, *57*(5). https://doi.org/10.1111/psyp.13549

Sjouwerman, R., Niehaus, J., & Lonsdorf, T. B. (2015). Contextual Change After Fear Acquisition Affects Conditioned Responding and the Time Course of Extinction Learning—Implications for Renewal Research. *Frontiers in Behavioral Neuroscience*, *9*. https://doi.org/10.3389/fnbeh.2015.00337

Smith, K. E., & Pollak, S. D. (2021). Rethinking Concepts and Categories for Understanding the Neurodevelopmental Effects of Childhood Adversity. *Perspectives on Psychological Science*, *16*(1), 67–93. https://doi.org/10.1177/1745691620920725

Spielberger, C. D. (1983). *Manual for the State-Trait Inventory STAI (Form Y)*. Palo Alto, CA: Mind Garden.

Spinhoven, P., Elzinga, B. M., Hovens, J. G. F. M., Roelofs, K., van Oppen, P., Zitman, F. G., & Penninx, B. W. J. H. (2011). Positive and negative life events and personality traits in predicting course of depression and anxiety. *Acta Psychiatrica Scandinavica*, *124*(6), 462–473. https://doi.org/10.1111/j.1600-0447.2011.01753.x

Stegmann, Y., Schiele, M. A., Schümann, D., Lonsdorf, T. B., Zwanzger, P., Romanos, M., … Pauli, P. (2019). Individual differences in human fear generalization—pattern identification and implications for anxiety disorders. *Translational Psychiatry*, *9*(1), 307. https://doi.org/10.1038/s41398-019-0646-8

Teicher, M. H., Gordon, J. B., & Nemeroff, C. B. (2022). Recognizing the importance of childhood maltreatment as a critical factor in psychiatric diagnoses, treatment, research, prevention, and education. *Molecular Psychiatry*, *27*(3), 1331–1338. https://doi.org/10.1038/s41380-021-01367-9

Thome, J., Hauschild, S., Koppe, G., Liebke, L., Rausch, S., Herzog, J. I., … Lis, S. (2018). Generalisation of fear in PTSD related to prolonged childhood maltreatment: An experimental study. *Psychological Medicine*, *48*(13), 2223–2234. https://doi.org/10.1017/S0033291717003713

Tiedemann, F. (2020). *Gghalves: Compose half-half plots using your favourite geoms*. Retrieved from https://CRAN.R-project.org/package=gghalves

Torchiano, M. (2020). *Effsize: Efficient effect size computation*. https://doi.org/10.5281/zenodo.1480624

Vervliet, B., Craske, M. G., & Hermans, D. (2013). Fear extinction and relapse: State of the art. *Annual Review of Clinical Psychology*, *9*, 215–248. https://doi.org/10.1146/annurev-clinpsy-050212-185542

Wickham, H. (2007). Reshaping data with the reshape package. *Journal of Statistical Software*, *21*(12), 1–20. Retrieved from http://www.jstatsoft.org/v21/i12/

Wickham, H. (2016). *ggplot2: Elegant graphics for data analysis*. Springer-Verlag New York. Retrieved from https://ggplot2.tidyverse.org

Wickham, H. (2019). *Stringr: Simple, consistent wrappers for common string operations*. Retrieved from https://CRAN.R-project.org/package=stringr

Wickham, H. (2020). *Forcats: Tools for working with categorical variables (factors)*. Retrieved from https://CRAN.R-project.org/package=forcats

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L. D., François, R., … Yutani, H. (2019). Welcome to the tidyverse. *Journal of Open Source Software*, *4*(43), 1686. https://doi.org/10.21105/joss.01686

Wickham, H., François, R., Henry, L., & Müller, K. (2022). *Dplyr: A grammar of data manipulation*. Retrieved from https://CRAN.R-project.org/package=dplyr

Wickham, H., & Girlich, M. (2022). *Tidyr: Tidy messy data*. Retrieved from https://CRAN.R-project.org/package=tidyr

Wickham, H., & Hester, J. (2020). *Readr: Read rectangular text data*. Retrieved from https://CRAN.R-project.org/package=readr

Wickham, H., & Miller, E. (2020). *Haven: Import and export 'SPSS', 'stata' and 'SAS' files*. Retrieved from https://CRAN.R-project.org/package=haven

Wingenfeld, K., Spitzer, C., Mensebach, C., Grabe, H., Hill, A., Gast, U., … Driessen, M. (2010). Die deutsche Version des Childhood Trauma Questionnaire (CTQ): Erste Befunde zu den psychometrischen Kennwerten. *PPmP - Psychotherapie · Psychosomatik · Medizinische Psychologie*, *60*(11), 442–450. https://doi.org/10.1055/s-0030-1247564

Xie, Y. (2015). *Dynamic documents with R and knitr* (2nd ed.). Boca Raton, Florida: Chapman; Hall/CRC. Retrieved from https://yihui.org/knitr/

Yasumoto, A. (2023). *ftExtra: Extensions for 'flextable'*. Retrieved from https://CRAN.R-project.org/package=ftExtra

Young, D. A., Neylan, T. C., O'Donovan, A., Metzler, T., Richards, A., Ross, J. A., & Inslicht, S. S. (2018). The interaction of BDNF Val66Met, PTSD, and child abuse on

psychophysiological reactivity and HPA axis function in a sample of Gulf War Veterans. *Journal of Affective Disorders*, *235*, 52–60. https://doi.org/10.1016/j.jad.2018.04.004

Young, E. S., Farrell, A. K., Carlson, E. A., Englund, M. M., Miller, G. E., Gunnar, M. R., … Simpson, J. A. (2019). The Dual Impact of Early and Concurrent Life Stress on Adults' Diurnal Cortisol Patterns: A Prospective Study. *Psychological Science*, *30*(5), 739–747. https://doi.org/10.1177/0956797619833664

Zeileis, A. (2004). Econometric computing with HC and HAC covariance matrix estimators. *Journal of Statistical Software*, *11*(10), 1–17. https://doi.org/10.18637/jss.v011.i10

Zeileis, A. (2006). Object-oriented computation of sandwich estimators. *Journal of Statistical Software*, *16*(9), 1–16. https://doi.org/10.18637/jss.v016.i09

Zeileis, A., & Grothendieck, G. (2005). Zoo: S3 infrastructure for regular and irregular time series. *Journal of Statistical Software*, *14*(6), 1–27. https://doi.org/10.18637/jss.v014.i06

Zeileis, A., & Hothorn, T. (2002). Diagnostic checking in regression relationships. *R News*, *2*(3), 7–10. Retrieved from https://CRAN.R-project.org/doc/Rnews/

Zeileis, A., Köll, S., & Graham, N. (2020). Various versatile variances: An object-oriented implementation of clustered covariances in R. *Journal of Statistical Software*, *95*(1), 1–36. https://doi.org/10.18637/jss.v095.i01

Zhu, H. (2020). *kableExtra: Construct complex table with 'kable' and pipe syntax*. Retrieved from https://CRAN.R-project.org/package=kableExtra

Zuo, X.-N., Xu, T., & Milham, M. P. (2019). Harnessing reliability for neuroscience research. *Nature Human Behaviour*, *3*(8), 768–771. https://doi.org/10.1038/s41562-019-0655-x