Prolonged delays in human microbiota transmission after a controlled antibiotic perturbation

Katherine S. Xue[1,2]†, Sophie Jean Walton[1,3]*, Doran A. Goldman[1]*, Maike L. Morrison[1], Adrian J. Verster[4], Autumn B. Parrott[5], Feiqiao Brian Yu[6], Norma F. Neff[6], Noah A. Rosenberg[1], Benjamin D. Ross[4], Dmitri A. Petrov[1,6], Kerwyn Casey Huang[5,6], Benjamin H. Good[1,6,7]†, David A. Relman[2,8,9]†

[1]Department of Biology, Stanford University, Stanford, CA 94305, USA
[2]Department of Microbiology and Immunology, Stanford University School of Medicine, Stanford, CA 94305, USA
[3]Biophysics Training Program, Stanford, CA 94305, USA
[4]Department of Microbiology and Immunology, Geisel School of Medicine at Dartmouth College, Hanover, NH 03755, USA
[5]Department of Bioengineering, Stanford, CA 94305, USA
[6]Chan Zuckerberg Biohub, San Francisco, CA 94158, USA
[7]Department of Applied Physics, Stanford, CA 94305, USA
[8]Department of Medicine, Stanford University School of Medicine, Stanford, CA 94305, USA
[9]Infectious Diseases Section, Veterans Affairs Palo Alto Health Care System, Palo Alto, CA 94304, USA

* These authors contributed equally.
† These authors contributed equally.

Correspondence: kxue@stanford.edu (K.S.X.), bhgood@stanford.edu (B.H.G.), relman@stanford.edu (D.A.R.)

Abstract

Humans constantly encounter new microbes, but few become long-term residents of the adult gut microbiome. Classical theories predict that colonization is determined by the availability of open niches, but it remains unclear whether other ecological barriers limit commensal colonization in natural settings. To disentangle these effects, we used a controlled perturbation with the antibiotic ciprofloxacin to investigate the dynamics of gut microbiome transmission in 22 households of healthy, cohabiting adults. Colonization was rare in three-quarters of antibiotic-taking subjects, whose resident strains rapidly recovered in the week after antibiotics ended. In contrast, the remaining subjects exhibited lasting responses to antibiotics, with extensive species losses and transient expansions of potential opportunistic pathogens. These subjects experienced elevated rates of commensal colonization, but only after long delays: many new colonizers underwent sudden, correlated expansions months after the antibiotic perturbation. Furthermore, strains that had previously transmitted between cohabiting partners rarely recolonized after antibiotic disruptions, showing that colonization displays substantial historical contingency. This work demonstrates that there remain substantial ecological barriers to colonization even after major microbiome disruptions, suggesting that dispersal interactions and priority effects limit the pace of community change.

Introduction

New microbes constantly enter the digestive tract from our diets and environments, but few of these microbes become long-term residents of the adult gut microbiome[1–3]. To colonize the gut, microbes must overcome a series of ecological barriers: they must disperse into a new host, survive the harsh environment of the stomach[4,5], and compete against resident microbes to occupy an ecological niche[3,6–11]. Understanding these ecological barriers is crucial for understanding how gut bacteria spread between hosts[12–14] and for the design and maintenance of microbiome therapeutics[3,11,15–17].

Despite these substantial ecological barriers to colonization, people who live together frequently carry pairs of nearly identical microbial strains[12–14,18,19], suggesting that commensal microbes can transmit between cohabiting partners and colonize established communities. However, few studies have conducted the longitudinal sampling necessary to capture these transmission events in action[13,20], and it remains unclear what ecological contexts allow microbes to successfully colonize new hosts.

One prominent view is that the colonization of the gut is primarily determined by the availability of open niches[21–25]. In mouse models, for example, the presence of a single resident strain can prevent other, closely related species from colonizing, suggesting that niche availability could play a major role in the colonization resistance of the gut microbiota[26–31]. This simple view of colonization predicts that any perturbation that opens niches in an established community should lead to rapid colonization and extensive community change. But in natural human settings, it remains unclear how ecological factors like dispersal limitation[32] and priority effects[33] constrain the introduction and initial expansion of new microbes, especially because their effects are difficult to distinguish from a lack of available niches. Disentangling these ecological barriers requires controlled perturbations that open niches in the human gut, along with longitudinal sampling to link these open niches to subsequent colonization dynamics.

To address this gap, we tracked the dynamics of colonization in a household cohort before and after a controlled antibiotic perturbation. Antibiotics are a common perturbation that can disrupt the colonization resistance of established gut communities[23,34–36]. While this property is often used to promote the engraftment of deliberately introduced strains[37,38], previous work has also shown that gut microbiota can be remarkably resilient to antibiotic perturbations over longer timescales, recovering much of their initial species-level composition in the weeks following treatment[35,39–41]. It remains unclear to what extent this resilience is driven by the recovery of resident strains[41] or the colonization of new strains, and whether strains shared with cohabiting partners can serve as a reservoir to promote microbiome recovery[42]. We sought to address these questions by using strain-resolved metagenomics to profile the microbiomes of antibiotic-taking subjects and their cohabiting partners at high temporal and genetic resolution. These data allow us to observe the colonization of niches opened by antibiotics, providing a unique window into the ecological forces governing colonization in native human gut communities.

Results

Heterogeneous responses to antibiotic perturbation in a household cohort

To measure the dynamics of colonization in a naturalistic human setting, we performed a longitudinal study of 48 healthy adults from 22 households (**Fig. 1a** and **Extended Data Fig. 1**). Subjects collected stool samples at a mixture of weekly and daily intervals for two months, and in the middle of the study, one subject in each household took the broad-spectrum antibiotic ciprofloxacin orally for five days as a controlled perturbation to open niches in the established,

94   adult gut microbiome[23,34,35]. After this initial sampling period, subjects also collected follow-up
95   samples every few months for up to two years after the antibiotic perturbation, allowing us to
96   examine colonization dynamics over longer timescales. We performed metagenomic
97   sequencing of these stool samples and used a reference-based pipeline to profile species
98   abundance and strain-level variation[41,43,44].
99
Consistent with previous work[12–14,19], we found that many pairs of cohabiting, unrelated adults
shared gut microbial strains at the beginning of the study, indicating that they were acquired
through previous horizontal transmission (**Fig. 1b**). To identify shared strains, we first identified
species shared between cohabiting partners, and we classified the strains of that species as
"shared" if they had higher genetic similarity than 99% of strain pairs of the same species from
unrelated subjects (**Extended Data Fig. 2a-e; Methods**; median threshold of >99.9% average
nucleotide identity across species, or >75% of private marker SNPs detected). The total amount
of strain sharing varied widely across households: in 7 of 30 (23%) cohabiting pairs, we
detected no shared strains, whereas in the remaining subjects, we identified a median of four
shared strains that comprised a median of 23% of the relative abundance of the gut microbiome
(**Fig. 1b** and **Extended Data Fig. 2f,g**). The total abundance of shared strains was not strongly
correlated with the length of cohabitation (**Fig. 1c**; Pearson p=0.68). Two subject pairs shared
>30% of their gut microbiomes after living together for less than one year, but we detected no
shared strains in two households who had lived together for 6 and 30+ years, respectively (**Fig.**
1c). These data show that shared strains do not accumulate linearly with time, and that even
prolonged cohabitation is not sufficient to fully homogenize human gut microbiomes. We
hypothesized that these varied amounts of strain sharing could provide a backdrop for
investigating the responses to the antibiotic perturbation.
118
We first characterized the community-level effects of ciprofloxacin on gut microbiome
composition. Consistent with previous work[45,46], we found that ciprofloxacin perturbation caused
rapid changes in community composition that exceeded the magnitude of day-to-day variability
in the control subjects, who did not take antibiotics (**Fig. 1d,e, Extended Data Fig. 3a**).
However, the magnitude and duration of the responses varied widely across subjects, as
measured by the Jensen-Shannon divergence (JSD) of their species-level composition relative
to the initial timepoint. Most antibiotic-taking subjects (n=17/22, 77%) had only transient
antibiotic responses and declines in community diversity (**Fig. 1d** and **Extended Data Fig. 3b;**
Methods), in line with previous work showing that gut microbiomes often display resilience after
antibiotic perturbation[35,39,40]. In these subjects, community composition shifted during the
antibiotic perturbation but recovered quickly after the perturbation ended, such that, one month
after antibiotics, the gut microbiome had a JSD<0.4 from the pre-antibiotic community, a range
comparable to the control subjects.
132
Although most subjects experienced transient antibiotic responses, we found that five subjects
(n=5/22, 23%) experienced major changes in species composition that persisted for >1 month
after ciprofloxacin perturbation (**Fig. 1e** and **Fig. 2**, **Extended Data Figs. 4-7**). In two of these
subjects (XDA and XEA), the dominant bacterial family shifted from *Prevotellaceae* to
Bacteroidaceae, mirroring microbiome changes observed in industrialized populations[47]. In the
remaining three subjects (XAA, XBA, and XKA), species-level changes within *Bacteroidaceae*
drove these lasting shifts in microbiome composition (**Extended Data Fig. 3c,h**). Despite these
changes in microbiome composition, subjects with lasting antibiotic responses underwent
changes in absolute abundance, as estimated by colony plating, that were broadly comparable
to those in subjects with transient responses (**Extended Data Fig. 3d; Methods**).
143

We investigated various host and microbial factors that might explain these heterogeneous
responses. However, we found no consistent relationship with prior antibiotic usage (**Extended**
Data Fig. 3e) or pre-perturbation community diversity (**Extended Data Fig. 3f**; Wilcoxon
p=0.22). We also found no relationship with the amount of strain sharing at the beginning of the
study (**Extended Data Fig. 3g**). Instead, we found that the lasting antibiotic responses were
most strongly predicted by the magnitude of the initial perturbation (**Fig. 1d**). This link between
perturbation magnitude and duration suggests that sufficiently large perturbations can cross a
"tipping point" that overcomes the resilience of the gut microbiome and shifts the community into
a new state[39,40,45,46].
153

Rapid recovery of resident strains in subjects with transient antibiotic responses

The heterogeneous responses to ciprofloxacin present an opportunity to link these large-scale
changes in community composition to subsequent colonization dynamics. Prior work has shown
that antibiotics cause species extinctions that clear niches in the gut and increase colonization
by opportunistic pathogens[21–23,36,48] and introduced strains[37,38]. We therefore sought to quantify
these species-level dynamics within each community (**Fig. 3**, **Extended Data Fig. 8**),
hypothesizing that subjects who experienced more extensive species disruptions would
experience elevated levels of colonization and transmission.
162

Many species experienced large declines in relative abundance (>32-fold) during the 5-day
course of ciprofloxacin (**Fig. 2a-e**, **3a**, and **Extended Data Fig. 8a**). These disrupted species
accounted for ~20-80% of the initial gut microbiome composition in subjects with transient
ciprofloxacin responses and ~70-95% in lasting responders (**Extended Data Fig. 8b**). In
contrast, we observed few disruptions in control subjects: disrupted species comprised a
median relative abundance of 0.4% at the beginning of the study, showing that species
disruptions are uncommon in the absence of perturbations[18,44,49,50].
170

Bacterial families had varying responses to antibiotics: across all ciprofloxacin-taking subjects,
the *Rikenellaceae*, *Porphyromonadaceae*, and *Ruminococcaceae* families were significantly
enriched among disrupted species, whereas the *Lachnospiraceae* and *Eubacteriaceae* were
significantly less likely to experience disruption (**Fig. 3b**; **Methods**; permutation test, Bonferroni-
corrected p<0.05). The initial proportions of these taxa were not significantly different between
subjects with transient and lasting antibiotic responses, suggesting that the ciprofloxacin
susceptibility of individual taxa does not fully account for the heterogeneity we observe across
subjects (**Extended Data Fig. 8c**).
179

After the ciprofloxacin course ended, most disrupted species in subjects with transient antibiotic
responses recovered to within an order of magnitude of their pre-antibiotic relative abundance
(n=247/821, 70%; **Fig. 2d** and **3a**; **Extended Data Fig. 8b**). By calculating the genetic similarity
of the pre- and post-antibiotic strains, we determined that this microbiome recovery was driven
primarily by the recovery of the resident strain, or recolonization by a close relative, rather than
colonization by an unrelated strain of the same species (**Fig. 3c**). Of the 133 recovered
populations in subjects with transient responses that had sufficient sequencing coverage to
analyze strain-level composition, 129 (97%) involved the recovery of the pre-antibiotic strain,
and none of the remaining populations showed evidence of transmission from the cohabiting
partner (**Methods**).
190

191 We investigated whether strain transmission from cohabiting partners could contribute to these
192 high levels of resident strain recovery. Surprisingly, we found that the rate of recovery was

independent of any prior household sharing, both among all antibiotic-taking subjects (**Fig. 3d**; chi-square p=0.53) and among subjects with transient antibiotic responses only (**Extended Data Fig. 8g**, chi-square p=0.58). Most resident populations recovered without sharing strains with the cohabiting partner. The high rate of recovery of these populations (n=123/195, 75%) is inconsistent with ecological drift, suggesting that selection to fill open niches drives species recovery after antibiotic perturbation. However, we also observed many cases where a disrupted species failed to recover more than a month after antibiotics, even when the strain was shared with the cohabiting partner (**Fig. 2c** and **Fig. 3d; Extended Data Fig. 8d** and **8e**). These minimal rates of recolonization are especially striking given how rapidly most of the resident strains recovered (**Fig. 3c**). These data show that there can be substantial ecological barriers to transmission between cohabiting subjects, even when niches appear to be available in the gut microbiome.

Resident strains restore colonization resistance in subjects with transient antibiotic responses

To quantify the dynamics of colonization after these niche-opening perturbations, we searched for newly colonizing species that were not detected at the beginning of the study (**Fig. 2f** and **2g**) as well as newly colonizing strains of an existing species that were not detected at the beginning of the study and had high genetic divergence from the initial resident strain (**Fig. 2e**; **Methods**). In subjects with transient responses, the rapid recovery of resident strains appeared to limit the opportunities for colonization in the month after antibiotic perturbation. No new colonizing strains or species were detected in over half of these subjects (n=9/17, 53%; **Extended Data Fig. 8h**), and the total relative abundance of newly colonizing strains and species was not significantly different from the control subjects (**Extended Data Fig. 8i**; Wilcoxon rank-sum test p=0.10).

One explanation for the limited colonization in the first month after ciprofloxacin is that it could take new colonizers a longer time to reach detectable relative abundances. We tested this idea by investigating the longer-term dynamics of colonization in follow-up samples collected for up to two years after the antibiotic perturbation. Of the 22 households who completed the main, two-month sampling design, 18 households collected a median of 3 follow-up samples per subject over a median of 12 months after the end of the main study. This expanded dataset allowed us to quantify the dynamics of colonization over longer timescales.

These data revealed that, even after two years of follow-up sampling, colonization remained limited in subjects with transient antibiotic responses (**Fig. 3c**). The number (median: 0; Wilcoxon rank-sum test, p=0.61) and relative abundance (median: 0.4%; Wilcoxon rank-sum test, p=0.36) of new colonizers in these subjects remained similar to the control subjects, who also experienced minimal new colonization even over these longer timescales (**Extended Data Fig. 9a,b**). The main exception to this trend occurred in subject XIA: a new strain of *Prevotella copri* that was not previously detected in this subject's household colonized 5-8 months after the antibiotic perturbation and reached a maximum relative abundance of nearly 70% (**Extended Data Fig. 8f**). This example shows that new colonizers can occasionally have major effects on gut microbiome composition. Overall, however, the limited colonization we observe up to two years after ciprofloxacin perturbation suggests that the rapid recovery of resident strains in subjects with transient antibiotic responses is sufficient to restore the natural colonization resistance of the gut microbiome.

Extensive species losses and disruptions of colonization resistance in subjects with lasting antibiotic responses

We hypothesized that subjects with lasting antibiotic responses might experience higher levels of new colonization. Although the proportion of disrupted species was only slightly higher in these subjects compared to those with transient responses (**Fig. 3a, Extended Data Fig. 8b**), these species exhibited much lower rates of recovery. Only ~25% (n=78/309) of disrupted species recovered in the month after the ciprofloxacin course ended, compared to ~70% (n=574/821) of disrupted species in subjects with transient responses (**Fig. 3a**). Most disrupted species did not recover and instead appeared to be lost from the gut microbiome, even when they were shared with a cohabiting partner (**Fig. 3d**).

In contrast to the subjects with transient responses, many new strains and species reached detectable abundances in the month after antibiotics in subjects with lasting responses (**Fig. 3c** and **Extended Data Fig. 8h**). Most of these events appeared to involve the introduction of new species rather than a new strain of an existing species (**Fig. 3c**; n=47/53, 89%), though sequencing coverage limits our ability to identify new strains in rare species. Compared to resident species, these new colonizers had significantly lower prevalence (**Fig. 3e**; Wilcoxon rank-sum test, $p<10^{-15}$) and median abundance (**Extended Data Fig. 8j**) among subjects in the Human Microbiome Project, suggesting that these new species are uncommon in healthy gut microbiomes. The *Campylobacteraceae* and *Veillonellaceae* families were enriched among new colonizers (**Fig. 3b**; **Methods**; permutation test, Bonferroni-corrected p<0.05), which included common oral bacteria like *Campylobacter concisus*, *Dialister invisus*, and *Megasphaera micronuciformis*[51–54] (**Extended Data Figs. 4b** and **5b**). These findings are consistent with prior work showing that oral microbes are frequently transmitted into the gut[55] and can reach high relative abundances when gut commensals decrease in absolute abundance[56]. We also identified potential opportunistic pathogens[21,48,57] like *Klebsiella pneumoniae*, *Citrobacter freundii*, and *Streptococcus gallolyticus* (**Extended Data Fig. 5b**), although no subjects reported side effects or illnesses after taking ciprofloxacin. Altogether, many species expansions in the month after antibiotic perturbation were consistent with overgrowth of potential pathogens, a phenomenon that is common after antibiotic perturbations of the gut microbiome and is often ascribed to a disruption of colonization resistance[48].

Although we detected many new species in the month after antibiotics in subjects with lasting antibiotic responses, these new species had limited effects on long-term gut microbiome composition. New species were often transient (**Fig. 2f**), and only 25 of 47 (53%) new species were still detectable a month after the ciprofloxacin course ended (**Fig. 3c**). The primary exception was *Bacteroides stercoris* in subject XKA, which transmitted from the cohabiting partner after ciprofloxacin perturbation and reached a relative abundance of ~60% by day 64, which it maintained throughout the >1.5 years of follow-up sampling (**Extended Data Fig. 8k**). Overall, however, these examples of commensal colonization were surprisingly rare in the month after antibiotics, suggesting that substantial ecological barriers to colonization can remain even after major microbiome disruptions.

Commensal strains abruptly colonize subjects with lasting antibiotic responses, but only after long delays

Ecological theory[58,59] suggests that the low levels of commensal colonization in the subjects with lasting antibiotic responses could be driven by the functional redundancy of the gut microbiome. If the surviving resident species in these subjects can fill the niches cleared by antibiotics, we would expect that colonization would remain limited over longer timescales as well because the gut microbiome has reached a new stable state.

Surprisingly, we found that even though commensal colonization was limited in the month after antibiotics, subjects with lasting responses experienced elevated colonization more than a year after the original perturbation (**Fig. 4a,c** and **Extended Data Fig. 9a,b**). In four of these five subjects, new strains and species comprised ~15-70% of the gut microbiome one to two years after antibiotic perturbation, compared to only 0.4% in their cohabiting partners (**Extended Data Fig. 9c**; Wilcoxon rank-sum test, p=0.01). Of the 25 new colonizers that were still detectable at the end of the study and had sufficient sequencing coverage for strain analysis, only one-third (n=8, 32%) were detected in the cohabiting partner, while the rest were acquired from other sources (**Methods**). This elevated colonization in subjects with lasting antibiotic responses suggests that major, antibiotic-mediated disruptions can create niches for new microbes to colonize.

However, while the overall rates of colonization were elevated in these subjects, our longitudinal data revealed that the colonization events only took place after long delays (**Fig. 4a,b**). We continued to detect new colonization events more than nine months after the ciprofloxacin perturbation ended, and in subjects XAA and XDA, more than half of the colonization events detected during follow-up sampling occurred more than a year after antibiotics (**Fig. 4a**). This lengthy timescale of colonization far exceeds the 1-2 week timescale of resident strain recovery (**Fig. 3c**), even though these resident strains frequently recovered from undetectable ($<10^{-5}$) relative abundances after antibiotics (**Fig. 2a,d**). The large gap between these timescales suggests that new species face substantial ecological barriers to colonization. These long delays are also reminiscent of ecological literature showing that species introduced into new environments often display invasion lags, persisting at low abundances for long periods of time before undergoing exponential expansions[60,61].

Some delays in the detection of new colonizers are naturally expected due to the time it takes for a species to grow from its initial inoculum to reach detectable relative abundances. Due to the large number of cells in the human gut[62], these growth lags can be substantial even under strong ecological selection (**Methods**). We can estimate the strength of this effect using our longitudinal data, by comparing the relative growth rates of the invading species with the total time it takes for them to invade (**Fig. 4b, Methods**).

We fit the relative abundance trajectories of new colonizers to a simple ecological model where each species grows exponentially until it reaches its inherent carrying capacity (**Fig. 4b, Methods**). These data revealed that new colonizers expanded rapidly despite their lengthy colonization delays. Of the 112 new strains and species detected more than one month after antibiotic perturbation, 76 (68%) new colonizers increased from undetectable ($<10^{-5}$) to maximum relative abundances between consecutive sampling timepoints (**Fig. 4d**). While the total time between samples varied across our cohort (**Extended Data Fig. 9d**), some of these intervals comprised <5% of the total colonization time and were consistent with a net growth advantage for the colonizer of at least 0.5 doublings per day. We used these inferred relative growth rates to calculate the maximum time required for a newly introduced species to reach

carrying capacity, starting from a single cell. Our conservative estimates suggest at least 33 of the 112 (29%) new colonizers were introduced at least six months after the end of antibiotics, with the remaining examples mostly limited by the temporal resolution of our follow-up sampling (**Extended Data Fig. 9d**). This rapid growth shows that new colonizers experience strong ecological selection, and that detection lags cannot fully explain the large colonization delays we observe.

What ecological forces could delay colonization in the face of such strong ecological selection? One hypothesis is that the influx of viable microbes into the gut is extremely low, and the rate of colonization is determined by the rate of these rare microbial introductions. In this case, the inferred introduction times provide a corresponding bound on the overall rates of introduction. For example, the ~6 month delay after antibiotics and before the inferred introduction of *Parabacteroides johnsonii* in XDA (**Fig. 4b**) implies an average influx of <0.01 successful migrants per day (**Methods**). However, we also observed strong temporal clustering of colonization events (**Fig. 4a**), suggesting that the simplest models of dispersal limitation do not apply. In subjects XKA, XDA, and XAA, we identified 2, 5, and 5 colonization events respectively at a single timepoint—more than would be expected to occur by chance if colonization events occurred independently (**Extended Data Fig. 9e**; **Methods**; permutation test, $p < 0.05$).

These clusters of new colonizers often contained groups of closely related species. For example, *Bacteroides caccae* and *Bacteroides clarus* were both present in the cluster of 5 colonizers in XDA (**Fig. 2a**), while *Alistipes onderdonkii, Alistipes putredinis,* and *Alistipes shahii* were present in the cluster of 5 colonizers in XAA (**Extended Data Fig. 4b**). We also detected two new strains of *Phocaeicola vulgatus* that simultaneously colonized XAA at a different timepoint (**Extended Data Fig. 9f**). This clustering of closely related strains raises the intriguing possibility that facilitative interactions between "guilds" of gut microbial strains[63] can promote their joint dispersal or colonization and contribute to invasional meltdown[64], in which invasive species facilitate each other's establishment.

We wondered whether the strains that were previously shared with the cohabiting partner might play an outsized role in this long-term recovery, since they might have greater compatibility with the rest of the community or increased opportunities to transmit and recolonize. However, in 4 of the 5 subjects with lasting antibiotic responses, we found that the recovery of resident strains always occurred within the same narrow time window as in subjects with transient responses (**Fig. 3c**). With the exception of subject XBA (see below), we did not identify any instances in which a shared strain that failed to recover in the month after antibiotics recolonized at a later date (**Fig. 3c** and **4a**). Instead, in each of these subjects, we identified at least one strain that was shared before antibiotics but failed to recolonize in the 1-2 years after the antibiotic perturbation. Over the same period, we also identified at least one new transmission of a strain from the cohabiting partner that was not previously shared, suggesting that there is still ongoing microbial transmission (**Fig. 4e**).

For example, before antibiotics, subjects XDA and XDB shared a strain of *Eubacterium eligens* that was present at a relative abundance of ~2% in each subject (**Fig. 2c**). In XDA, *E. eligens* was disrupted by ciprofloxacin and was never detected in follow-up sampling, even though the same strain was maintained at constant relative abundance in XDB. During this time, five new strains transmitted from XDB to XDA (**Fig. 2a, 4e**), suggesting that dispersal limitation alone is unlikely to explain the absence of *E. eligens* recolonization. Instead, these findings suggest that ecological factors like priority effects and changing selection pressures limit recolonization, creating historical contingency in microbiome composition.

Rapid transitions between colonization-resistant alternative states in a subject with a lasting antibiotic response

While four of the five subjects with lasting antibiotic responses experienced elevated levels of colonization, the remaining subject (XBA) exhibited markedly different dynamics. New colonizers accounted for only ~3% of this subject's gut microbiome nearly two and a half years after antibiotics, prompting us to more closely examine the dynamics of this community.

By sequencing 106 timepoints over a two-year period, we found that XBA's gut microbiome underwent sudden, striking transitions between three distinct community states (**Fig. 5a** and **Extended Data Fig. 10a**). Prior to antibiotics, the subject had a diverse gut microbiome, with multiple co-resident *Bacteroidaceae* species (state 1; days -28-0). The ciprofloxacin perturbation led to the monodominance of a single species, *Bacteroides fragilis*, which maintained a relative abundance of nearly 50% for more than a year (state 2; days 0-401). After day 390, community diversity sharply increased over three timepoints in a two-week span, as eight *Bacteroidaceae* species simultaneously underwent dramatic increases in relative abundance (**Fig. 5a** and **Extended Data Fig. 10b**). This third state (days 401-881) was stably maintained through the end of follow-up sampling. The transition from state 2 to state 3 was not associated with any known changes in the diet, lifestyle, or health status of XBA. These long periods of community stability, punctuated by sudden shifts in species abundance, suggest that the gut microbiome can undergo rapid transitions between alternative states[65,66].

We investigated whether the transition to state 3 in XBA was driven by 1) strain persistence, in which strains present before antibiotics drive species recovery, or 2) new colonization, including recolonization from the cohabiting partner (**Fig. 5b-d**). Among species with sufficient sequencing coverage to analyze strains, we identified cases of both strain persistence and new colonization. *Bifidobacterium longum,* which became undetectable for several months after antibiotics, became detectable again three months before the transition to state 3. This post-antibiotic strain of *Bifidobacterium longum* was more genetically similar to the pre-antibiotic strain in XBB than the pre-antibiotic strain in XBA, suggesting that its expansion was driven by colonization rather than strain persistence (**Fig. 5b,c** and **Extended Data Fig. 10c; Methods**). *B. longum* can inhibit *B. fragilis* in co-culture[67] and reduce its microbial load *in vivo*[68], suggesting that the colonization of *B. longum* may have facilitated the return of other *Bacteroidaceae*. However, in contrast to *B. longum*, the strains of *B. uniformis* and *P. vulgatus* that we detected in state 3 were more similar to the pre-antibiotic strains in XBA than the pre-antibiotic strains in XBB (**Fig. 5d** and **Extended Data Fig. 10d-f**), indicating that these strains persisted through the *B. fragilis*-dominated state 2 before undergoing rapid, simultaneous expansions. These dynamics bear striking similarities to the delayed, rapid, clustered colonization dynamics that we observed in the other four subjects with lasting antibiotic responses, suggesting that they may be driven by similar ecological forces.

Curiously, we found that the changes in community composition in XBA coincided with sharp shifts in the relative abundance of Type VI secretion system (T6SS) genes (**Fig. 5e** and **Extended Data Fig. 10g**). Many *Bacteroidaceae* carry T6SS gene cassettes, which contain effector and immunity genes that allow cells to kill neighboring cells that lack the immunity gene[69–72]. Strains that carry distinct T6SS effector-immunity (E-I) pairs can antagonize one another[69–72], which may contribute to the maintenance of distinct community states[69]. Intriguingly, although XBA and XBB shared many *Bacteroidaceae* strains, their communities had different dominant E-I pairs (**Fig. 5e**), and we speculate that this incompatibility may have limited the ability of *Bacteroidaceae* to transmit and colonize XBA after ciprofloxacin perturbation. We also observed changes in the T6SS composition of subjects XAA and XKA,

who had lasting antibiotic responses and underwent major shifts in the abundance of *Bacteroidaceae* species, suggesting that T6SS may play a role in other subjects as well (**Extended Data Fig. 10h** and **10i**). From this observational study alone, we cannot determine whether the changing abundances of T6SS genes are the cause or consequence of the rapid switches between alternative community states, but these analyses raise the possibility that interbacterial antagonism could contribute to community stability and colonization resistance in the gut microbiome[57,73].

Discussion

Our results show that strong ecological barriers limit colonization in the adult gut microbiome, even after antibiotic perturbations that cause extensive species losses. These findings counter the prevailing view that colonization is primarily limited by the availability of open niches, suggesting instead that the introduction and initial expansion of new microbes also present major barriers to colonization in natural settings.

What ecological forces are responsible for these prolonged delays in colonization after antibiotic perturbation? The rapid colonization dynamics we observe (**Fig. 4b**), together with the elevated levels of colonization in subjects with lasting antibiotic responses (**Fig. 4c**), suggest that new colonizers experience strong ecological selection. A common explanation for colonization delays is that dispersal limitation strongly restricts the pool of available colonizers, slowing the pace of colonization[7]. However, the coordinated clusters of colonization events that we observe are unlikely to arise from random introductions of individual species (**Fig. 4a** and **Extended Data Fig. 9e**), evidence against the simplest models of dispersal limitation.

We argue instead that collective effects shape colonization dynamics by causing microbial fitness to depend strongly on aspects of community context like population size and interactions with other microbes. These collective effects could arise during dispersal—for example, if microbial species migrate together in larger propagules, a form of community coalescence[74] (**Fig. 4a** and **Extended Data Fig. 9e**). They can also arise after dispersal, creating priority effects[33,75] in the resident community. Priority effects can occur through inhibitory interactions like interbacterial antagonism, which allows established strains to kill new arrivals[57,76] (**Fig. 5e**), or through facilitative interactions like quorum sensing[77], cross-feeding[78], and the production of public goods[79]. Regardless of their mechanism, strong collective effects in the gut microbiome complicate the design of microbiome therapeutics by creating historical contingency in colonization outcomes (**Fig. 4e**). Large inoculation doses and tailored microbial consortia may be necessary to overcome these ecological barriers and successfully colonize established microbial communities.

This work shows how the combination of controlled perturbations, longitudinal sampling, and strain-resolved metagenomic sequencing can disentangle the ecological forces that shape colonization in natural settings. Our conclusions are limited by the observational nature of our study, as well as the difficulty of detecting colonization events at low relative abundances. Despite these limitations, *in situ* studies like ours are necessary for investigating how colonization is shaped by uniquely human aspects of lifestyle and gastrointestinal anatomy, factors that are difficult to replicate in mouse models[42,80] or *in vitro* communities[81]. Future work that combines our approaches with barcoded strain libraries[82] or spatially resolved sampling devices[83] can further illuminate the ecological and evolutionary forces that shape colonization in the human gut.

Materials and methods

Data availability

Sequencing data will be made publicly available from the NCBI SRA upon manuscript publication. The computer code that performs the analysis is available at https://github.com/ksxue/household-transmission-after-abx.

Study design

This study was approved by the Stanford University Institutional Review Board as Protocol 54715, and written, informed consent was obtained from all participants. Healthy, cohabiting adults were recruited from Stanford University and the surrounding community (**Extended Data Fig. 1a-d**). Exclusion criteria included antibiotic usage in the past three months, past reactions to fluoroquinolone antibiotics, chronic illness, pregnancy or nursing, and age under 18. During the main, two-month sampling period, participants collected weekly stool samples for 9 weeks (days -28, -21, -14, -7, 0, 7, 14, 21, 28, 35) and daily stool samples for 19 days in the middle of the study (days -7 to 11) (**Extended Data Fig. 1e**). One participant in each household took a 500mg oral dose of ciprofloxacin twice daily for 5 days (days 0-4). At each time point, participants collected one stool swab that was stored in 1mL RNAlater (Thermo Fisher AM7021) and collected approximately 20mL of raw stool in a sterile vial. Stool samples were frozen immediately after collection in subjects' home freezers and were stored there for up to four weeks until they were transferred to storage at -80 degrees C in the laboratory. A total of 48 participants in 22 households completed the sampling protocol. Participants from 18 households collected follow-up samples using the same methods at occasional intervals for up to two years after the main sampling period (**Extended Data Fig. 1e**).

Participants also completed a questionnaire at the beginning of the study that included questions about demographics, diet, lifestyle, recent travel and antibiotic use, household structure, and cohabitation history. At the end of the study and every three months during follow-up sampling, participants were asked to report changes in diet, lifestyle, and living arrangements, as well as recent travel, illnesses, injuries, and antibiotic usage.

DNA extraction and metagenomic sequencing

DNA was extracted from stool swabs stored in RNAlater using the DNeasy PowerSoil HTP 96 Kit (Qiagen 12955-4). To avoid cross-contamination that could lead to erroneous inferences of strain sharing[84], samples collected from cohabiting partners during the main, two-month sampling period were arrayed on different 96-well plates. When processing follow-up samples, which were fewer in number, samples were arrayed so that no sample was adjacent to a sample from a cohabiting partner. For stool swabs that yielded less than 1.5ng/uL of DNA, we aliquoted a stool fragment from the accompanying sample of frozen, raw stool and redid the DNA extraction using the DNeasy PowerSoil Kit (Qiagen 12888-100).

Libraries were prepared for metagenomic sequencing using the Nextera DNA Flex Library Prep kit (Illumina 20018705). Samples collected at four "key" timepoints (days -28, 0, 7, and 35) from both control and antibiotic-taking subjects were sequenced to a target depth of 10Gbp. All remaining samples collected from antibiotic-taking subjects during the main, two-month sampling period were sequenced to a target depth of 2Gbp. All follow-up samples collected from both control and antibiotic-taking subjects were sequenced to a target depth of 2Gbp. In total, we sequenced 942 samples collected from the 48 subjects in our study (**Extended Data Fig. 1e**).

CFU counts

We drilled cores from stool samples frozen at -80 degrees C using the CXT 353 Benchtop
Frozen Aliquotter (Basque Engineering) and aliquoted 50-100mg of frozen stool from each
sample. We resuspended each stool fragment in 0.5mL reduced PBS and plated 10-fold
dilutions on Brain-Heart Infusion agar plates supplemented with 5% sheep's blood in an
anaerobic chamber (Coy Instruments). We incubated plates at 37 degrees C and counted
colonies after 48 hours.
540

Species and SNP profiling of metagenomic samples

We first used skewer[85] to trim sequencing adapters and bowtie2[86] to filter out reads that mapped
to the human genome. Remaining reads will be made available in the SRA upon manuscript
publication. To obtain species abundances, we used the MIDAS[43] species module to map
sequencing reads against a database of universal, single-copy bacterial genes. We used these
sequencing reads to calculate species abundances, which were used to compute Jensen-
Shannon divergence and normalized F_{ST}[87], a metric of compositional variability.
548
Using these species abundance estimates, we generated a list of species that were detected in
a given subject across all sequenced timepoints. We then used the MIDAS snps module to map
sequencing reads against a set of dereplicated reference genomes representing the detected
species. For each population, defined by the set of reads in a sample that mapped to a species'
reference genome, we calculated the relative abundance of single-nucleotide polymorphisms
(SNPs) at every genome site. To limit the influence of reads with high similarity to multiple
reference genomes, we discarded sites that had sequencing coverage more than two-fold
above or below the median genome coverage.
557

Identifying cross-contamination and other sample abnormalities

To identify cross-contamination during DNA extraction and library preparation that could
confound our identification of shared strains between subjects, we calculated the number of
fixed differences between each pair of populations with a median sequencing coverage >5, and
we identified all unusually similar population pairs, defined as population pairs from non-
cohabiting subjects that had fewer than 3 fixed differences. We identified samples that had
unusually similar population pairs for multiple species, suggesting that this population similarity
was likely to be due to cross-contamination rather than genetic similarity between strains.
566
In an initial analysis, we identified 24 samples that had signatures of cross-contamination. Most
of these samples were located along the edge of a single DNA extraction plate, suggesting that
improper sealing of this plate during DNA extraction may have contributed to cross-
contamination[84]. The remaining samples with signatures of cross-contamination were in
adjacent wells on either their DNA extraction or metagenomic library preparation plates. We
removed these samples from downstream analyses.
573
We re-sequenced these samples by aliquoting a stool fragment from the banked sample of
frozen, raw stool, extracting DNA using the DNeasy PowerSoil Kit (Qiagen 12888-100), and
preparing libraries for metagenomic sequencing using the Nextera DNA Flex Library Prep kit
(Illumina 20018705). We re-calculated the number of fixed differences between each pair of
populations with median sequencing coverage >5 after including these re-sequenced samples

in our dataset. We identified no samples from non-cohabiting subjects that had unusually similar
population pairs for multiple species, suggesting that the level of cross-contamination between
our remaining samples was lower than our limit of detection.
582
We also identified one case in which we suspected that subjects swapped identifiers.
Household XH reported that subject XHC took ciprofloxacin. However, XHC had fewer species
disruptions than any other antibiotic-taking subject, and we detected more disrupted species in
XHB than in XHC (**Extended Data Fig. 8a** and **8b**). We therefore classified subject XHB as the
antibiotic-taking subject from this household. The main conclusions of our study are not affected
by whether we designate XHB or XHC as the antibiotic-taking subject.
589

Classifying subject responses to antibiotic perturbation

For each subject, we calculated the Jensen-Shannon divergence (JSD) between the species
abundances in the initial sampling timepoint and each subsequent sequenced timepoint. We
calculated the maximum JSD from the initial timepoint between days -28 and 35, as well as the
final JSD from the initial timepoint to day 35. Antibiotic-taking subjects were classified as having
a transient antibiotic response if they had a final JSD <0.4 and a lasting antibiotic response if
they had final JSD >0.4.
597

Identifying shared strains

We used two complementary methods to infer whether a pair of populations shared strains. In
the first, "fixed differences" method, which we adapted from InStrain[88], we calculated the
number of fixed differences between each pair of populations with median sequencing coverage
>5. We define a fixed difference as a site at which the two populations carried distinct major
alleles (frequency>0.5), and the major allele in each population was detected in the other
population at a frequency below the Illumina sequencing error rate of 10^{-3}. We determined that
two populations shared strains if there were fewer fixed differences between them than in 99%
of population pairs from non-cohabiting subjects, and we classified two populations as not
sharing strains if the number of fixed differences between them exceeded this threshold
(**Extended Data Fig. 2a,b**). We classified strain sharing as unknown if neither population in a
pair had median sequencing coverage >5.
610
We also developed a second, "strain fishing" method for identifying shared strains using "private
marker" SNPs[41,44]. We first identified "quasi-phaseable" (QP) populations in which we could infer
the genotype of the dominant strain[44]. We classified a population as QP if it had median
sequencing coverage >5 and we detected SNPs with frequency between 0.2 and 0.8 at <0.1%
of the genome, meaning that the population had low genetic variation. In each QP population,
we identified private marker sites at which the major allele in the QP population was the minor
allele (frequency<0.5) in all populations of the same species analyzed by Garud, Good *et al.*[44],
which includes the Human Microbiome Project and several additional cross-sectional cohorts.
We determined that two populations shared strains if either population was QP and we detected
the QP allele at any frequency above the Illumina sequencing error rate of 10^{-3} at >75% of
private marker sites in the other population. We classified two populations as not sharing strains
if the QP allele was detected above a frequency of 10^{-3} at <75% of private marker sites
(**Extended Data Fig. 2c,d**). We classified strain sharing as unknown if neither population in a
pair was QP.
625

We found that the fixed differences and strain fishing methods agreed on >90% of population
pairs that could be analyzed using both methods (**Extended Data Fig. 2e**), and we combined
the calls from both methods to generate our final classifications of strain sharing. Strain sharing
in a population pair was classified as "unknown" if neither method was able to classify strain
sharing or if the two methods returned discordant results; "not shared" if one or both methods
classified the population pair as not sharing strains; and "shared" if one or both methods
classified the population pair as sharing strains.
633

Species disruption, recovery, and colonization

We classified species responses to antibiotics using their relative abundance dynamics. A
species was classified as "disrupted" if its median pre-antibiotic relative abundance exceeded
10^{-3} and its minimum post-antibiotic abundance declined by more than 1.5 orders of magnitude
below its median pre-antibiotic relative abundance. A disrupted species was also classified as
"recovered" if its median post-antibiotic relative abundance in a three-sample window exceeded
one-tenth of its median pre-antibiotic relative abundance. We calculated the time of recovery to
be the first timepoint in this window during which the species exceeded one-tenth of its median
pre-antibiotic relative abundance. Species whose median pre-antibiotic relative abundance
exceeded 10^{-3} and were not classified as "disrupted" were classified as "maintained."
644

645 A species was classified as "colonizing" if it was absent at the first three sequenced timepoints
646 and its median relative abundance in a three-sample window subsequently exceeded 10^{-3}. We
647 calculated the time of detection to be the first timepoint in this window during which the species
648 exceeded 10^{-3} relative abundance.
649

Colonization of new strains of an existing species

We also sought to identify cases in which new strains of existing species colonized the gut
microbiome. We used the methods described above to determine whether strains were shared
between a population at the initial and subsequent timepoints from each subject ("**Identifying**
shared strains"). Populations were classified as candidate colonizing strains when all
timepoints past a certain timepoint (designated as the time of strain colonization) were classified
as strains "not shared" or "unknown" when compared with the initial timepoint, suggesting that
the initial strain was no longer detected.
658

We reverified candidate colonizing strains through longitudinal strain analysis of candidate new
colonization cases. As before, we calculated the relative abundance of single-nucleotide
polymorphisms (SNPs) at every genome site for each population. To limit the influence of reads
with high similarity to multiple reference genomes, we discarded sites that had sequencing
coverage more than 2.5- fold above or below the median genome coverage. We only kept sites
that passed this criterion for at least 80% of samples measured for that population. We found
SNPs that underwent selective sweeps by identifying SNPs that grew from a frequency below .2
to frequency above .8 between a pair of timepoints. We excluded sweeping SNPs where the
coverage ratio of the site to the median depth of the genome changed by more than a factor of
two between the two timepoints. A strain turnover event occurred if at least 1000 SNPs swept
between a pair of timepoints, indicating that a minor strain rose from low to high frequency.
670

671 We searched each population for strain turnover events in which a low-frequency strain that was
672 not initially present swept to high frequency. We calculated the frequency of the minor strain as

673 the median frequency of the SNPs associated with the sweeping strain. We accepted the new
674 colonization call if and only if the frequency of the minor strain was zero at the initial timepoint.
675

Mathematical modeling of colonization delays

To interpret the colonization and recovery dynamics we observed, we fit the relative abundance
trajectory of each species to a simple mathematical model, in which the resident population
grows exponentially at a constant rate r before reaching an intrinsic carrying capacity K. We
allowed the growth rates and carrying capacities to vary between species, and between different
populations of the same species in different hosts. Each trajectory in this model can therefore
be specified by a triplet of parameters, r, K, and τ, where τ is the time that the population first
reaches carrying capacity (the "colonization time" in **Fig. 3c** and **4a**).
684

We estimated the model parameters by minimizing the sum of squared errors between the
observed and predicted values of the logarithm of the relative abundance. We only included
timepoints where the observed relative abundance of the species exceeded our limit of
detection, which we estimated as the inverse of the total marker gene coverage summed across
all species in that sample. We also restricted the model fit to the time interval where the first
colonization (or re-colonization) event was estimated to have occurred. To eliminate potential
degeneracies in the inferred parameters, we added a weak regularization term to the loss
function,
693

695 $$R(r, K) = 0.1 \log r + 0.0002 (\log K + 11)^2$$
694

696 which favors the smallest values of r and K that are consistent with the observed data. Error
697 minimization was performed in a custom Python script using the minimize function from the
698 SciPy library[89].
699

700 Using these inferred parameters, we estimated the total time required for each species to grow
701 to its final carrying capacity starting from a single founding cell. Under simple models of
702 demographic noise[90], this time can be approximated by the asymptotic formula,
703

709 $$\Delta t = \frac{1}{r} \log(N_e r K)$$
710

704 where N_e is the effective population size of a single person's gut microbiome. To be
705 conservative, we used an effective population size of $N_e \sim 10^{13}$, which corresponds to the total
706 number of bacterial cells in a typical person's gut[62]. This large effective population size implies
707 that the total growth times can be quite long. For example, a growth advantage of $r = 1$ per day
708 and a carrying capacity of $K \sim 0.01$ corresponds to a total growth time of $\Delta t \sim 25$ days.
711

712 If the estimated value of Δt is less than τ, then the difference $\tau - \Delta t$ represents the earliest
713 possible time that the colonizing cell was first introduced. Under the assumption that successful
714 colonizers arrive as a Poisson process, the expected value of $\tau - \Delta t$ is equal to the inverse of
715 the rate of successful introductions. We used this relation to convert the minimum possible
716 introduction time into a corresponding bound on the rate of introduction.
717
718

Statistical tests of taxonomic enrichment

We used permutations to test whether certain bacterial families were enriched among disrupted species (**Fig. 3b**). To test for taxonomic enrichment among disrupted species, we began with a simple null model in which species from all bacterial families are equally likely to be disrupted by ciprofloxacin. To generate a null distribution of the number of disrupted species in each family, we identified all populations with median pre-antibiotic relative abundance above 10^{-3} in the ciprofloxacin-taking subjects, and we drew populations at random without replacement, matching the total number of disrupted populations observed across all ciprofloxacin-taking subjects. We calculated the number of disrupted populations in each bacterial family in this simulated dataset, and we compared this distribution to the number of disrupted populations in each bacterial family observed in our ciprofloxacin-taking subjects. To calculate the p-values shown in the text, we calculated the fraction of simulations with a more extreme number of disrupted populations in each family than the observed value and performed a Bonferroni correction by dividing by the number of families tested.

We used an analogous approach to test for taxonomic enrichment among colonizing species (**Fig. 3b**) and species that were disrupted by antibiotics but failed recover (**Extended Data Fig. 8d**).

Comparing the prevalence and abundance of resident species and new colonizers in the Human Microbiome Project

We downloaded the species abundance profiles analyzed in Good and Rosenfeld[91], which were generated by using the MIDAS[43] species module to map sequencing reads from the Human Microbiome Project[92] (HMP) against a database of universal, single-copy bacterial genes. We calculated the prevalence of each species, defined as the number of subjects in which the species was detected at any non-zero frequency, as well as the median relative abundance of each species in subjects that carry the species at a non-zero frequency.

We identified new colonizers as described above. We also identified a set of resident species, which were detected at any non-zero frequency at the beginning of the study and subsequently attained a median relative abundance $>10^{-3}$ in a three-sample window. These resident species fulfill the same abundance criteria that we used to identify new species, but, in contrast to new species, they were detected at the beginning of the study. We identified the HMP prevalence (**Fig. 3e**) and median relative abundance (**Extended Data Fig. 8j**) of each resident species and new colonizer in each subject.

Inferring co-colonization of P. vulgatus strains in subject XAA.

We observed that the pre- and post- antibiotic *P vulgatus* populations in XAA harbored thousands of SNPs at intermediate frequencies, suggesting co-colonization by two or more strains in both the pre- and post-antibiotic intervals. Our strain-sharing analysis above demonstrated that the major strains turned over between the pre- and post antibiotic timepoints, so we sought to determine whether the minor strains were new colonizers as well.

To carry out this analysis, we assigned individual SNPs to 4 strain haplotypes by manually clustering their allele frequency trajectories over time. We employed the same filters on SNPs and timepoints that we used for verifying the pre-existing strain calls above. We also polarized the SNPs such that the allele frequency referred to the minor allele in the cohort used for calling private marker SNPs above. A manual examination of the allele frequency spectrum at the initial

timepoint suggested that the resident population was composed of two diverged strains in a 60:40 ratio. We assigned SNPs to the major strain (strain 1) if their frequency in the initial timepoint fell between 60% and 95%, and to the minor strain (strain 2) if their frequency was between 5% and 40%.

We assigned SNPs to the post-antibiotic strains based on their frequencies in two key timepoints. Day 459 was a quasi-phaseable (QP) timepoint, and is thus well-suited for identifying the major strain. We assigned SNPs to the major strain in the post-antibiotic period (strain 3) if their frequency was >80% this timepoint. Similarly, we found that one day in the post-antibiotic period (day 425) had a large number of SNPs between 30-70%, and is thus well-suited for identifying candidate SNPs that might belong to the minor strain. We therefore assigned SNPs to the minor strain in the post-antibiotic time window (strain 4) if their frequency at day 425 was >20%, but their frequency in all subsequent timepoints was <50%. We estimated the total number of genetic differences between strains (**Extended Data Fig. 9f**) by taking the differences between the lists of SNPs assigned to each strain.

We estimated the relative frequencies of the strains by taking the characteristic frequencies of the SNPs assigned to each cluster. Since a fraction of these SNPs will be shared across multiple strains, we restricted these estimates to the private marker SNPs in each cluster, which were inferred using the same procedure as above ("**Identifying shared strains**"). To mitigate the effects of mapping errors, we only considered SNPs in the list of core genomes, as previously described[44]. We also excluded SNPs that were present in both the major and minor strains in the post-antibiotic period. This procedure yielded a total of 32, 7, 17, and 23 private SNPs for strains 1-4, respectively. We estimated the frequencies of the respective strains as the median frequency of the private SNPs corresponding to that strain. The median frequency of the pre-antibiotic strains was zero in the post-antibiotic timepoints, and vice versa, confirming that the re-emergence of the *P. vulgatus* population was caused by the colonization of two new strains.

Distinguishing between strain re-emergence and strain replacement in subject XBA

We sought to determine whether increases in the relative abundance of *Bifidobacterium longum* and several *Bacteroidaceae* species in subject XBA after antibiotics was caused by the reemergence of pre-antibiotic strains or recolonization from XBA's cohabiting partner, XBB. We first identified species that had at least one QP population (median sequencing coverage >5, SNPs with frequency between 0.2 and 0.8 at <0.1% of the genome; see **Identifying shared strains**) before antibiotics and after antibiotics in both XBA and XBB. Of the 13 species shown in **Extended Data Fig. 10a**, *Bacteroides uniformis, Phocaeicola vulgatus, Bacteroides faecis,* and *Bifidobacterium longum* met these criteria.

Bacteroides uniformis. We identified 3800 sites in the core genome of *B. uniformis* at which the major allele in XBA (frequency>0.8) was the minor allele in XBB (frequency<0.2) in all pre-antibiotic, QP populations. At these sites, the major allele in XBA before antibiotics, which we called the XBA allele, had median frequency 1 in XBA and median frequency 0 in XBB before antibiotics. We called the strain carrying the XBA allele at these sites "strain 1," and we called the strain carrying the alternate allele "strain 2." The XBA allele was present at a median frequency of 1 at these sites after antibiotics in XBA and 0 in XBB, suggesting that the increase in relative abundance of *B. uniformis* was driven by the re-emergence of strain 1 rather than recolonization by strain 2 (**Fig. 5d** and **Extended Data Fig. 9d**). We identified zero sites in the

816 core genome at which the major allele in XBA before antibiotics (frequency>0.8) became the
817 minor allele after antibiotics (frequency<0.2), suggesting that there were no SNPs that reached
818 fixation between the end of antibiotics at day 5 and the re-emergence of *B. uniformis* around
819 day 401.
820
Phocaeicola vulgatus. We identified two sites in the core genome of *P. vulgatus* at which the
major allele in XBA (frequency>0.8) was the minor allele in XBB (frequency<0.2) in all pre-
antibiotic, QP populations. At these two sites, the major allele in XBA before antibiotics, which
we called the XBA allele, had median frequency ~0.85 in XBA and median frequency 0 in XBB
before antibiotics. We called the strain carrying the XBA allele at these two sites "strain 1," and
we called the strain carrying the alternate alleles "strain 2." The XBA allele was present at a
median frequency of 1 after antibiotics in XBA and 0 in XBB, suggesting that the increase in the
relative abundance of *P. vulgatus* was driven by the re-emergence of strain 1 (**Fig. 5d** and
Extended Data Fig. 9e).
830
We also identified and manually verified 7 sites in the core genome at which the major allele in
XBA before antibiotics (frequency>0.8) became the minor allele after antibiotics
(frequency<0.2). At these 7 sites, the major allele in XBA after antibiotics had a median
frequency of 1 in XBA and 0 in XBB, suggesting that these SNPs reached fixation on the
background of strain 1 between the end of antibiotics at day 5 and the re-emergence of *P.*
vulgatus around day 401 (**Fig. 5d** and **Extended Data Fig. 9f**). These sites were widely
dispersed across the genome, suggesting that they represent distinct mutations rather than a
single recombination event.
839
Based on the site-frequency spectrum of the *P. vulgatus*, we inferred that the XBB population
consisted of at least two distinct, co-colonizing strains. We identified 16011 sites in the genome
that had intermediate allele frequencies (0.2<frequency<0.8) in all non-QP populations from
XBB and had correlated frequencies across samples, suggesting that these sites distinguish the
two co-colonizing strains. Earlier, we determined that the majority strain in XBB differed from the
majority strain in XBA at only two segregating sites. We designated the majority strain in XBB,
which carried the XBA-like alleles at these 16011 sites, "strain 2," and we called the minor strain
in XBB "strain 3." To calculate the frequency of strain 3, we calculated the median frequency of
SNPs at these 16011 sites (**Fig. 5d**).
849
Bifidobacterium longum. We identified 3667 sites in the core genome of *Bifidobacterium*
longum at which the major allele in XBA (frequency>0.8) was the minor allele in XBB
(frequency<0.2) in all pre-antibiotic, QP populations. At 1026 of these sites, the major allele in
XBB before antibiotics, which we called the XBB allele, had median frequency 0 in XBA and 1 in
XBB before antibiotics, suggesting that these sites distinguish the dominant strains in each
subject. (We suspect that the remaining sites distinguish a minor, co-colonizing strain, but our
sequencing data was insufficient to estimate the frequency of this strain.) We called the strain
carrying the XBA allele at these 1026 sites "strain 1," and we called the strain carrying the XBB
allele at these sites "strain 2." The XBB allele was present at a frequency of 1 after antibiotics in
XBA and 1 in XBB, suggesting that colonization of XBA by strain 2 was responsible for the
increase in the relative abundance of *B. longum* after antibiotics (**Fig. 5c** and **Extended Data**
Fig. 9c).
862
863 ***Bacteroides faecis***. We identified 9 sites in the core genome of *Bacteroides faecis* at which the
864 major allele in XBA (frequency>0.8) was the minor allele in XBB (frequency<0.2) in all pre-
865 antibiotic, QP populations. These 9 sites were widely dispersed across the genome, suggesting
866 that they represent distinct mutations rather than a single recombination event. One group of

four SNPs had correlated frequency trajectories, suggesting that they are carried on a single haplotype (SNP group 1). The remaining five SNPs also had correlated frequency trajectories, suggesting that they are carried on a second, distinct haplotype (SNP group 2). Based on the SNP frequency trajectories and the pigeonhole principle, we inferred that the XBA and XBB populations were comprised of three distinct strains: strain 1, which carried the initial XBA major allele at all 9 sites; strain 2, which carried the initial XBB major allele at all 9 sites; and strain 3, which carried the initial XBA major allele at the four sites in SNP group 1 and the initial XBB major allele at the five sites in SNP group 2. XBA was co-colonized by strains 1 and 3 before antibiotics, but strain 3 appears to have reached fixation after antibiotics. Strain 3 was also present at a median frequency of ~0.1 in the *B. faecis* population of XBB throughout the sampling period, so we were unable to determine whether the increase in abundance of *B. faecis* after antibiotics was driven by the re-emergence of strain 3 in XBA or colonization from XBB. Aside from the sites in SNP group 2, we identified zero sites in the core genome at which the major allele in XBA before antibiotics (frequency>0.8) became the minor allele after antibiotics (frequency<0.2), suggesting that there were no SNPs that reached fixation between the end of antibiotics at day 5 and the re-emergence of *B. faecis* around day 401.

Dynamics of Type VI Secretion Systems

Metagenomic sequencing reads were mapped using bowtie2[86] against a database of 104 Type VI effector, immunity, and structural genes encompassing three distinct T6SS subtypes from *Bacteroidales* species[69]. To prevent cross-mapping, we included only the final 400 bases of the C-terminal domains of the effector genes. To calculate the normalized abundance of each T6SS gene, we divided the number of reads mapping to the gene by the length of the gene and the total number of reads in the sample and multiplied this number by 10^9.

Acknowledgements

The authors thank Susan Mello and Robert Pesich for their assistance with study coordination and Ania Robaczewska for technical assistance with library preparation. We also thank Emily Ebel, Jaime Lopez, Saria McKeithen-Mead, Matthew Olm, Tadashi Fukami, and members of the Petrov, Huang, Good, and Relman labs for valuable discussions and comments on the manuscript. Sequencing support for this project was provided as part of the Chan Zuckerberg Biohub Microbiome Initiative, and additional samples were sequenced at the DNA Services Lab, Roy J. Carver Biotechnology Center, University of Illinois at Urbana-Champaign. K.S.X. has been supported by a James McDonnell Foundation Postdoctoral Fellowship in Understanding Dynamic and Multi-Scale Systems and a Jane Coffin Childs Memorial Fund Postdoctoral Fellowship. S.J.W. is supported by a Hertz Foundation Fellowship and Stanford Graduate Fellowship. The authors acknowledge funding from NIH/NIGMS R35-GM142685 (to B.D.R.); NIH/NIGMS R35-GM118165 (to D.A.P.); NSF EF-2125383 (to K.C.H.); Alfred P. Sloan Foundation grant FG-2021-15708 (to B.H.G.) and NIH/NIGMS R35-GM146949 (to B.H.G.); the Thomas C. and Joan M. Merigan Endowment at Stanford University (to D.A.R.); NIH/NIAID R21-AI168860 (to D.A.R.); and NIH/NIAID R01-AI147023 (to D.A.R.). D.A.P., K.C.H., and B.H.G. are Chan Zuckerberg Biohub Investigators.

Author contributions

K.S.X, D.A.G., D.A.P., K.C.H., B.H.G., and D.A.R. designed the research; K.S.X., D.A.G, A.B.P., F.B.Y., and N.F.N. performed the research; K.S.X., S.J.W., D.A.G., M.L.M., A.J.V., A.B.P., N.A.R., B.D.R, and B.H.G. analyzed the data; K.S.X., D.A.P., K.C.H., B.H.G., and D.A.R. wrote the paper; and all authors reviewed it before submission.

 Figures

Fig. 1 | Heterogeneous responses to antibiotic perturbation in a longitudinal household cohort.

a, Study design. Healthy, cohabiting adults collected weekly stool samples for 9 weeks before and after one member of each household took a 5-day course of ciprofloxacin. In 18 households, subjects collected occasional follow-up samples for up to two years after the initial 9-week sampling period. **b**, Cohabiting subjects share a range of gut microbial strains. Histograms show the distribution of the total relative abundance of populations with and without shared strains among cohabiting pairs at the initial sampling timepoint. **c**, The total abundance of shared strains is not correlated with the length of cohabitation. Points show the longest-cohabiting pair in each household. **d,** Heterogeneous responses to antibiotic perturbation. The antibiotic responses of each community can be classified as transient (blue) or lasting (red) based on the Jensen-Shannon divergence (JSD) of their species composition between the beginning and end of the main study. **e**, Examples of transient and lasting antibiotic responses, relative to a control subject. Left panels show the species-level community composition over time, while right panels show the JSD from the initial timepoint. Gold bars indicate the ciprofloxacin perturbation, while the gray shading illustrates the range of JSDs between non-cohabiting subjects (2.5 to 97.5 percentiles).

Fig. 2 | Species disruption, recovery, and colonization in a subject with lasting antibiotic responses.

a, Relative abundances of species in subject XDA (left) and their cohabiting partner XDB (right). Species are shown if they were present in XDA before antibiotics (median relative abundance >0.1%) or if they newly colonized XDA (**Methods**). Points at the first and last timepoints indicate whether the strains at that timepoint are shared (closed) or not shared (open) with any timepoint from the cohabiting subject (**Methods**); no point is shown if the species was not shared between

subjects, or if the sequencing depth was insufficient to determine whether strains were shared. Timepoints are outlined in black if a new strain was detected relative to the initial timepoint. **b-g,** Examples of species from XDA that (**b**) were maintained through ciprofloxacin perturbation; (**c**) were disrupted by ciprofloxacin and did not recover; (**d**) were disrupted by ciprofloxacin and experienced recovery of the resident strain; (**e**) were disrupted by ciprofloxacin and experienced recovery via the transmission of a new strain from the cohabiting partner; (**f**) transiently colonized after ciprofloxacin perturbation; and (**g**) were newly transmitted from the cohabiting partner after ciprofloxacin perturbation. Points indicate whether the strains at that timepoint are shared (closed) or not shared (open) with the strains at any timepoint in the cohabiting partner; points are not shown if the species was not shared between subjects, or if the sequencing depth was insufficient to determine whether strains were shared (**Methods**). The gold and gray boxes denote the ciprofloxacin perturbation and the period of follow-up sampling, respectively. The dashed line indicates the limit of detection of 10^{-5}.

Fig. 3 | Dynamics of species disruption and colonization in the month after antibiotic perturbation.

a, Number of species classified as maintained, disrupted, recovered, and colonized after ciprofloxacin in each antibiotic-taking subject. **b**, Enrichment and depletion of bacterial families among disrupted species (left) and new colonizers (right). Families with significant enrichment (red) or depletion (blue) are labeled (Bonferroni-corrected p<0.05). **c**, Timeline of species recovery and colonization after antibiotic perturbation. Colonization times are inferred by fitting relative abundance trajectories to a simple model of exponential growth with saturation (**Methods**). Strains and species are annotated as transient colonizers if they no longer had relative abundance >10^{-3} one month after antibiotics. Species that were present before antibiotics but had insufficient coverage after antibiotics to determine strain-level composition are not shown. The gold bar shows the timing of the ciprofloxacin perturbation. **d**, Relationship between strain sharing and species recovery for all antibiotic taking subjects. Strain sharing between cohabiting subjects does not increase the likelihood of species recovery after ciprofloxacin perturbation. **e**, Prevalence of resident species and new colonizers in the Human Microbiome Project.

Fig. 4 | Colonization is elevated in subjects with lasting antibiotic responses, but only after long delays.

a, Timeline of species recovery and colonization in subjects with lasting antibiotic responses. Colonization times are inferred by fitting relative abundance trajectories to a simple model of exponential growth with saturation (see panel b; **Methods**). Strains and species are annotated as transient colonizers if they no longer had relative abundance >10^{-3} at the end of the study. Timepoints with statistically significant cohorts of colonization events are annotated with an asterisk (permutation test, $p<0.05$). Species that were present before antibiotics but had insufficient coverage after antibiotics to determine strain-level composition are not shown. Due to space constraints, timepoints with no new colonization are subsampled to once per month for subject XBA. **b**, The earliest introduction time, transit time, and colonization time of each new colonizer can be inferred from relative abundance trajectories (**Methods**). The gold bar shows the timing of the ciprofloxacin perturbation. These two cases depict the colonization of a new strain of species that was disrupted by antibiotics. **c**, Colonization is elevated in subjects with lasting antibiotic responses in the two years after ciprofloxacin perturbation. Lines show the total relative abundance of newly colonizing strains and species in each subject over time. **d**, New colonizers rapidly reach carrying capacity, even when they colonize long after antibiotics. Histograms show the number of intervals required for each new strain or species to increase from undetectable relative abundances to its inferred carrying capacity. **e**, Strains that are shared with a cohabiting partner can fail to recolonize after antibiotics, even as new strains are acquired from the cohabiting partner. The number of shared strains lost and gained after antibiotic perturbation is shown for each subject with lasting antibiotic responses and in aggregate for all other antibiotic-taking subjects.

Fig. 5 | Rapid transitions between alternative states in a subject with a lasting antibiotic response.

a, Subject XBA undergoes rapid shifts in community composition around days 0 and 401. **b**, Number of mutations distinguishing strains of *B. uniformis, P. vulgatus,* and *B. longum* subject XBA and their cohabiting partner XBB (**Methods**). Strain relative abundances in (**c,d**) are calculated based on the frequencies of these distinguishing SNPs and are shown in corresponding colors. **c**, *Bifidobacterium longum* transmits from XBB to XBA after antibiotics. **d**, Strains of *Bacteroides uniformis* and *Phocaeicola vulgatus* present before antibiotics re-emerge around day 401. Triangles in (**c,d**) indicate timing of the transition from state 2 to 3. **e**, Type VI secretion system (T6SS) effector genes undergo sharp shifts in abundance that correlate with shifts in community composition.

1013 **Extended Data Figures**

1014

Extended Data Figure 1 | Subject demographics and sampling timelines

Subject demographics, including **a**, age at the beginning of the study and self-reported **b**,
gender **c**, race, and **d**, ethnicity according to US Census Bureau categories. **e**, Sampling
timelines show each sample collected during the study and the target depth to which each
sample was sequenced. The second letter of each subject identifier represents the subject's
household; for instance, subjects XAA, XAC, and XAD live in household A.
1021

Extended Data Figure 2 | Identifying strains shared between cohabiting subjects.

a, Number of fixed differences between population pairs for three representative species. Using the "fixed differences" method, we classified two populations from cohabiting subjects as sharing strains if they had fewer fixed differences than 99% of population pairs from non-cohabiting subjects, as shown by the dashed line (**Methods**). Population pairs from the same subject are shown as a control. **b**, Distribution of genetic distances used as a threshold to annotate strain sharing for each species using the "fixed differences" method. **c**, Percent of private SNPs detected in population pairs for three representative species. Using this "strain fishing" method, we classified two populations as sharing strains if we detected >75% of private marker SNPs from one population in the other population, as shown by the dashed line (**Methods**). **d**, Percent of non-cohabiting population pairs for each species that share >75% of private marker SNPs. **e**, The fixed differences and strain fishing methods agreed on >90% of population pairs that could be analyzed using both methods. **f**, Total abundance of species and strain sharing in pairs of cohabiting and non-cohabiting subjects at the beginning of the study. **g**, Gut microbiome composition at the beginning of the study, with each species colored based on strain and species sharing between cohabiting subjects. In subject pair XLA-XLB, for example,

1038   the gut microbiome of subject XLA is shown and colored based on species and strain sharing
1039   with subject XLB.
1040

Extended Data Figure 3 | Community responses to antibiotic perturbation.

a, Antibiotic perturbation increases the compositional variability of the microbiome. Points show the compositional variability of each community calculated at days -28, 0, 7, and 35 using normalized F_{ST}[87]. **b**, Antibiotic perturbation causes mostly transient declines in community diversity. Points show the effective number of species calculated from the Shannon diversity

index before antibiotics, after antibiotics, and one month post-antibiotics. **c**, Maximum and final family-level Jensen-Shannon divergence of each community from the initial timepoint during the main study. **d**, Change in CFUs from the initial timepoint. **e**, Most recent antibiotic usage (self-reported) for each subject, colored by subject responses to ciprofloxacin perturbation. **f**, Subjects with transient and lasting responses have similar levels of pre-antibiotic diversity, as quantified by the effective number of species calculated from the Shannon diversity index. **g**, Subjects with transient and lasting responses have similar proportions of the gut microbiome that are comprised of shared strains. **h**, Species-level community composition over time for each antibiotic-taking subject. Subjects are arranged in order of increasing maximum Jensen-Shannon divergence from the initial timepoint.

Extended Data Figure 4 | Species dynamics in subject XAA

a, Species-level community composition over time in subject XAA, a subject with lasting antibiotic responses, and two cohabiting controls, subjects XAC and XAD. **b**, Analogous version of **Fig. 2a** for subjects XAA, XAC, and XAD. Species are shown if they were present in XAA before antibiotics (median relative abundance >0.1%) or if they newly colonized XAA (**Methods**). Points at the first and last timepoints indicate whether the strains at that timepoint are shared (closed) or not shared (open) with any timepoint from the cohabiting subject (**Methods**); no point is shown if the species was not shared between subjects, or if the sequencing depth was insufficient to determine whether strains were shared. Timepoints are outlined in black if a new strain was detected relative to the initial timepoint.

Extended Data Figure 5 | Species dynamics in subject XBA

Analogous version of **Extended Data Fig. 4** for subject XBA. **a**, Species-level community composition over time in subject XBA, a subject with lasting antibiotic responses, and a cohabiting control, subject XBB. **b**, Relative abundances of species in subjects XBA and XBB. Species are shown if they were present in XBA before antibiotics (median relative abundance >0.1%) or if they newly colonized XBA (**Methods**). Points at the first and last timepoints indicate whether the strains at that timepoint are shared (closed) or not shared (open) with any timepoint from the cohabiting subject (**Methods**); no point is shown if the species was not shared between subjects, or if the sequencing depth was insufficient to determine whether strains were shared. Timepoints are outlined in black if a new strain was detected relative to the initial timepoint.

1076

Extended Data Figure 6 | Species dynamics in subject XEA

Analogous version of **Extended Data Fig. 4** for subject XEA. **a**, Species-level community composition over time in subject XEA, a subject with lasting antibiotic responses, and a cohabiting control, subject XEB. **b**, Relative abundances of species in subjects XEA and XEB. Species are shown if they were present in XEA before antibiotics (median relative abundance >0.1%) or if they newly colonized XEA (**Methods**). Points at the first and last timepoints indicate whether the strains at that timepoint are shared (closed) or not shared (open) with any timepoint from the cohabiting subject (**Methods**); no point is shown if the species was not shared between subjects, or if the sequencing depth was insufficient to determine whether strains were shared. Timepoints are outlined in black if a new strain was detected relative to the initial timepoint.

1077

1088

Extended Data Figure 7 | Species dynamics in subject XKA

Analogous version of **Extended Data Fig. 4** for subject XKA. **a**, Species-level community composition over time in subject XKA, a subject with lasting antibiotic responses, and a cohabiting control, subject XKB. **b**, Relative abundances of species in subjects XKA and XKB. Species are shown if they were present in XKA before antibiotics (median relative abundance >0.1%) or if they newly colonized XKA (**Methods**). Points at the first and last timepoints indicate whether the strains at that timepoint are shared (closed) or not shared (open) with any timepoint from the cohabiting subject (**Methods**); no point is shown if the species was not shared between subjects, or if the sequencing depth was insufficient to determine whether strains were shared. Timepoints are outlined in black if a new strain was detected relative to the initial timepoint.

1099

Extended Data Figure 8 | Species disruption, recovery, and colonization in the first month after ciprofloxacin perturbation.

a, Number of species classified as maintained, disrupted, recovered, and colonized after ciprofloxacin in each control subject. **b**, Total abundance of species classified as maintained, disrupted, recovered, and colonized in each subject at the initial (day -28) and final (day 35)

timepoints in the main study. **c**, Total abundance of bacterial families that are significantly less
(left) or more (right) likely to be disrupted in each antibiotic-taking subject at the initial timepoint.
d, Enrichment and depletion of bacterial families among disrupted species that fail to recover.
Families with significant enrichment (red) or depletion (blue) are labeled (Bonferroni-corrected
p<0.05). **e**, Total abundance of bacterial families that are significantly less (left) or more (right)
likely to be disrupted without recovery in each antibiotic-taking subject at the initial timepoint. **f**,
Prevotella copri colonizes subject XIA (blue) during follow-up sampling. Points indicate whether
the strains at that timepoint are shared (closed) or not shared (open) with the strains at any
timepoint in the cohabiting partner; points are not shown if the species was not shared between
subjects, or if the sequencing depth was insufficient to determine whether strains were shared
(**Methods**). The gold and gray boxes denote the ciprofloxacin perturbation and the period of
follow-up sampling, respectively. The dashed line indicates the limit of detection of 10^{-5}. **g**,
Relationship between strain sharing and species recovery for subjects with transient antibiotic
responses. Strain sharing between cohabiting subjects does not increase the likelihood of
species recovery after ciprofloxacin perturbation. **h**, Number of colonizing strains and species in
each subject during the first month after ciprofloxacin perturbation. **i**, Total relative abundance of
colonizing strains and species in each subject during the first month after ciprofloxacin
perturbation. **j**, Median abundance of resident species and new colonizers in the Human
Microbiome Project. **k**, *Bacteroides stercoris* transmits from subject XKB (gray) to subject XKA
(red) after antibiotic perturbation.
1125
1126

1127

Extended Data Figure 9 | Colonization is elevated over longer timescales in subjects with lasting antibiotic responses.

a, Number of strains and species that colonized each subject more than one month after the end of the ciprofloxacin course. **b**, Total abundance of all new colonizers in each subject at the end of the study. **c**, Total abundance of new colonizers in each subject with lasting antibiotic responses and their cohabiting partners at the end of the study. **d**, Distribution of transit times (the number of days between when a colonizing strain or species was last undetectable and when it reaches steady-state relative abundance; **Methods**) across all new colonizers. Transit times were rounded up to the full time interval, regardless of the number of days between samples. Many species increased from undetectable to steady-state relative abundances within a single time interval and could be consistent with transit times as short as 1 day. **e**, Colonization events sometimes occur in coordinated cohorts. Crosses show the maximum number of colonization events detected at a single follow-up sampling timepoint for each subject with lasting antibiotic responses. Colonization times are inferred by fitting relative abundance trajectories to a simple model of exponential growth (**Methods**). The gray distributions show the expected distribution if colonization were distributed uniformly over time (**Methods**). **f**, Two new strains of *Phocaeicola vulgatus* are detected simultaneously in subject XAA after antibiotic perturbation (**Methods**). Left panel shows the relative abundances of pre- and post-antibiotic strains (colored regions), while the right panel shows the genetic distance between them. The gold bar shows the timing of the ciprofloxacin perturbation.

1148

1149

Extended Data Figure 10 | Strain and species dynamics in subject XBA.

a, Subject XBA undergoes rapid switches in community states. Relative abundances are shown for *Eggerthella lenta, Bifidobacterium longum,* and all *Bacteroidaceae* species detected in subject XBA. Strain sharing is annotated as in **Fig. 2** (**Methods**). **b**, Several *Bacteroidaceae* species undergo sudden, simultaneous increases in relative abundance around day 400. **c,d**, Lines show the median abundance of SNPs that distinguish the initial strains of *Bifidobacterium longum* (**c**) and *Bacteroides uniformis* (**d**) in XBA and XBB. Shaded regions show the 25 and 75 percentiles of these SNP frequencies. **e,f**, Lines show the frequencies of the two SNPs that distinguish the initial strains of *Phocaeicola vulgatus* in XBA and XBB at the initial timepoint (**e**) and the seven SNPs that fix in *Phocaeicola vulgatus* in XBA after the antibiotic perturbation (**f**). **g**, The relative abundance of *Bacteroides fragilis* is strongly correlated with the abundance of the T6SS GA3-E13 effector gene in subject XBA. **h,i**, Subjects XAA (**h**) and XKA (**i**) undergo major shifts in the abundance of T6SS genes after antibiotic perturbation.

References

1. Robinson, C. D., Bohannan, B. J. & Britton, R. A. Scales of persistence: transmission and the microbiome. *Curr. Opin. Microbiol.* **50**, 42–49 (2019).
2. Miller, E. T., Svanbäck, R. & Bohannan, B. J. M. Microbiomes as Metacommunities: Understanding Host-Associated Microbes through Metacommunity Ecology. *Trends Ecol. Evol.* **33**, 926–935 (2018).
3. Walter, J., Maldonado-Gómez, M. X. & Martínez, I. To engraft or not to engraft: an ecological framework for gut microbiome modulation with live microbes. *Curr. Opin. Biotechnol.* **49**, 129–139 (2018).
4. Zmora, N. *et al.* Personalized Gut Mucosal Colonization Resistance to Empiric Probiotics Is Associated with Unique Host and Microbiome Features. *Cell* (2018) doi:10.1016/j.cell.2018.08.041.
5. Bezkorovainy, A. Probiotics: determinants of survival and growth in the gut. *Am. J. Clin. Nutr.* **73**, 399S-405S (2001).
6. Vellend, M. *The Theory of Ecological Communities (MPB-57)*. *The Theory of Ecological Communities (MPB-57)* (2016). doi:10.1515/9781400883790.
7. Skellam, J. G. Random Dispersal in Theoretical Populations. *Biometrika* **38**, 196–218 (1951).
8. Stephens, P. A., Sutherland, W. J. & Freckleton, R. P. What Is the Allee Effect? *Oikos* **87**, 185–190 (1999).
9. Lewis, M. A. & Kareiva, P. Allee Dynamics and the Spread of Invading Organisms. *Theor. Popul. Biol.* **43**, 141–158 (1993).
10. Taylor, C. M. & Hastings, A. Allee effects in biological invasions. *Ecol. Lett.* **8**, 895–908 (2005).
11. Dini-Andreote, F. & Custer, G. F. Ecological principles of fecal microbiota transplantation. *Trends Microbiol.* **31**, 776–779 (2023).
12. Valles-Colomer, M. *et al.* The person-to-person transmission landscape of the gut and oral microbiomes. *Nature* **614**, 125–135 (2023).
13. Korpela, K. *et al.* Selective maternal seeding and environment shape the human gut microbiome. *Genome Res.* **28**, 561–568 (2018).
14. Brito, I. L. *et al.* Transmission of human-associated microbiota along family and social networks. *Nat. Microbiol.* **4**, 964–971 (2019).
15. Suez, J., Zmora, N., Segal, E. & Elinav, E. The pros, cons, and many unknowns of probiotics. *Nat. Med.* **25**, 716–729 (2019).
16. Smillie, C. S. *et al.* Strain Tracking Reveals the Determinants of Bacterial Engraftment in the Human Gut Following Fecal Microbiota Transplantation. *Cell Host Microbe* (2018) doi:10.1016/j.chom.2018.01.003.
17. Aggarwala, V. *et al.* Precise quantification of bacterial strains after fecal microbiota transplantation delineates long-term engraftment and explains outcomes. *Nat. Microbiol.* **6**, 1309–1318 (2021).
18. Faith, J. J. *et al.* The Long-Term Stability of the Human Gut Microbiota. *Science* **341**, 1237439 (2013).
19. Rothschild, D. *et al.* Environment dominates over host genetics in shaping human gut microbiota. *Nature* **555**, 210–215 (2018).
20. Siranosian, B. A. *et al.* Rare transmission of commensal and pathogenic bacteria in the gut microbiome of hospitalized adults. *Nat. Commun. 2022 131* **13**, 1–17 (2022).
21. Buffie, C. G. & Pamer, E. G. Microbiota-mediated colonization resistance against intestinal pathogens. *Nat. Rev. Immunol.* **13**, 790–801 (2013).
22. Lawley, T. D. & Walker, A. W. Intestinal colonization resistance. *Immunology* **138**, 1–11 (2013).

23. Fishbein, S. R. S., Mahmud, B. & Dantas, G. Antibiotic perturbations to the gut microbiome.
Nat. Rev. Microbiol. 1–17 (2023) doi:10.1038/s41579-023-00933-y.
24. Maldonado-Gómez, M. X. *et al.* Stable Engraftment of Bifidobacterium longum AH1206 in
the Human Gut Depends on Individualized Features of the Resident Microbiome. *Cell Host*
Microbe (2016) doi:10.1016/j.chom.2016.09.001.
25. Conwill, A. *et al.* Anatomy promotes neutral coexistence of strains in the human skin
microbiome. *Cell Host Microbe* **30**, 171-182.e7 (2022).
26. Cheng, A. G. *et al.* Design, construction, and in vivo augmentation of a complex gut
microbiome. *Cell* **185**, 3617-3636.e19 (2022).
27. Segura Munoz, R. R. *et al.* Experimental evaluation of ecological principles to understand
and modulate the outcome of bacterial strain competition in gut microbiomes. *ISME J.* **16**,
1594–1604 (2022).
28. Shepherd, E. S., Deloache, W. C., Pruss, K. M., Whitaker, W. R. & Sonnenburg, J. L. An
exclusive metabolic niche enables strain engraftment in the gut microbiota. *Nature* (2018)
doi:10.1038/s41586-018-0092-4.
29. Lee, S. M. *et al.* Bacterial colonization factors control specificity and stability of the gut
microbiota. *Nature* (2013) doi:10.1038/nature12447.
30. Buffie, C. G. *et al.* Precision microbiome reconstitution restores bile acid mediated
resistance to Clostridium difficile. *Nature* (2015) doi:10.1038/nature13828.
31. Whitaker, W. R., Shepherd, E. S. & Sonnenburg, J. L. Tunable Expression Tools Enable
Single-Cell Strain Distinction in the Gut Microbiome. *Cell* **169**, 538-546.e12 (2017).
32. Custer, G. F., Bresciani, L. & Dini-Andreote, F. Ecological and Evolutionary Implications of
Microbial Dispersal. *Front. Microbiol.* **13**, (2022).
33. Fukami, T. Historical Contingency in Community Assembly: Integrating Niches, Species
Pools, and Priority Effects. *Annu. Rev. Ecol. Evol. Syst.* (2015) doi:10.1146/annurev-
ecolsys-110411-160340.
34. de Nies, L., Kobras, C. M. & Stracy, M. Antibiotic-induced collateral damage to the
microbiota and associated infections. *Nat. Rev. Microbiol.* 1–16 (2023) doi:10.1038/s41579-
023-00936-9.
35. Schwartz, D. J., Langdon, A. E. & Dantas, G. Understanding the impact of antibiotic
perturbation on the human microbiome. *Genome Med.* **12**, 82 (2020).
36. Bohnhoff, M. & Miller, C. P. Enhanced susceptibility to salmonella infection in streptomycin-
treated mice. *J. Infect. Dis.* (1962) doi:10.1093/infdis/111.2.117.
37. Frazão, N., Sousa, A., Lässig, M. & Gordo, I. Horizontal gene transfer overrides mutation in
Escherichia coli colonizing the mammalian gut. *Proc. Natl. Acad. Sci.* **116**, 17906–17915
(2019).
38. Ianiro, G. *et al.* Variability of strain engraftment and predictability of microbiome composition
after fecal microbiota transplantation across different diseases. *Nat. Med.* **28**, 1913–1923
(2022).
39. Lozupone, C. A., Stombaugh, J. I., Gordon, J. I., Jansson, J. K. & Knight, R. Diversity,
stability and resilience of the human gut microbiota. *Nature* **489**, 220–230 (2012).
40. Fassarella, M. *et al.* Gut microbiome stability and resilience: elucidating the response to
perturbations in order to modulate gut health. *Gut* **70**, 595–605 (2021).
41. Roodgar, M. *et al.* Longitudinal linked-read sequencing reveals ecological and evolutionary
responses of a human gut microbiome during antibiotic treatment. *Genome Res.* **31**, 1433–
1446 (2021).
42. Ng, K. M. *et al.* Recovery of the Gut Microbiota after Antibiotics Depends on Host Diet,
Community Context, and Environmental Reservoirs. *Cell Host Microbe* **26**, 650-665.e4
(2019).

43. Nayfach, S., Rodriguez-Mueller, B., Garud, N. & Pollard, K. S. An integrated metagenomics
pipeline for strain profiling reveals novel patterns of bacterial transmission and
biogeography. *Genome Res.* **26**, 1612–1625 (2016).
44. Garud, N. R., Good, B. H., Hallatschek, O. & Pollard, K. S. Evolutionary dynamics of
bacteria in the gut microbiome within and across hosts. *PLOS Biol.* **17**, e3000102 (2019).
45. Dethlefsen, L., Huse, S., Sogin, M. L. & Relman, D. A. The pervasive effects of an antibiotic
on the human gut microbiota, as revealed by deep 16s rRNA sequencing. *PLoS Biol.* (2008)
doi:10.1371/journal.pbio.0060280.
46. Dethlefsen, L. & Relman, D. A. D. A. Incomplete recovery and individualized responses of
the human distal gut microbiota to repeated antibiotic perturbation. *Proc. Natl. Acad. Sci.*
(2010) doi:10.1073/pnas.1000087107.
47. Sonnenburg, E. D. & Sonnenburg, J. L. The ancestral and industrialized gut microbiota and
implications for human health. *Nat. Rev. Microbiol.* **17**, 383–390 (2019).
48. de Nies, L., Kobras, C. M. & Stracy, M. Antibiotic-induced collateral damage to the
microbiota and associated infections. *Nat. Rev. Microbiol.* 1–16 (2023) doi:10.1038/s41579-
023-00936-9.
49. David, L. A. *et al.* Host lifestyle affects human microbiota on daily timescales. *Genome Biol.*
15, R89 (2014).
50. Zhao, S. *et al.* Adaptive Evolution within Gut Microbiomes of Healthy People. *Cell Host*
Microbe **25**, 656-667.e8 (2019).
51. Dewhirst, F. E. *et al.* The Human Oral Microbiome. *J. Bacteriol.* **192**, 5002–5017 (2010).
52. Deo, P. N. & Deshmukh, R. Oral microbiome: Unveiling the fundamentals. *J. Oral Maxillofac.*
Pathol. **23**, 122 (2019).
53. Liu, F., Ma, R., Wang, Y. & Zhang, L. The Clinical Importance of Campylobacter concisus
and Other Human Hosted Campylobacter Species. *Front. Cell. Infect. Microbiol.* **8**, (2018).
54. Zhou, P., Manoil, D., Belibasakis, G. N. & Kotsakis, G. A. Veillonellae: Beyond Bridging
Species in Oral Biofilm Ecology. *Front. Oral Health* **2**, 774115 (2021).
55. Schmidt, T. S. *et al.* Extensive transmission of microbes along the gastrointestinal tract.
eLife **8**, e42693 (2019).
56. Liao, C. *et al.* Relative enrichment of oral bacteria in feces indicates loss of intestinal
commensals with implications for host health. 2022.10.24.513595 Preprint at
https://doi.org/10.1101/2022.10.24.513595 (2022).
57. Sorbara, M. T. & Pamer, E. G. Interbacterial mechanisms of colonization resistance and the
strategies pathogens use to overcome them. *Mucosal Immunol.* **12**, 1–9 (2019).
58. Posfai, A., Taillefumier, T. & Wingreen, N. S. Metabolic Trade-Offs Promote Diversity in a
Model Ecosystem. *Phys. Rev. Lett.* **118**, 028103 (2017).
59. Newton, D. P., Ho, P.-Y. & Huang, K. C. Modulation of antibiotic effects on microbial
communities by resource competition. *Nat. Commun.* **14**, 2398 (2023).
60. Crooks, J. & Soulé, M. Lag Times in Population Explosions of Invasive Species: Causes and
Implications. in 103–125 (1999). doi:10.1007/978-94-011-4523-7_7.
61. Crooks, J. A. Lag times and exotic species: The ecology and management of biological
invasions in slow-motion. *Écoscience* **12**, 316–329 (2005).
62. Sender, R., Fuchs, S. & Milo, R. Revised Estimates for the Number of Human and Bacteria
Cells in the Body. *PLoS Biol.* (2016) doi:10.1371/journal.pbio.1002533.
63. Wu, G., Zhao, N., Zhang, C., Lam, Y. Y. & Zhao, L. Guild-based analysis for understanding
gut microbiome in human health and diseases. *Genome Med.* **13**, 22 (2021).
64. Simberloff, D. & Von Holle, B. Positive Interactions of Nonindigenous Species: Invasional
Meltdown? *Biol. Invasions* **1**, 21–32 (1999).
65. Fukami, T. & Nakajima, M. Community assembly: alternative stable states or alternative
transient states? *Ecol. Lett.* **14**, 973–984 (2011).

66. Beisner, B. E., Haydon, D. T. & Cuddington, K. Alternative Stable States in Ecology. *Front. Ecol. Environ.* **1**, 376–382 (2003).

67. Rios-Covián, D. *et al.* A proteomic approach towards understanding the cross talk between Bacteroides fragilis and Bifidobacterium longum in coculture. *Can. J. Microbiol.* **62**, 623–628 (2016).

68. Odamaki, T. *et al.* Effect of the oral intake of yogurt containing Bifidobacterium longum BB536 on the cell numbers of enterotoxigenic Bacteroides fragilis in microbiota. *Anaerobe* **18**, 14–18 (2012).

69. Verster, A. J. *et al.* The Landscape of Type VI Secretion across Human Gut Microbiomes Reveals Its Role in Community Composition. *Cell Host Microbe* (2017) doi:10.1016/j.chom.2017.08.010.

70. Coyne, M. J., Roelofs, K. G. & Comstock, L. E. Type VI secretion systems of human gut Bacteroidales segregate into three genetic architectures, two of which are contained on mobile genetic elements. *BMC Genomics* **17**, 58 (2016).

71. Chatzidaki-Livanis, M., Geva-Zatorsky, N. & Comstock, L. E. Bacteroides fragilis type VI secretion systems use novel effector and immunity proteins to antagonize human gut Bacteroidales species. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 3627–3632 (2016).

72. Hecht, A. L. *et al.* Strain competition restricts colonization of an enteric pathogen and prevents colitis. *EMBO Rep.* **17**, 1281–1291 (2016).

73. Russell, A. B. *et al.* A Type VI Secretion-Related Pathway in Bacteroidetes Mediates Interbacterial Antagonism. *Cell Host Microbe* **16**, 227–236 (2014).

74. Rillig, M. C. *et al.* Interchange of entire communities: microbial community coalescence. *Trends Ecol. Evol.* **30**, 470–476 (2015).

75. Debray, R. *et al.* Priority effects in microbiome assembly. *Nat. Rev. Microbiol.* **20**, 109–121 (2022).

76. Granato, E. T., Meiller-Legrand, T. A. & Foster, K. R. The Evolution and Ecology of Bacterial Warfare. *Curr. Biol.* **29**, R521–R537 (2019).

77. Oliveira, R. A., Cabral, V., Torcato, I. & Xavier, K. B. Deciphering the quorum-sensing lexicon of the gut microbiota. *Cell Host Microbe* **31**, 500–512 (2023).

78. Culp, E. J. & Goodman, A. L. Cross-feeding in the gut microbiome: Ecology and mechanisms. *Cell Host Microbe* **31**, 485–499 (2023).

79. Rakoff-Nahoum, S., Foster, K. R. & Comstock, L. E. The evolution of cooperation within the gut microbiota. *Nature* **533**, 255–259 (2016).

80. Zhou, W., Chow, K., Fleming, E. & Oh, J. Selective colonization ability of human fecal microbes in different mouse gut environments. *ISME J.* **13**, 805–823 (2019).

81. Wu, L. *et al.* Data-driven prediction of colonization outcomes for complex microbial communities. 2023.04.19.537502 Preprint at https://doi.org/10.1101/2023.04.19.537502 (2023).

82. Vasquez, K. S. *et al.* Quantifying rapid bacterial evolution and transmission within the mouse intestine. *Cell Host Microbe* **29**, 1454-1468.e4 (2021).

83. Shalon, D. *et al.* Profiling the human intestinal environment under physiological conditions. *Nature* **617**, 581–591 (2023).

84. Lou, Y. C. *et al.* Using strain-resolved analysis to identify contamination in metagenomics data. *Microbiome* **11**, 36 (2023).

85. Jiang, H., Lei, R., Ding, S.-W. & Zhu, S. Skewer: a fast and accurate adapter trimmer for next-generation sequencing paired-end reads. *BMC Bioinformatics* **15**, 182 (2014).

86. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **9**, 357–359 (2012).

87. Morrison, M. L., Alcala, N. & Rosenberg, N. A. FSTruct: An FST-based tool for measuring ancestry variation in inference of population structure. *Mol. Ecol. Resour.* **22**, 2614–2626 (2022).

88. Olm, M. R. *et al.* inStrain profiles population microdiversity from metagenomic data and
sensitively detects shared microbial strains. *Nat. Biotechnol.* (2021) doi:10.1038/s41587-
020-00797-0.
89. Virtanen, P. *et al.* SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nat.*
Methods **17**, 261–272 (2020).
90. Ghosh, O. M. & Good, B. H. Emergent evolutionary forces in spatial models of luminal
growth and their application to the human gut microbiota. *Proc. Natl. Acad. Sci.* **119**,
e2114931119 (2022).
91. Good, B. H. & Rosenfeld, L. B. Eco-evolutionary feedbacks in the human gut microbiome.
2022.01.26.477953 Preprint at https://doi.org/10.1101/2022.01.26.477953 (2023).
92. Huttenhower, C. *et al.* Structure, function and diversity of the healthy human microbiome.
Nature **486**, 207–214 (2012).
1375