CRISPR-Cas inhibits plasmid transfer and immunizes bacteria against antibiotic

resistance acquisition in manure

Chahat Upreti[1], Pranav Kumar[1], Lisa M. Durso[2*], and Kelli L. Palmer[1*]

[1] Department of Biological Sciences, The University of Texas at Dallas, Richardson, Texas

[2] U.S. Department of Agriculture, Agricultural Research Service, Agroecosystem Management

Unit, 251 Filley Hall, UNL East Campus, Lincoln, Nebraska

*Co-corresponding authors

Dr. Lisa Durso: lisa.durso@usda.gov

Dr. Kelli Palmer: kelli.palmer@utdallas.edu

Abstract

The horizontal transfer of antibiotic resistance genes among bacteria is a pressing global issue.

The bacterial defense system CRISPR-Cas acts as a barrier to the spread of antibiotic

resistance plasmids, and CRISPR-Cas-based antimicrobials can be effective to selectively

deplete antibiotic-resistant bacteria. While significant surveillance efforts monitor the spread of

antibiotic-resistant bacteria in the clinical context, a major, often overlooked aspect of the issue

is resistance emergence in agriculture. Farm animals are commonly treated with antibiotics, and

antibiotic resistance in agriculture is on the rise. Yet, CRISPR-Cas efficacy has not been

investigated in this setting. Here, we evaluate the prevalence of CRISPR-Cas in agricultural

Enterococcus faecalis strains and its anti-plasmid efficacy in an agricultural niche – manure. We

show that the prevalence of CRISPR-Cas subtypes is similar between clinical and agricultural *E.*

faecalis strains. CRISPR-Cas was found to be an effective barrier against resistance plasmid

transfer in manure, with improved effect as time progressed. CRISPR-based antimicrobials to

cure resistant *E. faecalis* of erythromycin resistance was limited by delivery efficiency of the

CRISPR antimicrobial in manure. However, immunization of bacteria against resistance gene

acquisition in manure was highly effective. Together, our results show that *E. faecalis* CRISPR-

Cas is prevalent and effective in an agricultural setting, and has the potential to be utilized for

depleting antibiotic-resistant populations. Our work has broad implications for tackling antibiotic

resistance in the increasingly relevant agricultural setting, in line with a OneHealth approach.

Importance

Antibiotic resistance is a growing global health crisis in human and veterinary medicine.

Previous work has shown technologies based on CRISPR-Cas - a bacterial defense system - to

be effective in tackling antibiotic resistance. Here we test if CRISPR-Cas is present and

effective in agricultural niches, specifically in the ubiquitously present bacterium – *Enterococcus*

faecalis. We show that CRISPR-Cas is prevalent, functional in manure, and has the potential to

be used to specifically kill bacteria carrying antibiotic resistance genes. This study demonstrates

the utility of CRISPR-Cas based strategies for control of antibiotic resistance in agricultural

settings.

1. Introduction

A widely prevalent adaptive immunity system possessed by bacteria and archaea is Clustered

Regularly Interspaced Short Palindromic Repeats (CRISPR)-Cas [1,2]. Although it has now been

programmed as a tool to edit genomes of a large number of organisms[3–6], in its native

environment, it confers bacteria and archaea defense against mobile genetic elements (MGE)

like viruses and plasmids[1,2,7,8]. Upon entry of the MGE, a portion of its DNA, known as a

protospacer, is integrated into the host genome in the form of a spacer[8]. Later, the spacer is

transcribed (called crRNA) and loaded on Cas-encoded nuclease(s) to surveille for future MGE

entries. If a region of an incoming MGE is complementary to the crRNA, the MGE is cleaved by

the nuclease[9–11]. Since its discovery, a large majority of studies on the CRISPR-Cas mechanism

of action have focused on its antiviral activity. However approx. 10% of CRISPR protospacers

identified in bacteria and archaea target plasmids, suggesting that CRISPR-Cas naturally

provides defense against invading plasmids as well[12]. Indeed, CRISPR-Cas has been shown to

act as a natural barrier against conjugative plasmid transfer[13,14].

A highly relevant clinical organism in the study of CRISPR-Cas function is *Enterococcus*

faecalis. E. faecalis is a Gram-positive bacterium routinely isolated from a large range of hosts

including animals, humans and environmental samples such as plants, sand, and water[15,16,17]. It

is an opportunistic pathogen and causes diseases including urinary tract infections, infective

endocarditis and bacteremia[18–21]. Owing in part to its significant repertoire of intrinsic antibiotic

resistance genes and its ability to efficiently acquire resistance genes from external sources, *E.*

faecalis is one of the leading causes of hospital-acquired infections[22,23]. Since *E. faecalis*

colonizes several food animals, strains from these sources can also carry both antibiotic

resistance and virulence genes [24,25]. Further, due to the generalist nature of *E. faecalis*,

transmission of resistant *E. faecalis* from animals to humans (zoonosis) can occur. Indeed,

there have been several cases of probable zoonotic transmission leading to major outbreaks in

the past[26,27], and enterococci, including *E. faecalis*, are routinely monitored in retail meat as part

of the National Antimicrobial Resistance Monitoring System[28].

Some *E. faecalis* strains encode Type II CRISPR-Cas systems which employ a crRNA-Cas9

nuclease-tracrRNA complex to surveille and cleave MGEs[29–31]. Since plasmid-mediated transfer

of antibiotic resistance genes is a major contributor to resistance transmission in *E. faecalis*[23,32],

it is important to test if CRISPR-Cas could act as a barrier against this process, especially in

conditions where zoonotic transmission may occur. Additionally, the prevalence and function of

CRISPR-Cas in strains of animal origin should be investigated, as most focus has been on

human-origin strains. Some recent studies have looked at *E. faecalis* samples isolated from

wastewater treatment plants, animal feces and soil. However, these studies have either small

sample sizes, low source diversity or unclear origin of strains[33–35].

Previous work in our laboratory has shown *E. faecalis* CRISPR-Cas systems confer defense

against plasmids and can be programmed as antimicrobials against specific antibiotic-resistant

strains, both *in vitro* and *in vivo* in the mouse gut [14,36]. During in vitro mating experiments,

CRISPR-Cas acted as a moderately effective barrier against plasmid transfer, while in vivo its

efficiency was strikingly robust[14], suggesting that the conjugation environment impacts CRISPR-

Cas function. Given the importance of agriculturally relevant niches in *E. faecalis* biology, it is

crucial to test how effective CRISPR-Cas is in this setting. An example of such a niche is

manure. Animal feces are a known reservoir of antibiotic resistance gene-encoding pathogenic

bacteria including *E. faecalis* and are thus potential hubs for zoonotic transmission[37–39], as well

as potential critical control points. Since previous work in *E. faecalis* has demonstrated

environmental impacts on CRISPR-Cas efficacy[14], discerning its efficacy in manure could be of

major significance.

In this study, we first identified the prevalence of CRISPR-Cas in a large cohort of agricultural *E.*

faecalis strains. Next, we evaluated CRISPR-Cas efficacy in sterilized manure using previously

established model *E. faecalis* strains. Finally, we tested the effectiveness of CRISPR-

antimicrobials designed to eliminate antibiotic resistance genes from *E. faecalis* in manure.

2. Results

CRISPR-Cas prevalence in agricultural *Enterococcus faecalis* is similar to that in human-

derived strains

Before testing the efficacy of CRISPR-Cas in a model agricultural setting, we wanted to

determine how prevalent CRISPR-Cas is in agriculturally-derived *E. faecalis* and how it

compares to its prevalence in strains derived from humans. For this, we downloaded all *E.*

faecalis genomes from NCBI Refseq available as of September 1, 2021, along with genomes

from a recent large-scale OneHealth investigation of Enterococci[25]. We segregated the

genomes based on source of isolation – human or animal (the term animal here is an umbrella

term for non-human sources like food animals, birds, pets etc. *Supplementary figure 1*).

Together, these comprised 1653 genomes sourced from humans (80.47%) and 401 sourced

from animals (19.53%). To identify CRISPR-Cas loci in these genomes, we performed a BLAST

search using the *cas9* sequences of the *E. faecalis* CRISPR1-Cas and CRISPR3-Cas loci as

queries, and then searched for CRISPR arrays within the set of genomes containing *cas9*. We

found that 27.57% of animal-derived strains possessed CRISPR1-Cas, and 9.52% strains

possessed CRISPR3-Cas. This was similar to their prevalence in human-derived strains –

29.36% for CRISPR1-Cas and 8.5% for CRISPR3-Cas (Figure 1). This suggests that CRISPR-

Cas prevalence is consistent across human and animal niches, with the caveat that our study is

limited to the *E. faecalis* strains that have been selected for sequencing by other groups, which

may have biases (i.e. prioritization of antibiotic-resistant strains for sequencing).

Figure 1

FIG 1 Relative abundances of CRISPR-Cas types in *E. faecalis* strains from animal and human

sources. Percentages were calculated as the number of strains in each source category

(animal/human) containing a particular CRISPR-Cas type (CRISPR1-Cas or CRISPR3-Cas)

divided by the total number of strains in that source category. p-values were calculated using

the Z test for equality of two percentages using independent samples.

CRISPR-Cas blocks plasmid transfer in manure, and manure-specific effects on CRISPR

defense are observed

We next tested the hypothesis that *E. faecalis* CRISPR-Cas is effective in an agricultural niche

(manure). We tested the efficacy of CRISPR-Cas in blocking the dissemination of pAM714 from

E. faecalis OG1SSp to *E. faecalis* T11RF (Fig 2). For this, we performed conjugation assays on

solid brain heart infusion (BHI) agar, in liquid BHI, and in liquified autoclaved dairy manure (Fig

3). Two different recipients were used, T11RF with a functional CRISPR-Cas system, and

T11RF*Δcas9* which lacks functional CRISPR-Cas defense against pAM714 (Table 1)[13,14]. For

148   these experiments, donor and recipient cells were mixed together and either spread on BHI

149   agar to form a biofilm or inoculated into liquid BHI or manure. After 18 hours incubation, agar

150   plate biofilms were harvested and resuspended in buffer, and liquid cultures were sampled for

151   determination of colony forming units (CFU) of donors, recipients, and transconjugant cells. This

152   timepoint is referred to as passage 0.

155   **Figure 2**

FIG 2 A schematic representation of the concept. The donor strain OG1SSp contains a

pheromone-responsive plasmid which harbors a *repB* gene (blue). The recipient strain T11RF

encodes a Type-II CRISPR-Cas containing a spacer with perfect complementarity to *repB* (red).

If CRISPR-Cas is effective in a particular niche, it will block the entry of the incoming plasmid. If

not, the plasmid enters the recipient, which is then termed a transconjugant.

Table 1 *E. faecalis* strains used in this study

Strain	Description	Reference(s)
T11RF	Rifampin-fusidic acid-resistant derivative of strain T11	40,41
T11RFΔcas9	T11RF with an in-frame deletion of cas9	40
OG1SSp(pAM714)	Spectinomycin-streptomycin-resistant derivative of strain OG1 harboring pAM714, conferring erythromycin resistance via Tn*917* insertion upstream of the par locus; *cyl*+	42,43
CK135RF	Spontaneous fusidic acid-resistant derivative of CK135	36

In passage 0, we found transconjugants in the range of 10^4–10^7 CFU/mL across the three

environments and the two recipients, suggesting widespread plasmid transfer across all 6

combinations (Fig 4). However, there was a significant difference in pAM714 conjugation

frequency (number of transconjugants divided by the number of donors) on BHI agar in the

absence vs presence of *cas9*, with a higher conjugation frequency in the absence of functional

CRISPR-Cas (Fig 5a). This was expected since it has been reported previously, demonstrating

that CRISPR-Cas is a moderately effective barrier against plasmid transfer on solid agar[14]. This

effect was also seen in liquid BHI medium, and importantly, also in manure, although the

difference in conjugation frequency in manure for T11RF versus T11RFΔ*cas9* recipients was

less than the other conditions at this time point.

Figure 3

FIG 3 A schematic of the experimental design. The donor strain (OG1SSp(pAM714)) was mixed

with two versions of the recipient (T11RF, T11RFΔ*cas9*) in three conditions – liquid BHI,

manure and BHI agar plate. After 18 hours, the mixture (Passage 0) was plated to quantify

donor (spectinomycin, streptomycin, and erythromycin selection), recipient + transconjugants

(rifampin and fusidic acid selection) and transconjugant (rifampin, fusidic acid, and erythromycin

selection) populations. At the same time, Passage 0 was also diluted into fresh liquid BHI

(1:1000 dilution) and incubated for 24 hours to give Passage 1. The Passage 1 culture was

similarly used to generate Passage 2.

We then performed two successive passages of these initial conjugation mixtures. For passage

1, we diluted the passage 0 conjugation mixture 1:1000 into fresh BHI broth and incubated it at

37°C for 24 hours, followed by plating on selective agars. Transferring the conjugation mixture

from manure exposure to liquid BHI was designed to be a crude way to model zoonotic

transmission from an agricultural environment to a human GI tract. Here, we found the

difference in conjugation frequencies in the presence versus absence of CRISPR-Cas

continued to be significantly lower across all three conditions (Fig 5b). This demonstrates that

CRISPR-Cas continued to be effective at blocking plasmid transfer in this sample set.

Passage 1 conjugation mixtures were diluted and incubated as described above to obtain

passage 2 cultures. By the end of passage 2, the transconjugant populations in agar and liquid

BHI media were almost completely absent (limit of detection of 10^2 CFU/mL), irrespective of

CRISPR-Cas. As a result, no differences in conjugation frequencies across the two conditions

were observed (Fig 5b). Surprisingly, for conjugation mixtures that originated from manure,

there was a sizeable transconjugant population following passage 2, but only in the absence of

functional CRISPR-Cas (Fig 5, top right). As a result, the conjugation frequency in the absence

of CRISPR-Cas was significantly higher than in the presence of CRISPR-Cas (Fig 5c). These

results suggest a manure-specific effect on either CRISPR-Cas function and/or the

competitiveness of T11RF(pAM714) transconjugants.

206   **Figure 4**

FIG 4 Bacterial populations across time and niche. Each graph shows the number of donors,

recipients, and transconjugants in different environments. The conjugation experiments shown

on the left had T11RF as the recipient and those on right had T11RF*Δcas9*. Colors in the

background represent the condition in which the bacteria were growing: manure (dark green),

liquid BHI (yellow) and solid agar plate (light orange). Dashed lines represent a CFU/ml count of

10^2 which is the limit of detection.

214   **Figure 5**

215   **FIG 5** CRISPR-Cas conjugation frequency under different conditions. Each point represents log

216   conjugation frequency (calculated as number of transconjugants divided by number of donors)

217   in each of the three conditions where the initial conjugation took place (manure, BHI broth, BHI

218   agar). Student's t-test was used to calculate p-values after determining that log conjugation

219   frequency values followed a normal distribution.

CRISPR antimicrobials appear effective in manure, but are limited by low conjugation frequency

Having shown that CRISPR-Cas can function in an agriculturally-relevant *in vitro* setting, we next wanted to see if CRISPR-Cas-based antimicrobials can be used to selectively deplete antibiotic-resistant *E. faecalis* in that niche. Previous studies have tested their antimicrobial efficacy in solid agar and the murine gastrointestinal tract[36]. Solid agar is a popular choice for conjugation experiments due to the simplicity of its make-up and convenience, but for CRISPR-based antimicrobials to be widely used, they need to be tested in settings that better mimic the *in situ* environment. Manure is one such environmental niche. For this we used a previously designed CRISPR antimicrobial referred to as pKH88[36]. pKH88 is a derivative of the pheromone-responsive plasmid, pPD1, and has been modified to express *cas9* and spacers targeting either *tetM* (for tetracycline resistance) or *ermB* (for erythromycin resistance). For our experiments, we delivered pKH88[sp-*tetM*] or pKH88[sp-*ermB*] from the *E. faecalis* strain CK135RF (donor strain), which is a derivative of the strain OG1. Our target (recipient) strain was OG1SSp(pAM714), which encodes *ermB* on pAM714. Figure 6 shows a schematic diagram of our approach. In this scenario, our goals were to deliver pKH88[sp-*ermB*] to OG1SSp(pAM714), and then either (a) kill or inhibit the recipients' growth by Cas9-mediated DNA damage, or (b) promote the loss of pAM714 via Cas9-mediated DNA damage. We expected no effects of delivering pKH88[sp-*tetM*] to OG1SSp(pAM714), as OG1SSp(pAM714) does not encode *tetM*; pKH88[sp-*tetM*] is our control.

Figure 6

FIG 6 A schematic representation of the CRISPR antimicrobial. The strain CK135RF contains

the pheromone-responsive plasmid pKH88, which encodes *cas9* and a CRISPR guide RNA

under a constitutively active *bacA* promoter. The guide RNA consists of a spacer that targets

either the *ermB* gene or the *tetM* gene, and a repeat sequence. The *cat* gene, encoding

chloramphenicol resistance, is inserted in the plasmid only for tracking purposes. The strain

OG1SSp contains the plasmid pAM714 which harbors the *ermB* gene. In scenario 1, the

CRISPR antimicrobial pKH88[sp-*ermB*] enters OG1SSp and cleaves pAM714. In scenario 2,

OG1SSp(pAM714) donates pAM714 to CK135RF(pKH88[sp-*ermB*]) and the CRISPR

antimicrobial cleaves the incoming plasmid.

With this setup, we tested the hypothesis that the pKH88[sp-*ermB*] CRISPR antimicrobial

plasmid could selectively deplete erythromycin resistance from *E. faecalis* populations cultured

in manure. CRISPR antimicrobial donor strains and OG1SSp(pAM714) recipients were mixed

and plated on solid agar or inoculated into liquid BHI or manure, and passages 1, and 2 were

assessed for donor, recipient, and transconjugant CFU/mL as described above. We observed a

significant decrease in the erythromycin resistant population by passage 2 for agar plate

conjugations (Fig 7a), in line with previously published results[36]. However, a similar decrease in

the erythromycin-resistant population was not observed for liquid BHI or manure conditions in

any of the passages (Fig 7a). To investigate a possible cause for this, we measured the

transconjugant population in each case. We found that the transconjugant populations were

substantially lower in liquid BHI and manure across all passages (10^2 (limit of detection) to 10^7),

as compared to agar (10^9 to 10^{10}) (Fig 7b). A possible reason for this could be the absence of a

solid surface for the bacteria to form a biofilm in the case of liquid BHI and manure, which

promotes conjugation. However, in both conditions and across all passages (with the exception

of passage 0 in liquid BHI), there were fewer transconjugants in the case of the experimental

antimicrobial pKH88-*ermB* as compared to the control pKH88[sp-*tetM*] (Fig 7b), and this

reduction was significant in passages 1 and 2 in manure. Assuming similar plasmid transfer

levels for both pKH88[sp-*ermB*] and pKH88[sp-*tetM*], this could be explained as the action of the

antimicrobial: OG1SSp(pAM714) recipients that received pKH88[sp-*ermB*] had their pAM714

cleaved due to CRISPR-Cas targeting and hence were not detected in transconjugant-selective

plates. This suggests that the antimicrobial was effective in manure when conjugation did occur.

To further test the efficacy of the antimicrobial, we looked at the subset of the transconjugant

population that was erythromycin-resistant. These would be OG1SSp(pAM714) recipient cells

that received the pKH88[sp-*ermB*] or [sp-*tetM*] antimicrobial plasmid and also maintained

erythromycin resistance. We found that in almost every passage for both manure and agar, this

cell population was significantly lower in the case of the experimental antimicrobial pKH88[sp-

ermB] as compared to the control pKH88[sp-*tetM*] (Fig 7c). This points to the effectiveness of

the antimicrobial.

A erm resistant population

B OG1SSp with pKH88 acquired

C OG1SSp with acquired pKH88 and native pAM714

Figure 7

FIG 7 Evaluating CRISPR antimicrobial efficiency. (A) shows the number of total erythromycin-

resistant bacteria in the mixture, across the three conditions, 3 passages and with the control

(tet, in brown) or experimental (erm, in green) CRISPR antimicrobial. A lower number on this

graph signifies depletion of the erythromycin-resistant population. (B) shows the number of

transconjugants i.e. the recipient (OG1SSp) cells that have accepted the antimicrobial plasmid

pKH88. (C) shows the number of OG1SSp(pKH88) transconjugants that still maintain the

pAM714 plasmid. All values were log transformed and t-test with Welch's correction was used to

generate p-values. 'na' represents cases where all values in both groups were the same and

hence a t-test could not be performed. Dashed lines represent a CFU/ml of 10^2 which is the limit

of detection.

Finally, we tested if the presence of the CRISPR antimicrobial could immunize the donor

CK135RF(pKH88) bacteria against acquiring the *ermB* gene from OG1SSp(pAM714). For this

we calculated the number of CK135RF(pKH88) bacteria that were erythromycin-resistant, which

could be the result of pAM714 transferring from OG1SSp to CK135RF. We observed a stark

difference in this population between the experimental and control antimicrobial. The number of

erythromycin-resistant CK135RF(pKH88) was significantly lower in the case of the *erm* targeting

antimicrobial as compared to the one that targeted the *tet* gene (Fig 8). This shows that the

pKH88[sp-*ermB*] CRISPR antimicrobial immunized its host CK135RF against acquisition of

erythromycin resistance . This effect was seen in both manure and agar.

307   **Figure 8**

FIG 8 Immunization of bacteria against antibiotic resistance acquisition in manure. Bars

represent number of CK135RF cells that have become erythromycin-resistant, likely via

accepting pAM714 from OG1SSp, in addition to carrying pKH88. A lower number represents

immunization via the antimicrobial plasmid pKH88. All values were log transformed and t-test

with Welch's correction was used to generate p-values. Dashed lines represent a CFU/ml of 10^2

which is the limit of detection.

3. Materials and Methods

Bacteria and reagents used: E. faecalis strains were grown in brain heart infusion (BHI) broth or

on agar plates. Antibiotic concentrations used were as follows: rifampin, 50 µg/ml; fusidic acid,

25 µg/ml; spectinomycin, 500 µg/ml; streptomycin, 500 µg/ml; erythromycin, 50 µg/ml;

tetracycline, 10 µg/ml; chloramphenicol, 15 µg/ml.

Database collection and genome analysis: NCBI Genome was used to download _E. faecalis_

genomes publicly available as of September 1, 2021. Based on the host (source of isolation),

the downloaded genomes were categorized as originating from human or animal sources.

323 Genomes with hosts that did not match either of these categories were excluded. A recent study

324 provided a significant number of additional genomes to our study that were not part of NCBI

325 Genome (Refseq)[25]. Genomes were subject to filtering based on genome size to eliminate

326 anomalies. The average genome size was calculated, and genomes with sizes less than half or

327 greater than 1.5 times of this average size were discarded (following Refseq recommendation).

328 Strains were queried on BLAST against the *cas9* sequences of CRISPR1-Cas

329 (GCF_000172575.2) and CRISPR3-Cas (GCF_000157475.1) to discover strains with potentially

330 active CRISPR-Cas (percent identity >= 98.5%). CRISPRCasFinder was used to identify

331 CRISPR arrays in these genomes, and the known repeat sequences of CRISPR 1,2,3 were

332 used to filter bona-fide CRISPR arrays from the result[30,44].

333 *Manure microcosm:* Dairy cattle manure was obtained from a settling basin in Kimberley, Idaho,

334 and autoclaved to destroy resident flora. The solid manure was filtered through a cheesecloth to

335 remove large fecal solids, and 1 mL of the resultant liquid manure was used per conjugation

336 reaction.

337 *Conjugation experiments:* Donors and recipients were grown overnight (from freezer stocks) in

338 liquid BHI medium, followed by a 1:10 subculture. The subsample was statically incubated at

339 37°C for 1.5 hours after which time optical density (OD) measurement at 600 nm was taken to

340 ensure similar ODs for both donor and recipient. For each microcosm, 100 µl of the donor

341 culture and 900 µl of a recipient culture were centrifuged, 90% of the supernatant was removed,

342 and then cells were resuspended. Three experimental conditions were used to study

343 conjugation outcomes – solid brain heart infusion (BHI) agar plate, liquid BHI and liquid manure.

344 Conjugation reactions were incubated for 18 hours at 37°C. After this, the conjugation mixture

345 was collected, serial dilutions were performed and plated on agar plates with appropriate

346 antibiotic selection. Where appropriate, the recipient population was calculated by subtracting

347 CFU/mL of transconjugants from that of recipients+transconjugants. Plates were incubated at

37°C for 72 hours followed by counting the number of colony forming units/mL (CFU/mL).

Conjugation frequency was calculated as $\frac{CFU/mL \ of \ transconjugants}{CFU/mL \ of \ donors}$. For multi-passage

experiments, a 1:1000 dilution subculture was prepared and allowed to incubate in liquid BHI for

24 hours before another set of plating. For CRISPR antimicrobial experiments, the pheromone

cAD1 (500 ng/mL) was added to the conjugation mixture to increase conjugation.

4. Discussion

In this study, we assessed the prevalence and function of CRISPR-Cas in agriculturally relevant

niches. We discovered that the prevalence of both CRISPR1-Cas and CRISPR3-Cas (the two

independently functional CRISPR-Cas types in *E. faecalis*) was very similar when comparing

strains of animal origin to those of human origin. These two CRISPR-Cas systems differ in

terms of their protospacer-adjacent motif (PAM) used by the Cas nuclease to cleave the

protospacer in the invading genome, along with differences in the corresponding *cas* genes[13].

The strong similarity in their prevalence suggests that mechanistic differences between the two

subtypes do not have an impact on their distributions in *E. faecalis* from diverse hosts.

We also assessed how effective CRISPR-Cas is an agricultural niche, using dairy manure in a

microcosm design. Previous studies reported that CRISPR-Cas efficacy varies with the

environment it is tested in, with differences observed *in vitro* versus *in vivo*[14]. Our data show *E.*

faecalis CRISPR-Cas is able to block plasmid transfer following exposure to manure. The effect

of manure on CRISPR function is observable even after two post-manure passages in BHI

medium, something that was not observed in the control BHI broth and agar microcosms. We

speculate that even though the manure was autoclaved and most of its solids removed before

using, it still contained factors that could be altering *E. faecalis* physiology. Specifically,

T11RFΔ*cas9* remains at high levels in BHI-passaged cultures post-manure exposure (Figure 4).

This is not observed for T11RF post-manure, or any of the cultures originating in BHI. Teasing

apart the mechanism by which this happens and the relative contributions of manure, CRISPR-

Cas, and pAM714 would be the next step.

Overall, the microcosm studies provided clear evidence of *E. faecalis* CRISPR-Cas functioning

in an agricultural (manure) environment, and highlighted future challenges to be considered in

experiments exploring potential field applications of the concept. First, our manure model is

highly simplified – we autoclaved it to remove all resident microbiota and then filtered and

strained it to remove most of the solids. The former was done to avoid complications that could

arise from the resident microbiota affecting CRISPR-Cas function and the latter to help

standardize quantification of manure amounts. The solids, if present, could provide additional

surface for the bacteria to form biofilms on, which would promote conjugation and hence,

possibly, enhance the effect of CRISPR antimicrobials. Although we saw CRISPR antimicrobials

have some effect, we were unable to bring up the conjugation rates to a sufficient extent to

robustly test the effectiveness of the antimicrobial. It is possible that the large size of our

antimicrobial plasmid was the reason for its low conjugation rate; engineering the antimicrobial

using a smaller plasmid could potentially help improve its conjugation rate and performance in

agricultural niches. Secondly, the experiments could be performed with the resident microbiota

intact by using a combination of antibiotic-selective and species-selective plates. Thirdly, after

letting the bacteria conjugate in the three respective microcosms, we subsequently passaged

the conjugation mixture in liquid BHI in all three conditions, as a crude way to model zoonotic

transfer from manure to the human GI tract while keeping conditions consistent across the three

arms of the experiment. An alternative approach could be to passage each of the three arms in

the same microcosm that they were exposed to during the initial period of conjugation (solid

agar, liquid BHI and manure).

Our work has several important implications both in terms of basic CRISPR biology and tackling

antibiotic resistance. Our analysis looking at CRISPR-Cas prevalence in *E. faecalis* is perhaps

the largest of its kind, and the insight that the distribution of CRISPR-Cas and its subtypes is

invariant across animal and human sources points to a lack of CRISPR-mediated host

colonization preference. Analysis of CRISPR spacers and their targets would provide additional

refinement, illuminating MGEs and corresponding spacers that might be unique to agricultural

isolates. The *E. faecalis* microcosm studies offer a proof-of-concept on the potential for

CRISPR-Cas to limit horizonal gene transfer in dairy manures. Finally, while conjugation

efficiency limited the ability of our CRISPR antimicrobial to spread in manure to antibiotic-

resistant target cells, we did successfully immunize an *E. faecalis* strain against antibiotic

resistance acquisition, which was observed to be more robust in manure compared to liquid BHI

(Figure 8). Overall, our study demonstrates that CRISPR-Cas is a barrier to antibiotic resistance

dissemination in the manure environment, and identifies areas for future study of CRISPR-Cas

efficacy in the agricultural setting.

ACKNOWLEDGEMENT

We thank US Department of Agriculture for their collaboration. We also thank members of the

Palmer lab for their constant guidance and feedback. We are grateful to Dr. Robert Dungan

(USDA ARS, Kimberly, Idaho) for the dairy manure sample that was used for the experiments.

This study was funded by the USDA-ARS Office of National Programs (Project No. 58-3042-8-

012) and administered via National Program 202 Soil and Air. This work was also supported by

R01AI116610 from the National Institutes of Health and the Cecil H. and Ida Green Chair in

Systems Biology Science to K.P.

Supplementary figure 1

5. References

1. Barrangou, R. *et al.* CRISPR provides acquired resistance against viruses in prokaryotes. *Science* **315**, 1709–1712 (2007).

2. Ishino, Y., Krupovic, M. & Forterre, P. History of CRISPR-Cas from Encounter with a Mysterious Repeated Sequence to Genome Editing Technology. *J. Bacteriol.* **200**, e00580-17 (2018).

3. Jinek, M. *et al.* RNA-programmed genome editing in human cells. *eLife* **2**, e00471 (2013).

4. Cong, L. *et al.* Multiplex Genome Engineering Using CRISPR/Cas Systems. *Science* **339**, 819–823 (2013).

5. Sander, J. D. & Joung, J. K. CRISPR-Cas systems for genome editing, regulation and targeting. *Nat. Biotechnol.* **32**, 347–355 (2014).

6. Pickar-Oliver, A. & Gersbach, C. A. The next generation of CRISPR–Cas technologies and applications. *Nat. Rev. Mol. Cell Biol.* **20**, 490–507 (2019).

7. Mojica, F. J. M., Díez-Villaseñor, C., Soria, E. & Juez, G. Biological significance of a family of regularly spaced repeats in the genomes of Archaea, Bacteria and mitochondria. *Mol. Microbiol.* **36**, 244–246 (2000).

8. Pourcel, C., Salvignol, G. & Vergnaud, G. CRISPR elements in Yersinia pestis acquire new repeats by preferential uptake of bacteriophage DNA, and provide additional tools for evolutionary studies. *Microbiol. Read. Engl.* **151**, 653–663 (2005).

9. Swarts, D. C., Mosterd, C., Passel, M. W. J. van & Brouns, S. J. J. CRISPR Interference Directs Strand Specific Spacer Acquisition. *PLOS ONE* **7**, e35888 (2012).

10. Hale, C. R. *et al.* RNA-guided RNA cleavage by a CRISPR RNA-Cas protein complex. *Cell* **139**, 945–956 (2009).

11. Jore, M. M. *et al.* Structural basis for CRISPR RNA-guided DNA recognition by Cascade. *Nat. Struct. Mol. Biol.* **18**, 529–536 (2011).

12. Shmakov, S. A. *et al.* The CRISPR Spacer Space Is Dominated by Sequences from

Species-Specific Mobilomes. *mBio* **8**, e01397-17 (2017).

13. Price, V. J., Huo, W., Sharifi, A. & Palmer, K. L. CRISPR-Cas and Restriction-Modification

Act Additively against Conjugative Antibiotic Resistance Plasmid Transfer in Enterococcus

faecalis. *mSphere* **1**, e00064-16.

14. Price, V. J. *et al.* Enterococcus faecalis CRISPR-Cas Is a Robust Barrier to Conjugative

Antibiotic Resistance Dissemination in the Murine Intestine. *mSphere* **4**, e00464-19 (2019).

15. Sghir, A. *et al.* Quantification of Bacterial Groups within Human Fecal Flora by

Oligonucleotide Probe Hybridization. *Appl. Environ. Microbiol.* **66**, 2263–2266 (2000).

16. Mundt, J. O. Occurrence of enterococci in animals in a wild environment. *Appl. Microbiol.*

11, 136–140 (1963).

17. *Enterococci: From Commensals to Leading Causes of Drug Resistant Infection.*

(Massachusetts Eye and Ear Infirmary, 2014).

18. Codelia-Anjum, A. *et al.* Enterococcal Urinary Tract Infections: A Review of the

Pathogenicity, Epidemiology, and Treatment. *Antibiotics* **12**, 778 (2023).

19. Dahl, A. & Bruun, N. E. Enterococcus faecalis infective endocarditis: focus on clinical

aspects. *Expert Rev. Cardiovasc. Ther.* **11**, 1247–1257 (2013).

20. Arias, C. A. & Murray, B. E. The rise of the Enterococcus: beyond vancomycin resistance.

Nat. Rev. Microbiol. **10**, 266–278 (2012).

21. Suppli, M. *et al.* Mortality in enterococcal bloodstream infections increases with

inappropriate antimicrobial therapy. *Clin. Microbiol. Infect.* **17**, 1078–1083 (2011).

22. Weiner, L. M. *et al.* Antimicrobial-Resistant Pathogens Associated With Healthcare-

Associated Infections: Summary of Data Reported to the National Healthcare Safety

Network at the Centers for Disease Control and Prevention, 2011-2014. *Infect. Control*

Hosp. Epidemiol. **37**, 1288–1301 (2016).

23. Hollenbeck, B. L. & Rice, L. B. Intrinsic and acquired resistance mechanisms in

enterococcus. *Virulence* **3**, 421–433 (2012).

24. Výrostková, J. *et al.* Antimicrobial Resistance of Enterococcus sp. Isolated from Sheep and

Goat Cheeses. *Foods* **10**, 1844 (2021).

25. Zaheer, R. *et al.* Surveillance of Enterococcus spp. reveals distinct species and

antimicrobial resistance diversity across a One-Health continuum. *Sci. Rep.* **10**, 3937

(2020).

26. Rushton-Green, R. *et al.* Agricultural Origins of a Highly Persistent Lineage of Vancomycin-

Resistant Enterococcus faecalis in New Zealand. *Appl. Environ. Microbiol.* **85**, e00137-19

(2019).

27. Bager, F., Madsen, M., Christensen, J. & Aarestrup, F. M. Avoparcin used as a growth

promoter is associated with the occurrence of vancomycin-resistant Enterococcus faecium

on Danish poultry and pig farms. *Prev. Vet. Med.* **31**, 95–112 (1997).

28. Tyson, G. H. *et al.* Prevalence and Antimicrobial Resistance of Enterococci Isolated from

Retail Meats in the United States, 2002 to 2014. *Appl. Environ. Microbiol.* **84**, e01902-17

(2018).

29. Bourgogne, A. *et al.* Large scale variation in Enterococcus faecalis illustrated by the

genome analysis of strain OG1RF. *Genome Biol.* **9**, R110 (2008).

30. Palmer, K. L. & Gilmore, M. S. Multidrug-Resistant Enterococci Lack CRISPR-cas. *mBio* **1**,

e00227-10.

31. Makarova, K. S. *et al.* An updated evolutionary classification of CRISPR–Cas systems. *Nat.*

Rev. Microbiol. **13**, 722–736 (2015).

32. Zou, J. *et al.* Dissemination of Linezolid Resistance Through Sex Pheromone Plasmid

Transfer in Enterococcus faecalis. *Front. Microbiol.* **11**, (2020).

33. Katyal, I., Chaban, B., Ng, B. & Hill, J. E. CRISPRs of Enterococcus faecalis and E. hirae

Isolates from Pig Feces Have Species-Specific Repeats But Share Some Common Spacer

Sequences. *Microb. Ecol.* **66**, 182–188 (2013).

34. Lyons, C., Raustad, N., Bustos, M. A. & Shiaris, M. Incidence of Type II CRISPR1-Cas

Systems in Enterococcus Is Species-Dependent. *PLOS ONE* **10**, e0143544 (2015).

35. He, Q. *et al.* Comparative genomic analysis of Enterococcus faecalis: insights into their

environmental adaptations. *BMC Genomics* **19**, 527 (2018).

36. Rodrigues, M., McBride, S. W., Hullahalli, K., Palmer, K. L. & Duerkop, B. A. Conjugative

Delivery of CRISPR-Cas9 for the Selective Depletion of Antibiotic-Resistant Enterococci.

Antimicrob. Agents Chemother. **63**, e01454-19 (2019).

37. Jacobsen, A. M. & Halling-Sørensen, B. Multi-component analysis of tetracyclines,

sulfonamides and tylosin in swine manure by liquid chromatography-tandem mass

spectrometry. *Anal. Bioanal. Chem.* **384**, 1164–1174 (2006).

38. Laube, H., Friese, A., von Salviati, C., Guerra, B. & Rösler, U. Transmission of ESBL/AmpC-

producing Escherichia coli from broiler chicken farms to surrounding areas. *Vet. Microbiol.*

172, 519–527 (2014).

39. He, L.-Y. *et al.* Dissemination of antibiotic resistance genes in representative broiler feedlots

environments: identification of indicator ARGs and correlations with environmental variables.

Environ. Sci. Technol. **48**, 13120–13129 (2014).

40. Cox, C. R., Coburn, P. S. & Gilmore, M. S. Enterococcal Cytolysin: A Novel Two

Component Peptide System that Serves as a Bacterial Defense Against Eukaryotic and

Prokaryotic Cells. *Curr. Protein Pept. Sci.* **6**, 77–84.

41. McBride, S. M., Fischetti, V. A., LeBlanc, D. J., Jr, R. C. M. & Gilmore, M. S. Genetic

Diversity among Enterococcus faecalis. *PLOS ONE* **2**, e582 (2007).

42. Bolotin, A., Quinquis, B., Sorokin, A. & Ehrlich, S. D. Clustered regularly interspaced short

palindrome repeats (CRISPRs) have spacers of extrachromosomal origin. *Microbiology* **151**,

2551–2561 (2005).

43. Ike, Y., Clewell, D. B., Segarra, R. A. & Gilmore, M. S. Genetic analysis of the pAD1

hemolysin/bacteriocin determinant in Enterococcus faecalis: Tn917 insertional mutagenesis

and cloning. *J. Bacteriol.* **172**, 155–163 (1990).

44. Couvin, D. *et al.* CRISPRCasFinder, an update of CRISRFinder, includes a portable

version, enhanced performance and integrates search for Cas proteins. *Nucleic Acids Res.*

46, W246–W251 (2018).