Convergent generation of atypical prions in knock-in mouse models of genetic prion disease

Surabhi Mehra[1,*], Matthew E.C. Bourkas[1,2,*], Lech Kaczmarczyk[3,4], Erica Stuart[1], Hamza Arshad[1,2], Jennifer K. Griffin[1], Kathy L. Frost[5], Daniel J. Walsh[6], Surachai Supattapone[6,7], Stephanie A. Booth[5,8], Walker S. Jackson[3,4], and Joel C. Watts[1,2,§]

[1]Tanz Centre for Research in Neurodegenerative Diseases, University of Toronto, Toronto, Ontario, Canada

[2]Department of Biochemistry, University of Toronto, Toronto, Ontario, Canada

[3]Wallenberg Center for Molecular Medicine, Department of Biomedical and Clinical Sciences, Linköping University, Linköping, Sweden

[4]German Center for Neurodegenerative Diseases (DZNE), Bonn, Germany

[5]One Health Division, National Microbiology Laboratory, Public Health Agency of Canada, Winnipeg, Manitoba, Canada

[6]Department of Biochemistry and Cell Biology, Geisel School of Medicine at Dartmouth, Hanover, New Hampshire, USA

[7]Department of Medicine, Geisel School of Medicine at Dartmouth, Hanover, New Hampshire, USA

[8]Department of Medical Microbiology and Infectious Diseases, Faculty of Health Sciences, University of Manitoba, Winnipeg, Manitoba, Canada

*These authors contributed equally to this study

§To whom correspondence should be addressed at: Krembil Discovery Tower, Rm. 4KD481, 60 Leonard Ave., Toronto, ON, Canada, M5T 0S8; Tel: (416) 507-6891; Fax: (416) 603-6435; joel.watts@utoronto.ca

Acknowledgements: We are grateful to Sue Plyte and the staff at the UHN Animal Resources Centre for their assistance with facilitating the prion inoculation experiments. The authors thank Dr. Stanley Prusiner (University of California San Francisco) for providing the HuM-D13 antibody and the HRP-labeled HuM-P antibody. This work was funded by a grant from the Canadian Institutes of Health Research (PJT-169048). SM was supported by a postdoctoral fellowship from Parkinson Canada.

Abstract

Most cases of human prion disease arise due to spontaneous misfolding of wild-type or mutant prion protein. Though recapitulating spontaneous prion conversion in animal models has proven challenging, transgenic mice expressing the misfolding-prone bank vole prion protein (BVPrP) recreate certain key aspects of sporadic and genetic prion disease. However, it remains unclear whether spontaneous prion generation can occur in the absence of protein over-expression and how disease-causing mutations affect prion strain properties. To address these issues, we generated knock-in mice expressing physiological levels of either wild-type or mutant BVPrP with isoleucine at codon 109. While mice expressing wild-type BVPrP remained free from neurological disease, a subset of knock-in mice expressing BVPrP with mutations that cause either fatal familial insomnia (D178N) or familial Creutzfeldt-Jakob disease (E200K) developed progressive neurological illness. Brains from spontaneously ill knock-in mice contained prion disease-specific neuropathological changes as well as atypical protease-resistant prion protein. Moreover, brain extracts from spontaneously ill D178N- or E200K-mutant BVPrP knock-in mice transmitted disease to mice expressing wild-type BVPrP. Surprisingly, the properties of the D178N- and E200K-mutant prions appeared identical both pre- and post-transmission, suggesting that both mutations guide the formation of a highly similar atypical prion strain. These findings imply that knock-in mice expressing mutant BVPrP spontaneously develop a *bona fide* prion disease and that mutations causing prion diseases may share a uniform initial mechanism of action. Therefore, these mice represent useful tools for studying the early stages of genetic prion diseases.

Keywords: prion, knock-in mice, protein misfolding, Creutzfeldt-Jakob disease, fatal familial insomnia, transmission

Introduction

Human prion diseases such as Creutzfeldt-Jakob disease (CJD) are caused by misfolding of the cellular prion protein (PrP^C) into PrP^{Sc}, a pathological conformation that aggregates and deposits in the brain [17]. In addition to PrP^{Sc} deposition, the neuropathological hallmarks of prion disease include spongiform degeneration of the brain parenchyma as well as prominent astrocytic gliosis [71]. PrP^C is expressed at its highest levels in neurons and astrocytes, can be modified by the addition of up to two N-linked glycans, and is attached to the outer leaflet of the plasma membrane by a glycosylphosphatidylinositol (GPI) anchor. While PrP^C is predominantly α-helical, PrP^{Sc} aggregates adopt a parallel in-register β-sheet structure and are partially resistant to degradation by proteases such as proteinase K (PK) [42, 49, 51, 70]. Structural variation within PrP^{Sc} aggregates encodes distinct prion strains, which produce unique disease phenotypes [30-32, 48]. Aggregates of PrP^{Sc} can act as seeds to template the conformational conversion of PrP^C into additional PrP^{Sc}, allowing prions to spread from cell-to-cell within the brain and from the periphery into the central nervous system.

The ability of PrP^{Sc} to self-propagate underlies the infectious nature of the prion diseases. However, most cases of prion disease in humans do not manifest due to an infectious etiology. Instead, spontaneous misfolding of PrP^C into PrP^{Sc} within the brain is thought to be the initiating event in sporadic prion diseases, such as sporadic CJD (sCJD). Similarly, in genetic prion disorders such as familial CJD (fCJD), fatal familial insomnia (FFI), and Gerstmann-Sträussler-Scheinker disease (GSS), mutations within the *PRNP* gene encoding PrP are believed to promote the spontaneous formation of PrP^{Sc}. Different disease-causing mutations in human PrP can lead to the formation of distinct prion strains that exhibit differential susceptibility to cleavage by PK [80]. The biochemical hallmark of sCJD, fCJD, and FFI is "stereotypical" protease-resistant PrP (PrP^{res}), also called PrP27-30, which is a C-terminal fragment characterized by multiple PK-resistant glycoforms with molecular weights between 19 and 30

kDa [6, 64]. In contrast, "atypical" N- and C-terminally truncated PrP^{res} fragments with molecular

weights between ~6 and 11 kDa are found in most cases of GSS [66, 77, 78].

Although ~99% of human prion disease cases arise due to the spontaneous generation of PrP^{Sc}

within the brain, little is known about the molecular mechanisms that govern this process.

Inoculation of wild-type or transgenic mice with prions induces a neurodegenerative disorder

that recapitulates all the biochemical and pathological hallmarks of human prion diseases [88].

However, it remains unknown whether the earliest events that occur in the sporadic and genetic

prion diseases are the same as those in the infectious prion diseases, which require a pre-

existing source of PrP^{Sc}. Thus, there has been considerable interest in developing mouse

models that exhibit spontaneous prion formation within the brain. While some success has been

achieved, no existing model fully recapitulates the key features present in the human prion

diseases [11, 87]. Certain lines of transgenic mice overexpressing mouse PrP (MoPrP)

containing prion disease-causing pathogenic mutations develop spontaneous neurological

illness but fail to generate infectious PrP^{Sc} that is highly resistant to PK digestion [10, 15, 22, 27,

34, 35, 52, 56, 79, 92]. Similarly, knock-in mouse models of fCJD, FFI, and GSS that express

physiological levels of mutant MoPrP with the correct spatiotemporal expression pattern within

the brain do not exhibit overt signs of neurological illness nor highly PK-resistant PrP, despite

the presence of prion disease-specific neuropathological changes within the brains of certain

lines [38, 39, 50]. Interestingly, transgenic mice overexpressing mutant human PrP also fail to

develop neurological illness, suggesting that sequence elements within human PrP prevent

spontaneous misfolding within the lifespan of a mouse [4, 5].

In recent years, bank voles (*Myodes glareolus*) have been increasingly used in prion disease

research since they are susceptible to prion strains from several different species, including

humans [1, 2, 21, 58, 59, 67]. Bank vole PrP (BVPrP) facilitates prion replication and formation

in various *in vitro*, cellular, and animal paradigms, indicating that BVPrP is a highly permissive substrate for conversion into PrP^{Sc} [3, 13, 19, 23-26, 53, 60, 73, 85]. Moreover, transgenic mice overexpressing wild-type BVPrP containing isoleucine at polymorphic codon 109 (I109) develop a spontaneous and transmissible neurological illness characterized by prion disease-specific neuropathological changes as well as the presence of a highly PK-resistant PrP fragment in the brain [61, 84, 86]. Addition of the D178N or E200K mutations, which respectively cause FFI and fCJD, to the sequence of BVPrP(I109) hastens the onset of neurological disease in transgenic mice [84]. Collectively, these findings suggest that BVPrP(I109) is intrinsically prone to adopting misfolded, infectious conformations, potentially making it an ideal substrate for studying spontaneous prion formation under conditions of physiological protein expression.

In this study, we sought to generate improved mouse models of spontaneous prion formation by taking advantage of the unique misfolding propensity of BVPrP as well as the more translational nature of knock-in mice. Whereas knock-in mice expressing wild-type BVPrP(I109) remained healthy for their entire lifespan, knock-in mice expressing either D178N- or E200K-mutant BVPrP(I109) developed progressive signs of neurological illness, which were accompanied by the emergence of atypical PrP^{res} within the brain. Furthermore, brain extracts from spontaneously ill D178N or E200K BVPrP(I109) knock-in mice transmitted disease to knock-in mice expressing wild-type BVPrP(I109), confirming the generation of authentic prion infectivity. Surprisingly, stark differences in disease phenotype were not observed between the D178N- and E200K-mutant lines, implying that both mutations may initially act to promote the formation of a similar, perhaps identical atypical prion strain.

Materials and Methods

Generation and characterization of knock-in mice

Gene targeting in V6.5 embryonic stem cells was performed at the DZNE/Bonn University using CRISPR/Cas9 as described previously [40]. Plasmids containing the open reading frames (ORF) of either wild-type, D178N-mutant, or E200K-mutant BVPrP (I109 polymorphic variant) were used as a starting point [84]. Targeting constructs were generated by ligating the respective variants of the BVPrP ORF between EagI and ClaI sites of the intermediate vector pWJPrP101 [40] containing homology regions and a neomycin selection cassette removable by Flp recombinase. The Cas9 vector used for double-strand break generation in the *Prnp* gene is available from Addgene (plasmid #78621) [40]. Expansion of gene-edited embryonic stem cells and aggregation with diploid CD-1(ICR) mouse embryos was performed at The Centre for Phenogenomics (Toronto, Canada). Chimeric mice were identified by their mixed coat colors and then bred with B6(Cg)-*Tyr^{c-2J}*/J mice ("B6-albino mice"; Jackson Lab #000058) to identify those that underwent germline transmission events. Chimeras were crossed with a Flp deleter strain (B6.129S4-*Gt(ROSA)26Sor^{tm1(FLP1)Dym}*/RainJ; Jackson Lab #009086) to remove the selection cassette and then backcrossed with wild-type C57BL/6 mice to remove the Flp transgene. Mice that were positive for the BVPrP knock-in allele and negative for the Flp transgene were then intercrossed to create BVPrP homozygotes. All knock-in lines were maintained by crossing homozygous female with homozygous male mice.

Mice were housed at 4-5 animals per cage and were maintained on a 12 h light-12 h dark cycle, while given unlimited access to food and water. All mouse experiments were performed in accordance with the guidelines set by the Canadian Council on Animal Care under protocols (AUP #4263.17 and 6322.3) approved by the University Health Network Animal Care Committee. The presence of spontaneous neurological illness was assessed in mice 3 times per week until symptom onset. From this point, mice were assessed daily and euthanized upon

progression to end-stage disease symptoms. Symptomatic profiles were determined using the standard diagnostic criteria for prion disease in mice [14]. Alternatively, mice were euthanized upon the appearance of intercurrent illness or when they reached an advanced age, typically around 600 days. Upon collection, mouse brains were divided parasagitally and then either snap frozen using dry ice and stored at -80 °C or immersed in 10% neutral buffered formalin for fixation. Mice were not perfused prior to brain collection.

Brain homogenization and detergent extraction

Frozen mouse hemibrains were homogenized using a Minilys bead beater (PreCellys) to generate 10% (w/v) brain homogenates in Dulbecco's phosphate-buffered saline (DPBS). For the preparation of detergent-extracted protein samples, 9 parts of 10% brain homogenate were mixed with 1 part of 10X detergent extraction buffer [5% (w/v) sodium deoxycholate and 5% (v/v) NP-40 prepared in DPBS]. The samples were incubated on ice for 20 min with vortexing every 5 min and then centrifuged at 5,000x *g* for 5 min at 4 °C. The protein concentration in the supernatant was determined using the bicinchoninic acid (BCA) assay.

Prion transmission assays

Groups of 7-9 knock-in mice expressing wild-type BVPrP(I109) at ~6 weeks of age were anaesthetized using isoflurane gas and then intracerebrally inoculated with 30 µL of 1% brain homogenate prepared from frozen hemibrains and diluted in PBS containing 5% (w/v) BSA. Inoculations were performed into the right cerebral hemisphere to a depth of ~3 mm using a tuberculin syringe with an attached 27 gauge, 0.5-inch needle (BD Biosciences #305945). Inoculated mice were housed in groups of 3-4 animals in disposable cages and monitored for the development of neurological illness as described above. Once progressive signs of neurological illness were apparent, mice were euthanized, and their brains were removed and divided into hemispheres using the sagittal plane. Alternatively, the brains of asymptomatic mice

were collected at the experimental endpoint (540 days post-inoculation). The left hemisphere was frozen and stored at -80 °C while the right hemisphere was fixed in 10% neutral buffered formalin and stored at 22 °C (room temperature) for neuropathological examination. Mice were not perfused prior to brain collection. All inoculation experiments utilized roughly equal numbers of male and female mice, except for one experiment involving inoculation of brain homogenate from an asymptomatic 20-month-old kiBVI^wt mouse, which utilized only female mice.

Detergent insolubility assays

Detergent-extracted brain homogenates containing 50-100 µg total protein were diluted in 1X detergent extraction buffer [0.5% (w/v) sodium deoxycholate, 0.5% (v/v) NP-40 prepared in DPBS] and then ultracentrifuged at 100,000x g for 1 h at 4 °C using a TLA-55 rotor (Beckman). The supernatant was removed, and the pellet was resuspended in 1X Bolt LDS sample buffer (Thermo Fisher #B0007) containing 2.5% (v/v) β-mercaptoethanol. The samples were boiled at 95 °C for 10 min and then analyzed by immunoblotting.

Enzymatic digestions

PNGase F digestions (New England Biolabs #P0704S) were performed according to the manufacturer's recommendation. Briefly, 50 µg of detergent-extracted brain homogenate was mixed with 5 µL of Glycoprotein Denaturing Buffer (10X) and incubated at 95 °C for 10 min. The samples were allowed to cool down on ice, and then 5 µL each of 10% NP-40, 10X GlycoBuffer 2, and 0.5 µL of PNGase F were added to make a final reaction volume of 50 µL. Following incubation at 37 °C for 1-2 h, the reaction was stopped by adding LDS sample buffer (1X final concentration) and incubating the samples at 95 °C for 10 min. Samples were then analyzed by immunoblotting or ELISA.

Thermolysin (TL) was purchased from MilliporeSigma (#T7902) and dissolved in dH_2O to

generate a stock concentration of 1 mg/mL. For TL digestions, 500 µg of detergent-extracted

brain homogenate was diluted into a final volume of 100 µL 1X detergent extraction buffer

containing 100 µg/µL TL for a final TL:protein ratio of 1:50. The reaction mixture was incubated

on a thermomixer at 37 °C for 1 h at 600 rpm, and digestions were stopped by the addition of

EDTA to a final concentration of 5 mM. Sarkosyl was then added to a final concentration of 2%

(v/v). This was followed by ultracentrifugation at 100,000x g for 1 h at 4 °C. Finally, the

supernatant was gently removed, and the pellet was resuspended in 1X Bolt LDS sample buffer

containing 2.5% (v/v) β-mercaptoethanol, boiled, and analyzed by immunoblotting. Alternatively,

detergent-extracted brain homogenate was treated with different concentrations of TL at 37 °C

for 1 h and then analyzed directly by immunoblotting without isolating the insoluble fraction. A

stock solution of proteinase K (PK) at a concentration of 20 mg/mL was obtained from Thermo

Scientific (#EO0491). For PK digestions, a similar protocol to that used for TL digestions was

employed, except that 1 mg of detergent-extracted brain homogenate was digested with 50

µg/µL PK for a final PK:protein ratio of 1:50, and the reaction was stopped by adding PMSF to a

final concentration of 2 mM.

Immunoblotting

Proteins were separated on 10% Bolt Bis-Tris gels (Thermo Fisher Scientific) and then

transferred onto 0.45 mm polyvinylidene fluoride (PVDF) membranes using Tris-Glycine transfer

buffer containing 20% (v/v) methanol. Membranes were blocked in 5% (w/v) skim milk prepared

in 1X Tris-buffered saline containing 0.05% (v/v) Tween-20 (TBST) overnight at 4 °C or for at

least 1 h at 22 °C. The following day, membranes were incubated with primary antibody for 1 h

at 22 °C. The primary antibodies that were used include anti-PrP antibodies HuM-D18 (1:5,000

dilution), HuM-P (1:10,000 dilution), HuM-D13 (1:10,000 dilution), POM1 (MilliporeSigma

#MABN2285; 1:5,000 dilution), HuM-R1 (1:10,000 dilution), EP1802Y (Abcam #ab52604; 1:10,000 dilution), and SAF-32 (1:5,000 dilution); and an anti-GFAP antibody (Thermo Scientific #A-21282, 1:10,000 dilution). Blots were washed 3 times with TBST (10 min each), incubated with horseradish peroxidase (HRP)-linked secondary antibodies diluted in blocking buffer for 1 h at 22 °C, and then washed an additional 3 times with TBST. Membranes were developed using Western Lightning ECL Pro (PerkinElmer Inc.) or SuperSignal West Dura Extended Duration Substrate (Thermo Fisher Scientific) and exposure to X-ray films. For reprobing, blots were washed with TBST and then incubated in blocking buffer containing 0.05% (w/v) sodium azide overnight at 4 °C to inactivate the HRP. The next day, blots were reprobed with actin 20–33 antibody (MilliporeSigma #A5060; 1:10,000 dilution).

Enzyme-linked immunosorbent assays (ELISAs)

Relative BVPrP expression levels in the knock-in mice were determined by ELISA. Various amounts of recombinant BVPrP(I109) [3] and detergent-extracted brain homogenates from kiBVI^{wt} mice were used as standards. Immulon 4 HBX 96-well plates (VWR #62402-959) were coated with the anti-PrP antibody HuM-D18 at 5 µg/mL in coating buffer (200 mM NaH$_2$PO$_4$, pH 7.5) overnight at 4 °C. The plate was blocked with 1% BSA diluted in PBS with 0.05 % Tween-20 (PBS-T) at 22 °C for > 2 h, followed by washing with PBS-T. Thereafter, the standards and test samples prepared in PBS containing 0.5 % Triton-X were added in triplicates and the plate was incubated overnight at 4 °C with shaking. Following 4 washes with PBS-T, the HRP-labeled HuM-P detection antibody was added at 1:50,000 dilution in blocking buffer and incubated at 22 °C for 2 h with shaking. The plate was thoroughly washed 5 times with PBS-T and 100 µL of TMB-Blue substrate (DAKO) was added followed by incubation in the dark for 5-10 min. The reaction was stopped by adding 100 µL of 1 M HCl to each well. Finally, the absorbance at 450 nm was read using a BMG CLARIOstar microplate reader.

Conformational stability assays

Assays were performed using a concentration gradient of guanidine hydrochloride (GdnHCl). 20 µL of detergent-extracted brain homogenates were added to an equal volume of 2X GdnHCl stocks to create final concentrations of 1, 1.5, 2, 2.5, 3, 3.5, and 4 M GdnHCl. To generate the 0 M sample, an equal volume of DPBS was added. The samples were incubated at 22 °C for 2 h with shaking (800 rpm). The concentration of GdnHCl was then diluted to 0.4 M by adding detergent extraction buffer (1% final concentration) and DPBS. The samples were then subjected to TL digestion (100 µg/mL) as described above and ultracentrifuged at 100,000x g for 1 h at 4 °C. The supernatants were gently removed and the pellets were resuspended in 1X Bolt LDS sample buffer containing 2.5% (v/v) β-mercaptoethanol and boiled for 10 min at 95 °C. Densitometry analysis was carried out using ImageJ from three independent replicates and [GdnHCl]$_{50}$ values (the concentration of GdnHCl at which 50% of the PrP aggregates are solubilized) were calculated using a variable slope (four parameter) dose-response model in GraphPad Prism as described previously [45].

Neuropathology

Formalin-fixed hemibrains were processed using a Leica Pearl automated tissue processor and then embedded in paraffin. Sections (5 µm) were cut, mounted on positively charged glass slides, and then dried overnight at 37 °C. Slides were deparaffinized using xylene, rehydrated through a graded series of ethanol, and then either stained with hematoxylin and eosin (H&E) or processed for immunohistochemistry. For H&E staining, slides were stained with Hematoxylin 560 MX (Leica #3801575) for 2 min, rinsed with dH$_2$O, incubated with Blue Buffer 8 (Leica #3802915) for 90 s, rinsed with dH$_2$O, and then incubated with Define MX-AQ (Leica #3803595) for 30-45 s. After rinsing with dH$_2$O, the slides were incubated with Eosin 515 LT (Leica #3801619) for 2 min, washed 3X with 100% ethanol (3 min each), incubated with 3 changes of xylenes (5 min each), and then coverslipped and mounted with Permount. For GFAP

immunostaining, the Polink-2 Plus HRP Rabbit DAB Detection kit (GBI Labs) was used. Slides were treated with 3% (v/v) hydrogen peroxide for 10 min and then washed with dH$_2$O. Antigen retrieval was performed using 10 mM sodium citrate, pH 6 for 30 min at 95 °C, and then slides were cooled and washed twice (2 min each) with TBST. The rabbit polyclonal GFAP antibody (Dako #Z0334; 1:4,000 dilution) was applied overnight at 4 °C. After rinsing with TBST, development, and counterstaining with hematoxylin, slides were dehydrated and then coverslipped and mounted using Permount Mounting Medium (ThermoFisher Scientific). For PrP immunohistochemistry, slides were treated with 98% formic acid for 10-15 min, rinsed 3 times with dH$_2$O (5 min each), and then processed using the M.O.M kit (Vector Laboratories). Slides were incubated with the mouse monoclonal PrP antibody 9A2 (Wageningen Bioveterinary Research; 1:500 dilution) for 30 min at 22 °C. Slides were developed using the NovaRed system (Vector Laboratories), counterstained with hematoxylin, dehydrated, and then coverslipped and mounted using VectaMount. All slides were digitized using the Zeiss Axio Scan.Z1 slide scanner and then representative images were captured using PMA.start.

For quantification of vacuolation, snapshots of scanned slides from H&E-stained tissue were taken and then converted to 8-bit black and white images using ImageJ. Freehand regions of interest were drawn around the desired brain region, and then the threshold was set to 0-210 to reveal areas of the brain without stain. Following conversion to binary and creation of a binary mask, the Analyze Particles function in ImageJ was used to determine the percentage brain area covered by vacuolation. A size range of 8 to infinity and a circularity of 0.8 to 1.0 was used to ensure that the interiors of cerebral blood vessels were not counted as vacuoles.

For quantification of astrocytic gliosis, snapshots of scanned slides from tissue stained with a GFAP antibody were taken and converted to 8-bit black and white images using ImageJ. The H-DAB model in the IHC Toolbox plugin in ImageJ was used to remove non-stained regions.

Freehand regions of interest were then drawn around the desired brain region and then the threshold was set to 0-100. The percentage area covered by GFAP staining was then measured.

Statistical analysis

All statistical analysis was conducted using GraphPad Prism (version 10.0.0) with a significance threshold of $P < 0.05$. No tests for data normality were performed. Survival curves were compared by the log-rank (Mantel-Cox) test and sex-specific differences were analyzed by the Mann-Whitney test. ELISA measurements were compared using Welch ANOVA followed by Dunnett's T3 multiple comparisons test. Detergent-insoluble PrP levels were compared using one-way ANOVA followed by Tukey's multiple comparisons test. Neuropathological indicators were compared using a Kruskal-Wallis test followed by Dunn's multiple comparisons test.

Results

Generation of BVPrP knock-in mice

To generate mice expressing physiological levels of wild-type or mutant BVPrP, we replaced the open reading frame encoding MoPrP, which is located entirely within the third exon of the *Prnp* gene, with that of BVPrP using a CRISPR/Cas9-based gene editing approach (**Fig. 1a**) [40]. Following removal of N- and C-terminal signal sequences, the amino acid sequence of mature BVPrP differs from MoPrP at 8 positions. To promote spontaneous prion formation in the brain, we used the I109 polymorphic variant of BVPrP since transgenic mice over-expressing wild-type (wt) BVPrP(I109) develop spontaneous disease much more rapidly than mice over-expressing BVPrP with methionine at codon 109 (M109) [41, 86]. We have previously found that addition of the D178N and E200K mutations to BVPrP(I109) hastens the onset of spontaneous neurological illness in transgenic mice [84]. Thus, we generated knock-in mice expressing either wt, D178N-mutant, or E200K-mutant BVPrP(I109), which will be referred to as kiBVI^wt,

kiBVI[D178N], and kiBVI[E200K] mice, respectively. All mice were homozygous for the knock-in alleles and thus only express wt or mutant BVPrP(I109). The E200K mutation causes fCJD when paired with either methionine (M) or valine (V) at polymorphic codon 129 in human PrP [28, 33]. In contrasting, the D178N mutation causes either FFI if it occurs *in cis* to M129 or fCJD if it occurs *in cis* to V129 [29]. Since BVPrP contains methionine at codon 129, the kiBVI[D178N] mice are more reflective of FFI than fCJD.

Levels of BVPrP(I109) expression in the brain differed among the three lines, with relative levels dependent on the anti-PrP antibody used for detection by immunoblotting (**Fig. 1b, S1a**). PrP levels in kiBVI[E200K] mice were similar to or slightly lower than levels in kiBVI[wt] mice, whereas levels in kiBVI[D178N] mice were considerably lower. Quantification of relative BVPrP(I109) expression levels in the brain by ELISA revealed that PrP levels were ~50-60% lower in kiBVI[D178N] mice than in kiBVI[wt] mice but similar between kiBVI[E200K] and kiBVI[wt] mice (**Fig. 1c**). This is consistent with results from MoPrP-based knock-in mice and our previous finding that D178N-mutant BVPrP(I109) has a shorter half-life than either wt or E200K-mutant BVPrP(I109), resulting in lower steady-state levels [39, 84]. Brain BVPrP(I109) levels in kiBVI[wt] mice were comparable to MoPrP levels in the brains of wild-type C57BL/6 mice, and PrP was not detected in any of the knock-in lines when using an antibody that recognizes MoPrP but not BVPrP (**Fig. 1d**). Following removal of N-linked glycans, the endoproteolytic processing of BVPrP(I109) to produce C1 and C2 fragments was observed in all 3 lines (**Fig. 1e, S1b**). This indicates that the synthesis and processing of BVPrP(I109) in the knock-in lines occurs similarly to MoPrP in wild-type mice.

Mutant BVPrP knock-in mice develop spontaneous disease
Groups of kiBVI[wt], kiBVI[D178N], and kiBVI[E200K] mice were monitored longitudinally for the development of signs of spontaneous neurological illness consistent with prion disease. Mice

were analyzed up to 20 months of age since non-specific causes of morbidity or mortality in aged mice appear more frequently after this point. Whereas all kiBVI^{wt} mice remained free of neurological illness for the duration of the experiment, a subset of kiBVI^{D178N} and kiBVI^{E200K} mice developed progressive signs of neurological disease beginning around 400 days of age (**Fig. 2a, Table 1**). In spontaneously sick kiBVI^{E200K} and kiBVI^{D178N} mice, the most common signs of illness were prominent kyphosis and weight loss with the variable presence of dehydration, limb clasping, and gait issues. Some mice also exhibited a severe dermatitis phenotype characterized by neurotic scratching/over-grooming of the head region, and this was more common in the kiBVI^{D178N} line. Approximately 60% of kiBVI^{D178N} and kiBVI^{E200K} mice developed spontaneous disease by 20 months of age, and there was no difference in kinetics of disease onset between the two mutant BVPrP-expressing lines. For the kiBVI^{D178N} and kiBVI^{E200K} mice that developed spontaneous neurological illness, there was no difference in the age of disease onset between male and female animals (**Fig. 2b**).

Progressive accumulation of misfolded PrP species in mutant BVPrP knock-in mice
We have previously shown that spontaneously ill transgenic mice overexpressing wild-type or mutant BVPrP(I109) contain atypical PrP^{res} species in their brains [84]. Thus, we analyzed brain extracts from the knock-in mice for the presence of detergent-insoluble and protease-resistant PrP (**Fig. 3a**). Initially, we used the protease thermolysin (TL), which has been shown to effectively discriminate between normal and aggregated forms of proteins in prion disease and other protein misfolding disorders [20, 44, 52, 62]. In brains from healthy 3-month-old knock-in mice, full-length PrP species were fully degraded by TL concentrations of higher than 2 µg/mL, whereas complete digestion of the C1 endoproteolytic fragment in the kiBVI^{wt} and kiBVI^{E200K} lines required TL concentrations of at least 20 µg/mL (**Fig. S2**). Interestingly, the C1 fragment in kiBVI^{D178N} mice was much more sensitive to TL digestion, potentially suggesting that the D178N mutation destabilizes the C-terminal domain of BVPrP(I109). To ensure complete digestion of all

properly folded PrP species, we used TL at a concentration of 100 µg/mL in all subsequent protease digestion experiments.

Despite expressing similar or lower levels of BVPrP, levels of detergent-insoluble PrP species were substantially elevated in 3-month-old kiBVI^{E200K} and kiBVI^{D178N} mice compared to kiBVI^{wt} mice (**Fig. 3b**). At this age, all detergent-insoluble PrP species were sensitive to TL digestion. Compared to 20-month-old kiBVI^{wt} mice, increased amounts of detergent-insoluble PrP species were also present in spontaneously ill kiBVI^{E200K} and kiBVI^{D178N} mice, and these species were resistant to TL digestion (**Fig. 3c**). TL-resistant PrP was not detected in any of the brains from 20-month-old kiBVI^{wt} mice (**Fig. 3c, d, Table 1**). In contrast, 100% of the brains from kiBVI^{D178N} mice that developed spontaneous neurological illness displayed TL-resistant PrP (**Fig. 3d, Table 1**). Moreover, all brains from asymptomatic kiBVI^{D178N} mice collected at 20 months of age also contained TL-resistant PrP (**Fig. S3**). The presence of TL-resistant PrP was more variable in the kiBVI^{E200K} line with only 64% of brains examined exhibiting clear evidence of TL-resistant species (**Table 1**). Not all kiBVI^{E200K} mice that developed spontaneous neurological illness possessed TL-resistant PrP (**Fig. 3d**), possibly due to misdiagnosis, and TL-resistant PrP was variably present in the brains from asymptomatic kiBVI^{E200K} mice collected at 20 months of age (**Fig. S3**). The main TL-resistant PrP species appeared to be similar in molecular weight to undigested PrP (**Fig. 3c, d**), and the slight difference in appearance between the kiBVI^{E200K} and kiBVI^{D178N} lines could reflect differences in N-glycosylation efficiency, which was also observed in PrP^C species from young mice (**Fig. 1b**), or mutation-specific differences in protein charge. A smaller and less abundant ~17 kDa TL-resistant fragment was also observed in some of the spontaneously ill kiBVI^{E200K} and kiBVI^{D178N} mice (**Fig. 3c**).

For all three knock-in lines, total PrP levels remained largely unchanged between young and old mice (**Fig. S4a**), whereas levels of detergent-insoluble PrP were modestly increased in

spontaneously ill kiBVI^{E200K} and kiBVI^{D178N} mice compared to younger mice (**Fig. S4b, c**). TL-resistant PrP was absent in kiBVI^{E200K} and kiBVI^{D178N} mice up to 12 months of age, suggesting that it may arise relatively late in the disease course (**Fig. 3e**). The brains of mice with high levels of TL-resistant PrP were also examined for the presence of PK-resistant PrP species. A PK concentration of 50 μg/mL was utilized, as this results in a rigorous digestion of PrP^C. A subset of mice examined exhibited a PK-resistant PrP fragment with a molecular weight of ~10 kDa (**Fig. 3d**). As with TL digestion, PK-resistant PrP species were more commonly found in brains from the kiBVI^{D178N} line than the kiBVI^{E200K} line (**Table 1**). Collectively, these results suggest that PrP^{res} progressively accumulate with age in the brains of kiBVI^{E200K} and kiBVI^{D178N} mice, but not kiBVI^{wt} mice.

Prion disease-specific neuropathological changes in mutant BVPrP knock-in mice

Brains from spontaneously ill kiBVI^{E200K} and kiBVI^{D178N} mice exhibited prominent grey matter vacuolation in the hippocampus, cortex, and thalamus, whereas no vacuolation was observed in the brains of asymptomatic 20-month-old kiBVI^{wt} mice (**Fig. 4a, b**). Age-related white matter vacuolation was present in the brains of all older mice, regardless of symptomatic status or genotype (**Fig. S5**). There was no difference in the extent or localization of vacuolation between symptomatic kiBVI^{E200K} and kiBVI^{D178N} mice. The brains of symptomatic kiBVI^{E200K} and kiBVI^{D178N} mice also exhibited increased astrocytic gliosis compared to aged asymptomatic kiBVI^{wt} mice in the brain regions with vacuolation, as evidenced by GFAP immunolabeling (**Fig. 4c, d**). Although the increase in GFAP staining was less prominent for the kiBVI^{D178N} line, there were no statistically significant differences in the extent of GFAP immunolabeling between the kiBVI^{E200K} and kiBVI^{D178N} mice (**Fig. 4d**). Increased GFAP levels in aged symptomatic kiBVI^{E200K} and kiBVI^{D178N} mice compared to aged asymptomatic kiBVI^{wt} mice and young asymptomatic kiBVI^{E200K} and kiBVI^{D178N} mice were also observed by immunoblotting (**Fig. 4e, f**). Endoproteolytic cleavage products of GFAP similar to those found in the brains of prion-infected

mice and believed to be generated by calpain cleavage were also observed in brain homogenates from symptomatic kiBVI^{E200K} and kiBVI^{D178N} mice [91].

Immunohistochemical staining for PrP using the 9A2 antibody failed to detect any extracellular PrP^{Sc} deposition in regions of the brain that exhibited spongiform degeneration from spontaneously ill kiBVI^{D178N} and kiBVI^{E200K} mice. However, intracellular PrP deposition within glial cells was selectively present in the striatum of sick kiBVI^{D178N} and kiBVI^{E200K} mice (**Fig. 4g**), and there were no apparent differences in PrP deposition patterns between the two lines. We hypothesize that this intracellular staining may represent uptake of misfolded PrP species by glial cells and subsequent failure of the endosomal/lysosomal system to clear these aggregates effectively.

Transmission properties of spontaneously formed prions

To test if prion infectivity was present in the brains of spontaneously ill kiBVI^{E200K} and kiBVI^{D178N} mice, we performed transmission studies in kiBVI^{wt} mice, which do not themselves develop spontaneous disease (**Fig. 5a**). Two inocula each from spontaneously ill kiBVI^{E200K} and kiBVI^{D178N} mice were selected based on the presence of high amounts of TL-resistant PrP in brain extracts. As a negative control, kiBVI^{wt} mice were inoculated with brain extracts prepared from either of two distinct asymptomatic 20-month-old kiBVI^{wt} mice. All the mice inoculated with kiBVI^{wt} extract remained free of neurological illness for up to 18 months post-inoculation (**Fig. 5b, Table 2**). To the contrary, 100% of kiBVI^{wt} mice inoculated with brain extracts from spontaneously ill kiBVI^{E200K} mice developed neurological illness with mean incubation periods of ~10-11 months. The most common clinical signs were kyphosis, weight loss, tremor, and bradykinesia. Disease transmission was less efficient when using inocula prepared from spontaneously ill kiBVI^{D178N} mice, potentially due to a slight transmission barrier introduced by

the D178N mutation, with only ~40% of inoculated mice exhibiting neurological illness within the 18-month experimental timeframe (**Fig. 5b, Table 2**).

Whereas brains from kiBVI^{wt} mice inoculated with kiBVI^{wt} extract did not contain PrP^{res}, both TL- and PK-resistant PrP species were detected in the brains of all mice inoculated with kiBVI^{E200K} extract (**Fig. 5c**). Likewise, TL- and PK-resistant PrP were found in the brains of some, but not all kiBVI^{wt} mice inoculated with kiBVI^{D178N} extract. Surprisingly, of the 5 kiBVI^{wt} brains that displayed TL-resistant PrP following inoculation with kiBVI^{D178N} extract, 4 were from mice collected at 540 days post-inoculation in the absence of clinical signs of neurological illness. The TL- and PK-resistant PrP species in the inoculated kiBVI^{wt} mice were reminiscent of those present in spontaneously ill kiBVI^{E200K} and kiBVI^{D178N} mice, and there were no obvious differences in molecular weights of the PrP^{res} fragments induced by the kiBVI^{E200K} and kiBVI^{D178N} lines. The brains of kiBVI^{wt} mice inoculated with kiBVI^{E200K} or kiBVI^{D178N} extract also contained neuropathological indicators of prion disease such as vacuolation and astrocytic gliosis as well as increased endoproteolytic cleavage of GFAP (**Fig. 5d-f**). Unlike in the spontaneously sick mice, vacuolation in the kiBVI^{wt} mice inoculated with kiBVI^{E200K} or kiBVI^{D178N} samples was largely restricted to the hippocampus. GFAP staining in the hippocampus was less pronounced in the mice inoculated with samples from kiBVI^{D178N} mice, likely reflecting the reduced rate of symptomatic transmission. Intracellular PrP deposition within striatal glial cells of inoculated kiBVI^{wt} mice was observed in animals inoculated with brain extract from kiBVI^{E200K} or kiBVI^{D178N} mice (**Fig. 5d**). There were no obvious differences in the neuropathological signatures of kiBVI^{wt} mice inoculated with kiBVI^{E200K} or kiBVI^{D178N} samples. Collectively, these results demonstrate that transmissible prions can form spontaneously in the brains of kiBVI^{E200K} and kiBVI^{D178N} mice, but not in the brains of kiBVI^{wt} mice.

Conformational characterization of spontaneous and transmitted prions

To characterize the conformational properties of the prions formed spontaneously in the brains of kiBVI^{E200K} and kiBVI^{D178N} mice, we employed a conformational stability assay that measures the relative resistance of protein aggregates to denaturation with guanidine hydrochloride (**Fig. S6**) [45]. This assay is commonly used to discriminate between different prion strains [46, 65]. There was no difference in the conformational stability of the TL-resistant PrP aggregates in the brains of spontaneously ill kiBVI^{E200K} and kiBVI^{D178N} mice (**Fig. 6a, b**). Similarly, no differences in conformational stability were observed following transmission to kiBVI^{wt} mice (**Fig. 6c, d**).

Discussion

To the best of our knowledge, the kiBVI^{D178N} and kiBVI^{E200K} mice described herein are the first mouse models that exhibit all the cardinal features of human prion diseases without requiring the use of PrP overexpression, a non-native promoter, or injection with a pre-existing source of PrP^{Sc}. The mice developed progressive signs of neurological illness, prion disease-specific neuropathological changes such as spongiform degeneration and astrocytic gliosis, as well as TL- and PK-resistant PrP species in their brains. Most importantly, brain extracts from spontaneously sick mice expressing mutant BVPrP transmitted disease to kiBVI^{wt} mice expressing wild-type BVPrP, which did not develop spontaneous disease themselves, confirming the generation of authentic prion infectivity in the kiBVI^{D178N} and kiBVI^{E200K} lines. Thus, these mice will be useful for interrogating the biological mechanisms governing spontaneous prion formation in the brain during FFI and fCJD as well as uncovering therapeutic strategies for counteracting these processes. For instance, we show in a complementary study that small molecules known to interfere with templated prion replication do not prolong the onset of spontaneous disease in either the kiBVI^{D178N} or kiBVI^{E200K} lines [83]. A limitation of our study is that while the D178N and E200K mutations are most commonly heterozygous in cases of genetic prion disease (i.e., wt and mutant PrP are simultaneously present), we employed mice

that were homozygous for the mutations and thus expressed higher relative levels of mutant PrP. However, it should be noted that individuals with homozygous E200K mutations have been identified and develop typical fCJD but at a younger age than those with a heterozygous mutation [57, 76].

We utilized the I109 polymorphic variant of BVPrP for our studies because it more readily forms prions spontaneously when overexpressed in transgenic mice than the M109 polymorphic variant [86]. Indeed, transgenic mice that display modest overexpression of sheep PrP with isoleucine at position 112, which is analogous to position 109 in BVPrP, also develop a spontaneous and transmissible prion disease [82]. Transgenic mice expressing very high levels of the M109 BVPrP polymorphic variant developed a late-onset, but non-transmissible spontaneous neurological illness characterized by spongiform degeneration and PrP deposition in the absence of PrP[res] [41]. Overexpression of mouse or hamster PrP in transgenic mice can also lead to a proteinopathy that resembles authentic prion disease but is non-transmissible [37, 89]. Thus, the presence of isoleucine at codon 109 of BVPrP may be critical for permitting the formation of PrP assemblies that are able to self-propagate. However, we did not observe any signs of spontaneous prion formation in aged kiBVI[wt] mice, which is consistent with the lack of reports describing spontaneous disease in I109 bank voles. Therefore, PrP overexpression is required to initiate spontaneous prion formation in mice expressing wt PrP, even when a permissive substrate such as BVPrP is utilized.

Knock-in mouse models that express D178N- or E200K-mutant MoPrP have previously been generated and spontaneously develop prions that can be transmitted to mice expressing wild-type MoPrP [38, 39]. However, unlike the BVPrP-based knock-ins, they do not develop overt signs of progressive neurological illness and fail to exhibit the PrP[res] species that typify authentic prion disease. Besides the presence of isoleucine at codon 109, the sequence determinants of

BVPrP that allow it to function as a superior substrate for studying spontaneous prion formation compared to MoPrP remain unknown. One study found that the addition of two BVPrP-specific C-terminal residues (E227 and S230) to MoPrP was sufficient to elicit spontaneous disease in mice [41], and this region has been implicated as a "linchpin domain" responsible for BVPrP's enhanced susceptibility to prion strains from other species [12]. Other studies have identified residues within the α-helical domain of BVPrP, specifically asparagine at positions 155 and 170, that may facilitate cross-species prion transmission [1, 43]. How any of these specific residue differences may promote prion formation is unclear, since the structure of bank vole PrP^C is very similar to that of PrP^C from other mammals, other than the existence of a more rigid loop immediately prior to the second α-helix [16]. The presence of a rigid loop in the structure of PrP^C has been linked to the appearance of spontaneous prion generation in transgenic mice [74, 75].

While brains from most aged knock-in mice expressing mutant BVPrP exhibited TL-resistant PrP, a smaller fraction contained PK-resistant PrP. This suggests that a spectrum of PrP assemblies develop in the knock-in mice, with the formation of TL-resistant species likely preceding the generation of potentially larger or more densely packed PK-resistant aggregates (**Fig. 7**). Whereas FFI and fCJD patients develop stereotypical PrP^{res} with a molecular weight of 19-21 kDa following removal of N-linked glycans [55, 80], the brains of spontaneously sick kiBVI^{D178N} and kiBVI^{E200K} mice exhibited a smaller PK-resistant PrP fragment with a molecular weight of ~10 kDa, which is more reminiscent of those found in GSS [66, 77, 78]. Short PK-resistant PrP fragments can also be found in Nor98/atypical scrapie in sheep and in variably protease-sensitive prionopathy (VPSPr) in humans [8, 69, 93]. Interestingly, these "atypical" prion diseases, all of which are thought to result from spontaneous misfolding of PrP, can be efficiently transmitted to bank voles with the I109 polymorphism [59, 67, 68], and ~7 kDa non-fibrillar PrP^{res} species purified from GSS brains are sufficient for disease transmission [81]. Thus, the presence of the I109 residue may selectively stabilize these misfolded PrP species,

allowing them to propagate and spread within the brain. The presence of prion disease-associated neuropathological changes such as spongiform degeneration in spontaneously ill kiBVI^{D178N} and kiBVI^{E200K} mice despite the absence of extracellular PrP^{Sc} deposition and stereotypical PrP^{res} is consistent with the hypothesis that stereotypical PrP^{res} may not be the principal neurotoxic species in prion disease [9, 72].

Potential relationships between atypical PrP^{res} and the stereotypical PrP^{res} species found in most animal and human prion diseases remain to be fully investigated. Repeated passage of Nor98/atypical scrapie in transgenic mice expressing bovine PrP leads to a molecular phenotype indistinguishable from classical bovine spongiform encephalopathy, which is typified by the presence of stereotypical PrP^{res} [36]. Moreover, whereas I109 bank voles faithfully propagate the atypical short PrP^{res} fragments upon transmission of Nor98 scrapie, Nor98-inoculated M109 bank voles exhibit classical PrP^{res} [68]. Intriguingly, cases of GSS caused by the P102L mutation exhibit a short ~8 kDa PrP^{res} fragment, either alone or in combination with stereotypical PrP^{res} [54, 63]. Therefore, stereotypical PrP^{res} may emerge from or along with atypical PrP^{res}, potentially due to selection of rare conformational species within a mixture or deformed templating [18, 47].

The striking similarities in the neuropathological profiles, PrP^{res} signatures, conformational stabilities, and transmission properties of the prions present in kiBVI^{D178N} and kiBVI^{E200K} mice imply that both lines spontaneously develop a highly similar, if not identical atypical prion strain. This was unexpected given that BVPrP-based transgenic models of FFI and fCJD exhibit mutation-specific pathological signatures, although this could potentially be an artifact of PrP over-expression or differences in spatial expression patterns [84]. MoPrP-based knock-in models of FFI and fCJD also exhibit different pathological phenotypes, perhaps due to the lack of residue I109 since MoPrP contains leucine at corresponding residue 108 [38, 39]. However,

translatome studies have shown that the molecular phenotypes in the MoPrP-based FFI and fCJD knock-in lines are more similar than expected [7]. Although we cannot rule out a scenario where the properties of BVPrP(I109) override mutation-specific strain generation, our data raise the possibility that the D178N and E200K mutations may uniformly act to promote the initial misfolding of PrP into a similar self-propagating species (**Fig. 7**). The key properties of this initial misfolded species are likely determined by its atypical PK-resistant core rather than by the specific mutation, since similar fragments are known to be both N- and C-terminally truncated and do not contain either residue 178 or 200 [30, 69, 81]. The mechanism by which the D178N and E200K mutations promote the accumulation of misfolded BVPrP remains to be established. The mutations may either directly promote spontaneous misfolding of BVPrP, stabilize the formation of prion assemblies, and/or reduce the clearance rates of aggregates. An increased turnover rate for the mutant proteins implies a mutation-induced decrease in protein stability [84], and the D178N mutation may further accelerate misfolding by destabilization of the C1 endoproteolytic fragment, which is thought to be an endogenous inhibitor of prion conversion [90].

In humans, disease-causing mutations within PrP clearly influence prion strain generation in the brain. For instance, individuals with the D178N mutation coupled to M129 only develop FFI. In contrast, those with the E200K-M129 haplotype develop CJD with one of three distinct PrP[res] signatures [6]. We propose a dual-hit model in which PrP mutations play two roles in genetic prion disease. First, they act by promoting the accumulation of an atypical misfolded PrP species and second, later in the disease course, they bias the emergence of classical PrP[res] towards conformations that are compatible with the specific mutation. Given that this latter phase might require a rare secondary misfolding event, our two-phase model can provide insights into why genetic prion diseases manifest later in life, even though the mutations exist from birth. Facilitated by the inherent misfolding propensity of BVPrP and the capacity of I109 to

stabilize atypical prion species, kiBVI^{D178N} and kiBVI^{E200K} mice may effectively recapitulate

pivotal early PrP misfolding events that occur in the brain during genetic prion disease.

Declarations

Ethics approval

All mouse experiments were performed in accordance with guidelines set by the Canadian

Council on Animal Care under protocols (AUP #4263.17 and 6322.3) approved by the University

Health Network Animal Care Committee.

Availability of data and material

All data generated or analyzed during this study are included in this published article.

Competing interests

The authors have no competing interests to declare that are relevant to the content of this

article.

Funding

This work was funded by a grant from the Canadian Institutes of Health Research to JCW (PJT-

169048). SM was partially funded by a fellowship from Parkinson Canada. The funding bodies

had no role in the design of the study, the collection, analysis, or interpretation of data, or the

writing of the manuscript.

Author's contributions

All authors contributed to the study conception and design. Material preparation, data collection

and analysis were performed by Surabhi Mehra, Matthew E.C. Bourkas, Lech Kaczmarczyk,

Erica Stuart, Hamza Arshad, Jennifer K. Griffin, Kathy L. Frost, Daniel J. Walsh, Surachai

References

Agrimi U, Nonno R, Dell'Omo G, Di Bari MA, Conte M, Chiappini B *et al* (2008) Prion protein amino acid determinants of differential susceptibility and molecular feature of prion strains in mice and voles. PLoS Pathog 4: e1000113 Doi 10.1371/journal.ppat.1000113

Arshad H, Bourkas MEC, Watts JC (2020) The utility of bank voles for studying prion disease. Prog Mol Biol Transl Sci 175: 179-211 Doi 10.1016/bs.pmbts.2020.08.009

Arshad H, Patel Z, Amano G, Li LY, Al-Azzawi ZAM, Supattapone S *et al* (2023) A single protective polymorphism in the prion protein blocks cross-species prion replication in cultured cells. J Neurochem 165: 230-245 Doi 10.1111/jnc.15739

Asante EA, Gowland I, Grimshaw A, Linehan JM, Smidak M, Houghton R *et al* (2009) Absence of spontaneous disease and comparative prion susceptibility of transgenic mice expressing mutant human prion proteins. J Gen Virol 90: 546-558 Doi 10.1099/vir.0.007930-0

Asante EA, Linehan JM, Smidak M, Tomlinson A, Grimshaw A, Jeelani A *et al* (2013) Inherited prion disease A117V is not simply a proteinopathy but produces prions transmissible to transgenic mice expressing homologous prion protein. PLoS Pathog 9: e1003643 Doi 10.1371/journal.ppat.1003643

Baiardi S, Rossi M, Mammana A, Appleby BS, Barria MA, Cali I *et al* (2021) Phenotypic diversity of genetic Creutzfeldt-Jakob disease: a histo-molecular-based classification. Acta Neuropathol 142: 707-728 Doi 10.1007/s00401-021-02350-y

Bauer S, Dittrich L, Kaczmarczyk L, Schleif M, Benfeitas R, Jackson WS (2022) Translatome profiling in fatal familial insomnia implicates TOR signaling in somatostatin neurons. Life Sci Alliance 5: Doi 10.26508/lsa.202201530

Benestad SL, Sarradin P, Thu B, Schonheit J, Tranulis MA, Bratberg B (2003) Cases of scrapie with unusual features in Norway and designation of a new type, Nor98. Vet Rec 153: 202-208 Doi 10.1136/vr.153.7.202

Benilova I, Reilly M, Terry C, Wenborn A, Schmidt C, Marinho AT *et al* (2020) Highly infectious prions are not directly neurotoxic. Proc Natl Acad Sci U S A 117: 23815-23822 Doi 10.1073/pnas.2007406117

Bouybayoune I, Mantovani S, Del Gallo F, Bertani I, Restelli E, Comerio L *et al* (2015) Transgenic fatal familial insomnia mice indicate prion infectivity-independent mechanisms of pathogenesis and phenotypic expression of disease. PLoS Pathog 11: e1004796 Doi 10.1371/journal.ppat.1004796

Brandner S, Jaunmuktane Z (2017) Prion disease: experimental models and reality. Acta Neuropathol 133: 197-222 Doi 10.1007/s00401-017-1670-5

Burke CM, Mark KMK, Walsh DJ, Noble GP, Steele AD, Diack AB *et al* (2020) Identification of a homology-independent linchpin domain controlling mouse and bank vole prion protein conversion. PLoS Pathog 16: e1008875 Doi 10.1371/journal.ppat.1008875

Burke CM, Walsh DJ, Steele AD, Agrimi U, Di Bari MA, Watts JC *et al* (2019) Full restoration of specific infectivity and strain properties from pure mammalian prion protein. PLoS Pathog 15: e1007662 Doi 10.1371/journal.ppat.1007662

Carlson GA, Kingsbury DT, Goodman PA, Coleman S, Marshall ST, DeArmond S *et al* (1986) Linkage of prion protein and scrapie incubation time genes. Cell 46: 503-511 Doi 10.1016/0092-8674(86)90875-5

Chiesa R, Piccardo P, Ghetti B, Harris DA (1998) Neurological illness in transgenic mice expressing a prion protein with an insertional mutation. Neuron 21: 1339-1351 Doi 10.1016/s0896-6273(00)80653-4

Christen B, Perez DR, Hornemann S, Wuthrich K (2008) NMR Structure of the Bank Vole Prion Protein at 20 degrees C Contains a Structured Loop of Residues 165-171. Journal of Molecular Biology 383: 306-312 Doi 10.1016/j.jmb.2008.08.045

Colby DW, Prusiner SB (2011) Prions. Cold Spring Harb Perspect Biol 3: a006833 Doi 10.1101/cshperspect.a006833

Collinge J (2010) Medicine. Prion strain mutation and selection. Science 328: 1111-1112 Doi 10.1126/science.1190815

Cosseddu GM, Nonno R, Vaccari G, Bucalossi C, Fernandez-Borges N, Di Bari MA *et al* (2011) Ultra-efficient PrP(Sc) amplification highlights potentialities and pitfalls of PMCA technology. PLoS Pathog 7: e1002370 Doi 10.1371/journal.ppat.1002370

Cronier S, Gros N, Tattum MH, Jackson GS, Clarke AR, Collinge J *et al* (2008) Detection and characterization of proteinase K-sensitive disease-related prion protein with thermolysin. Biochem J 416: 297-305 Doi 10.1042/BJ20081235

Di Bari MA, Nonno R, Castilla J, D'Agostino C, Pirisinu L, Riccardi G *et al* (2013) Chronic wasting disease in bank voles: characterisation of the shortest incubation time model for prion diseases. PLoS Pathog 9: e1003219 Doi 10.1371/journal.ppat.1003219

Dossena S, Imeri L, Mangieri M, Garofoli A, Ferrari L, Senatore A *et al* (2008) Mutant prion protein expression causes motor and memory deficits and abnormal sleep patterns in a transgenic mouse model. Neuron 60: 598-609 Doi 10.1016/j.neuron.2008.09.008

Erana H, Charco JM, Di Bari MA, Diaz-Dominguez CM, Lopez-Moreno R, Vidal E *et al* (2019) Development of a new largely scalable in vitro prion propagation method for the production of infectious recombinant prions for high resolution structural studies. PLoS Pathog 15: e1008117 Doi 10.1371/journal.ppat.1008117

Erana H, Diaz-Dominguez CM, Charco JM, Vidal E, Gonzalez-Miranda E, Perez-Castro MA *et al* (2023) Understanding the key features of the spontaneous formation of bona fide prions through a novel methodology that enables their swift and consistent generation. Acta neuropathologica communications 11: 145 Doi 10.1186/s40478-023-01640-8

Espinosa JC, Nonno R, Di Bari M, Aguilar-Calvo P, Pirisinu L, Fernandez-Borges N *et al* (2016) PrPC Governs Susceptibility to Prion Strains in Bank Vole, While Other Host Factors Modulate Strain Features. J Virol 90: 10660-10669 Doi 10.1128/JVI.01592-16

Fernandez-Borges N, Di Bari MA, Erana H, Sanchez-Martin M, Pirisinu L, Parra B *et al* (2018) Cofactors influence the biological properties of infectious recombinant prions. Acta Neuropathol 135: 179-199 Doi 10.1007/s00401-017-1782-y

Friedman-Levi Y, Meiner Z, Canello T, Frid K, Kovacs GG, Budka H *et al* (2011) Fatal prion disease in a mouse model of genetic E200K Creutzfeldt-Jakob disease. PLoS Pathog 7: e1002350 Doi 10.1371/journal.ppat.1002350

Goldfarb LG, Korczyn AD, Brown P, Chapman J, Gajdusek DC (1990) Mutation in codon 200 of scrapie amyloid precursor gene linked to Creutzfeldt-Jakob disease in Sephardic Jews of Libyan and non-Libyan origin. Lancet 336: 637-638 Doi 10.1016/0140-6736(90)93443-s

Goldfarb LG, Petersen RB, Tabaton M, Brown P, LeBlanc AC, Montagna P *et al* (1992) Fatal familial insomnia and familial Creutzfeldt-Jakob disease: disease phenotype determined by a DNA polymorphism. Science 258: 806-808 Doi 10.1126/science.1439789

Hallinan GI, Ozcan KA, Hoq MR, Cracco L, Vago FS, Bharath SR *et al* (2022) Cryo-EM structures of prion protein filaments from Gerstmann-Straussler-Scheinker disease. Acta Neuropathol 144: 509-520 Doi 10.1007/s00401-022-02461-0

Hoyt F, Alam P, Artikis E, Schwartz CL, Hughson AG, Race B *et al* (2022) Cryo-EM of prion strains from the same genotype of host identifies conformational determinants. PLoS Pathog 18: e1010947 Doi 10.1371/journal.ppat.1010947

Hoyt F, Standke HG, Artikis E, Schwartz CL, Hansen B, Li K *et al* (2022) Cryo-EM structure of anchorless RML prion reveals variations in shared motifs between distinct strains. Nature communications 13: 4005 Doi 10.1038/s41467-022-30458-6

Hsiao K, Meiner Z, Kahana E, Cass C, Kahana I, Avrahami D *et al* (1991) Mutation of the prion protein in Libyan Jews with Creutzfeldt-Jakob disease. N Engl J Med 324: 1091-1097 Doi 10.1056/NEJM199104183241604

Hsiao KK, Groth D, Scott M, Yang SL, Serban H, Rapp D *et al* (1994) Serial transmission in rodents of neurodegeneration from transgenic mice expressing mutant prion protein. Proc Natl Acad Sci U S A 91: 9126-9130 Doi 10.1073/pnas.91.19.9126

Hsiao KK, Scott M, Foster D, Groth DF, DeArmond SJ, Prusiner SB (1990) Spontaneous neurodegeneration in transgenic mice with mutant prion protein. Science 250: 1587-1590 Doi 10.1126/science.1980379

Huor A, Espinosa JC, Vidal E, Cassard H, Douet JY, Lugan S *et al* (2019) The emergence of classical BSE from atypical/Nor98 scrapie. Proc Natl Acad Sci U S A 116: 26853-26862 Doi 10.1073/pnas.1915737116

Jackson GS, Linehan J, Brandner S, Asante EA, Wadsworth JDF, Collinge J (2022) Overexpression of mouse prion protein in transgenic mice causes a non-transmissible spongiform encephalopathy. Sci Rep 12: 17198 Doi 10.1038/s41598-022-21608-3

Jackson WS, Borkowski AW, Faas H, Steele AD, King OD, Watson N *et al* (2009) Spontaneous generation of prion infectivity in fatal familial insomnia knockin mice. Neuron 63: 438-450 Doi 10.1016/j.neuron.2009.07.026

Jackson WS, Borkowski AW, Watson NE, King OD, Faas H, Jasanoff A *et al* (2013) Profoundly different prion diseases in knock-in mice carrying single PrP codon substitutions associated with human diseases. Proc Natl Acad Sci U S A 110: 14759-14764 Doi 10.1073/pnas.1312006110

Kaczmarczyk L, Mende Y, Zevnik B, Jackson WS (2016) Manipulating the Prion Protein Gene Sequence and Expression Levels with CRISPR/Cas9. PLoS One 11: e0154604 Doi 10.1371/journal.pone.0154604

Kobayashi A, Matsuura Y, Takeuchi A, Yamada M, Miyoshi I, Mohri S *et al* (2019) A domain responsible for spontaneous conversion of bank vole prion protein. Brain Pathol 29: 155-163 Doi 10.1111/bpa.12638

Kraus A, Hoyt F, Schwartz CL, Hansen B, Artikis E, Hughson AG *et al* (2021) High-resolution structure and strain comparison of infectious mammalian prions. Mol Cell 81: 4540-4551 e4546 Doi 10.1016/j.molcel.2021.08.011

Kurt TD, Aguilar-Calvo P, Jiang L, Rodriguez JA, Alderson N, Eisenberg DS *et al* (2017) Asparagine and glutamine ladders promote cross-species prion conversion. J Biol Chem 292: 19076-19086 Doi 10.1074/jbc.M117.794107

Lau A, So RWL, Lau HHC, Sang JC, Ruiz-Riquelme A, Fleck SC *et al* (2020) alpha-Synuclein strains target distinct brain regions and cell types. Nat Neurosci 23: 21-31 Doi 10.1038/s41593-019-0541-x

Lau HHC, Lau A, Watts JC (2018) Discriminating Strains of Self-Propagating Protein Aggregates Using a Conformational Stability Assay. Methods Mol Biol 1777: 339-354 Doi 10.1007/978-1-4939-7811-3_22

Legname G, Nguyen HO, Peretz D, Cohen FE, DeArmond SJ, Prusiner SB (2006) Continuum of prion protein structures enciphers a multitude of prion isolate-specified phenotypes. Proc Natl Acad Sci U S A 103: 19105-19110 Doi 10.1073/pnas.0608970103

Makarava N, Baskakov IV (2013) The evolution of transmissible prions: the role of deformed templating. PLoS Pathog 9: e1003759 Doi 10.1371/journal.ppat.1003759

Manka SW, Wenborn A, Betts J, Joiner S, Saibil HR, Collinge J *et al* (2023) A structural basis for prion strain diversity. Nat Chem Biol 19: 607-613 Doi 10.1038/s41589-022-01229-7

Manka SW, Zhang W, Wenborn A, Betts J, Joiner S, Saibil HR *et al* (2022) 2.7 A cryo-EM structure of ex vivo RML prion fibrils. Nature communications 13: 4004 Doi 10.1038/s41467-022-30457-7

Manson JC, Jamieson E, Baybutt H, Tuzi NL, Barron R, McConnell I *et al* (1999) A single amino acid alteration (101L) introduced into murine PrP dramatically alters incubation time of transmissible spongiform encephalopathy. Embo Journal 18: 6855-6864 Doi DOI 10.1093/emboj/18.23.6855

Mckinley MP, Bolton DC, Prusiner SB (1983) A Protease-Resistant Protein Is a Structural Component of the Scrapie Prion. Cell 35: 57-62 Doi Doi 10.1016/0092-8674(83)90207-6

Mercer RCC, Daude N, Dorosh L, Fu ZL, Mays CE, Gapeshina H *et al* (2018) A novel Gerstmann-Straussler-Scheinker disease mutation defines a precursor for amyloidogenic 8 kDa PrP fragments and reveals N-terminal structural changes shared by other GSS alleles. PLoS Pathog 14: e1006826 Doi 10.1371/journal.ppat.1006826

Mok TH, Nihat A, Luk C, Sequeira D, Batchelor M, Mead S *et al* (2021) Bank vole prion protein extends the use of RT-QuIC assays to detect prions in a range of inherited prion diseases. Sci Rep 11: 5231 Doi 10.1038/s41598-021-84527-9

Monaco S, Fiorini M, Farinazzo A, Ferrari S, Gelati M, Piccardo P *et al* (2012) Allelic origin of protease-sensitive and protease-resistant prion protein isoforms in Gerstmann-Straussler-Scheinker disease with the P102L mutation. PLoS One 7: e32382 Doi 10.1371/journal.pone.0032382

Monari L, Chen SG, Brown P, Parchi P, Petersen RB, Mikol J *et al* (1994) Fatal familial insomnia and familial Creutzfeldt-Jakob disease: different prion proteins determined by a DNA polymorphism. Proc Natl Acad Sci U S A 91: 2839-2842 Doi 10.1073/pnas.91.7.2839

Nazor KE, Kuhn F, Seward T, Green M, Zwald D, Purro M *et al* (2005) Immunodetection of disease-associated mutant PrP, which accelerates disease in GSS transgenic mice. Embo Journal 24: 2472-2480 Doi 10.1038/sj.emboj.7600717

Nitsan Z, Cohen OS, Chapman J, Kahana E, Korczyn AD, Appel S *et al* (2020) Familial Creutzfeldt-Jakob disease homozygous to the E200K mutation: clinical characteristics and disease course. J Neurol 267: 2455-2458 Doi 10.1007/s00415-020-09826-z

Nonno R, Di Bari MA, Cardone F, Vaccari G, Fazzi P, Dell'Omo G *et al* (2006) Efficient transmission and characterization of Creutzfeldt-Jakob disease strains in bank voles. PLoS Pathog 2: e12 Doi 10.1371/journal.ppat.0020012

Nonno R, Notari S, Di Bari MA, Cali I, Pirisinu L, d'Agostino C *et al* (2019) Variable Protease-Sensitive Prionopathy Transmission to Bank Voles. Emerg Infect Dis 25: 73-81 Doi 10.3201/eid2501.180807

Orru CD, Groveman BR, Raymond LD, Hughson AG, Nonno R, Zou W *et al* (2015) Bank Vole Prion Protein As an Apparently Universal Substrate for RT-QuIC-Based Detection and Discrimination of Prion Strains. PLoS Pathog 11: e1004983 Doi 10.1371/journal.ppat.1004983

Otero A, Hedman C, Fernandez-Borges N, Erana H, Marin B, Monzon M *et al* (2019) A Single Amino Acid Substitution, Found in Mammals with Low Susceptibility to Prion Diseases, Delays Propagation of Two Prion Strains in Highly Susceptible Transgenic Mouse Models. Mol Neurobiol 56: 6501-6511 Doi 10.1007/s12035-019-1535-0

Owen JP, Maddison BC, Whitelam GC, Gough KC (2007) Use of thermolysin in the diagnosis of prion diseases. Mol Biotechnol 35: 161-170 Doi 10.1007/BF02686111

Parchi P, Chen SG, Brown P, Zou W, Capellari S, Budka H *et al* (1998) Different patterns of truncated prion protein fragments correlate with distinct phenotypes in P102L Gerstmann-Straussler-Scheinker disease. Proc Natl Acad Sci U S A 95: 8322-8327 Doi 10.1073/pnas.95.14.8322

Parchi P, Giese A, Capellari S, Brown P, Schulz-Schaeffer W, Windl O *et al* (1999) Classification of sporadic Creutzfeldt-Jakob disease based on molecular and phenotypic analysis of 300 subjects. Ann Neurol 46: 224-233

Peretz D, Scott MR, Groth D, Williamson RA, Burton DR, Cohen FE *et al* (2001) Strain-specified relative conformational stability of the scrapie prion protein. Protein Sci 10: 854-863 Doi 10.1110/ps.39201

Piccardo P, Liepnieks JJ, William A, Dlouhy SR, Farlow MR, Young K *et al* (2001) Prion proteins with different conformations accumulate in Gerstmann-Straussler-Scheinker disease caused by A117V and F198S mutations. Am J Pathol 158: 2201-2207 Doi 10.1016/S0002-9440(10)64692-5

Pirisinu L, Di Bari MA, D'Agostino C, Marcon S, Riccardi G, Poleggi A *et al* (2016) Gerstmann-Straussler-Scheinker disease subtypes efficiently transmit in bank voles as genuine prion diseases. Sci Rep 6: 20443 Doi 10.1038/srep20443

Pirisinu L, Di Bari MA, D'Agostino C, Vanni I, Riccardi G, Marcon S *et al* (2022) A single amino acid residue in bank vole prion protein drives permissiveness to Nor98/atypical scrapie and the emergence of multiple strain variants. PLoS Pathog 18: e1010646 Doi 10.1371/journal.ppat.1010646

Pirisinu L, Nonno R, Esposito E, Benestad SL, Gambetti P, Agrimi U *et al* (2013) Small ruminant nor98 prions share biochemical features with human gerstmann-straussler-scheinker disease and variably protease-sensitive prionopathy. PLoS One 8: e66405 Doi 10.1371/journal.pone.0066405

Riek R, Hornemann S, Wider G, Billeter M, Glockshuber R, Wuthrich K (1996) NMR structure of the mouse prion protein domain PrP(121-231). Nature 382: 180-182 Doi 10.1038/382180a0

Ritchie DL, Ironside JW (2017) Neuropathology of Human Prion Diseases. Prog Mol Biol Transl Sci 150: 319-339 Doi 10.1016/bs.pmbts.2017.06.011

Sandberg MK, Al-Doujaily H, Sharps B, De Oliveira MW, Schmidt C, Richard-Londt A *et al* (2014) Prion neuropathology follows the accumulation of alternate prion protein isoforms after infective titre has peaked. Nature communications 5: 4347 Doi 10.1038/ncomms5347

Schwenke KA, Walzlein JH, Bauer A, Thomzig A, Beekes M (2022) Primary glia cells from bank vole propagate multiple rodent-adapted scrapie prions. Sci Rep 12: 2190 Doi 10.1038/s41598-022-06198-4

Sigurdson CJ, Joshi-Barr S, Bett C, Winson O, Manco G, Schwarz P *et al* (2011) Spongiform encephalopathy in transgenic mice expressing a point mutation in the beta2-alpha2 loop of the prion protein. J Neurosci 31: 13840-13847 Doi 10.1523/JNEUROSCI.3504-11.2011

Sigurdson CJ, Nilsson KP, Hornemann S, Heikenwalder M, Manco G, Schwarz P *et al* (2009) *De novo* generation of a transmissible spongiform encephalopathy by mouse transgenesis. Proc Natl Acad Sci U S A 106: 304-309 Doi 10.1073/pnas.0810680105

Simon ES, Kahana E, Chapman J, Treves TA, Gabizon R, Rosenmann H *et al* (2000) Creutzfeldt-Jakob disease profile in patients homozygous for the PRNP E200K mutation. Ann Neurol 47: 257-260

Tagliavini F, Lievens PM, Tranchant C, Warter JM, Mohr M, Giaccone G *et al* (2001) A 7-kDa prion protein (PrP) fragment, an integral component of the PrP region required for infectivity, is the major amyloid protein in Gerstmann-Straussler-Scheinker disease A117V. J Biol Chem 276: 6009-6015 Doi 10.1074/jbc.M007062200

Tagliavini F, Prelli F, Ghiso J, Bugiani O, Serban D, Prusiner SB *et al* (1991) Amyloid Protein of Gerstmann-Straussler-Scheinker Disease (Indiana Kindred) Is an 11-Kd Fragment of Prion Protein with an N-Terminal Glycine at Codon-58. Embo Journal 10: 513-519 Doi 10.1002/j.1460-2075.1991.tb07977.x

Telling GC, Haga T, Torchia M, Tremblay P, DeArmond SJ, Prusiner SB (1996) Interactions between wild-type and mutant prion proteins modulate neurodegeneration in transgenic mice. Genes Dev 10: 1736-1750 Doi 10.1101/gad.10.14.1736

Telling GC, Parchi P, DeArmond SJ, Cortelli P, Montagna P, Gabizon R *et al* (1996) Evidence for the conformation of the pathologic isoform of the prion protein enciphering and propagating prion diversity. Science 274: 2079-2082 Doi 10.1126/science.274.5295.2079

Vanni I, Pirisinu L, Acevedo-Morantes C, Kamali-Jamil R, Rathod V, Di Bari MA *et al* (2020) Isolation of infectious, non-fibrillar and oligomeric prions from a genetic prion disease. Brain 143: 1512-1524 Doi 10.1093/brain/awaa078

Vidal E, Sanchez-Martin MA, Erana H, Lazaro SP, Perez-Castro MA, Otero A *et al* (2022) Bona fide atypical scrapie faithfully reproduced for the first time in a rodent model. Acta neuropathologica communications 10: 179 Doi 10.1186/s40478-022-01477-7

Walsh DJ, Rees JR, Jackson WS, Watts JC, Supattapone S (Submitted) Anti-prion drugs do not improve survival in knock-in models of inherited prion disease.

Watts JC, Giles K, Bourkas ME, Patel S, Oehler A, Gavidia M *et al* (2016) Towards authentic transgenic mouse models of heritable PrP prion diseases. Acta Neuropathol 132: 593-610 Doi 10.1007/s00401-016-1585-6

Watts JC, Giles K, Patel S, Oehler A, Dearmond SJ, Prusiner SB (2014) Evidence that bank vole PrP is a universal acceptor for prions. PLoS Pathog 10: e1003990 Doi 10.1371/journal.ppat.1003990

Watts JC, Giles K, Stohr J, Oehler A, Bhardwaj S, Grillo SK *et al* (2012) Spontaneous generation of rapidly transmissible prions in transgenic mice expressing wild-type bank vole prion protein. Proc Natl Acad Sci U S A 109: 3498-3503 Doi 10.1073/pnas.1121556109

Watts JC, Prusiner SB (2017) Experimental Models of Inherited PrP Prion Diseases. Cold Spring Harbor perspectives in medicine 7: Doi 10.1101/cshperspect.a027151

Watts JC, Prusiner SB (2014) Mouse models for studying the formation and propagation of prions. J Biol Chem 289: 19841-19849 Doi 10.1074/jbc.R114.550707

Westaway D, DeArmond SJ, Cayetano-Canlas J, Groth D, Foster D, Yang SL *et al* (1994) Degeneration of skeletal muscle, peripheral nerves, and the central nervous system in transgenic mice overexpressing wild-type prion proteins. Cell 76: 117-129 Doi 10.1016/0092-8674(94)90177-5

Westergard L, Turnbaugh JA, Harris DA (2011) A naturally occurring C-terminal fragment of the prion protein (PrP) delays disease and acts as a dominant-negative inhibitor of PrPSc formation. J Biol Chem 286: 44234-44242 Doi 10.1074/jbc.M111.286195

Williams D, Mehrabian M, Arshad H, Eid S, Sackmann C, Zhao W *et al* (2021) The cellular prion protein interacts with and promotes the activity of Na,K-ATPases. PLoS One 16: e0258682 Doi 10.1371/journal.pone.0258682

Yang W, Cook J, Rassbach B, Lemus A, DeArmond SJ, Mastrianni JA (2009) A New Transgenic Mouse Model of Gerstmann-Straussler-Scheinker Syndrome Caused by the A117V Mutation of PRNP. J Neurosci 29: 10072-10080 Doi 10.1523/JNEUROSCI.2542-09.2009

Zou WQ, Puoti G, Xiao X, Yuan J, Qing L, Cali I *et al* (2010) Variably protease-sensitive prionopathy: a new sporadic disease of the prion protein. Ann Neurol 68: 162-172 Doi 10.1002/ana.22094

Table 1. Spontaneous neurological illness in knock-in mice expressing mutant BVPrP

Line	Mean age of disease onset ± SEM (d)	Signs of neurological illness (n/n_0)	Mice showing TL-resistant PrP (n/n_0)	Mice showing PK-resistant PrP (n/n_0)
kiBVI^{wt}	> 598-603	0/22	0/21	0/2
kiBVI^{E200K}	524 ± 15	14/23	14/22	2/6
kiBVI^{D178N}	537 ± 15	15/23	22/22	7/11

n, number of mice; n_0, number of mice examined

Table 2. Transmission of brain homogenates from BVPrP knock-in mice to kiBVI^{wt} mice

Inoculum			Mean incubation period ± SEM (d)	Signs of neurological illness (n/n_0)	Mice showing TL-resistant PrP (n/n_0)	Mice showing PK-resistant PrP (n/n_0)
Line	Age (d)	Spontaneously ill?				
kiBVI^{wt}	600	No	> 540	0/8	0/6	0/6
	602	No	> 540	0/9	0/5	0/5
kiBVI^{E200K}	560	Yes	312 ± 13	8/8	8/8	8/8
	467	Yes	324 ± 23	8/8	8/8	8/8
kiBVI^{D178N}	491	Yes	248 ± 94	2/6	2/6*	2/6^
	522	Yes	287 ± 67	3/6	3/6*	2/6^

n, number of mice; n_0, number of mice examined
* 4 of the 5 brains from kiBVI^{wt} mice inoculated with kiBVI^{D178N} samples that showed TL-resistant PrP were from asymptomatic mice collected at 541 dpi
^ All of the brains from kiBVI^{wt} mice inoculated with kiBVI^{D178N} samples that showed PK-resistant PrP were from asymptomatic mice collected at 541 dpi

Figure 1. Generation and characterization of knock-in mice expressing wild-type or mutant bank vole PrP. a) Schematic of the gene-targeted alleles in knock-in mice expressing either wild-type (kiBVI^wt), E200K-mutant (kiBVI^E200K), or D178N-mutant (kiBVI^D178N) BVPrP(I109). The eight amino acid residue differences between the mature forms of bank vole and mouse PrP are shown, as are the approximate epitopes for the anti-PrP antibodies used in this study. **b**) Immunoblots for PrP in brain extracts from 3 mice each for the indicated knock-in lines. BVPrP was detected using the antibodies HuM-P and HuM-D18, and both blots were reprobed with an anti-actin antibody. **c**) ELISA-based quantification of relative BVPrP levels (mean ± SEM) in brain extracts from knock-in mice (n = 3 per line). Statistical significance was assessed using Welch ANOVA followed by Dunnett's T3 multiple comparisons test. **d**) Immunoblots for PrP in brain extracts from the indicated mouse lines probed with antibodies that recognize both mouse and bank vole PrP (POM1) or only mouse PrP (HuM-R1). Both blots were reprobed with an antibody against actin. **e**) Immunoblots for PrP in PNGase F-treated brain extracts from 3 mice each for the indicated knock-in lines. BVPrP was detected using the antibodies HuM-P and HuM-D18. Full-length BVPrP as well as the C1 and C2 endoproteolytic products are indicated.

Figure 2. Knock-in mice expressing mutant bank vole PrP develop spontaneous neurological illness. a) Kaplan-Meier curves for the development of signs of neurological illness in kiBVI^wt (black, n = 22), kiBVI^E200K (red, n = 23), and kiBVI^D178N (blue, n = 23) mice. The onset of neurological illness in kiBVI^E200K and kiBVI^D178N mice was significantly different from kiBVI^wt mice (*P* < 0.0001 as assessed by the Log-rank test), whereas there was no difference in disease onset between the kiBVI^E200K and kiBVI^D178N lines (*P* = 0.97). **b**) Analysis of sex-specific effects on the age of onset of spontaneous neurological illness (mean ± SEM) in kiBVI^E200K (left graph; n = 11 for males, n = 3 for females) and kiBVI^D178N (right graph; n = 8 for males, n = 7 for females). Statistical significance was assessed using unpaired, two-tailed Mann-Whitney tests.

Figure 3. Protease-resistant PrP in brains from spontaneously ill knock-in mice expressing mutant bank vole PrP. a) Schematic of protocols used for the analysis of detergent-insoluble and protease-resistant PrP species in the brains of knock-in mice. **b**) Immunoblots for detergent-insoluble PrP species, with (+) or without (-) digestion with 100 μg/mL thermolysin (TL), in brain homogenates from 3-month-old kiBVI[wt], kiBVI[E200K], and kiBVI[D178N] mice. **c**) Immunoblots for detergent-insoluble PrP species, with or without TL digestion, in brain homogenates from asymptomatic 20-month-old kiBVI[wt] mice and spontaneously ill kiBVI[E200K] and kiBVI[D178N] mice. The bottom panel displays a longer exposure of the blot shown in the top panel. **d**) Immunoblots for detergent-insoluble PrP species in brain extracts from aged asymptomatic kiBVI[wt] mice as well as spontaneously ill kiBVI[E200K] and kiBVI[D178N] mice following digestion with either 100 μg/mL TL (top panel) or 50 μg/mL proteinase K (PK, bottom panel). **e**) Immunoblots for detergent-insoluble, TL-resistant PrP species in brain extracts from kiBVI[wt] (left), kiBVI[E200K] (middle), and kiBVI[D178N] (right) mice at the indicated ages. Two independent mice per age are shown. For the kiBVI[wt] line, a sample without TL digestion is also shown. In all panels, PrP was detected using the antibody HuM-D13.

Figure 4. Spontaneously ill knock-in mice expressing mutant bank vole PrP exhibit prion disease-specific neuropathology. a) Representative H&E-stained brain sections of the

hippocampus, cortex, and thalamus from spontaneously ill kiBVI^{E200K} and kiBVI^{D178N} mice as well as 20-month-old asymptomatic kiBVI^{wt} mice. Scale bar = 100 μm (applies to all sections). **b)** Quantification of percent area covered by vacuolation (mean ± SEM) in the indicated brain regions from aged knock-in mice (n = 5-6 samples per line). **c)** Representative GFAP-stained brain sections of the hippocampus, cortex, and thalamus from spontaneously ill kiBVI^{E200K} and kiBVI^{D178N} mice as well as 20-month-old asymptomatic kiBVI^{wt} mice. Scale bar = 100 μm (applies to all sections). **d)** Quantification of percent area covered by GFAP staining in the indicated brain regions from aged knock-in mice. n = 5-6 samples per line. **e)** Immunoblot of GFAP levels in brain homogenates from the three lines of knock-in mice at the indicated ages. The blot was reprobed with an antibody against actin. **f)** Immunoblots of GFAP levels in brain extracts from kiBVI^{E200K} (left) and kiBVI^{D178N} (right) mice at the indicated ages. Two independent mice per age were analyzed, and the blots were reprobed with an antibody against actin. **g)** Representative PrP-stained brain sections of the striatum from spontaneously ill kiBVI^{E200K} and kiBVI^{D178N} mice as well as 20-month-old asymptomatic kiBVI^{wt} mice. Scale bar = 100 μm (applies to all sections). Molecular weight markers in panels e and f are in kDa. In panels b and d, statistical significance was assessed using a Kruskal-Wallis test followed by Dunn's multiple comparison test.

Figure 5. Transmission of prions from the brains of spontaneously ill knock-in mice expressing mutant bank vole PrP to knock-in mice expressing wild-type bank vole PrP. a) Schematic of transmission experiments in kiBVI[wt] mice. **b**) Kaplan-Meier curves for the development of signs of neurological illness in kiBVI[wt] mice inoculated with brain extract from aged, asymptomatic kiBVI[wt] mice (black, n = 17), symptomatic kiBVI[E200K] mice (red, n = 16), or symptomatic kiBVI[D178N] mice (blue, n = 13). The onset of neurological illness in mice inoculated with kiBVI[E200K] or kiBVI[D178N] extract was significantly different from mice inoculated with kiBVI[wt] extract ($P < 0.0001$ and $P = 0.0036$, respectively, as assessed by the Log-rank test). **c**) Immunoblots for detergent-insoluble PrP species in brain extracts from kiBVI[wt] mice at the indicated days post-inoculation (DPI) with brain extract from kiBVI[wt] mice, symptomatic kiBVI[E200K] mice, or symptomatic kiBVI[D178N] mice following digestion with either 100 µg/mL TL (top panel) or 50 µg/mL PK (bottom panel). TL- and PK-resistant PrP species were detected using the antibody HuM-D13. **d**) Representative H&E- and GFAP-stained sections of the hippocampus and PrP-stained sections of the striatum from kiBVI[wt] mice at the indicated DPI with brain extract from either spontaneously ill kiBVI[E200K] or kiBVI[D178N] mice or 20-month-old asymptomatic kiBVI[wt] mice. Scale bars = 100 µm. **e**) Quantification of percent area covered by vacuolation or GFAP staining in the inoculated kiBVI[wt] mice. n = 7 samples per line. Statistical significance was assessed using a Kruskal-Wallis test followed by Dunn's multiple comparison test. **f**) Immunoblot of GFAP levels in brain extracts from kiBVI[wt] mice at the indicated DPI with brain extract from kiBVI[wt] mice, symptomatic kiBVI[E200K] mice, or symptomatic kiBVI[D178N] mice.

The blot was reprobed with an antibody against actin. Molecular weight markers in panels c and f are in kDa.

Figure 6. Conformational characterization of spontaneously formed and transmitted prions. a) Representative immunoblots of detergent-insoluble, TL-resistant PrP species following treatment of brain extracts from spontaneously ill kiBVI[E200K] mice (top blot) or kiBVI[D178N] mice (bottom blot) with the indicated concentrations of guanidine hydrochloride (GdnHCl). **b**) Quantification of residual detergent-insoluble, TL-resistant PrP levels in brain extracts from spontaneously ill kiBVI[E200K] mice (red, n = 3) or kiBVI[D178N] mice (blue, n = 3) treated with the indicated concentrations of GdnHCl. **c**) Representative immunoblots of detergent-insoluble, TL-resistant PrP species in brain homogenates from kiBVI[wt] mice inoculated with either kiBVI[E200K] (top blot) or kiBVI[D178N] (bottom blot) brain extract treated with the indicated concentrations of GdnHCl. **d**) Quantification of residual detergent-insoluble, TL-resistant PrP levels in brain homogenates from kiBVI[wt] mice inoculated with brain extract from either kiBVI[E200K] (red, n = 3) or kiBVI[D178N] (blue, n = 3) mice following treatment with the indicated concentrations of GdnHCl. In panels a and c, PrP was detected using the antibody HuM-D13. In panels b and d, the calculated $[GdnHCl]_{50}$ values are also shown.

Figure 7. Model for the spontaneous generation of prions in kiBVI^E200K and kiBVI^D178N mice. (1) Properly folded PrP is present in the brains of young, healthy knock-in mice expressing mutant BVPrP(I109). (2) As the mice age, the E200K and D178N mutations promote the generation of a very similar misfolded BVPrP species. (3) Polymerization of this misfolded species results in BVPrP aggregates that are resistant to thermolysin digestion and the appearance of clinical signs of neurological illness in the knock-in mice. (4) As disease progresses, the BVPrP aggregates undergo conformational maturation to produce aggregates that contain a core region that is resistant to proteinase K digestion, producing an atypical PrP^res fragment of ~10 kDa. This PrP^res fragment does not contain either the D178N- or E200K-mutant residues.