Cardiac Specific Overexpression of Adenylyl Cyclase 8 Reprograms the Mouse Sinoatrial Node Transcriptome

Jia-Hua Qu, Kirill V. Tarasov, Khalid Chakir, and Edward G. Lakatta *

Laboratory of Cardiovascular Science, Intramural Research Program, National Institute on Aging, National Institutes of Health, Baltimore, MD 21224, USA.

Address correspondence to:

Dr. Edward G. Lakatta*

Laboratory of Cardiovascular Science

Intramural Research Program

National Institute on Aging

National Institutes of Health

251 Bayview Boulevard

Suite 100

Baltimore, MD 21224

USA.

Tel: +1 410-558-8202

lakattae@grc.nia.nih.gov

Abstract

A coupled-clock system intrinsic to sinoatrial node (SAN) pacemaker cells that regulates the rate and rhythm of spontaneous action potential firing, is activated by Ca2+/calmodulin-stimulated Adenylyl Cyclase (AC) types 8 and 1. Our previous work in mice with cardiac specific overexpression of human AC8 gene (TGAC8) discovered that compared to its wild-type (WT) littermates, the heart rate (HR) of TGAC8 is elevated (by about 30%, 24 hours a day, 7days a week), and that the TGAC8 heart rhythm is markedly coherent, i.e., the HR Variability (HRV) lacks complexity, similar to that associated with aging or cardiac pathology. Reprogramming of molecular mechanisms, particularly SAN transcriptomic regulation that underlies the remarkable chronic shift in HR and HRV, however, has not been delineated. We conducted deep RNA sequencing (RNA-seq) in TGAC8 and WT SANs, using Mm10plus with human ADCY8 DNA sequence as the reference genome. Utilizing multiple bioinformatic techniques, we not only profiled the expression of marker genes related to SAN functions and AC-cAMP-PKA signaling, but also discovered negatively enriched hub pathways that differed in TGAC8 vs. WT, the top three being OXPHOS, ribosome, and cardiac muscle contraction. In contrast, signaling pathways related to inositol phosphate and its metabolism were positively enriched in TGAC8. Further, we identified two transcription regulators, KDM5A and PPARGC1A, that mediate effects of TGAC8 on ribosome and mitochondria. In summary, reprogrammed transcriptional regulation increases the HR of TGAC8 at the cost of impairment of other SAN cell functions, i.e., altered ribosome and inositol phosphate signaling and its crosstalk with mitochondria. Because these cell signaling alterations are associated with cardiac aging and age-associated CVDs, the TGAC8 mouse appears to be

49   an ideal model in which to probe for potential therapeutic targets for cardiac aging and

50   age-associated CVDs.

Key Words

53   Sinoatrial Node, Adenylyl Cyclase 8, Transcriptome, Coupled-Clock System, Ribosome,

54   Mitochondria, Inositol Phosphate

Introduction

According to the report from World Health Organization

(https://www.who.int/cardiovascular_diseases/about_cvd/en/), cardiovascular diseases

(CVDs) are the leading cause of death globally. The decline of cardiac performance in

CVDs may result from an impaired contractile function[1,2]. As one of the most important

regulators of contraction, cAMP is catalyzed by adenylyl cyclases (ACs) upon activation

by catecholamines binding to β-adrenergic receptors (β-ARs), and degraded by

phosphodiesterases (PDEs)[3]. The activated ACs-cAMP-PKA signaling pathway promotes

the phosphorylation of key proteins, contributing to strong positive inotropic, lusitropic,

and chronotropic responses[3]. Therefore, the desensitization of the β-adrenergic signaling

pathway, with the down-regulation of β1-ARs and ACs, disrupts the excitation-

contraction coupling, resulting in CVDs[4].

Contrary to two main AC isoforms expressed in the heart, AC5 and AC6, two additional

isoforms in sinoatrial node (SAN), AC1 and AC8, are activated by Ca^{2+}/calmodulin and

insensitive to subunit of heterotrimeric G proteins (Gαs) in vitro[5,6]. In pacemaker cells,

the AC8-cAMP-PKA signaling pathway induces the phosphorylation of Ca^{2+} cycling

proteins to generate the action potential[5]. In our previous work using the mouse model

with cardiac specific overexpression of human AC8 gene (TGAC8), we have discovered

that the heart rate (HR) of TGAC8 is increased approximately 30% compared with the

wild-type (WT) littermate, and the intrinsic coupled-clock system (in the absence of β1-

ARs stimulation) within the SAN cells is crucially dependent on the activation of AC8,

but the overall pattern of marked coherency or loss of complexity (HR variability) within

the TGAC8 heart rhythm is similar to that associated with aging or cardiac pathology[7].

Although we have already profiled the changes in the TGAC8 mouse left ventricle (LV)

at RNA, protein and phosphorylation levels[8,9], and have characterized the various cellular

functions at single nucleus resolution in C57/BL6 mouse SAN[10], the molecular

mechanisms, especially the transcriptomic regulation in SAN, by which TGAC8

regulates SAN functions and its long-term effect, remain elusive.

To unravel the underlying molecular mechanisms, in the present research, we conducted

deep RNA sequencing (RNA-seq) to profile the SAN transcriptome in mouse (TGAC8

vs. WT). Utilizing multiple bioinformatic techniques, we enriched representative

pathways and discovered an unexpected downregulation of components and functions of

mitochondria and ribosome, which may be associated with CVDs in long term of chronic

activation of AC8 in the heart. Our results provide valuable transcriptomic data for

further research on the regulation in cardiac function, especially through the ACs-cAMP-

PKA signaling pathway.

Materials and Methods

1. Animals.

The cardiac-specific overexpression of AC8 (TGAC8) mouse, generated by inserting a

cDNA coding for human AC8 after the murine a-myosin heavy chain promoter[6], was a

gift from Nicole Defer/Jacques Hanoune, Unite de Recherches, INSERM U-99, Hôpital

Henri Mondor, F-94010 Créteil, France. 12-week old male TGAC mice were used as

study cases, and age matched WT littermates, in the C57/BL6 background, were used as

controls. All studies were performed in accordance with the Guide for the Care and Use

of Laboratory Animals published by the National Institutes of Health (NIH Publication

no. 85-23, revised 1996). The experimental protocols were approved by the Animal Care

and Use Committee of the National Institutes of Health (protocol #441-LCS-2016).

2. Sinoatrial node (SAN) and SAN cell isolation.

SAN and SAN cells were isolated in the same way described in our previous paper[7].

Briefly, after anesthetization of TGAC8 and WT mice, the hearts were quickly removed

and SANs were dissected between inferior and superior vena cava, crista terminalis, and

intra-atrial septum and cut into strips perpendicular to the crista terminalis. The isolated

SANs were washed (X3) in the modified low Ca^{2+} Tyrode solution before enzymatic

digestion. The digested SANs were washed (X3) in modified high potassium solution

before dispersion by gentle pipetting. The dispersed SAN cells were stored at 4°C for

following experiments.

3. RNA extraction, cDNA library preparation, and RNA-seq.

By company. We sent eight TGAC8 and eight WT mouse SANs to DNA Link USA

company to conduct total RNA-seq. RNAs were extracted from each SNAs separately

and used as template for cDNA library preparation following the instruction manual of

SMARTer Stranded Total RNA-Seq Kit-Pico/Clontech Laboratories, Lnc. The

sequencing was conducted on the Illumina NextSeq 500 platform and the data were

obtained within the basecaling software RTA2.0.

4. RNA-seq data processing.

The raw reads were trimmed within Cutadapt and aligned within SAMtools. Using

Mm10plus and ADCY8 as the reference genome, the aligned reads were mapped to

calculate the raw counts. After quality control and filtration, we selected six WT samples

(WS1, WS2, WS3, WS4, WS5 and WS 8) and five TGAC8 samples (MS1, MS2, MS4,

MS5 and MS8) for the subsequent differential expression gene (DEG) analysis using

DESeq2 package in R language, and identified 16575 genes. The normalized counts by

DESeq2 were imported into Partek Flow to perform PCA and generate expression heat

map. Furtherly, we calculated sample distances using the Euclidean method for rows and

columns distance clustering and generated matrix using the ward.D method for matrix

clustering in R language. The adjusted p-value and fold change calculated by DESeq2

were used to draw volcano plot to figure out the expression changes of TGAC8 SAN

transcriptome. Specifically, 352 genes were downregulated (-log10(adjusted p-value) >

1.3 and log2(fold change) < 0) and 323 upregulated ((-log10(adjusted p-value) > 1.3 and

log2(fold change) > 0).

5. Marker genes selection and analysis.

The expressions of several families of genes encoding proteins related to ion channels

and AC8-cAMP-PKA pathway, i.e., sodium/calcium exchanger in solute carrier family

(Slc), hyperpolarization activated cyclic nucleotide gated potassium channel (Hcn), S100

calcium binding protein (S100), sodium voltage-gated channel (Na+), potassium voltage-

gated channel (K+), calcium voltage-gated channel (Ca2+), stromal interaction molecule

(Stim), phosphodiesterase (Pde), protein kinase CAMP-dependent regulation proteins

(PKA), and sarcolipin (Sln), were extracted from TGAC8 SAN RNA-seq data and

visualized in heat map. The significant changes of these genes were further displayed in

volcano plot and bar plot. All analyses were performed and all plots were produced in R

language.

6. Ingenuity pathway analysis (IPA).

All identified genes were uploaded to IPA software (QIAGEN, March 2020)[11]. The

Ingenuity knowledge base (gene only) about all species and all tissues was selected as the

reference set. All Ingenuity supported third party information and Ingenuity expert

information were adopted as the data sources, and only experimentally observed

relationships were selected to increase confidence. Setting the cutoff as adjusted p-value

< 0.05, IPA recognized 636 differential expressed genes, including 321 down and 315 up,

which were used for subsequent canonical pathway analysis, upstream regulator

prediction, disease state analysis match, and regulatory pathway construction.

Specifically, OXPHOS pathway and apelin cardiomyocyte signaling pathway were

161   visualized, and the regulatory pathway starting from KDM5A and PPARGC1A was

162   generated in IPA.

7. KEGG pathway analysis using pathfindR package.

164
To reduce the redundancy and find the most important pathways, KEGG pathway

analysis was performed using pathfindR package in R language[12]. The KEGG mouse

(mmu) database was used as background gene set. Besides normal KEGG enrichment, we

generated term-gene graph to cluster all pathways into seven clusters and select the hub

pathway in each cluster. We also displayed the pathway level heat map for individual

sample to clarify the contribution of each sample to the pathway enrichment.

8. Gene set enrichment analysis (GSEA).

172
To take all identified genes into consideration and preclude bias due to arbitrary cutoff,

we uploaded all identified genes into GSEA software[13] for analyses within four

databases, KEGG, GO_BP, GO_CC, and GO_MF. The software can recognize 16571

native features. After collapsing features into gene symbols, there are 12873 genes

remained. The calculation parameters in GSEA were as below: (1) gene-set permutation;

(2) permutation number: 1000; (3) scoring_scheme: weighted; (4) norm: meandiv; (5)

mode: Max_probe; (6) include_only_symbols: true; (7) set_max: 500; (8) set_min: 15;

(9) metric: Signal2Noise; (10) order: descending. The top 20 significantly enriched

pathways sorted by normalized enrichment score (NES) within each database were used

for generation of bubble graphs and representative GSEA enrichment plots.

9. RNA extraction, cDNA synthesis, qRT-PCR, and data analysis.

The preparation for and performance of qRT-PCR were done as stated in our previous

paper in detail[7]. Briefly, RNAs were extracted from isolated mouse SAN with RNeasy

Mini Kit (Qiagen, Valencia, CA, United States) and DNAse. 2 mg of total RNA was used

for cDNA synthesis with MMLV reverse transcriptase (Promega) in final 50 mL volume.

Primers used for each transcript assessed are listed in Supplementary Table (in

preparation). qRT-PCR was performed on QuantStudio 6 Flex Real-Time PCR System

(Thermo Fisher Scientific) with 384-well platform. Reaction was performed with

FastStart Universal SYBR Green Master Kit with Rox (Roche). Normalized to expression

of HPRT level, qRT-PCR analysis was performed using ddCt method.

10. Statistical analyses.

Transcriptomic data were processed in Partek® Flow® software, version 9.0 Copyright ©;

2020 Partek Inc., St. Louis, MO, USA, by utilizing the computational resources of the

NIH HPC Biowulf cluster (http://hpc.nih.gov). Most statistical analyses were performed

using RStudio (version: 1.1.463) in R language (version: 3.5.3). Some other data

processing and statistics were conducted in GraphPad Prism (version: 7) and Microsoft

Excel (version: 2019). Adobe Illustrator (version: CC 2019) were also used for graphics

without changing data trend or structure. For qRT-PCR analysis, the two-sided unpaired

Student's t-test was used to compare the mRNA expression between TGAC8 and WT

SAN, and p-value < 0.05 was taken as statistically significant.

Results

1. RNA sequencing on mouse SAN tissue.

To study the molecular mechanism, we isolated SANs from twelve-week-old male

TGAC8 mice and their WT littermates to conduct deep RNA sequencing (RNA-seq). TG

and WT samples were separated into two groups in primary components analysis (PCA)

(Figure A1). Furtherly, we calculated sample distances and observed that the inner group

sample distance was smaller than inter group sample distance (Figure 1B). We also scaled

and visualized the expressions of all identified genes in heatmap, and observed two

hierarchical clusters, indicating the differential regulation between the two groups of

samples (Figure 1C). Setting the cutoff as -log10(adjusted p-value) > 1.3, i.e., adjusted p-

value < 0.05, we obtained 675 differentially regulated (up and down) genes, among

which 352 genes were downregulated (down) and 323 upregulated (up) (Figure 1D).

As SAN is the main pacemaker of the heart and AC8-cAMP-PKA axis is pivotal to

excitation-contraction coupling, we profiled the expression of several families of genes

encoding proteins related to ion channels and AC8-cAMP-PKA pathway, i.e.,

sodium/calcium exchanger in solute carrier family (Slc), hyperpolarization activated

cyclic nucleotide gated potassium channel (Hcn), S100 calcium binding protein (S100),

sodium voltage-gated channel (Na^+), potassium voltage-gated channel (K^+), calcium

voltage-gated channel (Ca^{2+}), stromal interaction molecule (Stim), phosphodiesterase

(Pde), protein kinase cAMP-dependent regulation proteins (PKA), and sarcolipin (Sln)

(Figure 2A), among which there were 13 genes regulated significantly by overexpression

of AC8 (Figure 2B). Particularly, one member in solute carrier family, Slc8a3, was

increased most significantly (Figure 1D and Figure 2C), and two members in

phosphodiesterase were also upregulated as feedback to the overexpression of AC8 to

230 finetune the content of cAMP inside SAN (Figure 2C). The characterization of RNA-seq

231 data demonstrated that the cardiac-specific overexpression of AC8 reprogramed the

232 transcriptome in SAN and regulated expression of genes related to cardiac functions.

2. Functional enrichment of TGAC8 SAN transcriptome.

To reveal the functions of the reprogramed SAN transcriptome, we performed functional

analysis using Ingenuity Pathway Analysis (IPA) software. The top 25 enriched canonical

pathways sharing same genes interacted with each other. As expected, protein kinase A

signaling was in the center of the aggregated pathways (Figure 3A), confirming the

activation of AC8-cAMP-PKA signaling pathway in our transgenic mouse model

(TGAC8). With enrichment p-value and activation prediction z-score, pathways related to

mitochondrial functions were unexpectedly enriched and ranked at the top, sorted by -

log10(p-value) (Figure 3B).

We also conducted KEGG pathway analysis using pathfindR package in R language to

study the functions of TGAC8 SAN transcriptome from another respect. Similar to

results from IPA, oxidative phosphorylation (OXPHOS) pathway was also enriched and

ranked as the top pathway among all enriched KEGG pathways (Figure 4A). Many genes

were shared by multiple pathways (Figure 4B-4C), so to reduce the redundancy and find

the most important molecular pathways, we clustered the enriched pathways into seven

clusters, from which we selected one hub pathway as the representative for each pathway

cluster (Figure 4D). The OXPHOS pathway was still the most significant pathway as the

hub in cluster 1. In addition, ribosome and cardiac muscle contraction were also enriched

and served as the hub in cluster 2 and cluster 3 respectively (Figure 4D). Interestingly,

from the term-gene graph and pathway heatmap, we observed that the majority of

associated genes were downregulated, especially genes in the top three hub pathways,

i.e., OXPHOS, ribosome, and cardiac muscle contraction (Figure 4B-4C, 4E). Therefore,

the regulation of these three pathways may be the molecular mechanism underlying the

TGAC8 SAN transcriptome, and we continued to analyze them in depth.

3. Suppressed mitochondria and mitochondrial function pathways.
The fitness of SAN relies on normal mitochondrial functions to generate enough energy

and maintain energy homeostasis. The above analyses in IPA and KEGG, with the cutoff

p-value < 0.05, had already implied the inhibition of mitochondrial function in TGAC8

mouse. To take all identified genes into consideration, we made use of gene set

enrichment analysis (GSEA). In congruent with the above analyses, many mitochondrial

components and mitochondrial functions were enriched without the requirement of any

cutoff of p-value or fold change in GSEA (Figure 5A). Regardless of the cutoff, more

genes in the gene set of OXPHOS were used for calculation and most genes were still

enriched as downregulated in TGAC8 mouse, as shown in the representative GSEA

enrichment plots (Figure 5B). Therefore, multiple functional enrichment methods,

including IPA, pathfindR, and GSEA, drew the same conclusion: suppressed

mitochondria and mitochondrial function in TGAC8 vs. WT.

To visualize individual differentially regulated gene related to mitochondria and

mitochondrial function, we visualized the OXPHOS pathway within IPA. Significantly

regulated genes (adjusted p-value < 0.05) encode subunits in OXPHOS complex I, III, IV

and V (colored with purple outlines), and all of them were downregulated (filled in green)

(Figure 5C).

To confirm the RNA-seq results, we would perform q-RT PCR to evaluate the expression

of the transcribed mRNAs. Mitochondrial function was also analyzed via WB, activity

kits, and seahorse analysis, especially for complex I, III, IV, and V.

4. Suppressed ribosome and translation pathways.

The translation process is one stage of the cell protein quality control, which consumes

large amount of energy produced in mitochondria. Mitochondria, as a semi-autonomic

organelle, consists of proteins that are synthesized in cytoplasm and mitochondria itself.

The majority of mitochondrial proteins are translated by cytosolic ribosome, and a small

portion is translated by mitochondrial ribosome. The enrichment of ribosome as the hub

in cluster 2 of KEGG pathways (Figure 4D) prompted us to further analyze ribosome and

translation. In agreement with the enrichments of both cytosolic ribosome and

mitochondrial ribosome subunits, translation inside and outside of mitochondria were

also enriched in GSEA (Figure 6A). In line with the downregulation of genes related to

ribosome that were displayed in term-gene graph (Figure 3B), the peaks in GSEA

enrichment plots of ribosome and translation were downward, indicating the suppression

of translation (Figure 6B). Particularly, the inhibited mitochondrial ribosome and

mitochondrial translation implied the possible reciprocal regulation between

mitochondria and ribosome: suppressed mitochondria produce less energy for translation,

while downregulated ribosome and weakened translation generate fewer mitochondrial

proteins.

We would perform qRT-PCR to confirm the downregulated transcription of ribosomal

components and do protein synthesis detection and WB to verify the suppressed

translation.

5. Association with impaired cardiac functions.
Intriguingly, only inositol phosphate related pathways were significantly positively

enriched within GSEA (Figure 7). As a family of calcium channels, inositol 1,4,5-

trisphosphate receptors (IP3Rs) are ubiquitously expressed in all tissues[8]. In the heart,

IP3Rs have been associated with regulation of cardiomyocyte function ranging from the

regulation of pacemaking, excitation–contraction and excitation–transcription

coupling[14,15]. The upregulated inositol phosphate increases the inner cellular Ca^{2+} to

enhance cardiac contraction[14,15], which perhaps mediates the effect of overexpression of

AC8 to increase the HR[7]. However, the abnormally activated IP3Rs signaling may

contribute to CVDs, like arrhythmias, hypertrophy and heart failure[14,15]. Mechanistically,

inositol 1,4,5-trisphosphate (IP3) is involved in the flow of Ca^{2+} between mitochondria

and endoplasmic/sarcoplasmic reticulum[16]. In cardiomyocytes, IP3 regulates

mitochondrial Ca^{2+} uptake through the mitochondrial ryanodine receptor (RyR) to

influence adenosine triphosphate (ATP) production, which may be one mechanism how

abnormal inositol phosphate metabolism impairs the cardiac function[17].

In terms of cardiac function, we enriched apelin cardiomyocyte signaling pathway in IPA

(Figure 3B). Apelin plays an important role in the regulation of cardiac functions,

including inducing vasodilation, raising contractility and HR, and reducing reperfusion

injury[18-20]. From the apelin cardiomyocyte signaling pathway generated in IPA, we

observed that the transcriptional expression of myosin light chain genes was decreased in

TGAC8 vs. WT, but the transcriptional expression of genes encoding several enzymes,

including phosphatidylinositol-4,5-bisphosphate 3-kinase, phospholipase C, and protein

kinase C, was increased in TGAC8 vs. WT (Figure 8A). To corroborate the enrichment of

KEGG pathway, cardiac muscle contraction (Figure 4), we searched the enriched

pathways within GSEA and found expression of most genes in the cardiac muscle

contraction pathway was declined and the peak was downward in TGAC8 mouse (Figure

8B). The two enrichments in IPA and KEGG prompted us to further study the possibility

of cardiac function impairment.

By the utilization of databases in IPA, we compared the TGAC8 mouse SAN

transcriptome with all available public projects and displayed the analysis match in

Figure 7C. Obviously, there were more matched disease projects with positive z-score,

indicating that TGAC8 SAN transcriptome was more comparable to/like heart disease

states than normal state (Figure 8C). Therefore, the reprogramed SAN transcriptome

implied that the chronic overexpression of AC8 in the heart increases the HR at the cost

of cardiac function and may result in/be associated with CVDs in long term.

6. Regulatory pathway of TGAC8 SAN transcriptome.

To clarify the mechanistic pathway, we performed the prediction of possible upstream

regulators mediating the effect of overexpression of AC8 to reprogram the SAN

transcriptome. As RNA-seq data mainly reflects the regulation at transcriptional level, we

first set the filter to screen all transcription regulators. Many predicted upstream

regulators were also identified as significantly changed genes in the TGAC8 SAN

transcriptome, so we set the second filter to select those genes whose transcriptional

expression was changed in the same direction to the trend of predicted activation. Finally,

we ranked all remained genes by B-H corrected p-value of prediction. Two regulators,

PPARGC1A and KDM5A, were the most significantly inhibited and activated factors

respectively. Peroxisome proliferator-activated receptor gamma coactivator 1-alpha

(PGC-1α), encoded by the *PPARGC1A* gene[21], interacts with many other regulators,

including peroxisome proliferator-activated receptor gamma (PPAR-γ or PPARG), cAMP

response element-binding protein (CREB), and nuclear respiratory factors (NRFs)[22-24], to

regulate mitochondrial biogenesis and energy metabolism[25]. Lysine-specific demethylase

5A (KDM5A) is encoded by the *KDM5A* gene[26], and its loss restores mitochondrial

function in cells lacking Rb1, partly by activating PGC-1α[27]. Therefore, we presumed the

two transcription regulators as the mediators of the effect of overexpression of AC8 in

SAN, and thus we established a regulatory pathway starting from the two regulators

within IPA (Figure 9A). In the established pathway, the majority of genes under the

control of the two regulators encode subunits in mitochondrial complex I, III, IV and V,

whose downregulations contribute to mitochondrial dysfunction and disrupted OXPHOS

process. In addition, the protein, Mitochondrial Ribosome Protein L53 (MRPL53), which

is translated from one downstream gene whose expression is under the control of

KDM5A, is a subunit of mitochondrial ribosome, and another protein, Translocase Of

Outer Mitochondrial Membrane 7 (TOMM7), which is translated from another gene, is

important in import of molecules into mitochondria. Therefore, the activated KDM5A

could also suppress other mitochondrial functions such as mitochondrial translation and

transport.

367

Discussion

369 In SAN, the ACs-cAMP-PKA signaling pathway plays an important role in

370 pacemaking[3,5]. Previously, we discovered that the cardiac-specific overexpression of AC8

371 elevates the HR, but the overall pattern of marked coherency or loss of complexity (HR

372 variability) within the TGAC8 heart rhythm is similar to that associated with aging or

373 cardiac pathology[7]. In this research, we conducted RNA-seq to analyze the SAN

374 transcriptome to reveal the underlying molecular mechanism and envisioned a possible

375 long-term effect of chronic overexpression of AC8 in the heart. As expected, we observed

376 obvious reprogramming in the SAN transcriptome (Figure 1-2). Beyond our expectation,

377 it was mitochondrial function that was enriched most significantly and consistently in all

378 analyses using multiple bioinformatic methods. We also enriched the suppressed

379 ribosome and translation, and promoted inositol metabolism. Integrating all analyses, we

380 deduced a schematic (Figure 9B). The cardiac-specific overexpression of AC8 suppresses

381 ribosome and translation, and mitochondria and their functions, while promotes inositol

382 phosphate and its metabolism. Two transcription regulators, KDM5A and PPARGC1A,

383 may mediate the effect of overexpression of AC8 on ribosome and mitochondria. The

384 abnormal ribosome and inositol phosphate could also cross talk with mitochondria to

385 disrupt mitochondrial functions. In long term, the chronic overexpression of AC8

386 increases HR overtime at the cost of impaired SAN functions, perhaps resulting

387 in/associated with CVDs.

388

1. Why does TGAC8 mouse SAN inhibit mitochondria and ribosome while activating inositol phosphate metabolism?

In our previous work, we observed the elevated HR in TGAC8 mouse relative to WT mouse[7]. The high-intensive and restless cardiac contraction lays a heavy load on energy production in mitochondria, which is the hub of energy homeostasis and metabolic homeostasis. The energy in shortage is hard to meet the demand for protein synthesis and cardiac contraction at the same time, so the translation could be sacrificed, as shown in the enrichments of suppressed ribosome components and translation in this research. The temporary or moderate decline of translation could induce several beneficial stress responses, including endoplasmic reticulum (ER) unfolded protein response (UPR) (UPR[er]) and mitochondrial UPR (UPR[mt]), to improve the protein quality control[28]. Interrogating the processed TGAC8 SAN RNA-seq data, we found some factors involved in UPR were upregulated, e.g., Atf6, Mbtps1, Mbtps2, Ern1, etc.

On the other hand, the regulated mitochondrial function is often accompanied with altered metabolism. Inositol phosphate, a type of metabolic derivative from lipids, is often localized in plasma membrane[29], where it affects Ca^{2+} release from intracellular stores[30] and regulates Ca^{2+} flow between mitochondria and endoplasmic/sarcoplasmic reticulum[16]. In the heart, the upregulated inositol phosphate increases the inner cellular Ca^{2+} to enhance cardiac contraction[14,15], which may be an additional arm to PKA pathway in mediating the effect of activation of AC8 to increase the HR[7]. Therefore, TGAC8 mouse SAN prefers to inhibit mitochondria and ribosome while activate inositol phosphate metabolism to maintain high-intensive cardiac contraction.

2. How do the regulated pathways mediate the effect of TGAC8 in long term?

The three hub pathways, mitochondrial function (Figure 5), ribosome and translation

(Figure 6), and inositol metabolism (Figure 7), were not distinct from each other.

Mitochondria not only produce energy for protein synthesis, but also control metabolism

to impact inositol phosphate indirectly. In turn, the assembly of complete functional

mitochondrial complex relies on proteins synthesized in cytosol and mitochondria. IP3

regulates mitochondrial Ca^{2+} uptake to influence ATP production[17]. Although the

temporary or moderate protein stress and mitochondrial stress are beneficial to health, the

long-term and vigorous stress is detrimental to health[31]. For example, the elevated HR

increases blood supply in response to exercise or emergency, but it aggravates the burden

of heart in the context of chronic heart diseases. Similarly, the abnormally and

persistently activated IP3Rs signaling may result in arrhythmias, hypertrophy and even

heart failure[14,15]. Under chronic CVDs states, UPR may be maladaptive[32], exemplified by

its arrhythmic effect during human heart failure by affecting cardiac ion channels

expression[33]. The persistent UPR could also be an upstream signal to promote cancer[34].

In our TGAC8 mouse model, we observed elevated HR as a chronic stress laid on mouse

heart[7]. The enriched cardiac function pathways, especially downregulated cardiac

contraction pathway, predict a coming cardiac disorder, in line with a recently published

work where aggravated age-related myocardial dysfunction was reported in TGAC8 mice

compared with their WT littermates[35]. Therefore, our results not only support the concept

of detrimental role of chronic stress, but also indicate the association of early-stage

reprogramming of SAN transcriptome with the late-stage cardiac pathology.

3. Can we regulate these pathways to ameliorate the side-effect of TGAC8 in long term?

As the robust functions of ribosome and mitochondria are pivotal to health, the ideal way

to improve cardiac fitness is to concurrently enhance cardiac contraction and biological

processes, i.e., OXPHOS, translation, and metabolism, to maintain cardiac homeostasis at

a higher level. The TGAC8 has already elevated the HR, so we need to increase the

functions of ribosome and mitochondria to maintain health. Two transcription regulators,

KDM5A and PPARGC1A (PGC-1α), were predicted to be the mediators in TGAC8 to

inhibit ribosome and mitochondria. The expression of KDM5A was reported to be

negatively correlated with subunits in mitochondrial complexes[27], and PGC-1α is well

known as a positive regulator of mitochondrial biogenesis[25]. Therefore, the two regulators

have the potential to be therapeutic targets in improving SAN functions, particularly in

aged TGAC8 mouse, or patients under similar chronic stresses.

4. Limitation.

We unraveled the reprogramed transcriptome in TGAC8 vs. WT SAN, which indicates

the suppressed components and function of ribosome and mitochondria, along with

promoted inositol phosphate metabolism, but more wet lab experiments were not

included in this research due to the scarcity of isolated SAN samples. In addition, the

complete and clear signaling pathway, from AC8 to mediators, to target genes, and to

downstream effects, need further validation. For example, the double gene mutation

mouse models, e.g., TG^{AC8}-TG^{PPARGC1A} or TG^{AC8}-KO^{KDM5A} mouse, could be used to in-

depth study the regulatory mechanisms of overexpression of AC8 in SAN and to reveal

potential therapeutic targets of chronic CVDs, especially during aging.

Acknowledgements

We thank all members in Laboratory of Cardiovascular Science, National Institute on

Aging, National Institutes of Health, for technical assistance and discussion. This work

utilized the computational resources of the NIH HPC Biowulf cluster (http://hpc.nih.gov).

Funding

Intramural Research Program, National Institute on Aging, National Institutes of Health.

Disclosures

None.

Reference

1   Chiao, Y. A. & Rabinovitch, P. S. The Aging Heart. *Cold Spring Harb Perspect Med* **5**, a025148 (2015). https://doi.org:10.1101/cshperspect.a025148

2   Strait, J. B. & Lakatta, E. G. Aging-associated cardiovascular changes and their relationship to heart failure. *Heart Fail Clin* **8**, 143-164 (2012). https://doi.org:10.1016/j.hfc.2011.08.011

3   Guellich, A., Mehel, H. & Fischmeister, R. Cyclic AMP synthesis and hydrolysis in the normal and failing heart. *Pflugers Arch* **466**, 1163-1175 (2014). https://doi.org:10.1007/s00424-014-1515-1

4   de Lucia, C., Eguchi, A. & Koch, W. J. New Insights in Cardiac beta-Adrenergic Signaling During Heart Failure and Aging. *Front Pharmacol* **9**, 904 (2018). https://doi.org:10.3389/fphar.2018.00904

5   Vinogradova, T. M. & Lakatta, E. G. Regulation of basal and reserve cardiac pacemaker function by interactions of cAMP-mediated PKA-dependent Ca2+ cycling with surface membrane channels. *Journal of molecular and cellular cardiology* **47**, 456-474 (2009). https://doi.org:10.1016/j.yjmcc.2009.06.014

6   Lipskaia, L. *et al.* Enhanced cardiac function in transgenic mice expressing a
Ca(2+)-stimulated adenylyl cyclase. *Circ. Res.* 86, 795-801 (2000).
https://doi.org:10.1161/01.res.86.7.795

7   Moen, J. M. *et al.* Overexpression of a Neuronal Type Adenylyl Cyclase (Type
8) in Sinoatrial Node Markedly Impacts Heart Rate and Rhythm. *Front
Neurosci* 13, 615 (2019). https://doi.org:10.3389/fnins.2019.00615

8   Tarasov, K. V. *et al.* A remarkable adaptive paradigm of heart performance
and protection emerges in response to the constitutive challenge of marked
cardiac-specific overexpression of adenylyl cyclase type 8. *eLife* 11, e80949
(2022). https://doi.org:10.7554/eLife.80949

9   Qu, J.-H. *et al.* Reprogramming of the cardiac phosphoproteome in
conjunction with proteome and transcriptome creates the enhanced
performance and protection circuitry in response to chronic adenylyl cyclase-
driven stress. *bioRxiv*, 2022.2004.2029.488779 (2023).
https://doi.org:10.1101/2022.04.29.488779

10   Qu, J. H., Telljohann, R., Byshkov, R. & Lakatta, E. G. Characterization of
diverse populations of sinoatrial node cells and their proliferation potential
at single nucleus resolution. *Heliyon* 9, e12708 (2023).
https://doi.org:10.1016/j.heliyon.2022.e12708

11   Kramer, A., Green, J., Pollard, J., Jr. & Tugendreich, S. Causal analysis
approaches in Ingenuity Pathway Analysis. *Bioinformatics (Oxford, England)*
30, 523-530 (2014). https://doi.org:10.1093/bioinformatics/btt703

12   Ulgen, E., Ozisik, O. & Sezerman, O. U. pathfindR: An R Package for
Comprehensive Identification of Enriched Pathways in Omics Data Through
Active Subnetworks. *Front Genet* 10, 858 (2019).
https://doi.org:10.3389/fgene.2019.00858

13   Subramanian, A. *et al.* Gene set enrichment analysis: A knowledge-based
approach for interpreting genome-wide expression profiles. *Proceedings of
the National Academy of Sciences* 102, 15545-15550 (2005).
https://doi.org:10.1073/pnas.0506580102

14   Garcia, M. I. & Boehning, D. Cardiac inositol 1,4,5-trisphosphate receptors.
Biochim Biophys Acta Mol Cell Res 1864, 907-914 (2017).
https://doi.org:10.1016/j.bbamcr.2016.11.017

15   Kockskamper, J. *et al.* Emerging roles of inositol 1,4,5-trisphosphate
signaling in cardiac myocytes. *Journal of molecular and cellular cardiology*
45, 128-147 (2008). https://doi.org:10.1016/j.yjmcc.2008.05.014

16   Jaconi, M. *et al.* Inositol 1,4,5-trisphosphate directs Ca2+ flow between
mitochondria and the endoplasmic/sarcoplasmic reticulum: A role in
regulating cardiac autonomic Ca2+ spiking. *Mol. Biol. Cell* 11, 1845-1858
(2000). https://doi.org:10.1091/mbc.11.5.1845

17   Seidlmayer, L. K. *et al.* Inositol 1,4,5-trisphosphate-mediated sarcoplasmic
reticulum-mitochondrial crosstalk influences adenosine triphosphate
production via mitochondrial Ca2+ uptake through the mitochondrial
ryanodine receptor in cardiac myocytes. *Cardiovascular research* 112, 491-
501 (2016). https://doi.org:10.1093/cvr/cvw185

18   Berry, M. F. *et al.* Apelin has in vivo inotropic effects on normal and failing
534   hearts. *Circulation* 110, Ii187-193 (2004).
535   https://doi.org:10.1161/01.CIR.0000138382.57325.5c
536   19   Ashley, E. A. *et al.* The endogenous peptide apelin potently improves cardiac
537   contractility and reduces cardiac loading in vivo. *Cardiovascular research* 65,
538   73-82 (2005). https://doi.org:10.1016/j.cardiores.2004.08.018
539   20   Szokodi, I. *et al.* Apelin, the novel endogenous ligand of the orphan receptor
540   APJ, regulates cardiac contractility. *Circ. Res.* 91, 434-440 (2002).
541   https://doi.org:10.1161/01.res.0000033522.37861.69
542   21   Esterbauer, H., Oberkofler, H., Krempler, F. & Patsch, W. Human
543   peroxisome proliferator activated receptor gamma coactivator 1 (PPARGC1)
544   gene: cDNA sequence, genomic organization, chromosomal localization, and
545   tissue expression. *Genomics* 62, 98-102 (1999).
546   https://doi.org:10.1006/geno.1999.5977
547   22   Wallberg, A. E., Yamamura, S., Malik, S., Spiegelman, B. M. & Roeder, R. G.
548   Coordination of p300-mediated chromatin remodeling and TRAP/mediator
549   function through coactivator PGC-1alpha. *Molecular cell* 12, 1137-1149
550   (2003). https://doi.org:10.1016/s1097-2765(03)00391-5
551   23   Puigserver, P. *et al.* Activation of PPARgamma coactivator-1 through
552   transcription factor docking. *Science* 286, 1368-1371 (1999).
553   https://doi.org:10.1126/science.286.5443.1368
554   24   Wu, Z. *et al.* Mechanisms controlling mitochondrial biogenesis and
555   respiration through the thermogenic coactivator PGC-1. *Cell* 98, 115-124
556   (1999). https://doi.org:10.1016/s0092-8674(00)80611-x
557   25   Dorn, G. W., 2nd, Vega, R. B. & Kelly, D. P. Mitochondrial biogenesis and
558   dynamics in the developing and diseased heart. *Genes & development* 29,
559   1981-1991 (2015). https://doi.org:10.1101/gad.269894.115
560   26   Defeo-Jones, D. *et al.* Cloning of cDNAs for cellular proteins that bind to the
561   retinoblastoma gene product. *Nature* 352, 251-254 (1991).
562   https://doi.org:10.1038/352251a0
563   27   Váraljai, R. *et al.* Increased mitochondrial function downstream from
564   KDM5A histone demethylase rescues differentiation in pRB-deficient cells.
565   *Genes & development* 29, 1817-1834 (2015).
566   https://doi.org:10.1101/gad.264036.115
567   28   Zhang, R. *et al.* Caloric Restriction Induces MicroRNAs to Improve
568   Mitochondrial Proteostasis. *iScience* 17, 155-166 (2019).
569   https://doi.org:https://doi.org/10.1016/j.isci.2019.06.028
570   29   Lyall, F. in *Basic Science in Obstetrics and Gynaecology (Fourth Edition)* (eds
571   Phillip Bennett & Catherine Williamson) 143-171 (Churchill Livingstone,
572   2010).
573   30   Hughes, A. R. & Putney, J. W., Jr. Inositol phosphate formation and its
574   relationship to calcium signaling. *Environmental health perspectives* 84, 141-
575   147 (1990). https://doi.org:10.1289/ehp.9084141
576   31   Li, X. *et al.* Targeting mitochondrial reactive oxygen species as novel therapy
577   for inflammatory diseases and cancers. *J. Hematol. Oncol.* 6, 19-19 (2013).
578   https://doi.org:10.1186/1756-8722-6-19

32 Arrieta, A., Blackwood, E. A. & Glembotski, C. C. ER Protein Quality
Control and the Unfolded Protein Response in the Heart. *Current topics in*
microbiology and immunology 414, 193-213 (2018).
https://doi.org:10.1007/82_2017_54
33 Liu, M. & Dudley, S. C., Jr. Role for the Unfolded Protein Response in Heart
Disease and Cardiac Arrhythmias. *Int. J. Mol. Sci.* 17, 52 (2015).
https://doi.org:10.3390/ijms17010052
34 Santamaría, P. G., Mazón, M. J., Eraso, P. & Portillo, F. UPR: An Upstream
Signal to EMT Induction in Cancer. *J Clin Med* 8, 624 (2019).
https://doi.org:10.3390/jcm8050624
35 Mougenot, N. *et al.* Cardiac adenylyl cyclase overexpression precipitates and
aggravates age-related myocardial dysfunction. *Cardiovascular research* 115,
1778-1790 (2019). https://doi.org:10.1093/cvr/cvy306

Figures and Legends

Figure1. Conduction of RNA sequencing on samples of TGAC8 vs. WT mouse SAN.

A-C. PCA (A), Sample distance matrix (B), and hierarchial cluster (C) of all samples

used in the TGAC8 SAN transcriptome (WT: WS1, WS2, WS3, WS4, WS5, WS8;

TGAC8: MS1, MS2, MS4, MS5, MS8). **D**. Volcano plot of TGAC8 SAN RNA-seq data.

Genes with adjusted p-value less than 0.05 are regarded as significantly regulated by the

overexpression of AC8, among which genes with log2(TG/WT) more than zero and less

than zero are marked as red (up) and blue dots (down) respectively. Other genes are

nonsignificantly regulated and marked as grey dots (ns). One of the most significantly

upregulated genes, Slc8a3, is labeled besides its dot.

616

617 **Figure 2. Marker genes in TGAC8 SAN transcriptome. A**. Heatmap of selected

618 families of genes related to ion channels and AC8-cAMP-PKA pathway. **B**. Volcano plot

619 of all selected genes in A. Red dots represent upregulated genes (adjusted p-value < 0.05

620 and log2FoldChange > 0); blue dots represent downregulated genes (adjusted p-value <

621 0.05 and log2FoldChange < 0); grey dots represent nonsignificantly regulated genes

622 (adjusted p-value ≥ 0.05). **C**. Bar plot of expressions of genes that are significantly

623 regulated in AC8 SAN (up and down).

Figure 3. Ingenuity Pathway Analysis of TGAC8 SAN transcriptome. A. Overlapping network of top 25 pathways. Three representative pathways, Oxidative Phosphorylation, Mitochondrial Dysfunction, and Protein Kinase A Signaling, are marked in blue rectangles. **B**. Top 21 canonical pathways enriched in IPA, ranked by -log10(p-value). The 20th and 21st pathways have the same -log10(p-value). Bar color indicates activation state based on the uploaded transcriptome into IPA. Orange bars with positive z-scores and blue bars with negative z-scores denote activated and inhibited pathways respectively. The darker the more significant. White bars denote pathways insignificantly activated or inhibited with z-score equal to zero. Grey bars denote pathways for which no activity pattern is available within IPA knowledgebase.

Figure 4. KEGG pathway clustering analysis of TGAC8 SAN transcriptome. A.

Bubble graph of KEGG pathway analysis of TGAC8 SAN transcriptome. X-axis is fold

enrichment of pathway, bubble size represents number of genes in a pathway, and bubble

color gradient indicates significance with -log10(p-value). **B**. Term-gene graph of top 10

terms, ranked by fold enrichment. Grey dot denotes enriched term. Red and green dots

denote upregulated and downregulated genes respectively. Some genes are enriched in

more than one pathway. The pathways on the top right are aggregated and thus zoomed in

as Figure **C**. **D**. Bubble graph of hub KEGG pathways in each pathway cluster using

pathfindR package. Legend same to Figure A. **E**. Heat map of term score in each sample.

Color gradient indicates scaled enrichment score. The more red, the more enriched.

647 **Figure 5. Downregulated mitochondria and mitochondrial functions in TGAC8**

648 **SAN. A**. Bubble graph of GSEA of TGAC8 SAN transcriptome. X-axis is normalized

649 enrichment score (NES) in GSEA, bubble size represents number of genes in a term, and

650 bubble color indicates enrichment using four different databases (KEGG, GO_CC,

651 GO_MF, GO_BP). **B**. Two representative GSEA enrichment plots. Top: enriched using

652 KEGG database; bottom: GO_BP. **C**. Visualization of changed genes in the term of

653 oxidative phosphorylation, generated in IPA. Genes with purple outline are changed

654 significantly. All genes are in green, indicating decreased.

655

656

657

658 **Figure 6. Downregulated ribosome and translation in TGAC8 SAN. A**. Bubble graph

659 of GSEA of TGAC8 SAN transcriptome. X-axis is normalized enrichment score (NES) in

660 GSEA, bubble size represents number of genes in a term, and bubble color indicates

661 enrichment using four different databases (KEGG, GO_CC, GO_MF, GO_BP). **B**. Two

662 representative GSEA enrichment plots. Top: enriched using KEGG database; bottom:

663 GO_BP.

664

665 Figure 7. Upregulated inositol phosphate and its metabolism in TGAC8 SAN. A.

666 Bubble graph of GSEA of TGAC8 SAN transcriptome. X-axis is normalized enrichment

667 score (NES) in GSEA, bubble size represents number of genes in a term, and bubble

668 color indicates enrichment using four different databases (KEGG, GO_CC, GO_MF,

669 GO_BP). **B**. Two representative GSEA enrichment plots. Top: enriched using KEGG

670 database; bottom: GO_BP.

A

Apelin Cardiomyocyte
Signaling Pathway

B

C

Ananlysis Match

673 **Figure 8. Association with impaired cardiac functions in TGAC8 SAN. A**.

674 Visualization of changed genes in the term of apelin cardiomyocyte signaling pathway,

675 generated in IPA. Genes with purple outline are changed significantly. Green represents

676 downregulated, and red represents upregulated. **B**. GSEA enrichment plot of cardiac

677 muscle contraction. **C**. Bar plot of analysis match with public projects in IPA. X-axis is z-

678 score overall score. Positive and negative scores indicate positive and negative

679 correlations respectively. Color gradient represents -log10(p-value overall score),

680 indicating the confidence of the analysis match. The darker, the high confidence. Y-axis is

681 public project's name. Blue indicates disease state, and red indicates normal state.

Figure 9

Figure 9. Regulatory pathway of TGAC8 SAN transcriptome. A. Mechanistic

pathway from predicted upstream regulators to regulated downstream pathways. Two

oval dots in the center denote the upstream regulators predicted on the basis of TGAC8

SAN transcriptome. Blue indicates inhibited, and orange indicates activated. Surrounding

dots denote genes that are under the control of the two upstream regulators and changed

significantly in TGAC8 SAN transcriptome. Green indicates downregulation, and red

indicates upregulation. Edges between dots represent relationships between genes.

Arrows denotes promotion, and short bar denotes suppression. Blue and orange edges

indicate that the predicted activation state of upstream regulator leads to the gene

downregulation and upregulation respectively in TGAC8 SAN transcriptome. Yellow

edges represent findings inconsistent with state of downstream molecule. Two text boxes

on the right represent downstream pathways that are enriched from differentially

expressed genes. **B**. Deduced schematic: Transgenic overexpression of AC8 in SAN

suppresses ribosome and translation, and mitochondria and their functions, while

promotes inositol phosphate and its metabolism, thus associated with impaired SAN

functions. Two transcription regulators, KDM5A and PPARGC1A, are possible mediators

of overexpression of AC8.