ElegansBot: Development of equation of motion deciphering locomotion including omega turns of *Caenorhabditis elegans*

Taegon Chung, Iksoo Chang, and Sangyeol Kim
Department of Brain Sciences,
Daegu Gyeongbuk Institute of Science and Technology, Korea

2023 September

Keywords

Undulatory locomotion, *C. elegans*, Kinetics, Newton's equation of motion, From synapse to behavior

Correspondence

Sangyeol Kim (Email: sykim@dgist.ac.kr, Fax: 82-053-785-1829), Department of Brain Sciences, Daegu Gyeongbuk Institute of Science and Technology, Daegu 42988, Korea

Abstract

Locomotion is a fundamental behavior of *Caenorhabditis elegans* (*C. elegans*). Previous works on kinetic simulations of animals helped researchers understand the physical mechanisms of locomotion and the muscle-controlling principles of neuronal circuits as an actuator part. It has yet to be understood how *C. elegans* utilizes the frictional forces caused by the tension of its muscles to perform sequenced locomotive behaviors. Here, we present a two-dimensional rigid body chain model for the locomotion of *C. elegans* by developing Newtonian equations of motion for each body segment of *C. elegans*. Having accounted for friction-coefficients of the surrounding environment, elastic constants of *C. elegans*, and its kymogram from experiments, our kinetic model (ElegansBot) reproduced various locomotion of *C. elegans* such as, but not limited to, forward-backward-(omega turn)-forward locomotion constituting escaping behavior and delta-turn navigation. Additionally, ElegansBot precisely quantified the forces acting on each body segment of *C. elegans* to allow investigation of the force distribution. This model will facilitate our understanding of the detailed mechanism of various locomotive behaviors at any given friction-coefficients of the surrounding environment. Furthermore, as the model ensures the performance of realistic behavior, it can be used to research actuator-controller interaction between muscles and neuronal circuits.

Introduction

With only a few hundred neurons, *Caenorhabditis Elegans*(*C. elegans*) performs various behaviors such as locomotion, sleeping, reproduction, and hunting (Hall & Altun, 2008). The connectome structure among 302 neurons and 165 somatic cells of *C. elegans* was discovered by pioneering works (Cook et al., 2019; White et al., 1986). *C. elegans* is a cost-efficient and widely-used model in neuronal research. Its small body size and minimal nutritional requirements contribute to its cost efficiency. The organism matures in a shorter period, about three days, compared to other model animals such as fruit flies or mice. Its transparent body allows for easy microscopic observation of its internal structures or artificially expressed green fluorescent proteins. Moreover, due to the hermaphroditic nature of *C. elegans*, offspring mostly share the same genotype as the parent, which simplifies the multiplication of the worm population for research purposes (Hall & Altun, 2008).

C. elegans bends its body with a sinusoidal wave pattern when moving forward or backward. The driving force for this movement comes from the difference between perpendicular and parallel frictional forces, which it experienced from a surrounding environment. This thrust force pushes the worm along the ground surface with which the worm contacts(Boyle, 2009; Boyle et al., 2012; Hu et al., 2009; Niebur & Erdös, 1991). Even if a worm has a sinusoidal modulation generated inside it, it has difficulties in forward and backward locomotion if it does not feel the difference in frictional forces from its surroundings.

Kinetic simulators of rod-shaped animals such as *C. elegans* (Boyle et al., 2012; Niebur & Erdös, 1991), fish (Ekeberg, 1993), and snakes (Hu et al., 2009) have been used in various studies. These simulators demonstrate how the activities of muscle cells are represented as behavioral phenotypes, which are determined by signals from a neuronal circuit simulator (Boyle et al., 2012; Ekeberg, 1993; Niebur & Erdös, 1991). They also show how muscle cells return proprioceptive signals back to the neuronal circuit simulator and how animals intentionally distribute body weight for locomotion patterns (Hu et al., 2009). Similarly, the kinetic simulator of fish(Ekeberg, 1993), which has a locomotion pattern in that the animal mostly undulates in a particular direction, was used with a neuronal network simulator to model the undulation of swimming behavior. This combination of kinetic simulator and neuronal network simulator was also used to model how the locomotion pattern changes due to a surrounding environment(Boyle et al., 2012) and how the central pattern generator arises from a few cells(Boyle et al., 2012; Izquierdo & Beer, 2018).

Even though there were studies on kinetic simulation of rod-shaped animals
(Boyle et al., 2012; Ekeberg, 1993; Hu et al., 2009), to our best knowledge, there was
no kinetic model that reproduces complex locomotion behavior of *C. elegans,* which
includes all of the various modes of locomotion of *C. elegans* such as forward
locomotion, backward locomotion, and turn from experimental observations. Instead,
muscle cell activities from Ansatz (Hu et al., 2009), a hypothesis of the solution, or
signals from a neuronal circuit simulator (Boyle et al., 2012; Ekeberg, 1993) were
applied to the kinetic simulators. A simulator should have an operational structure
that imitates physical quantities from an experimental video to reproduce the motion
of *C. elegans*. However, until now, no kinetic simulation has such a structure. If there
is a simulator that reproduces the motion of individual experiments, analysis of the
kinetics of motion of specific experiments, which provides information on the
individual force that exerts on each body part of the animal, will be enabled. Also, as
the kinetic simulation reproduces the motion of *C. elegans*, the behavioral phenotype
that emerged from the muscle activity of neuronal circuit simulation will be more
credible.

We built a Newtonian-mechanics 2-dimensional rigid body chain model of *C.
elegans* to reproduce its locomotion. We incorporated its body angle, related to the
contraction of the body wall muscle of *C. elegans*, into the primary operating principle
of our kinetic model so that the model simulates measurable physical quantities of *C.
elegans* from its experimental video. The model includes a chain of multiple rod rigid
bodies, a damped torsional spring between the rigid bodies, and a control angle,
which is the dynamic baseline angle from the value of the kymogram of a physical
experiment. We formulated Newtonian equations of translational and rotational
motion of the rigid body model and computed the numerical solution of the equations
by numerical integration using semi-implicit Euler method. As a result, we were able
to demonstrate trajectories and kinetics of the general locomotion of *C. elegans*, such
as crawling, swimming (Vidal-Gadea et al., 2011), omega-turn, and delta-turn
(Broekmans et al., 2016).

Results

Newton's equation of motion for locomotion of *Caenorhabditis elegans:* How does EleganBot work? We introduce the simple chain model of *C. elegans*' body. *C. elegans* has an elongated body along the head-to-tail axis. Thus, the worm's body can be approximated as a midline extended along the anterial-posterial axis in the xy-coordinate plane (Fig. 1A). Let M (=2 μg) be the mass and L (=1 mm) be the length of the worm. Midline was approximated as n (=25) straight rods, whose ends are connected to the ends of neighboring rods (Fig. 1A). The mass, length, and moment of inertia of each rod is $m = M/n$, $2r = L/n$, and $I = mr^2/3$ respectively. When numbering the rods in order, with the rod at the end of the head being labeled as "1-rod" and the rod at the end of the tail being labeled as "n-rod", let us designate the i-th rod as the "i-rod". The point where i-rod and (i+1)-rod meets is "i-joint".

The motion of the worm corresponds to the motion of all the rods. To describe the motion of each rod (i-rod), we need to determine the displacement vector ($\mathbf{d}_i$), velocity vector ($\mathbf{v}_i$), the angle measured counterclockwise from the positive x-axis to the tangential direction of the rod (s_i) (Fig. 1B), and angular velocity (ω_i) of i-rod at a given time t. However, the minimum information required to describe the motion of all rods includes the displacement vector ($\mathbf{d}_c$) and velocity vector ($\mathbf{v}_c$) of the worm's center of mass, s_i and ω_i for each rod (Details in Supplementary Information).

Value of time-dependent variable such as $\mathbf{d}_c$, $\mathbf{v}_c$, s_i, and ω_i at given time, t will be expressed as $*^{(t)}$. When initial values, $\mathbf{d}_c^{(0)}$, $\mathbf{v}_c^{(0)}$, $s_i^{(0)}$, and $\omega_i^{(0)}$ are given, the Newtonian equation of motion for acceleration, $\mathbf{a}_c$ and angular acceleration, $\{\alpha_i\}_{i \in \{1, \cdots, n\}}$ must be acquired and numerically integrated twice to find $\mathbf{d}_c^{(t)}$, $\mathbf{v}_c^{(t)}$, $s_i^{(t)}$, and $\omega_i^{(t)}$ at a given time, t. To obtain the Newtonian equations of motion, we must find every force and torque acting on each rod. There are frictional force, muscle force, and joint force among types of forces acting on the rod, and there are frictional torque, muscle torque, and joint torque among types of torques whose descriptions are as follows.

The only external force acting on the worm is frictional force from a ground surface such as an agar plate or water. The frictional force is an anisotropic Stokes frictional force, with a magnitude proportional to the speed and assumed different friction coefficients in perpendicular and parallel directions (Boyle et al., 2012), which guarantees that linearity in velocity is preserved in frictional force as well (Details in Supplementary Information). Because of this preservation of linearity, the

frictional forces of translational motion (Fig. 1D) and rotational motion (Fig. 1E) can be calculated separately and added together to find total frictional force and torque. Previously known values of the friction coefficients in perpendicular and parallel directions are used (Boyle et al., 2012).

Figure 1. Components of ElegansBot. (A) Chain model for *C. elegans* body. (B) Rods in chain model. (C) i-actuator, which is a damped torsional spring. (D) Frictional force (black arrow) due to the translation motion of a rod. (E) Frictional force (black arrow) due to the rotational motion of a rod. (F) Joint force (black arrows) acting on i-rod and (i+1)-rod.

Let the perpendicular and parallel friction coefficients be $b_\perp$ and $b_\parallel$ for a straightened worm, respectively. Each rod experiences 1/n of the frictional force the worm gets. Thus, the perpendicular and parallel friction coefficients of each rod are $b_\perp/n$, $b_\parallel/n$, respectively. The ratio of perpendicular friction coefficient to parallel friction coefficient ($b_\perp/b_\parallel$) is 40 in agar plate and 1.5 in water (Boyle, 2009; Boyle et al., 2012). This ratio is an important determining factor in whether the locomotion would be crawling or swimming(Boyle et al., 2012). The total frictional force that i-rod receives is $\mathbf{F}_{b,i} = -\frac{b_\parallel}{n}(\mathbf{v}_i \cdot \hat{\mathbf{r}}_i)\hat{\mathbf{r}}_i - \frac{b_\perp}{n}(\mathbf{v}_i \cdot \hat{\mathbf{N}}_i)\hat{\mathbf{N}}_i$ ("·": dot product of vectors, $\hat{\mathbf{r}}_i \equiv [\cos(s_i)\quad \sin(s_i)]^\mathsf{T}$: unit vector parallel to i-rod, $\hat{\mathbf{N}}_i \equiv [-\sin(s_i)\quad \cos(s_i)]^\mathsf{T}$: unit vector perpendicular to i-rod), and the total frictional torque that i-rod receives is

$\tau_{b,i} = -\frac{1}{3}\frac{b_\perp}{n}r^2\omega_i$ (positive or negative values are for torque pointing away from or
into the paper plane, respectively.) (The proof is in Supplementary Information).

Mature hermaphrodite *C. elegans* has four muscle strands at the left dorsal,
right dorsal, left ventral, and right ventral part of the body, and each muscle strand
has 24, 24, 23, and 24 muscle cells, respectively (White et al., 1986). Muscle cells at
similar positions on the anterior-posterior axis have an activity pattern in that
muscles on one side (either dorsal or ventral) cooperate, and those on the opposite
side have alternative activities. (Hall & Altun, 2008).

Therefore we modeled a group of about four muscle cells, which are left dorsal,
right dorsal, left ventral, and right ventral, at the same position on the anterior-
posterior axis as one actuator (Fig. 1C) so that there are a total of 24 ($\simeq$
$(24 + 24 + 23 + 24)/4$) actuators in the worm. On i-joint of the chain, there is an
actuator labeled as i-actuator. As the number of actuators is 24, we set the number of
rod(n) as 25, which is one more than the number of actuators. The actuator was
modeled as a damped torsional spring due to the viscoelastic characteristics of
muscle(Boyle et al., 2012; Hill, 1937). If the dorsal muscles of i-actuator contract more
than the ventral muscles, i-actuator will bend to the dorsal direction and vice versa.
To express this phenomenon by an equation, we defined the torque that i-actuator
exerts on i-rod as $\tau_i = \tau_{\kappa,i} + \tau_{c,i}$ that the elastic term is $\tau_{\kappa,i} = \kappa\left(\theta_i - \theta_{\text{ctrl},i}\right)$ and the
damping term is $\tau_{c,i} = c(\omega_{i+1} - \omega_i)$ where $\theta_{\text{ctrl},i}$ is control angle, $\theta_i = s_{i+1} - s_i$, and
κ and c are the elasticity and damping coefficients of an actuator, respectively.

Control angle ($\theta_{\text{ctrl},i}$) is a variable inside the elastic part of the muscle torque
($\tau_{\kappa,i}$), to which $\tau_{\kappa,i}$ drives θ_i close. Also, control angle ($\theta_{\text{ctrl},i}$), which can be expressed
by a heatmap (Fig. 2A, 2C, 3A, 3B), is an input value based on experimental data, a
numerical model, or a neuronal network model. $\tau_{c,i}$ represents the damping effect of
muscle cells and somatic cells near i-actuator. The elasticity coefficient (κ) and
damping coefficient (c) of an actuator were induced from previously known
values(Boyle et al., 2012) (Details in Supplementary Information).

By assuming that i-rod receives torque (τ_i) from i-actuator, and (i+1)-rod
receives torque ($-\tau_i$), we can depict the bending that arises from the differential
contraction of dorsal and ventral muscles in i-actuator. The total muscle torque that
i-rod receives from damped torsional springs on both ends is $\tau_{c\kappa,i} = \tau_i - \tau_{i-1}$. The
total muscle force ($\mathbf{F}_{c\kappa,i}$) that i-rod receives from both of its ends is as follows (Details
in Supplementary Information, Fig. S2).

$$\mathbf{F}_{c\kappa,i} = \sum_{j=-1}^{0} \frac{(-1)^{j-1}\tau_{i+j}\sin\frac{\theta_{i+j}}{2}}{r\cos^2\frac{\theta_{i+j}}{2}} \begin{bmatrix} \cos\left(\frac{s_{i+j}+s_{i+j+1}}{2}\right) \\ \sin\left(\frac{s_{i+j}+s_{i+j+1}}{2}\right) \end{bmatrix}$$

Two neighboring rods (i-rod and (i+1)-rod) are connected at i-joint. Therefore, when a force is applied to i-rod, (i+1)-rod also receives distributed force (Fig. 1F) which we name as "joint force". The joint force that (i+1)-rod exerts on i-rod is symbolized as $\mathbf{F}_i \equiv \mathbf{F}_{(i+1)i}$. By Newton's third law of motion about action and reaction, the joint force that i-rod exerts on (i+1)-rod is $\mathbf{F}_{i(i+1)} = -\mathbf{F}_{(i+1)i} = -\mathbf{F}_i$ (Fig. 1F). Joint force ($\mathbf{F}_i$) can be calculated from the previously introduced given values (s_i, $\mathbf{F}_{c\kappa,i}$, $\mathbf{F}_{b,i}$, $\tau_{c\kappa,i}$, $\tau_{b,i}$) (Details in Supplementary Information). When $\mathbf{F}_0 = \mathbf{F}_n = \mathbf{0}$, then the total joint force that i-rod receives is $\mathbf{F}_{\text{joint},i} = \mathbf{F}_i - \mathbf{F}_{i-1}$ and the total torque caused by joint force is $\boldsymbol{\tau}_{\text{joint},i} = [\mathbf{r}_i \times (\mathbf{F}_i + \mathbf{F}_{i+1})]$ where "$\times$" between two vectors means cross product.

As all forces and torques are found, $\mathbf{d}_c^{(t)}$, $\mathbf{v}_c^{(t)}$, $s_i^{(t)}$, $\omega_i^{(t)}$ can be calculated by solving translational and rotational Newtonian equations of motion with numerical integration. The time-step (Δt) used in this work is 10^{-5} sec. Because the only external force exerts on the worm is the frictional force, the equation of translational motion is $\mathbf{a}_c = \frac{\sum_i \mathbf{F}_{b,i}}{M}$. If friction coefficients ($b_\perp$, $b_\parallel$) are significantly greater than $\frac{M}{\Delta t}$, numerical integration using explicit Euler method ($\mathbf{v}_c^{(t+\Delta t)} = \mathbf{v}_c^{(t)} + \mathbf{a}_c^{(t)}\Delta t = \mathbf{v}_c^{(t)} + \frac{\sum_i \mathbf{F}_{b,i}^{(t)}}{M}\Delta t$) becomes unstable (Butcher, 2004). So, we tackled this instability of numerical integration by developing semi-implicit Euler Method ($\mathbf{v}_c^{(t+\Delta t)} = \mathbf{v}_c^{(t)} + \mathbf{a}_c^{(t+\Delta t)}\Delta t \simeq \mathbf{v}_c^{(t)} + \frac{1}{1+\frac{b_\perp \Delta t}{M}}\frac{\sum_i \mathbf{F}_{b,i}^{(t)}}{M}\Delta t$), which makes numerical integration stable when any frictional coefficients greater than or equal to 0 is given (Details in Supplementary Information).

The equation of rotational motion of i-rod is $I\alpha_i = \tau_{\text{total},i} = \tau_{b,i} + \tau_{c\kappa,i} + \tau_{\text{joint},i}$. When the friction-related value ($b_\parallel r^2$), elasticity-related value ($\kappa\Delta t$), or damping coefficient(c) is significantly larger than $\frac{I}{\Delta t}$, numerical integration using explicit Euler method ($\omega_i^{(t+\Delta t)} = \omega_i^{(t)} + \alpha_i^{(t)}\Delta t = \omega_i^{(t)} + \frac{\tau_{\text{total},i}^{(t)}}{I}\Delta t$) becomes unstable (Butcher, 2004). To solve this instability, we constructed a semi-implicit Euler method for

rotational motion and an error-corrected equation for angular momentum (Details in
Supplementary Information). By using these semi-implicit Euler methods, solutions
for $\mathbf{d}_c^{(t)}$, $\mathbf{v}_c^{(t)}$, $s_i^{(t)}$, $\omega_i^{(t)}$ of a worm at a given time can be available for the ground surface
of agar whose $b_\perp$, $b_\parallel$ are significantly larger than $\frac{M}{\Delta t}$, water which has smaller friction
coefficients than agar, or frictionless ground surface.

Can *C. elegans* in ElegansBot crawl or swim? A kymogram is a heatmap that shows
body angle, $\theta_i^{(t)}$ ($i \in \{1, \cdots, n-1\}$) at a given time, t. By fitting a sine function to the
kymogram of previous work (Vidal-Gadea et al., 2011), we obtained linear-
wavenumber(after this referred to as wavenumber) and period of *C. elegans* crawling
on the agar plate and swimming in water. The wavenumber (v) and the period (T)
are, respectively, 1.832 and 1.6 (sec) on the agar plate and 0.667 and 0.4 (sec) in water.
For both crawling and swimming, amplitude (A) was set to 0.6 (rad) arbitrarily to
match the trajectory shown in the experimental video (Vidal-Gadea et al., 2011). Each
kymogram of crawling (Fig. 2A) and swimming (Fig. 2C) was calculated by
substituting amplitude (A), wavenumber (v), and period (T) into $\theta_{\mathrm{ctrl},i}^{(t)} =$
$A\cos\big(2\pi(v(i-1)/(n-2) - t/T)\big)$.

Crawling trajectory, which performs sinusoidal locomotion in the positive x-
axis direction, was obtained by inputting a crawling kymogram as $\theta_{\mathrm{ctrl},i}^{(t)}$ input to
ElegansBot (Fig. 2B). Regarding crawling, the head track and the tail track have
similar shapes. However, the tail track is more toward the negative x-axis direction
than the head track. The difference between the head and tail tracks indicates that the
worm pushes the ground surface by the distance between the head track and tail
track to obtain thrust (Fig. 2B). Indeed, we found that the body part placed diagonally
with respect to the direction of the worm's locomotion is pushing along the ground
surface (Fig. S3A). The thrust force of the worm cancels out most of the drag force,
which enables the worm to move at nearly constant velocity. The average velocity of
the worm is 0.208 (mm/sec), which is consistent with the known values (Cohen et al.,
2012; Jung et al., 2016; Omura et al., 2012; Shen et al., 2012).

Figure 2. Simulated locomotion from a sine kymogram. (A) Crawling kymogram. Kymogram indicates the angle of i-joint which is located between i-rod and (i+1)-rod. Red and blue color mean i-joint bend in the dorsal and ventral directions, respectively. (B) Crawling trajectory. The yellow circle indicates the position of the worm's head. The Orange and sky-blue lines show the worm's head and tail trajectories, respectively. (C) Swimming kymogram. (D) Swimming trajectory

In the previous work, the worm showed swimming behavior in a water droplet on an agar plate (Vidal-Gadea et al., 2011). As the friction coefficient of water is smaller than that of agar, even though the area that the worm swept was wider during swimming than crawling, the worm did not move forward much in comparison to the area it swept (Fig. 2D). The worm gained significant momentum in the forward direction of locomotion when the body bent in the c-shape (Fig. S3B). In contrast to crawling, during swimming, the worm did not receive constant thrust force over time. Thus, the speed of the worm exhibited significant oscillations over time (Fig. S3B), and the average velocity was 0.223 (mm/sec).

ElegansBot exhibits more complex behavior including the turn motion. Unlike previous *C. elegans* body kinetic simulation studies, our simulation can replicate the worm's behavior using a kymogram (Fig. 3A, 3B) derived from experimental videos. We utilized open-source software, Tierpsy Tracker (Javer et al., 2018) and WormPose (Hebert et al., 2021), to obtain the kymogram input for the ElegansBot. Through simulation, we aimed to reproduce the omega-turn and delta-turn behaviors

observed in the experimental videos (Broekmans et al., 2016). When we used the
vertical and horizontal friction coefficients $b_\perp$ and $b_\parallel$ on agar, as proposed in the
paper (Boyle et al., 2012), the trajectory was not accurately replicated. Given that the
friction coefficients could vary depending on the concentration of the agar gel, we
used $b_\perp/100$ and $b_\parallel/100$ for the vertical and horizontal friction coefficients,
respectively, which resulted in a better trajectory replication.

The trajectory (Fig. 3C, 3D, Video S1, S2) obtained from ElegansBot accurately
reproduces the experimental video (Broekmans et al., 2016). The changes in the
direction of movement caused by turns are well replicated. Additionally, during the
omega-turn or delta-turn, the body briefly performs a deep bend, and we newly
discovered the mechanism that gains significant propulsion from the deep bend
region to change direction using ElegansBot (Fig. 3C, 3D). Moreover, the ElegansBot
accurately reproduces not only the turns but also complex behaviors like the
sequence of forward-backward-turn-forward, also known as escaping behavior.
296

Figure 3. Simulated locomotion from a kymogram of a real worm locomotion video. The length and direction of a black arrow indicate the magnitude and direction of the frictional force that the corresponding body part, which is the starting point of the arrow, exerts on the surface. (A) escaping behavior kymogram. Triangles over the heatmap indicate the corresponding time of snapshots shown in Figure (C). (B) Delta-turn kymogram. Triangles over the heatmap indicate the corresponding time of snapshots shown in Figure (D). (C) Escaping behavior trajectory. (D) Delta-turn trajectory. (Those videos are in Supplementary Information)

ElegansBot presents body shape ensembles of *C. elegans* from a shape in water en route to agar. While there have been studies on how locomotion patterns change in agar and water by merging neural and kinetic simulations (Boyle et al., 2012), there have been none that solely used kinetic simulation to analyze how speed manifests depending on the frequency and period of locomotion. We demonstrate this aspect. We studied the locomotion speed of the worm under different friction coefficients, which represent the influence of water, agar, and intermediate frictional environment, using ElegansBot. The vertical and horizontal friction coefficients in water are $b_{\text{water},\perp} = 5.2 \times 10^3$ (μg/sec) and $b_{\text{water},\parallel} = b_{\text{water},\perp}/1.5$, respectively, while in agar, these values are $b_{\text{agar},\perp} = 1.28 \times 10^8$ (μg/sec) and $b_{\text{agar},\parallel} = b_{\text{agar},\perp}/40$ (Boyle et al., 2012). For environmental index $\sigma \in$ [0,1], we have defined the vertical and horizontal friction coefficients in the environment between water ($\sigma = 0$) and agar ($\sigma = 1$) as $b_{\sigma,\perp} = b_{\text{water},\perp}^{1-\sigma} b_{\text{agar},\perp}^{\sigma}$ and $b_{\sigma,\parallel} = b_{\text{water},\parallel}^{1-\sigma} b_{\text{agar},\parallel}^{\sigma}$, respectively.

Under an environmental index σ, for various pairs of frequency-period (ν, T) when the control angle is $\theta_{\text{ctrl},i}^{(t)} = A\cos\left(2\pi\left(\nu\frac{i-1}{23} - \frac{t}{T}\right)\right)$ (with A = 0.6 (rad)), we have found the (ν, T) that maximizes the worm's average speed(optimal (ν, T)) (Fig. 4A). The optimal (ν, T) exhibits nearly linear distribution (Fig. 4B). We noticed a transition from swimming body shape to crawling body shape as σ varies (Fig. 4, S4). The optimal (ν, T) for σ=0(water) is (0.65, 0.4 sec), matching the actual (ν, T) value of swimming behavior (Vidal-Gadea et al., 2011). The optimal (ν, T) for σ=1(agar) is (1.9, 0.8 sec), and the optimal ν (1.9) matches the actual ν value (1.832) for crawling behavior (Vidal-Gadea et al., 2011), with the optimal T (0.8 sec) being half the actual T value (1.6 sec).

Figure 4. Body shape transition from the shape in water to the shape in agar.
(A) Average speed of the worm as a function of wavenumber (v) and period (T) for a given friction coefficient. The star symbol indicates the pair of (v, T) that maximizes the worm's average speed. (B) For each environmental index σ, the pair of (v, T) that maximizes the worm's average speed. The worm figures inside the small rectangles pointed to by the arrows represent the body shape corresponding to the respective (v, T) pair.

Discussion

ElegansBot is an advanced kinetic simulator that reproduces *C. elegans*' various locomotion. The known crawling speed range of *C. elegans* (Cohen et al., 2012; Jung et al., 2016; Omura et al., 2012; Shen et al., 2012) matches the speed in our simulation. The force dispersion pattern of the forward movement of a snake (Hu et al., 2009) is similar to the force dispersion pattern of crawling in our model, where the body part placed diagonally to the direction of movement generates thrust. The head and tail tracks of our simulation resemble the trace left on the agar plate by *C. elegans* during locomotion (Yeon et al., 2018), providing evidence of the mechanism where *C. elegans* moves forward by pushing along the ground surface. Given that friction and elasticity coefficients can vary between experiments, the appropriate selection of these values allows the trajectories of omega-turns and delta-turns in our simulations to match

the experimental videos (Broekmans et al., 2016). Previous work (Boyle, 2009; Boyle
et al., 2012) eliminated inertia from the equations of motion, but our simulation
includes it, allowing calculation even in cases where inertia is significant due to low
friction coefficients. Using the crawling and swimming wavenumbers and periods
from the experiments (Vidal-Gadea et al., 2011), we computed sine functions to create
trajectories for crawling and swimming. We also analyzed how friction forces act on
the worm during crawling and swimming, studying how the worm gains propulsion.
We demonstrated that we could reproduce various locomotion observed in
experimental videos, such as forward-backward-omega turn-forward constituting
escaping behavior and delta-turn navigation, by providing the kymogram obtained
from representative physical values from the experimental videos, as well as the
kymogram obtained from a program (Hebert et al., 2021; Javer et al., 2018) extracting
the body angles from actual experimental videos into ElegansBot. Our established
Newtonian equations of motion are accurate and robust, suggesting that not only
does our simulation replicate the experimental videos, but it also provides credible
estimates for detailed forces.
370
ElegansBot will serve as a strong bridge for enhancing the knowledge in "from-
synapse-to-behavior" research. Our method could be used for kinetic analysis of
behaviors not covered in this paper. It could also be used when analyzing behavior
changes caused by mutation or ablation experiments. Given that our simulation
allows for kinetic analysis, it could be used to calculate the energy expended by the
worm during locomotion, serving as an activity index. Our simulation only requires
the body angles as input data, so even if the video angle shifts and trajectory
information is lost, the trajectory can be recovered from the kymogram. Our
simulation could also be used when studying neural circuit models of *C. elegans*. It
could be used to check how signals from neural network models manifest as
behaviors which is a needed function from previous work (Sakamoto et al., 2021),
and it could be used when studying compound models of neural circuits and bodies.
For example, when creating models that receive proprioception input based on body
shape (Boyle et al., 2012; Ekeberg, 1993; Izquierdo & Beer, 2018; Niebur & Erdös,
1991), our method could be used. Finally, our method could be used in general for the
broad utility to analyze the motion of rod-shaped animals like snakes or eels and to
simulate the motion of rod-shaped robots.

Methods

1. Frequency and wavelength of *C. elegans* locomotion

Sine function fitting was applied to the crawling and swimming kymograms (Vidal-Gadea et al., 2011) to determine the frequency and wavelength of *C. elegans* locomotion on agar and water.

2. *C. elegans* locomotion videos

Videos of *C. elegans*' escaping behavior and foraging behavior were obtained from previous work (Broekmans et al., 2016). A single representative video out of a total of one hundred escaping behavior videos was used as data in this paper. Additionally, only the delta-turning portion of the foraging behavior videos was cut out and used as data in this paper.

3. Obtaining kymograms from video

The following method was used to extract the kymogram from the video of *C. elegans*: The body angles and midline were extracted using Tierpsy Tracker (Javer et al., 2018) from the original video where the worm is locomoting. Tierpsy Tracker failed to extract the midline of the worm when the body parts meet or the worm is coiled. The midline information of the frames successfully predicted by Tierpsy Tracker and the original video information were used as ground truth training data for a program called WormPose (Hebert et al., 2021). WormPose trained an artificial neural network to extract the body angles and midline of a coiled worm using a generative method based on the input data. The body angles were extracted from the original video using the trained WormPose program. For the frames where Tierpsy Tracker failed to extract the body angles, it was replaced with the body angles extracted by WormPose. Nonetheless, there were frames where the body angle extraction failed. If the period of failed body angle prediction was continuously less than three frames (about 0.01 seconds), the body angles for that period was predicted using linear interpolation.

4. Program code and programming libraries

Equations for the chain model, friction model, muscle model, and numerical integration that constitute the ElegansBot were designed from the body angle information and kymogram that change every moment of time. Python (van

Rossum, 1995) version 3.8 was used to implement the equations constituting ElegansBot as a program. NumPy (Harris et al., 2020) version 1.19 was used for numerical calculations, and Numba (Lam et al., 2015) version 0.54 was used for CPU calculation acceleration. The Matplotlib (Hunter, 2007) library was used to represent *C. elegans*' body pose and trajectory in figures and videos. The program code used in the research can be obtained from the open database GitHub (https://github.com/taegonchung/elegansbot), and the demo can be found at the laboratory website (https://biophysics.dgist.ac.kr/elegansbot/demo). This code calculated the ElegansBot simulation of 10 seconds of simulation time in about 10 seconds of run-time on an Intel E3-1230v5 CPU.

5. **Physical constants of the ground surface**

The friction coefficient values for the ground surface where *C. elegans* crawled and swam and the elastic and damping coefficients of *C. elegans* muscles were obtained from previous work (Boyle et al., 2012). The muscle elasticity and damping coefficients were converted into coefficients for the damped torsional spring to be used in our model (Details in Supplementary Information).

References

Boyle, J. H. (2009). *C. elegans locomotion: An integrated approach*.

Boyle, J. H., Berri, S., & Cohen, N. (2012). Gait Modulation in C. elegans: An Integrated Neuromechanical Model. *Frontiers in Computational Neuroscience*, *6*. https://doi.org/10.3389/fncom.2012.00010

Broekmans, O. D., Rodgers, J. B., Ryu, W. S., & Stephens, G. J. (2016). Resolving coiled shapes reveals new reorientation behaviors in C. elegans. *ELife*, *5*, e17227. https://doi.org/10.7554/eLife.17227

Butcher, J. C. (2004). *Numerical Methods for Ordinary Differential Equations*. John Wiley & Sons.

Cohen, E., Yemini, E., Schafer, W., Feitelson, D. G., & Treinin, M. (2012). Locomotion analysis identifies roles of mechanosensory neurons in governing locomotion dynamics of C. elegans. *Journal of Experimental Biology*, *215*(20), 3639–3648. https://doi.org/10.1242/jeb.075416

Cook, S. J., Jarrell, T. A., Brittin, C. A., Wang, Y., Bloniarz, A. E., Yakovlev, M. A., Nguyen,
K. C. Q., Tang, L. T.-H., Bayer, E. A., Duerr, J. S., Bülow, H. E., Hobert, O., Hall, D.
H., & Emmons, S. W. (2019). Whole-animal connectomes of both
Caenorhabditis elegans sexes. *Nature*, *571*(7763), 63–71.
https://doi.org/10.1038/s41586-019-1352-7

Ekeberg, O. (1993). A combined neuronal and mechanical model of fish swimming.
Biological Cybernetics, 12.

Hall, D. H., & Altun, Z. F. (2008). *C. elegans Atlas*.

Harris, C. R., Millman, K. J., van der Walt, S. J., Gommers, R., Virtanen, P., Cournapeau,
D., Wieser, E., Taylor, J., Berg, S., Smith, N. J., Kern, R., Picus, M., Hoyer, S., van
Kerkwijk, M. H., Brett, M., Haldane, A., del Río, J. F., Wiebe, M., Peterson, P., …
Oliphant, T. E. (2020). Array programming with NumPy. *Nature*, *585*(7825),
Article 7825. https://doi.org/10.1038/s41586-020-2649-2

Hebert, L., Ahamed, T., Costa, A. C., O'Shaughnessy, L., & Stephens, G. J. (2021).
WormPose: Image synthesis and convolutional networks for pose estimation
in C. elegans. *PLOS Computational Biology*, *17*(4), e1008914.
https://doi.org/10.1371/journal.pcbi.1008914

Hill, A. V. (1937). The heat of shortening and the dynamic constants of muscle.
Proceedings of the Royal Society of London. Series B - Biological Sciences,
126(843), 136–195. https://doi.org/10.1098/rspb.1938.0050

Hu, D. L., Nirody, J., Scott, T., & Shelley, M. J. (2009). The mechanics of slithering
locomotion. *Proceedings of the National Academy of Sciences*, *106*(25), 10081–
10085. https://doi.org/10.1073/pnas.0812533106

Hunter, J. D. (2007). Matplotlib: A 2D Graphics Environment. *Computing in Science &*
Engineering, *9*(3), 90–95. https://doi.org/10.1109/MCSE.2007.55

Izquierdo, E. J., & Beer, R. D. (2018). From head to tail: A neuromechanical model of
forward locomotion in Caenorhabditis elegans. *Philosophical Transactions of*
the Royal Society B: Biological Sciences, *373*(1758), 20170374.
https://doi.org/10.1098/rstb.2017.0374

Javer, A., Currie, M., Lee, C. W., Hokanson, J., Li, K., Martineau, C. N., Yemini, E.,
Grundy, L. J., Li, C., Ch'ng, Q., Schafer, W. R., Nollen, E. A. A., Kerr, R., & Brown,
A. E. X. (2018). An open-source platform for analyzing and sharing worm-
behavior data. *Nature Methods*, *15*(9), Article 9.
https://doi.org/10.1038/s41592-018-0112-1

Jung, J., Nakajima, M., Takeuchi, M., Najdovski, Z., Huang, Q., & Fukuda, T. (2016).
Microfluidic Device to Measure the Speed of C. elegans Using the Resistance

Change of the Flexible Electrode. *Micromachines*, *7*(3), 50.

https://doi.org/10.3390/mi7030050

Lam, S. K., Pitrou, A., & Seibert, S. (2015). Numba: A LLVM-based Python JIT

compiler. *Proceedings of the Second Workshop on the LLVM Compiler*

Infrastructure in HPC, 1–6. https://doi.org/10.1145/2833157.2833162

Niebur, E., & Erdös, P. (1991). Theory of the locomotion of nematodes: Dynamics of

undulatory progression on a surface. *Biophysical Journal*, *60*(5), 1132–1146.

https://doi.org/10.1016/S0006-3495(91)82149-X

Omura, D. T., Clark, D. A., Samuel, A. D. T., & Horvitz, H. R. (2012). Dopamine

Signaling Is Essential for Precise Rates of Locomotion by C. elegans. *PLOS*

ONE, *7*(6), e38649. https://doi.org/10.1371/journal.pone.0038649

Sakamoto, K., Soh, Z., Suzuki, M., Iino, Y., & Tsuji, T. (2021). Forward and backward

locomotion patterns in C. elegans generated by a connectome-based model

simulation. *Scientific Reports*, *11*(1), Article 1.

https://doi.org/10.1038/s41598-021-92690-2

Shen, X. N., Sznitman, J., Krajacic, P., Lamitina, T., & Arratia, P. E. (2012). Undulatory

Locomotion of Caenorhabditis elegans on Wet Surfaces. *Biophysical Journal*,

102(12), 2772–2781. https://doi.org/10.1016/j.bpj.2012.05.012

van Rossum, G. (1995). *Python reference manual* (Issue R 9525). CWI.

Vidal-Gadea, A., Topper, S., Young, L., Crisp, A., Kressin, L., Elbel, E., Maples, T.,

Brauner, M., Erbguth, K., Axelrod, A., Gottschalk, A., Siegel, D., & Pierce-

Shimomura, J. T. (2011). Caenorhabditis elegans selects distinct crawling and

swimming gaits via dopamine and serotonin. *Proceedings of the National*

Academy of Sciences, *108*(42), 17504–17509.

https://doi.org/10.1073/pnas.1108673108

White, J. G., Southgate, E., Thomson, J. N., & Brenner, S. (1986). The Structure of the

Nervous System of the Nematode Caenorhabditis elegans. *Philosophical*

Transactions of the Royal Society of London. Series B, Biological Sciences,

314(1165,), 1–340.

Wood, W. B. (Ed.). (1988). *The Nematode Caenorhabditis elegans*.

Yeon, J., Kim, J., Kim, D.-Y., Kim, H., Kim, J., Du, E. J., Kang, K., Lim, H.-H., Moon, D., &

Kim, K. (2018). A sensory-motor neuron type mediates proprioceptive

coordination of steering in C. elegans via two TRPC channels. *PLOS Biology*,

16(6), e2004929. https://doi.org/10.1371/journal.pbio.2004929

Acknowledgement

This work is supported by the p-CoE program of DGIST 23-CoE-BT-01. We also acknowledge Prof. Kyuhyung Kim for fruitful discussions on experiments of C. elegans.