ElegansBot: Development of equation of motion deciphering locomotion including omega turns of *Caenorhabditis elegans*

Taegon Chung, Iksoo Chang, and Sangyeol Kim
Department of Brain Sciences,
Daegu Gyeongbuk Institute of Science and Technology, Korea

2024 January

Keywords

Undulatory locomotion, *C. elegans*, Kinetics, Newton's equation of motion, From synapse to behavior

Correspondence

Sangyeol Kim (Email: sykim@dgist.ac.kr, Fax: 82-053-785-1829), Department of Brain Sciences, Daegu Gyeongbuk Institute of Science and Technology, Daegu 42988, Korea

Abstract

Locomotion is a fundamental behavior of *Caenorhabditis elegans* (*C. elegans*). Previous works on kinetic simulations of animals helped researchers understand the physical mechanisms of locomotion and the muscle-controlling principles of neuronal circuits as an actuator part. It has yet to be understood how *C. elegans* utilizes the frictional forces caused by the tension of its muscles to perform sequenced locomotive behaviors. Here, we present a two-dimensional rigid body chain model for the locomotion of *C. elegans* by developing Newtonian equations of motion for each body segment of *C. elegans*. Having accounted for friction-coefficients of the surrounding environment, elastic constants of *C. elegans*, and its kymogram from experiments, our kinetic model (ElegansBot) reproduced various locomotion of *C. elegans* such as, but not limited to, forward-backward-(omega turn)-forward locomotion constituting escaping behavior and delta-turn navigation. Additionally, ElegansBot precisely quantified the forces acting on each body segment of *C. elegans* to allow investigation of the force distribution. This model will facilitate our understanding of the detailed mechanism of various locomotive behaviors at any given friction-coefficients of the surrounding environment. Furthermore, as the model ensures the performance of realistic behavior, it can be used to research actuator-controller interaction between muscles and neuronal circuits.

Introduction

With only a few hundred neurons, *Caenorhabditis Elegans*(*C. elegans*) performs various behaviors such as locomotion, sleeping, reproduction, and hunting (Hall & Altun, 2008). The connectome structure among 302 neurons and 165 somatic cells of *C. elegans* was discovered by pioneering works (Cook et al., 2019; White et al., 1986). *C. elegans* is a cost-efficient and widely-used model in neuronal research. Its small body size and minimal nutritional requirements contribute to its cost efficiency. The organism matures in a shorter period, about three days, compared to other model animals such as fruit flies or mice. Its transparent body allows for easy microscopic observation of its internal structures or artificially expressed green fluorescent proteins. Moreover, due to the hermaphroditic nature of *C. elegans*, offspring mostly share the same genotype as the parent, which simplifies the multiplication of the worm population for research purposes (Hall & Altun, 2008).

C. elegans bends its body with a sinusoidal wave pattern when moving forward or backward. The driving force for this movement comes from the difference between perpendicular and parallel frictional forces, which it experienced from a surrounding environment. This thrust force pushes the worm along the ground surface with which the worm contacts(Boyle, 2009; Boyle et al., 2012; Hu et al., 2009; Niebur & Erdös, 1991). Even if a worm has a sinusoidal modulation generated inside it, it has difficulties in forward and backward locomotion if it does not feel the difference in frictional forces from its surroundings.

Mechanical simulators of rod-shaped animals such as *C. elegans* (Boyle et al., 2012; Niebur & Erdös, 1991), fish (Ekeberg, 1993), and snakes (Hu et al., 2009) have been used in various studies. These simulators demonstrate how the activities of muscle cells are represented as behavioral phenotypes, which are determined by signals from a neuronal circuit simulator (Boyle et al., 2012; Ekeberg, 1993; Niebur & Erdös, 1991). They also show how muscle cells return proprioceptive signals back to the neuronal circuit simulator and how animals intentionally distribute body weight for locomotion patterns (Hu et al., 2009). Similarly, the kinematic simulator of fish(Ekeberg, 1993), which has a locomotion pattern in that the animal mostly undulates in a particular direction, was used with a neuronal network simulator to model the undulation of swimming behavior. This combination of kinematic simulator and neuronal network simulator was also used to model how the locomotion pattern changes due to a surrounding environment(Boyle et al., 2012) and how the central pattern generator arises from a few cells(Boyle et al., 2012; Izquierdo & Beer, 2018).

Even though there were studies on kinematic simulation of rod-shaped
animals (Boyle et al., 2012; Ekeberg, 1993; Hu et al., 2009), to our best knowledge,
there was no kinetic model that reproduces complex locomotion behavior of *C.*
elegans, which includes all of the various modes of locomotion of *C. elegans* such as
forward locomotion, backward locomotion, and turn from experimental observations.
Instead, muscle cell activities from Ansatz (Hu et al., 2009), a hypothesis of the
solution, or signals from a neuronal circuit simulator (Boyle et al., 2012; Ekeberg,
1993) were applied to the kinematic simulators. A simulator should have an
operational structure that imitates physical quantities from an experiment to
reproduce the motion of *C. elegans* of the experiment. However, until now, no kinetic
simulation has such a structure. If there is a simulator that reproduces the motion of
individual experiments, analysis of the kinetics of motion of specific experiments,
which provides information on the individual force that exerts on each body part of
the animal, will be enabled. Also, as the kinetic simulation reproduces the motion of
C. elegans, the behavioral phenotype that emerged from the muscle activity of
neuronal circuit simulation will be more credible.

We built a Newtonian-mechanics 2-dimensional rigid body chain model of *C.*
elegans to reproduce its locomotion. We incorporated its body angle, related to the
contraction of the body wall muscle of *C. elegans*, into the primary operating principle
of our kinetic model so that the model simulates measurable physical quantities of *C.*
elegans from its experimental video. The model includes a chain of multiple rod rigid
bodies, a damped torsional spring between the rigid bodies, and a control angle,
which is the dynamic baseline angle from the value of the kymogram of a physical
experiment. We formulated Newtonian equations of translational and rotational
motion of the rigid body model and computed the numerical solution of the equations
by numerical integration using semi-implicit Euler method. As a result, we were able
to demonstrate trajectories and kinetics of the general locomotion of *C. elegans*, such
as crawling, swimming (Vidal-Gadea et al., 2011), omega-turn, and delta-turn
(Broekmans et al., 2016).

Results

Newton's equation of motion for locomotion of *Caenorhabditis elegans:* How does ElegansBot work? We introduce the simple chain model of *C. elegans*' body. *C. elegans* has an elongated body along the head-to-tail axis. Thus, the worm's body can be approximated as a midline extended along the anterial-posterial axis in the xy-coordinate plane (Fig. 1A). Let M (=2 μg, details in "Worm's mass, actuator elasticity coefficient, and damping coefficient" of Supplementary Information) be the mass and L (=1 mm) be the length of the worm. Midline was approximated as n (=25) straight rods, whose ends are connected to the ends of neighboring rods (Fig. 1A). The mass, length, and moment of inertia of each rod is $m = M/n$, $2r = L/n$, and $I = mr^2/3$ respectively. When numbering the rods in order, with the rod at the end of the head being labeled as "1-rod" and the rod at the end of the tail being labeled as "n-rod", let us designate the i-th rod as "i-rod". The point where i-rod and (i+1)-rod meets is "i-joint".

The motion of the worm corresponds to the motion of all the rods. To describe the motion of each rod (i-rod), we need to determine the displacement vector ($\mathbf{d}_i$), velocity vector ($\mathbf{v}_i$), the angle measured counterclockwise from the positive x-axis to the tangential direction of the rod (s_i) (Fig. 1B), and angular velocity (ω_i) of i-rod at a given time t. However, the minimum information required to describe the motion of all rods includes the displacement vector ($\mathbf{d}_c$) and velocity vector ($\mathbf{v}_c$) of the worm's center of mass, s_i and ω_i for each rod (Details in "Minimum information required to describe the motion of each rod" of Supplementary Information, Fig. S1).

Value of time-dependent variable such as $\mathbf{d}_c$, $\mathbf{v}_c$, s_i, and ω_i at given time, t will be expressed as $*^{(t)}$. When initial values, $\mathbf{d}_c^{(0)}$, $\mathbf{v}_c^{(0)}$, $s_i^{(0)}$, and $\omega_i^{(0)}$ are given, the Newtonian equation of motion for acceleration, $\mathbf{a}_c$ and angular acceleration, $\{\alpha_i\}_{i \in \{1, \cdots, n\}}$ must be acquired and numerically integrated twice to find $\mathbf{d}_c^{(t)}$, $\mathbf{v}_c^{(t)}$, $s_i^{(t)}$, and $\omega_i^{(t)}$ at a given time, t. To obtain the Newtonian equations of motion, we must find every force and torque acting on each rod. There are frictional force, muscle force, and joint force among types of forces acting on the rod, and there are frictional torque, muscle torque, and joint torque among types of torques whose descriptions are as follows.

The only external force acting on the worm is frictional force from a ground surface such as an agar plate or water. The frictional force is an anisotropic Stokes frictional force, with a magnitude proportional to the speed and assumed different

friction coefficients in perpendicular and parallel directions (Boyle et al., 2012), which guarantees that linearity in velocity is preserved in frictional force as well (Details in "Preservation of linearity in friction" of Supplementary Information). Because of this preservation of linearity, the frictional forces of translational motion (Fig. 1D) and rotational motion (Fig. 1E) can be calculated separately and added together to find total frictional force and torque. Previously known values of the friction coefficients in perpendicular and parallel directions are used (Boyle et al., 2012).

Figure 1. Components of ElegansBot. (A) Chain model for *C. elegans* body. (B) Rods in chain model. (C) i-actuator, which is a damped torsional spring. (D) Frictional force (black arrow) due to the translation motion of a rod. (E) Frictional force (black arrow) due to the rotational motion of a rod. (F) Joint force (black arrows) acting on i-rod and (i+1)-rod.

Let the perpendicular and parallel friction coefficients be $b_\perp$ and $b_\parallel$ for a straightened worm, respectively. Each rod experiences 1/n of the frictional force the worm gets. Thus, the perpendicular and parallel friction coefficients of each rod are $b_\perp/n$, $b_\parallel/n$, respectively. The ratio of perpendicular friction coefficient to parallel friction coefficient ($b_\perp/b_\parallel$) is 40 in agar plate and 1.5 in water (Boyle, 2009; Boyle et al., 2012). This ratio is an important determining factor in whether the locomotion would be crawling or swimming(Boyle et al., 2012). The total frictional force that i-

rod receives is $\mathbf{F}_{b,i} = -\frac{b_\parallel}{n}(\mathbf{v}_i \cdot \hat{\mathbf{r}}_i)\hat{\mathbf{r}}_i - \frac{b_\perp}{n}(\mathbf{v}_i \cdot \widehat{\mathbf{N}}_i)\widehat{\mathbf{N}}_i$ (".": dot product of vectors, $\hat{\mathbf{r}}_i \equiv$

$[\cos(s_i)\quad \sin(s_i)]^\top$: unit vector parallel to i-rod, $\widehat{\mathbf{N}}_i \equiv [-\sin(s_i)\quad \cos(s_i)]^\top$: unit

vector perpendicular to i-rod), and the total frictional torque that i-rod receives is

$\tau_{b,i} = -\frac{1}{3}\frac{b_\perp}{n}r^2\omega_i$ (positive or negative values are for torque pointing away from or

into the paper plane, respectively.) (The proof is in "Frictional torque by rotational

motion" of Supplementary Information).

Mature hermaphrodite *C. elegans* has four muscle strands at the left dorsal,

right dorsal, left ventral, and right ventral part of the body, and each muscle strand

has 24, 24, 23, and 24 muscle cells, respectively (White et al., 1986). Muscle cells at

similar positions on the anterior-posterior axis have an activity pattern in that

muscles on one side (either dorsal or ventral) cooperate, and those on the opposite

side have alternative activities. (Hall & Altun, 2008).

Therefore, we modeled a group of about four muscle cells, which are left dorsal,

right dorsal, left ventral, and right ventral, at the same position on the anterior-

posterior axis as one actuator (Fig. 1C) so that there is a total of 24 ($\simeq$

$(24 + 24 + 23 + 24)/4$) actuators in the worm. On i-joint of the chain, there is an

actuator labeled as i-actuator. As the number of actuators is 24, we set the number of

rod(n) as 25, which is one more than the number of actuators. The actuator was

modeled as a damped torsional spring due to the viscoelastic characteristics of

muscle(Boyle et al., 2012; Hill, 1937). If the dorsal muscles of i-actuator contract more

than the ventral muscles, i-actuator will bend to the dorsal direction and vice versa.

To express this phenomenon by an equation, we defined the torque that i-actuator

exerts on i-rod as $\tau_i = \tau_{\kappa,i} + \tau_{c,i}$ that the elastic term is $\tau_{\kappa,i} = \kappa(\theta_i - \theta_{\text{ctrl},i})$ and the

damping term is $\tau_{c,i} = c(\omega_{i+1} - \omega_i)$ where $\theta_{\text{ctrl},i}$ is control angle, $\theta_i = s_{i+1} - s_i$, and

κ and c are the elasticity and damping coefficients of an actuator, respectively.

Control angle ($\theta_{\text{ctrl},i}$) is a variable inside the elastic part of the muscle torque

($\tau_{\kappa,i}$), to which $\tau_{\kappa,i}$ drives θ_i close. Also, control angle ($\theta_{\text{ctrl},i}$), which can be expressed

by a heatmap (Fig. 2A, 2C, 3A, 3B), is an input value based on experimental data, a

numerical model, or a neuronal network model. $\tau_{c,i}$ represents the damping effect of

muscle cells and somatic cells near i-actuator. The elasticity coefficient (κ) and

damping coefficient (c) of an actuator were induced from previously known

values(Boyle et al., 2012) (Details in "Worm's mass, actuator elasticity coefficient, and

damping coefficient" of Supplementary Information).

By assuming that i-rod receives torque (τ_i) from i-actuator, and (i+1)-rod

receives torque ($-\tau_i$), we can depict the bending that arises from the differential

contraction of dorsal and ventral muscles in i-actuator. The total muscle torque that
i-rod receives from damped torsional springs on both ends is $\tau_{c\kappa,i} = \tau_i - \tau_{i-1}$. The
total muscle force ($\mathbf{F}_{c\kappa,i}$) that i-rod receives from both of its ends is as follows (Details
in "Proof of muscle force" of Supplementary Information, Fig. S2).

207
$$\mathbf{F}_{c\kappa,i} = \sum_{j=-1}^{0} \frac{(-1)^{j-1}\tau_{i+j}\sin\dfrac{\theta_{i+j}}{2}}{r\cos^2\dfrac{\theta_{i+j}}{2}} \begin{bmatrix} \cos\left(\dfrac{s_{i+j}+s_{i+j+1}}{2}\right) \\ \sin\left(\dfrac{s_{i+j}+s_{i+j+1}}{2}\right) \end{bmatrix}$$

Two neighboring rods (i-rod and (i+1)-rod) are connected at i-joint. Therefore,
when a force is applied to i-rod, (i+1)-rod also receives distributed force (Fig. 1F)
which we name as "joint force". The joint force that (i+1)-rod exerts on i-rod is
symbolized as $\mathbf{F}_i \equiv \mathbf{F}_{(i+1)i}$. By Newton's third law of motion about action and reaction,
the joint force that i-rod exerts on (i+1)-rod is $\mathbf{F}_{i(i+1)} = -\mathbf{F}_{(i+1)i} = -\mathbf{F}_i$ (Fig. 1F). Joint
force ($\mathbf{F}_i$) can be calculated from the previously introduced given values (s_i, $\mathbf{F}_{c\kappa,i}$, $\mathbf{F}_{b,i}$,
$\tau_{c\kappa,i}$, $\tau_{b,i}$) (Details in "Joint force calculation method" of Supplementary Information).
When $\mathbf{F}_0 = \mathbf{F}_n = \mathbf{0}$, then the total joint force that i-rod receives is $\mathbf{F}_{\text{joint},i} = \mathbf{F}_i - \mathbf{F}_{i-1}$
and the total torque caused by joint force is $\boldsymbol{\tau}_{\text{joint},i} = [\mathbf{r}_i \times (\mathbf{F}_i + \mathbf{F}_{i+1})]$ where "$\times$"
between two vectors means cross product.

As all forces and torques are found, $\mathbf{d}_c^{(t)}$, $\mathbf{v}_c^{(t)}$, $s_i^{(t)}$, $\omega_i^{(t)}$ can be calculated by
solving translational and rotational Newtonian equations of motion with numerical
integration. The time-step (Δt) used in this work is 10^{-5} sec unless otherwise noted.
Because the only external force exerts on the worm is the frictional force, the equation
of translational motion is $\mathbf{a}_c = \frac{\sum_i \mathbf{F}_{b,i}}{M}$. If friction coefficients ($b_\perp$, $b_\parallel$) are significantly
greater than $\frac{M}{\Delta t}$, numerical integration using explicit Euler method ($\mathbf{v}_c^{(t+\Delta t)} = \mathbf{v}_c^{(t)} +$
$\mathbf{a}_c^{(t)}\Delta t = \mathbf{v}_c^{(t)} + \frac{\sum_i \mathbf{F}_{b,i}^{(t)}}{M}\Delta t$) becomes unstable (Butcher, 2004). So, we tackled this
instability of numerical integration by developing semi-implicit Euler Method
($\mathbf{v}_c^{(t+\Delta t)} = \mathbf{v}_c^{(t)} + \mathbf{a}_c^{(t+\Delta t)}\Delta t \simeq \mathbf{v}_c^{(t)} + \frac{1}{1+\frac{b_\perp \Delta t}{M}}\frac{\sum_i \mathbf{F}_{b,i}^{(t)}}{M}\Delta t$), which makes numerical
integration stable when any frictional coefficients greater than or equal to 0 is given
(Details in "Proof of numerical integration for the translational motion of a worm
using semi-implicit Euler method" of Supplementary Information).

The equation of rotational motion of i-rod is $I\alpha_i = \tau_{\text{total},i} = \tau_{b,i} + \tau_{c\kappa,i} + \tau_{\text{joint},i}$.
When the friction-related value ($b_\parallel r^2$), elasticity-related value ($\kappa \Delta t$), or damping
coefficient(c) is significantly larger than $\frac{I}{\Delta t}$, numerical integration using explicit Euler
method ($\omega_i^{(t+\Delta t)} = \omega_i^{(t)} + \alpha_i^{(t)} \Delta t = \omega_i^{(t)} + \frac{\tau_{\text{total},i}^{(t)}}{I} \Delta t$) becomes unstable (Butcher,
2004). To solve this instability, we constructed a semi-implicit Euler method for
rotational motion and an error-corrected equation for angular momentum (Details in
"Numerical integration of the rotational motion of i-rod using semi-implicit Euler
method" and "Correction formula for the rotational inertia of the entire worm" of
Supplementary Information). By using these semi-implicit Euler methods, solutions
for $\mathbf{d}_c^{(t)}, \mathbf{v}_c^{(t)}, s_i^{(t)}, \omega_i^{(t)}$ of a worm at a given time can be available for the ground surface
of agar whose $b_\perp$, $b_\parallel$ are significantly larger than $\frac{M}{\Delta t}$, water which has smaller friction
coefficients than agar, or frictionless ground surface.

Can *C. elegans* in ElegansBot crawl or swim? A kymogram is a heatmap that shows
body angle, $\theta_i^{(t)}$ ($i \in \{1, \cdots, n-1\}$) at a given time, t. By fitting a sine function to the
kymogram of previous work (Vidal-Gadea et al., 2011), we obtained linear-
wavenumber(after this referred to as wavenumber) and period of *C. elegans* crawling
on the agar plate and swimming in water. The wavenumber (ν) and the period (T)
are, respectively, 1.832 and 1.6 (sec) on the agar plate and 0.667 and 0.4 (sec) in water.
For both crawling and swimming, amplitude (A) was set to 0.6 (rad) arbitrarily to
match the trajectory shown in the experimental video (Vidal-Gadea et al., 2011). Each
kymogram of crawling (Fig. 2A) and swimming (Fig. 2C) was calculated by
substituting amplitude (A), wavenumber (ν), and period (T) into　$\theta_{\text{ctrl},i}^{(t)} =$
$A\cos\big(2\pi(\nu(i-1)/(n-2) - t/T)\big)$.

Crawling trajectory, which performs sinusoidal locomotion in the positive x-
axis direction, was obtained by inputting a crawling kymogram as $\theta_{\text{ctrl},i}^{(t)}$ input to
ElegansBot (Fig. 2B). Regarding crawling, the head track and the tail track have
similar shapes. However, the tail track is more toward the negative x-axis direction
than the head track. The difference between the head and tail tracks indicates that the
worm pushes the ground surface by the distance between the head track and tail
track to obtain thrust (Fig. 2B). Indeed, we found that the body part placed diagonally
with respect to the direction of the worm's locomotion is pushing along the ground
surface (Fig. S3A). The thrust force of the worm cancels out most of the drag force,

which enables the worm to move at nearly constant velocity. The average velocity of
the worm is 0.208 (mm/sec), which is consistent with the known values (Cohen et al.,
2012; Jung et al., 2016; Omura et al., 2012; Shen et al., 2012).

Figure 2. Simulated locomotion from a sine kymogram. (A) Crawling kymogram.
Kymogram indicates the angle of i-joint which is located between i-rod and (i+1)-rod.
Red and blue color mean i-joint bend in the dorsal and ventral directions, respectively.
(B) Crawling trajectory. The yellow circle indicates the position of the worm's head.
The Orange and sky-blue lines show the worm's head and tail trajectories,
respectively. (C) Swimming kymogram. (D) Swimming trajectory

In the previous work, the worm showed swimming behavior in a water droplet
on an agar plate (Vidal-Gadea et al., 2011). As the friction coefficient of water is
smaller than that of agar, even though the area that the worm swept was wider during
swimming than crawling, the worm did not move forward much in comparison to the
area it swept (Fig. 2D). The worm gained significant momentum in the forward
direction of locomotion when the body bent in the c-shape (Fig. S3B). In contrast to
crawling, during swimming, the worm did not receive constant thrust force over time.
Thus, the speed of the worm exhibited significant oscillations over time (Fig. S3B),
and the average velocity was 0.223 (mm/sec).

ElegansBot exhibits more complex behavior including the turn motion. Unlike
previous *C. elegans* body kinematic simulation studies, our simulation can replicate

the worm's behavior using a kymogram (Fig. 3A, 3B) derived from experimental
videos. We utilized open-source software, Tierpsy Tracker (Javer et al., 2018) and
WormPose (Hebert et al., 2021), to obtain the kymogram input ($\theta_{\text{ctrl},i}^{(t)}$) for the
ElegansBot. Through simulation, we aimed to reproduce the omega-turn and delta-
turn behaviors observed in the experimental videos (Broekmans et al., 2016). When
we used the vertical and horizontal friction coefficients $b_\perp$ and $b_\parallel$ on agar, as
proposed in the previous work (Boyle et al., 2012), the trajectory was not accurately
replicated. Given that the friction coefficients could vary depending on the
concentration of the agar gel, we used $b_\perp/100$ and $b_\parallel/100$ for the vertical and
horizontal friction coefficients, respectively, which resulted in a better trajectory
replication (Details in "Proper Selection of Friction Coefficients" in Supplementary
Information, Fig. S4~7).

The trajectory (Fig. 3C, 3D, Video S1, S2) obtained from ElegansBot accurately
reproduces the experimental video (Broekmans et al., 2016). The changes in the
direction of movement caused by turns are well replicated. Additionally, during the
omega-turn or delta-turn, the body briefly performs a deep bend, and we newly
discovered the mechanism that gains significant propulsion from the deep bend
region to change direction using ElegansBot (Fig. 3C, 3D). Moreover, the ElegansBot
accurately reproduces not only the turns but also complex behaviors like the
sequence of forward-backward-turn-forward, also known as escaping behavior.

Additionally, we calculated the mechanical power of the worm as a
quantitative indicator to explain its locomotion during sequenced locomotive
behavior, based on behavior classification (forward, backward locomotion, or turn,
as defined in Methods). During escaping behavior, the worm produced an average
power of 2,094 fW in the initial forward locomotion, followed by an average of 16,437
fW (7.85 times that of the initial forward locomotion) in backward locomotion, and
an average of 11,118 fW (5.31 times that of the initial forward locomotion) during
turning (Fig. 4A). After turning and resuming forward locomotion, it produced an
average power of 5,480 fW (2.62 times that of the initial forward locomotion). This
indicates that the worm produced more power than that of initial forward locomotion
to escape sudden threats. Let's denote the average of a quantity for all given i as $\langle * \rangle_i$.
At the moment the worm formed a deep bend (t=11.2 sec), the average magnitude of
frictional force of the body part forming the deep bend ($= \langle |F_{b,i}^{(t)}| \rangle_i$ where i=4~15)
was 3,536 pN, compared to the average magnitude of the remaining parts ($\langle |F_{b,i}^{(t)}| \rangle_i =$
1,737 pN where i=1~3 or i=16~25), which was 2.04 times greater (Fig. 3C, Fig. 4A).

We analyzed delta-turn in the same manner. The worm produced an average power
of 3,514 fW in the initial forward locomotion, followed by an average of 11,176 fW
(3.18 times that of the initial forward locomotion) in subsequent backward
locomotion. In the relatively short duration of forward locomotion following the
backward locomotion, the worm produced average power of 17,544 fW (4.99 times
that of the initial forward locomotion), and an average of 13,046 fW (3.71 times that
of the initial forward locomotion) during turns (Fig. 4B). After the turn, when
resuming forward locomotion, the worm produced an average power of 6,429 fW
(1.83 times that of the initial forward locomotion). At the moment the worm formed
a deep bend (t=6.1 sec), the average magnitude of frictional force of the body part
($\langle |F_{b,i}^{(t)}| \rangle_i$ where i=16~25) was 10,497 pN, compared to the average magnitude of the
remaining parts ($\langle |F_{b,i}^{(t)}| \rangle_i$= 2,677 pN where i=1~15), which was 3.92 times greater
(Fig. 3D, Fig. 4B). In both escaping behavior and delta-turn, the worm consistently
produced more power in the subsequent backward locomotion and turn than in the
initial forward locomotion.
338

Figure 3. Simulated locomotion from a kymogram of a real worm locomotion video. The length and direction of a black arrow indicate the magnitude and direction of the frictional force $(-F_{b,i}^{(t)})$ that the corresponding body part, which is the starting point of the arrow, exerts on the surface. (A) escaping behavior kymogram. Triangles over the heatmap indicate the corresponding time of snapshots shown in Figure (C). (B) Delta-turn kymogram. Triangles over the heatmap indicate the corresponding time of snapshots shown in Figure (D). (C) Escaping behavior trajectory. (D) Delta-

348 turn trajectory. The arrow length scale is different from Figure (C) to clearly show the
349 arrows' directions and head and tail tracks. (Those videos are in Supplementary
350 Information)

Figure 4. Frictional force on each rod (A) Escaping behavior. The top panel represents the frictional force $\mathbf{F}_{b,i}^{(t)}$ experienced by i-rod. As indicated on the color wheel to the right, the hue of this heatmap represents the direction of the force, and the saturation represents the magnitude of the force. The second panel from the top shows the magnitude of the frictional force $\left|\mathbf{F}_{b,i}^{(t)}\right|$. The third panel from the top represents the average $\left\langle\left|\mathbf{F}_{b,i}^{(t)}\right|\right\rangle_i$ (black solid line) of each column in the middle panel and the power (red dotted line), which is the amount of energy the worm consumes per unit time. The bottom panel represents the classification of the worm's behavior (blue: forward locomotion, red: backward locomotion, green: turn) (The definitions of behavioral categories are in Methods). The triangles over each panel indicate the corresponding time of the snapshots depicted in Fig. 3C. (B) Delta-turn. The triangles over each panel indicate the corresponding time of the snapshots depicted in Fig. 3D.

ElegansBot presents body shape ensembles of *C. elegans* from a shape in water en route to agar. While there have been studies on how locomotion patterns change in agar and water by merging neural and kinematic simulations (Boyle et al., 2012), there have been none that solely used kinetic simulation to analyze how speed manifests depending on the frequency and period of locomotion. We demonstrate this aspect. We studied the locomotion speed of the worm under different friction coefficients, which represent the influence of water, agar, and intermediate frictional environment, using ElegansBot. The vertical and horizontal friction coefficients in water are $b_{\text{water},\perp} = 5.2 \times 10^3$ (μg/sec) and $b_{\text{water},\parallel} = b_{\text{water},\perp}/1.5$, respectively, while in agar, these values are $b_{\text{agar},\perp} = 1.28 \times 10^8$ (μg/sec) and $b_{\text{agar},\parallel} = b_{\text{agar},\perp}/40$ (Boyle et al., 2012). For environmental index $\sigma \in [0,1]$, we have defined the vertical and horizontal friction coefficients in the environment between water ($\sigma = 0$) and agar ($\sigma = 1$) as $b_{\sigma,\perp} = b_{\text{water},\perp}^{1-\sigma} b_{\text{agar},\perp}^{\sigma}$ and $b_{\sigma,\parallel} = b_{\text{water},\parallel}^{1-\sigma} b_{\text{agar},\parallel}^{\sigma}$, respectively.

Under an environmental index σ, for various pairs of frequency-period (ν, T) when the control angle is $\theta_{\text{ctrl},i}^{(t)} = A\cos\left(2\pi\left(\nu\frac{i-1}{23} - \frac{t}{T}\right)\right)$ (with A = 0.6 (rad)), we have found the (ν, T) that maximizes the worm's average velocity(optimal (ν, T)) (Fig. 5A). The optimal (ν, T) exhibits nearly linear distribution (Fig. 5B). We noticed a transition from swimming body shape to crawling body shape as σ varies (Fig. 5, S8). The optimal (ν, T) for σ=0(water) is (0.65, 0.4 sec), matching the actual (ν, T) value of swimming behavior (Vidal-Gadea et al., 2011). The optimal (ν, T) for σ=1(agar) is (1.9, 0.8 sec), and the optimal ν (1.9) matches the actual ν value (1.832) for crawling behavior (Vidal-Gadea et al., 2011), with the optimal T (0.8 sec) being half the actual T value (1.6 sec).

We wanted to understand the impact of the environmental index σ not only on forward locomotion but also on sequenced locomotive behavior. First, we analyzed the effect of the environmental index σ on escaping behavior as follows. Let's denote the set of a quantity for all pair of index i and time t as $\{*\}_{i,t}$. When the escaping behavior kymogram input $\{\theta_{\text{ctrl},i}^{(t)}\}_{i,t}$ was same as Fig. 3A, we explored the effect of vertical and horizontal friction coefficients on the worm's motion. Where σ ranged from 1.0 to 0, the trajectory varied with σ (Fig. S9A), and $E_\theta = \left\langle\left|\theta_i^{(t)} - \theta_{\text{ctrl},i}^{(t)}\right|\right\rangle_{i,t}$ decreased as σ decreased (Fig. S9B). From σ = 1.0 to 0.1, the total absolute angular change ($S = \sum_{t=0}^{T-\Delta t}\left|\left\langle s_i^{(t+\Delta t)}\right\rangle_i - \left\langle s_i^{(t)}\right\rangle_i\right|$ where T is total time of the experimental video.) increased as σ decreased. However, from σ = 0.7 to 0, S

remained constant within the error of 0.33 rad, and the total traveled distance($\sum_t$ $\left|\mathbf{v}_\mathrm{c}^{(t)}\Delta t\right|$) decreased as σ decreased. The maximum total traveled distance was at σ = 0.8. Using the same analysis method with the kymogram input $\{\theta_{\mathrm{ctrl},i}^{(t)}\}_{i,t}$ same as Fig. 3B, we analyzed the impact of the environmental index σ on delta-turn. Where σ ranged from 1.0 to 0, the trajectory varied with σ (Fig. S10A), and E_θ also decreased as σ decreased (Fig. S10B). From σ = 1 to 0.6, S increased as σ decreased. From σ = 0.6 to 0, S decreased as σ decreased. The maximum total traveled distance was at σ = 0.9. From σ = 0.9 to 0, the total traveled distance decreased as σ decreased.

Figure 5. Body shape transition from the shape in water to the shape in agar. (A) Average velocity of the worm as a function of wavenumber (ν) and period (T) for a given friction coefficient. The star symbol indicates the pair of (ν, T) that maximizes the worm's average velocity. (B) For each environmental index σ, the pair of (ν, T) that maximizes the worm's average velocity. The worm figures inside the small rectangles pointed to by the arrows represent the body shape corresponding to the respective (ν, T) pair.

Discussion

ElegansBot is an advanced kinetic simulator that reproduces *C. elegans*' various locomotion. The known crawling speed range of *C. elegans* (Cohen et al., 2012; Jung et al., 2016; Omura et al., 2012; Shen et al., 2012) matches the speed in our simulation.

The force dispersion pattern of the forward movement of a snake (Hu et al., 2009) is similar to the force dispersion pattern of crawling in our model, where the body part placed diagonally to the direction of movement generates thrust. The head and tail tracks of our simulation resemble the trace left on the agar plate by *C. elegans* during locomotion (Yeon et al., 2018), providing evidence of the mechanism where *C. elegans* moves forward by pushing along the ground surface. Given that friction and elasticity coefficients can vary between experiments, the appropriate selection of these values allows the trajectories of omega-turns and delta-turns in our simulations to match the experimental videos (Broekmans et al., 2016). Previous work (Boyle, 2009; Boyle et al., 2012) eliminated inertia from the equations of motion, but our simulation includes it, allowing calculation even in cases where inertia is significant due to low friction coefficients. Using the crawling and swimming wavenumbers and periods from the experiments (Vidal-Gadea et al., 2011), we computed sine functions to create trajectories for crawling and swimming. We also analyzed how friction forces act on the worm during crawling and swimming, studying how the worm gains propulsion. We demonstrated that we could reproduce various locomotion observed in experimental videos, such as forward-backward-(omega turn)-forward constituting escaping behavior and delta-turn navigation, by providing the kymogram obtained from representative physical values from the experimental videos, as well as the kymogram obtained from a program (Hebert et al., 2021; Javer et al., 2018) extracting the body angles from actual experimental videos into ElegansBot. Our established Newtonian equations of motion are accurate and robust, suggesting that not only does our simulation replicate the experimental videos, but it also provides credible estimates for detailed forces.

ElegansBot will serve as a strong bridge for enhancing the knowledge in "from-synapse-to-behavior" research. Our method could be used for kinetic analysis of behaviors not covered in this paper. It could also be used when analyzing behavior changes caused by mutation or ablation experiments. Given that our simulation allows for kinetic analysis, it could be used to calculate the energy expended by the worm during locomotion, serving as an activity index. Our simulation only requires the body angles as input data, so even if the video angle shifts and trajectory information is lost, the trajectory can be recovered from the kymogram. Our simulation could also be used when studying neural circuit models of *C. elegans*. It could be used to check how signals from neural network models manifest as behaviors which is a needed function from previous work (Sakamoto et al., 2021),

and it could be used when studying compound models of neural circuits and bodies.
For example, when creating models that receive proprioception input based on body
shape (Boyle et al., 2012; Ekeberg, 1993; Izquierdo & Beer, 2018; Niebur & Erdös,
1991), our method could be used. Finally, our method could be used in general for the
broad utility to analyze the motion of rod-shaped animals like snakes or eels and to
simulate the motion of rod-shaped robots.

Methods

1. Frequency and wavelength of *C. elegans* locomotion

Sine function fitting was applied to the crawling and swimming kymograms (Vidal-Gadea et al., 2011) to determine the frequency and wavelength of *C. elegans* locomotion on agar and water.

2. *C. elegans* locomotion videos

Videos of *C. elegans*' escaping behavior and foraging behavior were obtained from previous work (Broekmans et al., 2016). A single representative video out of a total of one hundred escaping behavior videos was used as data in this paper. Additionally, only the delta-turning portion of the foraging behavior videos was cut out and used as data in this paper.

3. Obtaining kymograms from video

The following method was used to extract the kymogram from the video of *C. elegans*: The body angles and midline were extracted using Tierpsy Tracker (Javer et al., 2018) from the original video where the worm is locomoting. Tierpsy Tracker failed to extract the midline of the worm when the body parts meet or the worm is coiled. The midline information of the frames successfully predicted by Tierpsy Tracker and the original video information were used as ground truth training data for a program called WormPose (Hebert et al., 2021). WormPose trained an artificial neural network to extract the body angles and midline of a coiled worm using a generative method based on the input data. The body angles were extracted from the original video using the trained WormPose program. For the frames where Tierpsy Tracker failed to extract the body angles, it was replaced with the body angles extracted by WormPose. Nonetheless, there were frames where the body angle extraction failed. If the period of failed body angle prediction was continuously less than three frames

| 488 | (about 0.01 seconds), the body angles for that period was predicted using |
| 489 | linear interpolation. |

4. Program code and programming libraries

Equations for the chain model, friction model, muscle model, and numerical integration that constitute the ElegansBot were designed from the body angle information and kymogram that change every moment of time. Python (van Rossum, 1995) version 3.8 was used to implement the equations constituting ElegansBot as a program. NumPy (Harris et al., 2020) version 1.19 was used for numerical calculations, and Numba (Lam et al., 2015) version 0.54 was used for CPU calculation acceleration. SciPy (Virtanen et al., 2020) version 1.5 was used for curve fitting and Savitzky-Golay filter (Savitzky & Golay, 1964) to classify the worm's behavioral categories. The Matplotlib (Hunter, 2007) library was used to represent *C. elegans*' body pose and trajectory in figures and videos. The program code used in the research can be obtained from the open database GitHub (https://github.com/taegonchung/elegansbot) or Python software repository PyPI (https://pypi.org/project/ElegansBot/), and the web live demo can be found at GitHub Page (https://taegonchung.github.io/elegansbot/). This code calculated the ElegansBot simulation of 10 seconds of simulation time in about 10 seconds of run-time on an Intel E3-1230v5 CPU.

5. Physical constants of the ground surface

The friction coefficient values for the ground surface where *C. elegans* crawled and swam and the elastic and damping coefficients of *C. elegans* muscles were obtained from previous work (Boyle et al., 2012). The muscle elasticity and damping coefficients were converted into coefficients for the damped torsional spring to be used in our model (Details in "Worm's mass, actuator elasticity coefficient, and damping coefficient" of Supplementary Information).

6. Defining Behavioral Categories

We determined the classification of the worm's behavior over time as follows. Let $\xi_i \equiv (i - 1)/(n - 2)$ (i.e., $0 \leq \xi_i \leq 1$). We defined a sine function fitting for the body angle $\theta_i^{(t)}$ as $\hat{\theta}_i^{(t)} = A^{(t)}\sin\left(\left(2\pi/\lambda^{(t)}\right)\left(\xi_i - \xi_0^{(t)}\right)\right) + \theta_0^{(t)}$ (where $A^{(t)} \geq 0$ and $0 \leq \xi_0^{(t)} < \lambda^{(t)}$). Let us denote the set of a quantity for all i as $\{*\}_i$.

For a given time t, by curve fitting the function $\hat{\theta}_i^{(t)}$ to the set of body angles $\{\theta_i^{(t)}\}_i$, we can obtain the parameters $(A^{(t)}, \lambda^{(t)}, \xi_0^{(t)}, \theta_0^{(t)})$ (Fig. S11). For curve fitting, we used 'curve_fit' from the scipy library (Virtanen et al., 2020). 'curve_fit' requires the function to be fitted, the data, and the initial guess values of the function parameters. Therefore, we determined the initial guess values $(\hat{A}^{(t)}, \hat{\lambda}^{(t)}, \hat{\xi}_0^{(t)}, \hat{\theta}_0^{(t)})$ as follows. For $t = 0$, we calculated $(\hat{A}^{(0)}, \hat{\lambda}^{(0)}, \hat{\xi}_0^{(0)}, \hat{\theta}_0^{(0)})$ using the following equations:

$$\hat{A}^{(0)} = \frac{1}{2}\left(\max_i(\theta_i^{(0)}) - \min_i(\theta_i^{(0)})\right)$$

$$\hat{\lambda}^{(0)} = \frac{2}{n-2}\left(\operatorname*{argmax}_i(\theta_i^{(0)}) - \operatorname*{argmin}_i(\theta_i^{(0)})\right)$$

$$\hat{\xi}_0^{(0)} = \frac{1}{n-2}\left(\operatorname*{argmax}_i(\theta_i^{(0)})\right) - \frac{\hat{\lambda}^{(0)}}{4}$$

$$\hat{\theta}_0^{(0)} = \frac{1}{n-1}\left(\sum_i \theta_i^{(0)}\right)$$

Using these initial guess values, we curve fitted $\hat{\theta}_i^{(0)}$ to $\{\theta_i^{(0)}\}_i$ to obtain $(A^{(0)}, \lambda^{(0)}, \xi_0^{(0)}, \theta_0^{(0)})$. For $t \geq \Delta t$, we obtained the initial guess values for curve fitting as $\hat{A}^{(t)} = A^{(t-\Delta t)}$, $\hat{\lambda}^{(t)} = \lambda^{(t-\Delta t)}$, $\hat{\xi}_0^{(t)} = \xi_0^{(t-\Delta t)}$, $\hat{\theta}_0^{(t)} = \xi_0^{(t-\Delta t)}$. Then, using these initial guess values, we curve fitted $\hat{\theta}_i^{(t)}$ to $\{\theta_i^{(t)}\}_i$ to obtain $(A^{(t)}, \lambda^{(t)}, \xi_0^{(t)}, \theta_0^{(t)})$. Since the phase $\xi_0^{(t)}$ is not continuous for all time t, we defined a continuous value $\tilde{\xi}_0^{(t)}$ for all time t as follows (Fig. S11):

$$\tilde{\xi}_0^{(t)} = \begin{cases} \xi_0^{(t)} & \text{if } t = 0 \\ \xi_0^{(t)} - \xi_0^{(t-\Delta t)} + \tilde{\xi}_0^{(t-\Delta t)} & \text{if } t > 0 \text{ and } -\frac{\lambda^{(t)}}{2} \leq \xi_0^{(t)} - \xi_0^{(t-\Delta t)} \leq \frac{\lambda^{(t)}}{2} \\ \xi_0^{(t)} - \xi_0^{(t-\Delta t)} + \tilde{\xi}_0^{(t-\Delta t)} + \lambda^{(t)} & \text{if } t > 0 \text{ and } \xi_0^{(t)} - \xi_0^{(t-\Delta t)} < -\frac{\lambda^{(t)}}{2} \\ \xi_0^{(t)} - \xi_0^{(t-\Delta t)} + \tilde{\xi}_0^{(t-\Delta t)} - \lambda^{(t)} & \text{if } t > 0 \text{ and } \frac{\lambda^{(t)}}{2} < \xi_0^{(t)} - \xi_0^{(t-\Delta t)} \end{cases}$$

To obtain the derivatives of the noise-reduced smoothed values $\bar{\tilde{\xi}}_0^{(t)}$ and $\bar{\theta}_0^{(t)}$

for the raw data $\tilde{\xi}_0^{(t)}$ and $\theta_0^{(t)}$, respectively, we applied a Savitzky-Golay filter

(Savitzky & Golay, 1964; Virtanen et al., 2020). This filter, set with a smoothing

time window of 0.5 second, a polynomial order of 2, and a derivative order of

1, yielded $\frac{d\bar{\tilde{\xi}}_0^{(t)}}{dt}$ and $\frac{d\bar{\theta}_0^{(t)}}{dt}$. We then calculated the temporal integrals $\xi'^{(\zeta)}_0 \equiv$

$\sum_{t=0}^{\zeta} \frac{d\bar{\tilde{\xi}}_0^{(t)}}{dt} \Delta t$ and $\theta'^{(\zeta)}_0 \equiv \sum_{t=0}^{\zeta} \frac{d\bar{\theta}_0^{(t)}}{dt} \Delta t$. Let us denote the average of a quantity

for all time t as $\langle * \rangle_t$. We calculated $\bar{\tilde{\xi}}_0^{(t)} = \xi'^{(t)}_0 - \langle \xi'^{(t)}_0 \rangle_t + \langle \tilde{\xi}_0^{(t)} \rangle_t$ and $\bar{\theta}_0^{(t)} =$

$\theta'^{(t)}_0 - \langle \theta'^{(t)}_0 \rangle_t + \langle \theta_0^{(t)} \rangle_t$ (Fig. S11). Finally, we defined the worm's behavior

classification as turn when $\bar{\theta}_0^{(t)} < -0.07$, as forward locomotion when $\bar{\theta}_0^{(t)} \geq$

-0.07 and $\frac{d\bar{\tilde{\xi}}_0^{(t)}}{dt} > 0$, and as backward locomotion when $\bar{\theta}_0^{(t)} \geq -0.07$ and

$\frac{d\bar{\tilde{\xi}}_0^{(t)}}{dt} \leq 0$ (Fig. S11).

References

Boyle, J. H. (2009). *C. elegans locomotion: An integrated approach*.

Boyle, J. H., Berri, S., & Cohen, N. (2012). Gait Modulation in C. elegans: An Integrated

Neuromechanical Model. *Frontiers in Computational Neuroscience, 6*.

https://doi.org/10.3389/fncom.2012.00010

Broekmans, O. D., Rodgers, J. B., Ryu, W. S., & Stephens, G. J. (2016). Resolving coiled

shapes reveals new reorientation behaviors in C. elegans. *eLife, 5*, e17227.

https://doi.org/10.7554/eLife.17227

Butcher, J. C. (2004). *Numerical Methods for Ordinary Differential Equations*. John

Wiley & Sons.

Cohen, E., Yemini, E., Schafer, W., Feitelson, D. G., & Treinin, M. (2012). Locomotion

analysis identifies roles of mechanosensory neurons in governing locomotion

dynamics of C. elegans. *Journal of Experimental Biology*, *215*(20), 3639–3648.

https://doi.org/10.1242/jeb.075416

Cook, S. J., Jarrell, T. A., Brittin, C. A., Wang, Y., Bloniarz, A. E., Yakovlev, M. A., Nguyen,

K. C. Q., Tang, L. T.-H., Bayer, E. A., Duerr, J. S., Bülow, H. E., Hobert, O., Hall, D.

H., & Emmons, S. W. (2019). Whole-animal connectomes of both

Caenorhabditis elegans sexes. *Nature*, *571*(7763), 63–71.

https://doi.org/10.1038/s41586-019-1352-7

Ekeberg, O. (1993). A combined neuronal and mechanical model of fish swimming.

Biological Cybernetics, 12.

Hall, D. H., & Altun, Z. F. (2008). *C. elegans Atlas*.

Harris, C. R., Millman, K. J., van der Walt, S. J., Gommers, R., Virtanen, P., Cournapeau,

D., Wieser, E., Taylor, J., Berg, S., Smith, N. J., Kern, R., Picus, M., Hoyer, S., van

Kerkwijk, M. H., Brett, M., Haldane, A., del Río, J. F., Wiebe, M., Peterson, P., …

Oliphant, T. E. (2020). Array programming with NumPy. *Nature*, *585*(7825),

Article 7825. https://doi.org/10.1038/s41586-020-2649-2

Hebert, L., Ahamed, T., Costa, A. C., O'Shaughnessy, L., & Stephens, G. J. (2021).

WormPose: Image synthesis and convolutional networks for pose estimation

in C. elegans. *PLOS Computational Biology*, *17*(4), e1008914.

https://doi.org/10.1371/journal.pcbi.1008914

Hill, A. V. (1937). The heat of shortening and the dynamic constants of muscle.

Proceedings of the Royal Society of London. Series B - Biological Sciences,

126(843), 136–195. https://doi.org/10.1098/rspb.1938.0050

Hu, D. L., Nirody, J., Scott, T., & Shelley, M. J. (2009). The mechanics of slithering

locomotion. *Proceedings of the National Academy of Sciences*, *106*(25), 10081–

10085. https://doi.org/10.1073/pnas.0812533106

Hunter, J. D. (2007). Matplotlib: A 2D Graphics Environment. *Computing in Science &*

Engineering, *9*(3), 90–95. https://doi.org/10.1109/MCSE.2007.55

Izquierdo, E. J., & Beer, R. D. (2018). From head to tail: A neuromechanical model of

forward locomotion in Caenorhabditis elegans. *Philosophical Transactions of*

the Royal Society B: Biological Sciences, *373*(1758), 20170374.

https://doi.org/10.1098/rstb.2017.0374

Javer, A., Currie, M., Lee, C. W., Hokanson, J., Li, K., Martineau, C. N., Yemini, E.,

Grundy, L. J., Li, C., Ch'ng, Q., Schafer, W. R., Nollen, E. A. A., Kerr, R., & Brown,

A. E. X. (2018). An open-source platform for analyzing and sharing worm-

behavior data. *Nature Methods*, *15*(9), Article 9.

https://doi.org/10.1038/s41592-018-0112-1

Jung, J., Nakajima, M., Takeuchi, M., Najdovski, Z., Huang, Q., & Fukuda, T. (2016).

Microfluidic Device to Measure the Speed of C. elegans Using the Resistance

Change of the Flexible Electrode. *Micromachines*, *7*(3), 50.

https://doi.org/10.3390/mi7030050

Lam, S. K., Pitrou, A., & Seibert, S. (2015). Numba: A LLVM-based Python JIT

compiler. *Proceedings of the Second Workshop on the LLVM Compiler*

Infrastructure in HPC, 1–6. https://doi.org/10.1145/2833157.2833162

Niebur, E., & Erdös, P. (1991). Theory of the locomotion of nematodes: Dynamics of

undulatory progression on a surface. *Biophysical Journal*, *60*(5), 1132–1146.

https://doi.org/10.1016/S0006-3495(91)82149-X

Omura, D. T., Clark, D. A., Samuel, A. D. T., & Horvitz, H. R. (2012). Dopamine

Signaling Is Essential for Precise Rates of Locomotion by C. elegans. *PLOS*

ONE, *7*(6), e38649. https://doi.org/10.1371/journal.pone.0038649

Sakamoto, K., Soh, Z., Suzuki, M., Iino, Y., & Tsuji, T. (2021). Forward and backward

locomotion patterns in C. elegans generated by a connectome-based model

simulation. *Scientific Reports*, *11*(1), Article 1.

https://doi.org/10.1038/s41598-021-92690-2

Savitzky, Abraham., & Golay, M. J. E. (1964). Smoothing and Differentiation of Data

by Simplified Least Squares Procedures. *Analytical Chemistry*, *36*(8), 1627–

1639. https://doi.org/10.1021/ac60214a047

Shen, X. N., Sznitman, J., Krajacic, P., Lamitina, T., & Arratia, P. E. (2012). Undulatory

Locomotion of Caenorhabditis elegans on Wet Surfaces. *Biophysical Journal*,

102(12), 2772–2781. https://doi.org/10.1016/j.bpj.2012.05.012

van Rossum, G. (1995). *Python reference manual* (Issue R 9525). CWI.

Vidal-Gadea, A., Topper, S., Young, L., Crisp, A., Kressin, L., Elbel, E., Maples, T.,

Brauner, M., Erbguth, K., Axelrod, A., Gottschalk, A., Siegel, D., & Pierce-

Shimomura, J. T. (2011). Caenorhabditis elegans selects distinct crawling and

swimming gaits via dopamine and serotonin. *Proceedings of the National*

Academy of Sciences, *108*(42), 17504–17509.

https://doi.org/10.1073/pnas.1108673108

Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D.,

Burovski, E., Peterson, P., Weckesser, W., Bright, J., van der Walt, S. J., Brett, M.,

Wilson, J., Millman, K. J., Mayorov, N., Nelson, A. R. J., Jones, E., Kern, R., Larson,

E., … van Mulbregt, P. (2020). SciPy 1.0: Fundamental algorithms for scientific

computing in Python. *Nature Methods*, *17*(3), Article 3.

https://doi.org/10.1038/s41592-019-0686-2

White, J. G., Southgate, E., Thomson, J. N., & Brenner, S. (1986). The Structure of the

Nervous System of the Nematode Caenorhabditis elegans. *Philosophical*

Transactions of the Royal Society of London. Series B, Biological Sciences,

314(1165,), 1–340.

Yeon, J., Kim, J., Kim, D.-Y., Kim, H., Kim, J., Du, E. J., Kang, K., Lim, H.-H., Moon, D., &

Kim, K. (2018). A sensory-motor neuron type mediates proprioceptive

coordination of steering in C. elegans via two TRPC channels. *PLOS Biology*,

16(6), e2004929. https://doi.org/10.1371/journal.pbio.2004929

Acknowledgement

This work is supported by the p-CoE program of DGIST 23-CoE-BT-01. We also acknowledge Prof. Kyuhyung Kim for fruitful discussions on experiments of C. elegans.