Systematic perturbations of SETD2, NSD1, NSD2, NSD3 and ASH1L reveals their distinct contributions to H3K36 methylation

Gerry A. Shipman[1,2,5], Reinnier Padilla[1,2,5], Cynthia Horth[1,2], Bo Hu[1,2], Eric Bareke[1,2], Francisca N. Vitorino[3], Joanna M. Gongora[3], Benjamin A. Garcia[3], Chao Lu[4], Jacek Majewski[1,2*]

[1]Department of Human Genetics, McGill University, Montreal, Quebec, H3A 1B1, Canada

[2]McGill University Genome Centre, Montreal, Quebec, H3A 0G1, Canada

[3]Department of Biochemistry & Molecular Biophysics, Washington University School of Medicine, St. Louis, Missouri 63110, USA

[4]Department of Genetics and Development, Columbia University Irving Medical Center, New York, New York 10032, USA

[5]These authors contributed equally to this work.

*Correspondence: jacek.majewski@mcgill.ca

Abstract

Background: Methylation of histone 3 lysine 36 (H3K36me) has emerged as an essential epigenetic component for the faithful regulation of gene expression. Despite its importance in development, disease, and cancer, how the molecular agents collectively shape the H3K36me landscape is unclear.

Results: We use a mouse mesenchymal stem cell model to perturb the H3K36me deposition machinery and infer the activities of the five most prominent players: SETD2, NSD1, NSD2, NSD3, and ASH1L. We find that H3K36me2 is the most abundant of the three methylation states and is predominantly deposited at intergenic regions by NSD1, and partly by NSD2. In contrast, H3K36me1/3 are most abundant within exons and are positively correlated with gene expression. We demonstrate that while SETD2 deposits most H3K36me3, it also deposits H3K36me2 within transcribed genes. Additionally, loss of SETD2 results in an increase of exonic H3K36me1, suggesting other H3K36 methyltransferases (K36MTs) prime gene bodies with lower methylation states ahead of transcription. Through a reductive approach, we uncover the distribution patterns of NSD3- and ASH1L-catalyzed H3K36me2. While NSD1/2 establish broad intergenic H3K36me2 domains, NSD3 deposits H3K36me2 peaks on active promoters and enhancers. Meanwhile, the activity of ASH1L is restricted to the regulatory elements of developmentally relevant genes, and our analyses implicate PBX2 as a potential recruitment factor.

Conclusions: Within genes, SETD2 deposits both H3K36me2/3, while the other

K36MTs are capable of depositing H3K36me1/2 independently of SETD2 activity. For

the deposition of H3K36me1/2, we find a hierarchy of K36MT activities where

NSD1>NSD2>NSD3>ASH1L. While NSD1 and NSD2 are responsible for most

genome-wide propagation of H3K36me2, the activities of NSD3 and ASH1L are

confined to active regulatory elements.

Keywords: SETD2, NSD1, NSD2, NSD3, ASH1L, H3K36 methylation, epigenetics,

transcription

Background

Epigenetic control of gene expression relies on a fine balance between the deposition

and maintenance of a myriad of chromatin modifications. The amino terminal tail of

histone 3 is a heavily post-translationally modified region of the nucleosome, with the

various modifications having diverse roles and permutations that maintain a balance

between heterochromatin, euchromatin, and open chromatin regions. Methylation at the

lysine 36 position of histone 3 (H3K36) is known to be associated with transcriptionally

active regions of the genome. The highest methylation state, H3K36me3, is a mark of

actively transcribed gene bodies (1,2). More recently, the intermediate methylation

state, H3K36me2, has garnered increased interest, with evidence supporting its

importance in preventing compaction of intergenic regions by restricting expansion of

antagonistic silencing marks, such as H3K27me2/3, thereby maintaining a balance

between gene expression and silencing (3,4). The lowest methylation state, H3K36me1,

has received limited attention (5,6), and its significance remains unclear.

Several constituents governing the deposition of H3K36 methylation (H3K36me)

are known. H3K36me3, which exists almost entirely within transcribed gene bodies, is

deposited by transcriptional coupling of the methyltransferase SETD2 to the elongating

RNA polymerase II complex (RNAPII) (7,8). In simple eukaryotes, such as yeast, the

presence of H3K36me3 itself has been shown to be important for repressing cryptic

transcription, facilitating histone turnover, and regulating pre-mRNA splicing (9,10,11).

In contrast, H3K36 mono- and di-methylation are the products of at least four other

histone methyltransferases, NSD1/2/3, and possibly ASH1L. While all four of these

enzymes have demonstrated H3K36 methyltransferase activity, they appear to have

non-redundant functional properties, and their implications in genetic disease and

cancer are markedly different. NSD1 loss of function mutations have been identified in

most patients with Sotos syndrome, a developmental overgrowth disorder, and also in a

subgroup of HPV-negative head and neck squamous cell carcinomas (HNSCC). Here,

loss of NSD1 results in depletion of intergenic H3K36me2, and a concomitant decrease

of H3K27ac and DNA methylation (DNAme), thereby reducing the activity of associated

cis-regulatory elements (CREs) and the expression of their putative target genes

(12,13,14,15). In contrast, NSD2 truncating and missense variants have been causally

associated with Wolf-Hirschhorn syndrome, a genetic disorder characterized by

intellectual and developmental delay (16,17,18). Translocation-mediated

overexpression and hyperactive variants of NSD2 and NSD3 have been identified in

several types of cancer, notably in multiple myeloma, lung squamous cell carcinoma,

and pancreatic ductal adenocarcinoma, with elevated H3K36me2 being implicated in

tumour progression and survival outcomes (19,20,21). As an H3K36 methyltransferase,

ASH1L remains relatively understudied. However, nonsense and missense variants

have been linked with autism spectrum disorder and brain development, and its

overexpression has been identified in both anaplastic thyroid cancer and acute

leukemia (22,23,24).

H3K36me is an essential epigenetic modification for the establishment and

maintenance of gene regulatory programs, and impacts the deposition of many other

histone marks, as well as DNAme (25,26,27,28). Clearly, there are considerable

pathological consequences when it is aberrantly regulated, underscoring the necessity

to better understand its functional significance. Here, we use a mouse mesenchymal

stem cell model to dissect the roles of the most well-established H3K36

methyltransferases (K36MTs) in the deposition and maintenance of this modification.

Mesenchymal stem cells, which are multipotent, can be induced into oncogenic

transformation (29,30,31). In this study, we use CRISPR-Cas9 gene editing to assemble

a panel of single and combinatorial knockout cell lines, allowing us to deconstruct the

individual contributions of each methyltransferase to the overall H3K36me landscape.

Our work provides new insights into the distinct and overlapping genomic preferences of

lysine 36 methyltransferases and consolidates existing knowledge in the context of a

structured analysis.

Results

H3K36 methylation states have distinct distribution patterns

In order to characterize the effects of individual K36MTs in shaping the H3K36me

landscape, we first sought to describe the deposition patterns of H3K36me in wild-type

C3H10T1/2 mouse mesenchymal stem cells (mMSCs). Therefore, we used liquid

chromatography - tandem mass spectrometry (MS) to quantify the genome-wide

abundance and chromatin immunoprecipitation with sequencing (ChIP-seq) to profile

the genomic distribution of all three H3K36me states. Consistent with previous

observations in this cell type, we find that H3K36me2 is broadly distributed both within

genes and intergenic regions (IGRs), while H3K36me3 is restricted specifically to

actively transcribed genes (Fig. 1a) (4,11). Interestingly, we find that H3K36me1 is also

found predominantly within genes, with lower levels also found in IGRs (Fig. 1a). As

quantified by MS, H3K36me2 is the most abundant of the three methylation states,

marking approximately 30% of all H3 peptides (Fig. 1b). In comparison, H3K36me1 and

H3K36me3 occupy approximately 14% and 7% of H3 peptides, respectively (Fig. 1b).

To assess the global distribution and abundance of H3K36me within IGRs, we centered

ChIP-seq signals on these regions to generate aggregate profiles. Expectedly, we find

that H3K36me3 is essentially absent and confined to flanking genic regions (Fig. 1c). In

contrast, the levels of H3K36me2 are greater within IGRs, reflecting its functional

importance within these regions (Fig. 1c) (4), while low levels of H3K36me1 are also

found within IGRs.

Upon examination of genome browser coverage tracks within transcribed genes,

we find that the levels of H3K36me1 and H3K36me2 are highest in 5' regions, plateau

within the first intron, and then decrease towards the 3' end, where they are replaced by

H3K36me3 (Fig. 1d). To closely examine the gene body distribution patterns of

H3K36me, we divided actively transcribed genes into expression quantiles and

assessed the differences of individual H3K36me marks within introns and exons across

varying levels of gene expression (Fig. 1e). We find that bulk enrichment of H3K36me1

and H3K36me3 signals are greater in exons compared to introns, whereas the inverse

relationship is true for H3K36me2 (Fig. S1a). Although the 5' to 3' profiles of H3K36me1

and H3K36me2 are similar, they correlate differently with gene expression levels:

H3K36me1 and H3K36me3 are positively correlated with gene expression whereas

H3K36me2 is not, likely reflecting the conversion efficiency of H3K36me2 to H3K36me3

within the exons of highly transcribed genes (Fig. S1b).

In summary, the dynamics of the three methylation states within gene bodies are

influenced by the levels of transcription: H3K36me1/2 decrease from 5' to 3' in gene

bodies where they are replaced by H3K36me3; H3K36me1/3 are positively correlated

with gene expression, whereas H3K36me2 is not; and H3K36me2 is the most abundant

methylation state and is broadly distributed in IGRs, whereas H3K36me1/3 are

predominantly found within genes.

Individual and combined H3K36 methyltransferase contributions to bulk

H3K36me

Each of the most well-established K36MTs - SETD2, NSD1, NSD2, NSD3 and ASH1L -

have been shown biochemically to exhibit methyltransferase activity specific for H3K36

via their catalytic Su(var)3-9, Enhancer-of-zeste and Trithorax (SET) domains

(9,32,33,34). However, their distinct contributions to the H3K36me landscape is

currently unknown. To better understand and decompose these contributions, we used

CRISPR-Cas9 gene editing to establish a panel of single and combinatorial knockout

mMSC lines, targeting the SET domain of each K36MT (Fig. S2). We used MS and

Western blots to quantify the overall changes to bulk levels of H3K36me across different

conditions. As described, H3K36me3 is the least abundant methylation state in parental

cells, marking approximately 7% of all H3 peptides (Fig. 1a), and consistent with

previous reports in mMSCs, it is almost entirely deposited by SETD2 (31). As shown by

MS, cells lacking functional SETD2 (SETD2-KO) have at least a 6 fold-reduction of

H3K36me3 in comparison to parental cells (Fig. 2a, Fig. S3a). Interestingly, in SETD2-

KO cells, the levels of H3K36me1/2 increase relative to parental cells, which is

consistent with fewer H3K36 residues being upgraded to higher orders of methylation

(Fig. 2a).

To investigate the individual contributions of each K36MT to global levels of

H3K36me1/2, we generated individual NSD1, NSD2, NSD3, and ASH1L knockout cells,

with three biological replicates for each cell line (Fig. S2a). Strikingly, ablation of NSD3

and ASH1L appears to have no discernible effect on the global abundance of

H3K36me1/2 (Fig. 2a). While both of these enzymes have previously been reported to

deposit these modifications (33,34), these results suggest that their effects may be

restricted to specific loci or masked by the effects of other K36MTs, such as NSD1 or

NSD2. In contrast, loss of NSD1 results in the greatest depletion of both H3K36me1/2,

especially H3K36me2 which is reduced more than 2-fold, while in NSD2-KO cells,

H3K36me2 levels are reduced by approximately 12% (Fig. 2a).

Further decreases of H3K36me2 are observed in multi-knockout conditions. The

concurrent loss of NSD1 and NSD2 (DKO) in mMSCs leads to a 3-fold reduction in

global H3K36me2 levels (Fig. 2a/b, Fig. S3a), which is consistent with previous studies

(31). In triple knockout cells lacking NSD1/2-SETD2 (hereafter referred to as TKO), bulk

levels of H3K36me2 are further reduced, indicating that SETD2 also contributes to the

deposition of this mark (Fig. 2a/b, Fig. S3a). Similar to SETD2-KO cells, TKO cells are

nearly devoid of H3K36me3. Ablation of NSD3 in NSD1/2/3-SETD2 quadruple knockout

cells and subsequently ASH1L in NSD1/2/3-SETD2-ASH1L quintuple knockout cells

(hereafter referred to as QKO and QuiKO, respectively) results in a total loss of

H3K36me2 (Fig. 2a/b), and these observations are further supported by ChIP-Rx (Fig.

S3a).

We have previously explored and characterized mMSCs overexpressing the

H3K36M oncohistone (H3K36M-OE), a mutation frequently identified in

chondroblastoma patients, and found that one of the primary effects of this mutation is

global H3K36 hypomethylation (31). Comparing the effects of this mutation to our

combinatorial knockouts, we find that the lowest levels of H3K36me are observed in the

H3K36M-OE, QKO, and QuiKO cell lines, confirming that this mutation exerts a global

effect on the activity of multiple K36MTs (Fig. 2a, Fig. S3a). Indeed, principal

component analysis (PCA) of the three methylation states reveals that cells with

multiple K36MT deletions cluster more closely with cells overexpressing the H3K36M

mutation (Fig. 2c, Fig. S3c). Interestingly, for H3K36me2, the NSD1-KO samples also

cluster more closely with the multi-knockout and H3K36M-OE samples, supporting that

NSD1 deposits the bulk of H3K36me2. In comparison, the other individual knockout

conditions cluster more closely with parental cells (Fig. 2c, Fig. S3c).

Overall, as it pertains to the global abundance of the three H3K36me states in

mMSCs, SETD2 deposits nearly all H3K36me3, while NSD1 is responsible for the

majority of bulk H3K36me1/2 levels. In comparison, NSD2 also deposits modest

amounts of H3K36me1/2, while NSD3 and ASH1L have the smallest contributions.

H3K36me1 is primarily associated with gene bodies

In both disease and development, nearly all previous investigations of H3K36me and its

respective writers have focused on the higher orders of methylation, H3K36me2/3, with

H3K36me1 remaining comparatively understudied. As revealed by MS and ChIP-seq,

the global abundance of H3K36me1 is significantly reduced in both NSD1/2-DKO and

H3K36M-OE cells (Fig. 2a, Fig. 3a). In QKO and QuiKO cells, H3K36me1 is virtually

absent. However, close analysis of the ChIP-seq profiles demonstrates that the

distribution pattern of the extremely low levels of remaining H3K36me1 closely

resembles that of parental cells (Fig. 2b, Fig. 3a). Given that the five most well-

established K36MTs are ablated in the QuiKO condition, this intriguingly suggests that

there may be another K36MT capable of depositing this modification.

Interestingly, in SETD2-KO cells, the global abundance of H3K36me1 increases

(Fig. 2a/b, Fig. 3a). We hypothesized that the increase of H3K36me1 after loss of

SETD2 is, at least in part, due to mono-methylated peptides no longer being upgraded

to higher orders of methylation. Since SETD2 is known to act primarily within gene

bodies, we assessed whether the increase of H3K36me1 occurs primarily within genes

or IGRs. Indeed, in comparing the genic to intergenic ratios of H3K36me1 signal

between parental and SETD2-KO cells, we find that the increase of H3K36me1 occurs

predominantly within genes (Fig. 3b). Subsequent in-depth analysis of the changes

within gene bodies reveals that the increase of H3K36me1 occurs primarily within exons

in comparison to introns (Fig. 3c). Overall, these results support that when the activity of

SETD2 is lost, the cascade of methylation from H3K36me1 to H3K36me3 is disrupted

primarily in genes, and more specifically, within exons, leading to an increase of

H3K36me1 in these regions. Furthermore, these results suggest that other K36MTs

may prime gene bodies with H3K36me1 in advance of SETD2 activity, which is tightly

associated with transcription.

Intergenic H3K36me2 is predominantly deposited by NSD1 and in part by NSD2 in

mMSCs

We and others have previously found that one of the predominant effects of the

H3K36M mutation is a marked reduction of intergenic H3K36me2, an effect that can be

largely recapitulated from the combined loss of NSD1 and NSD2 (Fig. 4a) (31). In this

context, however, the individual contributions of NSD1 and NSD2 to the intergenic

H3K36me2 landscape in mMSCs remains unclear. Therefore, we evaluated changes to

intergenic H3K36me2 in both NSD1-KO and NSD2-KO cells. Consistent with reports in

other cell types (4,15), loss of NSD1 results in the greatest depletion of intergenic

H3K36me2, reducing it nearly 2-fold (Fig. 4a, Fig. 4b). In contrast, the loss of NSD2 also

reduces intergenic H3K36me2, although to a much lower extent. Comparing these

results to NSD1/2-DKO cells, we report that while both NSD1-KO and NSD2-KO cells

have an effect on intergenic H3K36me2, neither condition alone can recapitulate the

depletion observed in DKO cells, suggesting that NSD1 and NSD2 may propagate this

mark in IGRs through an additive effect, with NSD1 playing the predominant role (Fig.

4a, Fig. 4b). In comparing the effects of the other K36MTs, no discernible changes to

intergenic H3K36me2 are observed in SETD2-KO, NSD3-KO, or ASH1L-KO cells

(Fig.S4a).

Intergenic H3K36me2 domains span megabases and are essential components

of the gene regulatory network, often demarcating enhancers and other CREs (15).

Therefore, it is worth noting that while most of the individual K36MTs do not exert a

global effect on the broad intergenic distribution of H3K36me2, the distribution does

become confined to smaller regions upon loss of NSD1 (Fig.4a, Fig.4c), supporting that

NSD1 is the predominant K36MT in these regions and that the other K36MTs may act

with more specificity. Indeed, even in the absence of NSD1, the combined catalytic

activities of the other K36MTs are insufficient to rescue the loss of these broad

H3K36me2 domains (Fig.4a, Fig.4b). The H3K36me2 signal is further attenuated in

DKO cells, and becomes progressively more punctate in subsequent multi-knockout

conditions (i.e. in TKO, QKO, and QuiKO cells). To visualize this, we generated

"peakiness" scores for all KO conditions, which represent the average H3K36me2 ChIP

signal in the top 1% of 1-kb bins and where a higher score indicates a more punctate

distribution (Fig.4c, Fig.S4b).

Taken together, NSD1 is consistently the primary K36MT responsible for the

establishment and maintenance of broad intergenic H3K36me2 domains, while in

mMSCs where it is expressed less, NSD2 also significantly contributes to the deposition

of this mark.

Genic H3K36me distribution patterns are highly dependent on SETD2 occupancy

In the presence of the H3K36M mutation, which affects all K36MTs including SETD2,

the profile of H3K36me2 within genes exhibits a shift towards 3' regions and becomes

elevated within exons, as compared to introns (Fig. 5a-d). In NSD1/2-DKO cells, we

found a similar trend, where although the preference for the 5' region of genes remains

unchanged, the absence of NSD1/2 results in a remarkable change in the dependence

on expression levels: H3K36me2 is now positively correlated with gene activity (Fig. 5b-

c). While the distribution of H3K36me2 in the highly transcribed genes of DKO cells

remains higher within introns than exons, in lowly transcribed genes there is now

greater signal in exons compared to introns, similar to the distribution pattern of

H3K36me3 (Fig. 1e), which may result from reduced occupancy of SETD2 in genes

transcribed at lower frequencies (Fig. 5b). The observed distinction between the highly

and lowly transcribed genes in DKO cells can likely be attributed to the increased

prominence of SETD2 as the primary K36MT for H3K36me2 in genic regions, leading to

the conversion of H3K36me2 to H3K36me3, specifically in highly transcribed genes. In

genes transcribed with less frequency, however, SETD2 may primarily deposit

H3K36me2 without further conversion.

Furthermore, in SETD2-KO cells, H3K36me2 is no longer converted to

H3K36me3, and a slight positive shift in the correlation between H3K36me2 and gene

expression is observed (Fig. 5b, Fig. 5c). In contrast, the individual knockouts of

ASH1L, NSD1, NSD2 and NSD3 do not change the dependency of H3K36me2 on gene

expression (Fig. S4c). Nonetheless, the strong positive correlations between

H3K36me2 and gene expression quantiles observed in H3K36M-OE, NSD1/2-DKO,

and SETD2-KO cells are consistently observed when assessing genes on a genome-

wide scale without segregating them based on quantiles, and in this context the overall

signal demonstrates an exonic enrichment in comparison to introns. (Fig. 5d, Fig. S4d).

NSD3-mediated H3K36me2 is targeted to active regulatory regions

Although depletion of NSD1, NSD2 and SETD2 results in a progressive loss of

H3K36me2, with NSD1/2 predominantly affecting intergenic regions, and SETD2

affecting gene-bodies, a non-negligible amount of H3K36me2 still remains in TKO cells

(Fig. 2a, Fig. 6a). The profile of H3K36me2 becomes progressively more punctate,

starting from megabase size euchromatic domains in parental cells, and ending with

kilobase size broad "peaks" in the TKO cells (Fig. 6a). Close examination of the

distribution in TKO cells reveals that most of these residual peaks are centered on TSS

and enhancer regions (Fig. 6b, Fig. 6c). Adding more epigenetic information, we note

that the majority of these broad peaks straddle regions of open chromatin, as

demonstrated by ATAC-seq and the presence of H3K27ac (Fig. 6b, Fig. 6c).

In a wildtype context, it is currently unknown where NSD3 exerts its catalytic activity and

to what extent. As previously described, in NSD3-KO cells there is almost no discernible

change to the distribution or bulk levels of H3K36me2 (Fig. 2a & Fig. S4a). Therefore,

we hypothesized that the effects of NSD3 may be concealed by the activity of other

K36MTs. In QKO cells, lacking NSD1/2/3-SETD2, nearly all remaining H3K36me2 is

depleted (Fig. 6a), indicating that the residual peaks remaining in TKO cells are

deposited by NSD3. We used genomic element annotations and carried out enrichment

analysis to characterize the NSD3-deposited H3K36me2 regions (35). Confirming our

previous observations, we find that the strongest functional H3K36me2 enrichment

categories are at enhancer and promoter-flanking regions (Fig. 6d). After centering MS-

normalized H3K36me2 signal on annotated enhancers and TSS', we find that the signal

is abolished following loss of NSD3 in QKO cells (Fig. 6e). Thus, it appears that the

residual H3K36me2 found in TKO cells is deposited by NSD3 and its activity is targeted

specifically to active promoters and enhancers.

This finding implicates NSD3 in the targeted deposition of H3K36me2 around

active regulatory elements (Fig. 6b-e). Because of the broad distribution of H3K36me2

in parental cells, this role only becomes evident in the absence of more catalytically

active K36MTs. Indeed, in NSD3-KO cells, it appears that a compensatory effect may

exist, where one or more of the other K36MTs may act to deposit H3K36me2 in regions

where NSD3 would generally localize. Thus, while NSD3 is able to deposit substantial

levels of H3K36me2, its mode of activity differs from NSD1/2: NSD1/2 establish broad

H3K36me2 domains both within genes and IGRs, while NSD3 establishes more

localized broad peaks centered on promoters and enhancers.

ASH1L selectively deposits H3K36me2 at the regulatory elements of developmentally relevant genes

Previous studies of ASH1L have demonstrated that it is capable of depositing H3K36me2, and that it has been found to be enriched at the promoters of certain developmentally important genes (34,36). However, its global contributions to H3K36me2 and the precise genomic regions at which it exerts its catalytic activity remains to be elucidated. As described, we depleted ASH1L in parental mMSCs, and detected no discernible changes to the bulk levels or distribution of H3K36me2 (Fig. 2a and Fig. S4a). The QKO cells, lacking NSD1/2/3 and SETD2, exhibit extremely low levels of H3K36me2, reduced by nearly 30x compared to parental cells (Fig. 2a), however, we detected a few (n=119) clearly defined broad (10-70 kb) peaks remaining across the genome (Fig. 7a). These peaks are again centered on regions of open chromatin accessibility, as determined by ATAC-seq, with approximately 60 promoter and 20 deep intergenic sites (Fig. 7a, Fig. 7b). Based on our previous findings, we hypothesized that these remaining peaks are deposited by ASH1L, and indeed, depletion of ASH1L in the QuiKO cells results in a total loss of all remaining H3K36me2 (Fig. 2a, Fig. 7a). It is remarkable that ASH1L exhibits such high specificity for a few selected regulatory regions. To identify possible associations between ASH1L and known DNA binding factors, we carried out motif analysis under the ATAC-seq peaks within the ASH1L-associated regions. We identified enrichment of binding motifs for several developmentally important transcription factors (Fig. 7c). Using mMSC RNA-seq data (37), we find that PBX2 is the only one of these transcription factors that is

expressed, implicating PBX2 as the best candidate for recruiting ASH1L to these

regions (Fig. 7c, Fig. S5a). Overall, in mMSCs, ASH1L appears to be the least prolific of

the K36MTs, and exhibits high specificity for its recruitment to regulatory elements.

Discussion

Over the past decade, a growing body of evidence has demonstrated that

H3K36me is essential for the establishment and maintenance of gene regulatory

programs (6). Chromosomal abnormalities and loss or gain of function mutations

affecting the writers of these marks leads to many different diseases and cancer,

underscoring the need to understand how the family of H3K36 methyltransferases

regulate their deposition. Our study provides new insights into the distinct genomic

preferences of the K36MTs – SETD2, NSD1/2/3, and ASH1L - and helps to consolidate

a wealth of previous observations (Fig. 8).

Of the five K36MTs in our study, SETD2 has the most established role in

depositing H3K36 methylation. As has been previously reported, it is the dominant

H3K36me3-catalyzing enzyme in mammals, but still possesses the ability to deposit

H3K36me1/2 (11) (Fig.8). Here, we find that in the absence of SETD2, the levels of

H3K36me3 decrease significantly and its genic enrichment disappears. At the same

time, the genic distribution of the lower H3K36me2 modification loses its intronic versus

exonic enrichment and becomes more correlated with transcription – reflecting its

SETD2-independent deposition patterns. This observation highlights the coupling of

SETD2 to the phosphorylated RNAPII complex and the transcriptional dependence of

its catalytic products. 5' genic regions generally contain long, rapidly transcribed introns,

not allowing SETD2 sufficient time to catalyze the final H3K36me3 step and resulting in

predominantly H3K36me1/2 marked nucleosomes. The co-transcriptional process of

pre-mRNA splicing slows down the RNAPII complex, thereby allowing more H3K36me3

to be deposited at exons and towards the 3' ends of genes, as exon density becomes

higher. This takes place at the expense of H3K36me1/2, which display the opposite 5'

to 3' and exon/intron trends. All of the above trends become more pronounced in the

absence of NSD1/2, where SETD2 remains the dominant K36MT within gene bodies.

Finally, the absence of SETD2 results in an overall increase of H3K36me1/2, reflecting

the ability of other enzymes to deposit these two marks and their ineffectiveness in

catalyzing the last methylation step.

NSD1 is a demonstrated agent of intergenic H3K36me2 deposition. In HNSCC

and mouse embryonic stem cells (mESCs), loss of NSD1 results in a near complete

depletion of intergenic H3K36me2 (4,15). Here, in mMSCs, we also find NSD1 to

significantly affect intergenic H3K36me2 levels, demonstrating the largest decrease

among all individual K36MT KOs. The effects of NSD1 depletion on H3K36me1/3 are

much less pronounced. We find that the decrease in H3K36me2 is largest in intergenic

regions, however, comparing the single NSD1-KO to NSD1/2-DKO shows that this

intergenic depletion is not complete, and that in this cell type, NSD2 and perhaps other

enzymes, play a notable role in the deposition of H3K36me2. A recent study

demonstrates that NSD1 is recruited to active CGIs, implying that NSD1-associated

H3K36me1/2 may then spread outwards into both gene bodies and intergenic regions

(Fig. 8) (38). Within gene bodies, it is then upgraded to H3K36me3, suggesting that

NSD1 may prime nucleosomes for the deposition of H3K36me3 by providing

H3K36me1/2 as the initial substrate.

NSD2 is known to be able to create broad domains of H3K36me2, but this has

primarily been shown in the context of oncogenic overexpression (19,39). The catalytic

properties of NSD2 under normal physiological conditions have not been well

characterized. Here, we find that the individual knockout of NSD2 results in a notable

decrease of all H3K36 methylation levels, with the largest fractional decrease occurring

for H3K36me3, followed by H3K36me2/1. The depletion of H3K36me2/1 is less

pronounced than that observed in NSD1-KO cells, suggesting that similar to other

model systems, NSD1 is the dominant H3K36me1/2 methyltransferase operating in

euchromatic intergenic regions. However, in mMSCs, it appears that both NSD1 and

NSD2 cooperate to produce the intergenic distribution of H3K36me1/2 and the results of

individual KOs, as compared to the NSD1/2-DKO is consistent with their additive effect.

Interestingly, NSD2-KO appears to result in a slightly greater decrease of H3K36me3,

as compared to NSD1-KO. This is consistent with published data showing enrichment of

NSD2 within actively transcribed gene bodies (40), leading to the possibility that NSD2

is particularly effective in depositing H3K36me1/2 within genes, possibly – more

specifically than NSD1 - acting as a priming substrate for the deposition of H3K36me3

by SETD2 (Fig. 8).

As expected, the NSD1/2-SETD2-TKO cells exhibit very low levels of all three

H3K36 methylation marks. However, levels of H3K36me1/2 in particular remain

detectable, and a clear enrichment signal is detectable by ChIP-seq. In the case of

H3K36me2, there is a remarkable change in the distribution of the signal: while in

parental cells H3K36me2 is broadly distributed, and in NSD1/2-DKO cells the signal is

mostly restricted to active gene bodies, in TKO cells the signal becomes constrained to

far narrower peaks centered on open chromatin and surrounding enhancer and

promoter regions. Although the individual KO of NSD3 has little discernible effect on

H3K36 methylation, we propose that the other K36MTs can compensate for its

absence, and its activity only becomes discernible in the TKO background. As

predicted, further deletion of NSD3, creating a QKO cell line results in a near total

depletion of all H3K36me levels. Previous work shows that the short isoform of NSD3 is

recruited to active regulatory elements by BRD4 (41), and it is likely that the same is

true of the full-length isoform of NSD3. Our results are consistent with those

observations, suggesting that NSD3, like NSD1, spreads outwards from CpG rich

regulatory elements. However, NSD3 is not able to create very broad euchromatic

H3K36me2 domains, resulting in the formation of much narrower peaks (Fig. 8).

Finally, we find ASH1L to be the least prolific, and most specific K36MT. Similar

to NSD3, the individual knockout of ASH1L has no detectable effect on H3K36me levels

and distribution, implying that in this cell type, other K36MTs can compensate for its

loss. By comparing the QKO to the QuiKO cells, we show that ASH1L is only

responsible for depositing H3K36me2 at the promoters and enhancers of approximately

100 developmentally related genes, and its ability to spread the modification is even

lower than that of NSD3 (Fig. 8). It remains to be seen what mechanisms govern the

recruitment of ASH1L to those specific regions. TFBS analysis suggests that specific

transcription factors, particularly from the PBX family may be involved.

Our results illuminate some of the intricate dependencies governing the K36MTs

and the deposition of H3K36 methylation. The overall outcomes depend on several key

variables: genomic recruitment and the distribution of K36MTs; their ability to spread

and read existing modifications through their PWWP domains; their catalytic activities;

and finally, expression levels in various cellular contexts. Many of these parameters are

being continually refined. In this study, we find a hierarchy of K36MT activities

pertaining to the deposition of H3K36me1/2, with NSD1>NSD2>NSD3>ASH1L. We find

that in contrast to mESCs, the function of NSD1 is less prominent and NSD2 contributes

significantly to the normal deposition of H3K36me2. SETD2 retains its position as the

dominant H3K36me3 methyltransferase. Furthermore, trace levels of H3K36me1

remain in the QuiKO cells, and the distribution patterns largely mirror those observed in

parental cells. Other protein families outside of the scope of this study have also been

implicated in the deposition of H3K36me, such as the SMYD family (42,43), and this

residual H3K36me1 indicates that another enzyme may be responsible for its

deposition, albeit at extremely low levels.

H3K36 methylation also has profound impacts on the deposition of other histone

marks and DNAme. In the context of active transcription, H3K36me2/3 coexist with

marks such as H3K4me1/3, H3K27ac, and DNAme either within genes and/or at CREs.

Conversely, H3K36me2/3 tend not to coexist with marks that define constitutive or

facultative heterochromatin, such as H3K9me3 and H3K27me3, respectively. In the

presence of mutations affecting the deposition of H3K36 methylation, such as H3K36M,

the downstream consequences affect the establishment and maintenance of all these

marks, and in turn result in aberrant gene expression (4,14,15,31). Therefore, through

better understanding the regulation of H3K36 methylation, our work helps pave the way

for future studies to further define these and other relationships that exist between

epigenetic marks.

Conclusions

Here, we identify both unique and overlapping roles by the family of K36MTs in their

regulation of H3K36me. Our findings demonstrate that within transcribed genes, SETD2

deposits both H3K36me2/3, while the other K36MTs are capable of depositing

H3K36me1/2 independently of SETD2 activity. In most cell types, NSD1 is responsible

for most propagation of intergenic H3K36me2, however, in mMSCs, NSD2 also

contributes to its deposition in these regions. We also identified that the catalytic

products of NSD3 are primarily focused on active promoters and enhancers, and that

the activity of ASH1L is even more restricted to specific developmental genes and their

regulatory elements. Given their implications in both development and disease, our

study helps to illuminate how the family of K36MTs collectively regulate the deposition

of H3K36me. Despite the overlap in the regions where each K36MT exerts its catalytic

functions, they each have unique properties in the propagation of these marks.

References

1. Pokholok DK, Harbison CT, Levine S, Cole M, Hannett NM, Lee TI, Bell GW,

Walker K, Rolfe PA, Herbolsheimer E, et al: Genome-wide map of nucleosome

acetylation and methylation in yeast. *Cell* 2005, 122:517-527.

2. Edmunds JW, Mahadevan LC, Clayton AL: Dynamic histone H3 methylation

during gene induction: HYPB/Setd2 mediates all H3K36 trimethylation. *EMBO J*

2008, 27:406-420.

3. Streubel G, Watson A, Jammula SG, Scelfo A, Fitzpatrick DJ, Oliviero G, McCole

R, Conway E, Glancy E, Negri GL, et al: The H3K36me2 Methyltransferase Nsd1

Demarcates PRC2-Mediated H3K27me2 and H3K27me3 Domains in Embryonic

Stem Cells. *Mol Cell* 2018, 70:371-379.e375.

4. Chen H, Hu B, Horth C, Bareke E, Rosenbaum P, Kwon SY, Sirois J, Weinberg

DN, Robison FM, Garcia BA, et al: H3K36 dimethylation shapes the epigenetic

interaction landscape by directing repressive chromatin modifications in

embryonic stem cells. *Genome Res* 2022, 32:825-837.

5. Wagner EJ, Carpenter PB: Understanding the language of Lys36 methylation at

histone H3. *Nat Rev Mol Cell Biol* 2012, 13:115-126.

6. Huang C, Zhu B: Roles of H3K36-specific histone methyltransferases in

transcription: antagonizing silencing and safeguarding transcription fidelity.

Biophys Rep 2018, 4:170-177.

7. Li B, Howe L, Anderson S, Yates JR, Workman JL: The Set2 histone

methyltransferase functions through the phosphorylated carboxyl-terminal

domain of RNA polymerase II. *J Biol Chem* 2003, 278:8897-8903.

8. Schaft D, Roguev A, Kotovic KM, Shevchenko A, Sarov M, Neugebauer KM,

Stewart AF: The histone 3 lysine 36 methyltransferase, SET2, is involved in

transcriptional elongation. *Nucleic Acids Res* 2003, 31:2475-2482.

9. Strahl BD, Grant PA, Briggs SD, Sun ZW, Bone JR, Caldwell JA, Mollah S, Cook

RG, Shabanowitz J, Hunt DF, Allis CD: Set2 is a nucleosomal histone H3-

selective methyltransferase that mediates transcriptional repression. *Mol Cell Biol*

2002, 22:1298-1306.

10. Carrozza MJ, Li B, Florens L, Suganuma T, Swanson SK, Lee KK, Shia WJ,

Anderson S, Yates J, Washburn MP, Workman JL: Histone H3 methylation by

Set2 directs deacetylation of coding regions by Rpd3S to suppress spurious

intragenic transcription. *Cell* 2005, 123:581-592.

11. Molenaar TM, van Leeuwen F: SETD2: from chromatin modifier to multipronged

regulator of the genome and beyond. *Cell Mol Life Sci* 2022, 79:346.

12. Kurotaki N, Imaizumi K, Harada N, Masuno M, Kondoh T, Nagai T, Ohashi H,

Naritomi K, Tsukahara M, Makita Y, et al: Haploinsufficiency of NSD1 causes

Sotos syndrome. *Nat Genet* 2002, 30:365-366.

13. Network CGA: Comprehensive genomic characterization of head and neck

squamous cell carcinomas. *Nature* 2015, 517:576-582.

14. Papillon-Cavanagh S, Lu C, Gayden T, Mikael LG, Bechet D, Karamboulas C,

Ailles L, Karamchandani J, Marchione DM, Garcia BA, et al: Impaired H3K36

methylation defines a subset of head and neck squamous cell carcinomas. *Nat*

Genet 2017, 49:180-185.

15. Farhangdoost N, Horth C, Hu B, Bareke E, Chen X, Li Y, Coradin M, Garcia BA,

Lu C, Majewski J: Chromatin dysregulation associated with NSD1 mutation in

head and neck squamous cell carcinoma. *Cell Rep* 2021, 34:108769.

16. Lozier ER, Konovalov FA, Kanivets IV, Pyankov DV, Koshkin PA, Baleva LS,

Sipyagina AE, Yakusheva EN, Kuchina AE, Korostelev SA: De novo nonsense

mutation in WHSC1 (NSD2) in patient with intellectual disability and dysmorphic

features. *J Hum Genet* 2018, 63:919-922.

17. Boczek NJ, Lahner CA, Nguyen TM, Ferber MJ, Hasadsri L, Thorland EC, Niu Z,

Gavrilova RH: Developmental delay and failure to thrive associated with a loss-

of-function variant in WHSC1 (NSD2). *Am J Med Genet A* 2018, 176:2798-2802.

18. Derar N, Al-Hassnan ZN, Al-Owain M, Monies D, Abouelhoda M, Meyer BF,

Moghrabi N, Alkuraya FS: De novo truncating variants in WHSC1 recapitulate the

Wolf-Hirschhorn (4p16.3 microdeletion) syndrome phenotype. *Genet Med* 2019,

21:185-188.

19. Lhoumaud P, Badri S, Rodriguez-Hernaez J, Sakellaropoulos T, Sethia G,

Kloetgen A, Cornwell M, Bhattacharyya S, Ay F, Bonneau R, et al: NSD2

overexpression drives clustered chromatin and transcriptional changes in a

subset of insulated domains. *Nat Commun* 2019, 10:4843.

20. Yuan G, Flores NM, Hausmann S, Lofgren SM, Kharchenko V, Angulo-Ibanez M,

Sengupta D, Lu X, Czaban I, Azhibek D, et al: Elevated NSD3 histone

methylation activity drives squamous cell lung cancer. *Nature* 2021, 590:504-

508.

21. Sun Y, Xie J, Cai S, Wang Q, Feng Z, Li Y, Lu JJ, Chen W, Ye Z: Elevated

expression of nuclear receptor-binding SET domain 3 promotes pancreatic

cancer cell growth. *Cell Death Dis* 2021, 12:913.

22. Satterstrom FK, Kosmicki JA, Wang J, Breen MS, De Rubeis S, An JY, Peng M,

Collins R, Grove J, Klei L, et al: Large-Scale Exome Sequencing Study

Implicates Both Developmental and Functional Changes in the Neurobiology of

Autism. *Cell* 2020, 180:568-584.e523.

23. Xu B, Qin T, Yu J, Giordano TJ, Sartor MA, Koenig RJ: Novel role of ASH1L

histone methyltransferase in anaplastic thyroid carcinoma. *J Biol Chem* 2020,

295:8834-8845.

24. Zhu L, Li Q, Wong SH, Huang M, Klein BJ, Shen J, Ikenouye L, Onishi M,

Schneidawind D, Buechele C, et al: ASH1L Links Histone H3 Lysine 36

Dimethylation to MLL Leukemia. *Cancer Discov* 2016, 6:770-783.

25. Yuan W, Xu M, Huang C, Liu N, Chen S, Zhu B: H3K36 methylation antagonizes

PRC2-mediated H3K27 methylation. *J Biol Chem* 2011, 286:7983-7989.

26. Li Y, Chen X, Lu C: The interplay between DNA and histone methylation:

molecular mechanisms and disease implicationsT. *EMBO Rep* 2021, 22:e51803.

27. Weinberg DN, Rosenbaum P, Chen X, Barrows D, Horth C, Marunde MR,

Popova IK, Gillespie ZB, Keogh MC, Lu C, et al: Two competing mechanisms of

DNMT3A recruitment regulate the dynamics of de novo DNA methylation at

PRC1-targeted CpG islands. *Nat Genet* 2021, 53:794-800.

28. Shirane K, Lorincz M: Epigenetic Mechanisms Governing Female and Male

Germline Development in Mammals. *Sex Dev* 2022, 16:365-387.

29. Reznikoff CA, Brankow DW, Heidelberger C: Establishment and characterization

of a cloned line of C3H mouse embryo cells sensitive to postconfluence inhibition

of division. *Cancer Res* 1973, 33:3231-3238.

30. Tang QQ, Otto TC, Lane MD: Commitment of C3H10T1/2 pluripotent stem cells

to the adipocyte lineage. *Proc Natl Acad Sci U S A* 2004, 101:9607-9611.

31. Lu C, Jain SU, Hoelper D, Bechet D, Molden RC, Ran L, Murphy D, Venneti S,

Hameed M, Pawel BR, et al: Histone H3K36 mutations promote sarcomagenesis

through altered histone methylation landscape. *Science* 2016, 352:844-849.

32. Rayasam GV, Wendling O, Angrand PO, Mark M, Niederreither K, Song L,

Lerouge T, Hager GL, Chambon P, Losson R: NSD1 is essential for early post-

implantation development and has a catalytically active SET domain. *EMBO J*

2003, 22:3153-3163.

33. Li Y, Trojer P, Xu CF, Cheung P, Kuo A, Drury WJ, Qiao Q, Neubert TA, Xu RM,

Gozani O, Reinberg D: The target of the NSD family of histone lysine

methyltransferases depends on the nature of the substrate. *J Biol Chem* 2009,

284:34283-34295.

34. Tanaka Y, Katagiri Z, Kawahashi K, Kioussis D, Kitajima S: Trithorax-group

protein ASH1 methylates histone H3 lysine 36. *Gene* 2007, 397:161-168.

35. Yates AD, Achuthan P, Akanni W, Allen J, Alvarez-Jarreta J, Amode MR,

Armean IM, Azov AG, Bennett R, Bhai J, et al: Ensembl 2020. *Nucleic Acids Res*

2020, 48:D682-D688.

36. Miyazaki H, Higashimoto K, Yada Y, Endo TA, Sharif J, Komori T, Matsuda M,

Koseki Y, Nakayama M, Soejima H, et al: Ash1l methylates Lys36 of histone H3

independently of transcriptional elongation to counteract polycomb silencing.

PLoS Genet 2013, 9:e1003897.

37. Rajagopalan KN, Chen X, Weinberg DN, Chen H, Majewski J, Allis CD, Lu C:

Depletion of H3K36me2 recapitulates epigenomic and phenotypic changes

induced by the H3.3K36M oncohistone mutation. *Proc Natl Acad Sci U S A* 2021,

118.

38. Sun Z, Lin Y, Islam MT, Koche R, Hedehus L, Liu D, Huang C, Vierbuchen T,

Sawyers CL, Helin K: Chromatin regulation of transcriptional enhancers and cell

fate by the Sotos syndrome gene NSD1. *Mol Cell* 2023, 83**:**2398-2416.e2312.

39. Sengupta D, Zeng L, Li Y, Hausmann S, Ghosh D, Yuan G, Nguyen TN, Lyu R,

Caporicci M, Morales Benitez A, et al: NSD2 dimethylation at H3K36 promotes

lung adenocarcinoma pathogenesis. *Mol Cell* 2021, 81**:**4481-4492.e4489.

40. Tanaka H, Igata T, Etoh K, Koga T, Takebayashi SI, Nakao M: The

NSD2/WHSC1/MMSET methyltransferase prevents cellular senescence-

associated epigenomic remodeling. *Aging Cell* 2020, 19**:**e13173.

41. Shen C, Ipsaro JJ, Shi J, Milazzo JP, Wang E, Roe JS, Suzuki Y, Pappin DJ,

Joshua-Tor L, Vakoc CR: NSD3-Short Is an Adaptor Protein that Couples BRD4

to the CHD8 Chromatin Remodeler. *Mol Cell* 2015, 60**:**847-859.

42. Brown MA, Sims RJ, Gottlieb PD, Tucker PW: Identification and characterization

of Smyd2: a split SET/MYND domain-containing histone H3 lysine 36-specific

methyltransferase that interacts with the Sin3 histone deacetylase complex. *Mol*

Cancer 2006, 5:26.

43. Zhang Y, Fang Y, Tang Y, Han S, Jia J, Wan X, Chen J, Yuan Y, Zhao B, Fang

D: SMYD5 catalyzes histone H3 lysine 36 trimethylation at promoters. *Nat*

Commun 2022, 13:3190.

44. Wickham, H: ggplot2: Elegant Graphics for Data Analysis. Use R! Springer New

York, NY; 2009. doi:10.1007/978-0-387-98141-3

45. Lopez-Delisle L, Rabbani L, Wolff J, Bhardwaj V, Backofen R, Grüning B,

Ramírez F, Manke T: pyGenomeTracks: reproducible plots for multivariate

genomic datasets. *Bioinformatics* 2021, 37:422-423.

46. Li H, Durbin R: Fast and accurate short read alignment with Burrows-Wheeler

transform. *Bioinformatics* 2009, 25:1754-1760.

47. Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A,

Keane T, McCarthy SA, Davies RM, Li H: Twelve years of SAMtools and

BCFtools. *Gigascience* 2021, 10.

48. Seemann, T. (2020). Samclip. https://github.com/tseemann/samclip

49. Quinlan AR, Hall IM: BEDTools: a flexible suite of utilities for comparing genomic

features. *Bioinformatics* 2010, 26:841-842.

50. Ramírez F, Dündar F, Diehl S, Grüning BA, Manke T: deepTools: a flexible

platform for exploring deep-sequencing data. *Nucleic Acids Res* 2014, 42:W187-

191.

51. Amemiya HM, Kundaje A, Boyle AP: The ENCODE Blacklist: Identification of

Problematic Regions of the Genome. *Sci Rep* 2019, 9:9354.

52. Sheffield, N. C. & Bock, C. LOLA: enrichment analysis for genomic region sets

and regulatory elements in R and Bioconductor. *Bioinformatics* 2016. 32:587-

589.

53. Zerbino DR, Wilder SP, Johnson N, Juettemann T, Flicek PR: The ensembl

regulatory build. *Genome Biol* 2015, 16:56.

54. Shpynov O, Dievskii A, Chernyatchik R, Tsurinov P, Artyomov MN: Semi-

supervised peak calling with SPAN and JBR genome browser. *Bioinformatics*

2021, 37:4235-4237.

55. Yu G, Wang LG, He QY: ChIPseeker: an R/Bioconductor package for ChIP peak

annotation, comparison and visualization. *Bioinformatics* 2015, 31:2382-2383.

56. Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, Cheng JX, Murre C,

Singh H, Glass CK: Simple combinations of lineage-determining transcription

factors prime cis-regulatory elements required for macrophage and B cell

identities. *Mol Cell* 2010, 38:576-589.

57. Liao Y, Smyth GK, Shi W: featureCounts: an efficient general purpose program

for assigning sequence reads to genomic features. *Bioinformatics* 2014, 30:923-

930.

58. Love MI, Huber W, Anders S: Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. *Genome Biol* 2014, 15:550.

59. Corces MR, Trevino AE, Hamilton EG, Greenside PG, Sinnott-Armstrong NA,

Vesuna S, Satpathy AT, Rubin AJ, Montine KS, Wu B, et al: An improved ATAC-

seq protocol reduces background and enables interrogation of frozen tissues.

Nat Methods 2017, 14:959-962.

60. Bolger AM, Lohse M, Usadel B: Trimmomatic: a flexible trimmer for Illumina

sequence data. *Bioinformatics* 2014, 30:2114-2120.

61. Langmead B, Salzberg SL: Fast gapped-read alignment with Bowtie 2. *Nat*

Methods 2012, 9:357-359.

62. Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, Nusbaum

C, Myers RM, Brown M, Li W, Liu XS: Model-based analysis of ChIP-Seq

(MACS). *Genome Biol* 2008, 9:R137.

63. Li, Q. H., Brown, J. B., Huang, H. Y. & Bickel, P. J. Measuring Reproducibility of

High-Throughput Experiments. *Ann Appl Stat 5* 2011. 3:1752-1779.

64. Dobin, A. et al: STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 2013.

29:15-21.

65. Lawrence, M. et al. Software for computing and annotating genomic ranges.

PLoS Lawrence M, Huber W, Pagès H, Aboyoun P, Carlson M, Gentleman R,

Morgan MT, Carey VJ: Software for computing and annotating genomic ranges.

PLoS Comput Biol 2013, 9:e1003118.

66. Yuan ZF, Sidoli S, Marchione DM, Simithy J, Janssen KA, Szurgot MR, Garcia

BA: EpiProfile 2.0: A Computational Platform for Processing Epi-Proteomics

Mass Spectrometry Data. *J Proteome Res* 2018, 17:2533-2541.

67. MacLean B, Tomazela DM, Shulman N, Chambers M, Finney GL, Frewen B,

Kern R, Tabb DL, Liebler DC, MacCoss MJ: Skyline: an open source document

editor for creating and analyzing targeted proteomics experiments. *Bioinformatics*

2010, 26:966-968.

Materials and methods

Cell culture, CRISPR-Cas9 gene editing and generation of stable cell lines

C3H10T1/2 mouse mesenchymal stem cells (ATCC) were cultured in DMEM with 10%

fetal bovine serum (FBS,) and supplemented with 1% GlutaMax. *Drosophila* S2 cells

were cultured in Schneider's *Drosophila* medium (ThermoFisher) containing 10% heat-

inactivated FBS. All cell lines tested negative for mycoplasma contamination. To

generate knockout cell lines, 3.5 x 10^5 C3H10T1/2 cells were nucleofected with

ribonucleoprotein (RNP)-mediated CRISPR-Cas9 using the Alt-R CRISPR-Cas9

System (IDT) and Lonza Amaxa® SE Cell Line 4D Nucleofector kit (V4XC-1032,

program CA-133). Synthetic crRNA guides were designed (Supplementary Table 1) and

combined with Alt-R® CRISPR-Cas9 tracrRNA, ATTO 550 (IDT) and coupled to Alt-R®

S.p. Cas9 Nuclease V3 following the IDT protocol "Delivery of ribonucleoprotein

complexes using the Lonza® Nucleofector System" prior to nucleofection. The

transfected cells were incubated for 48h. Single ATTO 550 positive cells were then

sorted into 96-well plates. Individual clones were then expanded and validated by

MiSeq sequencing using specific primers for target loci (Supplementary Table 1). Three

biological clones were chosen and expanded for each knockout cell line, and used in all

subsequent assays. C3H10T1/2 cells overexpressing the H3K36M mutation were

provided by the lab of Dr. Chao Lu (Stanford University), and C3H10T1/2 NSD1/2-DKO

and C3H10T1/2 NSD1/2-SETD2-TKO cells were provided by the lab of Dr. C. David

Allis (Rockefeller University).

Crosslinking and ChIP-seq

Approximately 20 million mMSC cells per cell line were used. Crosslinking was performed in 150 mm cell culture plates using 1% formaldehyde (Sigma) at room temperature with gentle rocking for 10 minutes. The crosslinking reaction was quenched using 1.25 M Glycine with gentle rocking at room temperature for 5 minutes. Fixed cell preparations were washed with ice-cold PBS, scraped, and then washed twice more with ice-cold PBS. Crosslinked pellets were resuspended in 500 µl cell lysis buffer (5 mM pH 8.5 PIPES, 85 mM KCl, 1% (v/v) IGEPAL CA-630, 50 mM NaF, 1 mM PMSF, 1 mM phenylarsine oxide, 5 mM sodium orthovanadate, EDTA-free protease inhibitor tablet) and incubated for 30 minutes on ice. Samples were centrifuged and pellets were resuspended in 500 µl of nuclei lysis buffer (50 mM pH 8.0 Tris-HCl, 10 mM EDTA, 1% (w/v) SDS, 50 mM NaF, 1 mM PMSF, 1 mM phenylarsine oxide, 5 mM sodium orthovanadate, EDTA-free protease inhibitor tablet) and incubated for 30 minutes on ice. Sonication of lysed nuclei was performed using the BioRuptor UCS-300 at maximum intensity for 75-90 cycles (10s on, 20s breaks). Sonication efficiency to achieve fragments between 150 - 500 bp was evaluated using gel electrophoresis on a reversed-crosslinked and purified aliquot from each sample. After sonication, chromatin was diluted to reduce SDS levels to 0.1% and concentrated using Nanosep 10K OMEGA (Pall) columns. Prior to the chromatin immunoprecipitation (ChIP) reaction, 2% sonicated *Drosophila* S2 cell chromatin was spiked into each sample for quantification of total levels of histone marks. The ChIP reactions were performed using the Diagenode SX-8G IP-Star Compact and Diagenode automated iDeal ChIP-seq Kit for Histones. Dynabeads Protein A (Invitrogen) were washed, then incubated with specific antibodies (Supplementary Table 2), 1.5 million cells of sonicated cell lysate, and

protease inhibitors for 10 hrs, followed by a 20 min wash cycle using the provided wash

buffers (Diagenode Immunoprecipitation Buffers, iDeal ChIP-seq Kit for Histones).

Reverse cross-linking was then performed using 5M NaCl at 65☐ for 4 hrs. ChIP

samples were then treated with 2 µl RNase Cocktail at 65☐ for 30 min followed by 2 µl

Proteinase K at 65☐ for 30 min. Samples were then purified with QIAGEN MinElute

PCR purification kit (QIAGEN) as per the manufacturer's protocol. In parallel, input

samples (chromatin from about 50,000 cells) were reverse crosslinked and DNA was

isolated following the same protocol. Library preparation was performed using the Kapa

Hyper Prep library preparation reagents following the manufacturer's protocol (Kapa

Hyper Prep Kit, Roche 07962363001). ChIP libraries were sequenced using the Illumina

HiSeq 4000 at 50 bp single reads or NovaSeq 6000 at 100 bp single reads.

Visualization

Unless otherwise stated, figures were created using ggplot2 (44) v3.3.0.

Coverage/alignment tracks were visualized using pyGenomeTracks (45) v3.2.1.

ChIP-seq processing and analysis

Publicly-available NSD1/2-SETD2-TKO H3K27ac ChIP-seq data was used in this study

(37).

Reads were processed as described (4), where they were aligned using BWA

(46) to a combined reference of mm10 and dm6 and afterwards filtered using a cut-off

of MAPQ < 3 using Samtools (47). Samclip v.0.2 (samtools view -h in.bam | samclip --

ref ref.fa | samtools sort > out.bam) was used to filter contaminated sequences for

H3K36me2 NSD1/2/3-SETD2-QKO and NSD1/2/3-SETD2-ASH1L-QuiKO replicates

prior to alignment (48).

Raw tag counts were binned into windows using bedtools (49) v2.29.0 as

previously described (15) in different sized bins (1kb and 10kb). deepTools (50) v3.3.1

was used to normalize ChIP signals either by dividing the total alignments (in millions)

(bamCoverage -b $BAM -o $OUTPUT.bigWig --normalizeUsing CPM --centerReads -e

200) or additionally taking the log2 ratio of ChIP signals by those of the input

(bamCompare -b1 $BAM_ChIP -b $BAM_Input -o OUTPUT.bigWig -e 200 --

centerReads -bs 200 --blackListFileName $BL_bed --normalizeUsing CPM --

scaleFactorsMethods None). PCA and hierarchical clustering based on the Pearson's

correlation matrix were generated using the log2 ratio of ChIP signals over input values

from 10kb bins. To generate coverage tracks, following CPM normalization as indicated

above, replicates were merged in a stepwise fashion using bigwigCompare from

deepTools with parameters '-b1 rep1 -b2 rep2 $outdir --operation mean -bs 200 -o

$merged.step1.cpm.bw' and '-b1 merged.step1.cpm.bw -b2 rep3 $outdir --operation

mean -bs 200 -o $merged.final.cpm.bw'. Additionally, quantitative normalization entailed

multiplying the original signal (either in CPM or log2 ratio over input) by the genome-

wide modification percentage values obtained from mass spectrometry (these values

were averaged if normalizing merged bigwig tracks), as previously described (15). The

ENCODE blacklist (51) was used.

Gene body distribution plots were generated using computeMatrix using genes

with at least 50000 bp width and an exon count of at least 6. Furthermore, to avoid low

gene counts, only the top 50th percentile (mean counts > 0.012 after normalization

using DESeq2 and adjusted for gene length). Only the first and last three exon/intron

pairs are shown in the plots. Annotated promoters were defined as 1000 bp upstream of

the transcription start sites. The depth-normalized mean signal found in each annotation

was calculated using computeMatrix with parameters 'scale-regions -R $feature.bed -S

$sample.bw -bl $BL_bed -b 0 -a 0 -m 1000 -bs 5 --samplesLabel $sample_name'.

Correlation plots depicting the relationship between H3K36me2 and gene expression

quantiles were created by plotting the average depth-normalized H3K36me signal within

gene bodies, taking into account the exclusion criteria mentioned above. This was

plotted against the average gene expression signal within each quantile.

Enrichment matrices for aggregate plots and heatmaps were generated using

computeMatrix   from   deepTools   for   intergenic   regions   (scale-regions   --

regionBodyLength 20000 --beforeRegionStartlength 20000 --afterRegionStartLength

20000 --binSize 1000), for promoters (reference-point -R $Promoters -bl $BL_bed --

referencePoint center --binSize 50 -a 3000 -b 3000 --missingDataAsZero --skipZeros)

and for enhancers (reference-point -R $Enhancers -bl $BL_bed --referencePoint center

--binSize 50 -a 3000 -b 3000 --missingDataAsZero --skipZeros). Genic regions were

taken as the union of any intervals having the "gene" annotation in Ensembl and

intergenic regions thus defined as the complement of genic ones. The ratio of intergenic

enrichment over neighboring genes was calculated as previously described (15). The

exon-to-intron ratio was determined by utilizing the deepTools plotEnrichment tool to

quantify H3K36me levels in annotated exons and introns sourced from Ensembl.

Subsequently, the counts were normalized based on sample library depth and total

length   of   annotations.   Active   enhancers   were   identified   as   those   containing

parental/wildtype ATAC-seq peaks, while inactive enhancers lacked such peaks. To

ensure equal numbers of active and inactive enhancers, random sampling without

replacement was performed, resulting in a sample size of n = 1758 for active enhancers

and n = 1758 for inactive enhancers. computeMatrix using the parameters 'reference-

point -R $inactive_enhancers $active_enhancers -bl $BL_bed --referencePoint center --

binSize 50 -a 3000 -b 3000 --missingDataAsZero --skipZeros --samplesLabel

$samples_label -o $mat.gz'. The same parameters for computeMatrix were used to

generate heatmaps for inactive and active promoters.

"Peakiness" scores were computed as the average read-depth normalized

coverage of the top 1% most covered 1-kb windows across the genome, excluding

those overlapping blacklisted regions, as previously described (4).

10 kb bins with H3K36me2 signal in NSD1/2-SETD2-TKO not found in

NSD1/2/3-SETD2-QKO were used for overlap enrichment analysis. Overlap enrichment

was determined with all the bins as the background set as implemented in LOLA (52)

v1.16.0 for Ensembl (35) 97 annotations (genes and regulatory build (53)).

SPAN (54) v1.1, a semi-supervised peak-caller, was used to call peaks for

NSD1/2/3-SETD2-QKO samples (java -Xmx8G -jar span.jar analyze -t $ChIP.bam -c

$Control.bam --cs $mm10.chrom.sizes -p $Results.peak). ChIPseeker was applied

(with options: genomicAnnotationPriority = c("Intergenic","Promoter", "Exon","Intron"),

tssRegion = c(-1000, 1000)) to obtain the genomic feature distribution for the intersect

of peaks from NSD1/2/3-SETD2-QKO replicates (55).

Motifs were obtained using HOMER (56) v4.11 for ATAC-seq peaks within

H3K36me2 peaks in the intersection of peaks from NSD1/2/3-SETD2-QKO replicates.

To generate genome-wide correlation plots between genes and H3K36me2

signal, we utilized featureCounts (57) (version 1.5.3) to count H3K36me2 reads within

exons, which were then aggregated for all exons belonging to their corresponding

genes. Subsequently, the raw counts underwent normalization using DESeq2's median

of ratios method (58), scaled according to the total length of exons for each gene and

log transformed after removing any normalized read counts equal to zero.

ChIP-Rx was processed as previously described (31). ChIP-Rx was calculated

for each sample by dividing the ratio of the total ChIP-seq reads mapping to the mm10

over the total ChIP-seq reads mapping to dm6 over the ratio of the total input reads

mapping to mm10 over the total input reads mapping to dm6 (see below). See

Supplementary Table 3 for calculated ChIP-Rx values.

$$ChIP-Rx = \frac{ratio(ChIP)}{ratio(Input)} \quad ratio(X) = \frac{\#\ of\ X\ reads\ mapped\ to\ mm10\ genome}{\#\ of\ X\ reads\ mapped\ to\ dm6\ genome}$$

ATAC-seq

ATAC-Seq library preparation was performed according to the Omni-ATAC protocol

(59). 50,000 C3H10T1/2 cultured cells were resuspended in 1 ml of cold ATAC-seq

resuspension buffer (RSB; 10 mM Tris-HCl pH 7.4, 10 mM NaCl, and 3 mM MgCl2 in

water). Cells were centrifuged at 500 rcf for 5 min in a pre-chilled (4°C) fixed-angle

centrifuge. After centrifugation, supernatant was aspirated and cell pellets were then

resuspended in 50 µl of ATAC-seq RSB containing 0.1% IGEPAL, 0.1% Tween-20, and

0.01% digitonin by pipetting up and down three times. This cell lysis reaction was

incubated on ice for 3 min. After lysis, 1 ml of ATAC-seq RSB containing 0.1% Tween-

20 (without IGEPAL and digitonin) was added, and the tubes were inverted to mix.

Nuclei were then centrifuged for 10 min at 500 rcf in a pre-chilled (4°C) fixed-angle

centrifuge. Supernatant was removed and nuclei were resuspended in 50 uL

transposition mix (2x TD Buffer, 100 nM final transposase, 16.5 uL PBS, 0.5 uL 1%

digitonin, 0.5 uL 10% Tween-20, 5 uL H2O) (Illumina Tagment DNA Enzyme and Buffer

Small Kit, 20034197). Transposition reactions were incubated at 37 °C for 30 min in a

thermomixer with shaking at 1000 rpm. Reactions were cleaned up with DNA Clean and

Concentrator 5 columns (Zymo Research). Illumina Nextera DNA Unique Dual Indexes

Set C (Illumina, 20027215) were added and amplified (12 cycles) using NEBNext 2x

MasterMix. Sequencing of the ATAC-Seq libraries was performed on the Illumina

NovaSeq 6000 system using 100-bp paired-end sequencing.

ATAC-seq processing and analysis

Publicly available parental/wildtype ATAC-Seq data from Rajagopalan et al. (2021) (37)

was used to distinguish between inactive and active enhancers. Raw reads were

trimmed and filtered for quality using Trimmomatic v0.39 (60). Trimmed reads were

mapped to the mm10 genome assembly using Bowtie2 v2.5.1 (61), and non-uniquely

mapping reads were removed. Afterwards, the reads were adjusted by shifting all

positive-strand reads 4 bp downstream and all negative-strand reads 5 bp upstream to

the center of the reads on the transposase binding event. Peak calling was performed

on each replicate using MACS2 v2.2.6 (62) with '-extsize 200 -shift -100 -nomodel'

parameters. To find a set of reproducible peaks across replicates, we calculated the

irreproducible discovery rate (IDR) (63) and excluded peaks with an IDR greater than

0.05 across every pair of replicates. Subsequently, the ENCODE blacklist (51) was

used to filter the peaks. Coverage tracks were generated using bamCoverage with

parameters '-b $BAM -o $OUTPUT.bigWig --normalizeUsing CPM --centerReads -e 200

--minMappingQuality 5 -bs 200''.

RNA-seq processing and analysis

Publicly available RNA-seq data in mMSC was used for this study, specifically three

replicates of parental C3H10T1/2 mouse mesenchymal stem cells (37). Raw reads were

aligned to mm10 genome build using STAR version 2.5.3a (64). Afterwards,

featureCounts (57) (version 1.5.3) was used to count exonic reads from the GTF

annotation (GENCODE version from UCSC). Expression was normalized using

DESeq2's median of ratios (58), and divided by the aggregate length of exons per gene.

GENCODE VM25 gene annotation was used to filter for transcripts that are at least

50000 bp and have at least 6 exons. Genes consisting of these transcripts were

categorized into five groups based on their expression levels, which was calculated by

taking the average of parental replicates. After filtering for genes below the 50th

percentile (normalized counts greater than 0.01), the remaining genes were divided into

four quantiles and hence four groups of genes based on expression level. A fifth group

of genes were designated that had zero expression. The transcripts for each of these

genes were then further subdivided into 10 kb upstream of the promoter, promoter

region, the first three exon/intron pairs, the last three exon/introns pairs, and lastly, 10

kb downstream of the last exon. Enrichment matrices were then generated for each

feature and for each gene expression quantile using deepTools' computeMatrix

(computeMatrix scale-regions -R $Feature.bed -S $Sample.bigWig -bl $BL.bed -b 0 -a 0

-m 1000 -bs 5 -o $output.mat.gz).

For obtaining "Active promoters" and "Inactive promoters", the remaining genes

after filtering for zero reads were divided into three quantiles. The four groups were

randomly downsampled to the group with the lowest number of genes. Finally,

promoters were obtained using GenomicRanges::promoters() v3.17 (65) for these four

groups of genes. Here, the "Active promoters" are referred to as an aggregate of the top

two groups with the highest gene expression and hence most active promoters,

"Inactive promoters" are the remaining two groups with low gene expression and least

active promoters. To ensure equal numbers of active and inactive promoters, random

sampling without replacement was performed, resulting in a sample size of n = 18,290

for active promoters and n = 18,290 for inactive promoters.

Histone acid extraction, histone derivatization, and analysis of post-translational

modifications by nano-LC-MS

4 million cells from each clonal mMSC line were collected and frozen at -80□. Thawed

pellets were lysed in nuclear isolation buffer (15 mM Tris pH 7.5, 60 mM KCl, 15 mM

NaCl, 5 mM MgCl2, 1 mM CaCl2, 250 mM sucrose, 10 mM sodium butyrate, 0.1% (v/v)

beta-mercaptoethanol, commercial phosphatase and protease inhibitor cocktail tablets)

containing 0.3% NP-40 alternative on ice for 5 min. Nuclei were subsequently washed

twice in the same buffer without NP-40, and pellets were resuspended using gentle

vortexing in chilled 0.4 N H2SO4, followed by a 3 hr incubation while rotating at 4□.

After centrifugation, supernatants were collected and proteins were precipitated in 20%

TCA overnight at 4□, washed with 0.1% HCl (v/v) acetone once, followed by two

washes with acetone alone. Histones were resuspended in deionized water. Acid-

extracted histones (20 µg) were resuspended in 50 mM ammonium bicarbonate (pH

8.0), derivatized using propionic anhydride and digested with trypsin as previously

described (15). After the second round of propionylation, the resulting histone peptides

were desalted using C18 Stage Tips, dried using a centrifugal evaporator and

reconstituted using 0.1% formic acid in preparation for LC-MS analysis. Nanoflow liquid

chromatography was performed using a Thermo Fisher Scientific, Vanquish Neo

UHPLC equipped with an Easy-Spray™ PepMap™ Neo nano-column (2 µm, C18, 75

µm X 150 mm). Buffer A was 0.1% formic acid and Buffer B was 0.1% formic acid in

80% acetonitrile. Peptides were resolved using at room temperature with a mobile

phase consisting of a linear gradient from 1 to 45% solvent B (0.1% formic acid in 100%

acetonitrile) in solvent A (0.1% formic acid in water) over 85 mins and then 45 to 98%

solvent B over 5 mins at a flow rate of 300 nL/min. The HPLC was coupled online to an

Orbitrap Exploris 240 (Thermo Scientific) mass spectrometer operating in the positive

mode using a Nanospray Flex Ion Source (Thermo Fisher Scientific) at 1.9 kV. A full

MS scan was acquired in the Orbitrap mass analyzer across 350–1050 m/z at a

resolution of 120,000 in positive profile mode with an auto maximum injection time and

an AGC target of 300%. Parallel reaction monitoring experiments were followed for

monitoring the targeted peptides based on the inclusion list. Targeted ions were

fragmented using HCD fragmentation. These scans typically used an NCE of 30, an

AGC target standard, and an auto maximum injection time. Raw files were analyzed

using EpiProfile 2.0 (66) and Skyline (67). The area for each modification state of a

peptide was normalized against the total signal for that peptide to give the relative

abundance of the histone modification. See Supplementary Tables 4 and 5 for raw m/z

data per modification and calculated MS values, respectively.

Western blotting

For each C3H10T1/2 Western blot sample, 1 million cells were collected and counted

using the automated Countess II cell counter (ThermoFisher). Each cell pellet was

washed with PBS. For histone marks, each pellet was resuspended in 100 µl of 1X

Laemmli lysis buffer, 1:100 proteinase inhibitor cocktail, and 0.1 mM PMSF (Laemmli

6X stock contains 0.35M Tris-HCl pH 6.8, 30% glycerol, 10% SDS, 20% beta-

mercaptoethanol, 0.04% bromophenol blue, in water). Samples were then sonicated

using the Bioruptor UCD-300 for 10 cycles at max intensity (30s on, 20s breaks), and

ran on stain-free TGX 4%-15% gradient pre-cast gels (Bio-Rad, 4568084). Semi-dry

electrotransfer onto a PVDF membrane was completed using the trans-blot Turbo

Transfer system (Bio-Rad, Trans-blot® Turbo RTA Mini LF PVDF Transfer Kit,

1704274). PVDF membranes were blocked with 5% BSA in TBSt, and probed overnight

with the respective primary antibody (Supplementary Table 2). Following washes with

TBSt, membranes were incubated for 1h with anti-rabbit horseradish peroxidase-

conjugated secondary antibody in 2% BSA-TBSt. Imaging was completed using the

ECL Clarity or Clarity Max solutions (Bio-Rad, 1705060, 1705062). Relative intensities

quantified using the Bio-Rad Image Lab™ Software (RRID:SCR_014210).

Additional files:

Supplementary Table 1 (.xlsx): CRISPR gRNA and validation primer sequences

Supplementary Table 2 (.xlsx): Antibody names and relevant manufacturer information

Supplementary Table 3 (.xlsx): Calculated ChIP-Rx values

Supplementary Table 4 (.xlsx): Raw mass spectrometry m/z data

Supplementary Table 5 (.xlsx): Calculated mass spectrometry values

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

The publicly-available NSD1/2-SETD2-TKO H3K27ac ChIP-seq data, Parental ATAC-

seq and Parental RNA-seq data were obtained from the National Center for

Biotechnology Information Gene Expression Omnibus with accession number

GSE160266 (37). Newly generated mMSC sequencing data (ChIP-Seq and ATAC-Seq)

are available in the National Center for Biotechnology Information Gene Expression

Omnibus with accession number GSE243566. Link to access data:

https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE243566. Reviewer token to

access data: yngpggymtzuvjgj

Competing interests

The authors declare that they have no competing interests.

Funding

J.M. B.G. and C.L. are supported by the United States National Institutes of Health (NIH) Grant P01-CA196539. B.G. C.L. is supported by the following NIH grants: R35GM138181 and R01CA266978. B.G. is supported by the St. Jude Chromatin Consortium grant and NIH grant R01HD106051. G.S. was supported by the Gershman Memorial and Kangles Fellowships through the McGill University Faculty of Medicine and Health Sciences. R.P. is supported by the Fonds de Recherche du Québec Santé.

Authors' contributions

G.S. and R.P. contributed equally to this work. Study design, G.S., R.P., and J.M. Writing (original draft), G.S., R.P., and J.M. Laboratory experiments, G.S. Bioinformatic analyses, R.P. and B.H. Data processing, R.P. and E.B. Mass spectrometry and subsequent analysis, F.V. and J.G. Supervision, J.M., B.G., and C.L. Funding acquisition, J.M., B.G., and C.L. All author(s) read and approved the final manuscript.

Acknowledgements

We would like to thank all members of the J.M., B.G., and C.L. groups for their input and feedback. We thank the McGill University Genome Centre for their expertise in sample quality verification and sequencing services. We thank the group of C.L. for generation

1020   of C3H10T1/2 cells overexpressing the H3K36M mutation and Dr. Daniel Weinberg for

1021   earlier work that generated many of the resources that became the core of this

1022   research. We owe an unmeasurable debt of gratitude to Dr. David Allis who, although

1023   no longer with us, continues to inspire our investigations of chromatin biology.

Figures

Fig. 1 The three H3K36me states have distinct genome-wide distributions. **a.** Genome-

browser representation of the three H3K36me states in wildtype C3H10T1/2 mMSCs,

illustrating the broad distribution of H3K36me2 in genic and intergenic regions, while

H3K36me1 and H3K36me3 are predominantly distributed within genic regions. **b.** Bulk

levels of H3K36me1/2/3 as quantified by MS, with H3K36me2 showing the highest

abundance followed by H3K36me1 and H3K36me3. The error bars indicate mean ±

standard deviation. Individual data points represent biological replicates (n=3 per

methylation state). **c.** Heatmaps showing H3K36me2 (input- and depth-normalized)

enrichment patterns ± 20 kb flanking intergenic regions (IGRs), indicating that at IGRs

(n = 9551), there are high levels of H3K36me2, low levels of H3K36me1 and lack of

H3K36me3. **d.** Genome-browser coverage tracks demonstrating that H3K36me1/2 are

highest at 5' regions, and then decrease towards the 3' end, where they are replaced by

H3K36me3 in regions of high exon density. **e.** Representative view of genic

H3K36me1/2/3 distributions, highlighting the 5' to 3' distribution patterns, exonic versus

intronic signal, and dependence on gene expression for the three H3K36 methylation

states. Only transcripts with at least 50000 base-pairs (bp) and 6 exons were used in

the analysis. An aggregate of H3K36me signals on the first three exons and the last

three exons are shown. The expression quantiles were calculated based on normalized

(reads per kilobases) expression counts from the parental samples. Expression quantile

4 comprises transcripts with the highest expression, expression quantile 1 comprises

transcripts with the lowest expression greater than zero, and expression quantile 0

comprises transcripts with zero counts.

1087

Fig. 2 Distinct alterations in H3K36me global abundance are found following single and

combinatorial knockouts of K36MTs. **a.** Genome-wide changes to H3K36me1/2/3 in

various conditions as quantified by MS. Individual dots denote biological replicates

within a condition (n=3 per condition, except for QuiKO where an outlier was omitted for

H3K36me1). Error bars indicate mean ± standard deviation. **b.** Quantitative Western

blots demonstrating changes to H3K36me1 (left) and H3K36me2 (right) in the various

knockout conditions relative to parental cells. Immunoblot and protein gel cropped for

relevant regions. Full total protein loading gel used for quantification of relative intensity

illustrated in Fig.S3b. **c.** Principal component analysis (PCA) plot of depth-normalized

log2 H3K36me ChIP-Seq signal over input based on 10 kilobase (kb) binned signals

(after scaling and centering). Bins with raw read counts consistently lower than 100

across all samples were excluded. Each data point denotes a biological replicate. The

colors represent the different conditions. Circles represent the parental and single-KO

samples, and the triangles represent the multiple-KO samples and H3K36M-OE.

1104
1105
1106

1107

Fig. 3 Effects of K36MT ablations and H3K36M-OE on H3K36me1 distribution patterns

within genes. **a** Genome-browser tracks of MS-normalized H3K36me1 signal showing

depletion in H3K36M-OE and multi-knockout conditions, and an increase in SETD2-KO

cells. The two bottom tracks illustrates that despite NSD1/2/3-SETD2-QKO and

NSD1/2/3-SETD2-ASH1L-QuiKO displaying reduced amounts of H3K36me1, the

distribution patterns remain similar to that observed in parental cells. **b**. Boxplots of

H3K36me1 genic to intergenic ratios, indicating that the increase in SETD2-KO cells is

found primarily within genic regions. (n=3 per condition) **c**. Boxplots showing the exon

versus intron H3K36me1 signal in parental and SETD2-KO cells, indicating increased

signal at exons following SETD2-KO. (n=3 per condition).

1118

1119
Fig. 4 Intergenic H3K36me2 decreases following NSD1-KO, and is further reduced in

H3K36M-OE and multi-knockout conditions. **a.** Coverage tracks of MS-normalized

H3K36me2 signal centered on intergenic regions flanked by genic regions, highlighting

reduced intergenic signal following loss of NSD1, more pronounced reductions in

NSD1/2-DKO and H3K36M-OE cells, and the most significant depletions in subsequent

multi-knockout conditions. **b.** Heatmaps showing H3K36me2 (input- and depth-

normalized) enrichment patterns ± 20 kb flanking IGRs, summarizing the H3K36me2

trends for multiple cell lines at IGRs. **c**. H3K36me2 "peakiness scores", representing the

average ChIP signal in the top 1% of 1-kb bins over the total signal, indicating more

punctate distributions of H3K36me2 in multi-knockout conditions. Individual data points

represent biological replicates for each condition (n=3 per condition, except for DKO

and K36M-OE where n=2). Error bars represent mean ± standard deviation.

1132
1133
1134
1135

1136

Fig. 5 Genic H3K36me2 distribution patterns are altered following SETD2-KO, NSD1/2-

DKO, and H3K36M-OE. **a.** Coverage tracks of MS-normalized H3K36me2 signal

showing changes in genic distribution following specific K36MT knockouts. **b**. Zoomed

in view of H3K36me2 distribution in gene bodies, highlighting changes in exon versus

intron signal and dependence on gene expression following K36MT deletions. Only

transcripts with at least 50000 base-pairs (bp) and 6 exons were used in the analysis.

An aggregate of H3K36me signals on the first three exons and the last three exons are

shown. The expression quantiles were calculated based on normalized (reads per

kilobases) expression counts from the parental samples. Expression quantile 4

comprises transcripts with the highest expression, expression quantile 1 comprises

transcripts with the lowest expression greater than zero whereas expression quantile 0

comprises transcripts with zero counts. **c.** Correlation plots of H3K36me2 CPM and

gene expression quantiles, indicating a shift to significant positive correlations between

H3K36me2 and gene expression following SETD2-KO, NSD12-DKO and H3K36M-OE.

d. Boxplots of mean H3K36me2 signal at exons versus introns, indicating a trend

towards greater signal at exons compared to introns following SETD2-KO, NSD12-DKO

and H3K36M-OE.

1154

Fig. 6 NSD3 deposits H3K36me2 specifically at active promoters and enhancers. **a.** Genome-browser tracks displaying MS-normalized H3K36me2 signal where the distribution becomes more punctate with additional K36MT deletions. **b.** Genome-

browser coverage tracks showing NSD1/2-SETD2-TKO (TKO) H3K36me2 signal at

active enhancer regions, which are identified by the presence of ATAC-seq and

H3K27ac peaks. **c.** Genome-browser coverage tracks displaying H3K36me2 signal for

NSD1/2-SETD2-TKO (TKO) at active promoter regions, which are identified by the

presence of ATAC-seq and H3K27ac peaks. **d.** Overlap enrichment results of Ensembl

annotations with bins found in TKO but not in QKO (i.e., identified as NSD3-catalyzed

H3K36me2). The size of the dots corresponds to the number of bins overlapping the

corresponding annotation. **** represents p-value < 1e-4 whereas *** represents p-

value < 1e-3 based on Fisher's exact test of bins overlapping a specific class of

annotated regions versus a background of all bins in TKO. Only the top four most

significant annotations are shown. **e.** Heatmaps showing H3K36me2 (input- and depth-

normalized as well as MS-scaled) signal centered on transcription start site (TSS) in

inactive (n=18,290) and active promoters (n=18,290) as well as H3K36me2 signal

centered on inactive (n=1758) and active enhancers (n=1758), highlighting the loss of

H3K36me2 signal at these active regions following NSD3 knockout.

1174

Fig. 7 ASH1L-mediated H3K36me2 deposition is targeted to regulatory elements of developmentally-relevant genes. **a** Coverage tracks of MS-normalized H3K36me2 signal for QKO and QuiKO, depicting a region of punctate peaks remaining in QKO cells that are depleted upon subsequent loss of ASH1L. Zooming into one of the peaks shows that they are still quite broad (10-70 kb). **b** Pie chart depicting the distribution of remaining H3K36me2 peaks in QKO cells. Only overlapping peaks from three biological

1183   replicates were used in the analysis. **c** Enriched motifs found in open chromatin regions

1184   (as marked by ATAC-seq peaks) within H3K36me2 domains in QKO cells, indicating

1185   potential recruitment factors for ASH1L.

1186

Fig. 8 The K36MTs have distinct genomic preferences in the deposition of H3K36me. **a.** SETD2 is coupled to the phosphorylated RNAPII complex (RNA Pol II) and deposits

H3K36me3 at exons with higher enrichment in 3' regions where exon density becomes

higher. This takes place at the expense of H3K36me2, which display the opposite 5' to

3' and exon/intron trends. **b.** NSD1 is recruited to active CpG islands and then spreads

outwards into both gene bodies and intergenic regions, depositing broad H3K36me2

domains. **c.** NSD2 has a specific affinity for depositing H3K36me2 within genes, while

also depositing broad intergenic H3K36me2 domains. **d.** NSD3 spreads outwards from

CpG-rich regulatory elements and deposits narrower H3K36me2 peaks. **e.** ASH1L

primarily deposits H3K36me2 at the promoters and enhancers of specific genes.

Created with BioRender.com.

1199
1200 **Supplementary figures**

1201

1202
Fig. S1 Genome-wide characterization of H3K36me indicates distinct patterns within

gene bodies and dependence on transcription. **a**. Boxplots of H3K36me1/2/3

enrichment at exons versus introns, illustrating higher H3K36me1 and H3K36me3

signals at exons whereas the inverse is found for H3K36me2. **b.** Correlation plots

illustrating the relationship between H3K36me1/2/3 signal and gene expression

quantiles. Reported values are Pearson's correlation coefficient (*R).*

1209

Fig. S2 Representative genome-browser coverage tracks validating CRISPR gene editing as assessed by MiSeq. gRNAs were designed to target the catalytic SET domain of each H3K36 methyltransferase. Gene names and genomic coordinates of edits are indicated at the bottom of each set of coverage tracks. WHSC1 and WHSC1L1 are the mouse orthologs of human NSD2 and NSD3, respectively. Similar edits were

1217 achieved in multi-knockout conditions (i.e. NSD1/2/3-SETD2-QKO and NSD1/2/3-

1218 SETD2-ASH1L-QuiKO). Three biological clones were obtained for each condition.

1219

1220

Fig. S3 Various K36MT deletions lead to distinct genome-wide alterations to H3K36me.

a. Global abundance of H3K36me1/2/3 as quantified using ChIP-seq with *Drosophila*

spike-in reference chromatin (ChIP-Rx) (as described in methods). Error bars represent

mean ± standard deviation. **b.** Western blots for H3K36me1 (left) and H3K36me2

(middle). Total protein loaded and used for quantification shown below each blot.

Western blot comparing parental and SETD2-KO H3K36me1 levels (right).

Quantification performed as described in methods. **c.** Pearson's correlation matrix-

based hierarchical clustering of H3K36me1/2, illustrating the clustering of single

knockouts with PA whereas H3K36M-OE clusters with the multiple knockouts.

1230

Fig. S4 Distinct H3K36me2 distribution patterns are found following knockout of

K36MTs. **a.** Heatmaps showing H3K36me2 (input- and depth-normalized) enrichment

patterns ± 20 kb flanking IGRs, summarizing the H3K36me2 trends at IGR in parental

and knockout cells. **b.** H3K36me2 "peakiness score", representing the average ChIP

signal in the top 1% of 1-kb bins. More punctate distributions of H3K36me2 are found

following multiple deletions whereas individual knockouts have less effect on peakiness

score. Individual data points represent replicates for each condition. **c.** Correlation plots

illustrating the relationship between H3K36me2 signal and gene expression quantiles.

Reported values are Pearson's correlation coefficient (*R).* With the exception of SETD2-

KO, single knockouts have no effect on the correlation between H3K36me2 and gene

expression. **d.** Genome-wide correlation analysis between exonic H3K36me2 and gene

expression, illustrating stronger correlations following SETD2-KO, NSD1/2-DKO and

H3K36M-OE compared to parental. Only genes with H3K36me2 and gene expression

counts greater than zero were included in the analysis. Both the H3K36me2 and gene

expression exonic read counts (aggregated per gene) were normalized using DESeq2,

scaled based on the cumulative length of all exons within each gene, and then log

transformed.

1249

1250
Fig. S5 NSD1/2/3 are expressed at different relative levels across three different cell

lines. **a-b** Bar graphs showing DESeq2 counts that have been normalized, with each

gene scaled according to its specific gene length. **a.** Only the Pbx gene family, including

Pbx2, is expressed in 10T-mMSCs, as shown in both parental and NSD1/2/3-SETD2-

QKO cell lines. Error bars represent median ± standard error. **b.** Relative expression of

NSD1/2/3 in mouse mesenchymal stem cells (mMSCs), mouse embryonic stem cells

(mESCs), and human head and neck squamous cell carcinoma (HNSCC) cells.