Development of novel genetically-encoded fluorescent probes to track ceramides during phagocytosis

Vladimir Girik[1], Larissa Van Ek[2,3], Isabelle Dentand Quadri[1], Maral Azam[1], Maria Cruz Cobo[1], Marion Mandavit[1], Isabelle Riezman[4], Howard Riezman[4], Anne-Claude Gavin[2,3], Paula Nunes-Hasler[1].

[1]Department of Pathology and Immunology, Geneva Center for Inflammation Research, Faculty of Medicine, University of Geneva, Geneva Switzerland

[2]Department of Cellular Physiology and Metabolism, Faculty of Medicine, University of Geneva, Geneva Switzerland

[3]Diabetes Center, Faculty of Medicine, University of Geneva, Geneva, Switzerland

[4]Department of Biochemistry, NCCR Chemical Biology, Faculty of Science, University of Geneva, Geneva Switzerland

*Corresponding Author:

Paula Nunes-Hasler

Department of Pathology and Immunology

Geneva Center for Inflammation Research

Faculty of Medicine

University of Geneva,

1 Rue Michel-Servet

Geneva Switzerland 1211

Paula.Nunes@unige.ch

Abstract

Ceramides (Cer) and their derivatives have been suggested to regulate phagocytosis, however their exact function remains poorly understood. Genetically-encoded fluorescent biosensors have been immensely useful in deciphering the critical role of phagosomal phosphoinositide regulation, and thus we sought to develop similar tools to examine Cer dynamics during phagocytosis. EGFP fusion of several known Cer-binding domains were generated and expressed in cells. While most constructs localized to the nucleus or cytosol, the CA3 cysteine-rich domain of the pseudokinase KSR1 displayed additionally plasma membrane (PM) or endolysosomal localization, depending on the position of the fluorescent tag and linker. C-terminally tagged KSR1-CA3 (C-KSR) and a more stable version C-KSR-GS responded as expected to Cer depletion and accumulation using a panel of Cer modifying drugs. In contrast, N-terminally tagged N-KSR did not respond to Cer induction by palmitate or glucosylceramide (GlcCer) synthase inhibitor Genz-123346. Liposome microarray analysis suggested N-KSR binding to the Cer derivative GlcCer. Interestingly, the three probes showed distinct dynamics during phagocytosis in model phagocytic fibroblasts. Despite partial lysosomal degradation, C-KSR robustly but transiently re-localized to the PM during phagocytosis, whereas N-KSR re-localized from the PM to the cytoplasm at later timepoints. Moreover, occasional weak recruitment of C-KSR-GS to endoplasmic reticulum and phagosome was enhanced upon overexpression of the endoplasmic reticulum proteins STIM1 and Sec22b, and was more salient in dendritic cells. The data suggest these novel probes may be valuable new members of the growing molecular toolkit used to analyze sphingolipid dynamics and function in living cells.

Introduction

Phagocytosis is a fundamental process whereby external foreign particles are internalized by cells, and the resulting membrane-engulfed compartment, called the phagosome, undergoes a series of maturation steps that lead to the formation of the digestive milieu and degradation of the foreign material (Levin, Grinstein, and Canton 2016). A large body of literature has highlighted the central role of dynamic phosphoinositide lipid remodeling in controlling this process through the recruitment of a variety of effectors including cytoskeletal, motor or vesicular docking proteins that help guide phagosomes to their fate (Levin, Grinstein, and Canton 2016; Yeung and Grinstein 2007; Levin, Grinstein, and Schlam 2015). Sphingolipids are a diverse group of lipids crucial for maintaining cellular membrane structure, fluidity, curvature, that have been long recognized to additionally play bioactive signaling roles (Quinville et al. 2021). Ceramides (Cer), a central hub of the sphingolipid network (Hannun and Obeid 2008; 2018), have also been implicated in various steps of the phagocytic process. These span from facilitation of receptor clustering and activation (Abdel Shakor, Kwiatkowska, and Sobota 2004; Avota, Gulbins, and Schneider-Schaulies 2011), to initial membrane deformation of phagocytic cups (Magenau et al. 2011; Niekamp et al. 2021), to promotion of phago-lysosome fusion (Schramm et al. 2008; Wu et al. 2018; Hinkovska-Galcheva et al. 2005), and inhibition of antigen presentation in dendritic cells (Sallusto et al. 1996; Eva Fuglsang, Louise Boye, and Hanne Frokiaer 2017). However, compared to phosphoinositides, less is known about the dynamics, regulation and roles of Cer and its related lipids during this critical immune process (Saharan, Mehendale, and Kamat 2022).

Arguably, part of the challenge may lie in the difficulty tracking these lipids in living cells (Canals, Salamone, and Hannun 2018). The most reliable methods such as mass spectrometry do not have the spatial resolution required for subcellular tracking (Buchberger et al. 2018), or rely on labor and material intensive organelle isolation that still suffer from contamination of sphingolipid-rich endoplasmic reticulum (ER) membranes due to tightly tethered membrane contact sites (MCS) (Scorrano et al. 2019). In addition, tools such as fluorescent sphingolipid analogs are rapidly metabolized and may partition in membranes differentially (Canals, Salamone, and Hannun 2018). Moreover, permeabilization steps necessary for allowing access of anti-Cer antibodies to intracellular membranes such as phagosomes may also disrupt natural lipid localization (Bjørnestad and Lund 2023). Therefore, an approach enabling detection of Cer in living cells with high spatial resolution is desirable. Genetically encoded lipid probes, typically composed lipid-binding domains coupled to a fluorescent tag, provide an opportunity to investigate subcellular localization, relative accessibility, and dynamics in living cells, and have been extensively employed to dissect phospholipid dynamics during phagocytosis (Hammond et al. 2022). Few described lipid

probes specific for sphingolipids have been published (Yamaji et al. 1998; Bakrač et al. 2010).
Moreover no fluorescent Cer biosensor based on isolated Cer-binding domains have been
described so far, although a range of full-length Cer-binding proteins have been employed
(Bidlingmaier et al. 2016; Bockelmann et al. 2018; Deng et al. 2021; K. Zhou et al. 2018)., and
a GST-fusion of the Cer binding domain of the pseudokinase KSR1 has been used *in vitro*
(Yin et al. 2009). In the current report, we document the development and characterization of
novel putative Cer-binding genetically encoded fluorescent probes. Out of several potential
Cer-binding biosensors, the cysteine-rich domain of the human pseudokinase KSR1 (Yin et
al. 2009) was the most promising candidate. KSR1-based probes responded robustly to
treatments modifying Cer levels in cells, though differences were noted between C- terminally
N-terminally tagged constructs, and probe instability appeared to contribute to this response.
Although weak *in vitro* binding precluded in-depth biochemical characterization, liposome
microarray data indicate that N-KSR constructs may instead be sensitive to glycosylated Cer.
Interestingly, differential probe localization depending on cell type and expression of ER-
localized constructs were documented to occur during phagocytosis in model phagocytes.
Together our data indicate that these constructs may be useful tools for dissecting subcellular
sphingolipid dynamics and warrant further characterization.

Results

KSR1-based probes show membrane-associated localization

To identify ceramide-specific biosensors, four domains that have been previously
demonstrated to bind ceramides *in vitro* and/or *in vivo* were selected: 1) the earmuff domain
(EMD) of the human SET protein [amino acids (aa) 70-226] (Mukhopadhyay et al. 2009); 2)
the N-terminal fragment of atypical human protein kinase C zeta (PRKCZ, also called PKCζ)
containing a lipid-binding C1 domain (aa 123-193); 3) the C20-kD carboxy-terminal domain
(amino acids 405-646) of PRKCZ (Wang et al. 2005; 2009); and 4) the C1 cysteine-rich
domain (CA3, also called CRD, amino acids 317-400) of the human pseudokinase KSR1 (Yin
et al. 2009) (Fig 1A). The selected domains were tagged with EGFP at the N- or C-terminus
(prefixed with N- and C- respectively) and expressed in mouse embryonic fibroblasts (MEFs).
All SET and PRKCZ-based fusion proteins were primarily localized to the nucleus (Fig 1B-C).
A nuclear export signal (NES) was added to C-PRKCZ-C20 and C-SET-EMD chimeras to
release proteins trapped in the nucleus, a strategy previously used for lipid biosensor
construction (Goulden et al. 2019). C-NES-PRKCZ-C20 and C-NES-SET-EMD were mainly
found in the cytosol, though nuclear localization was still noticeable (Fig 1B-C). In contrast,
the expression pattern of KSR1-based constructs depended on the position of the fluorescent

tag. N-KSR1(317-400) (abbreviated N-KSR) was found primarily at the plasma membrane

(PM), while the C-terminally tagged probe (C-KSR) localized to intracellular vesicles some of

which appeared enlarged (Fig 1D). The localization of both N- and C-KSR constructs was

dependent on two conserved cysteines in the C1 domain (C359/C362) whose mutation to

serine was shown previously to decrease ceramide binding *in vitro* (Yin et al. 2009) (Fig 1D).

To potentially increase the affinity for ceramides, a construct expressing two KSR1 CRDs in

tandem fused at the C-terminus to EGFP was also created (abbreviated 2X-C-KSR), following

another strategy previously used for lipid biosensor construction (Hammond et al. 2022).

However, the localization appeared similar to that of N-KSR. When expressed in human Hela

cells, KSR1-, SET-, and PRKCZ-based displayed an expression pattern similar to that of MEFs

(Fig S1). In previous studies, anti-ceramide antibodies primarily stained the plasma membrane

and the Golgi (Krishnamurthy, Dasgupta, and Bieberich 2007; Wang et al. 2009) whereas

exogenously added azido-functionalized ceramides partitioned into the PM and organellar

membranes, including mitochondria and lysosomes (Walter et al. 2017; Götz et al. 2020). Of

all candidate probes tested, only KSR1-based constructs showed salient membrane-

associated staining and were therefore selected for further validation.

KSR probes respond to the changes in ceramide levels

Next, the behavior of KSR probes in response to ceramide depletion or accumulation was

investigated. To induce ceramide depletion, cells were exposed to well-known inhibitors of

sphingolipid biosynthesis: myriocin (Myr), a potent inhibitor of serine-palmitoyltransferase

(SPT) (Miyake et al. 1995), the enzyme which catalyzes the first step in SL biosynthesis; and

fumonisin B1 (FumB1), a selective inhibitor of ceramide synthases (Merrill et al. 2001).

Transfected MEF cells showed significant variation in the expression of KSR probes from one

cell to another, making it challenging to assess the effect of SL inhibitors on probe localization

and abundance. To circumvent this issue, the KSR probes were re-cloned into Sleeping

Beauty transposon vectors (Ivics et al. 2009) and co-transfected with the transposase to

generate MEF stable cell lines. Surprisingly, C-KSR-expressing cells demonstrated a drastic

5-fold decrease in the intracellular fluorescence after a 3-day incubation with Myr (0.5 µM) or

FumB1 (2.5 µM), accompanied by some re-localization to the cytosol in a subset of cells (Fig

2A-B). In comparison, only a 2.5-fold decrease in the probe intensity was observed in N-KSR1

expressing cells after sphingolipid depletion with either drug (Fig 2C-D). As expected, all major

sphingolipids including ceramides (Cer), glycosylated ceramides (including glucosyl and

galactosyl ceramides, collectively termed hexosyl-ceramides, HexCer) and sphingomyelins

(SM), were markedly depleted by Myr treatment as measured by targeted quantitative tandem

mass-spectrometry lipidomic analysis (Fig 2E, Supplementary Data S1) (Guri et al. 2017). In addition, Myr treatment of MEF cells transiently transfected with C-KSR and cytosolic TagRFP did not produce any noticeable effect on TagRFP expression (Fig S2), confirming a specific effect on KSR constructs.

To induce Cer accumulation two methods were employed: treatment with the Cer precursor C16 acyl-chain fatty acid palmitate (Tran et al. 2016; Kiefer et al. 2015) or incubation of purified bacterial sphingomyelinase from *Staphylococcus aureus* (SMase), which releases ceramides by hydrolyzing sphingomyelin on the outer leaflet of the PM (Devaux 1991; Abe and Kobayashi 2014), where ceramides are believed to partition efficiently in both leaflets in eukaryotic cells (Pohl et al. 2009). After a 4h incubation in palmitate complexed to BSA (Palm) a 2.7-fold increase in C-KSR signal intensity was observed compared to cells treated with delipidated BSA alone (Fig 2A-B). However, the N-KSR fluorescence intensity was unchanged, and no obvious changes in localization were observed (Fig2C-D). Lipidomic analysis confirmed that Palm induced a 2-fold accumulation of Cer species, driven mainly by the 4-fold increase in C16-Cer (Fig 2F, Supplementary Data S1). However, unexpectedly depletion of C24-SM (p<0.001), and C24-HexCer (p=0.0144) was also observed (Fig 2F, Supplementary Data S1).

We then treated C-KSR expressing cells with SMase (0.5 U/ml) for 30 min. This led to C-KSR recruitment to the PM (Fig 3A-C), where both the fluorescence intensity at the PM, the total cellular fluorescence or blinded quantification of the fraction of cells displaying PM localization were higher in SMase treated cells compared to PBS-treated controls (p=0.0004, p=0.0007 and p= 0.0001 respectively, Fig 3B-C). Lipidomic analysis confirmed that SMase induced a dramatic 40-fold accumulation of Cer species, driven by C16-Cer (p<0.0001) (Fig 3D, Supplementary Data S1). Total HexCer were not different but increases in C10-HexCer (p=0.0406), a lipid of unknown biological significance, were observed (Fig 3D, Supplementary Data S1). Although total SM was not significantly different, reductions in longer chain C20-C22- and C24-SM were observed (p=0.0052, p=0.0117, p= 0.0052, respectively, Fig 3D, Supplementary Data S1). That cleavage of sphingomyelin was efficient was also confirmed by staining with a purified EGFP-tagged SM-specific probe lysenin (Kwiatkowska et al. 2007) (20 μg/ml) Fig S3A). Since SMase treatment triggered the largest Cer accumulation, as well as salient PM recruitment and global increase in fluorescence with the C-KSR probe, C-NES-PRKCZ-C20 and C-NES-SET-EMD were also examined under conditions of SMase treatment. However, no changes in localization or overall fluorescence were noted (Fig S3B). In addition, since diacylglycerol (DAG) is a molecule with close structural similarity to ceramide that could potentially be generated by activation of SM synthases upon large increases of Cer (Goñi and Alonso 1999; Ekman et al. 2015), the specificity of the C-KSR for ceramide was

further investigated by co-expression C-KSR-mRFP with the DAG probe GFP-PKC-C1(2) from the tandem C1 domains of the rat PKCγ (Codazzi, Teruel, and Meyer 2001). Cells were then treated with either 10 μM ionomycin (Iono) for 10 min, reported to induce PM DAG accumulation (Botelho et al. 2000) or with SMase as above. Whereas Iono treatment but not SMase induced PKC-C1(2) PM recruitment (p<0.0001 Iono, p=0.5214 SMase), the opposite pattern was observed for C-KSR (p=0.1854 Iono, p=0.0065 SMase, Fig 3E-F). Co-expression of the two probes could not explain these observations as treatment with SMase did not affect the localization of PKC-C1(2) when expressed alone, nor did it affect the localization of an alternate DAG probe YFP-DBD (C1b DAG-binding domain of PKCß) (Gallegos, Kunkel, and Newton 2006) (Fig S3C), ruling out potential DAG generation by SMases (Ekman et al. 2015). Collectively, the data suggest that C-KSR is sensitive to the changes in cellular levels of Cer and can detect locally generated ceramides at the PM.

C-KSR is degraded in lysosomes

De novo biosynthesis of Cer occurs in the ER, after which Cer are shipped to the Golgi for conversion into SM and complex sphingolipids. However, Cer can also be generated at the PM, in endolysosomal compartments, or mitochondria via hydrolysis of HexCer or SM (Hannun and Obeid 2018). Thus, to determine whether C-KSR might sense different subcellular pools of Cer we next sought to define in greater detail the organellar distribution of C-KSR. Immunostaining against various organelle markers revealed that C-KSR colocalized preferentially with late endosome Rab7 and lysosome markers LAMP1, but not with recycling endosome-resident Transferrin (Tfn), nor with early endosomal marker Rab5, or Golgi marker TGN38 (Fig 4A-B, Fig S4). Despite the fact that C-KSR appeared to reside on the membrane of enlarged vesicles (for example Fig 1D inset), a closer examination of the LAMP1 staining indicated that LAMP1 was still often peripheral to C-KSR, even in vesicular structures with a dark lumen (Fig 4A, bottom inset arrow). Such structures were reminiscent of autolysosomes, pointing to C-KSR transport into the degradative compartments. Immunoblots of lysates from MEFs stably expressing the KSR probes corroborated this hypothesis, showing extensive degradation of C-KSR1, while protein cleavage was much less evident for N-KSR1 (Fig 4C). Thus, if C-KSR is stabilized or protected from degradation upon membrane binding, pronounced degradation of C-KSR may explain the increase and decrease in global C-KSR fluorescence intensity after treatment with ceramide inducing or depleting drugs. Moreover, C-KSR degradation accounts for the observed colocalization with the late endosomes and lysosomes, though C-KSR was also present in the cytosol and at low levels at the PM in a subset of cells.

Differential response of N- and an improved C-KSR probe

The extensive degradation of C-KSR probe, but not N-KSR, prompted us to scrutinize the sequences of the constructs. The main difference was the short linker regions between the lipid-binding domains and the fluorescent tag. N-KSR contained flexible linkers rich in small or polar amino acids such as glycine and serine (GS linkers), which are widely used for constructing multidomain chimeric proteins (Chen, Zaro, and Shen 2013). For the construction of the C-KSR probe, the lipid-binding domain was subcloned into restriction sites of the pEGFP-N1 backbone, which inadvertently introduced an RLELKLRILQ amino acid motif that targets cytosolic proteins for lysosomal degradation via chaperone-mediated autophagy (CMA) (Majeski and Fred Dice 2004). To render the C-KSR probe more stable, the old linker was replaced with the short flexible GS sequence (GGSSGGGGA), as in the N-KSR (Fig 4D). The resulting chimeric probe was renamed C-KSR-GS. When transiently expressed in MEFs, C-KSR-GS was much less prone to degradation, as demonstrated in the representative immunoblot (Fig 4E). The subcellular localization of the novel C-KSR-GS-EGFP fusion was then examined by confocal microscopy. Images revealed localization to the PM and the cytosol, similar to the expression pattern of the N-KSR and the tandem KSR probes (Fig. 5A-C, compared to Fig 1D). Cells transfected with C-KSR-GS were then exposed to Cer depleting FumB1, or Cer inducing Palm and SMase to determine the response of this new probe to Cer modulation. Similar to the other probes, a reduction in global fluorescence upon treatment with FumB1 (Fig 5A, p=0.039) was observed. Interestingly an increase in global fluorescence upon incubation with Palm, observed with C-KSR but not N-KSR was discerned, though the fold difference was smaller (2.7-fold increase for C-KSR see Fig 2B, and 1.5-fold increase for C-KSR-GS, p=0.048, Fig 5B). Moreover, after treatment with the bacterial SMase, the fraction of cells displaying PM localization increased from approximately 35 to 54% (p<0.0001) indicating that C-KSR1-GS, like C-KSR also responds to ceramide upregulation at the PM (Fig 5C). To further explore potential differences in response to Cer modulation between N-KSR and C-KSR-GS we compared the response of the two probes to reagents previously shown to increase cellular ceramides. To this end cells expressing either probe were treated for 2h with 200 μM Cer analogue C6 (d18:1/6:0) (Kjellberg et al. 2015) or overnight with 200 μM of the GlcCer synthase inhibitor Genz-123346 (Genz) (H. Zhao et al. 2007) or their respective vehicle controls. Interestingly, whereas both probes showed a robust increase in cellular fluorescence upon treatment with C6 (p=0.011 for C-KSR-GS, p=0.0031 for N-KSR, Fig 5D-E) as compared to vehicle, they again displayed a striking differential response to Genz treatment, where an increase in fluorescence was observed wtih C-KSR-GS (p=0.0413) but not with N-KSR (p=0.3526) (Fig 5D-E). Together these data suggest that all 3 KSR-based

probes may have an inherent instability that is mitigated upon membrane binding, which explains the global fluorescent response to Cer modifying drugs, but that N- and C-terminal tagging may nevertheless affect the nature of the lipid (or lipids) bound.

Analysis of lipid binding by KSR probes

In light of the intriguing differences in localization and responses to Palm and Genz treatments, we next sought to further characterize the lipid-binding profile of N- and C-KSR-based probes. To this end we employed a high-throughput liposome microarray (LiMa) technology, where arrays of liposomes (small artificial vesicles) of various compositions are deposited inside a microfluidic chip, exposed to purified proteins or cell lysates, and binding is assessed by automated microscopy (Saliba et al. 2014). To obtain purified probes, the N-KSR and C-KSR probes were re-cloned into super-folding GFP (sfGFP) bacterial expression vectors (Pédelacq et al. 2006) and purified from *E.coli* cultures, where N-KSR was easily obtained but C-KSR had a tendency to aggregate, though smaller soluble quantities were still obtained. DOPC (dioleoyl-phosphatidylcholine), with fluorescent 0.1% phosphatidylethanolamine (PE)-Atto647 and 0.5% PE-PEG350 stabilizer (Magic lipid) served as a backbone for liposomes containing 2, 5 or 10% bovine Cer mixture (CER), chicken SM mixture (SM), C17-glucosyl(β)-ceramide (d18:1/17:0, GlcCer), Sphingosine (d18:1, Sph), Ceramide-1-Phosphate (16:0, C1P) or DO-PI(4,5)P2 (dioleoyl-phosphatidylionositol-(4,5)-bisphosphate). It should be noted that Cer and Sph have been previously reported to destabilize membranes or form aggregates (Veiga et al. 1999; Montes et al. 2002; Contreras et al. 2006) and in general, in our hands, liposome spots containing sphingolipids were prone to aggregation, deformation or wash-out, rendering many spots unusable. However, when small zones of characteristic spherical liposome structure where obtained, these were analyzed, with the caveat that lipid partitioning or segregation cannot be excluded. Nevertheless, observations were possible with all liposome compositions except for mixtures containing 5 and 10% GlcCer, where liposomes where never obtained. Overall, out of 5 chips containing 20 arrays, 3 arrays with purified N-KSR (missing 2 spots) and 2 arrays (missing 6 spots) with purified C-KSR were successfully analyzed (Fig 6A-B, Supplementary Data S2). Considering statistical analysis for liposomes represented in at least 3 spots, low but significant normalized binding intensity (NBI) scores were observed between N-KSR and liposomes containing 2% GlcCer as compared to DOPC liposomes (p=0.0018). Although sample size was too low for a meaningful statistical analysis, morphological examination of images also suggested a potential weaker binding of N-KSR to C1P and of C-KSR probe to liposomes with 10% CER (Fig 6B).

Although we were not able to obtain purified C-KSR-GS, we next performed LiMa analysis using lysates from transfected HEK293 cells. Since the *in vitro* binding of the probes was weak, the use of lysates might also provide missing auxiliary components such as co-factors and proteins to improve binding affinity. In addition, Since purified N-KSR did not appear to bind liposomes containing bovine CER mixtures, the liposome composition of the arrays was modified to include a broader variety of ceramides as well as additional lipids including: the intermediate fatty acid chain-length ceramide C18-ceramide (d18:1/18:0, C18CER); the ceramide precursor dihydroceramide (d18:0/d18:0, C18DHCER) which harbors an identical headgroup as ceramide; sphingosine-1-phosphate (d18:1, S1P); DAG (d18:1/18:1); as well as DO-phosphatidic acid (DOPA), though the latter was found to auto-fluoresce, and so was excluded from the analysis. Liposomes containing 10-15% DAG producing large brightly fluorescent aggregates were also excluded. C-KSR lysates were not included since the fraction of intact probe as compared to released EGFP was small (see Fig 4C). However, in 6 chips with 3 arrays for each probe (missing 12 spots out of 78 for both N-KSR and C-KSR-GS), neither N-KSR nor C-KSR-GS showed signs of binding to any of these liposome compositions (Fig S5A-B, Supplementary Data S2). Since in intact cells C-KSR-GS binds the PM and might require a more complex lipid combination for successful binding, we then we sought to mimic the composition of the inner plasma membrane (IPM) leaflet (Lorent et al. 2020). To this end, IPM liposome backbones were prepared using: 30.2% palmitoyloleoyl-phosphatidylcholine (POPC); 15% PO-phosphatidylethanolamine (POPE); 33% cholesterol; 21.2% DO-phosphatidylserine (DOPS); and 2,5, or 10% of the bovine CER mixture (CerMix), C18DHCER, C18CER, DAG, chicken SM mixture (SM) and Sph. Next, the binding profile of the C-KSR1-GS probe was examined and compared to lysates of soluble EGFP as a negative control and PKC-C1(2) as positive control. In total out of 4 chips, 3 arrays for C-KSR-GS (missing 14 out of 78 spots), 3 arrays of EGFP (missing 7 out of 78 spots) and 3 arrays of PKC-C1(2) (missing 6 out of 78 spots) were successfully quantified. Using the automated quantification pipeline, whereas robust and significant binding of PKC-C1(2) to DAG-containing liposomes as compared to IPM liposomes was readily observed ($p < 0.0001$ for both 2% and 5% DAG, Fig S5C-D, Supplementary Data S2), there were no significant differences in NBI scores of C-KSR-GS liposome binding as compared to liposomes of IPM composition alone (Fig 6C, Supplementary Data S2). However, a closer examination of the images revealed striking morphological differences between C-KSR-GS and EGFP (Fig 6D, Fig S5E). Whereas EGFP was often found encapsulated into liposomes regardless of composition, with no signs of external binding, C-KSR-GS was only rarely found encapsulated, and instead the images suggested a potential weak binding, particularly to Cer containing liposomes, though faint binding was also discernible for DAG and IPM alone (Fig 6D, Fig S5E). Although it difficult to draw definitive conclusions from these *in vitro* screens, which do not fully recapitulate

conditions in cells, together the data hint at a difference in the binding profiles of the KSR1-based probes that could explain the difference in their response to Cer induction by Palm and Genz (Fig 2B vs 2D, Fig 5D vs 5E). Thus, C-KSR and C-KSR-GS may require a more complex membrane context or other co-factors in order to bind Cer species with higher affinity. On the other hand, the data suggest that N-KSR might instead be sensitive to GlcCer which are decreased upon treatment with both Palm and Genz at the same time that Cer is increased (Fig 2F), which could result in no net change if some affinity for Cer exists.

Behavior of KSR probes during phagocytosis

In several reports, Cer generation has been associated with phagocytic ingestion and phagosomal maturation, and shown to be important for pathogen elimination, yet its precise role is still poorly understood (Pathak et al. 2018; Tafesse et al. 2015). Thus, to not only glean some hints at Cer dynamics (and eventually function) during this important immune process, but also to test our probes under a physiological stimulus, we next investigated the behavior of our three KSR probes during phagocytosis. To this end, we examined MEFs rendered phagocytic by overexpression of the FCGR2A-c-myc, a phagocytic cell model that are easier to transfect and manipulate than naturally phagocytic cells, and which we and others have observed to recapitulate many aspects of the phagocytic process, including phospholipids dynamics (Giodini, Rahner, and Cresswell 2009; Bodman-Smith et al. 2004; Hinkovska-Galcheva et al. 2005; Santos et al. 2022). Cells expressing each of the 3 KSR probes (N-, C- and C-KSR-GS) were exposed to 3 µm IgG-opsonized polystyrene beads for 30 min or 135 min, time-points representative of early and late stages of phagosomal maturation (Fig 7A-C). In parallel, sphingolipids levels were measured in C-KSR-expressing cells using targeted lipidomics as described above, where a large accumulation in total cellular Cer, SM and HexCer (4-, 2.6- and 6-fold, p=0.0002, 0.0204 and 0.0349, respectively) was observed at 30 min (Fig 7D, Supplementary Data S1). This effect was driven by changes in C24-Cer -SM and -HexCer species as well as C16-Cer and -HexCer (Fig 7Di-iii, Supplementary Data S1). It should be noted that large changes in total PC, PI and PS phospholipids were also observed (Fig S6A), which grossly biased the usual normalization procedure. Thus, rather than normalizing each individual sample by its own total glycerophospholipids, individual samples were first normalized by their corresponding total protein, and the entire dataset was normalized by the average daily total glycerophospholipid levels to obtain mol% values.

Interestingly, 30 min after the incubation with the IgG-coupled beads, accumulation of C-KSR could be observed at the PM, similar to what was observed after incubation with bacterial SMase (Fig 7A, see also Fig 3A). Exploring this PM localization further, we found PM staining

of C-KSR could still be detected at 45 min and 60 min after phagocytosis initiation, yet was

lost at later stages, after 90 and 135 min (Fig S6B). Similarly, PM localization of N-KSR and

C-KSR-GS was salient at 30 min (Fig 7B-C), even though both constructs were already found

at the PM under basal conditions, whereas at 135 min the N-KSR PM signal was not as clearly

discerned (Fig 7B). Moreover, whereas N-KSR was completely absent from phagosomes (Fig

7B, insets), in C-KSR-GS-expressing cells, accumulation around phagosomes was discerned

in a subset of phagosomes (Fig 7C, arrows). In some instances, a pattern of small flattened

cisternae or protruding tubular structures could be seen around phagosomes, which were

reminiscent of membrane contact sites (MCS) between the phagosomal membrane and ER,

structures that have been studied extensively by our laboratory in the past (Guido, Demaurex,

and Nunes 2015; Nunes et al. 2012; Nunes-Hasler et al. 2017; Santos et al. 2022) (Fig 7C,

arrows). To confirm whether such structures could indeed represent ER, C-KSR-GS was then

co-expressed with several ER-targeted constructs, including: a soluble ER-targeted RFP

TagRFP-KDEL, a calcium regulator and MCS marker mCh-STIM1 and mCh-Sec22b, an ER-

ERGIC protein that we have recently shown to be present in STIM1-independent MCS (Nunes

et al. 2012; Santos et al. 2022) (Figs 7E-F, Fig S6E). Interestingly, though overlap between

C-KSR-GS and KDEL was not easily discerned, likely due to low overall KDEL expression

(Fig S6E), overlap with STIM1 and Sec22b was observed in some but not all instances of

periphagosomal accumulation of the fluorescent proteins (Fig 7E-F). Finally, we expressed

the N-KSR and C-KSR-GS probes in a naturally phagocytic dendritic cell JAWS II. Similar to

MEFs, N-KSR did not localize to the phagosomal or other organellar membranes and

appeared to be depleted from the PM at late phagocytic time points (Fig 8A). In contrast,

robust C-KSR-GS accumulation at the phagosomal membrane could be discerned on a subset

of phagosomes, as well as on internal structures reminiscent of the ER (Fig 8B). Together

these intriguing observations suggest the following: first, that the large changes in Cer during

the first 30 min of phagocytosis detected in whole-cell MEF extracts are likely occurring at the

PM and are best detected by C-KSR which is the most sensitive of the 3 probes at detecting

changes of PM Cer. Second, that C-KSR-GS may more be useful to detect Cer species on

intracellular membranes including phagosomal and ER membranes. Finally the data also hint

at dynamic Cer fluxes, potentially between phagosomal membrane and ER at MCS.

Discussion

Despite a large number of proteins reported as Cer binders (Canals, Salamone, and

Hannun 2018; Bockelmann et al. 2018; K. Zhou et al. 2018), to our knowledge no genetically

encoded Cer-specific probe per se have not been reported, though whole GFP-tagged Cer-

binding proteins have been employed to track subcellular Cer in a limited number of studies (Canals, Salamone, and Hannun 2018). Out of 14 candidate biosensor constructs reported here, we focused on 3 based on the KSR1(317-400) Cer-binding domain which displayed morphology of membrane localization. In contrast, constructs based on PRKCZ and SET were either targeted to the nucleus, or, for those bearing a nuclear export signal (NES), did not respond to SMase treatment (Fig S3). Since sphingomyelinases and ceramidases have been reported in the nucleus, and indeed SET has a defined nuclear activity (Adachi, Pavlakis, and Copeland 1994), it is possible that these proteins may detect nuclear Cer rather simply being misfolded (Albi and Viola Magni 1997; Shiraishi, Imai, and Uda 2003), although the diffuse rather than nuclear membrane localization is unexpected. Future studies employing stimuli known to induce nuclear Cer may still help to verify the utility of these constructs.

In contrast, though nuclear shuttling has also been reported for KSR1, this scaffolding regulator MAPK signaling normally resides in the cytosol in resting conditions (Brennan et al. 2002). It is recruited to the PM upon growth factor stimulation, which induces both changes in the phosphorylation status of multiple KSR1 sites, as well as inducing the activities of neutral and acidic SMases, generating Cer (Mathias, Peña, and Kolesnick 1998; Razidlo et al. 2004). Thus KSR1-based probes are particularly poised to detect Cer in the context of PM lipids, and the baseline localization of N-KSR, 2x-C-KSR as well as C-KSR-GS, as well as recruitment to PM upon SMase treatment of C-KSR and C-KSR-GS are all consistent with this notion. In addition to Cer binding, mutagenesis studies suggest that phosphorylation of T274 and S392 contribute to KSR1 stability, and in particular dephosphorylation of S392, the latter part of the CA3 domain, participates in membrane re-localization upon growth factor activation (Brennan et al. 2002; Razidlo et al. 2004). Whether S392 in the CA3 of our KSR constructs is constitutively dephosphorylated, or whether separation from other KSR1 domains is sufficient to allow conformational changes favoring Cer binding remains to be defined, though full-length KSR1-S392A localizes to the nucleus rather than PM (Brennan et al. 2002). Several studies have reported direct Cer binding, including *in vitro* studies using an N-terminal GST-fusion with a CA3 of similar composition (aa 320-388) as in our own study , though others have failed to detect direct binding (Pérez-rivas, Prieto, and Lozano 2010; Yin et al. 2009; Zhang et al. 1997). On the other hand, the structure of KSR1 has been resolved, with a shallow Cer binding pocket identified based on similarity to other C1 domains to involve particularly residues L342, V345, M353, I354 and F355 (M. Zhou et al. 2002). Thus, the KSR1 CA3 would not be expected to, on its own, interact with the fatty acid chain or to be able to extract lipids from membranes, but rather to interact with lipid head-groups. However, the same group challenged the Cer-binding ability of the CA3 domain though it should be noted that, in this case, a truncated CA3 (aa 334 to 377) was employed.

In the characterization of our probes, a surprising observation was that the most salient

change in the behavior of the KSR probes in response to Cer-modifying drugs was a change

in total cellular fluorescence. This is in contrast to similar phospholipid probes, which change

locations rather than total fluorescence. The effect was consistently observed in both induction

and depletion directions, across several drug treatments, including Myr, FumB1, Palm,

SMase, C6 and Genz (Figs 2, 3, 5). For Myr treatment, we confirmed it was not simply due to

toxicity (which often accompanies changing levels of Cer (Hannun and Obeid 2008)) as

cytosolic RFP levels were unchanged. Moreover, we tested different concentrations in pilot

studies and chose concentrations and timepoints where visible signs of toxicity were low or

not yet apparent. With SMase treatment, neither the DAG probes nor the PRKCZ or SET

probes reacted in such a manner (Fig 3, S3). As mentioned briefly above, a potential

explanation is that the CA3 domain might itself be somewhat unstable or prone to misfolding

and subject to certain level of degradation, and that membrane binding thus stabilizes the

protein, leading to a decrease in degradation and concomitant increase in total probe

fluorescence. Indeed, full-length KSR1 binds cytosolic chaperones Hsp70, 90 and 68,

(Stewart et al. 1999), and ubiquitination of KSR1 has been reported, though the site identified

was outside the CA3 domain (Matheny et al. 2004; Rinaldi et al. 2016). Moreover,

phosphorylation at both T274 and S392 were required to increase the half-life of KSR1 from

1hr to 6hr (Rinaldi et al. 2016). Such a fast turnover time is consistent with our results which

were observed even with treatments as short as 30 min. The PM re-localization accompanying

SMase treatment observed for both C-KSR and C-KSR-GS, is consistent with such a

hypothesis, but for Palm treatment for C-KSR and Palm, C6 and Genz treatments for C-KSR-

GS, PM recruitment was less obvious and instead a truly global fluorescence increase was

observed. Whether this reflects the simultaneous binding of several types of organellar

membranes that were not resolved at the confocal level (for example ER, mitochondria, Golgi

and endo-lysosomal all at the same time), or whether post-translational modifications such as

phosphorylation or ubiquitination, or chaperone binding might also be modified, increasing

stability also of the cytosolic fraction of the probe, will be important to define in future studies.

In addition to its potential influence on cellular probe behavior, an intrinsic instability,

or requirement for post-translational modification of the CA3 domain to efficiently bind Cer

could well explain the difficulties we encountered in detecting *in vitro* binding. In these studies,

the only significant binding scores was the surprising interaction of the N-KSR probe to

GlcCer-containing liposomes. Though unexpected, as interaction of full-length KSR1 to

GlcCer has to our knowledge not been reported, such a modification to the lipid specificity

interaction is consistent with the difference between N- and C- constructs to Palm, Genz as

well as phagocytic stimuli. Future studies with drugs or genetic manipulation of enzymes such

as glucosyl and galactosyl ceramidases or transporters such as TMEM16F (Suzuki et al. 2013)
would be useful to corroborate these finding and determine more precisely the specificity of
this probe for glycosylated Cer, or potentially phosphorylated Cer in the cellular context.
Though we could not unambiguously verify specific direct binding of our probes to Cer *in vitro,*
which was in part due to liposome instability, an effect of instability of the purified or solubilized
probe itself, or of a disruptive interaction between probe and liposome membranes cannot be
discounted. Indeed, liposome morphology in arrays exposed to lysates bearing PKC-C1(2) or
EGFP was much more consistent and less spots were lost, even with sphingolipid containing
liposomes. Nevertheless, in the few zones that did show typical liposome morphology,
qualitative evidence of Cer binding could still be gleaned. However, for future biochemical
analyses it would be worth to explore whether, for example, the creation of S392A mutants of
the C-KSR-GS construct, which lacks MEK-interaction domains that leads to nuclear
localization (Brennan et al. 2002), may improve probe stability in solution, allowing *in vitro* Cer-
binding to be more precisely measured. Despite these drawbacks, the examination of our 3
KSR1-based constructs under different stimuli demonstrated that these probes may each
have their own utility as such or serve as a basis for future studies dissecting Cer, and
potentially GlcCer, dynamics in cells. The exacerbated instability of C-KSR, which prevented
constitutive PM localization, proved in fact to be advantageous in the way that it allowed for a
greater dynamic range in response to Cer changes by Palm and SMase, and more salient PM
recruitment with a phagocytic stimulus than C-KSR-GS. Thus, like conditional instability
employed in nanobody engineering (Tang et al. 2016), a future design could build upon these
observations to include proteasome targeting signals which maybe less disruptive than the
current design to membrane homeostasis (L. Zhao et al. 2022) to obtain a sensitive on-off
biosensor for PM Cer.

Finally, our intriguing observations of the differential behavior of N-KSR and C-KSR-
GS during phagocytosis suggests that the two probes are indeed responding to different lipid
changes in the cell, though verification using chemical or genetic manipulation of key enzymes
and transporters that modify Cer and GlcCer will be instrumental in more firmly establishing
this. One important notion to consider moving forward is that such cytosolic probes will only
detect lipids that are present on the cytosolic leaflet, or only when luminal leaflets are exposed
to the cytosol for example as may occur during PM or lysosomal disruption, where SM probe
lysenin has recently contributed greatly to our understanding of the role of SM (and indirectly
Cer) in membrane repair and antigen cytosol transfer in dendritic cells (Niekamp et al. 2022;
Canton et al. 2021). Though several studies indicate that Cer readily flip-flops across lipid
bilayers (Goñi et al. 2005), this is less likely for complex sphingolipids such as GlcCer, and
trans-bilayer transporters might maintain or establish gradients in response to cellular signals

as occurs for PS (Williamson 2015). As such, changes such as the loss of N-KSR PM signal
might equally represent a loss of total GlcCer as it might simply reflect its removal from the
cytosolic leaflet, which could explain why global changes in HexCer was not detected by
lipidomic analysis, though compensating changes in GalCer could also provide an
explanation. However, that the exciting ER localization of C-KSR-GS upon certain conditions
is simply a result of misfolding is unlikely, and investigation of whether this truly represents
Cer flux is ongoing in our laboratory. Although clearly more work will be required to define the
behavior and lipid specificity of these novel biosensors, we believe these constructs represent
useful new tools to explore sphingolipid regulation and function in living cells.

Materials and Methods

Reagents

The following antibodies (antibody Name/catalog #/dilution) were from: Abcam (USA) – anti-
Rab7/ ab137029/1:100; anti-Rab5/ab18211/1:100; Enzo Life Sciences (USA) – anti-LAMP1/
ADI-VAM-EN001/1:50, BD Biosciences (USA) – anti-TGN38/ 610898 /1:200; Thermo - AF555
Goat anti-mouse IgG/ A-21422/1:500, AF555-goat-anti-rabbit IgG/A-21428/1:250; Santa-Cruz
Biotechnology - Goat anti-rabbit IgG-HRP/1:5000, BioRad (USA) – Goat anti-mouse IgG-
HRP/170-6516/1:3000. The following lipids were purchased from: Avanti (USA, Sigma/Merck)
(Name/catalog #): DOPC/ 850375P, POPC/ 850457P, DOPS/ 840035P, POPE/850757P, PE-
PEG350/880430, C18 Ceramide/ 860518P, C18 Dihydroceramide/860627P,
Sphingosine/860490P, Diacylglycerol 18:1/800811O; Sphingomyelin from egg yolk/85615,
ceramide mix from bovine spinal cord/22244, Cholesterol/C8667. PE-Atto647/ AD 647N-161
was from Atto-Tec (Germany). All lipid standards were from Avanti (Sigma/Merck) amounts
given per 6-cm cell culture dish: DLPC (12:0-12:0 PC, 0.4 nmol/dish, cat. #850335), PE31:1
(17:0-14:1 PE, 1 nmol/dish, cat # LM-1104), PI31:1 (17:0-14:1 PI, 1 nmol/dish, cat # LM-1504),
PS31:1 (17:0-14:1 PS, 3.3 nmol/dish, cat# LM-1304), CL56:0 (14:0 Cardiolipin, 0.7 nmol/dish,
cat#710332), C12SM (C12 Sphingomyelin d18:1-12:0, 0.1 nmol/dish, cat#860583), C17Cer
(C17 Ceramide, d18:1-17:0, 2.5 nmol/dish, cat #860517), C8GC (C8 Glucosyl(B)Ceramide,
d18:1-8:0, 0.5 nmol/dish, cat#860540). Chemicals were purchased from Sigma/Merck unless
otherwise stated, (Name/catalog #): HPLC grade Methanol/4860-1L-R, HPLC grade
Chloroform/650471, tert-Butyl Methyl Ether (MTBE)/650560, Methylamine/534102; LC-MS
grade water/1153331000. Thermo: HPLC grade n-butanol/11378197. Plasmids used and
generated in this study are listed in Supplementary Table 1.

DNA cloning

549 All newly made constructs were verified by sequencing. Constructs expressing candidate

550 ceramide probes were constructed by subcloning fragments encoding human KSR1 (aa 317-

551 400, XM_047436985), N-terminal fragment of human PRKCZ(aa 123-193 , NM_002744.6),

552 20 kD C-terminal fragment human PRKCZ (aa 405-646), or human SET(aa 70-226,

553 NM_001122821.2) into pEGFP-N1 using restriction sites NheI and XhoI or into pEGFP-C1

554 using XhoI and KpnI sites to create C- or N-terminally tagged proteins, respectively. Tandem

555 2x-C-KSR probe was made by subcloning human KSR1(317-400) fragment encoding a

556 flexible linker sequence (GGSSGGGGA) into pKSR1-EGFP using XhoI and KpnI sites. The

557 sequence encoding Nuclear Export Signal (NES) from mammalian MAPKK kinase –

558 LQKKLEELEL was inserted by site directed mutagenesis (Q5 Site-Directed Mutagenesis Kit,

559 NEB, USA) after the start codons of human PRKCZ(405-646) or SET(70-226) to yield

560 plasmids pC-NES-C20-PRKCZ-EGFP and pC-NES-SET-EMD-EGFP, respectively. A point

561 mutation in the CA3 domain of KSR1 (C359S, C362S) with substitution of the two conserved

562 cysteines with serine residues was introduced using the Q5 Site-Directed Mutagenesis Kit.

563 For protein purification from *E.coli,* the human KSR1(aa 317-400) fragment was cloned into

564 SfiI restriction sites of the pETM11-SUMO3-sfGFP (Saliba et al. 2014) to make pETM11-

565 SUMO3-sfGFP-KSR1. The sequence encoding KSR1(317-400)-GFP was subcloned into

566 pETM11-SUMO3 via Gibson assembly using Gibson Assembly Cloning Kit (NEB, USA).

567 Sequences encoding N- or C-terminally tagged KSR1(317-400) were subcloned into SfiI sites

568 of pSBbi-pur-H-2Kb (# 111623, Addgene, USA) replacing H-2kb fragment. Primers used are

569 listed in Supplementary Table 2.

570 *Cell culture, transfection, and stable cell line establishment*

571 Wild-type mouse embryonic fibroblasts (MEF, ATCC CRL-2991), HeLa cells (ECACC,

572 93021013, a gift from N. Demaurex), and HEK 293T (HEK, ATCC-CRL-11268, a gift from N.

573 Demaurex) were cultured in DMEM (22320, Thermo), MEM (41090, Thermo) and DMEM

574 glutaMAX (31966, Thermo) respectively, containing 10% fetal bovine serum (FBS, Thermo),

575 1% pen/strep (10000 U/ml penicillin, 10 mg/ml streptomycin, PAN Biotech, Germany). JAWS

576 II (JAWS, ATCC CRL-11904) were grown in alpha minimum essential medium with

577 ribonucleosides,Deoxyribonucleosides (Gibco 22571), supplemented with 4 mM L-glutamine

578 (Gibco 25030), 1 mM sodium pyruvate (Gibco 58636), 5 ng/ml GM-CSF (Peprotech, 315-03)

579 20% FBS, 1% pen/strep. Delipidated FBS was prepared by incubating FBS with 0.5 g/ml lipid

580 removal beads (13358-U, Sigma/Merck) at 4°C overnight with rotation followed by

581 centrifugation (27000g, 20 min, 4°C) and supernatant filtration using 0.2 μm PES filters

582 (Nalgene). All cell lines were cultured at 37°C, 5% CO_2, and passaged 1-2 times per week,

583 and used between passages 10 and 40. All cell lines were tested every 6-12 months either by

584 PCR (LookOut kit, Sigma/Merck) or Mycostrips (Invivogen, ep-mysnc-50) and were negative.

Cells were transfected upon reaching 60-70% confluency 2-3 days after seeding using
Lipofectamine 2000 (Thermo) in full culture medium without antibiotics for 4-6h. To prepare
DNA-lipofectamine complexes, equal volumes of serum- and antibiotic-free media containing
DNA or Lipofectamine were mixed and incubated at room temperature for 1h. To obtain stable
cell lines via transposition, MEF cells were first co-transfected with pSBbi-pur plasmids
encoding the protein of interest and pCMV-(CAT)T7-SB100 plasmid (#34879, Addgene, USA)
encoding Sleeping Beauty transposase SB100x. Puromycin (10 µg/mL) selection was
performed 2 days after co-transfection. Stable cell lines (N-KSR1 and C-KSR1) were
maintained in full culture medium containing puromycin (10 µg/ml). JAWS transfection was
achieved using 20 µg DNA and 20 x 10^6 million cells per 100 µl nucleofection solution from
the Amaxa Nucleofector Mouse Dendritic Cell kit (Lonza), and the Y-001 program. Cells were
diluted in 400 µl RPMI 1640 Glutamax medium, 25 mM HEPES medium (Thermo, 72400021)
immediately after electroporation, incubated for 10 min at RT, then centrifuged at 200 g, 5 min,
RT and resuspended in warm full medium and incubated overnight before imaging.

Immunofluorescence

MEF and HeLa cells seeded on 0.17 mm glass coverslips (Carl Roth) were fixed with 4%
paraformaldehyde (PFA, AlfaAesar) in PBS for 30 min at RT followed by 3X washes in PBS.
JAWS were fixed in 1%PFA for 10 min. For staining nuclei, fixed cells were incubated with 10
µg/ml Hoechst 33342 (Thermo) for 5 min at followed by three washes with PBS and mounting.
For immunostainings, cells were permeabilized with 0.1% Triton X-100 (in PBS) for 5 min at
RT, washed 3X by PBS, quenched with 25 mM Glycine/PBS for 10 min at RT, treated with
Image-IT-FX (Thermo) for 30 min followed by blocking with 2% bovine serum albumin
(BSA)/PBS for 30 min. After that the cells were incubated with primary antibodies at 4°C
overnight and with a secondary antibody for 1h in dark humid chambers, antibodies diluted in
blocking buffer. 3x washes in PBS were carried out between each step in the procedure.
Finally, the coverslips were mounted onto glass slides using 5 µl of ProLong Diamond
Mounting medium (Thermo) per coverslip. For labeling recycling endosomes, MEF cells were
incubated for 30 min at 37°C with 50 µg/ml Alexa555-conjugated human transferrin (Thermo)
in MEF full culture medium containing 0.2% (BSA). After 2x washed with PBS, cells were fixed
with 3% PFA for 20 min, quenched with 50 mM ammonium chloride for 15 min (RT), followed
by 3 washes in PBS and mounting. For staining the lysosomal lumen, MEF cells stably
expressing C-KSR were incubated overnight with 1 µg/ml Alexa Fluor 594 Hydrazide (Thermo)
in full culture medium with puromycin. The following day, medium was changed 5x to full
culture medium without Alexa 594 and the dye was chased for 2.5h before the start of
phagocytosis. Images were acquired either using Zeiss LSM700 or LSM800 confocal systems
/Plan apochromat 63x/1.4 objectives and Zeiss Zen 2010b version service pack1, or Nipkow

Okagawa Nikon spinning disk confocal imaging system Plan Apo 63x/1.4 DICIII objective and

Visiview 4.0 software (Visitron Systems).

Drug treatments

To induce sphingolipid depletion MEF cells were treated with: 0.5 µM myriocin (or methanol

vehicle) from *Mycelia sterilia* in full culture medium, with 2.5 µM fumonisin B1 (methanol

vehicle) from *Fusarium moniliforme* in culture medium containing 10% FBS delipidated as

above for 3 days, or with 10 µM fumonisin B1 in regular cell culture medium overnight. To

induce sphingolipid production MEF cells were treated with 0.5 mM BSA-conjugated palmitic

acid complex (Cayman Chemical, USA) for 4h at 37°C in delipidated-FBS medium, and in

regular medium with 100 µM Genz-123346 (DMSO vehicle) overnight, or 200 µM C6-ceramide

(d18:1/6:0) (Avanti/Merck) for 2h (ethanol vehicle). To induce DAG accumulation cells were

treated with 10 µM ionomycin in regular medium for 10 min. To release ceramides locally at

the plasma membrane cells were treated with 0.5 U/ml sphingomyelinase from

Staphylococcus aureus for 30 min in DMEM medium without FBS and antibiotics. After

treatment cells were either placed on ice, scraped and used for lipid extraction or fixed with

4% PFA and mounted for microscopy. Alternatively, after SMase treatment, cells were stained

with a sphingomyelin probe EGFP-Lysenin (Kiyokawa et al. 2005) (purified from *E.coli,* see

below) at 20 µg/ml in serum-free medium for 30 min at 37°C followed by 2 washed in PBS and

fixation as described above.

Image analysis

Image analysis was performed using ImageJ v1.52 (Schneider, Rasband, and Eliceiri 2012)

NIH. Colocalization analysis was conducted using the JACoP v2.0 plugin (Bolte and

Cordelières 2006) that returned Mander's coefficients for each region of interest (ROI). For

quantification of KSR probe fluorescence levels at the PM, ROIs for cells and background

were drawn manually or using Wand tool when possible on single confocal planes. The Make

Band... command, set to 2 µm width was used to transform the whole-cell ROIs into a PM

ROIs. Mean fluorescence intensities were measured from background-subtracted images.

Analyses determining the percentage of cells displaying plasma membrane localization were

conducted on images randomized by the Filename_Randomizer.txt macro

https://imagej.nih.gov/ij/macros/Filename_Randomizer.txt, by a blinded experimenter that

counted the number of PM-positive cells and total cells in each image using the Cell Counter

plugin. For whole-cell fluorescence intensity analyses, confocal z-stacks of 9 frames spaced

at 0.5 µm were acquired and sum projections of stacks were created using the Z-Projection

tool. Cell and background ROIs were drawn as above on the sum projections images to obtain

mean total cellular fluorescence intensities.

Immunoblotting

Cells grown to 80-90% confluency on 6-cm dishes were washed 3x with PBS, scraped off, pelleted, and lysed in RIPA buffer (Thermo 89900) with protease inhibitors (Halt Protease Inhibitor cocktail, 87785) for 30 min at 4°C. Cell debris was pelleted at max speed for 10 min (4°C), and the supernatant was used for loading into the gel. Protein concentration was quantitated using RotiQuant kit (Carl Roth). Lysates were diluted in the SDS Loading Buffer (Roti-Load, Carl Roth) and heated at 95°C for 5 min. 15-30 µg protein were loaded and separated by SDS-PAGE in 4–12% SurePAGE 12-well pre-cast gels (Genscript). Proteins were transferred onto PVDF membranes using iBlot or iBlot2 system (Thermo). After transfer, membranes were blocked in 5%milk/TBS-1%Tween-20 (TBST) for 1h (RT, agitation) followed by incubation with primary antibodies diluted in 3%-milk/TBST overnight at 4°C and 1h incubation at RT with HRP-conjugated secondary antibodies 3%-milk/TBST with multiple washes in PBS between incubations with primary and secondary antibodies. Signals were detected using Immobilon HRP substrate (Merck) on the ImageQuant LAS 4000 mini imaging system (GE, USA).

Lipid extraction and lipidomics

For lipidomics experiments, lipids were extracted following a modified MTBE protocol (Guri et al. 2017). MEF cells grown to confluency in lipid-free or full culture medium on 6-cm dishes were washed in cold PBS and scraped off in 1 ml PBS. 900 µl and 100 µl cell suspension were transferred to two separate tubes and the cells were pelleted at 400g for 5 min (4°C). The pellets were kept at -80°C or used for protein quantitation (100 µl sample) by resuspension in RIPA buffer, using RotiQuant as above, or for lipid extraction (900 µl sample) as follows. First, pellets were resuspended in 100 µl LC-MS grade water, then 360 µl methanol and the lipid standard mix (see Reagents section) were added, and the tubes vigorously vortexed. Next, 1.2 ml MTBE was added, and the samples were incubated for 1h at RT with shaking. 200 µl of water was added to induce phase separation; samples were incubated at RT for 10 min followed by centrifugation at 1000g, 10 min (RT). The upper organic phase was transferred to 13mm glass tube, 400 µl of the artificial upper phase (MTBE/methanol/water (10:3:1.5, v/v)) was added, the phase separation was repeated, and the collected second upper phase was combined with the first one. The total lipid extract was divided into 2 equal parts: one for sphingolipids (SL) and one for glycerophospholipids (TL). Extracts were dried in a speed vacuum concentrator. After drying, the samples were flushed with nitrogen unless used for further steps. The sphingolipid aliquot was treated with methylamine to remove phospholipids by de-acylation (Clarke method (Clarke and Dawson 1981)). 0.5 mL monomethylamine reagent (MeOH/H2O/n-butanol/Methylamine solution (4:3:1:5 v/v) was added to the dried

lipids, tubes sonicated for 6 min, vortexed, followed by the incubation for 1h at 53°C. Finally,

the methylamine-treated fraction was dried (as mentioned above). The monomethylamine-

treated lipids were subjected to desalting via extraction with *n*-butanol. In short, 300 µl water-

saturated *n*-butanol (*n*-butanol/water 2:1 v/v, mixed and prepared at least 1 day before use)

was added to the dried lipids. The samples were vortexed and sonicated for 6 min, and 150

µl LC-MS grade water was added. The mixture was vortexed thoroughly and centrifuged at

3200 x g for 10 min. The upper phase was transferred to a 2 mL amber vial. The lower phase

extraction was repeated as described above twice more, and the organic phases were

combined and dried.

Targeted lipidomic analysis by mass spectrometry was performed according to prior protocols

(Jiménez-Rojo et al. 2020) as follows: TL and SL aliquots were resuspended either in 250 µl

Chloroform: Methanol (LC-MS grade, 1:1 v/v) or directly in the sample solvent

((Chloroform/Methanol/Water (2:7:1 v/v) + 5mM Ammonium Acetate). After sonication for 6

min (RT), lipid mixtures were transferred into wells of a 96-well Twin tech PCR plate

(Eppendorf) using Hamilton syringes. Sphingolipid aliquots were diluted 1:4, and total lipids

were diluted 1:10 in the sample solvent. PCR plate was sealed with EasyPierce foil (Thermo),

and the samples were injected into the TSQ Vantage Triple Stage Quadrupole mass

spectrometer (Thermo Fisher Scientific) equipped with a robotic sample handler and nanoflow

ion source Nanomate HD (Advion Biosciences, USA). The collision energy was optimized for

each lipid class. Lipid detection parameters are listed in Supplementary Table 3. Samples

were injected at a gas pressure of 30 psi with a spray voltage of 1.2 kV and run on the mass

spectrometer operated with a spray voltage of 3.5 kV for positive mode and 3.0 kV for negative

mode and a capillary temperature set at 190oC. Sphingolipid and glycerophospholipid species

were identified and quantified using multiple reaction monitoring (MRM). Cer species were

also quantified with a loss of water in the first quadrupole. Each biological replicate was

measured twice (2 technical replicates, TR). Each TR consisted of 3 measurements for each

transition. Each lipid species was quantified using standard curves created from internal

standards. Data analysis was performed using R version 3.6.2 (R Foundation for Statistical

Computing). The values where then normalized to the total lipid content of each corresponding

lipid extract (mol% of the sum of total PC, PE, PI, PS), except for samples of phagocytosing

cells, were lipid concentrations were normalized first to total protein of each corresponding

individual sample, then to the average of the sum of total PC, PE, PI, PS for the entire sample

group as explained in the results section.

Protein purification from E.coli

Rosetta 2(DE3) *E.coli* cells (Novagen) were transformed with the plasmids expressing His6-

SUMO3 tagged KSR probes or GFP alone (control): pETM11-His6-SUMO3- -sfGFP-N-KSR,

pETM11-His6-SUMO3-C-KSR-sfGFP, pETM11-His6-SUMO3-sfGFP. Transformed bacteria

were grown at 37°C in 1L LB medium containing 100 µg/ml kanamycin, 34 µg/ml

chloramphenicol until OD600 0.6-0.8. Protein expression was induced by the overnight

incubation with 0.4 mM Isopropyl β-D-thiogalactoside (IPTG) at 16°C. The next day cells

bacteria were lysed using the Microfluidizer LM20 high-pressure homogenizer (15000 psi

pressure, 2 cycles) in lysis buffer (50 mM Tris pH 7.5, 500 mM NaCl, 20 mM imidazole,

protease inhibitors (1 tablet/50 mL), 0.5 mM DTT) and the soluble fraction was obtained by

centrifugation at 16000g for 1h (4°C). Proteins harboring the His6 tag were captured by

incubation with Ni-NTA resin (Qiagen) (2ml resin per bacterial lysate from 1L culture) for 2h,

4°C, 30 rpm rotation. Agarose resin with bound proteins was centrifuged at 1200 rpm, 8 min

(4°C), and the slurry was transferred to a column with a cotton filter. The column was washed

with 20 ml Lysis buffer, His6-tagged proteins were eluted with 6 ml Elution Buffer (50 mM Tris

pH 7.5, 250 mM NaCl, 500 mM imidazole), DTT was added to 5 mM to prevent protein

aggregation. Eluted proteins were dialyzed against 50 mM Tris pH 7.5, 200 mM NaCl, 1 mM

DTT overnight at 4°C. The His6-SUMO3 tag was cleaved off the proteins by incubation with

His-tagged SenP2 protease (EMBL Protein Expression and Purification Core Facility

European, Germany) at a molar ratio of SenP2 to protein 40:1. The undigested proteins and

the protease were removed using Ni-NTA resin, purified proteins were eluted by applying

Elution Buffer 2 (50 mM Tris pH 7.5, 250 mM NaCl, 10 mM imidazole) to Ni-NTA-packed

column. The quality of purification was assessed by SDS-PAGE followed by Coomassie

staining (RotiBlue kit, CarlRoth). sfGFP-N-KSR was further purified by size exclusion

chromatography on the S75 column pre-equilibrated with 50 mM mM Tris, pH 7.4 and 250 mM

NaCl.

Cell lysate preparation from HEK cells

HEK cells grown on 10-cm dishes in HEK full culture medium to 70-80% confluency were

transfected as above with plasmids encoding EGFP (control), KSR lipid probes, or PKC-

C1(2)-GFP, where 20 µg plasmid DNA/50 µl lipofectamine 2000 was used per dish. After

overnight incubation, the cells were trypsinized for 1 min (RT) and resuspended in PBS

containing 0.5 mM EDTA. Cells suspension was centrifuged 200g, 5 min (4°C), followed by a

wash in PBS. After pelleting, cells were resuspended in 120 µl hypotonic lysis buffer (10 mM

Tris-HCl, pH=7.5, Halt protease and phosphatase inhibitors) and incubated on ice for 20 min.

Cells were lysed by passing the suspension 15 times through a 30G syringe needle. Protein

concentration was measured using a NanoDrop (Thermo). Finally, NaCl and DTT were added

to 150 mM and 5 mM, respectively.

Liposome Microarray-based assay (LiMA)

Liposome-based protein-lipid interaction assay was performed following the protocol described in (Saliba et al. 2014). Lipid mixtures of different compositions were prepared in chloroform: methanol: water (20:9:1 v/v). Lipid mixtures containing fluorescent lipid PE-Atto647 (0.5 mol %) were spotted onto thin agarose layers covered by the protective membrane on top of a glass slide using automated lipid spotting under an inert atmosphere (Automatic TLC-spotter4, Camag, Switzerland). A microfluidic chamber with four channels was manufactured from poly(dimethylsiloxane) PDMS, as described in (Saliba et al. 2014). Then the agarose layers with spotted lipids were glued onto PDMS chambers using a 2:1 mixture of silicone elastomer: curing agent diluted 3× (v:v) in hexane as an adhesive, so the resulting microfluidic chamber contained four channels of 30 spots, 120 lipid spots per chamber. Liposomes were generated by manually injecting the liposome buffer (10 mM HEPES, 150 mM NaCl) into each channel with a Hamilton syringe. Then, 40 µl of the solution containing purified proteins at 5-10 µM (or HEK cell lysates) in liposome buffer with 5 mM DTT were injected into channels using an automated injection system with a flow rate of 8 µl/min. After incubation of freshly formed liposomes with the GFP-tagged proteins for 1h at RT, the channels were manually washed with 80 µl of liposome buffer, and all spots in the chamber were imaged with the Zeiss Axio Observer Z1 video time-lapse system. All images were taken with the 20x objective; liposomes were imaged with the Cy5 filter set (5 and 10 ms exposure), and GFP-tagged proteins were imaged at different exposure times (1, 5, 10, 25, 50, 75, 100, 150, 200, and 300 ms). Images were analyzed using a CellProfiler version 4.2.5 (Broad Institute) custom pipeline described in detail in (Saliba et al. 2014). A Custom-made R script was used to calculate the normalized binding intensity value (NBI) for each protein-lipid interaction. NBI represents a mean of the ratio of GFP to Atto647 fluorescence divided by exposure time at all exposure times for which spots were successfully segmented and is proportional to the amount of GFP-tagged interaction protein recruited to the liposome surface.

Preparation of phagocytic beads and phagocytosis

3 µm carboxyl polystyrene beads (Spherotech, CP30-10) were covalently coupled with purified human IgG (hIgG, Dunn Labortechnik, Germany) by washing 3 times in sterile PBS with centrifugation at maximum speed (18,000 g) at 4°C, activating with 50 mM 1-Ethyl-3-(3-dimethylaminopropyl) carbodiimide hydrochloride (EDC-HCl, Carl Roth/2156.1) for 15 min in PBS at room temperature (RT) with rigorous shaking, followed by 3 washes at 4°C in 0.1 M $Na_2B_4O_7$ buffer (pH 8.0). Finally, 2 µg/µl of hIgG (and 15 ng/µl iFluor647-amine (AAT Bioquest), for fluorescent beads) were added to the beads and incubated overnight at 4°C on

a shaker. Next day, the beads were washed 2x with 250 mM glycine/PBS followed by two

washes in PBS. To render MEF cells phagocytic MEF cells were transfected with 1 µg/2x10^5

cells pcDNA3-myc-FCGR2A (Fc receptor) plus KSR probes and where indicated TagRFP-

KDEL, mCherry-STIM1 or mCherry-Sec22b at 0.3 µg per 2x10^5 cells. Cells were washed in

full medium 4-6h after transfection. The following day beads were added to the cells at a bead-

to-cell ratio 10:1, cells were centrifuged at 200g, 2 min, RT, incubated with the beads for the

indicated times at 37°C, washed with PBS, fixed with 4% PFA/PBS for 30 min followed by 3x

washes with PBS and mounting as described above.

Statistics and reproducibility

All statistical analyses were done using Prism (GraphPad Software). P values that are

significant are shown above the bars. Two-tailed unpaired t-test were performed when

comparing two samples, Welch's correction was used if F test showed a significant difference

in variance. For three or more sample comparisons a one-way ANOVA was employed with in

case of negative F test, and Brown-Forsythe and Welch test in case F test showed significant

difference in variance, with Dunnett's post-hoc test. For lipidomic data multiple ratio paired t-

test was performed with a two-stage step-up method (Benjamini, Krieger, and Yekutieli), with

a false discovery rate set at 5%. Categorical quantification of plasma membrane recruitment

was performed by a blinded experimenter on randomized images. Contingency tables were

analysed using a Chi-squared test. Cells exhibiting abnormal morphology (multinucleated,

high vacuolation) or signs of death (blebbing) were excluded from analysis. A minimum of 5

replicate fields per coverslip are imaged for analysis of fixed cells. These may contain from 1

to tens of cells or 1 to tens of phagosomes which are considered as replicate values.

Data availability

Full datasets, including all microscopy and Western blot source images will be made available

on the University of Geneva's FAIR-compliant data repository Yareta under a CC BY 4.0

licence upon publication. Plasmids generated in this study will be deposited on Addgene upon

publication, to be distributed subject to a standard Material's Transfer Agreement.

Acknowledgements

We are grateful to the bioimaging and proteins and peptides core facilities at the University of

Geneva Medical Centre, Faculty of Medicine as well as the mass spectrometry core facility at

the Faculty of Science for their invaluable help. Thanks to Drs. Nicolas Demaurex, Maud

Frieden, Brenda Kwak and Amado Carreras-Sureda for their valuable advice. This work was

funded by a Professor Dr. Max Cloëtta Foundation Medical Researcher Grant (to PNH), a

Swiss National Science Foundation grant (310030_189094, to PNH), a Schmidheiney

Foundation subsidy (2022) to PNH. HR acknowledges the Swiss National Science Foundation

(310030_184949), the NCCR Chemical Biology (51NF40-185898) and the Leducq

Foundation. ACG. and LVE. acknowledge the financial support of the Louis-Jeantet

Foundation, Switzerland.

Author Contributions

VG designed and performed experiments, analysed data and wrote the manuscript.

LVE, IDQ, MA and MCC designed and performed experiments and analysed data. IR, and

MM performed experiments and analysed data. HR and ACG designed experiments,

provided guidance on analysis and interpretation of data as well as materials. PNH

conceived the project, designed and performed experiments and wrote the manuscript.

Competing Interests

The authors declare no competing interests.

References

Abdel Shakor, Abo Bakr, Katarzyna Kwiatkowska, and Andrzej Sobota. 2004. "Cell Surface

Ceramide Generation Precedes and Controls FcgammaRII Clustering and

Phosphorylation in Rafts." *The Journal of Biological Chemistry* 279 (35): 36778–87.

https://doi.org/10.1074/jbc.M402170200.

Abe, Mitsuhiro, and Toshihide Kobayashi. 2014. "Imaging Local Sphingomyelin-Rich

Domains in the Plasma Membrane Using Specific Probes and Advanced Microscopy."

Biochimica et Biophysica Acta - Molecular and Cell Biology of Lipids 1841 (5): 720–26.

https://doi.org/10.1016/j.bbalip.2013.07.003.

Adachi, Yoshifumi, George N. Pavlakis, and Terry D. Copeland. 1994. "Identification and

Characterization of SET, a Nuclear Phosphoprotein Encoded by the Translocation Break

Point in Acute Undifferentiated Leukemia." *Journal of Biological Chemistry* 269 (3):

2258–62. https://doi.org/10.1016/s0021-9258(17)42162-4.

Albi, E., and M. P. Viola Magni. 1997. "Chromatin Neutral Sphingomyelinase and Its Role in

Hepatic Regeneration." *Biochemical and Biophysical Research Communications* 236

(1): 29–33. https://doi.org/10.1006/BBRC.1997.6803.

Avota, Elita, Erich Gulbins, and Sibylle Schneider-Schaulies. 2011. "DC-SIGN Mediated Sphingomyelinase-Activation and Ceramide Generation Is Essential for Enhancement of Viral Uptake in Dendritic Cells." *PLoS Pathogens* 7 (2). https://doi.org/10.1371/journal.ppat.1001290.

Bakrač, Biserka, Aleš Kladnik, Peter Maček, Gavin McHaffie, Andreas Werner, Jeremy H. Lakey, and Gregor Anderluh. 2010. "A Toxin-Based Probe Reveals Cytoplasmic Exposure of Golgi Sphingomyelin." *Journal of Biological Chemistry* 285 (29): 22186–95. https://doi.org/10.1074/jbc.M110.105122.

Bidlingmaier, Scott, Kevin Ha, Nam Kyung Lee, Yang Su, and Bin Liu. 2016. "Proteome-Wide Identification of Novel Ceramide-Binding Proteins by Yeast Surface Cdna Display and Deep Sequencing." *Molecular and Cellular Proteomics* 15 (4): 1232–45. https://doi.org/10.1074/mcp.M115.055954.

Bjørnestad, Victoria Ariel, and Reidar Lund. 2023. "Pathways of Membrane Solubilization: A Structural Study of Model Lipid Vesicles Exposed to Classical Detergents." *Langmuir* 39 (11): 3914–33. https://doi.org/10.1021/acs.langmuir.2c03207.

Bockelmann, Svenja, John G.M. Mina, Sergei Korneev, Dina G. Hassan, Dagmar Müller, Angelika Hilderink, Hedwich C. Vlieg, et al. 2018. "A Search for Ceramide Binding Proteins Using Bifunctional Lipid Analogs Yields CERT-Related Protein StarD7." *Journal of Lipid Research* 59 (3): 515–30. https://doi.org/10.1194/jlr.M082354.

Bodman-Smith, Katherine B, Rachel E Gregory, Patrick T Harrison, and John G Raynes. 2004. "FcγRIIa Expression with FcγRI Results in C-Reactive Protein- and IgG-Mediated Phagocytosis." *Journal of Leukocyte Biology* 75 (6): 1029–35. https://doi.org/10.1189/jlb.0703306.

Bolte, S., and F. P. Cordelières. 2006. "A Guided Tour into Subcellular Colocalization Analysis in Light Microscopy." *Journal of Microscopy* 224 (3): 213–32. https://doi.org/10.1111/j.1365-2818.2006.01706.x.

Botelho, Roberto J., Mary Teruel, Renee Dierckman, Richard Anderson, Alan Wells, John D. York, Tobias Meyer, and Sergio Grinstein. 2000. "Localized Biphasic Changes in Phosphatidylinositol-4,5-Bisphosphate at Sites of Phagocytosis." *Journal of Cell Biology* 151 (7): 1353–67. https://doi.org/10.1083/jcb.151.7.1353.

Brennan, Jennifer A., Deanna J. Volle, Oleg V. Chaika, and Robert E. Lewis. 2002. "Phosphorylation Regulates the Nucleocytoplasmic Distribution of Kinase Suppressor of Ras." *Journal of Biological Chemistry* 277 (7): 5369–77. https://doi.org/10.1074/jbc.M109875200.

Buchberger, Amanda Rae, Kellen DeLaney, Jillian Johnson, and Lingjun Li. 2018. "Mass Spectrometry Imaging: A Review of Emerging Advancements and Future Insights." *Analytical Chemistry* 90 (1): 240–65. https://doi.org/10.1021/acs.analchem.7b04733.

Canals, Daniel, Silvia Salamone, and Yusuf A. Hannun. 2018. "Visualizing Bioactive
    Ceramides." *Chemistry and Physics of Lipids* 216 (September): 142–51.
    https://doi.org/10.1016/j.chemphyslip.2018.09.013.

Canton, Johnathan, Hanna Blees, Conor M Henry, Michael D Buck, Oliver Schulz, Neil C
    Rogers, Eleanor Childs, et al. 2021. "The Receptor DNGR-1 Signals for Phagosomal
    Rupture to Promote Cross-Presentation of Dead-Cell-Associated Antigens." *Nature
    Immunology* 22 (2): 140–53. https://doi.org/10.1038/s41590-020-00824-x.

Chen, Xiaoying, Jennica L. Zaro, and Wei Chiang Shen. 2013. "Fusion Protein Linkers:
    Property, Design and Functionality." *Advanced Drug Delivery Reviews* 65 (10): 1357.
    https://doi.org/10.1016/J.ADDR.2012.09.039.

Clarke, N. G., and R. M. Dawson. 1981. "Alkaline O Leads to N-Transacylation. A New
    Method for the Quantitative Deacylation of Phospholipids." *Biochemical Journal* 195
    (1): 301. https://doi.org/10.1042/BJ1950301.

Codazzi, Franca, Mary N. Teruel, and Tobias Meyer. 2001. "Control of Astrocyte Ca2+
    Oscillations and Waves by Oscillating Translocation and Activation of Protein Kinase
    C." *Current Biology* 11 (14): 1089–97. https://doi.org/10.1016/S0960-9822(01)00326-8.

Contreras, F. Xabier, Jesús Sot, Alicia Alonso, and Félix M. Goñi. 2006. "Sphingosine
    Increases the Permeability of Model and Cell Membranes." *Biophysical Journal* 90 (11):
    4085–92. https://doi.org/10.1529/biophysj.105.076471.

Deng, Yaqin, Lin You, Yong Lu, Sungwon Han, Jingcheng Wang, Nikitha Vicas, Chuo
    Chen, and Jin Ye. 2021. "Identification of TRAMs as Sphingolipid-Binding Proteins
    Using a Photoactivatable and Clickable Short-Chain Ceramide Analog." *Journal of
    Biological Chemistry* 297 (6): 101415. https://doi.org/10.1016/j.jbc.2021.101415.

Devaux, Philippe F. 1991. "Static and Dynamic Lipid Asymmetry in Cell Membranes."
    Biochemistry 30 (5): 1163–73.
    https://doi.org/10.1021/BI00219A001/ASSET/BI00219A001.FP.PNG_V03.

Ekman, Peik, Terhi Maula, Shou Yamaguchi, Tetsuya Yamamoto, Thomas K.M. Nyholm,
    Shigeo Katsumura, and J. Peter Slotte. 2015. "Formation of an Ordered Phase by
    Ceramides and Diacylglycerols in a Fluid Phosphatidylcholine Bilayer - Correlation
    with Structure and Hydrogen Bonding Capacity." *Biochimica et Biophysica Acta -
    Biomembranes* 1848 (10): 2111–17. https://doi.org/10.1016/j.bbamem.2015.06.019.

Eva Fuglsang, Louise Boye, and Hanne Frokiaer. 2017. "Enhancement of Ceramide
    Formation Increases Endocytosis of Lactobacillus Acidophilus and Leads to Increased
    IFN-Beta and IL-12 Production in Dendritic Cells. ." *Journal of Clinical Immunology
    Research* 1 (1): 1–9.

Gallegos, Lisa L, Maya T Kunkel, and Alexandra C Newton. 2006. "Targeting Protein
    Kinase C Activity Reporter to Discrete Intracellular Regions Reveals Spatiotemporal
    Differences in Agonist-Dependent Signaling." *The Journal of Biological Chemistry* 281

(41): 30947–56. https://doi.org/10.1074/jbc.M603741200.

Giodini, Alessandra, Christoph Rahner, and Peter Cresswell. 2009. "Receptor-Mediated Phagocytosis Elicits Cross-Presentation in Nonprofessional Antigen-Presenting Cells." *Proceedings of the National Academy of Sciences of the United States of America* 106 (9): 3324–29. https://doi.org/10.1073/pnas.0813305106.

Goñi, Félix M., and Alicia Alonso. 1999. "Structure and Functional Properties of Diacylglycerols in Membranes." *Progress in Lipid Research* 38 (1): 1–48. https://doi.org/10.1016/S0163-7827(98)00021-6.

Goñi, Félix M, F-Xabier Contreras, L-Ruth Montes, Jesús Sot, and Alicia Alonso. 2005. "Biophysics (and Sociology) of Ceramides." *Biochemical Society Symposium*, no. 72: 177–88. https://doi.org/10.1042/bss0720177.

Götz, Ralph, Tobias C. Kunz, Julian Fink, Franziska Solger, Jan Schlegel, Jürgen Seibel, Vera Kozjak-Pavlovic, Thomas Rudel, and Markus Sauer. 2020. "Nanoscale Imaging of Bacterial Infections by Sphingolipid Expansion Microscopy." *Nature Communications 2020 11:1* 11 (1): 1–9. https://doi.org/10.1038/s41467-020-19897-1.

Goulden, Brady D., Jonathan Pacheco, Allyson Dull, James P. Zewe, Alexander Deiters, and Gerald R.V. Hammond. 2019. "A High-Avidity Biosensor Reveals Plasma Membrane PI(3,4)P2 Is Predominantly a Class I PI3K Signaling Product." *Journal of Cell Biology* 218 (3): 1066–79. https://doi.org/10.1083/jcb.201809026.

Guido, Daniele, Nicolas Demaurex, and Paula Nunes. 2015. "Junctate Boosts Phagocytosis by Recruiting Endoplasmic Reticulum Ca 2 + Stores near Phagosomes," no. October: 4074–82. https://doi.org/10.1242/jcs.172510.

Guri, Yakir, Marco Colombi, Eva Dazert, Sravanth K. Hindupur, Jason Roszik, Suzette Moes, Paul Jenoe, et al. 2017. "MTORC2 Promotes Tumorigenesis via Lipid Synthesis." *Cancer Cell* 32 (6): 807-823.e12. https://doi.org/10.1016/J.CCELL.2017.11.011.

Hammond, Gerald R.V., Morgan M.C. Ricci, Claire C. Weckerly, and Rachel C. Wills. 2022. "An Update on Genetically Encoded Lipid Biosensors." *Molecular Biology of the Cell* 33 (5): 1–7. https://doi.org/10.1091/mbc.E21-07-0363.

Hannun, Yusuf A., and Lina M. Obeid. 2008. "Principles of Bioactive Lipid Signalling: Lessons from Sphingolipids." *Nature Reviews Molecular Cell Biology*. https://doi.org/10.1038/nrm2329.

———. 2018. "Sphingolipids and Their Metabolism in Physiology and Disease." *Nature Reviews Molecular Cell Biology*. Nature Publishing Group. https://doi.org/10.1038/nrm.2017.107.

Hinkovska-Galcheva, Vania, Laurence A Boxer, Andrei Kindzelskii, Miki Hiraoka, Akira Abe, Sravan Goparju, Sarah Spiegel, Howard R Petty, and James A Shayman. 2005. "Ceramide 1-Phosphate, a Mediator of Phagocytosis." *Journal of Biological Chemistry*

280 (28): 26612–21. https://doi.org/10.1074/jbc.M501359200.

Ivics, Zoltán, Meng Amy Li, Lajos Mátés, Jef D Boeke, Andras Nagy, Allan Bradley, and
Zsuzsanna Izsvák. 2009. "Transposon-Mediated Genome Manipulation in Vertebrates."
Nature Methods 6 (6): 415–22. https://doi.org/10.1038/nmeth.1332.

Jiménez-Rojo, Noemi, Manuel D. Leonetti, Valeria Zoni, Adai Colom, Suihan Feng, Namrata
R. Iyengar, Stefan Matile, et al. 2020. "Conserved Functions of Ether Lipids and
Sphingolipids in the Early Secretory Pathway." *Current Biology : CB* 30 (19): 3775-
3787.e7. https://doi.org/10.1016/J.CUB.2020.07.059.

Kiefer, Kerstin, Amado Carreras-Sureda, Roberto García-López, Fanny Rubio-Moscardó,
Josefina Casas, Gemma Fabriàs, and Rubén Vicente. 2015. "Coordinated Regulation of
the Orosomucoid-like Gene Family Expression Controls de Novo Ceramide Synthesis in
Mammalian Cells." *Journal of Biological Chemistry* 290 (5): 2822–30.
https://doi.org/10.1074/jbc.M114.595116.

Kiyokawa, Etsuko, Takeshi Baba, Naomi Otsuka, Asami Makino, Shinichi Ohno, and
Toshihide Kobayashi. 2005. "Spatial and Functional Heterogeneity of Sphingolipid-Rich
Membrane Domains." *Journal of Biological Chemistry* 280 (25): 24072–84.
https://doi.org/10.1074/JBC.M502244200.

Kjellberg, Matti A., Max Lönnfors, J. Peter Slotte, and Peter Mattjus. 2015. "Metabolic
Conversion of Ceramides in HeLa Cells - A Cholesteryl Phosphocholine Delivery
Approach." *PLoS ONE* 10 (11): 1–19. https://doi.org/10.1371/journal.pone.0143385.

Krishnamurthy, Kannan, Somsankar Dasgupta, and Erhard Bieberich. 2007. "Development
and Characterization of a Novel Anti-Ceramide Antibody." *Journal of Lipid Research*
48 (4): 968–75. https://doi.org/10.1194/jlr.D600043-JLR200.

Kwiatkowska, Katarzyna, Renata Hordejuk, Piotr Szymczyk, Magdalena Kulma, Abo-Bakr
Abdel-Shakor, Andrzej Płucienniczak, Krzysztof Dołowy, Adam Szewczyk, and
Andrzej Sobota. 2007. "Lysenin-His, a Sphingomyelin-Recognizing Toxin, Requires
Tryptophan 20 for Cation-Selective Channel Assembly but Not for Membrane Binding."
Molecular Membrane Biology 24 (2): 121–34.
https://doi.org/10.1080/09687860600995540.

Levin, Roni, Sergio Grinstein, and Johnathan Canton. 2016. "The Life Cycle of Phagosomes:
Formation, Maturation, and Resolution." *Immunological Reviews* 273 (1): 156–79.
https://doi.org/10.1111/imr.12439.

Levin, Roni, Sergio Grinstein, and Daniel Schlam. 2015. "Phosphoinositides in Phagocytosis
and Macropinocytosis." *Biochimica et Biophysica Acta - Molecular and Cell Biology of
Lipids* 1851 (6): 805–23. https://doi.org/10.1016/j.bbalip.2014.09.005.

Lorent, J. H., K. R. Levental, L. Ganesan, G. Rivera-Longsworth, E. Sezgin, M. Doktorova,
E. Lyman, and I. Leventhal. 2020. "Plasma Membranes Are Asymmetric in Lipid
Unsaturation, Packing and Protein Shape." *Nature Chemical Biology* 16 (6): 644–52.

https://doi.org/10.1038/s41589-020-0529-6.

Magenau, Astrid, Carola Benzing, Nicholas Proschogo, Anthony S. Don, Leila Hejazi,
Denuja Karunakaran, Wendy Jessup, and Katharina Gaus. 2011. "Phagocytosis of IgG-
Coated Polystyrene Beads by Macrophages Induces and Requires High Membrane
Order." *Traffic* 12 (12): 1730–43. https://doi.org/10.1111/j.1600-0854.2011.01272.x.

Majeski, Amy E., and J. Fred Dice. 2004. "Mechanisms of Chaperone-Mediated Autophagy."
The International Journal of Biochemistry & Cell Biology 36 (12): 2435–44.
https://doi.org/10.1016/J.BIOCEL.2004.02.013.

Mátés, Lajos, Marinee K L Chuah, Eyayu Belay, Boris Jerchow, Namitha Manoj, Abel
Acosta-Sanchez, Dawid P Grzela, et al. 2009. "Molecular Evolution of a Novel
Hyperactive Sleeping Beauty Transposase Enables Robust Stable Gene Transfer in
Vertebrates." *Nature Genetics* 41 (6): 753–61. https://doi.org/10.1038/ng.343.

Matheny, Sharon A, Chiyuan Chen, Robert L Kortum, Gina L Razidlo, Robert E Lewis, and
Michael A White. 2004. "Ras Regulates Assembly of Mitogenic Signalling Complexes
through the Effector Protein IMP." *Nature* 427 (6971): 256–60.
https://doi.org/10.1038/nature02237.

Mathias, Shalini, Louis A. Peña, and Richard N. Kolesnick. 1998. "Signal Transduction of
Stress via Ceramide." *Biochemical Journal* 335 (3): 465–80.
https://doi.org/10.1042/bj3350465.

Merrill, Alfred H., M. Cameron Sullards, Elaine Wang, Kenneth A. Voss, and Ronald T.
Riley. 2001. "Sphingolipid Metabolism: Roles in Signal Transduction and Disruption by
Fumonisins." *Environmental Health Perspectives* 109 (SUPPL. 2): 283–89.
https://doi.org/10.1289/ehp.01109s2283.

Miyake, Y., Y. Kozutsumi, S. Nakamura, T. Fujita, and T. Kawasaki. 1995. "Serine
Palmitoyltransferase Is the Primary Target of a Sphingosine-like Immunosuppressant,
ISP-1/Myriocin." *Biochemical and Biophysical Research Communications* 211 (2):
396–403. https://doi.org/10.1006/bbrc.1995.1827.

Montes, L Ruth, M Begoña Ruiz-Argüello, Félix M Goñi, and Alicia Alonso. 2002.
"Membrane Restructuring via Ceramide Results in Enhanced Solute Efflux." *The
Journal of Biological Chemistry* 277 (14): 11788–94.
https://doi.org/10.1074/jbc.M111568200.

Mukhopadhyay, Archana, Sahar A. Saddoughi, Pengfei Song, Iyad Sultan, Suriyan
Ponnusamy, Can E. Senkal, Christopher F. Snook, et al. 2009. "Direct Interaction
between the Inhibitor 2 and Ceramide via Sphingolipid-Protein Binding Is Involved in
the Regulation of Protein Phosphatase 2A Activity and Signaling." *FASEB Journal* 23
(3): 751–63. https://doi.org/10.1096/fj.08-120550.

Niekamp, Patrick, Gaelen Guzman, Hans C. Leier, Ali Rashidfarrokhi, Veronica Richina,
Fabian Pott, Caroline Barisch, Joost C. M. Holthuis, and Fikadu G. Tafesse. 2021.

"Sphingomyelin Biosynthesis Is Essential for Phagocytic Signaling during *Mycobacterium Tuberculosis* Host Cell Entry." Edited by Christina L. Stallings. *MBio* 12 (1). https://doi.org/10.1128/mBio.03141-20.

Niekamp, Patrick, Felix Scharte, Tolulope Sokoya, Laura Vittadello, Yeongho Kim, Yongqiang Deng, Elisabeth Südhoff, et al. 2022. "Ca2+-Activated Sphingomyelin Scrambling and Turnover Mediate ESCRT-Independent Lysosomal Repair." *Nature Communications* 13 (1). https://doi.org/10.1038/s41467-022-29481-4.

Nunes-Hasler, Paula, Sophia Maschalidi, Carla Lippens, Cyril Castelbou, Samuel Bouvet, Daniele Guido, Flavien Bermont, et al. 2017. "STIM1 Promotes Migration, Phagosomal Maturation and Antigen Cross-Presentation in Dendritic Cells." *Nature Communications* 8 (1): 1852. https://doi.org/10.1038/s41467-017-01600-6.

Nunes, Paula, Daniela Cornut, Vanessa Bochet, Udo Hasler, Masatsugu Oh-Hora, Jean Marc Waldburger, and Nicolas Demaurex. 2012. "STIM1 Juxtaposes ER to Phagosomes, Generating Ca2+ Hotspots That Boost Phagocytosis." *Current Biology* 22 (21): 1990– 97. https://doi.org/10.1016/j.cub.2012.08.049.

Pathak, Divya, Neelay Mehendale, Shubham Singh, Roop Mallik, and Siddhesh S. Kamat. 2018. "Lipidomics Suggests a New Role for Ceramide Synthase in Phagocytosis." *ACS Chemical Biology* 13 (8): 2280–87. https://doi.org/10.1021/acschembio.8b00438.

Pédelacq, Jean Denis, Stéphanie Cabantous, Timothy Tran, Thomas C. Terwilliger, and Geoffrey S. Waldo. 2006. "Engineering and Characterization of a Superfolder Green Fluorescent Protein." *Nature Biotechnology* 24 (1): 79–88. https://doi.org/10.1038/nbt1172.

Pérez-rivas, Luís G, Stephanie Prieto, and José Lozano. 2010. "1 . Modulation of Ras Signaling by the KSR Family of Molecular Scaffolds." *Emerging Signaling Pathways in Tumor Biology* 661 (2): 1–23. http://www.trnres.com/ebook/uploads//T_12700118641 Lazo corrected.pdf.

Pohl, Antje, Iván López-Montero, Florent Rouvière, Fabrice Giusti, and Philippe F. Devaux. 2009. "Rapid Transmembrane Diffusion of Ceramide and Dihydroceramide Spin-Labelled Analogues in the Liquid Ordered Phase." *Http://Dx.Doi.Org/10.1080/09687680902733815* 26 (3): 194–204. https://doi.org/10.1080/09687680902733815.

Quinville, Brianna M, Natalie M Deschenes, Alex E Ryckman, Jagdeep S Walia, N M ; Ryckman, and A E ; Walia. 2021. "Molecular Sciences A Comprehensive Review: Sphingolipid Metabolism and Implications of Disruption in Sphingolipid Homeostasis." https://doi.org/10.3390/ijms22115793.

Razidlo, Gina L., Robert L. Kortum, Jamie L. Haferbier, and Robert E. Lewis. 2004. "Phosphorylation Regulates KSR1 Stability, ERK Activation, and Cell Proliferation." *Journal of Biological Chemistry* 279 (46): 47808–14.

https://doi.org/10.1074/jbc.M406395200.

Rinaldi, L, R Delle Donne, M Sepe, M Porpora, C Garbi, F Chiuso, A Gallo, et al. 2016. "Praja2 Regulates KSR1 Stability and Mitogenic Signaling." *Cell Death & Disease* 7 (5): e2230. https://doi.org/10.1038/cddis.2016.109.

Saharan, Ojal, Neelay Mehendale, and Siddhesh S. Kamat. 2022. "Phagocytosis: A (Sphingo)Lipid Story." *Current Research in Chemical Biology* 2 (March): 100030. https://doi.org/10.1016/j.crchbi.2022.100030.

Saliba, Antoine Emmanuel, Ivana Vonkova, Stefano Ceschia, Greg M. Findlay, Kenji Maeda, Christian Tischer, Samy Deghou, et al. 2014. "A Quantitative Liposome Microarray to Systematically Characterize Protein-Lipid Interactions." *Nature Methods* 11 (1): 47–50. https://doi.org/10.1038/nmeth.2734.

Sallusto, Federica, Chiara Nicolò, Ruggero De Maria, Silvia Corinti, and Roberto Testi. 1996. "Ceramide Inhibits Antigen Uptake and Presentation by Dendritic Cells." *Journal of Experimental Medicine* 184 (6): 2411–16. https://doi.org/10.1084/jem.184.6.2411.

Santos, Nina Criado, Samuel Bouvet, Flavien Bermont, Cyril Castelbou, Farah Mansour, Maral Azam, Francesca Giordano, and Paula Nunes-Hasler. 2022. "The SNARE Sec22b Regulates Phagosome Maturation by Promoting ORP8-Mediated PI(4)P Exchange at ER-Phagosome Contact Sites." *BioRxiv*, January, 2022.04.12.487993. https://doi.org/10.1101/2022.04.12.487993.

Schneider, Caroline A., Wayne S. Rasband, and Kevin W. Eliceiri. 2012. "NIH Image to ImageJ: 25 Years of Image Analysis." *Nature Methods* 9 (7): 671–75. https://doi.org/10.1038/NMETH.2089.

Schramm, Michael, Jasmin Herz, Albert Haas, Martin Krönke, and Olaf Utermöhlen. 2008. "Acid Sphingomyelinase Is Required for Efficient Phago-Lysosomal Fusion." *Cellular Microbiology* 10 (9): 1839–53. https://doi.org/10.1111/j.1462-5822.2008.01169.x.

Scorrano, Luca, Maria Antonietta De Matteis, Scott Emr, Francesca Giordano, György Hajnóczky, Benoît Kornmann, Laura L. Lackner, et al. 2019. "Coming Together to Define Membrane Contact Sites." *Nature Communications* 10 (1): 1–11. https://doi.org/10.1038/s41467-019-09253-3.

Shiraishi, Takayuki, Shinichiro Imai, and Yutaka Uda. 2003. "The Presence of Ceramidase Activity in Liver Nuclear Membrane." *Biological & Pharmaceutical Bulletin* 26 (6): 775–79. https://doi.org/10.1248/BPB.26.775.

Stewart, Scott, Meera Sundaram, Yanping Zhang, Jeeyong Lee, Min Han, and Kun-Liang Guan. 1999. "Kinase Suppressor of Ras Forms a Multiprotein Signaling Complex and Modulates MEK Localization." *Molecular and Cellular Biology* 19 (8): 5523–34. https://doi.org/10.1128/mcb.19.8.5523.

Suzuki, Jun, Toshihiro Fujii, Takeshi Imao, Kenji Ishihara, Hiroshi Kuba, and Shigekazu

Nagata. 2013. "Calcium-Dependent Phospholipid Scramblase Activity of TMEM16
Protein Family Members." *The Journal of Biological Chemistry* 288 (19): 13305–16.
https://doi.org/10.1074/jbc.M113.457937.

Tafesse, Fikadu G., Ali Rashidfarrokhi, Florian I. Schmidt, Elizaveta Freinkman, Stephanie
Dougan, Michael Dougan, Alexandre Esteban, Takeshi Maruyama, Karin Strijbis, and
Hidde L. Ploegh. 2015. "Disruption of Sphingolipid Biosynthesis Blocks Phagocytosis
of Candida Albicans." Edited by Alex Andrianopoulos. *PLoS Pathogens* 11 (10):
e1005188. https://doi.org/10.1371/journal.ppat.1005188.

Tang, Jonathan C Y, Eugene Drokhlyansky, Behzad Etemad, Stephanie Rudolph, Binggege
Guo, Sui Wang, Emily G Ellis, Jonathan Z Li, and Constance L Cepko. 2016.
"Detection and Manipulation of Live Antigen-Expressing Cells Using Conditionally
Stable Nanobodies." Edited by Liqun Luo. *ELife* 5: e15312.
https://doi.org/10.7554/eLife.15312.

Tran, Thi Thu Trang, Bárbara Graziela Postal, Sylvie Demignot, Agnès Ribeiro, Céline
Osinski, Jean Paul Pais De Barros, Agnieszka Blachnio-Zabielska, et al. 2016. "Short
Term Palmitate Supply Impairs Intestinal Insulin Signaling via Ceramide Production."
The Journal of Biological Chemistry 291 (31): 16328.
https://doi.org/10.1074/JBC.M115.709626.

Veiga, M. Pilar, José Luis R. Arrondo, Félix M. Goñi, and Alicia Alonso. 1999. "Ceramides
in Phospholipid Membranes: Effects on Bilayer Stability and Transition to Nonlamellar
Phases." *Biophysical Journal* 76 (1 I): 342–50. https://doi.org/10.1016/S0006-
3495(99)77201-2.

Vieira, O V, R J Botelho, L Rameh, S M Brachmann, T Matsuo, H W Davidson, A Schreiber,
J M Backer, L C Cantley, and S Grinstein. 2001. "Distinct Roles of Class I and Class III
Phosphatidylinositol 3-Kinases in Phagosome Formation and Maturation." *The Journal
of Cell Biology* 155 (1): 19–25. https://doi.org/10.1083/jcb.200107069.

Walter, T., J. Schlegel, A. Burgert, A. Kurz, J. Seibel, and M. Sauer. 2017. "Incorporation
Studies of Clickable Ceramides in Jurkat Cell Plasma Membranes." *Chemical
Communications* 53 (51): 6836–39. https://doi.org/10.1039/C7CC01220A.

Wang, Guanghu, Kannan Krishnamurthy, Nagavedi S. Umapathy, Alexander D. Verin, and
Erhard Bieberich. 2009. "The Carboxyl-Terminal Domain of Atypical Protein Kinase Cζ
Binds to Ceramide and Regulates Junction Formation in Epithelial Cells." *Journal of
Biological Chemistry* 284 (21): 14469–75. https://doi.org/10.1074/jbc.M808909200.

Wang, Guanghu, Jeane Silva, Kannan Krishnamurthy, Eric Tran, Brian G Condie, and Erhard
Bieberich. 2005. "Direct Binding to Ceramide Activates Protein Kinase Czeta before the
Formation of a Pro-Apoptotic Complex with PAR-4 in Differentiating Stem Cells." *The
Journal of Biological Chemistry* 280 (28): 26415–24.
https://doi.org/10.1074/jbc.M501492200.

Williamson, Patrick. 2015. "Phospholipid Scramblases." *Lipid Insights* 8 (Suppl 1): 41–44. https://doi.org/10.4137/LPI.S31785.

Wu, Yuqing, Cao Li, Andrea Riehle, Barbara Pollmeier, Erich Gulbins, and Heike Grassmé. 2018. "Mycobacterial Infection Is Promoted by Neutral Sphingomyelinase 2 Regulating a Signaling Cascade Leading to Activation of B1-Integrin." *Cellular Physiology and Biochemistry* 51 (4): 1815–29. https://doi.org/10.1159/000495683.

Yamaji, A, Y Sekizawa, K Emoto, H Sakuraba, K Inoue, H Kobayashi, and M Umeda. 1998. "Lysenin, a Novel Sphingomyelin-Specific Binding Protein." *The Journal of Biological Chemistry* 273 (9): 5300–5306. https://doi.org/10.1074/jbc.273.9.5300.

Yeung, Tony, and Sergio Grinstein. 2007. "Lipid Signaling and the Modulation of Surface Charge during Phagocytosis." *Immunological Reviews* 219 (1): 17–36. https://doi.org/10.1111/j.1600-065X.2007.00546.x.

Yin, Xianglei, Mohammad Zafrullah, Hyunmi Lee, Adriana Haimovitz-Friedman, Zvi Fuks, and Richard Kolesnick. 2009. "A Ceramide-Binding C1 Domain Mediates Kinase Suppressor of Ras Membrane Translocation." *Cellular Physiology and Biochemistry* 24 (3–4): 219–30. https://doi.org/10.1159/000233248.

Zhang, Yuhua, Yao Bei, Sylvie Delikat, Shariff Bayoumy, Lin Xin-Hua, Subham Basu, Michael McGinley, Po Ying Chan-Hui, Henri Lichenstein, and Richard Kolesnick. 1997. "Kinase Suppressor of Ras Is Ceramide-Activated Protein Kinase." *Cell* 89 (1): 63–72. https://doi.org/10.1016/S0092-8674(00)80183-X.

Zhao, Hongmei, Malgorzata Przybylska, I. Huan Wu, Jinhua Zhang, Craig Siegel, Svetlana Komarnitsky, Nelson S. Yew, and Seng H. Cheng. 2007. "Inhibiting Glycosphingolipid Synthesis Improves Glycemic Control and Insulin Sensitivity in Animal Models of Type 2 Diabetes." *Diabetes* 56 (5): 1210–18. https://doi.org/10.2337/db06-0719.

Zhao, Lin, Jia Zhao, Kunhong Zhong, Aiping Tong, and Da Jia. 2022. "Targeted Protein Degradation: Mechanisms, Strategies and Application." *Signal Transduction and Targeted Therapy* 7 (1): 113. https://doi.org/10.1038/s41392-022-00966-4.

Zhou, Kecheng, Andrea Dichlberger, Hector Martinez-Seara, Thomas K.M. Nyholm, Shiqian Li, Young Ah Kim, Ilpo Vattulainen, Elina Ikonen, and Tomas Blom. 2018. "A Ceramide-Regulated Element in the Late Endosomal Protein LAPTM4B Controls Amino Acid Transporter Interaction." *ACS Central Science* 4 (5): 548–58. https://doi.org/10.1021/acscentsci.7b00582.

Zhou, Ming, David A. Horita, David S. Waugh, R. Andrew Byrd, and Deborah K. Morrison. 2002. "Solution Structure and Functional Analysis of the Cysteine-Rich C1 Domain of Kinase Suppressor of Ras (KSR)." *Journal of Molecular Biology* 315 (3): 435–46. https://doi.org/10.1006/jmbi.2001.5263.

Figure 1. *Localization of candidate ceramide (Cer) probes in mouse embryonic fibroblasts (MEFs).* **A)** Schematic representation of Cer-binding domains used for probe construction from the following source proteins: Human SET earmuff domain (EMD), amino acids (as) 69-226; atypical PKC type zeta C1 domain (PRKCZ-C1, aa 123-193) and C20 domain (encompassing also the kinase domain, PRKCZ-C20 aa 405-592); and KSR1 CA3 domain (containing an atypical C1 domain, KSR, aa 317-400). **B-D)** Confocal images of MEF cells transfected with N-terminally (N-) or C-terminally (C-) EGFP-tagged Cer-binding domains of SET (B), PRKCZ (C) and KSR1 (D). Constructs preceded by a nuclear export signal are labelled NES, the tandem construct is labelled 2X and the cysteine mutants C359S/C362S. The inset in C-KSR highlights enlarged vesicular structures. White bars = 10 μm, yellow bar = 3 μm.

Figure 2. *Response of C- and N-KSR to Cer-modifying drugs.* **A, C)** Confocal images of MEF cells stably expressing C-KSR (A) or N-KSR (B) (in green) treated with; 0.5 µM myriocin (Myr) or 2.5 µM fumonisin B1 (FumB1) FumB1 for 3d; or 0.5 mM BSA-complexed palmitate (Palm) for 4h (bottom panels), and respective vehicle controls (top panels). **B-D)** Quantification of total cellular mean fluorescence intensity (MFI) of cells in A and C respectively. Bars are means +/- SEM, (n= 3 coverslips for each condition). White bars = 10 µm. **E)** Targeted lipidomic analysis of MEF cells stably expressing C-KSR, treated with Myr as above. Left panel shows the quantification of the sum total mass of lipids classified as ceramides (Cer), hexosyl-ceramides (HexCer), and sphingomyelins (SM), normalized to the total mass glycerophospholipids in the same sample (expressed as percent mole, %mol). Inset shows the quantification of the 5 most abundant Cer (left) HexCer (middle) and SM (right) species grouped by fatty acid chain length. **F)** Similar analysis as in **E)** of cells treated for 4h with Palm. Bars are means + SEM. (n=3, see also Table S1).

Figure 3. *C-KSR responds to Sphingomyelinase-induced Cer.* **A)** Confocal images of MEF cells stably expressing C-KSR (green) treated with 0.5 U/ml bacterial sphingomyelinase (SMase) or PBS control for 30 min. **B)** Quantification of total cellular MFI of cells in A), as well as fluorescence of the plasma membrane (PM) only of cells in A). (n=4 coverslips) **C)** Blinded quantification of the fraction of cells displaying PM localization in PBS or SMase-treated condition in A). (Pooled cells from n=4 coverslips is 170 PBS/202 SMase-treated) A-B). **D)** Targeted lipidomic analysis of MEF cells stably expressing C-KSR, treated with SMase as above. Left upper panel shows the quantification of the sum total Cer, HexCer, SM. Insets show the quantification of the 5 most abundant Cer (i) HexCer (ii, right) and SM (ii, left) species grouped by fatty acid chain length. (n=3, see also Table S1) **E)** Confocal images of MEF cells co-transfected with C-KSR-mRFP (magenta) and diacylglycerol (DAG) probe PCK-C1(2)-EGFP (green), control (Ctl, top) or treated with 10 μM ionomycin (Iono) for 10 min (middle), or with SMase as above (bottom). **F)** Blinded quantification of cells in E displaying PM localization of either C-KSR or PKC-C1(2) probe. (n= 6/3/3 for Ctl/Iono/SMase). Bars are means + SEM. White bars = 10 μm, yellow bar = 3 μm.

Figure 4. *Generation of a more stable C-KSR probe, C-KSR-GS.* **A)** Confocal images of MEF cells transfected with C-KSR (green) and loaded for 30 min with recycling endosome marker Alexa555-conjugated human transferrin (Tfn, 50 µg/ml, magenta top); or immunostained with late endosome marker anti-Rab7 (magenta middle) or lysosomal marker anti-LAMP1 (magenta bottom). The arrow points to an enlarged vesicular structure showing peripheral C-KSR localization with a hollow lumen, that is surrounded yet by LAMP1 staining. **B)** Quantification of co-localization between C-KSR and organelle marker, expressed as Mander's coefficient (n=4/4/14 coverslips for Tfn/Rab7/LAMP1). **C)** Western blot of lysates of MEF cells stably expressing C-KSR or N-KSR or transfected with cytosolic EGFP and probed with anti-GFP antibody. **D)** Schematic representation of the linker mutagenesis employed to create the new probe C-KSR-GS. **E)** Western blot of MEF cells transfected with either C-KSR-GS or EGFP, and stained first with anti-GFP, then re-probed with anti-tubulin (Tub). Bars are means + SEM. White bars = 10 µm, yellow bars = 3 µm.

Figure 5. *C-KSR-GS responds to Cer modulation similarly to C-KSR.* **A-C)** Confocal images of MEFs transfected with C-KSR-GS and treated with 10 μM FumB1 (bottom) for 1d or methanol (MeOH) vehicle control (top) (A); 0.5 mM Palm for 4 h (bottom) or BSA control (top) (B); or 0.5 U/ml SMase for 30 min (bottom) or PBS control (top) (C). Right panels shows the quantification of total cellular fluorescence (MFI) (n=3 coverslips, for A-B) or blinded quantification of the fraction of cells displaying prominent PM localization (C, n=4 coverslips pooled, 168/237 cells PBS/SMase). **D-E)** Confocal images of MEFs expressing C-KSR-GS (D) or N-KSR (E) and treated (bottom) with either Cer analog C6 (200 μM, 2h, left) or 100 μM of the GlcCer synthase inhibitor Genz-123346 (Genz, 1d, right), or their respective ethanol (EtOH) or dimethylsulfoxide (DMSO) vehicle controls (top). Right panels show the quantification of total cellular fluorescence (MFI) (in D: n=3/3/4/3 coverslips for EtOH/C6/DMSO/Genz; in E: n = 4 coverslips for all conditions). Bars are means +(or +/-) SEM. White bars = 10 μm.

Figure 6. *Liposome microarray (LiMa) analysis of KSR probes.* **A)** Heatmap summarizing the automated quantification of liposome binding in LiMa analysis using purified N-KSR and C-KSR and DOPC-based liposomes. Values are expressed as Log(2) transformed ratios of the mean normalized binding indices (NBI) for each liposome composition tested as compared to the corresponding score for DOPC liposomes alone (n=3/1-2 for N-KSR/C-KSR, see also Table S2) **B)** Representative epifluorescence images of liposome array spots showing liposome membranes labelled in PE-Atto647 (magenta) or probe fluorescence (green, N-KSR left, C-KSR right) **C)** Heatmap of LiMa analysis as in A) of C-KSR-GS or cytosolic EGFP obtained from HEK293 lysates and inner plasma membrane (IPM)-mimicking liposomes (n=2-6, see also Table S2). **D)** Representative epifluorescence images as in B) for C-KSR-GS (left) and EGFP (right), in the context of IPM liposomes (see also Fig S5) . White bars = 10 µm.

Figure 7. *Behavior of KSR probes during phagocytosis in MEFs.* **A-C)** Confocal images of phagocytic MEFs expressing (in green) C-KSR (A), N-KSR (B) and C-KSR-GS (C), exposed to IgG-Alexa647-coupled beads at 1:10 cell:bead ratio for 0 (left), 30 (middle) and 135 (right) min. **D)** Targeted lipidomic analysis of MEF cells stably expressing C-KSR, treated exposed to IgG-beads as above. Top panel shows the quantification of the sum total Cer, HexCer, SM. Quantification of the 5 most abundant species grouped by fatty acid chain length is shown for Cer (i) SM (ii) and HexCer (iii) species. (n=3, see also Table S3). **E-F)** Confocal images of phagocytic MEFs co-transfected with C-KSR-GS (green) and proteins known to localize to membrane contact sites between ER and phagosomes mCh-STIM1 (E, magenta) and mCh-Sec22b (F, magenta), and exposed to IgG-beads as above. White arrows show periphagosomal regions where overlap of fluorescent protein signal is discerned, green arrow shows regions with green fluorescence only. Bars are means +SEM. White bars = 3 μm.

Figure 8. *Behavior of KSR probes during phagocytosis in dendritic cells.* **A-B)** Confocal images of dendritic cell line JAWS transfected with (in green) N-KSR (A) exposed to IgG-coupled beads at 1:10 cell:bead ratio for 30 (left) and 135 (right) min, and C-KSR-GS (B) exposed to beads for 30 min. Brightfield images are shown to confirm presence of internalized beads. Bars = 3 µm.