Maturation and detoxification of synphilin-1 inclusion bodies regulated by sphingolipids

Xiuling Cao[1,2]†, Xiang Wu[1]†, Lei Zhao[2]†, Ju Zheng[2,3], Xuejiao Jin[1], Xinxin Hao[2], Joris Winderickx[3], Shenkui Liu[1], Lihua Chen[2,4]*, Beidong Liu[1,2]*

[1] State Key Laboratory of Subtropical Silviculture, Zhejiang A&F University, Hangzhou, China. [2] Department of Chemistry and Molecular Biology, University of Gothenburg, Gothenburg, Sweden. [3] Functional Biology, KU Leuven, Leuven, Belgium. [4] Guangzhou Laboratory, Guangzhou, Guangdong, China.

† These authors contributed equally to this work.

* Corresponding author. Email: beidong.liu@cmb.gu.se (B.L.); chen_lihua@gzlab.ac.cn (L.C.)

Abstract

Due to proteostasis stress induced by aging or disease, misfolded proteins can form toxic intermediate species of aggregates and eventually mature into less toxic inclusion bodies (IBs). Here, using a yeast imaging-based screen, we identified 84 potential synphilin-1 (SY1) IB regulators and isolated the conserved sphingolipid metabolic components in the most enriched groups. Furthermore, we show that, in both yeast cells and mammalian cells, SY1 IBs are associated with mitochondria. Pharmacological inhibition of the sphingolipid metabolism pathway or knockout of its key genes results in a delayed IB maturation and increased SY1 cytotoxicity. We postulate that SY1 IB matures by association with the mitochondrion membrane, and that sphingolipids stimulate the maturation via their membrane-modulating function and thereby protecting cells from SY1 cytotoxicity. Our findings identify a conserved cellular component essential for IB maturation and suggest a mechanism by which cells may detoxify the pathogenic protein aggregates through forming mitochondrion-associated IBs.

Introduction

Proteins are vulnerable macromolecules and are often endangered by misfolding and aggregation during the

life of the cell due to destabilizing mutations, stress conditions or unique metabolic challenges that occur

during aging or disease (Buchberger et al., 2010; Hartl and Hayer-Hartl, 2009). To cope with protein damage

and in order to survive, the eukaryotic cell has evolved complex protein quality control systems that monitor

and maintain the integrity of its proteome, including refolding, degradation and sequestration (Buchberger

et al., 2010; Frydman, 2001; Hartl and Hayer-Hartl, 2009; Lim and Yue, 2015). The refolding of stress-

denatured proteins and the degradation of misfolded proteins are assisted by molecular chaperones (Chen et

al., 2011; Hartl and Hayer-Hartl, 2009). In addition to refolding and degradation, cells have another cellular

strategy to cope with an overload of aberrant proteins or toxic intermediate species of aggregates, namely

sequestration into specialized quality control IBs (Bagola and Sommer, 2008; Kaganovich et al., 2008;

Kopito, 2000).

It has become increasingly clear that the formation of IBs is like a "triage" center, thus being protective

machinery against proteostasis stress during aging or disease. Therefore, the inability to form IBs would

poison the intracellular environment and cause cell dysfunction or even death (Lim and Yue, 2015). The

toxicity of protein aggregates is more pronounced in neurons partly due to their unique cellular architecture.

Indeed, mounting evidence has linked a number of neurodegenerative diseases including Alzheimer's

disease (AD), Parkinson's disease (PD), Huntington's disease (HD) and amyotrophic lateral sclerosis (ALS),

to the formation of misfolded proteins aggregates containing IBs (Ross and Poirier, 2004). Although IBs

have been extensively studied, including their unique characteristics (Kaganovich *et al.*, 2008; Ogrodnik et

al., 2014; Weisberg et al., 2012) and the IB clearance machinery (Moldavski et al., 2015), the genes and

cellular components that regulate the IB maturation remain poorly understood. In addition, we have little

insight into the effects of IBs on cellular membrane-bound structures as IBs have been found to interact with

membrane domains (Büttner et al., 2010).

SY1 contributes to inclusion formation and has been implicated in the pathogenesis of PD, an incurable,

progressive neurodegenerative disorder. The biological functions of SY1 are not fully understood, and

whether SY1 is cytotoxic or protective remains controversial. In aged yeast cells, SY1 expression reduced

survival and triggered apoptosis and necrotic cell death in a Sir2 dependent manner (Büttner *et al.*, 2010).

However, SY1 exerted neuroprotective effects by inhibiting reactive oxygen species (ROS) production and

apoptosis in an *in vitro* PD model (Shishido et al., 2019), and SY1 attenuated mutant LRRK2-induced PD-

like phenotypes and played a neuronal protective role in cultured cells and *Drosophila* (Liu et al., 2016). Transgenic mice expressing SY1 caused pathological inclusions and degeneration of dopaminergic neurons in the substantia nigra (Krenz et al., 2009), and reduced motor skill learning and motor performance (Jin et al., 2008; Nuber et al., 2010). In contrast, SY1 overexpression in A53T human α-synuclein (α-Syn) transgenic mice reduced astrogliosis and neuronal degeneration, resulting in increased lifespan(Smith et al., 2010). In an aged A30P α-Syn transgenic mouse model, Casadei *et al* observed that overexpression of SY1 promoted clearance of soluble and misfolded α-Syn, but did not rescue the motor deficit phenotype (Casadei et al., 2014).

Given that a high-content imaging approach has been successfully conducted for SY1 (Zhao et al., 2016), it was possible to study on a genome-wide scale the cellular components that are involved in IB formation by using yeast as a model. Moreover, since SY1 binds phospholipids, membranes, lipid droplets and synaptic vesicles (Ribeiro et al., 2002; Takahashi et al., 2006) and given that the membrane-binding properties of α-Syn are thought to be relevant for its pathological activity, we speculated that also for SY1 the binding to different types of lipid structures might be related to its cytotoxicity. Here we combined the SGA approach with a high-content imaging-based screen to survey the genetic mutants with altered SY1 aggregation types using a yeast genome-wide mutant library. We identified the conserved sphingolipid metabolic pathway as an essential component for the maturation of SY1 IBs and for the detoxification of SY1 aggregates. In the sphingolipid-rich endomembrane system, SY1 aggregates tightly interact with mitochondria. Disruption of sphingolipid synthesis prevented this association, resulting in small, dispersed aggregates and enhanced SY1-induced cytotoxicity, notably mitochondrial toxicity. Furthermore, pharmacological inhibition of the sphingolipid pathway or knockdown of SPTLC2 altered the morphology of SY1 IBs and increased their cytotoxicity in mammalian cells, and this could be reversed by increasing the level of dihydrosphingosine (DHS), a sphingolipid synthesis intermediate.

Results

SY1 IB management in yeast cells

The formation of SY1 IBs in the yeast (*Saccharomyces cerevisiae*) has been studied by expressing the protein (DsRed-SY1) with a plasmid pYX212 under the control of a constitutive *TPI1* promoter(Zhao *et al.*, 2016).

Using a similar approach, we observed different phenotypes of SY1 aggregates in yeast cells. During the lag

phase (4 h), multiple small IBs presented in the cytoplasm but in the exponential growth phase (16 h), these

small aggregates merged into larger IBs, resulting in a decreased number of smaller aggregates. Subsequently,

when the cells entered the stationary phase (48 h), most of the SY1-expressing cells retained only one to a

few large IBs (Figure 1A). Thus, consistent with previously reported data (Büttner *et al.*, 2010), this time-

course study demonstrated a maturation process of SY1 IBs in yeast cells. To study the details of this process,

we defined the IB phenotypes as Class 1 (one aggregate in a cell), Class 2 (two aggregates in a cell), and

Class 3 (three or more aggregates in a cell), respectively (Figure 1B), as previously reported for mutant

huntingtin inclusions (Yang et al., 2016). We then monitored the SY1 IBs every 4 hours from 4 to 48 hours.

Most of the SY1 IB-forming cells exhibited a Class 3 phenotype (96%) at the 4-hour time-point. During

growth, the proportion of Class 3 ratio dropped over time, while the proportion of Class 1 and Class 2 ratios

increased inversely. After 24 hours, the proportions became stable and Class 1 IBs accounted for

approximately 50% when cells reached stationary phase (Figure 1C). All these observations supported the

notion that there is a compartmentalization process of SY1 IBs in the yeast cell. We therefore hypothesized

that the formation and maturation of IBs is a cytoprotective process that helps sequester more toxic

aggregates/oligomers into larger inert deposits (Kitamura et al., 2006).

Dysregulation of the formation of pathogenic protein inclusions is the major hallmark of age-related

neurodegenerative diseases. To investigate whether IBs formed by SY1 expression in old cells have the same

maturation process as that in young cells. We isolated old cells and examined the SY1 aggregates in old cells

($\geq$5 generations) and young cells ($\leq$1 generation) (Figure 1D). In young cells, about 27% cells had SY1

inclusions while in the old cells the ratio increased to 61% (Figure 1E). Furthermore, when counted the

number of aggregates in individual cells, we found that there were much more aggregates per cell in the old

cell population (Figure 1F, 4.91±3.32 of old cells vs 2.64±1.29 of young cells). These findings suggested

that there was a dysregulation of SY1 IB maturation in old cells. However, the underlying mechanism

remains unclear. This motivated us to further investigate the genes and cellular components that regulate

SY1 IB maturation.

Genome-wide screening for cellular components regulating the maturation of SY1 inclusions

To dissect the genes and cellular components that mediate the maturation of IBs, a genome-wide, high-

content imaging-based screening approach was performed (Zhao *et al.*, 2016) (Figure 2A). Prior to

performing the screen, we determined the appropriate time-point to assess the phenotypes of SY1 IBs. As

shown in Figure 1A-C, at 16-20 hours of growth (OD_{600} is 3.0 ~ 4.0), approximately 50% SY1-expressing

cells had Class 3 aggregates, and the ratios were stable thereafter. Therefore, we chose the time point 20-

hours for the following screening and other studies. We set the same criteria as in the previous study (Zhao

et al., 2016) to include a particular mutant strain as a potential hit if: 1) the difference in the number of cells

with Class 3 SY1 IBs between wild-type and mutant cells is statistically significant ($P \leq 0.05$, *t*-test) and 2)

the absolute difference of the percentage of wild-type and mutant cells carrying SY1 inclusions is at least

20%. Potential hits acquired from the screening were then confirmed by manual examination of the Class 3

inclusion phenotype using a microscopic approach.

The screen was performed three times independently. Based on the above criteria and subsequent manual

confirmation, a list of 84 mutants that showed an increase in cells carrying Class 3 SY1 IBs was obtained

(Supplementary file 1). GO enrichment analysis indicated that these mutants were enriched in the following

modules: serine C-palmitoyltransferase (SPT) complex, SPOTS complex, protein urmylation, TORC2

complex, palmitoyltransferase complex, and others (Figure 2B and 2C). The complete list of GO terms can

be found in Supplementary file 2. Among them, SPT complex and SPOTS complex, containing LCB1, LCB2

and TSC3 had the highest enrichment (Table S2, Figure 2B and 2C), corresponding to the mutants including

lcb1-4, *lcb2-1*, *lcb2-2* and *tsc3-2* in our screen. We performed gene complementation experiments on three

mutants *lcb1-4*, *lcb2-1* and *tsc3-2*. Restoring the corresponding gene almost completely rescued the SY1 IB

phenotype with significantly reduced Class 3 IBs (Figure 2D and 2E). Thus, our above screen suggested a

link between the highly conserved sphingolipid metabolic pathway and the maturation of SY1 IBs. A

substantial body of evidence has highlighted an important role of sphingolipid metabolism in several

neurodegenerative diseases, including Alzheimer's disease (Mielke and Lyketsos, 2010), amyotrophic lateral

sclerosis(Cutler et al., 2002), and Parkinson's disease (France-Lanord et al., 1997). Therefore, we aimed to

elucidate the functional role of sphingolipid metabolism in the regulation of SY1 IB maturation and

cytotoxicity.

Sphingolipids regulate SY1 IB maturation and cytotoxicity in yeast cells

The first step of intracellular sphingolipid de novo synthesis is identical in both yeast and humans, with

palmitoyl-CoA and serine catalyzed by the SPT complex to form 3-Ketosphinganine, which could be

blocked by myriocin (Figure 3A). 3-ketosphinganine is subsequently catalyzed to dihydrosphingosine

(DHS), and dihydroceramide, and further converted into complex sphingolipids. We first confirmed that

exogenous addition of DHS in SPT mutants (*lcb1-4*, *lcb2-1* and *tsc3-2*) could rescue the phenotype of

increased Class 3 SY1 IBs (Figure 3B and 3C). In addition, blockade of intracellular sphingolipid

metabolism pathways by myriocin strongly perturbed SY1 IB maturation process, resulting in elevated Class

3 IBs (Figure 3D and 3E). Again, addition of exogenous DHS could suppress the myriocin-induced Class 3

aggregates (Figure 3D and 3E). Next, we tested whether the SPT genes could regulate SY1 cytotoxicity in

yeast cells. Expressing SY1 did not cause dramatic growth defect in wild-type cells, but led to severe growth

defects in the SPT mutants (Figure 3F, left). Furthermore, this defect could also be rescued by the addition

of DHS (Figure 3F, right). Taken together, these data suggest that sphingolipids may help cells reduce the

cytotoxicity of protein aggregates by regulating the IB maturation process. These findings motivated us to

further unravel the underlying mechanism by which sphingolipids regulate the maturation process and

cytotoxicity of SY1 IBs.

SY1 IBs associate with mitochondria in yeast cells

Sphingolipids are structural molecules of cell membranes that play an important role in maintaining barrier

function and fluidity (Bento-Oliveira et al., 2020; Hannun and Obeid, 2008; Schlarmann et al., 2021). One

of the primary modes of action of sphingolipids is to modulate membrane fluidity, thickness and curvature.

In this context, we asked whether there is a direct association of SY1 IBs with certain membrane-bound

organelles in the yeast cell. To address this question, we examined the location of SY1 IBs (20 h) in cells

and their possible association with membranous organelles visualized by corresponding marker proteins.

Our results showed that SY1 IBs had no obvious association with the plasma membrane (Pma1-GFP),

vacuole (Vph1-GFP), or ER (Sec63-GFP), but were strongly associated with mitochondria (Tom70-GFP)

(Figure 4A and Figure 4-figure supplement 1A, about 80%). Since SY1 IBs have three different

morphologies (Class 1-3) and the Class 3 IBs transition to Class 1 during maturation (Figure 1A-C), we

wanted to confirm whether all these different IBs associate with mitochondria in a similar way. The results

confirmed that all the three different IBs were equivalently associated with mitochondria (Figure 4B and

Figure 4-figure supplement 1B). Also, super-resolution microscopy and 3D-SIM techniques demonstrated

that the SY1 IBs were surrounded by mitochondria (Figure 4C).

The next question is whether other pathogenic IBs are also associated with mitochondria. To address this

question, we used the same expression system to overexpress TDP43, FUS1 or Htt103Qp (GFP-labeled) in

yeast cells expressing Tom70-RFP and examined the possible association with mitochondria. Our results

showed that the IBs of these proteins indeed interacted with mitochondria, similar to what we found for the

IBs of SY1 (Figure 4-figure supplement 2). Hence, it appears that the association of IBs with mitochondria

may be a common biological event among these disease proteins. To further study the consequences of this

association and to decipher potential factors that regulate this association, we first investigated whether SY1

IBs physically associate with mitochondria. We performed four co-immunoprecipitation (Co-IP)

experiments to determine whether SY1 could be detected in pull-downs for the inner mitochondrial

membrane protein Tim44 or the outer mitochondrial membrane proteins Tom70, Tom37, and Sam35. The

SY1 protein was indeed the case in extracts obtained from the Tom70-GFP, Tom37-GFP and Tim44-GFP

strains (Figure 4D). Next, we performed a microscopic analysis and found that the association of SY1 IBs

with mitochondria was significantly hampered in *tom70Δ, tom37Δ* and *tim44* mutants (Figure 4E and 4F).

Collectively, these results confirm that SY1 IBs are in physical contact with mitochondrial membrane

components Tom70, Tom37 and Tim44, which is consistent with the fact that SY1 is a membrane lipid raft

interacting protein (Büttner *et al.*, 2010; O'Farrell et al., 2001; Takahashi *et al.*, 2006).

Sphingolipids regulate SY1-induced mitochondrial toxicity in yeast cells

Several lines of evidence have supported the notion that potentially toxic proteins such as α-Syn and TDP-

43 can associate with mitochondria and damage the mitochondrial membrane, leading to cytotoxicity

(Bäuerlein et al., 2017; Busciglio et al., 2002; Hashimoto et al., 2003; Hsu et al., 2000; Park et al., 2020;

Wang et al., 2019a; Wang et al., 2019b). For example, pathogenic α-Syn aggregates preferentially bind to

mitochondria, leading to decreased mitochondrial biogenesis (Park *et al*., 2020) and impaired cellular

respiration (Wang *et al*., 2019b). To explore whether the interaction between SY1 IBs and mitochondrial

membranes could impair mitochondrial function, we measured mitochondrial membrane potential,

intracellular ROS levels, and cell viability represented by ATP levels in SY1-expressing cells. The results

showed that mitochondrial membrane potential and cell viability of SY1-expressing cells were significantly

lower than that of control cells, while ROS levels were increased in SY1 expressing cells (Figure 5A). These

changes indicate a decrease in mitochondrial function caused by SY1 IBs. Given that SY1 IBs interact with

sphingolipid-enriched membrane raft domains and that sphingolipid metabolism regulates the maturation of

SY1 IBs, we wondered whether sphingolipids play a role in the observed SY1-induced mitochondrial

dysfunction. To address this question, we first blocked the sphingolipid synthesis by myriocin in the SY1-

expressing yeast cells and measured mitochondrial membrane potential, intracellular ROS and cell viability.

We found that myriocin treatment aggravated SY-1-induced cytotoxicity, as indicated by decreased

mitochondrial membrane potential and cell viability, and increased intracellular ROS (Figure 5B), without

altering SY1 protein expression (Figure 5-figure supplement 1A). Second, we added exogenous DHS to

SY1-expressing cells and observed ameliorated SY-1-induced cytotoxicity as indicated by increased

mitochondrial membrane potential and cell viability, as well as decreased intracellular ROS (Figure 5C),

without altering SY1 protein expression (Figure 5-figure supplement 1B). Collectively, these results showed

that sphingolipids regulated the mitochondrial toxicity induced by SY1 IBs, and suggested that sphingolipids

might detoxify SY1 IBs through mitochondria.

SPTLC2 knockout alters the morphology of SY1 IBs and increases the cytotoxicity in mammalian

cells

We then wondered whether the observed association with mitochondria and the cytotoxicity of SY1 IBs was

conserved in mammalian cells. Similarly, SY1 forms IBs in mammalian cells, which are known as

aggresomes (Zaarur et al., 2008). First, we expressed SY1 in HEK293t cells to study the association of SY1

IBs with mitochondria. A 3D structure between SY1 IB and mitochondria was constructed by fluorescence

3D reconstruction and it confirmed that in these mammalian cells the SY1 IBs are in close contact with

mitochondria, although the morphology of the mitochondrion is different (Figure 6A). Furthermore, the

mitochondrial membrane potential was significantly decreased in SY1-expressing cells, while the levels of

intracellular ROS and mitochondrial superoxide were significantly increased (Figure 6B), indicating

impaired mitochondrial function by SY1 IBs. As such, our data suggest that in yeast and mammalian cells

the formation and processing of SY1 IBs occur in a similar manner, leading to mitochondrial dysfunction

through influencing the mitochondrial membranes.

To further confirm this, we analyzed whether sphingolipids are indeed involved in the formation and

cytotoxicity of SY1 IBs in mammalian cells as well. To this end, we used the CRISPR/Cas9 approach to

knock out SPTLC2, the ortholog of yeast LCB2 and one of the subunits of the mammalian SPT complex

(Figure 6-figure supplement 1). Notably, knockout of SPTLC2 significantly reduced the accumulation of

intracellular sphingolipids, but had little effect on normal cell division and morphology (Tafesse et al., 2015).

Microscopic analysis showed that the morphology of SY1 IBs formed in mammalian cells can be divided

into two forms, one compact and spherical and the other more dispersed with an irregular structure as if

many small IBs stuck together, without clear boundaries or uniform centers (Figure 6C). In wild-type cells,

the SY1 IBs were predominantly spherical. However, in SPTLC2 knockout cells, the morphology of SY1

IBs was significantly altered, with an increased number of dispersed irregular IBs (Figure 6D) and an

increased number of SY1 IBs per cell (Figure 6-figure supplement 2). This is consistent with the phenotypes

observed in SPT mutant yeast cells (Figure 3B). Furthermore, mitochondrial function, as judged from cell

viability, intracellular ROS levels and mitochondrial membrane potential, was decreased in the SPTLC2-

deficient cells (Figure 6E), while the SY1 protein expression was not altered (Figure 6-figure supplement 3).

Taken together, these results further substantiate that SY1 IBs are processed and managed similarly in

mammalian cells and in yeast cells and that the sphingolipid pathway plays a role in the cytotoxicity of SY1

IBs in mammalian cells as well.

Sphingolipids regulate SY1-induced cytotoxicity and SY1 IB morphology in mammalian cells

Because the initial steps of sphingolipid biosynthesis are conserved from yeast to humans, myriocin also

inhibits the de novo synthesis of intracellular sphingolipids in mammalian cells (Ogretmen, 2018). We added

myriocin to SY1-expressing HEK293t cells and examined the corresponding changes in cytotoxicity. The

results showed that inhibition of the intracellular sphingolipid synthesis pathway by myriocin enhanced SY1

cytotoxicity based on ATP content, intracellular ROS and mitochondrial membrane potential assays (Figure

7A). Furthermore, the addition of exogenous DHS rescued the cytotoxicity caused by SY1 expression

(Figure 7B) without altering SY1 protein expression (Figure 7-figure supplement 1A and B).

We further examined the cytotoxicity of SY1 in SPTLC2 knockout cells with or without DHS. The results

showed that DHS restored the morphology of IBs (Figure 7C), as well as the number of SY1 IBs formed in

SPTLC2 knockout cells (Figure 7-figure supplement 2). Accordingly, DHS also ameliorated the SY1-

instigated cytotoxicity by normalizing cell viability, intracellular ROS, and mitochondrial membrane

potential (Figure 7D), without affecting SY1 expression (Figure 7-figure supplement 1C). These results

reinforce the notion that sphingolipid metabolism plays an important role in the formation of SY1 IBs and

SY1-induced cytotoxicity.

Discussion

In this study, we used a genome-wide high-content imaging-based screening approach to identify regulators

of SY1 IBs formation. Among the 5500 mutants, 84 mutants exhibited increased Class 3 SY1 inclusions.

GO enrichment analysis revealed the SPT complex as the most enriched functional group. Subsequent

studies revealed that the sphingolipid pathway plays an important role in the processing of SY1 IBs and the

detoxification of SY1 by mitochondria in yeast cells. Corresponding experiments in mammalian cells

recapitulated the role of sphingolipids in the detoxification of SY1-induced mitochondrial toxicity. Taken

together, our results suggest a role for sphingolipids in controlling the cytotoxicity of pathologically

aggregated proteins in cells.

IBs management and their cytotoxicity

The formation and accumulation of abnormal intracellular protein aggregates, including the pathological

aggregates observed in neurodegenerative diseases, are generally considered to be a double-edged sword.

On the one hand, their ability to co-aggregate and sequester basic cellular components, including molecular

chaperones, causes cytotoxicity (Gong et al., 2012; Olzscha et al., 2011; Park et al., 2013). On the other hand,

formation of protein aggregates can also protect cells against highly toxic, soluble oligomeric substances

(Arrasate et al., 2004; Douglas et al., 2008; Tanaka et al., 2004). Many studies have shown that converting

small aggregates/oligomers into IBs can reduce the cytotoxicity of various aggregated proteins (Miller et al.,

2015). In line with a previous report (Büttner *et al.*, 2010), our current study demonstrated that the merging

of smaller foci into IBs also occurred when SY1 was expressed in yeast and mammalian cells.

Association of mitochondrial with protein aggregates

The formation of large intracellular IBs leads to severe impairment of membrane function of IB-bound

organelles, thereby affecting organelle function. This has been confirmed in several experiments, such as the

disruptive effect of Huntingtin polyQ IBs on ER function (Bäuerlein *et al.*, 2017) and the effect of neuronal

C9orf72 poly-GA aggregates on the Proteasome (Guo et al., 2018). Several lines of evidence implicate

mitochondrial dysfunction as a primary cause of PD. Environmental toxins such as mitochondrial complex

inhibitors rotenone and 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine induce dopaminergic neuronal death

(Polito et al., 2016). In addition, mitochondrial complex I deficiency and mitochondrial electron transport

chain defects have been reported in the nigra of PD patients (Haelterman et al., 2014). However, it is unclear

how these protein aggregates affect the mitochondria. As previously reported, small foci formed by SY1

started to merge into larger deposits when cells approached the diauxic shift to reprogram metabolism from

fermentation to respiration (Büttner *et al.*, 2010), which was confirmed by our present study (Figure 1A). In

addition, we show that SY1 physically interacts with mitochondrial membrane proteins and that larger SY1

deposits are surrounded by mitochondrial structures in both yeast and mammalian cells (Figure 4A-D and

Figure 6A). However, it is unknown why these IBs preferentially interact with mitochondria. This preference

may be partly explained by a recent study in which the authors showed that the mitochondrial import receptor

Tom70 could nucleate stress-induced aggregation of cytosolic proteins (Liu, 2021). In addition, impaired

mitochondria could form cytosolic aggregates and thereby stimulate the aggregation of other proteins,

including many pathogenic proteins such as α-Syn and amyloid β (Bento-Oliveira *et al.*, 2020). In this study,

however, we did not investigate the influence of mitochondria on SY1 IBs formation. Nevertheless, our

current study shows a fine management of aggregated proteins In eukaryotic cells—protein IBs are tightly

associated with mitochondria once formed and continue to mature onto mitochondria.

In addition to SY1, there is a similar association of IBs and mitochondria with other aggregation-prone

proteins, including TDP43, FUS1 and Htt103Qp (Figure 4-figure supplement 2), suggesting that

mitochondrion-contact may be a common feature for the maturation of pathogenic protein IBs. Additional

experiments are needed to confirm this hypothesis, although an interaction with mitochondria has already

been described for the aforementioned and several other aggregation-prone proteins (reviewed in (Abramov

et al., 2017)). Furthermore, it would be of great value to illustrate how these IBs are sequestered to

mitochondria and how the smaller IBs merge into large IBs with the help of mitochondria.

The role of sphingolipids in SY1 IB maturation and mitochondrial function.

Sphingolipids are structural molecules of cell membranes with important roles in maintaining barrier

function and fluidity (Hannun and Obeid, 2008). In addition, bioactive sphingolipids have variety of

functions that are involved in almost all major aspects of cell biology, and key sphingolipid metabolites are

critical for normal brain development and function (Hannun and Obeid, 2018). Membranes and membrane

dynamics are required for these processes and an imbalance of lipids, including ceramide and sphingolipid

intermediates, may contribute to mitochondrial dysfunction that could lead to the demise of neurons (Lin et

al., 2019). SY1 inclusions co-localize with lipid droplets and endomembranes, and SY1 interacts with

cytoplasmic lipid rafts in mammalian cells and yeast (Büttner *et al.*, 2010; Takahashi *et al.*, 2006). The C-

terminus of SY1 was found to bind selectively to acidic phospholipids, including phosphatidic acid,

phosphatidylserine, and phosphatidylglycerol, but not to naturally charged phospholipids and the amino acid

sequence 610-640 was found to represent the major determinant of phospholipid binding (Takahashi *et al.*,
2006). The opening question is how sphingolipids contribute to the SY1 IB maturation: do they merely
provide a structural basis for docking or do they have an active function in triggering this process. Moreover,
the relationship between sphingolipids and mitochondrial function during aging is not fully understood. It
has long been appreciated that mitochondrial activity and quality decline with age (reviewed in(Amorim et
al., 2022; Lima et al., 2022)). On the other hand, sphingolipids play critical roles in the regulation of lifespan
in yeast, worms, and flies, and in the regulation of senescence in mammalian cells (Li and Kim, 2022; Singh
and Li, 2018; Trayssac et al., 2018). However, the changes in sphingolipid profiles during aging have not
been fully characterized. In mammalian cells, there is an accumulation of ceramide and sphingosine during
aging (Sacket et al., 2009), but the levels of other species including long-chain base sphingolipids are not
reported. In this regard, additional studies will be needed to profile the sphingolipid species in the aged cells
and to identify the major sphingolipids that regulate mitochondrial function.

Materials and methods

Plasmids

All plasmids used in this study are listed in Supplementary file 3. The plasmid pYX212-DsRed-SY1-His
was constructed by replacing the URA3 cassette with the His3MX6 cassette by homologous recombination
in yeast. The corresponding plasmids for complementation experiments were derived from the MoBY library
(Ho et al., 2009). The plasmids in this collection were constructed based on the p5472 plasmid with its native
promoter and terminator. To construct plasmid pcDNA3.1-GFP-SY1, the GFP fragment and the SY1
fragment were inserted into the Not I/Hind III digested pcDNA3.1 (-) vector using Exnase (Vazyme, C112).
For the construction of plasmid pcDNA3.1-SY1-Flag, the PCR-expanded SY1-Flag fragment was inserted
directly into the Not I/Hind III digested pcDNA3.1 (-) vector.

Yeast strains and growth conditions

The yeast strains used in this study are listed in Supplemental file 4. Experiments were mainly performed in
the BY4741 strain (MATa *his3Δ1 leu2Δ0 met15Δ0 ura3Δ0*) and its respective mutants. The *S. cerevisiae*
MATα strain Y7092 (Tong and Boone, 2007) was used for query strain constructions. The yeast single gene
knockout collection (SGA-V2) and the essential gene temperature sensitive allele collection (ts-V5) were

kindly provided by Prof. Charlie Boone, University of Toronto, Canada. Strains expressing Pma1-GFP,

Sec63-GFP, Vph1-GFP, Tom70-GFP, Tom37-GFP, Sam35-GFP, and Tim44-GFP were obtained from the

yeast GFP collection (Huh et al., 2003). Yeast strains were grown in YPD (1% yeast extract, 2% peptone,

and 2% glucose) or appropriate synthetic drop-out medium (0.17% yeast nitrogen base without amino acids

and ammonium sulfate, 0.5% ammonium sulfate and the appropriate synthetic drop-out amino acid, and 2%

glucose). Yeast cells were grown at 30°C or at 22°C in case of temperature sensitive (ts) alleles.

SGA analysis

The experimental methods are based on a previous report (Zhao *et al.*, 2016), and essentially consist of

transforming the plasmids pYX212-Dsred-SY1 and pYX212-Dsred (control), respectively, into the yeast

query strain Y7092. Subsequently, SGA mating was performed as previously described (Tong et al., 2001;

Tong et al., 2004) to introduce the plasmids pYX212-Dsred-SY1 or pYX212-Dsred into the SGA-V2 and

ts-V5 collections, respectively. All pinning steps for collection handling were performed using a SINGER

ROTOR HDA robot (Singer Instrument, Watchet UK).

For the genome-wide high content imaging screening of SY1 IB morphology, the constructed collections

expressing Dsred-SY1 were pre-cultured in 200 μL SD-Ura liquid medium in 96-well plates with antibiotics

at 30°C (22°C for ts mutants) for 3 days without shaking. Then, each pre-culture sample was diluted to

OD_{600nm} 0.1 to a final volume of 200 μL of SD-Ura medium. After 20 h (24 h for ts mutants) of incubation

with shaking at 30°C (for both single and ts mutants), cells were fixed by adding formaldehyde to a final

concentration of 3.7% for 30 min at room temperature and then washed twice with 1×PBS. For imaging,

appropriate amounts of fixed cells were transferred to new 96-well glass bottom plates (Matri-plate) with

200 μL of 1×PBS in each well and imaged using ImageXpress MICRO (Molecular Devices Corporation

(MDC), San Jose, USA), an automated cellular imaging system. To automatically quantify cells with

different SY1 aggregation phenotypes, a customized subprogram of the MetaXpress software (MDC) was

used (Zhao *et al.*, 2016). Image preprocessing with shading correction and background subtraction was first

performed; cells and IBs were separated from their background based on differences in signal intensity,

respectively. Then, the number of IBs per cell was extracted by combining the two masks (cell and inclusion

body). The percentages of the three types (class 1 to class 3) of cells were counted. All mutants that showed

statistically significant differences as compared to the wild-type were re-streaked, individually tested, and

further manually analyzed to confirm the phenotypic differences observed in the screen. At least 300 cells

were counted for manual confirmation. For manual confirmation, cells were grown from a starting OD_{600nm}

of 0.1 in 15 ml Falcon tubes for 20 h (24 h for ts mutants) and fixed and washed as described above. All

manual confirmations were verified using a Zeiss Axio Observer 7 microscope (Carl Zeiss, Jena, Germany).

Spot assays

374 For spot assays, the mutants carrying DsRed-SY1 were cultured in 3 mL SD-Ura liquid medium at 22°C for

375 24 h. The strains were diluted to $OD_{600\ nm} = 0.5$ and then serially diluted tenfold in sterile water in 96-well

376 plates. The dilutions were spotted on plates with or without DHS. Images were captured after incubation at

377 30°C for 2 days.

Bioinformatic analysis

379 Candidate hits were analyzed for enrichment in the Gene Ontology (GO) using ClueGO (V2.5.9) (Bindea et

380 al., 2009) in Cytoscape (V3.9.1) with a cut-off of $P < 0.05$. Bubble and network plots of significantly altered

381 GO terms and genes were generated using OmicStudio tools (https://www.omicstudio.cn/tool) and

382 Cytoscape.

Isolation of SY1-expressing old yeast cells

Old yeast cells overexpressing SY1 were obtained by magnetic sorting (Smeal et al., 1996). The wild-type

BY4741 cells expressing Dsred-SY1 were cultured in 100 mL SD-Ura liquid medium, and then harvested

after growth to 0.5 with an OD_{600nm} of 0.1, washed three times in ice-cold PBS pH7.4, and incubated with 1

mg/mL sulfo-NHS-LC biotin (Thermo Scientific, 21335) labeling for 20 min at room temperature. Cells

were further washed four times with cold PBS pH 8.0 and grown overnight in 100 mL new fresh medium.

Cells were harvested the next day and washed three times with cold PBS pH 7.4, and incubated with 250

mL streptavidin-conjugated paramagnetic beads (5 mg/ml, Thermo Scientific, 88817) for 2 h at 4°C.

Unlabeled cells were removed by magnetic sorting. The cells were then grown overnight in 100 mL fresh

SD-Ura medium. The isolation process was repeated two more times. At the final isolation, cells were

collected from the supernatant of the first wash and resuspended in PBS to serve as the young cells, while

cells labeled with magnetic beads were centrifuged and resuspended in PBS to serve as old cells. The yeast

bud scars were photographed through the DAPI channel using a Zeiss Axio Observer 7 fluorescence

microscope to confirm and calculate the degree of yeast senescence. Subsequently, the SY1 aggregates were

photographed through the Dsred channel to calculate the type of SY1 aggregates.

Complementation of yeast mutants with MoBY plasmid

The mutants expressing pYX212-DsRed-SY1 were transformed with the corresponding MoBY plasmid or empty control vector (p5472). For the morphological observation experiment of SY1 IBs, the cells were cultured in SD-Ura, His liquid medium at 30ºC starting from $OD_{600nm} = 0.1$ for 20 h. The cells were then inspected by a fluorescence microscope and photographed.

Myriocin treatment and DHS supplementation in yeast

Myriocin (MCE, HY-N6798) was dissolved in DMSO at a concentration of 10 mM. The stock solution was incubated in a 55°C water bath for 15 min prior to use. DHS (dihydrosphingosine, Sigma, D3314) was prepared in distilled DMSO to 30 mM stock solution. For treatments, cells were grown overnight and diluted in medium to an initial OD_{600nm} of 0.1 with myriocin or DHS at a final concentration of 25 or 20 μM, respectively.

Microscopic analysis of SY1 IBs in yeast

For the microscopic analysis of SY1 IBs in yeast and the co-localization analyses of SY1 IBs with fluorescently labeled proteins Pma1-GFP, Sec63-GFP, Vph1-GFP, and Tom70-GFP, pYX212-DsRed-SY1 was transformed into corresponding cells and yeast strains were grown overnight and diluted in culture medium to a starting OD_{600nm} of 0.1. After shaking at 30°C for 20 h, the cells were fixed by adding formaldehyde. Cell images were captured using Zeiss Axio Observer 7. For mitochondrial staining analysis, MitoTracker Green (Invitrogen, M7514) staining solution was prepared according to the manufacturer's protocol. After 20 h of growth from an OD_{600nm} of 0.1, $1×10^7$ cells were fixed with formaldehyde for 30 min and then washed twice in 1×PBS (pH 7.4). The cells were resuspended in 1× PBS (pH 7.4) containing 125 nM MitoTracker probe and incubated for 30 min at room temperature. After one wash with PBS, the cells were imaged by microscopy.

3D-SIM microscopy

3D-SIM microscopy images were obtained as previously reported (Song et al., 2014). Briefly, yeast strain Tom70-GFP expressing pYX212-DsRed-SY1 was grown overnight and diluted in medium to a starting OD_{600nm} of 0.1. After shaking at 30°C for 20 h, cells were fixed with 3.7% formaldehyde and washed twice with 1× PBS (pH 7.4). The ELYRA PS.1 LSM780 setup from Zeiss (Carl Zeiss, Jena, Germany) was used

for 3D-SIM. Images were acquired with a 100×/1.46 Plan-Apochromat oil immersion objective. Z-stacks

with 100 nm spacing were used to scan the whole yeast in 3D-SIM. Zeiss Zen microscope software was used

for acquisition, super-resolution processing and calculation, and 3D reconstruction. The ELYRA system was

corrected for chromatic aberration in x-, y- and z-directions using multi-color beads, and all obtained images

were inspected and aligned accordingly.

Co-IP for SY1 with mitochondrial membrane proteins in yeast

pYX212-DsRed-SY1 and pYX212-DsRed were transformed into GFP strains (Tom70-GFP, Tom37-GFP,

Sam35-GFP, and Tim44-GFP) respectively. The strains were pre-cultured and diluted to 100 mL medium

with OD_{600nm}=0.1. Cells were harvested when OD_{600nm} reached 1.0 by centrifugation at 4000 rpm for 5 min

at 4°C and washed twice with 20 mL ice-cold ddH$_2$O water. Cells were then washed with 1 mL ice-cold IP

buffer (115 mM KCl, 5 mM NaCl, 2 mM MgCl$_2$, 1 mM EDTA, 20 mM HEPES/KOH, pH 7.5), and

transferred to 1.5 mL Eppendorf tubes. Cell pellets were resuspended in 250 μL IP buffer containing protease

inhibitors, DTT, and glass beads and disrupted using a FastPrep-24 device (MP Biomedicals). Cells were

further lysed on ice for 20 min with additional IP buffer containing 0.5% Nonidet P-40 (NP-40, Beyotime,

ST2045). The lysates were centrifuged at 16,000 g for 20 min at 4°C, and 700-800 μL of supernatant was

transferred to a new tube. For the IP assay, 20 μL anti-GFP beads (chromoTek, GFP-Trap) were resuspended

in 100 μL IP buffer and mixed with the supernatant. The mixture was incubated for 1-1.5 h at 4°C on an

overhead rotator, centrifuged at 2000 rpm for 1 min, and washed 4 times with 1 mL IP buffer containing 0.5%

NP-40. The final eluted proteins were boiled with 2 x Laemmli buffer / 5% beta-mercaptoethanol for 10 min

at 95°C and the supernatant was used for Western blot analysis.

Detection of ROS and mitochondrial membrane potential in yeast cells

To examine ROS levels and mitochondrial membrane potential in yeast cells, BY4741 yeast cells expressing

pYX212-DsRed-SY1 or pYX212-DsRed were pre-cultured overnight and diluted to an OD_{600nm} of 0.1. Cells

were harvested at the endpoints, washed twice with 1× PBS (pH 7.4), and treated with DCFH-DA (Beyotime,

S0033S) or Rhodamine123 (Beyotime, C2007) at a final concentration of 10 μM. Cells were incubated for

20 min at 30°C in the dark. Cells were then washed at least three times with 1×PBS (pH 7.4) before flow

cytometric analysis using BD Accuri C6 (BD Biosciences, San Jose, USA) with FITC channel.

Cell lines, culture condition, transfection and drug treatment

The human embryonic kidney (HEK293t) cell line was obtained from the National Collection of

Authenticated Cell Cultures (NCACC; Shanghai, China). Cells were cultured in Dulbecco's modified Eagle's

medium (DMEM; Gibco, c11995500bt) supplemented with 10% fetal bovine serum (FBS; Biological

Industries, 04-001-1acs) and 1% penicillin-streptomycin (HycloneTM, SV30010) in a humidified incubator

containing 5% CO_2 at 37°C. Transfection was performed on six-well plates or confocal dishes for different

purposes with plasmids using X-tremeGENE HP DNA Transfection Reagent (Roche, 6366236001)

according to the manufacturer's protocol. For drug treatment, cells were seeded at appropriate density in cell

culture plates and cultured for 24 h. Cells were treated with 25 μM myriocin for 3 days or 20 μM DHS for 2

days before subsequent analysis.

Cell imaging in mammalian cells

To observe the association of SY1 IBs with mitochondria in mammalian cells, HEK293t cells were seeded

into 35 mm confocal cell culture dishes (BioSharp), transfected with GFP-SY1, and incubated for 24 h. The

cells were washed with PBS and then stained with 10 μM MitoTracker Deep Red FM (Invitrogen, M22426)

at 37°C for 20 min. After staining, the cells were washed twice with PBS, and rinsed with PBS. Fluorescence

images were captured using a Zeiss LSM 880 confocal microscope (Carl Zeiss, Jena, Germany), and Z-stack

images for 3D rendering were taken at 0.4 μm interlayer distance. 3D rendering was performed using the

Surface tool in the Imaris software (version 9.0.1; Bitplane AG).

For analysis of SY1 IBs formation under myriocin or DHS treatment, HEK293t cells were seeded in confocal

culture dishes and transfected with pcDNA3.1-GFP-SY1 after treatment. Cells were fixed with 4%

paraformaldehyde (Beyotime, P0099) for 20 min and permeabilized with 0.1% (v/v) Triton X-100 in PBS

(Beyotime, P0096) for 15 min on ice. The cytoskeleton was then stained with phalloidin-Atto 565 (Sigma,

94072), followed by nuclei with DAPI (Southern Biotech, 0100-20). Images were captured using a Zeiss

LSM 880 confocal microscope.

Measurement of mitochondrial function in mammalian cells

HEK293t cells were detached and transferred to 75-mm flow tubes and exposed to 5 μM MitoSOX

(Invitrogen, M36008) or 10 μM DCFH-DA (Beyotime, S0033S) in serum-free medium for 10 min

(MitoSOX) or 20 min (DCFH-DA) in a 37°C cell incubator. Cells were then washed three times with serum-

free cell culture medium and analyzed using a BD Accuri C6 flow cytometer.

Mitochondrial membrane potential (MMP) was measured with JC-1 (Beyotime, C2006) according to the

manufacturer's instructions. Briefly, treated cells were harvested, washed twice with PBS, and then stained

in a mixture of 500 μL medium and 500 μL JC-1 at 37°C for 20 min in the dark. Cells were washed 3 times

with staining buffer and analyzed by flow cytometry. Changes in MMP are represented by the ratio of red to

green fluorescence intensity.

Measurement of cell viability and cytotoxicity

Cell viability was measured by monitoring the cellular ATP content using a CellTiter-Lumi Luminescent

Cell Viability Assay Kit (Beyotime, C0065S). Cells were treated and further incubated with 100 μL of

detection reagent at room temperature for 2 min to promote cell lysis. Cells were further incubated at room

temperature for 10 min to stabilize the luminescence signal, and ATP levels were then analyzed by using a

Spark multimode microplate reader (Tecan Trading AG, Männedorf, Switzerland).

Western blot analysis

Cells were harvested and lysed in RIPA buffer (Beyotime, P0013C) supplemented with a protease inhibitor

of 1 mM PMSF (Beyotime, ST506). Total protein concentrations were then quantified using the BCA Protein

Quantification Kit (Beyotime, P0012S). A total of 20 μg of protein from each sample was separated by SDS-

PAGE and transferred to nitrocellulose membranes (Cytiva, 10600001). The membranes were blocked with

1% BSA for 1 h at 37°C and incubated with primary antibodies: anti-Flag (Cell Signaling, D6W5B, 1:1000

dilution), anti-GFP (Abcam, ab6556, 1:1000 dilution), anti-SY1 (Affinity, DF8619, 1:1000 dilution), anti-

PGK1 (Abcam, ab113687, 1:1000 dilution), anti-GAPDH (Wanleibio, WL01114, 1:1000 dilution), followed

by HRP-conjugated secondary antibody incubation: goat anti-rabbit IgG (Abcam, ab6721, 1:10000 dilution).

Protein bands were detected with ECL Prime Western Blotting Detection Reagent (GE, RPN2232) using an

Amersham imager 600 (GE, Piscataway, USA).

Generation of HEK293t SPTLC2 knockout cells by CRISPR/Cas9

For the generation of SPTLC2 knockout cell line, guide RNA sequences for CRISPR were designed at

CRISPR design web site (http://crispr.mit.edu/). Guide RNA sequences for human SPTLC2 were #1: 5'-

CGAATGGCTGCGTGGCGAAC-3' and #2: 5'-AAGTACGGAACGGGTACGTG-3', respectively. Both

guide RNAs target the exon 1 of the SPTLC2 gene (Figure 6-figure supplement 1). The DNA sequence of

the guide RNA (gRNA) was synthesized and cloned into the pX459 CRISPR/Cas9-Puro vector to construct

knockout plasmids px459-SPTLC2-gRNA1 and px459-SPTLC2-gRNA2.

Cells were seeded at appropriate densities in 35-mm dishes and grown for 24 h before transfection with the

plasmids. 48 h after transfection, cells were treated with 2 μg/mL Puromycin (Sangon Biotech, A610593)

for an additional three days. Single cells were then obtained by serial dilution and cultured for expansion.

Genomic DNA was extracted from 18 single-cell expansion cultures and primers 5'-

GCCACCGCCTACAGAGCCTGC-3' / 5'-CCGGGAGTAAGACCTCCAGGCGC-3' were used to amplify

the exon 1 sequence of SPTLC2 by PCR. Two cell lines with deletion mutations were identified by

sequencing (Figure 6-figure supplement 1).

Statistical analysis

Three or more biological replicates were analyzed unless explicitly stated. Statistical significance was

indicated as $p < 0.05$ unless explicitly stated. All graphs show mean and error bars representing standard

error of the deviation (S.D.), using Prism 8.0.0 software (GraphPad Software).

Acknowledgments

We would like to thank Professor Charles Boone (University of Toronto) for providing SGA-V2 and ts-V5

strain collections, as well as the Moby plasmids. We also thank the Centre for Cellular Imaging at University

of Gothenburg for microscopy support, and Center for Large-scale cell-based screening at University of

Gothenburg for assistance with SGA analysis.

Additional information

Funding

Funder	Grant reference number	Author
National Natural Science Foundation of China	32000387	Xiuling Cao
Zhejiang Provincial Natural Science Foundation of China	LY23C060001	Xuejiao Jin
Scientific Research Foundation of Zhejiang A&F University	2021LFR053	Xuejiao Jin

Fellowship from China Scholarship Council and KU Leuven	grant C14/17/063	Ju Zheng
Swedish Cancer Fund (Cancerfonden)	19 0069	Beidong Liu
Swedish Research Council (Vetenskapsrådet)	VR 2019-03604	Beidong Liu
Guangdong Basic and Applied Basic Research Foundation	2023A1515010483	Lihua Chen
FWO-Vlaanderen	G0C7222N	Joris Winderickx
KU Leuven	C14/21/095	Joris Winderickx

The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

Author contributions

Conceptualization: Beidong Liu; Methodology: Xiang Wu, Lei Zhao., Ju Zheng, Xinxin Hao; Investigation:

Xiuling Cao, Lei Zhao, Xuejiao Jin, Lihua Chen, Beidong Liu; Visualization: Xiuling Cao, Xiang Wu, Lei

Zhao, Lihua Chen. Supervision: Xiuling Cao, Shenkui Liu, Lihua Chen, Beidong Liu; Writing—original

draft: Xiuling Cao, Lei Zhao, Lihua Chen; Writing—review & editing: Xiuling Cao, Joris Winderickx, Lihua

Chen, Beidong Liu.

Additional files

Supplementary files

Supplementary file 1. Mutants with manually confirmed increased class 3 synphilin-1 aggregates.

Supplementary file 2. Distribution of GO terms among 84 candidate hits with significantly increased Class

3 aggregates via ClueGO analysis.

Supplementary file 3. Plasmids used in this study.

Supplementary file 4. Strains used in this study.

Data availability

All data generated or analyzed during this study are included in the manuscript and supplementary

information. No datasets have been submitted to public repositories.

References

Abramov, AY, Berezhnov, AV, Fedotova, EI, Zinchenko, VP, and Dolgacheva, LP. 2017. Interaction of misfolded
proteins and mitochondria in neurodegenerative disorders. *Biochemical Society Transactions* **45**: 1025-1033.
https://doi.org/10.1042/bst20170024

Amorim, JA, Coppotelli, G, Rolo, AP, Palmeira, CM, Ross, JM, and Sinclair, DA. 2022. Mitochondrial and
metabolic dysfunction in ageing and age-related diseases. *Nature Reviews Endocrinology* **18**: 243-258.
https://doi.org/10.1038/s41574-021-00626-7

Arrasate, M, Mitra, S, Schweitzer, ES, Segal, MR, and Finkbeiner, S. 2004. Inclusion body formation reduces
levels of mutant huntingtin and the risk of neuronal death. *Nature* **431**: 805-810.
https://doi.org/10.1038/nature02998

Bagola, K, and Sommer, T. 2008. Protein quality control: on IPODs and other JUNQ. *Current Biology* **18**: R1019-
R1021. https://doi.org/10.1016/j.cub.2008.09.036

Bäuerlein, FJB, Saha, I, Mishra, A, Kalemanov, M, Martínez-Sánchez, A, Klein, R, Dudanova, I, Hipp, MS, Hartl,
FU, Baumeister, W, and Fernández-Busnadiego, R. 2017. In situ architecture andcellular interactions of PolyQ
inclusions. *Cell* **171**: 179-187.e110. https://doi.org/10.1016/j.cell.2017.08.009

Bento-Oliveira, A, Santos, FC, Marquês, JT, Paulo, PMR, Korte, T, Herrmann, A, Marinho, HS, and de Almeida,
RFM. 2020. Yeast sphingolipid-enriched domains and membrane compartments in the absence of
mannosyldiinositolphosphorylceramide. *Biomolecules* **10**: 871. https://doi.org/10.3390/biom10060871

Bindea, G, Mlecnik, B, Hackl, H, Charoentong, P, Tosolini, M, Kirilovsky, A, Fridman, WH, Pagès, F, Trajanoski,
Z, and Galon, J. 2009. ClueGO: a Cytoscape plug-in to decipher functionally grouped gene ontology and pathway
annotation networks. *Bioinformatics* **25**: 1091-1093. https://doi.org/10.1093/bioinformatics/btp101

Buchberger, A, Bukau, B, and Sommer, T. 2010. Protein quality control in the cytosol and the endoplasmic
reticulum: brothers in arms. *Molecular Cell* **40**: 238-252. https://doi.org/10.1016/j.molcel.2010.10.001

Busciglio, J, Pelsman, A, Wong, C, Pigino, G, Yuan, M, Mori, H, and Yankner, BA. 2002. Altered metabolism of
the amyloid β precursor protein is associated with mitochondrial dysfunction in Down's Syndrome. *Neuron* **33**:
677-688. https://doi.org/10.1016/S0896-6273(02)00604-9

Büttner, S, Delay, C, Franssens, V, Bammens, T, Ruli, D, Zaunschirm, S, de Oliveira, RM, Outeiro, TF, Madeo, F,
Buée, L, et al. 2010. Synphilin-1 enhances α-synuclein aggregation in yeast and contributes to cellular stress and
cell death in a Sir2-dependent manner. *PLoS One* **5**: e13700. https://doi.org/10.1371/journal.pone.0013700

Casadei, N, Pöhler, AM, Tomás-Zapico, C, Torres-Peraza, J, Schwedhelm, I, Witz, A, Zamolo, I, De Heer, R,
Spruijt, B, Noldus, LP, et al. 2014. Overexpression of synphilin-1 promotes clearance of soluble and misfolded
alpha-synuclein without restoring the motor phenotype in aged A30P transgenic mice. *Human Molecular Genetics*
23: 767-781. https://doi.org/10.1093/hmg/ddt467

Chen, B, Retzlaff, M, Roos, T, and Frydman, J. 2011. Cellular strategies of protein quality control. *Cold Spring*
Harbor perspectives in biology **3**: a004374. https://doi.org/10.1101/cshperspect.a004374

Cutler, RG, Pedersen, WA, Camandola, S, Rothstein, JD, and Mattson, MP. 2002. Evidence that accumulation of
ceramides and cholesterol esters mediates oxidative stress–induced death of motor neurons in amyotrophic lateral

sclerosis. *Annals of Neurology* **52**: 448-457. https://doi.org/10.1002/ana.10312

Douglas, PM, Treusch, S, Ren, H-Y, Halfmann, R, Duennwald, ML, Lindquist, S, and Cyr, DM. 2008. Chaperone-
dependent amyloid assembly protects cells from prion toxicity. *Proceedings of the National Academy of Sciences,*
USA **105**: 7206-7211. https://doi.org/10.1073/pnas.0802593105

France-Lanord, V, Brugg, B, Michel, PP, Agid, Y, and Ruberg, M. 1997. Mitochondrial free radical signal in
ceramide-dependent apoptosis: a putative mechanism for neuronal death in Parkinson's disease. *Journal of*
Neurochemistry **69**: 1612-1621. https://doi.org/10.1046/j.1471-4159.1997.69041612.x

Frydman, J. 2001. Folding of newly translated proteins in vivo: the role of molecular chaperones. *Annual Review*
of Biochemistry **70**: 603-647. https://doi.org/10.1146/annurev.biochem.70.1.603

Gong, H, Romanova, NV, Allen, KD, Chandramowlishwaran, P, Gokhale, K, Newnam, GP, Mieczkowski, P,
Sherman, MY, and Chernoff, YO. 2012. Polyglutamine toxicity is controlled by prion composition and gene
dosage in yeast. *PLoS Genetics* **8**: e1002634. https://doi.org/10.1371/journal.pgen.1002634

Guo, Q, Lehmer, C, Martínez-Sánchez, A, Rudack, T, Beck, F, Hartmann, H, Pérez-Berlanga, M, Frottin, F, Hipp,
MS, Hartl, FU, et al. 2018. In situ structure of neuronal C9orf72 Poly-GA aggregates reveals proteasome
recruitment. *Cell* **172**: 696-705.e612. https://doi.org/10.1016/j.cell.2017.12.030

Haelterman, NA, Yoon, WH, Sandoval, H, Jaiswal, M, Shulman, JM, and Bellen, HJ. 2014. A mitocentric view
of Parkinson's disease. *Annual Review of Neuroscience* **37**: 137-159. https://doi.org/10.1146/annurev-neuro-
071013-014317

Hannun, YA, and Obeid, LM. 2008. Principles of bioactive lipid signalling: lessons from sphingolipids. *Nat Rev*
Mol Cell Biol **9**: 139-150. https://doi.org/10.1038/nrm2329

Hannun, YA, and Obeid, LM. 2018. Sphingolipids and their metabolism in physiology and disease. *Nature*
Reviews Molecular Cell Biology **19**: 175-191. https://doi.org/10.1038/nrm.2017.107

Hartl, FU, and Hayer-Hartl, M. 2009. Converging concepts of protein folding in vitro and in vivo. *Nature*
Structural & Molecular Biology **16**: 574. https://doi.org/10.1038/nsmb.1591

Hashimoto, M, Rockenstein, E, Crews, L, and Masliah, E. 2003. Role of protein aggregation in mitochondrial
dysfunction and neurodegeneration in Alzheimer's and Parkinson's diseases. *Neuromolecular Medicine* **4**: 21-36.
https://doi.org/10.1385/nmm:4:1-2:21

Ho, CH, Magtanong, L, Barker, SL, Gresham, D, Nishimura, S, Natarajan, P, Koh, JLY, Porter, J, Gray, CA,
Andersen, RJ, et al. 2009. A molecular barcoded yeast ORF library enables mode-of-action analysis of bioactive
compounds. *Nature Biotechnology* **27**: 369-377. https://doi.org/10.1038/nbt.1534

Hsu, LJ, Sagara, Y, Arroyo, A, Rockenstein, E, Sisk, A, Mallory, M, Wong, J, Takenouchi, T, Hashimoto, M, and
Masliah, E. 2000. α-Synuclein promotes mitochondrial deficit and oxidative stress. *The American Journal of*
Pathology **157**: 401-410. https://doi.org/10.1016/S0002-9440(10)64553-1

Huh, WK, Falvo, JV, Gerke, LC, Carroll, AS, Howson, RW, Weissman, JS, and O'Shea, EK. 2003. Global analysis
of protein localization in budding yeast. *Nature* **425**: 686-691. https://doi.org/10.1038/nature02026

Jin, HG, Yamashita, H, Nakamura, T, Fukuba, H, Takahashi, T, Hiji, M, Kohriyama, T, and Matsumoto, M. 2008.
Synphilin-1 transgenic mice exhibit mild motor impairments. *Neuroscience Letters* **445**: 12-17.

https://doi.org/10.1016/j.neulet.2008.08.073

Kaganovich, D, Kopito, R, and Frydman, J. 2008. Misfolded proteins partition between two distinct quality
control compartments. *Nature* **454**: 1088-1095. https://doi.org/10.1038/nature07195

Kitamura, A, Kubota, H, Pack, CG, Matsumoto, G, Hirayama, S, Takahashi, Y, Kimura, H, Kinjo, M, Morimoto,
RI, and Nagata, K. 2006. Cytosolic chaperonin prevents polyglutamine toxicity with altering the aggregation state.
Nature Cell Biology **8**: 1163-1170. https://doi.org/10.1038/ncb1478

Kopito, RR. 2000. Aggresomes, inclusion bodies and protein aggregation. *Trends in Cell Biology* **10**: 524-530.
https://doi.org/10.1016/S0962-8924(00)01852-3

Krenz, A, Falkenburger, BH, Gerhardt, E, Drinkut, A, and Schulz, JB. 2009. Aggregate formation and toxicity by
wild-type and R621C synphilin-1 in the nigrostriatal system of mice using adenoviral vectors. *Journal of*
Neurochemistry **108**: 139-146. https://doi.org/10.1111/j.1471-4159.2008.05755.x

Li, S, and Kim, H-E. 2022. Implications of sphingolipids on aging and age-related diseases. *Frontiers in aging* **2**:
797320. https://doi.org/10.3389/fragi.2021.797320

Lim, J, and Yue, Z. 2015. Neuronal aggregates: formation, clearance, and spreading. *Developmental Cell* **32**: 491-
501. https://doi.org/10.1016/j.devcel.2015.02.002

Lima, T, Li, TY, Mottis, A, and Auwerx, J. 2022. Pleiotropic effects of mitochondria in aging. *Nature Aging* **2**:
199-213. https://doi.org/10.1038/s43587-022-00191-2

Lin, G, Wang, L, Marcogliese, PC, and Bellen, HJ. 2019. Sphingolipids in the pathogenesis of Parkinson's disease
and   Parkinsonism.   *Trends   in   Endocrinology   and   Metabolism*   **30**:   106-117.
https://doi.org/10.1016/j.tem.2018.11.003

Liu, J, Li, T, Thomas, JM, Pei, Z, Jiang, H, Engelender, S, Ross, CA, and Smith, WW. 2016. Synphilin-1 attenuates
mutant LRRK2-induced neurodegeneration in Parkinson's disease models. *Human Molecular Genetics* **25**: 672-
680. https://doi.org/10.1093/hmg/ddv504

Liu, QaF, Benjamin and Unruh, Jay R. and Guo, Fengli and Si, Kausik and Li, Rong and Zhou, Chuankai. 2021.
Mitochondrial origin of cytosolic protein aggregation. *SSRN* **37**: 3808300. https://doi.org/10.2139/ssrn.3808300

Mielke, MM, and Lyketsos, CG. 2010. Alterations of the sphingolipid pathway in Alzheimer's disease: new
biomarkers and treatment targets? *Neuromolecular Medicine* **12**: 331-340. https://doi.org/10.1007/s12017-010-
8121-y

Miller, SB, Mogk, A, and Bukau, B. 2015. Spatially organized aggregation of misfolded proteins as cellular stress
defense strategy. *Journal of Molecular Biology* **427**: 1564-1574. https://doi.org/10.1016/j.jmb.2015.02.006

Moldavski, O, Amen, T, Levin-Zaidman, S, Eisenstein, M, Rogachev, I, Brandis, A, Kaganovich, D, and
Schuldiner, M. 2015. Lipid droplets are essential for efficient clearance of cytosolic inclusion bodies.
Developmental Cell **33**: 603-610. https://doi.org/10.1016/j.devcel.2015.04.015

Nuber, S, Franck, T, Wolburg, H, Schumann, U, Casadei, N, Fischer, K, Calaminus, C, Pichler, BJ, Chanarat, S,
Teismann, P, et al. 2010. Transgenic overexpression of the alpha-synuclein interacting protein synphilin-1 leads
to   behavioral   and   neuropathological   alterations   in   mice.   *Neurogenetics*   **11**:   107-120.
https://doi.org/10.1007/s10048-009-0212-2

O'Farrell, C, Murphy, DD, Petrucelli, L, Singleton, AB, Hussey, J, Farrer, M, Hardy, J, Dickson, DW, and Cookson, MR. 2001. Transfected synphilin-1 forms cytoplasmic inclusions in HEK293 cells. *Brain Res Mol Brain Res* **97**: 94-102. https://doi.org/10.1016/s0169-328x(01)00292-3

Ogretmen, B. 2018. Sphingolipid metabolism in cancer signalling and therapy. *Nature Reviews: Cancer* **18**: 33-50. https://doi.org/10.1038/nrc.2017.96

Ogrodnik, M, Salmonowicz, H, Brown, R, Turkowska, J, Średniawa, W, Pattabiraman, S, Amen, T, Abraham, A-c, Eichler, N, Lyakhovetsky, R, and Kaganovich, D. 2014. Dynamic JUNQ inclusion bodies are asymmetrically inherited in mammalian cell lines through the asymmetric partitioning of vimentin. *Proceedings of the National Academy of Sciences, USA* **111**: 8049-8054. https://doi.org/10.1073/pnas.1324035111

Olzscha, H, Schermann, SM, Woerner, AC, Pinkert, S, Hecht, MH, Tartaglia, GG, Vendruscolo, M, Hayer-Hartl, M, Hartl, FU, and Vabulas, RM. 2011. Amyloid-like aggregates sequester numerous metastable proteins with essential cellular functions. *Cell* **144**: 67-78. https://doi.org/10.1016/j.cell.2010.11.050

Park, JH, Burgess, JD, Faroqi, AH, DeMeo, NN, Fiesel, FC, Springer, W, Delenclos, M, and McLean, PJ. 2020. Alpha-synuclein-induced mitochondrial dysfunction is mediated via a sirtuin 3-dependent pathway. *Molecular neurodegeneration* **15**: 5. https://doi.org/10.1186/s13024-019-0349-x

Park, S-H, Kukushkin, Y, Gupta, R, Chen, T, Konagai, A, Hipp, Mark S, Hayer-Hartl, M, and Hartl, FU. 2013. PolyQ proteins interfere with nuclear degradation of cytosolic proteins by sequestering the Sis1p chaperone. *Cell* **154**: 134-145. https://doi.org/10.1016/j.cell.2013.06.003

Polito, L, Greco, A, and Seripa, D. 2016. Genetic profile, environmental exposure, and their interaction in Parkinson's Disease. *Parkinson's Disease* **2016**: 6465793. https://doi.org/10.1155/2016/6465793

Ribeiro, CS, Carneiro, K, Ross, CA, Menezes, JRL, and Engelender, S. 2002. Synphilin-1 is developmentally localized to synaptic terminals, and its association with synaptic vesicles is modulated by alpha-synuclein. *Journal of Biological Chemistry* **277**: 23927-23933. https://doi.org/10.1074/jbc.M201115200

Ross, CA, and Poirier, MA. 2004. Protein aggregation and neurodegenerative disease. *Nature Medicine* **10**: S10-17. https://doi.org/10.1038/nm1066

Sacket, SJ, Chung, H-y, Okajima, F, and Im, D-s. 2009. Increase in sphingolipid catabolic enzyme activity during aging. *Acta Pharmacologica Sinica* **30**: 1454-1461. https://doi.org/10.1038/aps.2009.136

Schlarmann, P, Ikeda, A, and Funato, K. 2021. Membrane contact sites in yeast: control hubs of sphingolipid homeostasis. *Membranes* **11**: 971. https://doi.org/10.3390/membranes11120971

Shishido, T, Nagano, Y, Araki, M, Kurashige, T, Obayashi, H, Nakamura, T, Takahashi, T, Matsumoto, M, and Maruyama, H. 2019. Synphilin-1 has neuroprotective effects on MPP(+)-induced Parkinson's disease model cells by inhibiting ROS production and apoptosis. *Neuroscience Letters* **690**: 145-150. https://doi.org/10.1016/j.neulet.2018.10.020

Singh, P, and Li, R. 2018. Emerging roles for sphingolipids in cellular aging. *Current Genetics* **64**: 761-767. https://doi.org/10.1007/s00294-017-0799-z

Smeal, T, Claus, J, Kennedy, B, Cole, F, and Guarente, L. 1996. Loss of transcriptional silencing causes sterility in old mother cells of *S. cerevisiae*. *Cell* **84**: 633-642. https://doi.org/10.1016/s0092-8674(00)81038-7

Smith, WW, Liu, Z, Liang, Y, Masuda, N, Swing, DA, Jenkins, NA, Copeland, NG, Troncoso, JC, Pletnikov, M,
Dawson, TM, et al. 2010. Synphilin-1 attenuates neuronal degeneration in the A53T alpha-synuclein transgenic
mouse model. *Human Molecular Genetics* **19**: 2087-2098. https://doi.org/10.1093/hmg/ddq086

Song, J, Yang, Q, Yang, J, Larsson, L, Hao, X, Zhu, X, Malmgren-Hill, S, Cvijovic, M, Fernandez-Rodriguez, J,
Grantham, J, et al. 2014. Essential genetic interactors of SIR2 required for spatial sequestration and asymmetrical
inheritance of protein aggregates. *PLoS Genetics* **10**: e1004539. https://doi.org/10.1371/journal.pgen.1004539

Tafesse, FG, Rashidfarrokhi, A, Schmidt, FI, Freinkman, E, Dougan, S, Dougan, M, Esteban, A, Maruyama, T,
Strijbis, K, and Ploegh, HL. 2015. Disruption of sphingolipid biosynthesis blocks phagocytosis of *Candida*
albicans. *PLoS Pathogens* **11**: e1005188. https://doi.org/10.1371/journal.ppat.1005188

Takahashi, T, Yamashita, H, Nagano, Y, Nakamura, T, Kohriyama, T, and Matsumoto, M. 2006. Interactions of
Synphilin-1   with   phospholipids   and   lipid   membranes.   *FEBS   Letters*   **580**:   4479-4484.
https://doi.org/10.1016/j.febslet.2006.07.019

Tanaka, M, Kim, YM, Lee, G, Junn, E, Iwatsubo, T, and Mouradian, MM. 2004. Aggresomes formed by alpha-
synuclein   and   synphilin-1   are   cytoprotective.   *Journal   of   Biological   Chemistry*   **279**:   4625-4631.
https://doi.org/10.1074/jbc.M310994200

Tong, AH, Evangelista, M, Parsons, AB, Xu, H, Bader, GD, Pagé, N, Robinson, M, Raghibizadeh, S, Hogue, CW,
Bussey, H, et al. 2001. Systematic genetic analysis with ordered arrays of yeast deletion mutants. *Science* **294**:
2364-2368. https://doi.org/10.1126/science.1065810

Tong, AH, Lesage, G, Bader, GD, Ding, H, Xu, H, Xin, X, Young, J, Berriz, GF, Brost, RL, Chang, M, et al. 2004.
Global   mapping   of   the   yeast   genetic   interaction   network.   *Science*   **303**:   808-813.
https://doi.org/10.1126/science.1091317

Tong, AHY, and Boone, C (2007). 16 High-throughput strain construction and systematic synthetic lethal
screening in *Saccharomyce scerevisiae*. In Methods in Microbiology, I. Stansfield, and M.J.R. Stark, eds.
(Academic Press), pp. 369-707. https://doi.org/10.1016/S0580-9517(06)36016-3

Trayssac, M, Hannun, YA, and Obeid, LM. 2018. Role of sphingolipids in senescence: implication in aging and
age-related diseases. *Journal of Clinical Investigation* **128**: 2702-2712. https://doi.org/10.1172/jci97949

Wang, P, Deng, J, Dong, J, Liu, J, Bigio, EH, Mesulam, M, Wang, T, Sun, L, Wang, L, Lee, AY, et al. 2019a. TDP-
43 induces mitochondrial damage and activates the mitochondrial unfolded protein response. *PLoS Genetics* **15**:
e1007947. https://doi.org/10.1371/journal.pgen.1007947

Wang, X, Becker, K, Levine, N, Zhang, M, Lieberman, AP, Moore, DJ, and Ma, J. 2019b. Pathogenic alpha-
synuclein aggregates preferentially bind to mitochondria and affect cellular respiration. *Acta neuropathologica*
communications **7**: 41. https://doi.org/10.1186/s40478-019-0696-4

Weisberg, SJ, Lyakhovetsky, R, Werdiger, A-c, Gitler, AD, Soen, Y, and Kaganovich, D. 2012.
Compartmentalization of superoxide dismutase 1 (SOD1G93A) aggregates determines their toxicity. *Proceedings*
of the National Academy of Sciences, USA **109**: 15811-15816. https://doi.org/10.1073/pnas.1205829109

Yang, J, Hao, X, Cao, X, Liu, B, and Nyström, T. 2016. Spatial sequestration and detoxification of Huntingtin by
the ribosome quality control complex. *eLife* **5**: e11792. https://doi.org/10.7554/eLife.11792

Zaarur, N, Meriin, AB, Gabai, VL, and Sherman, MY. 2008. Triggering aggresome formation. Dissecting

aggresome-targeting and aggregation signals in synphilin 1. *Journal of Biological Chemistry* **283**: 27575-27584.

https://doi.org/10.1074/jbc.M802216200

Zhao, L, Yang, Q, Zheng, J, Zhu, X, Hao, X, Song, J, Lebacq, T, Franssens, V, Winderickx, J, Nystrom, T, and

Liu, B. 2016. A genome-wide imaging-based screening to identify genes involved in synphilin-1 inclusion

formation in *Saccharomyces cerevisiae*. *Scientific Reports* **6**: 30134. https://doi.org/10.1038/srep30134

Figures and figure supplements

Figure 1 SY1 aggregates mature into large IBs over time in normal cells, but this process is defective

in aged cells.

(A) The maturation process of SY1 aggregates in yeast cells from scattered small aggregates to large single

IBs over time. DsRed-SY1 was expressed in yeast with the constitutive promoter TPI1, the phenotype of

SY1 aggregates was visualized at 4 h, 16 h and 48 h. Scale bar = 2 μm. **(B)** Cells with different phenotypes

of SY1 IBs are defined as three classes: Class 1 refers to only one IB in a cell, Class 2 refers to two IBs in a

cell, and Class 3 indicates three or more IBs in a cell. Scale bar = 2 μm. **(C)** Time-course study showing that

SY1 aggregates mature after 24 h of cultivation, when the cells have traversed the diauxic shift to enter the

respiratory and stationary growth phase. IBs were examined every 4 h from 4 to 48 h; green bars indicate

percentage of Class 3; white bars indicate percentage of Class 2; red bars indicate percentage of Class 1; the

purple curve indicates the optical density of the cell culture, indicated by the vertical axis on the right. **(D)**

Maturation of SY1 IBs was defective in old yeast cells. Scale bar = 5 μm. **(E)** The proportion of cells with

SY1 IBs was increased in old cells (61%) compared to young cells (27%). **(F)** The number of SY1 IBs in a

cell was increased in young cells (2.64±1.29) versus old cells (4.91±3.32).

Figure 2 Genome-wide high-content imaging screen identifying the serine C-palmitoyltransferase

complex (SPT) as a modulator of SY1 inclusion maturation.

(A) Schematic workflow for identifying mutants that regulate SY1 inclusion maturation. **(B)** The GO

enrichment analysis of genes whose mutants exhibited increased SY1 Class 3 aggregates (ClueGO, cut-off:

P < 0.05). **(C)** The network of the most enriched modules obtained above in 2B. **(D-E)** The increased SY1

Class 3 aggregates in the SPT mutant (*lcb1-4, lcb2-1, tsc3-2*) were rescued by backer-introducing the

corresponding wild-type gene (a MoBY plasmid carries a WT gene, and p5472 is the empty vector control).

Representative images from three independent experiments are shown in D, and quantifications were shown

in E. Scale bar = 2 μm.

Figure 3 Sphingolipids regulate SY1 IB maturation and cytotoxicity in yeast cells.

(A) A schematic diagram of the sphingolipid pathway in yeast cells. Myriocin blocks sphingolipid synthesis

via inhibition of the serine C-palmitoyltransferase (SPT) complex. **(B-C)** Dihydrosphingosine (DHS)

supplementation rescued the deficiency of SY1 IBs maturation in SPT mutants (*lcb1-4, lcb2-1, tsc3-2*).

Representative images from three independent experiments are shown in B, and quantifications are shown

in C. Scale bar = 2 μm. **(D-E)** Blocking intracellular sphingolipid synthesis with myriocin resulted in

defective maturation of SY1 IBs, and exogenous addition of DHS could reverse the effect of myriocin.

Representative images from three independent experiments are shown in D, and quantifications are shown

in E. Scale bar = 2 μm. **(F)** DHS ameliorated the cytotoxicity of SY1 in SPT mutants (*lcb1-4, lcb2-1, tsc3-*

2).

Figure 4 SY1 IBs associate with mitochondria in yeast cells.

(A) SY1 IBs had no obvious association with the plasma membrane (Pma1-GFP), vacuoles (Vph1-GFP), or

ER (Sec63-GFP), but were strongly associated with mitochondria (Tom70-GFP) (See also Figure 4-figure

supplement 1A). Scale bar = 2 μm. **(B)** MitoTracker probe confirmed that all the three classes of IBs were

closely associated with mitochondria (See also Figure 4-figure supplement 1B). Scale bar = 2 μm. **(C)** Super-

resolution microscopy and 3D-SIM construction demonstrated that the SY1 IBs are packed and surrounded

by mitochondria. Scale bar = 1 μm. **(D)** Co-immunoprecipitation assays showing that SY1 physically binds

to several mitochondrial membrane proteins, including Tom70, Tom37 and Tim44. The yeast mitochondrial

membrane proteins Tom70, Tom37, Sam35 and Tim44 were individually C-terminally tagged with GFP, and

the strains were then transformed with a control plasmid (Vector) or a SY1-overproducing plasmid (SY1),

respectively. Cells were grown for 24 h and harvested for subsequent IP experiments. **(E-F)** The association

of SY1 IBs with mitochondria decreases when the physical interaction between SY1 and mitochondrial

membrane proteins is disrupted (*tom70Δ, tom37Δ, tim44*). Representative images from three independent

experiments are shown in E and quantifications are shown in F. Scale bar = 2 μm.

Figure 4-figure supplement 1. Statistics of SY1 IBs surrounded by cellular organelles.

(A) Statistics of the proportion of SY1 IBs surrounded by different organelles, corresponding to Fig 4A. **(B)**

Statistics of the proportion of three different classes of SY1 IBs surrounded by mitochondria, corresponding

to Fig 4B. n.s., not statistically significant.

Figure 4-figure supplement 2. IBs of TDP43, FUS1 and Htt103Qp interact with mitochondria.

Shown are the IBs formed by TDP43, FUS1 and Htt103Qp when fused to GFP and their co-localization with

Tom70, which was genomically tagged with RFP. Scale = 2 μm.

Figure 5 Sphingolipids regulate SY1-induced mitochondrial toxicity in yeast cells.

(A) Overexpression of SY1 decreased mitochondrial membrane potential (indicated by Rho 123 fluor

intensity), increased cellular ROS levels (indicated by DCF mean intensity) and reduced cell viability

(indicated by ATP level) in yeast cells. **(B)** Effects of myriocin treatment on SY1 cytotoxicity in yeast cells

as judged by mitochondrial membrane potential / ROS levels / cell viability). **(C)** Effects of exogenous

addition of DHS on SY1 cytotoxicity in yeast cells based on mitochondrial membrane potential / ROS levels

/ cell viability.

Figure 5-figure supplement 1. The effect of myriocin (A) and DHS (B) treatment on SY1 protein

expression in yeast cells.

SY1 expression was examined by Western blot analysis in samples prepared from cells used in the

experiments corresponding to Fig. 5B and Fig. 5C respectively. n.s., not statistically significant.

Figure 6 SPTLC2 knockout alters the morphology of SY1 IBs and increases the cytotoxicity in

mammalian cells.

(A) The 3D reconstruction showed that SY1 IBs were associated with mitochondria in mammalian HEK293t

cells. GFP-SY1 was expressed in HEK293t cells and mitochondria were probed with MitoTracker. The 3D

structure was constructed using Imaris software. Scale bar = 10 μm. **(B)** SY1 expression decreased

mitochondrial function in HEK293t cells. Mitochondrial membrane potential was monitored by JC-1

staining, ROS levels were determined by DCF mean intensity values, mitochondrial superoxide was

analyzed by MitoSOX mean intensity values. **(C)** The 3D structures of spherical and irregular SY1 IBs, as

well as their interaction with mitochondria was visualized using Imaris software (See also Figure 6-figure

supplement 2). **(D)** SPTLC2 knockout HEK293t cells had impaired management of SY1 IBs, with increased

dispersed irregular IB structures. Scale bar = 20 μm. **(E)** SPTLC2 knockout increased SY1 cytotoxicity. Cell

viability (indicated by luminescence) / ROS (indicated by DCF mean intensity) / mitochondrial membrane

potential (indicated by JC-1 staining) assays were analyzed.

Figure 6-figure supplement 1. SPTLC2-Knockout by CRISPR/Cas9 genome editing.

(A) The experimental flow chart for SPTLC2 knockout in HEK293t cells using CRISPR/Cas9. **(B)** The

sequencing results of the corresponding deletion sites of two obtained knockout cell lines.

Figure 6-figure supplement 2. Statistics of SY1 IB formation.

Statistics of SY1 IB formation in wild type HEK293t and SPTLC2 knockout cells.

Figure 6-figure supplement 3. The expression of SY1 protein.

The expression of SY1 protein in HEK293t SPTLC2 knockout cells. n.s., not statistically significant.

Figure 7 Sphingolipids regulate SY1-induced cytotoxicity and SY1 IB morphology in mammalian cells.

(A) Inhibition of the intracellular sphingolipid synthesis pathway by myriocin enhanced SY1 cytotoxicity

based on ATP content, intracellular ROS levels and mitochondrial membrane potential assays. **(B)** Addition

of exogenous DHS rescued the cytotoxicity caused by SY1 expression. **(C)** Exogenous addition of DHS to

SPTLC2 knockout cells restored the typical spherical SY1 IBs in HEK293t. Statistics of the ratio of cells

with dispersed irregular IBs after exogenous addition of DMSO and DHS to HEK293t SPTLC2 knockout

cells (See also Figure 7-figure supplement 2). Scale bar = 20 μm. **(D)** DHS ameliorated SY1 cytotoxicity by

normalizing cell viability, intracellular ROS levels, and mitochondrial membrane potential.

Figure 7-figure supplement 1. The expression of SY1 protein in HEK293t and SPTLC2 knockout

cells

(A-B) The effect of myriocin (A) and DHS (B) treatment on SY1 expression in HEK293t cells. **(C)** The

effect of DHS treatment on SY1 expression in SPTLC2 knockout cells. n.s., not statistically significant.

Figure 7-figure supplement 2. Statistics of SY1 IB formation.

Statistics of the effect on SY1 IB formation in SPTLC2 knockout cells after exogenous addition of either

DMSO or DHS.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

A

GFP-SY1 MitoTracker Merge Inset

C

Spherical IBs Irregular IBs

GFP-SY1
MitoTracker

B

P = 0.0003

JC-1 Staining (Red/Green Ratio)

Ctrl SY1-FLAG

P = 0.0007

DCF Mean Intens

Ctrl SY1-FLAG

P = 0.0012

MitoSOX Mean Intens

Ctrl SY1-FLAG

D

GFP-SY1 Phalloidin DAPI Merge Inset

Ctrl

SPTLC2-KO

P < 0.0001

% of cells with irregular IBs

Ctrl SPTLC2-KO

E

P = 0.0007

Luminescence

Ctrl SPTLC2-KO

P < 0.0001

DCF Fluore Intens

Ctrl SPTLC2-KO

P < 0.0001

JC-1 Staining (Red/Green Ratio)

Ctrl SPTLC2-KO

Figure 7