Maturation and detoxification of synphilin-1 inclusion bodies regulated by sphingolipids

Xiuling Cao[1,2]†, Xiang Wu[1]†, Lei Zhao[2]†, Ju Zheng[2,3], Xuejiao Jin[1], Xinxin Hao[2], Joris Winderickx[3], Shenkui Liu[1], Lihua Chen[2,4]*, Beidong Liu[1,2]*

[1] State Key Laboratory of Subtropical Silviculture, Zhejiang A&F University, Hangzhou, China. [2] Department of Chemistry and Molecular Biology, University of Gothenburg, Gothenburg, Sweden. [3] Functional Biology, KU Leuven, Leuven, Belgium. [4] Guangzhou Laboratory, Guangzhou, Guangdong, China.

† These authors contributed equally to this work.

* Corresponding author. Email: beidong.liu@cmb.gu.se (B.L.); chen_lihua@gzlab.ac.cn (L.C.)

Abstract

Due to proteostasis stress induced by aging or disease, misfolded proteins can form toxic intermediate species of aggregates and eventually mature into less toxic inclusion bodies (IBs). Here, using a yeast imaging-based screen, we identified 84 potential synphilin-1 (SY1) IB regulators and isolated the conserved sphingolipid metabolic components in the most enriched groups. Furthermore, we show that, in both yeast cells and mammalian cells, SY1 IBs are associated with mitochondria. Pharmacological inhibition of the sphingolipid metabolism pathway or knockout of its key genes results in a delayed IB maturation and increased SY1 cytotoxicity. We postulate that SY1 IB matures by association with the mitochondrion membrane, and that sphingolipids stimulate the maturation via their membrane-modulating function and thereby protecting cells from SY1 cytotoxicity. Our findings identify a conserved cellular component essential for IB maturation and suggest a mechanism by which cells may detoxify the pathogenic protein aggregates through forming mitochondrion-associated IBs.

Introduction

Proteins are vulnerable macromolecules and are often endangered by misfolding and aggregation during the life of the cell due to destabilizing mutations, stress conditions or unique metabolic challenges that occur during aging or disease (Buchberger et al., 2010; Hartl and Hayer-Hartl, 2009). To cope with protein damage and in order to survive, the eukaryotic cell has evolved complex protein quality control systems that monitor and maintain the integrity of its proteome, including refolding, degradation and sequestration (Buchberger *et al.*, 2010; Frydman, 2001; Hartl and Hayer-Hartl, 2009; Lim and Yue, 2015). The refolding of stress-denatured proteins and the degradation of misfolded proteins are assisted by molecular chaperones (Chen et al., 2011; Hartl and Hayer-Hartl, 2009). In addition to refolding and degradation, cells have another cellular strategy to cope with an overload of aberrant proteins or toxic intermediate species of aggregates, namely sequestration into specialized quality control IBs (Bagola and Sommer, 2008; Kaganovich et al., 2008; Kopito, 2000).

It has become increasingly clear that the formation of IBs is like a "triage" center, thus being protective machinery against proteostasis stress during aging or disease. Therefore, the inability to form IBs would poison the intracellular environment and cause cell dysfunction or even death (Lim and Yue, 2015). The toxicity of protein aggregates is more pronounced in neurons partly due to their unique cellular architecture. Indeed, mounting evidence has linked a number of neurodegenerative diseases including Alzheimer's disease (AD), Parkinson's disease (PD), Huntington's disease (HD) and amyotrophic lateral sclerosis (ALS), to the formation of misfolded proteins aggregates containing IBs (Ross and Poirier, 2004). Although IBs have been extensively studied, including their unique characteristics (Kaganovich *et al.*, 2008; Ogrodnik et al., 2014; Weisberg et al., 2012) and the IB clearance machinery (Moldavski et al., 2015), the genes and cellular components that regulate the IB maturation remain poorly understood. In addition, we have little insight into the effects of IBs on cellular membrane-bound structures as IBs have been found to interact with membrane domains (Büttner et al., 2010).

SY1 contributes to inclusion formation and has been implicated in the pathogenesis of PD, an incurable, progressive neurodegenerative disorder. The biological functions of SY1 are not fully understood, and whether SY1 is cytotoxic or protective remains controversial. In aged yeast cells, SY1 expression reduced survival and triggered apoptosis and necrotic cell death in a Sir2 dependent manner (Büttner *et al.*, 2010). However, SY1 exerted neuroprotective effects by inhibiting reactive oxygen species (ROS) production and apoptosis in an *in vitro* PD model (Shishido et al., 2019), and SY1 attenuated mutant LRRK2-induced PD-

like phenotypes and played a neuronal protective role in cultured cells and *Drosophila* (Liu et al., 2016). Transgenic mice expressing SY1 caused pathological inclusions and degeneration of dopaminergic neurons in the substantia nigra (Krenz et al., 2009), and reduced motor skill learning and motor performance (Jin et al., 2008; Nuber et al., 2010). In contrast, SY1 overexpression in A53T human α-synuclein (α-Syn) transgenic mice reduced astrogliosis and neuronal degeneration, resulting in increased lifespan(Smith et al., 2010). In an aged A30P α-Syn transgenic mouse model, Casadei *et al* observed that overexpression of SY1 promoted clearance of soluble and misfolded α-Syn, but did not rescue the motor deficit phenotype (Casadei et al., 2014).

Given that a high-content imaging approach has been successfully conducted for SY1 (Zhao et al., 2016), it was possible to study on a genome-wide scale the cellular components that are involved in IB formation by using yeast as a model. Moreover, since SY1 binds phospholipids, membranes, lipid droplets and synaptic vesicles (Ribeiro et al., 2002; Takahashi et al., 2006) and given that the membrane-binding properties of α-Syn are thought to be relevant for its pathological activity, we speculated that also for SY1 the binding to different types of lipid structures might be related to its cytotoxicity. Here we combined the SGA approach with a high-content imaging-based screen to survey the genetic mutants with altered SY1 aggregation types using a yeast genome-wide mutant library. We identified the conserved sphingolipid metabolic pathway as an essential component for the maturation of SY1 IBs and for the detoxification of SY1 aggregates. In the sphingolipid-rich endomembrane system, SY1 aggregates tightly interact with mitochondria. Disruption of sphingolipid synthesis prevented this association, resulting in small, dispersed aggregates and enhanced SY1-induced cytotoxicity, notably mitochondrial toxicity. Furthermore, pharmacological inhibition of the sphingolipid pathway or knockdown of SPTLC2 altered the morphology of SY1 IBs and increased their cytotoxicity in mammalian cells, and this could be reversed by increasing the level of dihydrosphingosine (DHS), a sphingolipid synthesis intermediate.

Results

SY1 IB management in yeast cells

The formation of SY1 IBs in the yeast (*Saccharomyces cerevisiae*) has been studied by expressing the protein (DsRed-SY1) with a plasmid pYX212 under the control of a constitutive *TPI1* promoter (Zhao *et*

al., 2016). Using a similar approach, we observed different phenotypes of SY1 aggregates in yeast cells.

During the lag phase (4 h), multiple small IBs presented in the cytoplasm but in the exponential growth

phase (16 h), these small aggregates merged into larger IBs, resulting in a decreased number of smaller

aggregates. Subsequently, when the cells entered the stationary phase (48 h), most of the SY1-expressing

cells retained only one to a few large IBs (Figure 1A). Thus, consistent with previously reported data

(Büttner *et al.*, 2010), this time-course study demonstrated a maturation process of SY1 IBs in yeast cells.

To study the details of this process, we defined the IB phenotypes as Class 1 (one aggregate in a cell),

Class 2 (two aggregates in a cell), and Class 3 (three or more aggregates in a cell), respectively (Figure 1B),

as previously reported for mutant huntingtin inclusions (Yang et al., 2016). We then monitored the SY1 IBs

every 4 hours from 4 to 48 hours. Most of the SY1 IB-forming cells exhibited a Class 3 phenotype (96%)

at the 4-hour time-point. During growth, the proportion of Class 3 ratio dropped over time, while the

proportion of Class 1 and Class 2 ratios increased inversely. After 24 hours, the proportions became stable

and Class 1 IBs accounted for approximately 50% when cells reached stationary phase (Figure 1C). All

these observations supported the notion that there is a compartmentalization process of SY1 IBs in the

yeast cell. We therefore hypothesized that the formation and maturation of IBs is a cytoprotective process

that helps sequester more toxic aggregates/oligomers into larger inert deposits (Kitamura et al., 2006).

Dysregulation of the formation of pathogenic protein inclusions is the major hallmark of age-related

neurodegenerative diseases. To investigate whether IBs formed by SY1 expression in old cells have the

same maturation process as that in young cells. We isolated old cells and examined the SY1 aggregates in

old cells (≥5 generations) and young cells (≤1 generation) (Figure 1D). In young cells, about 27% cells had

SY1 inclusions while in the old cells the ratio increased to 61% (Figure 1E). Furthermore, when counted

the number of aggregates in individual cells, we found that there were much more aggregates per cell in

the old cell population (Figure 1F, 4.91±3.32 of old cells vs 2.64±1.29 of young cells). These findings

suggested that there was a dysregulation of SY1 IB maturation in old cells. However, the underlying

mechanism remains unclear. This motivated us to further investigate the genes and cellular components

that regulate SY1 IB maturation.

Genome-wide screening for cellular components regulating the maturation of SY1 inclusions

To dissect the genes and cellular components that mediate the maturation of IBs, a genome-wide, high-

content imaging-based screening approach was performed (Zhao *et al.*, 2016) (Figure 2A). Prior to

performing the screen, we determined the appropriate time-point to assess the phenotypes of SY1 IBs. As

shown in Figure 1A-C, at 16-20 hours of growth (OD_{600} is 3.0 ~ 4.0), approximately 50% SY1-expressing

cells had Class 3 aggregates, and the ratios were stable thereafter. Therefore, we chose the time point 20-

hours for the following screening and other studies. We set the same criteria as in the previous study (Zhao

et al., 2016) to include a particular mutant strain as a potential hit if: 1) the difference in the number of

cells with Class 3 SY1 IBs between wild-type and mutant cells is statistically significant ($P \leq 0.05$, *t*-test)

and 2) the absolute difference of the percentage of wild-type and mutant cells carrying SY1 inclusions is at

least 20%. Potential hits acquired from the screening were then confirmed by manual examination of the

Class 3 inclusion phenotype using a microscopic approach.

The screen was performed three times independently. Based on the above criteria and subsequent manual

confirmation, a list of 84 mutants that showed an increase in cells carrying Class 3 SY1 IBs was obtained

(Supplementary file 1). GO enrichment analysis indicated that these mutants were enriched in the

following modules: serine C-palmitoyltransferase (SPT) complex, SPOTS complex, protein urmylation,

TORC2 complex, palmitoyltransferase complex, and others (Figure 2B and 2C). The complete list of GO

terms can be found in Supplementary file 2. Among them, SPT complex and SPOTS complex, containing

LCB1, LCB2 and TSC3 had the highest enrichment (Table S2, Figure 2B and 2C), corresponding to the

mutants including *lcb1-4*, *lcb2-1*, *lcb2-2* and *tsc3-2* in our screen. We performed gene complementation

experiments on three mutants *lcb1-4*, *lcb2-1* and *tsc3-2*. We performed a time course monitoring of

aggregates in these mutants and found that there was no significant decrease in Class 3 aggregates in the

mutants as in the wild type (Figure 2-figure supplement 1). Restoring the corresponding gene almost

completely rescued the SY1 IB phenotype with significantly reduced Class 3 IBs (Figure 2D and 2E). Thus,

our above screen suggested a link between the highly conserved sphingolipid metabolic pathway and the

maturation of SY1 IBs. A substantial body of evidence has highlighted an important role of sphingolipid

metabolism in several neurodegenerative diseases, including Alzheimer's disease (Mielke and Lyketsos,

2010), amyotrophic lateral sclerosis(Cutler et al., 2002), and Parkinson's disease (France-Lanord et al.,

1997). Therefore, we aimed to elucidate the functional role of sphingolipid metabolism in the regulation of

SY1 IB maturation and cytotoxicity.

Sphingolipids regulate SY1 IB maturation and cytotoxicity in yeast cells

The first step of intracellular sphingolipid de novo synthesis is identical in both yeast and humans, with

palmitoyl-CoA and serine catalyzed by the SPT complex to form 3-Ketosphinganine, which could be

blocked by myriocin (Figure 3A). 3-ketosphinganine is subsequently catalyzed to dihydrosphingosine

(DHS), and dihydroceramide, and further converted into complex sphingolipids. Since downregulation of

SPT in yeast decreases sphingolipid levels (Huang et al., 2012), we first confirmed that exogenous addition

of DHS in SPT mutants (*lcb1-4*, *lcb2-1* and *tsc3-2*) could rescue the phenotype of increased Class 3 SY1

IBs (Figure 3B and 3C). In addition, blockade of intracellular sphingolipid metabolism pathways by

myriocin strongly perturbed SY1 IB maturation process, resulting in elevated Class 3 IBs (Figure 3D and

3E). Again, addition of exogenous DHS could suppress the myriocin-induced Class 3 aggregates (Figure

3D and 3E). Next, we tested whether the SPT genes could regulate SY1 cytotoxicity in yeast cells.

Expressing SY1 did not cause dramatic growth defect in wild-type cells, but led to severe growth defects

in the SPT mutants (Figure 3F, left). Furthermore, this defect could also be rescued by the addition of DHS

(Figure 3F, right). Taken together, these data suggest that sphingolipids may help cells reduce the

cytotoxicity of protein aggregates by regulating the IB maturation process. These findings motivated us to

further unravel the underlying mechanism by which sphingolipids regulate the maturation process and

cytotoxicity of SY1 IBs.

SY1 IBs associate with mitochondria in yeast cells

Sphingolipids are structural molecules of cell membranes that play an important role in maintaining barrier

function and fluidity (Bento-Oliveira et al., 2020; Hannun and Obeid, 2008; Schlarmann et al., 2021). One

of the primary modes of action of sphingolipids is to modulate membrane fluidity, thickness and curvature.

In this context, we asked whether there is a direct association of SY1 IBs with certain membrane-bound

organelles in the yeast cell. To address this question, we examined the location of SY1 IBs (20 h) in cells

and their possible association with membranous organelles visualized by corresponding marker proteins.

Our results showed that SY1 IBs had no obvious association with the plasma membrane (Pma1-GFP),

vacuole (Vph1-GFP), or ER (Sec63-GFP), but were strongly surrounded with mitochondria (Tom70-GFP)

(Figure 4A and Figure 4-figure supplement 1A, about 80%). Since SY1 IBs have three different

morphologies (Class 1-3) and the Class 3 IBs transition to Class 1 during maturation (Figure 1A-C), we

wanted to confirm whether all these different IBs surrounded with mitochondria in a similar way. The

results confirmed that all the three different IBs were equivalently associated with mitochondria (Figure

4B and Figure 4-figure supplement 1B). Furthermore, in myriocin-treated cells, most SY1 IBs were

similarly surrounded by mitochondria (Figure 4-figure supplement 1C). Also, super-resolution microscopy

and 3D-SIM techniques demonstrated that the SY1 IBs were surrounded by mitochondria (Figure 4C).

The next question is whether other pathogenic IBs are also associated with mitochondria. To address this

question, we used the same expression system to overexpress TDP43, FUS1 or Htt103Qp (GFP-labeled) in

yeast cells expressing Tom70-RFP and examined the possible association with mitochondria. Our results

showed that the IBs of these proteins indeed interacted with mitochondria, similar to what we found for the

IBs of SY1 (Figure 4-figure supplement 2). Hence, it appears that the association of IBs with mitochondria

may be a common biological event among these disease proteins. To further study the consequences of this

association and to decipher potential factors that regulate this association, we first investigated whether

SY1 IBs physically associate with mitochondria. We performed four co-immunoprecipitation (Co-IP)

experiments to determine whether SY1 could be detected in pull-downs for the inner mitochondrial

membrane protein Tim44 or the outer mitochondrial membrane proteins Tom70, Tom37, and Sam35. The

SY1 protein was indeed the case in extracts obtained from the Tom70-GFP, Tom37-GFP and Tim44-GFP

strains (Figure 4D). Next, we performed a microscopic analysis and found that the surrounding state of

SY1 IBs with mitochondria was significantly hampered in *tom70Δ, tom37Δ* and *tim44* mutants (Figure 4E

and 4F), as well as an increased proportion of Class 3 aggregates in the mutants (Figure 4-figure

supplement 1D). Furthermore, we expressed SY1 protein in BY4741 rho0 strain and found that the

maturation and mitochondrial surrounding state of SY1 IB was not affected by mitochondrial activity

(Figure 4-figure supplement 1E and 1F). Collectively, these results confirm that SY1 IBs are in physical

contact with mitochondrial membrane components Tom70, Tom37 and Tim44, which is consistent with the

fact that SY1 is a membrane lipid raft interacting protein (Büttner *et al.*, 2010; O'Farrell et al., 2001;

Takahashi *et al.*, 2006).

Sphingolipids regulate SY1-induced mitochondrial toxicity in yeast cells

Several lines of evidence have supported the notion that potentially toxic proteins such as α-Syn and TDP-

43 can associate with mitochondria and damage the mitochondrial membrane, leading to cytotoxicity

(Bäuerlein et al., 2017; Busciglio et al., 2002; Hashimoto et al., 2003; Hsu et al., 2000; Park et al., 2020;

Wang et al., 2019a; Wang et al., 2019b). For example, pathogenic α-Syn aggregates preferentially bind to

mitochondria, leading to decreased mitochondrial biogenesis (Park *et al.*, 2020) and impaired cellular

respiration (Wang *et al.*, 2019b). To explore whether the interaction between SY1 IBs and mitochondrial

membranes could impair mitochondrial function, we measured mitochondrial membrane potential and cell

viability represented by ATP levels in SY1-expressing cells. The results showed that mitochondrial

membrane potential and cell viability of SY1-expressing cells were significantly lower than that of control

cells (Figure 5A). These changes indicate a decrease in mitochondrial function caused by SY1 IBs. Given

that SY1 IBs interact with sphingolipid-enriched membrane raft domains and that sphingolipid metabolism

regulates the maturation of SY1 IBs, we wondered whether sphingolipids play a role in the observed SY1-

induced mitochondrial dysfunction. To address this question, we first blocked the sphingolipid synthesis by

myriocin in the SY1-expressing yeast cells and measured mitochondrial membrane potential and cell

viability. We found that myriocin treatment aggravated SY-1-induced cytotoxicity, as indicated by

decreased mitochondrial membrane potential and cell viability (Figure 5B), without altering SY1 protein

expression (Figure 5-figure supplement 1A). Second, we added exogenous DHS to SY1-expressing cells

and observed ameliorated SY-1-induced cytotoxicity as indicated by increased mitochondrial membrane

potential and cell viability (Figure 5C), without altering SY1 protein expression (Figure 5-figure

supplement 1B). Notably, our assay showed that the knocking out of TOM70 or TOM37 reduced the

mitochondrial toxicity caused by SY1 expression ((Figure 5-figure supplement 2). Collectively, these

results showed that sphingolipids regulated the mitochondrial toxicity induced by SY1 IBs, and suggested

that sphingolipids might detoxify SY1 IBs through mitochondria.

SPTLC2 knockout alters the morphology of SY1 IBs and increases the cytotoxicity in mammalian

cells

We then wondered whether the observed association with mitochondria and the cytotoxicity of SY1 IBs

was conserved in mammalian cells. Similarly, SY1 forms IBs in mammalian cells, which are known as

aggresomes (Zaarur et al., 2008). First, we expressed SY1 in HEK293t cells to study the association of

SY1 IBs with mitochondria. A 3D structure between SY1 IB and mitochondria was constructed by

fluorescence 3D reconstruction and it confirmed that in these mammalian cells the SY1 IBs are in close

contact with mitochondria, although the morphology of the mitochondrion is different (Figure 6A).

Furthermore, the mitochondrial membrane potential was significantly decreased in SY1-expressing cells,

while the levels of mitochondrial superoxide were significantly increased (Figure 6B), indicating impaired

mitochondrial function by SY1 IBs. As such, our data suggest that in yeast and mammalian cells the

formation and processing of SY1 IBs occur in a similar manner, leading to mitochondrial dysfunction

through influencing the mitochondrial membranes.

To further confirm this, we analyzed whether sphingolipids are indeed involved in the formation and

cytotoxicity of SY1 IBs in mammalian cells as well. To this end, we used the CRISPR/Cas9 approach to

knock out SPTLC2, the ortholog of yeast LCB2 and one of the subunits of the mammalian SPT complex

(Figure 6-figure supplement 1). Notably, knockout of SPTLC2 significantly reduced the accumulation of

intracellular sphingolipids, but had little effect on normal cell division and morphology (Tafesse et al.,

2015). Microscopic analysis showed that the morphology of SY1 IBs formed in mammalian cells can be

divided into two forms, one compact and spherical and the other more dispersed with an irregular structure

as if many small IBs stuck together, without clear boundaries or uniform centers (Figure 6C). In wild-type

cells, the SY1 IBs were predominantly spherical. However, in SPTLC2 knockout cells, the morphology of

SY1 IBs was significantly altered, with an increased number of dispersed irregular IBs (Figure 6D) and an

increased number of SY1 IBs per cell (Figure 6-figure supplement 2A). This is consistent with the

phenotypes observed in SPT mutant yeast cells (Figure 3B). We also performed SPTLC2 gene

complementation experiments in knockout cells, which reduced the number of cells forming IBs and the

percentage of dispersed irregular IBs (Figure 6-figure supplement 2B). Furthermore, mitochondrial

function, as judged from mitochondrial membrane potential and cell viability, was decreased in the

SPTLC2-deficient cells (Figure 6E), while the SY1 protein expression was not altered (Figure 6-figure

supplement 3). Taken together, these results further substantiate that SY1 IBs are processed and managed

similarly in mammalian cells and in yeast cells and that the sphingolipid pathway plays a role in the

cytotoxicity of SY1 IBs in mammalian cells as well.

Sphingolipids regulate SY1-induced cytotoxicity and SY1 IB morphology in mammalian cells

Because the initial steps of sphingolipid biosynthesis are conserved from yeast to humans, myriocin also

inhibits the de novo synthesis of intracellular sphingolipids in mammalian cells (Ogretmen, 2018). We

added myriocin to SY1-expressing HEK293t cells, which significantly increased the number of SY1 IBs as

the SPTLC2-deficient cells (Figure 7-figure supplement 1A), and then examined the corresponding

changes in cytotoxicity. The results showed that inhibition of the intracellular sphingolipid synthesis

pathway by myriocin enhanced SY1 cytotoxicity based on mitochondrial membrane potential and ATP

content assays (Figure 7A). Furthermore, the addition of exogenous DHS rescued the cytotoxicity caused

by SY1 expression (Figure 7B) without altering SY1 protein expression (Figure 7-figure supplement 1B

and C).

We further examined the cytotoxicity of SY1 in SPTLC2 knockout cells with or without DHS. The results

showed that DHS restored the morphology of IBs (Figure 7C), as well as the number of SY1 IBs formed in

SPTLC2 knockout cells (Figure 7-figure supplement 2). Accordingly, DHS also ameliorated the SY1-

instigated cytotoxicity by normalizing mitochondrial membrane potential and cell viability (Figure 7D),

without affecting SY1 expression (Figure 7-figure supplement 1D). These results reinforce the notion that

sphingolipid metabolism plays an important role in the formation of SY1 IBs and SY1-induced

cytotoxicity.

Discussion

In this study, we used a genome-wide high-content imaging-based screening approach to identify

regulators of SY1 IBs formation. Among the 5500 mutants, 84 mutants exhibited increased Class 3 SY1

inclusions. GO enrichment analysis revealed the SPT complex as the most enriched functional group.

Subsequent studies revealed that the sphingolipid pathway plays an important role in the processing of

SY1 IBs and the detoxification of SY1 by mitochondria in yeast cells. Corresponding experiments in

mammalian cells recapitulated the role of sphingolipids in the detoxification of SY1-induced

mitochondrial toxicity. Taken together, our results suggest a role for sphingolipids in controlling the

cytotoxicity of pathologically aggregated proteins in cells.

IBs management and their cytotoxicity

The formation and accumulation of abnormal intracellular protein aggregates, including the pathological

aggregates observed in neurodegenerative diseases, are generally considered to be a double-edged sword.

On the one hand, their ability to co-aggregate and sequester basic cellular components, including

molecular chaperones, causes cytotoxicity (Gong et al., 2012; Olzscha et al., 2011; Park et al., 2013). On

the other hand, highly toxic, soluble oligomeric substances form protein aggregates can also protect cells

(Arrasate et al., 2004; Douglas et al., 2008; Tanaka et al., 2004). Many studies have shown that converting

small aggregates/oligomers into IBs can reduce the cytotoxicity of various aggregated proteins (Miller et

al., 2015). In line with a previous report (Büttner *et al.*, 2010), our current study demonstrated that the

merging of smaller foci into IBs also occurred when SY1 was expressed in yeast and mammalian cells.

Association of mitochondrial with protein aggregates

The formation of large intracellular IBs leads to severe impairment of membrane function of IB-bound organelles, thereby affecting organelle function. This has been confirmed in several experiments, such as the disruptive effect of Huntingtin polyQ IBs on ER function (Bäuerlein *et al.*, 2017) and the effect of neuronal *C9orf72* poly-GA aggregates on the Proteasome (Guo et al., 2018). Several lines of evidence implicate mitochondrial dysfunction as a primary cause of PD. Environmental toxins such as mitochondrial complex inhibitors rotenone and 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine induce dopaminergic neuronal death (Polito et al., 2016). In addition, mitochondrial complex I deficiency and mitochondrial electron transport chain defects have been reported in the nigra of PD patients (Haelterman et al., 2014). However, it is unclear how these protein aggregates affect the mitochondria. As previously reported, small foci formed by SY1 started to merge into larger deposits when cells approached the diauxic shift to reprogram metabolism from fermentation to respiration (Büttner *et al.*, 2010), which was confirmed by our present study (Figure 1A). In addition, we show that SY1 physically interacts with mitochondrial membrane proteins and that larger SY1 deposits are surrounded by mitochondrial structures in both yeast and mammalian cells (Figure 4A-D and Figure 6A). However, it is unknown why these IBs preferentially interact with mitochondria. This preference may be partly explained by a recent study in which the authors showed that the mitochondrial import receptor Tom70 could nucleate stress-induced aggregation of cytosolic proteins (Liu, 2021). In addition, impaired mitochondrial protein import could form cytosolic aggregates and thereby stimulate the aggregation of other proteins, including many pathogenic proteins such as α-Syn and amyloid β (Nowicka et al., 2021). In this study, however, our current study shows a fine management of aggregated proteins in eukaryotic cells–protein IBs are tightly associated with mitochondria once formed and continue to mature onto mitochondria.

In addition to SY1, there is a similar association of IBs and mitochondria with other aggregation-prone proteins, including TDP43, FUS1 and Htt103Qp (Figure 4-figure supplement 2), suggesting that mitochondrion-contact may be a common feature for the maturation of pathogenic protein IBs. Additional experiments are needed to confirm this hypothesis, although an interaction with mitochondria has already been described for the aforementioned and several other aggregation-prone proteins (reviewed in (Abramov et al., 2017)). Furthermore, it would be of great value to illustrate how these IBs are sequestered to mitochondria and how the smaller IBs merge into large IBs with the help of mitochondria.

314 **The role of sphingolipids in SY1 IB maturation and mitochondrial function.**

315 Sphingolipids are structural molecules of cell membranes with important roles in maintaining barrier

316 function and fluidity (Hannun and Obeid, 2008). In addition, bioactive sphingolipids have variety of

317 functions that are involved in almost all major aspects of cell biology, and key sphingolipid metabolites are

318 critical for normal brain development and function (Hannun and Obeid, 2018). Membranes and membrane

319 dynamics are required for these processes and an imbalance of lipids, including ceramide and sphingolipid

320 intermediates, may contribute to mitochondrial dysfunction that could lead to the demise of neurons (Lin et

321 al., 2019). SY1 inclusions co-localize with lipid droplets and endomembranes, and SY1 interacts with

322 cytoplasmic lipid rafts in mammalian cells and yeast (Büttner *et al.*, 2010; Takahashi *et al.*, 2006). The C-

323 terminus of SY1 was found to bind selectively to acidic phospholipids, including phosphatidic acid,

324 phosphatidylserine, and phosphatidylglycerol, but not to naturally charged phospholipids and the amino

325 acid sequence 610-640 was found to represent the major determinant of phospholipid binding (Takahashi

326 *et al.*, 2006). The open question is how sphingolipids contribute to the SY1 IB maturation: do they

327 influence this process directly, or indirectly by affecting other pathways? There was no significant

328 enrichment for protein quality control or UPR-related pathways in our genome-wide screen, so it is

329 unlikely that sphingolipids indirectly cause maturation of IBs by affecting these two pathways. This

330 may require further investigation. Moreover, the relationship between sphingolipids and mitochondrial

331 function during aging is not fully understood. It has long been appreciated that mitochondrial activity and

332 quality decline with age (reviewed in(Amorim et al., 2022; Lima et al., 2022)). On the other hand,

333 sphingolipids play critical roles in the regulation of lifespan in yeast, worms, and flies, and in the

334 regulation of senescence in mammalian cells (Li and Kim, 2022; Singh and Li, 2018; Trayssac et al., 2018).

335 However, the changes in sphingolipid profiles during aging have not been fully characterized. In

336 mammalian cells, there is an accumulation of ceramide and sphingosine during aging (Sacket et al., 2009),

337 but the levels of other species including long-chain base sphingolipids are not reported. In this regard,

338 additional studies will be needed to profile the sphingolipid species in the aged cells and to identify the

339 major sphingolipids that regulate mitochondrial function.

340

341 **Materials and methods**

 Key Resources Table

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Antibody	anti-Flag	Cell Signaling	D6W5B	1:1000
Antibody	anti-GFP	Abcam	ab6556	1:1000
Antibody	anti-SY1	Affinity Biosciences	DF8619	1:1000
Antibody	anti-PGK1	Abcam	ab113687	1:1000
Antibody	anti-GAPDH	Wanleibio	WL01114	1:1000
Antibody	goat anti-rabbit IgG	Abcam	ab6721	1:10000
Chemical compound, drug	myriocin	MCE	HY-N6798	
Chemical compound, drug	dihydrosphingosine (DHS)	Sigma	D3314	
Chemical compound, drug	sulfo-NHS-LC biotin	Thermo Scientific	21335	
Chemical compound, drug	streptavidin-conjugated paramagnetic beads	Thermo Scientific	88817	
Chemical compound, drug	MitoTracker Green	Invitrogen	M7514	
Chemical compound, drug	Nonidet P-40	Beyotime	ST2045	
Chemical compound, drug	anti-GFP beads	chromoTek	GFP-Trap	
Chemical compound, drug	Laemmli buffer	Sigma	S3401	
Chemical compound, drug	beta-mercaptoethanol	Thermo Scientific	21985023	
Chemical compound, drug	Rhodamine123	Beyotime	C2007	
Chemical compound, drug	MitoTracker Deep Red FM	Invitrogen	M22426	
Chemical compound, drug	phalloidin-Atto 565	Sigma	94072	
Commercial assay or kit	MitoSOX	Invitrogen	M36008	
Commercial assay or kit	JC-1	Beyotime	C2006	
Commercial assay or kit	X-tremeGENE HP DNA Transfection Reagent	Roche	6366236001	
Commercial assay or kit	CellTiter-Lumi Luminescent Cell Viability Assay Kit	Beyotime	C0065S	
Commercial assay or kit	BCA Protein Quantification Kit	Beyotime	P0012S	
Cell line (*Homo-sapiens*)	HEK293t	NCACC	SCSP-502	
Cell line (*Homo-sapiens*)	HEK293t SPTLC2 knockout cells	This paper		
Strains, strain background (*S. cerevisiae*)	yeast single gene knockout collection (SGA-V2)	Charlie Boone Lab		
Strains, strain background (*S. cerevisiae*)	essential gene temperature sensitive allele collection (ts-V5)	Charlie Boone Lab		
Strain, strain background (*S. cerevisiae*)	BY4741 (MATa *his3Δ1 leu2Δ0 met15Δ0 ura3Δ0*)	Dharmacon Inc	YSC1048	
Strain, strain background (*S. cerevisiae*)	Y7092 (MATα *can1Δ::STE2pr-his5 lyp1Δ ura3Δ0 leu2Δ0 his3Δ1 met15Δ0*)	Tong and Boone, 2007		
Strains, strain background (*S. cerevisiae*)	Yeast GFP Collection	Invitrogen	95702	
Sequence-based reagent	Guide RNA for SPTLC2 knockout Guide RNA sequences #1:	This paper	Guide RNA	CGAATGGCTGCGTG GCGAAC

Sequence-based reagent	Guide RNA for SPTLC2 knockout Guide RNA sequences #2:	This paper	Guide RNA	AAGTACGGAACGGG TACGTG
Sequence-based reagent	Forward primer for SPTLC2 exon 1 sequence amplification:	This paper	PCR primers	GCCACCGCCTACAG AGCCTGC
Sequence-based reagent	Reverse primer for SPTLC2 exon 1 sequence amplification:	This paper	PCR primers	CCGGGAGTAAGACC TCCAGGCGC
Recombinant DNA reagent	pYX212-Dsred -SY1	Büttner et al., 2010		
Recombinant DNA reagent	pYX212-Dsred	Büttner et al., 2010		
Recombinant DNA reagent	pYX212-Dsred -SY1-His	This paper		
Recombinant DNA reagent	pcDNA3.1-GFP-SY1	This paper		
Recombinant DNA reagent	pcDNA3.1-SY1-Flag	This paper		
Recombinant DNA reagent	px459-SPTLC2-gRNA1	This paper		
Recombinant DNA reagent	px459-SPTLC2-gRNA2	This paper		
Software, algorithm	ImageJ	NIH	RRID:SCR_0 03070	https://imagej.net/image j-wiki-static/
Software, algorithm	Imaris	Bitplane AG	RRID:SCR_0 07370	https://imaris.oxinst.co m
Software, algorithm	ImageXpress software	Molecular Devices	RRID:SCR_0 16654	https://www.moleculard evices.com
Software, algorithm	ClueGO (V2.5.9)	Bindea et al., 2009	RRID:SCR_0 05748	https://apps.cytoscape.o rg/apps/cluego
Software, algorithm	Cytoscape (V3.9.1)	Cytoscape	RRID:SCR_0 03032	https://cytoscape.org/
Software, algorithm	GraphPad Prism	GraphPad Software	RRID:SCR_0 02798	https://www.graphpad.c om/

Plasmids

All plasmids used in this study are listed in Supplementary file 3. The plasmid pYX212-DsRed-SY1-His

was constructed by replacing the URA3 cassette with the His3MX6 cassette by homologous recombination

in yeast. The corresponding plasmids for complementation experiments were derived from the MoBY

library (Ho et al., 2009). The plasmids in this collection were constructed based on the p5472 plasmid with

its native promoter and terminator. To construct plasmid pcDNA3.1-GFP-SY1, the GFP fragment and the

SY1 fragment were inserted into the Not I/Hind III digested pcDNA3.1 (-) vector using Exnase (Vazyme,

C112). For the construction of plasmid pcDNA3.1-SY1-Flag, the PCR-expanded SY1-Flag fragment was

inserted directly into the Not I/Hind III digested pcDNA3.1 (-) vector.

Yeast strains and growth conditions

The yeast strains used in this study are listed in Supplemental file 4. Experiments were mainly performed

in the BY4741 strain (MATa *his3Δ1 leu2Δ0 met15Δ0 ura3Δ0*) and its respective mutants. The *S. cerevisiae*

MATα strain Y7092 (Tong and Boone, 2007) was used for query strain constructions. The yeast single

gene knockout collection (SGA-V2) and the essential gene temperature sensitive allele collection (ts-V5)

were kindly provided by Prof. Charlie Boone, University of Toronto, Canada. Strains expressing Pma1-

GFP, Sec63-GFP, Vph1-GFP, Tom70-GFP, Tom37-GFP, Sam35-GFP, and Tim44-GFP were obtained from

the yeast GFP collection (Huh et al., 2003). Yeast strains were grown in YPD (1% yeast extract, 2%

peptone, and 2% glucose) or appropriate synthetic drop-out medium (0.17% yeast nitrogen base without

amino acids and ammonium sulfate, 0.5% ammonium sulfate and the appropriate synthetic drop-out amino

acid, and 2% glucose). Yeast cells were grown at 30°C or at 22°C in case of temperature sensitive (ts)

alleles.

SGA analysis

The experimental methods are based on a previous report (Zhao *et al.*, 2016), and essentially consist of

transforming the plasmids pYX212-Dsred-SY1 and pYX212-Dsred (control), respectively, into the yeast

query strain Y7092. Subsequently, SGA mating was performed as previously described (Tong et al., 2001;

Tong et al., 2004) to introduce the plasmids pYX212-Dsred-SY1 or pYX212-Dsred into the SGA-V2 and

ts-V5 collections, respectively. All pinning steps for collection handling were performed using a SINGER

ROTOR HDA robot (Singer Instrument, Watchet UK).

For the genome-wide high content imaging screening of SY1 IB morphology, the constructed collections

expressing Dsred-SY1 were pre-cultured in 200 μL SD-Ura liquid medium in 96-well plates with

antibiotics at 30°C (22°C for ts mutants) for 3 days without shaking. Then, each pre-culture sample was

diluted to OD_{600nm} 0.1 to a final volume of 200 μL of SD-Ura medium. After 20 h (24 h for ts mutants) of

incubation with shaking at 30°C (for both single and ts mutants), cells were fixed by adding formaldehyde

to a final concentration of 3.7% for 30 min at room temperature and then washed twice with 1×PBS. For

imaging, appropriate amounts of fixed cells were transferred to new 96-well glass bottom plates (Matri-

plate) with 200 μL of 1×PBS in each well and imaged using ImageXpress MICRO (Molecular Devices

Corporation (MDC), San Jose, USA), an automated cellular imaging system. To automatically quantify

cells with different SY1 aggregation phenotypes, a customized subprogram of the MetaXpress software

(MDC) was used (Zhao *et al.*, 2016). Image preprocessing with shading correction and background

subtraction was first performed; cells and IBs were separated from their background based on differences

in signal intensity, respectively. Then, the number of IBs per cell was extracted by combining the two

masks (cell and inclusion body). The percentages of the three types (class 1 to class 3) of cells were

counted. All mutants that showed statistically significant differences as compared to the wild-type were re-

streaked, individually tested, and further manually analyzed to confirm the phenotypic differences

observed in the screen. At least 300 cells were counted for manual confirmation. For manual confirmation,

cells were grown from a starting OD_{600nm} of 0.1 in 15 ml Falcon tubes for 20 h (24 h for ts mutants) and

fixed and washed as described above. All manual confirmations were verified using a Zeiss Axio Observer

7 microscope (Carl Zeiss, Jena, Germany).

Spot assays

392　For spot assays, the mutants carrying DsRed-SY1 were cultured in 3 mL SD-Ura liquid medium at 22°C

393　for 24 h. The strains were diluted to $OD_{600\,nm} = 0.5$ and then serially diluted tenfold in sterile water in 96-

394　well plates. The dilutions were spotted on plates with or without DHS. Images were captured after

395　incubation at 30°C for 2 days.

Bioinformatic analysis

397　Candidate hits were analyzed for enrichment in the Gene Ontology (GO) using ClueGO (Bindea et al.,

398　2009) in Cytoscape with a cut-off of $P < 0.05$. Bubble and network plots of significantly altered GO terms

399　and genes were generated using OmicStudio tools (https://www.omicstudio.cn/tool) and Cytoscape.

Isolation of SY1-expressing old yeast cells

Old yeast cells overexpressing SY1 were obtained by magnetic sorting (Smeal et al., 1996). The wild-type

BY4741 cells expressing Dsred-SY1 were cultured in 100 mL SD-Ura liquid medium, and then harvested

after growth to 0.5 with an OD_{600nm} of 0.1, washed three times in ice-cold PBS pH7.4, and incubated with

1 mg/mL sulfo-NHS-LC biotin labeling for 20 min at room temperature. Cells were further washed four

times with cold PBS pH 8.0 and grown overnight in 100 mL new fresh medium. Cells were harvested the

next day and washed three times with cold PBS pH 7.4, and incubated with 250 mL streptavidin-

conjugated paramagnetic beads for 2 h at 4°C. Unlabeled cells were removed by magnetic sorting. The

cells were then grown overnight in 100 mL fresh SD-Ura medium. The isolation process was repeated two

more times. At the final isolation, cells were collected from the supernatant of the first wash and

resuspended in PBS to serve as the young cells, while cells labeled with magnetic beads were centrifuged

and resuspended in PBS to serve as old cells. The yeast bud scars were photographed through the DAPI

channel using a Zeiss Axio Observer 7 fluorescence microscope to confirm and calculate the degree of

yeast senescence. Subsequently, the SY1 aggregates were photographed through the Dsred channel to

calculate the type of SY1 aggregates.

Complementation of yeast mutants with MoBY plasmid

416 The mutants expressing pYX212-DsRed-SY1 were transformed with the corresponding MoBY plasmid or

417 empty control vector (p5472). For the morphological observation experiment of SY1 IBs, the cells were

418 cultured in SD-Ura, His liquid medium at 30ºC starting from $OD_{600nm} = 0.1$ for 20 h. The cells were then

419 inspected by a fluorescence microscope and photographed.

Myriocin treatment and DHS supplementation in yeast

421 Myriocin was dissolved in DMSO at a concentration of 10 mM. The stock solution was incubated in a

422 55°C water bath for 15 min prior to use. DHS was prepared in distilled DMSO to 30 mM stock solution.

423 For treatments, cells were grown overnight and diluted in medium to an initial OD_{600nm} of 0.1 with

424 myriocin or DHS at a final concentration of 25 or 20 μM, respectively.

Microscopic analysis of SY1 IBs in yeast

For the microscopic analysis of SY1 IBs in yeast and the co-localization analyses of SY1 IBs with

fluorescently labeled proteins Pma1-GFP, Sec63-GFP, Vph1-GFP, and Tom70-GFP, pYX212-DsRed-SY1

was transformed into corresponding cells and yeast strains were grown overnight and diluted in culture

medium to a starting OD_{600nm} of 0.1. After shaking at 30°C for 20 h, the cells were fixed by adding

formaldehyde. Cell images were captured using Zeiss Axio Observer 7. For mitochondrial staining

analysis, MitoTracker Green staining solution was prepared according to the manufacturer's protocol. After

20 h of growth from an OD_{600nm} of 0.1, 1×10^7 cells were fixed with formaldehyde for 30 min and then

washed twice in 1×PBS (pH 7.4). The cells were resuspended in 1× PBS (pH 7.4) containing 125 nM

MitoTracker probe and incubated for 30 min at room temperature. After one wash with PBS, the cells were

imaged by microscopy.

3D-SIM microscopy

437 3D-SIM microscopy images were obtained as previously reported (Song et al., 2014). Briefly, yeast strain

438 Tom70-GFP expressing pYX212-DsRed-SY1 was grown overnight and diluted in medium to a starting

439 OD_{600nm} of 0.1. After shaking at 30°C for 20 h, cells were fixed with 3.7% formaldehyde and washed twice

with $1\times$ PBS (pH 7.4). The ELYRA PS.1 LSM780 setup from Zeiss (Carl Zeiss, Jena, Germany) was used for 3D-SIM. Images were acquired with a $100\times/1.46$ Plan-Apochromat oil immersion objective. Z-stacks with 100 nm spacing were used to scan the whole yeast in 3D-SIM. Zeiss Zen microscope software was used for acquisition, super-resolution processing and calculation, and 3D reconstruction. The ELYRA system was corrected for chromatic aberration in x-, y- and z-directions using multi-color beads, and all obtained images were inspected and aligned accordingly.

Co-IP for SY1 with mitochondrial membrane proteins in yeast

pYX212-DsRed-SY1 and pYX212-DsRed were transformed into GFP strains (Tom70-GFP, Tom37-GFP, Sam35-GFP, and Tim44-GFP) respectively. The strains were pre-cultured and diluted to 100 mL medium with OD_{600nm}=0.1. Cells were harvested when OD_{600nm} reached 1.0 by centrifugation at 4000 rpm for 5 min at 4°C and washed twice with 20 mL ice-cold ddH_2O water. Cells were then washed with 1 mL ice-cold IP buffer (115 mM KCl, 5 mM NaCl, 2 mM $MgCl_2$, 1 mM EDTA, 20 mM HEPES/KOH, pH 7.5), and transferred to 1.5 mL Eppendorf tubes. Cell pellets were resuspended in 250 µL IP buffer containing protease inhibitors, DTT, and glass beads and disrupted using a FastPrep-24 device (MP Biomedicals). Cells were further lysed on ice for 20 min with additional IP buffer containing 0.5% Nonidet P-40. The lysates were centrifuged at 16,000 g for 20 min at 4°C, and 700-800 µL of supernatant was transferred to a new tube. For the IP assay, 20 µL anti-GFP beads were resuspended in 100 µL IP buffer and mixed with the supernatant. The mixture was incubated for 1-1.5 h at 4°C on an overhead rotator, centrifuged at 2000 rpm for 1 min, and washed 4 times with 1 mL IP buffer containing 0.5% NP-40. The final eluted proteins were boiled with 2 x Laemmli buffer / 5% beta-mercaptoethanol for 10 min at 95°C and the supernatant was used for Western blot analysis.

Detection of mitochondrial membrane potential in yeast cells

To examine mitochondrial membrane potential in yeast cells, BY4741 yeast cells expressing pYX212-DsRed-SY1 or pYX212-DsRed were pre-cultured overnight and diluted to an OD_{600nm} of 0.1. Cells were harvested at the endpoints, washed twice with $1\times$ PBS (pH 7.4), and treated with Rhodamine123 at a final concentration of 10 µM. Cells were incubated for 20 min at 30°C in the dark. Cells were then washed at least three times with $1\times$PBS (pH 7.4) before flow cytometric analysis using BD Accuri C6 (BD Biosciences, San Jose, USA) with FITC channel.

Cell lines, culture condition, transfection and drug treatment

The human embryonic kidney (HEK293t) cell line was obtained from the National Collection of Authenticated Cell Cultures (NCACC; Shanghai, China). Cells were cultured in Dulbecco's modified Eagle's medium (DMEM; Gibco, c11995500bt) supplemented with 10% fetal bovine serum (FBS; Biological Industries, 04-001-1acs) and 1% penicillin-streptomycin (HycloneTM, SV30010) in a humidified incubator containing 5% CO_2 at 37°C. Transfection was performed on six-well plates or confocal dishes for different purposes with plasmids using X-tremeGENE HP DNA Transfection Reagent according to the manufacturer's protocol. For drug treatment, cells were seeded at appropriate density in cell culture plates and cultured for 24 h. Cells were treated with 25 μM myriocin for 3 days or 20 μM DHS for 2 days before subsequent analysis.

Cell imaging in mammalian cells

To observe the association of SY1 IBs with mitochondria in mammalian cells, HEK293t cells were seeded into 35 mm confocal cell culture dishes (BioSharp), transfected with GFP-SY1, and incubated for 24 h. The cells were washed with PBS and then stained with 10 μM MitoTracker Deep Red FM at 37°C for 20 min. After staining, the cells were washed twice with PBS, and rinsed with PBS. Fluorescence images were captured using a Zeiss LSM 880 confocal microscope (Carl Zeiss, Jena, Germany), and Z-stack images for 3D rendering were taken at 0.4 μm interlayer distance. 3D rendering was performed using the Surface tool in the Imaris software.

For analysis of SY1 IBs formation under myriocin or DHS treatment, HEK293t cells were seeded in confocal culture dishes and transfected with pcDNA3.1-GFP-SY1 after treatment. Cells were fixed with 4% paraformaldehyde (Beyotime, P0099) for 20 min and permeabilized with 0.1% (v/v) Triton X-100 in PBS (Beyotime, P0096) for 15 min on ice. The cytoskeleton was then stained with phalloidin-Atto 565, followed by nuclei with DAPI (Southern Biotech, 0100-20). Images were captured using a Zeiss LSM 880 confocal microscope.

Measurement of mitochondrial function in mammalian cells

HEK293t cells were detached and transferred to 75-mm flow tubes and exposed to 5 μM MitoSOX in serum-free medium for 30 min in a 37°C cell incubator. Cells were then washed three times with HBSS buffer (Invitrogen,14025-092) and analyzed using a BD Accuri C6 flow cytometer.

Mitochondrial membrane potential (MMP) was measured with JC-1 according to the manufacturer's

instructions. Briefly, treated cells were harvested, washed twice with PBS, and then stained in a mixture of

500 µL medium and 500 µL JC-1 at 37° C for 20 min in the dark. Cells were washed 3 times with staining

buffer and analyzed by flow cytometry. Changes in MMP are represented by the ratio of red to green

fluorescence intensity.

Measurement of cell viability and cytotoxicity

Cell viability was measured by monitoring the cellular ATP content using a CellTiter-Lumi Luminescent

Cell Viability Assay Kit. Cells were treated and further incubated with 100 µL of detection reagent at room

temperature for 2 min to promote cell lysis. Cells were further incubated at room temperature for 10 min to

stabilize the luminescence signal, and ATP levels were then analyzed by using a Spark multimode

microplate reader (Tecan Trading AG, Männedorf, Switzerland).

Western blot analysis

Cells were harvested and lysed in RIPA buffer (Beyotime, P0013C) supplemented with a protease inhibitor

of 1 mM PMSF (Beyotime, ST506). Total protein concentrations were then quantified using the BCA

Protein Quantification Kit. A total of 20 µg of protein from each sample was separated by SDS-PAGE and

transferred to nitrocellulose membranes (Cytiva, 10600001). The membranes were blocked with 1% BSA

for 1 h at 37°C and incubated with primary antibodies: anti-Flag, anti-GFP, anti-SY1, anti-PGK1, anti-

GAPDH, followed by HRP-conjugated secondary antibody incubation: goat anti-rabbit IgG. Protein bands

were detected with ECL Prime Western Blotting Detection Reagent (GE, RPN2232) using an Amersham

imager 600 (GE, Piscataway, USA).

Generation of HEK293t SPTLC2 knockout cells by CRISPR/Cas9

For the generation of SPTLC2 knockout cell line, 2 guide RNA sequences for CRISPR were designed at

CRISPR design web site (http://crispr.mit.edu/). Both guide RNAs target the exon 1 of the SPTLC2 gene

(Figure 6-figure supplement 1). The DNA sequence of the guide RNA (gRNA) was synthesized and cloned

into the pX459 CRISPR/Cas9-Puro vector to construct knockout plasmids px459-SPTLC2-gRNA1 and

px459-SPTLC2-gRNA2.

Cells were seeded at appropriate densities in 35-mm dishes and grown for 24 h before transfection with the

plasmids. 48 h after transfection, cells were treated with 2 μg/mL Puromycin (Sangon Biotech, A610593)

for an additional three days. Single cells were then obtained by serial dilution and cultured for expansion.

Genomic DNA was extracted from 18 single-cell expansion cultures and the exon 1 sequence of SPTLC2

was amplified by PCR. Two cell lines with deletion mutations were identified by sequencing (Figure 6-

figure supplement 1).

Statistical analysis

Statistical analyses were performed with GraphPad Prism 8.0 software. Data were analyzed using the

Student's t-test and plotted as mean ± SD. Three or more biological replicates were analyzed unless

explicitly stated. Image J was used to count cells and compare bands of immunoblot. Results were

considered significant when $P < 0.05$. "n.s." stands for "not statistically significant."

Acknowledgments

We would like to thank Professor Charles Boone (University of Toronto) for providing SGA-V2 and ts-V5

strain collections, as well as the Moby plasmids. We also thank the Centre for Cellular Imaging at

University of Gothenburg for microscopy support, and Center for Large-scale cell-based screening at

University of Gothenburg for assistance with SGA analysis.

Additional information

Funding

Funder	Grant reference number	Author
National Natural Science Foundation of China	32000387	Xiuling Cao
Zhejiang Provincial Natural Science Foundation of China	LY23C060001	Xuejiao Jin
Scientific Research Foundation of Zhejiang A&F University	2021LFR053	Xuejiao Jin
Fellowship from China Scholarship Council and KU Leuven	grant C14/17/063	Ju Zheng
Swedish Cancer Fund (Cancerfonden)	19 0069 and 23 2769 Pj	Beidong Liu

Author contributions

Conceptualization: Beidong Liu; Methodology: Xiang Wu, Lei Zhao., Ju Zheng, Xinxin Hao; Investigation:

Xiuling Cao, Lei Zhao, Xuejiao Jin, Lihua Chen, Beidong Liu; Visualization: Xiuling Cao, Xiang Wu, Lei

Zhao, Lihua Chen. Supervision: Xiuling Cao, Shenkui Liu, Lihua Chen, Beidong Liu; Writing—original

draft: Xiuling Cao, Lei Zhao, Lihua Chen; Writing—review & editing: Xiuling Cao, Joris Winderickx,

Lihua Chen, Beidong Liu.

Additional files

Supplementary files

Supplementary file 1. Mutants with manually confirmed increased class 3 synphilin-1 aggregates.

Supplementary file 2. Distribution of GO terms among 84 candidate hits with significantly increased Class

3 aggregates via ClueGO analysis.

Supplementary file 3. Plasmids used in this study.

Supplementary file 4. Strains used in this study.

Data availability

All data generated or analyzed during this study are included in the manuscript and supplementary

information. No datasets have been submitted to public repositories.

References

Abramov, AY, Berezhnov, AV, Fedotova, EI, Zinchenko, VP, and Dolgacheva, LP. 2017. Interaction of misfolded
proteins and mitochondria in neurodegenerative disorders. *Biochemical Society Transactions* **45**: 1025-1033.
https://doi.org/10.1042/bst20170024

Amorim, JA, Coppotelli, G, Rolo, AP, Palmeira, CM, Ross, JM, and Sinclair, DA. 2022. Mitochondrial and
metabolic dysfunction in ageing and age-related diseases. *Nature Reviews Endocrinology* **18**: 243-258.
https://doi.org/10.1038/s41574-021-00626-7

Arrasate, M, Mitra, S, Schweitzer, ES, Segal, MR, and Finkbeiner, S. 2004. Inclusion body formation reduces
levels of mutant huntingtin and the risk of neuronal death. *Nature* **431**: 805-810.
https://doi.org/10.1038/nature02998

Bagola, K, and Sommer, T. 2008. Protein quality control: on IPODs and other JUNQ. *Current Biology* **18**:
R1019-R1021. https://doi.org/10.1016/j.cub.2008.09.036

Bäuerlein, FJB, Saha, I, Mishra, A, Kalemanov, M, Martínez-Sánchez, A, Klein, R, Dudanova, I, Hipp, MS,
Hartl, FU, Baumeister, W, and Fernández-Busnadiego, R. 2017. In situ architecture andcellular interactions of
PolyQ inclusions. *Cell* **171**: 179-187.e110. https://doi.org/10.1016/j.cell.2017.08.009

Bento-Oliveira, A, Santos, FC, Marquês, JT, Paulo, PMR, Korte, T, Herrmann, A, Marinho, HS, and de Almeida,
RFM. 2020. Yeast sphingolipid-enriched domains and membrane compartments in the absence of
mannosyldiinositolphosphorylceramide. *Biomolecules* **10**: 871. https://doi.org/10.3390/biom10060871

Bindea, G, Mlecnik, B, Hackl, H, Charoentong, P, Tosolini, M, Kirilovsky, A, Fridman, WH, Pagès, F,
Trajanoski, Z, and Galon, J. 2009. ClueGO: a Cytoscape plug-in to decipher functionally grouped gene ontology
and pathway annotation networks. *Bioinformatics* **25**: 1091-1093. https://doi.org/10.1093/bioinformatics/btp101

Buchberger, A, Bukau, B, and Sommer, T. 2010. Protein quality control in the cytosol and the endoplasmic
reticulum: brothers in arms. *Molecular Cell* **40**: 238-252. https://doi.org/10.1016/j.molcel.2010.10.001

Busciglio, J, Pelsman, A, Wong, C, Pigino, G, Yuan, M, Mori, H, and Yankner, BA. 2002. Altered metabolism of
the amyloid β precursor protein is associated with mitochondrial dysfunction in Down's Syndrome. *Neuron* **33**:
677-688. https://doi.org/10.1016/S0896-6273(02)00604-9

Büttner, S, Bitto, A, Ring, J, Augsten, M, Zabrocki, P, Eisenberg, T, Jungwirth, H, Hutter, S, Carmona-Gutierrez,
D, Kroemer, G, et al. 2008. Functional mitochondria are required for alpha-synuclein toxicity in aging yeast.
Journal of Biological Chemistry 283: 7554-7560. https://doi.org/10.1074/jbc.M708477200Büttner, S, Delay, C,
Franssens, V, Bammens, T, Ruli, D, Zaunschirm, S, de Oliveira, RM, Outeiro, TF, Madeo, F, Buée, L, et al. 2010.
Synphilin-1 enhances α-synuclein aggregation in yeast and contributes to cellular stress and cell death in a Sir2-
dependent manner. *PLoS One* **5**: e13700. https://doi.org/10.1371/journal.pone.0013700

Casadei, N, Pöhler, AM, Tomás-Zapico, C, Torres-Peraza, J, Schwedhelm, I, Witz, A, Zamolo, I, De Heer, R,
Spruijt, B, Noldus, LP, et al. 2014. Overexpression of synphilin-1 promotes clearance of soluble and misfolded
alpha-synuclein without restoring the motor phenotype in aged A30P transgenic mice. *Human Molecular*
Genetics **23**: 767-781. https://doi.org/10.1093/hmg/ddt467

Chen, B, Retzlaff, M, Roos, T, and Frydman, J. 2011. Cellular strategies of protein quality control. *Cold Spring*
Harbor perspectives in biology **3**: a004374. https://doi.org/10.1101/cshperspect.a004374

Cutler, RG, Pedersen, WA, Camandola, S, Rothstein, JD, and Mattson, MP. 2002. Evidence that accumulation of

ceramides and cholesterol esters mediates oxidative stress–induced death of motor neurons in amyotrophic lateral sclerosis. *Annals of Neurology* **52**: 448-457. https://doi.org/10.1002/ana.10312

Douglas, PM, Treusch, S, Ren, H-Y, Halfmann, R, Duennwald, ML, Lindquist, S, and Cyr, DM. 2008. Chaperone-dependent amyloid assembly protects cells from prion toxicity. *Proceedings of the National Academy of Sciences, USA* **105**: 7206-7211. https://doi.org/10.1073/pnas.0802593105

France-Lanord, V, Brugg, B, Michel, PP, Agid, Y, and Ruberg, M. 1997. Mitochondrial free radical signal in ceramide-dependent apoptosis: a putative mechanism for neuronal death in Parkinson's disease. *Journal of Neurochemistry* **69**: 1612-1621. https://doi.org/10.1046/j.1471-4159.1997.69041612.x

Frydman, J. 2001. Folding of newly translated proteins in vivo: the role of molecular chaperones. *Annual Review of Biochemistry* **70**: 603-647. https://doi.org/10.1146/annurev.biochem.70.1.603

Gong, H, Romanova, NV, Allen, KD, Chandramowlishwaran, P, Gokhale, K, Newnam, GP, Mieczkowski, P, Sherman, MY, and Chernoff, YO. 2012. Polyglutamine toxicity is controlled by prion composition and gene dosage in yeast. *PLoS Genetics* **8**: e1002634. https://doi.org/10.1371/journal.pgen.1002634

Guo, Q, Lehmer, C, Martínez-Sánchez, A, Rudack, T, Beck, F, Hartmann, H, Pérez-Berlanga, M, Frottin, F, Hipp, MS, Hartl, FU, et al. 2018. In situ structure of neuronal C9orf72 Poly-GA aggregates reveals proteasome recruitment. *Cell* **172**: 696-705.e612. https://doi.org/10.1016/j.cell.2017.12.030

Haelterman, NA, Yoon, WH, Sandoval, H, Jaiswal, M, Shulman, JM, and Bellen, HJ. 2014. A mitocentric view of Parkinson's disease. *Annual Review of Neuroscience* **37**: 137-159. https://doi.org/10.1146/annurev-neuro-071013-014317

Hannun, YA, and Obeid, LM. 2008. Principles of bioactive lipid signalling: lessons from sphingolipids. *Nat Rev Mol Cell Biol* **9**: 139-150. https://doi.org/10.1038/nrm2329

Hannun, YA, and Obeid, LM. 2018. Sphingolipids and their metabolism in physiology and disease. *Nature Reviews Molecular Cell Biology* **19**: 175-191. https://doi.org/10.1038/nrm.2017.107

Hartl, FU, and Hayer-Hartl, M. 2009. Converging concepts of protein folding in vitro and in vivo. *Nature Structural & Molecular Biology* **16**: 574. https://doi.org/10.1038/nsmb.1591

Hashimoto, M, Rockenstein, E, Crews, L, and Masliah, E. 2003. Role of protein aggregation in mitochondrial dysfunction and neurodegeneration in Alzheimer's and Parkinson's diseases. *Neuromolecular Medicine* **4**: 21-36. https://doi.org/10.1385/nmm:4:1-2:21

Ho, CH, Magtanong, L, Barker, SL, Gresham, D, Nishimura, S, Natarajan, P, Koh, JLY, Porter, J, Gray, CA, Andersen, RJ, et al. 2009. A molecular barcoded yeast ORF library enables mode-of-action analysis of bioactive compounds. *Nature Biotechnology* **27**: 369-377. https://doi.org/10.1038/nbt.1534

Hsu, LJ, Sagara, Y, Arroyo, A, Rockenstein, E, Sisk, A, Mallory, M, Wong, J, Takenouchi, T, Hashimoto, M, and Masliah, E. 2000. α-Synuclein promotes mitochondrial deficit and oxidative stress. *The American Journal of Pathology* **157**: 401-410. https://doi.org/10.1016/S0002-9440(10)64553-1

Huang, X, Liu, J, and Dickson, RC. 2012. Down-regulating sphingolipid synthesis increases yeast lifespan. *PLoS Genetics* **8**: e1002493. https://doi.org/10.1371/journal.pgen.1002493

Huh, WK, Falvo, JV, Gerke, LC, Carroll, AS, Howson, RW, Weissman, JS, and O'Shea, EK. 2003. Global

analysis of protein localization in budding yeast. *Nature* **425**: 686-691. https://doi.org/10.1038/nature02026

Jin, HG, Yamashita, H, Nakamura, T, Fukuba, H, Takahashi, T, Hiji, M, Kohriyama, T, and Matsumoto, M. 2008.
Synphilin-1 transgenic mice exhibit mild motor impairments. *Neuroscience Letters* **445**: 12-17.
https://doi.org/10.1016/j.neulet.2008.08.073

Kaganovich, D, Kopito, R, and Frydman, J. 2008. Misfolded proteins partition between two distinct quality
control compartments. *Nature* **454**: 1088-1095. https://doi.org/10.1038/nature07195

Kitamura, A, Kubota, H, Pack, CG, Matsumoto, G, Hirayama, S, Takahashi, Y, Kimura, H, Kinjo, M, Morimoto,
RI, and Nagata, K. 2006. Cytosolic chaperonin prevents polyglutamine toxicity with altering the aggregation
state. *Nature Cell Biology* **8**: 1163-1170. https://doi.org/10.1038/ncb1478

Kopito, RR. 2000. Aggresomes, inclusion bodies and protein aggregation. *Trends in Cell Biology* **10**: 524-530.
https://doi.org/10.1016/S0962-8924(00)01852-3

Krenz, A, Falkenburger, BH, Gerhardt, E, Drinkut, A, and Schulz, JB. 2009. Aggregate formation and toxicity
by wild-type and R621C synphilin-1 in the nigrostriatal system of mice using adenoviral vectors. *Journal of*
Neurochemistry **108**: 139-146. https://doi.org/10.1111/j.1471-4159.2008.05755.x

Li, S, and Kim, H-E. 2022. Implications of sphingolipids on aging and age-related diseases. *Frontiers in aging* **2**:
797320. https://doi.org/10.3389/fragi.2021.797320

Lim, J, and Yue, Z. 2015. Neuronal aggregates: formation, clearance, and spreading. *Developmental Cell* **32**:
491-501. https://doi.org/10.1016/j.devcel.2015.02.002

Lima, T, Li, TY, Mottis, A, and Auwerx, J. 2022. Pleiotropic effects of mitochondria in aging. *Nature Aging* **2**:
199-213. https://doi.org/10.1038/s43587-022-00191-2

Lin, G, Wang, L, Marcogliese, PC, and Bellen, HJ. 2019. Sphingolipids in the pathogenesis of Parkinson's
disease and Parkinsonism. *Trends in Endocrinology and Metabolism* **30**: 106-117.
https://doi.org/10.1016/j.tem.2018.11.003

Liu, J, Li, T, Thomas, JM, Pei, Z, Jiang, H, Engelender, S, Ross, CA, and Smith, WW. 2016. Synphilin-1
attenuates mutant LRRK2-induced neurodegeneration in Parkinson's disease models. *Human Molecular*
Genetics **25**: 672-680. https://doi.org/10.1093/hmg/ddv504

Liu, QaF, Benjamin and Unruh, Jay R. and Guo, Fengli and Si, Kausik and Li, Rong and Zhou, Chuankai. 2021.
Mitochondrial origin of cytosolic protein aggregation. *SSRN* **37**: 3808300. https://doi.org/10.2139/ssrn.3808300

Mielke, MM, and Lyketsos, CG. 2010. Alterations of the sphingolipid pathway in Alzheimer's disease: new
biomarkers and treatment targets? *Neuromolecular Medicine* **12**: 331-340. https://doi.org/10.1007/s12017-010-
8121-y

Miller, SB, Mogk, A, and Bukau, B. 2015. Spatially organized aggregation of misfolded proteins as cellular
stress defense strategy. *Journal of Molecular Biology* **427**: 1564-1574.
https://doi.org/10.1016/j.jmb.2015.02.006

Moldavski, O, Amen, T, Levin-Zaidman, S, Eisenstein, M, Rogachev, I, Brandis, A, Kaganovich, D, and
Schuldiner, M. 2015. Lipid droplets are essential for efficient clearance of cytosolic inclusion bodies.
Developmental Cell **33**: 603-610. https://doi.org/10.1016/j.devcel.2015.04.015

Nowicka, U, Chroscicki, P, Stroobants, K, Sladowska, M, Turek, M, Uszczynska-Ratajczak, B, Kundra, R,
Goral, T, Perni, M, Dobson, CM, et al. 2021. Cytosolic aggregation of mitochondrial proteins disrupts cellular
homeostasis by stimulating the aggregation of other proteins. *eLife* **10**: e65484.
https://doi.org/10.7554/eLife.65484

Nuber, S, Franck, T, Wolburg, H, Schumann, U, Casadei, N, Fischer, K, Calaminus, C, Pichler, BJ, Chanarat, S,
Teismann, P, et al. 2010. Transgenic overexpression of the alpha-synuclein interacting protein synphilin-1 leads
to behavioral and neuropathological alterations in mice. *Neurogenetics* **11**: 107-120.
https://doi.org/10.1007/s10048-009-0212-2

O'Farrell, C, Murphy, DD, Petrucelli, L, Singleton, AB, Hussey, J, Farrer, M, Hardy, J, Dickson, DW, and
Cookson, MR. 2001. Transfected synphilin-1 forms cytoplasmic inclusions in HEK293 cells. *Brain Res Mol*
Brain Res **97**: 94-102. https://doi.org/10.1016/s0169-328x(01)00292-3

Ogretmen, B. 2018. Sphingolipid metabolism in cancer signalling and therapy. *Nature Reviews: Cancer* **18**: 33-
50. https://doi.org/10.1038/nrc.2017.96

Ogrodnik, M, Salmonowicz, H, Brown, R, Turkowska, J, Średniawa, W, Pattabiraman, S, Amen, T, Abraham, A-
c, Eichler, N, Lyakhovetsky, R, and Kaganovich, D. 2014. Dynamic JUNQ inclusion bodies are asymmetrically
inherited in mammalian cell lines through the asymmetric partitioning of vimentin. *Proceedings of the National*
Academy of Sciences, USA **111**: 8049-8054. https://doi.org/10.1073/pnas.1324035111

Olzscha, H, Schermann, SM, Woerner, AC, Pinkert, S, Hecht, MH, Tartaglia, GG, Vendruscolo, M, Hayer-Hartl,
M, Hartl, FU, and Vabulas, RM. 2011. Amyloid-like aggregates sequester numerous metastable proteins with
essential cellular functions. *Cell* **144**: 67-78. https://doi.org/10.1016/j.cell.2010.11.050

Park, JH, Burgess, JD, Faroqi, AH, DeMeo, NN, Fiesel, FC, Springer, W, Delenclos, M, and McLean, PJ. 2020.
Alpha-synuclein-induced mitochondrial dysfunction is mediated via a sirtuin 3-dependent pathway. *Molecular*
neurodegeneration **15**: 5. https://doi.org/10.1186/s13024-019-0349-x

Park, S-H, Kukushkin, Y, Gupta, R, Chen, T, Konagai, A, Hipp, Mark S, Hayer-Hartl, M, and Hartl, FU. 2013.
PolyQ proteins interfere with nuclear degradation of cytosolic proteins by sequestering the Sis1p chaperone.
Cell **154**: 134-145. https://doi.org/10.1016/j.cell.2013.06.003

Polito, L, Greco, A, and Seripa, D. 2016. Genetic profile, environmental exposure, and their interaction in
Parkinson's Disease. *Parkinson's Disease* **2016**: 6465793. https://doi.org/10.1155/2016/6465793

Ribeiro, CS, Carneiro, K, Ross, CA, Menezes, JRL, and Engelender, S. 2002. Synphilin-1 is developmentally
localized to synaptic terminals, and its association with synaptic vesicles is modulated by alpha-synuclein.
Journal of Biological Chemistry **277**: 23927-23933. https://doi.org/10.1074/jbc.M201115200

Ross, CA, and Poirier, MA. 2004. Protein aggregation and neurodegenerative disease. *Nature Medicine* **10**: S10-
17. https://doi.org/10.1038/nm1066

Sacket, SJ, Chung, H-y, Okajima, F, and Im, D-s. 2009. Increase in sphingolipid catabolic enzyme activity
during aging. *Acta Pharmacologica Sinica* **30**: 1454-1461. https://doi.org/10.1038/aps.2009.136

Schlarmann, P, Ikeda, A, and Funato, K. 2021. Membrane contact sites in yeast: control hubs of sphingolipid
homeostasis. *Membranes* **11**: 971. https://doi.org/10.3390/membranes11120971

Shishido, T, Nagano, Y, Araki, M, Kurashige, T, Obayashi, H, Nakamura, T, Takahashi, T, Matsumoto, M, and

Maruyama, H. 2019. Synphilin-1 has neuroprotective effects on MPP(+)-induced Parkinson's disease model cells by inhibiting ROS production and apoptosis. *Neuroscience Letters* **690**: 145-150. https://doi.org/10.1016/j.neulet.2018.10.020

Singh, P, and Li, R. 2018. Emerging roles for sphingolipids in cellular aging. *Current Genetics* **64**: 761-767. https://doi.org/10.1007/s00294-017-0799-z

Smeal, T, Claus, J, Kennedy, B, Cole, F, and Guarente, L. 1996. Loss of transcriptional silencing causes sterility in old mother cells of *S. cerevisiae*. *Cell* **84**: 633-642. https://doi.org/10.1016/s0092-8674(00)81038-7

Smith, WW, Liu, Z, Liang, Y, Masuda, N, Swing, DA, Jenkins, NA, Copeland, NG, Troncoso, JC, Pletnikov, M, Dawson, TM, et al. 2010. Synphilin-1 attenuates neuronal degeneration in the A53T alpha-synuclein transgenic mouse model. *Human Molecular Genetics* **19**: 2087-2098. https://doi.org/10.1093/hmg/ddq086

Song, J, Yang, Q, Yang, J, Larsson, L, Hao, X, Zhu, X, Malmgren-Hill, S, Cvijovic, M, Fernandez-Rodriguez, J, Grantham, J, et al. 2014. Essential genetic interactors of SIR2 required for spatial sequestration and asymmetrical inheritance of protein aggregates. *PLoS Genetics* **10**: e1004539. https://doi.org/10.1371/journal.pgen.1004539

Tafesse, FG, Rashidfarrokhi, A, Schmidt, FI, Freinkman, E, Dougan, S, Dougan, M, Esteban, A, Maruyama, T, Strijbis, K, and Ploegh, HL. 2015. Disruption of sphingolipid biosynthesis blocks phagocytosis of *Candida albicans*. *PLoS Pathogens* **11**: e1005188. https://doi.org/10.1371/journal.ppat.1005188

Takahashi, T, Yamashita, H, Nagano, Y, Nakamura, T, Kohriyama, T, and Matsumoto, M. 2006. Interactions of Synphilin-1 with phospholipids and lipid membranes. *FEBS Letters* **580**: 4479-4484. https://doi.org/10.1016/j.febslet.2006.07.019

Tanaka, M, Kim, YM, Lee, G, Junn, E, Iwatsubo, T, and Mouradian, MM. 2004. Aggresomes formed by alpha-synuclein and synphilin-1 are cytoprotective. *Journal of Biological Chemistry* **279**: 4625-4631. https://doi.org/10.1074/jbc.M310994200

Tong, AH, Evangelista, M, Parsons, AB, Xu, H, Bader, GD, Pagé, N, Robinson, M, Raghibizadeh, S, Hogue, CW, Bussey, H, et al. 2001. Systematic genetic analysis with ordered arrays of yeast deletion mutants. *Science* **294**: 2364-2368. https://doi.org/10.1126/science.1065810

Tong, AH, Lesage, G, Bader, GD, Ding, H, Xu, H, Xin, X, Young, J, Berriz, GF, Brost, RL, Chang, M, et al. 2004. Global mapping of the yeast genetic interaction network. *Science* **303**: 808-813. https://doi.org/10.1126/science.1091317

Tong, AHY, and Boone, C (2007). 16 High-throughput strain construction and systematic synthetic lethal screening in *Saccharomyce scerevisiae*. In Methods in Microbiology, I. Stansfield, and M.J.R. Stark, eds. (Academic Press), pp. 369-707. https://doi.org/10.1016/S0580-9517(06)36016-3

Trayssac, M, Hannun, YA, and Obeid, LM. 2018. Role of sphingolipids in senescence: implication in aging and age-related diseases. *Journal of Clinical Investigation* **128**: 2702-2712. https://doi.org/10.1172/jci97949

Wang, P, Deng, J, Dong, J, Liu, J, Bigio, EH, Mesulam, M, Wang, T, Sun, L, Wang, L, Lee, AY, et al. 2019a. TDP-43 induces mitochondrial damage and activates the mitochondrial unfolded protein response. *PLoS Genetics* **15**: e1007947. https://doi.org/10.1371/journal.pgen.1007947

Wang, X, Becker, K, Levine, N, Zhang, M, Lieberman, AP, Moore, DJ, and Ma, J. 2019b. Pathogenic alpha-

synuclein aggregates preferentially bind to mitochondria and affect cellular respiration. *Acta neuropathologica*

communications **7**: 41. https://doi.org/10.1186/s40478-019-0696-4

Weisberg, SJ, Lyakhovetsky, R, Werdiger, A-c, Gitler, AD, Soen, Y, and Kaganovich, D. 2012.

Compartmentalization of superoxide dismutase 1 (SOD1G93A) aggregates determines their toxicity.

Proceedings of the National Academy of Sciences, USA **109**: 15811-15816.

https://doi.org/10.1073/pnas.1205829109

Yang, J, Hao, X, Cao, X, Liu, B, and Nyström, T. 2016. Spatial sequestration and detoxification of Huntingtin

by the ribosome quality control complex. *eLife* **5**: e11792. https://doi.org/10.7554/eLife.11792

Zaarur, N, Meriin, AB, Gabai, VL, and Sherman, MY. 2008. Triggering aggresome formation. Dissecting

aggresome-targeting and aggregation signals in synphilin 1. *Journal of Biological Chemistry* **283**: 27575-27584.

https://doi.org/10.1074/jbc.M802216200

Zhao, L, Yang, Q, Zheng, J, Zhu, X, Hao, X, Song, J, Lebacq, T, Franssens, V, Winderickx, J, Nystrom, T, and

Liu, B. 2016. A genome-wide imaging-based screening to identify genes involved in synphilin-1 inclusion

formation in *Saccharomyces cerevisiae*. *Scientific Reports* **6**: 30134. https://doi.org/10.1038/srep30134

764 **Figures and figure supplements**

Figure 1 SY1 aggregates mature into large IBs over time in normal cells, but this process is defective

in aged cells.

(**A**) The maturation process of SY1 aggregates in yeast cells from scattered small aggregates to large single

IBs over time. DsRed-SY1 was expressed in yeast with the constitutive promoter TPI1, the phenotype of

SY1 aggregates was visualized at 4 h, 16 h and 48 h. Scale bar = 2 μm. (**B**) Cells with different phenotypes

of SY1 IBs are defined as three classes: Class 1 refers to only one IB in a cell, Class 2 refers to two IBs in

a cell, and Class 3 indicates three or more IBs in a cell. Scale bar = 2 μm. (**C**) Time-course study showing

that SY1 aggregates mature after 24 h of cultivation, when the cells have traversed the diauxic shift to

enter the respiratory and stationary growth phase. IBs were examined every 4 h from 4 to 48 h; green bars

indicate percentage of Class 3; white bars indicate percentage of Class 2; red bars indicate percentage of

Class 1; the purple curve indicates the optical density of the cell culture, indicated by the vertical axis on

the right. Error bars indicate the SD. (**D**) Maturation of SY1 IBs was defective in old yeast cells. Scale bar

= 5 μm. (**E**) The proportion of cells with SY1 IBs was increased in old cells (61%) compared to young

cells (27%). Data were analyzed using the Student's t-test. Error bars indicate the SD. (**F**) The number of

SY1 IBs in a cell was increased in young cells (2.64±1.29) versus old cells (4.91±3.32). Data were

analyzed using the Student's t-test. Error bars indicate the SD.

Figure 2 Genome-wide high-content imaging screen identifying the serine C-palmitoyltransferase complex (SPT) as a modulator of SY1 inclusion maturation.

(**A**) Schematic workflow for identifying mutants that regulate SY1 inclusion maturation. (**B**) The GO enrichment analysis of genes whose mutants exhibited increased SY1 Class 3 aggregates (ClueGO, cut-off: $P < 0.05$). (**C**) The network of the most enriched modules obtained above in 2B. (**D-E**) The increased SY1 Class 3 aggregates in the SPT mutant (*lcb1-4, lcb2-1, tsc3-2*) were rescued by backer-introducing the corresponding wild-type gene (a MoBY plasmid carries a WT gene, and p5472 is the empty vector control). Representative images from three independent experiments are shown in D, and quantifications were shown in E. Scale bar = 2 μm. Data were analyzed using the Student's t-test. Error bars indicate the SD.

Figure 2-figure supplement 1. The time course of SY1 IB maturation in the SPT mutants

SPT mutants (*lcb1-4, lcb2-1, tsc3-2*) expressing the pYX212-DsRed-SY1 plasmid were harvested and observed every 4 hours.

Figure 3 Sphingolipids regulate SY1 IB maturation and cytotoxicity in yeast cells.

(**A**) A schematic diagram of the sphingolipid pathway in yeast cells. Myriocin blocks sphingolipid synthesis via inhibition of the serine C-palmitoyltransferase (SPT) complex. (**B-C**) Dihydrosphingosine (DHS) supplementation rescued the deficiency of SY1 IBs maturation in SPT mutants (*lcb1-4, lcb2-1, tsc3-2*). Representative images from three independent experiments are shown in B, and quantifications are shown in C. Scale bar = 2 μm. Data were analyzed using the Student's t-test. Error bars indicate the SD. (**D-E**) Blocking intracellular sphingolipid synthesis with myriocin resulted in defective maturation of SY1 IBs, and exogenous addition of DHS could reverse the effect of myriocin. Representative images from three independent experiments are shown in D, and quantifications are shown in E. Scale bar = 2 μm. Data were analyzed using the Student's t-test. Error bars indicate the SD. (**F**) DHS ameliorated the cytotoxicity of SY1 in SPT mutants (*lcb1-4, lcb2-1, tsc3-2*).

Figure 4 SY1 IBs associate with mitochondria in yeast cells.

(**A**) SY1 IBs had no obvious association with the plasma membrane (Pma1-GFP), vacuoles (Vph1-GFP),

or ER (Sec63-GFP), but were strongly associated with mitochondria (Tom70-GFP) (See also Figure 4-

figure supplement 1A). Scale bar = 2 μm. **(B)** MitoTracker probe confirmed that all the three classes of IBs

were closely associated with mitochondria (arrow heads indicate SY1 aggregates surrounded by

mitochondria). Scale bar = 2 μm. n.s., not statistically significant. Data were analyzed using the Student's

t-test. Error bars indicate the SD. **(C)** Super-resolution microscopy and 3D-SIM construction demonstrated

that the SY1 IBs are packed and surrounded by mitochondria. Scale bar = 1 μm. **(D)** Co-

immunoprecipitation assays showing that SY1 physically binds to several mitochondrial membrane

proteins, including Tom70, Tom37 and Tim44. The yeast mitochondrial membrane proteins Tom70, Tom37,

Sam35 and Tim44 were individually C-terminally tagged with GFP, and the strains were then transformed

with a control plasmid (Vector) or a SY1-overproducing plasmid (SY1), respectively. Cells were grown for

24 h and harvested for subsequent IP experiments. **(E-F)** The association of SY1 IBs with mitochondria

decreases when the physical interaction between SY1 and mitochondrial membrane proteins is disrupted

(*tom70Δ*, *tom37Δ*, *tim44*). Representative images from three independent experiments are shown in E and

quantifications are shown in F. Scale bar = 2 μm. Data were analyzed using the Student's t-test. Error bars

indicate the SD.

Figure 4-figure supplement 1. Statistics of SY1 IBs surrounded by cellular organelles.

(A) Statistics of the proportion of SY1 IBs surrounded by different organelles, corresponding to Fig 4A.

Data were analyzed using the Student's t-test. Error bars indicate the SD. **(B)** The 3D reconstruction using

Imaris software to demonstrate the interrelationship between SY1 Class 3 aggregates and mitochondria

(arrow heads indicate SY1 aggregates surrounded by mitochondria). BY4741 cells expressing DsRed-SY1

and mitochondria were stained using MitoTracker green dye. Scale = 2 μm. **(C)** Statistics of the proportion

of SY1 IBs surrounded by mitochondria in myriocin treatment and the DMSO treatment control. n.s., not

statistically significant. Data were analyzed using the Student's t-test. Error bars indicate the SD. **(D)**

Statistics of the proportion of Cells with SY1 Class 3 aggregates in *tom70Δ*, *tom37Δ*, and *tim44* mutants.

Data were analyzed using the Student's t-test. Error bars indicate the SD. **(E)** SY1 IBs are surrounded by

mitochondria in the rho0 strain. Mitochondria of the rho0 strain were labeled with Tom70-GFP and

photographed after expression of the pYX212-DsRed-SY1 plasmid. Scale = 2 μm. **(F)** Statistics of the

proportion of three different classes of SY1 IBs in BY4741 wild-type and rho0 strains, and there is no

significant difference between two strains. n.s., not statistically significant. Data were analyzed using the

Student's t-test. Error bars indicate the SD.

Figure 4-figure supplement 2. IBs of TDP43, FUS1 and Htt103Qp interact with mitochondria.

Shown are the IBs formed by TDP43, FUS1 and Htt103Qp when fused to GFP and their co-localization

with Tom70, which was genomically tagged with RFP. Scale = 2 μm.

Figure 5 Sphingolipids regulate SY1-induced mitochondrial toxicity in yeast cells.

(A) Overexpression of SY1 decreased mitochondrial membrane potential (indicated by Rho 123 fluor

intensity) and reduced cell viability (indicated by ATP level) in yeast cells. Data were analyzed using the

Student's t-test. Error bars indicate the SD. **(B)** Effects of myriocin treatment on SY1 cytotoxicity in yeast

cells as judged by mitochondrial membrane potential and cell viability. Data were analyzed using the

Student's t-test. Error bars indicate the SD. **(C)** Effects of exogenous addition of DHS on SY1 cytotoxicity

in yeast cells based on mitochondrial membrane potential and cell viability. Data were analyzed using the

Student's t-test. Error bars indicate the SD.

Figure 5-figure supplement 1. The effect of myriocin (A) and DHS (B) treatment on SY1 protein

expression in yeast cells.

SY1 expression was examined by Western blot analysis in samples prepared from cells used in the

experiments corresponding to Fig. 5B and Fig. 5C respectively. n.s., not statistically significant. Data were

analyzed using the Student's t-test.

Figure 5-figure supplement 2. SY1-induced mitochondrial toxicity decreased in *tom70Δ* and *tom37Δ*

cells.

Mitochondrial membrane potential was examined by expressing SY1 in *his3Δ*, *tom70Δ* and *tom37Δ* strains.

n.s., not statistically significant. Data were analyzed using the Student's t-test. Error bars indicate the SD.

Figure 6 SPTLC2 knockout alters the morphology of SY1 IBs and increases the cytotoxicity in

mammalian cells.

(A) The 3D reconstruction showed that SY1 IBs were associated with mitochondria in mammalian

HEK293t cells. GFP-SY1 was expressed in HEK293t cells and mitochondria were probed with

MitoTracker. The 3D structure was constructed using Imaris software. Scale bar = 10 μm. **(B)** SY1

expression decreased mitochondrial function in HEK293t cells. Mitochondrial membrane potential was

monitored by JC-1 staining, mitochondrial superoxide was analyzed by MitoSOX mean intensity values. Data were analyzed using the Student's t-test. Error bars indicate the SD. **(C)** The 3D structures of spherical and irregular SY1 IBs, as well as their interaction with mitochondria was visualized using Imaris software (See also Figure 6-figure supplement 2). **(D)** SPTLC2 knockout HEK293t cells had impaired management of SY1 IBs, with increased dispersed irregular IB structures. Scale bar = 20 μm. Data were analyzed using the Student's t-test. Error bars indicate the SD. **(E)** SPTLC2 knockout increased SY1 cytotoxicity. Mitochondrial membrane potential (indicated by JC-1 staining) assays / cell viability (indicated by luminescence) were analyzed. Data were analyzed using the Student's t-test. Error bars indicate the SD.

Figure 6-figure supplement 1. SPTLC2-Knockout by CRISPR/Cas9 genome editing.

(A) The experimental flow chart for SPTLC2 knockout in HEK293t cells using CRISPR/Cas9. **(B)** The sequencing results of the corresponding deletion sites of two obtained knockout cell lines.

Figure 6-figure supplement 2. Statistics of SY1 IB formation.

(A) Statistics of SY1 IB formation in wild type HEK293t and SPTLC2 knockout cells. Data were analyzed using the Student's t-test. Error bars indicate the SD. **(B)** SPTLC2 knockout HEK293t cells were simultaneously transfected with GFP-SY1 expression vector and SPTLC2 complementary plasmid (EX-S0597-Lv242- SPTLC2) or empty vector control (EX-NEG-Lv242), and the percentage of IB-forming cells and the percentage of irregular IB in IB-forming cells were counted. Data were analyzed using the Student's t-test. Error bars indicate the SD.

Figure 6-figure supplement 3. The expression of SY1 protein.

The expression of SY1 protein in HEK293t SPTLC2 knockout cells. n.s., not statistically significant. Data were analyzed using the Student's t-test.

Figure 7 Sphingolipids regulate SY1-induced cytotoxicity and SY1 IB morphology in mammalian cells.

(A) Inhibition of the intracellular sphingolipid synthesis pathway by myriocin enhanced SY1 cytotoxicity based on mitochondrial membrane potential and ATP content assays. Data were analyzed using the Student's t-test. Error bars indicate the SD. **(B)** Addition of exogenous DHS rescued the cytotoxicity caused by SY1 expression. Data were analyzed using the Student's t-test. Error bars indicate the SD. **(C)**

Exogenous addition of DHS to SPTLC2 knockout cells restored the typical spherical SY1 IBs in HEK293t.

Statistics of the ratio of cells with dispersed irregular IBs after exogenous addition of DMSO and DHS to

HEK293t SPTLC2 knockout cells (See also Figure 7-figure supplement 2). Scale bar = 20 μm. Data were

analyzed using the Student's t-test. Error bars indicate the SD. **(D)** DHS ameliorated SY1 cytotoxicity by

normalizing mitochondrial membrane potential and cell viability. Data were analyzed using the Student's t-

test. Error bars indicate the SD.

Figure 7-figure supplement 1. The expression of SY1 protein in HEK293t and SPTLC2 knockout

cells

(A) myriocin treatment significantly increased the number of SY1 IBs compared to controls. Data were

analyzed using the Student's t-test. Error bars indicate the SD. **(B-C)** The effect of myriocin (B) and DHS

(C) treatment on SY1 expression in HEK293t cells. n.s., not statistically significant. Data were analyzed

using the Student's t-test. **(D)** The effect of DHS treatment on SY1 expression in SPTLC2 knockout cells.

n.s., not statistically significant. Data were analyzed using the Student's t-test.

Figure 7-figure supplement 2. Statistics of SY1 IB formation.

Statistics of the effect on SY1 IB formation in SPTLC2 knockout cells after exogenous addition of either

DMSO or DHS. Data were analyzed using the Student's t-test. Error bars indicate the SD.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7