Title: Hypothalamic representation of the imminence of predator threat detected by the

vomeronasal organ in mice

Authors: Quynh Anh Thi Nguyen[1], Andrea Rocha[2], Yuna Yamashita[2], Ricky Chhor[2], Christian

Stadler[1], Crystal Pontrello[2], Sachiko Haga-Yamanaka[1,2*]

Affiliations: 1. Neuroscience Graduate Program, 2. Department of Molecular, Cell, and Systems

Biology, University of California, Riverside

*Corresponding author

Email: sachikoy@ucr.edu

Abstract

Animals have the innate ability to select optimal defensive behavioral outputs with an

appropriate intensity in response to predator threat in specific contexts. Such innate behavioral

decisions are thought to be computed in the medial hypothalamic nuclei that contain neural

populations directly controlling defensive behavioral outputs. The vomeronasal organ (VNO) is

one of the major sensory input channels through which predator cues are detected with ascending

inputs to the medial hypothalamic nuclei, especially to the ventromedial hypothalamus (VMH),

through the medial amygdala (MeA) and bed nucleus of the stria terminalis (BNST). Here, we

show that cat saliva contains predator cues that signal imminence of predator threat and regulate

the robustness of freezing behavior through the VNO in mice. Cat saliva activates neurons

expressing the V2R-A4 subfamily of sensory receptors, suggesting the existence of specific

receptor groups responsible for freezing behavior induced by the predator cues. The number of

VNO neurons activated in response to saliva correlates with the freshness of saliva and the

intensity of freezing behavior, while the downstream neurons in the accessory olfactory bulb

(AOB) and defensive behavioral circuit are quantitatively equally activated by fresh and old

saliva. Strikingly, however, only the number of VMH neurons activated by fresh saliva

positively correlates with the intensity of freezing behavior. Detailed analysis of the spatial

distribution of fresh and old saliva-responding neurons revealed a neuronal population within the

VMH that is more sensitive to fresh saliva than old saliva. Taken together, this study

demonstrates that predator cues in cat saliva change over time and differentially activate the

sensory-to-hypothalamus defensive behavioral pathway to modulate behavioral outputs.

Introduction

Prey animals make defensive behavioral decisions through prompt and careful analysis of their environment when encountering predators. Immediacy of the threat, determined by perceived distance from predators, is one of the key factors influencing defensive behavioral decisions, which regulates the intensity of behavioral outputs (Figure 1) (Blanchard and Blanchard, 1989). Predator-derived sensory signals, or predator cues, are detected through multiple sensory systems, including the visual system, auditory system, and olfactory system. These individual sensory cues are sufficient to induce innate defensive behavior such as fear-related freezing and flight, as well as anxiety-related risk assessment behavior. In nocturnal animals including rodents, olfaction is one of the major sensory modalities by which predator-derived chemical cues are detected (Apfelbach et al., 2005). Rodents respond to a variety of predator-derived chemical cues, such as odors from fox feces, snake skin, cat fur, and cat collars, in both the field and laboratory environments (Apfelbach et al., 2005; Kats and Dill, 1998; Papes et al., 2010; Pérez-Gómez et al., 2015; Takahashi, 2014). However, how animals use chemosensory cues to make appropriate behavioral decisions is not well understood.

The accessory olfactory system is one of the major sensory input channels through which predator-derived chemical cues are detected. It has been shown that some molecules derived from predator species activate sensory neurons in the VNO and induce defensive behavior (Isogai et al., 2011; Papes et al., 2010; Pérez-Gómez et al., 2015; Samuelsen and Meredith, 2009; Tsunoda et al., 2018). Moreover, a number of studies have revealed main components of the sensory neural circuits that mediate behavioral responses to chemical predator cues in rodents (Blanchard et al., 2003; Canteras, 2002; Choi et al., 2005; Dielenberg and McGregor, 2001; Markham et al., 2004; Petrovich et al., 2001). Predator signals detected by the VNO are transferred to the accessory olfactory bulb (AOB). The output neurons in the AOB (the mitral cells) then send the signals to the BNST and the posteroventral part of MeA (MeApv), which connects with the dorsomedial part of the VMH (VMHdm) that contains neural populations directly controlling defensive behavioral outputs (Gross and Canteras, 2012; LeDoux, 2012; Takahashi, 2014). It is, therefore, reasonable to hypothesize that the VNO detects predator cues and sends sensory signals to the VMH, triggering appropriate defensive behavioral decisions in

rodents. However, how the sensory signals detected through the VNO-to-VMH circuitry

modulate behavioral decisions in specific contexts remains elusive.

In this study, we investigated sensory-to-hypothalamic neural circuitries that regulate

defensive behavior induced by non-volatile chemical cues in cat saliva with different freshness.

We examined the role of the VNO in behavioral responses and sought sensory receptor genes for

the predator cues. Moreover, by comparing downstream neural populations activated by cat

saliva with different freshness, we investigated how the imminence of predator threat is

processed in the accessory olfactory circuitry to regulate behavioral outputs.

Results

Predator cues in cat saliva induce context-dependent defensive behavior in mice

To explore whether mice exhibit different intensities of defensive behavior in response to

chemosensory predator cues with varying threat immediacy, we examined their response to cat

saliva with different degrees of freshness, which potentially represents immediacy of the

predator threat. Cat saliva has been considered as a source of predator cues found on cat fur and

collars, which induce defensive behaviors in rodents (Engelke et al., 2021; Papes et al., 2010).

Fresh saliva is assumed to indicate a more immediate threat to the prey animal, which likely

triggers stronger defensive behavior. We exposed mice to a cotton swab containing cat saliva in

their home cage where they know a flight route is not available (Figure 1B). We found that mice

approached and sniffed the cotton swab, regardless of the presence of saliva, as part of their

exploratory behavior towards a novel object in their home cage (Supplemental movies 1-3,

Figure 1C; red boxes indicate the first contact). The latency to the first contact was slightly

longer for saliva collected immediately before introduction to the mouse cage (fresh saliva) when

compared to that for saliva collected a few hours prior to the behavior test (old saliva) (Figure

1D). The difference in latencies was statistically significant, indicating that mice can perceive the

qualitative change of cat saliva over time (Figure 1D). Baseline freezing episodes were observed

in all groups of mice when the sample was introduced into the cage due to the experimenter's

intervention (Figure 1C). However, the groups exposed to cat saliva continued to exhibit freezing

behavior throughout the 10-minute observation period, while mice exposed to a clean swab did

not exhibit freezing behavior after the first contact (Figure 1C, E). Moreover, we found a correlation between the freshness of cat saliva and duration of freezing behavior in mice. Mice exposed to fresh saliva exhibited considerably longer freezing episodes, accounting for over 50% of the total duration of observation, whereas those exposed to old saliva exhibited freezing episodes for less than 20% of the total duration on average (Figure 1E). We found no significant difference in the latency to the first freezing episode from the first contact with a swab between mice exposed to fresh and old saliva (Supplemental Figure 1A). Mice also exhibited stretch sniffing risk assessment behavior, which did not differ toward any of the swab types (Figure 1F, Supplemental Figure 1B). On the contrary, mice exposed to a swab containing either fresh or old saliva significantly avoided interacting with the swab, while mice exposed to a clean control swab spent a significant amount of time directly interacting with the swab to investigate it, including direct sniffing and chewing (Figure 1G). The comparison of temporal behavioral patterns revealed a slightly higher frequency of direct investigation behavior toward old saliva compared to fresh saliva at the beginning of the exposure period (Supplemental Figure 1C). Importantly, the stress hormone adrenocorticotropic hormone (ACTH) in plasma showed a trend towards two-fold upregulation after exposure to both fresh and old saliva (Figure 1H). These results suggest that mice can still recognize predator threat through old saliva, but freezing behavior in response to fresh saliva attenuates quickly over time.

Taken together, this behavior study indicates that cat saliva contains predator cues that signal the imminence of predator threat and differentially induce defensive behavioral outputs in mice. More specifically, fresh saliva is recognized as an immediate danger and induces more robust freezing behavior, while old saliva is recognized as a less immediate danger and induces less robust freezing behavioral responses.

Cat saliva activates VNO neurons and modulates freezing behavior

As the freezing behavior was induced when mice made direct contact with the saliva-containing swab, we postulated that direct contact with the predator cues in cat saliva is necessary for freezing behavior induction. To examine the necessity of direct contact, we placed a cotton swab with or without cat saliva into a porous container and investigated mouse's behavior toward the container. The pores on the container allow volatile chemicals in cat saliva

to diffuse out, while nonvolatile chemicals stay inside of the container with the swab.

Interestingly, we found that when we introduced a container containing a swab with fresh saliva

into the home cage, the mice no longer exhibited freezing behavior towards the container, but

instead directly investigated it similarly to the control swab-containing container (Figure 2A, B).

This suggests that freezing behavior requires direct contact with the saliva-containing swab and

that the freezing-inducing cues in cat saliva detected in this scheme are nonvolatile.

Mice have the ability to detect nonvolatile chemical cues through the VNO (Mohrhardt et

al., 2018). Ethologically relevant chemical cues from different animals, such as pheromones and

predator cues, are sucked up into the VNO and activate VNO sensory neurons when animals

contact the chemical source (Luo et al., 2003). We, therefore, explored the role of the VNO in

inducing freezing in response to cat saliva using a VNO-deficient laboratory mouse line that

lacks the primary signal transduction ion channel in VNO neurons (transient receptor potential

cation channel, subfamily C, member 2-knockout line or Trpc2-KO line) (Leypold et al., 2002).

In Trpc2-KO mice, chemosensory signals detected by sensory receptor proteins are expected to

be attenuated before being relayed to the brain, due to the lack of the downstream transduction

channel in VNO neurons (Leypold et al., 2002; Liman et al., 1999; Stowers et al., 2002). Thus,

Trpc2-KO mice have been widely utilized to examine roles of the VNO in various behaviors

(Beny-Shefer et al., 2017; Ferrero et al., 2013; Haga et al., 2010; Kimchi et al., 2007; Leypold et

al., 2002; Papes et al., 2010; Pérez-Gómez et al., 2015; Stowers et al., 2002). Upon exposure to a

fresh saliva-containing swab, Trpc2-KO mice failed to exhibit freezing behavior, in contrast to

their wild-type littermates who displayed freezing responses. Additionally, the Trpc2-KO mice

directly investigated the saliva swab with a similar level of intensity as the control swab, while

the wild-type littermates actively avoided it (Figure 2C, D). These findings suggest that VNO

activation, disrupted in Trpc2-KO mice, is necessary for the freezing behavior observed in

response to cat saliva in mice.

To investigate if VNO neurons are indeed activated by cat saliva, we analyzed

immunoreactivity (IR) of phosphorylated ribosomal protein S6 subunit, pS6, in the VNO of mice

exposed to cat saliva. As the cellular pS6 level is greatly upregulated upon activation of VNO

neurons, it has been used as a sensitive marker to examine neural activation in the VNO

(Carvalho et al., 2020; Itakura et al., 2022; Osakada et al., 2022; Tsunoda et al., 2018). We

observed that exposure to both fresh and old cat saliva induced an increase in pS6-IR in the basal

VNO neurons that express Gαo protein (Figure 2E, G). Conversely, pS6-IR was rarely observed

in the apical VNO neurons (Figure 2F). Notably, we found that the number of pS6-positive

neurons was more than three times higher in mice exposed to fresh saliva compared to that

exposed to old saliva (Figure 2G). Strikingly, when the duration of freezing was plotted as a

function of pS6-IR signals in the VNO of individual animals, we observed a significantly

positive correlation (R^2 = 0.3586, 95% CI [0.2458, 0.8116], p = 0.0020) between the numbers of

active sensory neurons and behavioral outputs (Figure 2H), suggesting that more intense freezing

behavior is induced when more VNO neurons are activated by cat saliva.

Taken together, these results indicate that freezing-inducing predator cues are detected by

Gαo-positive sensory neurons in the VNO, and these sensory neurons convey the imminence of

predator threat to higher brain centers.

Cat saliva activates the V2R-A4 subfamily of the vomeronasal receptors

The significant reduction of the freezing duration to aged saliva reflects possible

attenuation of signal inputs to a specific population of sensory receptors, or alternatively, it may

reflect a change of sensory receptor repertoires due to qualitative changes of predator cues over

time. To test these possibilities, we examined sensory receptors expressed in the neurons

activated by fresh and old saliva. The basal VNO neurons express sensory receptors that belong

to vomeronasal receptor type 2 (V2R) family of G protein-coupled receptors in mice (Figure 3A)

(Mohrhardt et al., 2018). The mouse V2R family consists of over 120 intact genes, which are

divided into V2R A1-6, A8-10, and B-D subfamilies based on sequence homology (Figure 3A)

(Francia et al., 2014; Silvotti et al., 2011, 2007). With the exception of the V2R C group that is

co-expressed with other groups of V2Rs in the basal VNO neurons, each sensory neuron

expresses only one member of the V2R receptor, allowing individual neurons to respond to

specific molecules (Mohrhardt et al., 2018).

To investigate which subfamilies of V2R are expressed in cat saliva-activated VNO

neurons, we designed *in situ* hybridization (ISH) probes that detect individual V2R subfamilies

and performed dual staining of V2R ISH with pS6-IR (Supplemental Figure 2). We found that

the majority of pS6-IR signals induced by cat saliva co-localized with signals of the ISH probe

against the V2R-A4 subfamily (Figure 3B, Supplemental Figure 2). Approximately 45% of pS6-

IR signals induced by fresh saliva (83 out of 181) co-localized with V2R-A4 probe signals, with

less than 1 % of pS6-IR (1 out of 120) co-localized with A8-4 probe signals (Figure 3C).

Notably, V2R-A4 probe signals also co-localized with approximately 45% of pS6-IR signals

induced by old saliva (37 out of 86). None of the other probe signals co-localized with pS6-IR

induced by both fresh and old saliva. These results indicate that members of the V2R-A4

subfamily are potential sensory receptors for the predator cues contained in both fresh and old

cat saliva, and varying levels of imminence of predator threat stimulate the same type of sensory

receptors in a quantitatively different manner to regulate the freezing behavior.

Cat saliva activates neurons in the AOB and defensive behavioral circuit

We next aimed to understand how the differential activation of sensory receptor neurons

in the VNO by different freshness of cat saliva influences behavioral decisions through

downstream neural circuits (Figure 4A). We investigated this by examining the induction of

cFos, an immediate early gene product, in the projection neurons (mitral cells) and local

inhibitory neurons (granule cells) in the AOB of mice exposed to cat saliva. The axons of apical

and basal VNO neurons project to the anterior and posterior regions of the AOB (aAOB and

pAOB), respectively (Mohrhardt et al., 2018). We found that the activation of the basal VNO

neurons by cat saliva leads to predominant induction of cFos-IR signals in the pAOB, whose

glomerulus layer is Gαi-IR negative (Figure 4B-F). The numbers of cFos-positive mitral and

granule cells in the pAOB in saliva-exposed mice are significantly higher than those in mice

exposed to the control swab (Figure 4D, F). However, contrary to our expectations, we did not

observe any differences in the numbers of cFos-positive mitral and granule cells in the AOB

between fresh and old saliva-exposed mice (Figure 4D, F). These results suggest that, although

the AOB processes saliva signals from the VNO following predator cue detection, the

relationship between the degree of AOB activation and freezing behavior intensity is non-linear.

To further investigate how signals of fresh and old saliva are processed in the

downstream circuitry, we examined induction of cFos in the neural substrates involved in innate

defensive behaviors (Figure 4A). We investigated the BNST and MeApv, as these nuclei receive

direct ascending inputs from the mitral cells in the AOB (Mohrhardt et al., 2018). We also investigated the VMH and dPAG, as these nuclei have been shown to control defensive behavioral outputs (Silva et al., 2013). Upon exposure to fresh and old saliva, cFos expression in the MeApv, VMH, and dPAG, but not in the BNST, appeared to be upregulated compared to the control stimulus (Figure 4G-J). Notably, cFos expression was significantly increased in the VMH of mice exposed to fresh and old saliva compared to the control (Figure 4I), consistent with previous observations that the dorsomedial and central subdivisions of VMH (VMHdm/c) contain subpopulations of neurons that control freezing behavior (Engelke et al., 2021; Kunwar et al., 2015; Silva et al., 2016, 2013; Wang et al., 2015) However, there was no difference in the number of cFos-positive cells between the two saliva-exposed groups in the neural substrates tested (Figure 4G-J). These results suggest that, although the defensive circuit is activated by cat saliva signals, the relationship between the degree of activation and freezing behavior intensity is non-linear, similar to the situation in the AOB.

Taken together, these results indicate that the relationship between neural activity and behavioral output is more complex in the downstream circuit than in the VNO. More specifically, the numbers of active neurons and intensity of freezing behavior do not appear to correlate in the downstream regions. This suggests that fresh and old cat saliva may differentially activate functionally distinct neural populations within these regions, which in turn leads to quantitatively different freezing behavioral responses.

Fresh and old cat saliva differentially activate VMH neurons to control defensive behavior

To investigate the potential differences between fresh and old saliva-activated neurons, we analyzed correlations of the duration of freezing behavior and numbers of cFos-IR signals in different cell types in individual animals from different exposure groups. To our surprise, we found that the numbers of cFos-IR signals in AOB mitral and granule cells, as well as those in neurons in the BNST and dPAG, did not correlate with the duration of freezing behavior in any of the exposure groups (Figure 5A-C, F). However, we observed a significant positive correlation between the number of cFos-IR signals and behavior intensity only in the fresh saliva-exposed group in the VMH ($R^2 = 0.5708$, 95% CI [-0.1449, 0.9714], $p = 0.0412$) while no such correlation was observed in the old saliva-exposed group (Figure 5E). We observed a

similar positive correlation trend in the MeApv ($R^2 = 0.3854$, 95% CI [-0.3845, 0.9525], $p = 0.0942$), although it was not statistically significant possibly due to low sample numbers (Figure 5D). Taken together, these findings support our hypothesis that fresh and old saliva may differently activate the downstream circuit.

We therefore examined whether fresh and old cat saliva activate neurons in spatially distinct regions within the VMH. We observed the distribution of cFos-IR neurons in five serial brain sections from rostral to caudal VMHdm regions. The VMH region within each hemisphere section was further divided into 10 bins along the dorsomedial-ventrolateral axis, and cFos-IR signals in each bin were counted (Figure 6A). The majority of cFos-IR signals induced by both fresh and old cat saliva were localized in the dorsomedial to central part of the VMH (bins 1 – 5). Strikingly, the numbers of cFos-IR signals induced by fresh and old saliva were significantly different in the section 2 bin 5, as well as the section 3 bin 3 and bin 4, which are adjacent to each other (Figure 6B). The cFos-IR signals in the other regions did not show significant difference between fresh and old saliva-exposed mice. These results suggest that there is a distinct population of neurons clustered in a close proximity within the VMH, which is more sensitive to fresh saliva than old saliva. This population of neurons in the VMHdm may play a role in inducing robust freezing behavior to fresh cat saliva. Moreover, these results also suggest that the neurons that induce robust freezing to fresh saliva and less robust freezing to old saliva reside in the same region of the VMH and may use different mechanisms to encode freezing intensity. Overall, our results suggest that the qualitative properties of predator signal inputs, such as their freshness, can induce distinct neural processing and ultimately lead to different defensive behavioral outcomes.

Discussion

This study provides evidence for the role of cat saliva as a source of predator cues that signal imminent predator threat and regulate the intensity of freezing behavior through the accessory olfactory pathway in mice (Figure 7). Notably, both fresh and old saliva activate neurons expressing the V2R-A4 subfamily of sensory receptors, suggesting the existence of a specific receptor group responding to the predator cues. Importantly, the number of vomeronasal neurons activated in response to saliva correlates with the freshness of saliva and the intensity of

freezing behavior. Moreover, the number of VMH neurons activated by immediate threat (fresh saliva) exhibits a positive correlation with the intensity of behavioral outputs, suggesting their direct involvement in the modulation of freezing behavior. In contrast, the activation of VMH neurons by less immediate threat (old saliva) does not show such a correlation. These contrasting responses to different freshness levels of cat saliva suggest the presence of distinct populations of neurons in the VMH controlling the intensity of freezing behavior. These findings establish a framework for understanding the neural mechanisms underlying innate defensive behavior decisions.

The VNO as the sensor of predator cues that induce fear-related behavior

This study demonstrates that the VNO is involved in detecting and processing chemosensory predator signals that induce fear-related freezing behavior. Previous studies have shown that the VNO mediates defensive behaviors against predators that are rather related to anxiety state, such as risk assessment, area avoidance, and reduced exploratory and locomotion behaviors (Isogai et al., 2011; Papes et al., 2010; Pérez-Gómez et al., 2015; Tsunoda et al., 2018). The difference between these studies and our current study is likely due to different sample types and/or their freshness, as we observed that fear-related behavior is correlated with freshness of predator samples. It is interesting to assume that the biological sources of predator cues, such as saliva, urine, and tears, contain varieties of unidentified cues that convey time-sensitive signals through the mouse accessory olfactory system. This is consistent with the fact that VNO neurons expressing more than half of vomeronasal receptors and a significant fraction of AOB mitral cells have been shown to be activated by predator-derived samples (Ben-Shaul et al., 2010; Isogai et al., 2011). Therefore, it is important to investigate how the sensory pathway in the accessory olfactory system processes varieties of predator cues, which eventually leads to defensive behavioral decisions in higher brain centers.

Moreover, our study demonstrates that time-dependent alterations of the chemosensory predator signals can be detected by the VNO, which induces altered defensive behavioral responses. For instance, 4-hour-old saliva induces reduced freezing behavior compared to fresh saliva. On the other hand, duration of direct investigation remains low toward both fresh and old saliva, and the stress hormone ACTH level is upregulated in both cases. A future study ought to

identify specific molecules—most likely proteins or peptides—in cat saliva responsible for inducing these distinct defensive responses in mice. A cat lipocalin allergen, Fel d 4, has been identified to induce risk assessment defensive behavior in mice (Papes et al., 2010). While Fel d 4 is one of the potential candidates, there exists a possibility of a different molecule or a combination of multiple molecules playing a role. Once the molecules are identified, it is imperative to investigate how their quantity and quality change over time and how these factors correlate with freezing behavior. Such an exploration will provide insights into how mice sense the imminence of threat using those chemical signals.

V2R-A4 subfamily as the receptor for predator cues in cat saliva

Screening of sensory receptors in the VNO neurons activated by fresh and old saliva demonstrated that members of the V2R-A4 subfamily are the candidate receptors for the predator cues in cat saliva. The V2R-A4 subfamily has been shown to be expressed in the VNO neurons that are activated by biological samples derived from various mammalian predators (Carvalho et al., 2020; Isogai et al., 2011; Tsunoda et al., 2018). Therefore, it is likely that specific members of the V2R-A4 subfamily are the receptors for the predator cues in cat saliva that induce freezing behavior.

Dual staining of vomeronasal receptors with immediate early gene products has been effectively utilized to identify receptors for several pheromones (Carvalho et al., 2020; Haga et al., 2010; Isogai et al., 2018, 2011; Itakura et al., 2022; Kimoto et al., 2005; Osakada et al., 2022, 2018; Tsunoda et al., 2018). However, due to very high sequence homologies among members of the V2R-A4 subfamily (Francia et al., 2014), alternative approaches need to be considered to further isolate the receptors responsible for the detection of predator cues in cat saliva. Potential approaches include receptor mRNA profiling with pS6 immunoprecipitation from the VNO of mice stimulated by cat saliva (Isogai et al., 2018; Jiang et al., 2015), as well as receptor mRNA profiling from isolated single cells activated by cat saliva in GcaMP imaging using the VNO slices *in vitro* (Haga-Yamanaka et al., 2014; Wong et al., 2020). Receptor candidates identified using either of the methods can be further confirmed by examining necessity and sufficiency for detecting cat saliva using genetically modified mouse lines (Haga et al., 2010; Haga-Yamanaka

et al., 2014; Isogai et al., 2018; Itakura et al., 2022; Osakada et al., 2022, 2018; Tsunoda et al., 2018).

Differential processing of fresh and old saliva signals in the VNO-to-VMH pathway

Our correlation analyses between the number of activated neurons and behavior intensity suggest that sensory signals encoding different imminence of predator threat are differentially processed in the VNO-to-VMH pathway. The differential behavioral responses do not appear to result from global neuromodulations of the neural circuit by stress hormones, as differential endocrinological responses were not observed (Figure 1H). This suggests that more specific neural processing happens in the circuitry. In the circuitry downstream of the VNO, sensory signals can converge and diverge from the AOB before reaching the VMH. The axon terminals of hundreds of VNO neurons expressing the same receptor converge into a few specific glomerulus structures (Belluscio et al., 1999; Del Punta et al., 2002; Rodriguez et al., 1999), and, in some cases, neurons expressing receptors in the same subfamily send axons into the same glomeruli (Wagner et al., 2006). Individual AOB mitral cells extend multiple primary dendrites into multiple glomeruli (Imamura et al., 2020; Larriva-Sahd, 2008; Takami and Graziadei, 1991; Wagner et al., 2006; Yonekura and Yokoi, 2008). In turn, efferent axons of the mitral cells project to multiple brain substrates, namely the BNST, the MEA, and the posteromedial cortical amygdaloid nucleus (PMCo) (Davis et al., 1978; von Campenhausen and Mori, 2000). These structural features indicate that sensory inputs to the AOB mitral cells are converged, and outputs from the mitral cells may be diverged.

These features of the ascending neural pathway within the AOB may contribute to the sorting of sensory signals derived from fresh and old saliva. Fresh saliva activates more V2R-A4-expressing VNO neurons than old saliva, which may result in stronger activation of V2R-A4 glomeruli in fresh saliva-exposed mice compared to those exposed to old saliva. This differential activation of V2R-A4 glomeruli by fresh and old saliva may lead to activation of the mitral cells innervating the V2R-A4 glomeruli with different excitation levels. We assume that such differential activations of the mitral cells between fresh and old saliva result in the differential activation of targeting neural substrates, possibly MeApv, which results in differential activation of VMH neurons (Figure 7). The methodology that we undertook in this study, cFos-

immunohistochemistry, only provides information about neurons responding to cat saliva above a given threshold. It is therefore important to examine how the mitral cells respond to fresh and old saliva using more quantitative techniques, such as *in vivo* electrophysiological recording, in future studies.

Differential activation of VMH neurons potentially underlying distinct intensities of freezing behavior

Our findings indicate that the imminent predator cues present in cat saliva activate a specific subpopulation of VMH neurons, whose overall activity positively correlates with the intensity of freezing behavioral outputs (Figure 5E). In contrast, the less imminent predator cues activate a subpopulation of VMHdm/c neurons, whose overall activity does not correlate with the intensity of freezing behavior (Figure 5E). These findings suggest that induction of robust freezing behavior is determined by a distinct set of VMH neurons, and the intensity of freezing behavior is encoded separately, possibly by the intensity or dynamics of the neuronal firing, which cannot be detected by the cFos-IR used in this study. The VMHdm/c region contains several subpopulations of neurons that coordinate various defensive behavioral responses (Engelke et al., 2021; Esteban Masferrer et al., 2020; Kennedy et al., 2020; Silva et al., 2016; Tobias et al., 2023). Previous studies have demonstrated that optogenetic stimulation of VMHdm/c neurons induces freezing/immobility responses, suggesting that these neurons mediate freezing behavior in response to predator cues (Kunwar et al., 2015; Wang et al., 2015). In our current study, the distribution of cFos-IR in the VMH revealed a population of neurons clustered in close proximity that are more sensitive to fresh saliva than old saliva (Figure 6B). Therefore, we propose that imminent predator cues in cat saliva may activate these specific neurons with certain excitation levels to directly induce robust freezing behavior, while less imminent predator cues in cat saliva may use a different mechanism to regulate freezing behavior.

What distinguishes the subpopulation of VMHdm/c neurons activated by fresh saliva from those activated by old saliva? One possibility is that these neurons stimulate distinct downstream neural circuitries. The VMHdm/c neurons project to two major nuclei, the anterior hypothalamic nucleus (AHN) and the periaqueductal gray (PAG), which directly influence

different aspects of defensive behavior (Canteras et al., 1994; Wang et al., 2015). Specifically, selective activation of the VMHdm/c-to-PAG pathway has been shown to cause freezing/immobility behavior, while activation of the VMHdm/c-to-AHN pathway rather promotes avoidance and induces immobility to a lesser extent (Wang et al., 2015). Although the number of cFos-positive cells in the dPAG did not seem to correlate with the intensity of freezing behavior in any saliva exposure group in our study (Figure 5F), it is possible that fresh saliva-activated neurons send inputs to the PAG, whereas old saliva-activated neurons send inputs to the AHN and/or other projection sites. Thus, the number of old saliva-activated cells in the VMH may correlate with different aspects of defensive responses, rather than freezing behavior.

Given the observation from a previous study that neurons projecting to the PAG also send collateral projections to the AHN, and vice versa, neuronal populations activated by fresh and old saliva may or may not overlap. If the two sets of neurons overlap, it is possible that the strength of presynaptic inputs to the VMHdm/c differs between fresh and old saliva stimulation, which leads to differential activation of downstream pathways. Previous studies have shown that optogenetic stimulation of the VMHdm/c with different stimulus frequencies can induce different types of defensive behaviors (Wang et al., 2015), suggesting that distinct activity patterns of fresh and old saliva-responsive neurons in the VMHdm/c may contribute to differences in defensive responses. Investigating these activity patterns would be an intriguing direction for future research.

In summary, our study proposes a predator defensive circuitry that involves the accessory olfactory system and the VMHdm/c. Understanding the precise neural mechanisms of how different degrees of imminence of predator threat are processed in this circuit will provide valuable insights into the decision-making processes underlying defensive behaviors in animals.

Materials and methods

Animals

The experimental procedures were approved by the Institutional Animal Care and Use Committee at the University of California, Riverside (UCR) and were conducted in accordance with the National Institutes of Health Guide for the Care and Use of Laboratory Animals. C57BL/6N mice originally obtained from Charles River (Catalog# 027) were bred at the UCR vivarium. The Trpc2 knock-out mouse line was generously provided by Dr. Richard Axel Lab at Columbia University. All mice were housed in the vivarium under a reversed light cycle (lights on at 21:00; lights off at 9:00) for a minimum of 2 weeks prior to the behavior and histology experiments. Cats used in this study were obtained from the Nutrition and Pet Care Center at the University of California, Davis (a class A dealer) and were also housed in the UCR vivarium.

Collection of saliva from cats

A person familiar with the cat gently held it for all procedures. A sterile cotton swab was inserted into the cat's mouth where the cat chewed and licked on the swab for at least 20 seconds or until the swab is saturated with saliva. The swabs containing saliva were then placed in microcentrifuge tubes and stored until used. Fresh saliva was collected within 15 minutes prior to its use in the mouse behavioral study, while old saliva was collected 4 hours prior to the study. The old saliva was kept at room temperature until it was used.

Behavioral assay

All behavioral experiments were conducted during the animals' dark cycle. Mice were placed in their experimental room and allowed approximately 4 hours to acclimate to the ambient conditions. A cat saliva-containing swab was manually introduced into the animal's home cage to assess the defensive behavioral responses of resident mice. Recording of mouse behaviors commenced 2 minutes prior to introducing the sample and concluded 10 minutes thereafter. If tissues were required for histology, the animals were euthanized after 60 minutes, while for plasma collection, euthanasia took place after 15 minutes. Following the introduction of the sample, the behavioral room remained vacant of any individuals until the animal was euthanized. Mouse behavior was captured by night vision cameras equipped with infrared lights.

Behavioral analysis

Recordings of each behavioral session were saved and provided in a blinded manner to a researcher for analysis. The analysis was performed using a behavioral analysis program called Behavioral Observation Research Interactive Software (BORIS) (Friard and Gamba, 2016). Freezing behavior was defined as immobility lasting more than 2 seconds and was scored accordingly. Stretch sniffing risk assessment behavior was scored when animals exhibited flat-back/stretch-attend posture. Direct investigation was scored when there was physical contact with the swab. Animals that exhibited inactivity in their home cage for more than 50% of the pre-drop analysis duration in each experiment were excluded from the data set.

Histology and Immunohistochemistry (IHC)

After being exposed to the cat specimen or control swab for 60 minutes, the animals were euthanized with an overdose of sodium pentobarbital. Intracardial perfusions were performed using 4% paraformaldehyde (PFA) in PBS. The brains were extracted and post-fixed in 4% PFA overnight at 4°C. Subsequently, the brains were cryoprotected by sequentially immersing them in 15% and 30% sucrose solutions in PBS. The VNOs were also extracted and underwent the same post-fixation procedure, followed by decalcification with EDTA pH 8.0. Cryoprotection of the VNOs was achieved using a 30% sucrose solution in PBS. The cryoprotected tissues were then embedded in OCT (Optimal Cutting Temperature) media and frozen over liquid nitrogen. Tissues embedded in OCT were sectioned at 14 μm (VNO) and 20 μm (brain) thicknesses using a cryostat set at -25°C. The sections were placed onto positively charged microscope slides. The slides were stored at -80°C until stained. A standard frozen IHC protocol was employed. For VNO sections, staining was performed using a rabbit anti-pS6 antibody (Cell Signaling Technology, #4858) at a dilution of 1:200, incubated overnight at 4°C, followed by a secondary antibody, Goat anti-rabbit IgG Alexa Fluor 647 (Life Technologies #a21245), for 60 minutes at room temperature. For AOB sections, staining was carried out using a mouse anti-Gαi2 antibody (Millipore #MAB3077MI) and a guinea pig anti-cFos antibody (Synaptic Systems #226 308) at dilutions of 1:500 and 1:1000, respectively. This was followed by incubation with a goat anti-

mouse IgG2b Alexa Fluor 546 (Life Technologies #a21143) and a Goat anti-guinea pig IgG

Alexa Fluor 647. For brain sections, staining was performed using a guinea pig anti-cFos

antibody at dilutions of 1:1000 incubated overnight at 4°C, followed by Goat anti-guinea pig IgG

Alexa Fluor 647. Finally, the slides were mounted using MS Shield Mounting Medium With 4,6-

diamidino-2-phenylindole (DAPI) and DABCO™ (EMS #17989-20).

Enzyme-linked Immunosorbent Assay (ELISA)

Plasma collected from mice exposed to cat saliva or control swabs was frozen at -80°C

before initiating the ELISA protocol. The ACTH ELISA kit (MD Bioproducts) was utilized,

following the instructions provided with the kit.

In situ hybridization (ISH) and dual staining

Detection of *GαO* RNA in fixed frozen VNO sections was conducted using the ACD

RNAscope® probe targeting GNAO1 (GαO) (ACD #444991) and the RNAscope® Multiplex

Fluorescent Reagent Kit v2 (ACD #323100). Probe binding was visualized using Akoya

Biosciences' Opal 570 (FP1488001KT) Dye at a dilution of 1:750 in RNAscope TSA Buffer.

Following RNAscope staining, the slides were incubated in anti-pS6 antibody at 1:200 dilution

overnight at 4°C, then detected by goat anti-rabbit IgG AlexaFluor 647 for one hour at room

temperature.

Vmn2r (V2R) probes for the following genes were used: *Vmn2r13* for subfamily A1,

Vmn2r118 for A2, *Vmn2r117* for A3, *Vmn2r42* for A4, *Vmn2r69* for A5-1, *Vmn2r65* for A5-2,

Vmn2r76 for A5-3, *Vmn2r120* for A6, *Vmn2r98* for A8-1, *Vmn2r105* for A8-2, *Vmn2r58* for A8-

3, *Vmn2r62* for A8-4, *Vmn2r81* for A9, *Vmn2r18* for A10, *Vmn2r24* for B, and *Vmn2r53* for D.

Each cRNA probe was synthesized using the PCR-based method (Hua et al., 2018).

For detection of *V2R* RNA in ISH, VNO tissues were sectioned and placed onto slides up

to one week before ISH and stored at -80 °C. Fluorescent ISH was performed following

published protocols (Ishii et al., 2004). Briefly, the VNO slides were baked at 60 °C for 5

minutes and fixed with 4% PFA. The slides were then treated with 10 μg/mL Proteinase K

(Sigma #3115828001), followed by being acetylated in 0.1 M Triethanolamine (Sigma #90279)-

HCl with acetic anhydride (Sigma #320102). *V2R* cRNA probes were hybridized with tissue

sections overnight at 65 ℃ in a hybridization solution consisting of 50 % formamide, 10 mM

Tris-HCl pH 8.0, 200 µg/mL yeast tRNA, 10 % dextran sulfate, 1x Denhardt's, 600 mM NaCl,

0.25 % SDS, 1 mM EDTA pH 8.0. After hybridization, the slides were washed with 2x SSC in

50 % formamide for 30 minutes, then 2x SSC, 0.2x SSC, and 0.1x SSC in H_2O for 20 minutes

each at 65℃. The slides were then blocked with blocking reagent (Perkin Elmer) and 0.5 %

peroxidase block sequentially. Anti-DIG-POD (Millipore Sigma #11207733910) was used to

detect DIG-labeled probes. TSA-Cy3 Plus kit (Akoya #NEL744001KT) was used to detect anti-

DIG antibody at 1:200. The slides were incubated with anti-pS6 antibody at 1:200 dilution

overnight at 4℃, then detected by goat anti-rabbit IgG AlexaFluor 647 for one hour at room

temperature.

Following staining, the slides were mounted with MS Shield Mounting Medium with

DAPI and DABCO™.

Image Quantification

Fluorescent images were captured at either 10X or 20X magnification using the Zeiss

Axio Imager.M2 microscope. Whole slide images of IHC-stained brains were scanned through

fluorescent imaging using Pannoramic SCAN (3D Histech) at Reveal Biosciences (San Diego,

CA). Automated counting of fluorescent positive cells was performed using a proprietary code

developed in our laboratory within the QuPath software.

VMH cFos IR Quantification and Distribution Analysis

All VMH regions were identified by clusters of nuclei visualized through DAPI staining

and manually annotated in QuPath using standard coronal mouse brain atlas as reference.

Annotations were processed with a proprietary script for immunoreactivity quantification.

Distribution analyses were performed in QuPath using a rectangular annotation set parallel to the

dorsomedial-ventrolateral axis of each VMH region. This rectangular annotation was then split

into 10 equal bins, and immunoreactivity quantification in each bin was automated in QuPath, where the detections were finally manually recorded.

Statistical Analysis

All statistical analyses were performed using Prism software (GraphPad). Behavioral analysis results and cFos-IR distribution in the VMH, involving two variables, were compared using a traditional Two-Way ANOVA with Tukey's multiple comparisons. Behavioral and immunohistological results were statistically compared using One-Way ANOVA with Tukey's multiple comparisons, assuming a parametric normal distribution. The correlation between immunoreactivity and behavior was analyzed using either Spearman's rank correlation coefficient or Pearson correlation coefficient: for data points that did not pass Shapiro-Wilk and Kolmogorov-Smirnov normality tests, Spearman's rank correlation coefficient was used, while for data points that pass Shapiro-Wilk and Kolmogorov-Smirnov normality tests, Pearson correlation coefficient was used.

Acknowledgments

We appreciate the generosity of Dr. Michael Rust for providing a vivarium cat colony and Drs C. Ron Yu, and Richard Axel for providing Trpc2 knock-out mice. We thank Drs C. Ron Yu, Jing Wang, Hongdian Yang, Martin M. Riccomagno, Edward Zagha, Takuya Osakada, and Naoki Yamanaka for helpful discussion and comments on the manuscript. We are grateful for technical assistance from Office of Campus Veterinarian at University of California, Riverside (UCR). This study was supported by funds from the National Institutes of Health (R01DC019135) and startup funds from UCR to S.H.-Y, and a fellowship from Graduate Assistance in Areas of National Need to Q.A.N.

Author contributions

Conceptualization, S.H.-Y.; Methodology, Q.A.N., C.P., and S.H.-Y.; Investigation, Q.A.N., A.R., Y.Y., R.C., and C.S., and S.H.-Y.; Writing – Original Draft, Q.A.N and S.H.-Y.;

Writing – Review & Editing, Q.A.N and S.H.-Y.; Supervision, S.H.-Y.; Funding Acquisition, Q.A.N and S.H.-Y.

Declaration of interests

The authors declare no competing interests.

Figure legends

Figure 1 Predator cues in cat saliva induce context-dependent defensive behavior in mice

(**A**) A schematic diagram illustrating the elicitation of "freeze" or "flight" defensive behaviors in relation to distance. (Adapted with permission from (Blanchard and Blanchard, 1989))

(**B**) A schematic illustration depicting a test chamber for evaluating defensive behaviors.

(**C**) Raster plots displaying freezing episodes of individual mice exposed to a clean control swab (gray, n = 18), fresh saliva (green, n = 18), or old saliva (yellow, n = 15). The introduction of a clean, fresh, or old saliva swab into a mouse's cage is denoted as time 0. Red lines indicate the first contact with a swab for each mouse.

(**D**) Latency to the first episode of contact with a swab. One-way ANOVA (F $(2,47)$ = 5.409, p = 0.0077) with Tukey's multiple comparisons test (Ctrl vs. Fresh, p = 0.0640; Ctrl vs Old, p = 0.5932; and Fresh vs Old, p = 0.0074).

(**E**) The percentage of total freezing episodes after the first contact. One-way ANOVA (F $(2,47)$ = 37.38, $p < 0.0001$) with Tukey's multiple comparisons test (Ctrl vs. Fresh, $p < 0.0001$; Ctrl vs Old, p = 0.0411; and Fresh vs Old, $p < 0.0001$).

(**F**) The percentage of total stretch sniff episodes. One-way ANOVA (F$(2,49)$ = 0.007016, p = 0.9930).

(**G**) The percentage of total direct investigation episodes. One-way ANOVA (F $(2,48)$ = 25.17, $p < 0.0001$) with Tukey's multiple comparisons test (Ctrl vs. Fresh, $p < 0.0001$; Ctrl vs Old, $p < 0.0001$; and Fresh vs Old, p = 0.5762).

(**H**) Plasma ACTH concentrations following exposure to clean (n = 10), fresh (n = 8), or old saliva (n = 10) swabs. One-way ANOVA (F (2,25) = 3.735, p = 0.0381) with Tukey's multiple comparisons test (Ctrl vs. Fresh, p = 0.0370; Ctrl vs Old, p = 0.1609; and Fresh vs Old, p = 0.6804).

In (**D**) - (**H**), the values are presented as means ± S.E.M., with individual dots representing individual mice. *, **, and **** denote significance levels of $p < 0.05$, 0.01, and 0.0001, respectively.

Figure 2 Cat saliva activates VNO neurons and modulates freezing behavior

(**A**) The percentage of total freezing episodes directed towards control (n = 12) or fresh saliva (n = 11) swabs in cassettes that prevent direct contact with the swabs. Two-tailed t test (t = 0.08745, df = 21, p = 0.9311).

(**B**) The percentage of total investigation episodes toward cassettes containing control or fresh saliva swabs. Two-tailed t test (t=1.662, df=21, p = 0.1114).

(**C**) The percentage of total freezing episodes directed toward control or fresh saliva swabs in Trpc2-WT (n = 10) or -KO (n = 9) mice. Two-way ANOVA ([genotype] F (1, 34) = 3.662, p = 0.0641, [swab type] F (1, 34) = 8.053, p = 0.0076, [genotype] and [swab type] F (1, 34) = 5.963, p = 0.0200) with Tukey's multiple comparisons test (WT-Ctrl vs KO-Ctrl, p = 0.9922; WT-Ctrl vs WT-Saliva, p = 0.0164; WT-Ctrl vs KO-Saliva, p = 0.9136; KO-Ctrl vs WT-Saliva, p = 0.0100; KO-Ctrl vs KO-Saliva, p = 0.9832; WT-Saliva vs KO-Saliva, p = 0.0037).

(**D**) The percentage of total direct investigation episodes toward control or fresh saliva swabs in Trpc2-WT or -KO mice. Two-way ANOVA ([genotype] F (1, 34) = 11.46, p = 0.0018, [swab type] F (1, 34) = 7.725, p = 0.0088, [genotype] and [swab type] F (1, 34) = 13.96, p = 0.0007) with Tukey's multiple comparisons test (WT-Ctrl vs KO-Ctrl, p = 0.9051; WT-Ctrl vs WT-Saliva, $p < 0.0001$; WT-Ctrl vs KO-Saliva, p = 0.9732; KO-Ctrl vs WT-Saliva, p = 0.0006; KO-Ctrl vs KO-Saliva, p = 0.9949; WT-Saliva vs KO-Saliva, p = 0.0003).

(**E**) Representative images illustrating the expression of pS6 (green) and Gα$_o$ (magenta) in the VNO of mice exposed to control, fresh, or old saliva swabs. DAPI was used as a nuclear counterstain. Scale bar: 50 μm.

(**F, G**) The number of pS6-IR positive cells in 1000 VNO cells in the apical (**F**) and basal (**G**) regions of the VNO neuroepithelium in mice stimulated with control (n = 8), fresh saliva (n = 8), and old saliva (n = 8) swabs. (**F**) One-way ANOVA (F $(2, 21)$ = 2.562, p = 0.1010). (**G**) One-way ANOVA (F $(2, 21)$ = 17.90, $p < 0.0001$) with Tukey's multiple comparisons test (Ctrl vs. Fresh, $p < 0.0001$; Ctrl vs Old, $p = 0.2262$; and Fresh vs Old, $p = 0.0014$).

(**H**) The percentage of total freezing episodes in relation to the number of pS6-positive cells in the VNO of individual mice (n = 8 per swab type). Correlation analysis was conducted using Spearman's rank correlation coefficient as total data points did not pass Shapiro-Wilk and Kolmogorov-Smirnov normality tests. Statistical results are reported in the graph.

(**A - D, F, G**) the values are presented as means ± S.E.M., with individual dots representing individual mice. *, **, ***, and **** denote significance levels of $p < 0.05, 0.01, 0.001,$ and 0.0001, respectively.

Figure 3 Cat saliva activates the V2R-A4 subfamily of the vomeronasal receptors

(**A**) Phylogenetic tree of the V2R family and its subfamilies used for V2R cRNA probe design.

(**B**) Representative images displaying the expression of *V2RA4* (magenta) and pS6 (green) in the VNO of a mouse exposed to a fresh saliva swab. DAPI was utilized as a nuclear counterstain. Scale bar: 50 μm.

(**C**) The percentage of cells co-labeled with V2R probes and anti-pS6 antibody among the total pS6-positive cells in the VNO of mice exposed to fresh saliva (top, from 3 – 6 animals) or old saliva (bottom, from 3 – 6 animals).

Figure 4 Cat saliva activates neurons in the AOB and defensive behavioral circuit

(**A**) Schematic diagrams illustrating the VNO sensory pathway that controls defensive behaviors.

(**B**) Representative images displaying the expression of cFos (magenta) in the mitral cell (MCL) and granule cell (GCL) layers of the AOB of mice exposed to control, fresh, or old saliva swabs. $G\alpha_{i2}$ (green) visualizes the anterior region of the glomerular layer (GL). DAPI was utilized as a nuclear counterstain. Scale bar: 200 μm.

(**C, D**) The number of cFos-IR positive mitral cells per 1000 mitral cells in the anterior (**C**) and posterior (**D**) regions of the AOB (n = 10 per swab type). (**C**) One-way ANOVA (F (2, 27) = 0.6106, p = 0.5504). (**D**) One-way ANOVA (F (2, 27) = 6.665, p = 0.0044) with Tukey's multiple comparisons test (Ctrl vs. Fresh, p = 0.0210; Ctrl vs Old, p = 0.0059; and Fresh vs Old, p = 0.8593).

(**E, F**) The number of cFos-IR positive granule cells per 1000 granule cells in the anterior (**E**) and posterior (**F**) regions of the AOB (n = 10 per swab type). (**E**) One-way ANOVA (F (2, 27) = 0.5831, p = 0.5651). (**F**) One-way ANOVA F (2, 27) = 5.996, p = 0.0070) with Tukey's multiple comparisons test (Ctrl vs. Fresh, p = 0.0262; Ctrl vs Old, p = 0.0099; and Fresh vs Old, p = 0.9105).

(**G**) The number of cFos-IR positive neurons per 1000 cells in the BNST in mice stimulated with control (n = 10), fresh saliva (n = 5), and old saliva (n = 6) swabs. One-way ANOVA (F (2, 18) = 0.4375, p = 0.6524).

(**H**) The number of cFos-IR positive neurons per 1000 cells in the MeApv in mice stimulated with control (n = 9), fresh saliva (n = 6), and old saliva (n = 7) swabs. One-way ANOVA (F (2, 19) = 2.454, p = 0.1128).

(**I**) The number of cFos-IR positive neurons per 1000 cells in the VMH in mice stimulated with control (n = 6), fresh saliva (n = 6), and old saliva (n = 6) swabs. One-way ANOVA (F (2, 15) = 6.368, p = 0.0100) with Tukey's multiple comparisons test (Ctrl vs. Fresh, p = 0.0116; Ctrl vs Old, p = 0.0374; and Fresh vs Old, p = 0.8247).

(**J**) The number of cFos-IR positive neurons per 1000 cells in the dPAG in mice stimulated with control (n =5), fresh saliva (n = 6), and old saliva (n = 8) swabs. One-way ANOVA (F (2, 16) = 2.136, p = 0.1506).

In (**C-J**), the values are presented as means ± S.E.M., with individual dots representing individual mice. * and ** denote significance levels of $p < 0.05$ and 0.01, respectively.

Figure 5 Fresh and old cat saliva differentially activate VMH neurons to control defensive behavior

(**A-F**) Percentage of the total freezing episodes as a function of the number of cFos-positive posterior AOB mitral cells (**A**) and granule cells (**B**), BNST (**C**), MeApv (**D**), VMH (**E**), and dPAG (**F**) neurons. Green, and yellow symbols represent values from mice exposed to fresh saliva and old saliva, respectively. Sample size is same as seen in figure 4. Correlation was analyzed using Pearson correlation coefficient as data points in each group pass Shapiro-Wilk and Kolmogorov-Smirnov normality tests. Statistical results are reported in the table below each graph. * denotes statistical significance with $p < 0.05$.

Figure 6 A population of VMH neurons is more sensitive to fresh saliva than old saliva

(**A**) Schematic diagram illustrating the cFos-IR distribution analysis. Each VMH region in five consecutive sections (140 μm apart) was prepared along the dorsomedial (dm)-ventrolateral (vl) axis and divided into 10 bins that are vertical to the dm-vl axis. cFos-IR signals in each bin were counted.

(**B**) The number of cFos-IR-positive cells in each bin in the VMH hemisphere sections. Sample images are the same as seen in figure 4 (n = 6 – 7 animals). The values are presented as means + S.E.M., with individual dots representing values from individual VMH hemisphere sections. Statistical analysis was performed using Two-Way ANOVA with Tukey's multiple comparisons. Statistical results are reported in Supplemental Table 1. Black * and ** denote significance levels of $p < 0.05$ and 0.01, respectively, between control and fresh or old saliva conditions. Red * and ** denote significance levels of $p < 0.05$ and 0.01, respectively, between fresh and old saliva conditions.

Figure 7 A model of the VMH-to-VMH pathway processing imminence of predator threat

The circuitry is selectively activated based on the varying imminence of predator cues. Fresh saliva activates more V2R-A4-expressing VNO neurons (blue), which results in stronger activation of the mitral cells innervating V2R-A4 glomeruli. This eventually triggers freezing-correlated neural activity in the VMH. In contrast, old saliva activates fewer V2R-A4-expressing VNO neurons (blue), which results in weaker activation of the mitral cells innervating V2R-A4 glomeruli, leading to freezing-non-correlated activity in the VMH.

Supplemental figure 1

(**A**) Latency to the first freezing episode in mice exhibited freezing after the first contact with a swab. Two-tailed t test (t=0.7172, df=27, $p = 0.4794$).

(**B, C**) Raster plots displaying stretch sniffing (**B**) and direct investigation (**C**) episodes of individual mice exposed to a clean control swab (gray), fresh saliva (green), or old saliva (yellow). The introduction of a clean, fresh, or old saliva swab into a mouse's cage is denoted as time 0. Red lines in B indicate the first contact with a swab for each mouse.

Supplemental figure 2

Representative images of *V2R* in situ hybridization and pS6 immunohistochemistry for each V2R subfamily.

Supplemental figure 3

Representative images displaying the expression of cFos (red) in the BNST, MeApv, VMH, and dPAG of mice exposed to control, fresh, or old saliva swabs. Each neural substrate is delineated by a dashed white line, defined by counterstaining with DAPI. Scale bar: 200 μm.

Supplemental Table 1

Statistical summary of the two-way ANOVA with Tukey's multiple comparisons for Figure 6 (B).

Supplemental movie 1 Mouse behavior toward a control swab.

Supplemental movie 2 Mouse behavior toward a swab containing fresh saliva.

Supplemental movie 3 Mouse behavior toward a swab containing old saliva.

References

Apfelbach R, Blanchard CD, Blanchard RJ, Hayes RA, McGregor IS. 2005. The effects of predator odors in mammalian prey species: a review of field and laboratory studies. *Neurosci Biobehav Rev* **29**:1123–1144. doi:10.1016/j.neubiorev.2005.05.005

Belluscio L, Koentges G, Axel R, Dulac C. 1999. A map of pheromone receptor activation in the mammalian brain. *Cell* **97**:209–220. doi:10.1016/s0092-8674(00)80731-x

Ben-Shaul Y, Katz LC, Mooney R, Dulac C. 2010. In vivo vomeronasal stimulation reveals sensory encoding of conspecific and allospecific cues by the mouse accessory olfactory bulb. *Proc Natl Acad Sci U S A* **107**:5172–5177. doi:10.1073/pnas.0915147107

Beny-Shefer Y, Zilkha N, Lavi-Avnon Y, Bezalel N, Rogachev I, Brandis A, Dayan M, Kimchi T. 2017. Nucleus Accumbens Dopamine Signaling Regulates Sexual Preference for Females in Male Mice. *Cell Reports*. doi:10.1016/j.celrep.2017.11.062

Blanchard DC, Li CI, Hubbard D, Markham CM, Yang M, Takahashi LK, Blanchard RJ. 2003. Dorsal premammillary nucleus differentially modulates defensive behaviors induced by different threat stimuli in rats. *Neurosci Lett* **345**:145–148. doi:10.1016/s0304-3940(03)00415-4

Blanchard RJ, Blanchard DC. 1989. Attack and defense in rodents as ethoexperimental models for the study of emotion. *Prog Neuropsychopharmacol Biol Psychiatry* **13 Suppl**:S3-14. doi:10.1016/0278-5846(89)90105-x

Canteras NS. 2002. The medial hypothalamic defensive system: hodological organization and functional implications. *Pharmacol Biochem Behav* **71**:481–491. doi:10.1016/s0091-3057(01)00685-2

Canteras NS, Simerly RB, Swanson LW. 1994. Organization of projections from the ventromedial nucleus of the hypothalamus: a Phaseolus vulgaris-leucoagglutinin study in the rat. *J Comp Neurol* **348**:41–79. doi:10.1002/cne.903480103

Carvalho VM de A, Nakahara TS, Souza MA de A, Cardozo LM, Trintinalia GZ, Pissinato LG, Venancio JO, Stowers L, Papes F. 2020. Representation of Olfactory Information in Organized Active Neural Ensembles in the Hypothalamus. *Cell Rep* **32**:108061. doi:10.1016/j.celrep.2020.108061

Choi GB, Dong H-W, Murphy AJ, Valenzuela DM, Yancopoulos GD, Swanson LW, Anderson DJ. 2005. Lhx6 delineates a pathway mediating innate reproductive behaviors from the amygdala to the hypothalamus. *Neuron* **46**:647–660. doi:10.1016/j.neuron.2005.04.011

Davis BJ, Macrides F, Youngs WM, Schneider SP, Rosene DL. 1978. Efferents and centrifugal afferents of the main and accessory olfactory bulbs in the hamster. *Brain Res Bull* **3**:59–72. doi:10.1016/0361-9230(78)90062-x

Del Punta K, Puche A, Adams NC, Rodriguez I, Mombaerts P. 2002. A divergent pattern of sensory axonal projections is rendered convergent by second-order neurons in the accessory olfactory bulb. *Neuron* **35**:1057–1066. doi:10.1016/s0896-6273(02)00904-2

Dielenberg RA, McGregor IS. 2001. Defensive behavior in rats towards predatory odors: a review. *Neurosci Biobehav Rev* **25**:597–609. doi:10.1016/s0149-7634(01)00044-6

Engelke DS, Zhang XO, O'Malley JJ, Fernandez-Leon JA, Li S, Kirouac GJ, Beierlein M, Do-Monte FH. 2021. A hypothalamic-thalamostriatal circuit that controls approach-avoidance conflict in rats. *Nat Commun* **12**:2517. doi:10.1038/s41467-021-22730-y

Esteban Masferrer M, Silva BA, Nomoto K, Lima SQ, Gross CT. 2020. Differential Encoding of Predator Fear in the Ventromedial Hypothalamus and Periaqueductal Grey. *J Neurosci* **40**:9283–9292. doi:10.1523/JNEUROSCI.0761-18.2020

Ferrero DM, Moeller LM, Osakada T, Horio N, Li Q, Roy DS, Cichy A, Spehr M, Touhara K, Liberles SD. 2013. A juvenile mouse pheromone inhibits sexual behaviour through the vomeronasal system. *Nature* **502**:368–371. doi:10.1038/nature12579

Francia S, Silvotti L, Ghirardi F, Catzeflis F, Percudani R, Tirindelli R. 2014. Evolution of spatially coexpressed families of type-2 vomeronasal receptors in rodents. *Genome Biol Evol* **7**:272–285. doi:10.1093/gbe/evu283

Friard O, Gamba M. 2016. BORIS : a free, versatile open‑source event‑logging software for video/audio coding and live observations. *Methods Ecol Evol* **7**:1325‑1330. doi:10.1111/2041-210x.12584

Gross CT, Canteras NS. 2012. The many paths to fear. *Nat Rev Neurosci* **13**:651–658. doi:10.1038/nrn3301

Haga S, Hattori T, Sato T, Sato K, Matsuda S, Kobayakawa R, Sakano H, Yoshihara Y, Kikusui T, Touhara K. 2010. The male mouse pheromone ESP1 enhances female sexual receptive behaviour through a specific vomeronasal receptor. *Nature* **466**:118–122. doi:10.1038/nature09142

Haga-Yamanaka S, Ma L, He J, Qiu Q, Lavis LD, Looger LL, Yu CR. 2014. Integrated action of pheromone signals in promoting courtship behavior in male mice. *Elife* **3**:e03025. doi:10.7554/eLife.03025

Hua R, Yu S, Liu M, Li H. 2018. A PCR-Based Method for RNA Probes and Applications in Neuroscience. *Front Neurosci* **12**:266. doi:10.3389/fnins.2018.00266

Imamura F, Ito A, LaFever BJ. 2020. Subpopulations of Projection Neurons in the Olfactory Bulb. *Front Neural Circuits* **14**:561822. doi:10.3389/fncir.2020.561822

Ishii T, Omura M, Mombaerts P. 2004. Protocols for two- and three-color fluorescent RNA in situ hybridization of the main and accessory olfactory epithelia in mouse. *J Neurocytol* **33**:657–669. doi:10.1007/s11068-005-3334-y

Isogai Y, Si S, Pont-Lezica L, Tan T, Kapoor V, Murthy VN, Dulac C. 2011. Molecular organization of vomeronasal chemoreception. *Nature* **478**:241–245. doi:10.1038/nature10437

Isogai Y, Wu Z, Love MI, Ahn MH-Y, Bambah-Mukku D, Hua V, Farrell K, Dulac C. 2018. Multisensory Logic of Infant-Directed Aggression by Males. *Cell* **175**:1827-1841.e17. doi:10.1016/j.cell.2018.11.032

Itakura T, Murata K, Miyamichi K, Ishii KK, Yoshihara Y, Touhara K. 2022. A single vomeronasal receptor promotes intermale aggression through dedicated hypothalamic neurons. *Neuron* **110**:2455-2469.e8. doi:10.1016/j.neuron.2022.05.002

Jiang Y, Gong NN, Hu XS, Ni MJ, Pasi R, Matsunami H. 2015. Molecular profiling of activated olfactory neurons identifies odorant receptors for odors in vivo. *Nat Neurosci* **18**:1446–1454. doi:10.1038/nn.4104

Kats LB, Dill LM. 1998. The scent of death: Chemosensory assessment of predation risk by prey animals. *Écoscience* **5**:361–394.

Kennedy A, Kunwar PS, Li L-Y, Stagkourakis S, Wagenaar DA, Anderson DJ. 2020. Stimulus-specific hypothalamic encoding of a persistent defensive state. *Nature* **586**:730–734. doi:10.1038/s41586-020-2728-4

Kimchi T, Xu J, Dulac C. 2007. A functional circuit underlying male sexual behaviour in the female mouse brain. *Nature* **448**:1009–1014. doi:10.1038/nature06089

Kimoto H, Haga S, Sato K, Touhara K. 2005. Sex-specific peptides from exocrine glands stimulate mouse vomeronasal sensory neurons. *Nature* **437**:898–901. doi:10.1038/nature04033

Kunwar PS, Zelikowsky M, Remedios R, Cai H, Yilmaz M, Meister M, Anderson DJ. 2015. Ventromedial hypothalamic neurons control a defensive emotion state. *Elife* **4**. doi:10.7554/eLife.06633

Larriva-Sahd J. 2008. The accessory olfactory bulb in the adult rat: a cytological study of its cell types, neuropil, neuronal modules, and interactions with the main olfactory system. *J Comp Neurol* **510**:309–350. doi:10.1002/cne.21790

LeDoux J. 2012. Rethinking the emotional brain. *Neuron* **73**:653–676. doi:10.1016/j.neuron.2012.02.004

Leypold BG, Yu CR, Leinders-Zufall T, Kim MM, Zufall F, Axel R. 2002. Altered sexual and social behaviors in trp2 mutant mice. *Proc Natl Acad Sci U S A* **99**:6376–6381. doi:10.1073/pnas.082127599

Liman ER, Corey DP, Dulac C. 1999. TRP2: a candidate transduction channel for mammalian pheromone sensory signaling. *Proc Natl Acad Sci U S A* **96**:5791–5796. doi:10.1073/pnas.96.10.5791

Luo M, Fee MS, Katz LC. 2003. Encoding pheromonal signals in the accessory olfactory bulb of behaving mice. *Science* **299**:1196–1201. doi:10.1126/science.1082133

Markham CM, Blanchard DC, Canteras NS, Cuyno CD, Blanchard RJ. 2004. Modulation of predatory odor processing following lesions to the dorsal premammillary nucleus. *Neurosci Lett* **372**:22–26. doi:10.1016/j.neulet.2004.09.006

Mohrhardt J, Nagel M, Fleck D, Ben-Shaul Y, Spehr M. 2018. Signal Detection and Coding in the Accessory Olfactory System. *Chem Senses* **43**:667–695. doi:10.1093/chemse/bjy061

Osakada T, Abe T, Itakura T, Mori H, Ishii KK, Eguchi R, Murata K, Saito K, Haga-Yamanaka S, Kimoto H, Yoshihara Y, Miyamichi K, Touhara K. 2022. Hemoglobin in the blood acts as a chemosensory signal via the mouse vomeronasal system. *Nat Commun* **13**:556. doi:10.1038/s41467-022-28118-w

Osakada T, Ishii KK, Mori H, Eguchi R, Ferrero DM, Yoshihara Y, Liberles SD, Miyamichi K, Touhara K. 2018. Sexual rejection via a vomeronasal receptor-triggered limbic circuit. *Nat Commun* **9**:4463. doi:10.1038/s41467-018-07003-5

Papes F, Logan DW, Stowers L. 2010. The vomeronasal organ mediates interspecies defensive behaviors through detection of protein pheromone homologs. *Cell* **141**:692–703. doi:10.1016/j.cell.2010.03.037

Pérez-Gómez A, Bleymehl K, Stein B, Pyrski M, Birnbaumer L, Munger SD, Leinders-Zufall T, Zufall F, Chamero P. 2015. Innate Predator Odor Aversion Driven by Parallel Olfactory Subsystems that Converge in the Ventromedial Hypothalamus. *Curr Biol* **25**:1340–1346. doi:10.1016/j.cub.2015.03.026

Petrovich GD, Canteras NS, Swanson LW. 2001. Combinatorial amygdalar inputs to hippocampal domains and hypothalamic behavior systems. *Brain Res Brain Res Rev* **38**:247–289. doi:10.1016/s0165-0173(01)00080-7

Rodriguez I, Feinstein P, Mombaerts P. 1999. Variable patterns of axonal projections of sensory neurons in the mouse vomeronasal system. *Cell* **97**:199–208. doi:10.1016/s0092-8674(00)80730-8

Samuelsen CL, Meredith M. 2009. The vomeronasal organ is required for the male mouse medial amygdala response to chemical-communication signals, as assessed by immediate early gene expression. *Neuroscience* **164**:1468–1476. doi:10.1016/j.neuroscience.2009.09.030

Silva BA, Mattucci C, Krzywkowski P, Cuozzo R, Carbonari L, Gross CT. 2016. The ventromedial hypothalamus mediates predator fear memory. *Eur J Neurosci* **43**:1431–1439. doi:10.1111/ejn.13239

Silva BA, Mattucci C, Krzywkowski P, Murana E, Illarionova A, Grinevich V, Canteras NS, Ragozzino D, Gross CT. 2013. Independent hypothalamic circuits for social and predator fear. *Nat Neurosci* **16**:1731–1733. doi:10.1038/nn.3573

Silvotti L, Cavalca E, Gatti R, Percudani R, Tirindelli R. 2011. A recent class of chemosensory neurons developed in mouse and rat. *PLoS One* **6**:e24462. doi:10.1371/journal.pone.0024462

Silvotti L, Moiani A, Gatti R, Tirindelli R. 2007. Combinatorial co-expression of pheromone receptors, V2Rs. *J Neurochem* **103**:1753–1763. doi:10.1111/j.1471-4159.2007.04877.x

Stowers L, Holy TE, Meister M, Dulac C, Koentges G. 2002. Loss of sex discrimination and male-male aggression in mice deficient for TRP2. *Science* **295**:1493–1500. doi:10.1126/science.1069259

Takahashi LK. 2014. Olfactory systems and neural circuits that modulate predator odor fear. *Front Behav Neurosci* **8**:72. doi:10.3389/fnbeh.2014.00072

Takami S, Graziadei PP. 1991. Light microscopic Golgi study of mitral/tufted cells in the accessory olfactory bulb of the adult rat. *J Comp Neurol* **311**:65–83. doi:10.1002/cne.903110106

Tobias BC, Schuette PJ, Maesta-Pereira S, Torossian A, Wang W, Sethi E, Adhikari A. 2023. Characterization of ventromedial hypothalamus activity during exposure to innate and conditioned threats. *Eur J Neurosci* **57**:1053–1067. doi:10.1111/ejn.15937

Tsunoda M, Miyamichi K, Eguchi R, Sakuma Y, Yoshihara Y, Kikusui T, Kuwahara M, Touhara K. 2018. Identification of an Intra- and Inter-specific Tear Protein Signal in Rodents. *Curr Biol* **28**:1213-1223.e6. doi:10.1016/j.cub.2018.02.060

von Campenhausen H, Mori K. 2000. Convergence of segregated pheromonal pathways from the accessory olfactory bulb to the cortex in the mouse. *Eur J Neurosci* **12**:33–46. doi:10.1046/j.1460-9568.2000.00879.x

Wagner S, Gresser AL, Torello AT, Dulac C. 2006. A multireceptor genetic approach uncovers an ordered integration of VNO sensory inputs in the accessory olfactory bulb. *Neuron* **50**:697–709. doi:10.1016/j.neuron.2006.04.033

Wang L, Chen IZ, Lin D. 2015. Collateral pathways from the ventromedial hypothalamus mediate defensive behaviors. *Neuron* **85**:1344–1358. doi:10.1016/j.neuron.2014.12.025

Wong WM, Cao J, Zhang X, Doyle WI, Mercado LL, Gautron L, Meeks JP. 2020. Physiology-forward identification of bile acid-sensitive vomeronasal receptors. *Sci Adv* **6**:eaaz6868. doi:10.1126/sciadv.aaz6868

Yonekura J, Yokoi M. 2008. Conditional genetic labeling of mitral cells of the mouse accessory olfactory bulb to visualize the organization of their apical dendritic tufts. *Mol Cell Neurosci* **37**:708–718. doi:10.1016/j.mcn.2007.12.016

Figure 1 Predator cues in cat saliva induce context-dependent defensive behavior in mice

(A) A schematic diagram illustrating the elicitation of "freeze" or "flight" defensive behaviors in relation to distance. (Adapted with permission from (Blanchard and Blanchard, 1989))

(B) A schematic illustration depicting a test chamber for evaluating defensive behaviors.

(C) Raster plots displaying freezing episodes of individual mice exposed to a clean control swab (gray, n = 18), fresh saliva (green, n = 18), or old saliva (yellow, n = 15). The introduction of a clean, fresh, or old saliva swab into a mouse's cage is denoted as time 0. Red lines indicate the first contact with a swab for each mouse.

(D) Latency to the first episode of contact with a swab. One-way ANOVA (F (2,47) = 5.409, p = 0.0077) with Tukey's multiple comparisons test (Ctrl vs. Fresh, p = 0.0640; Ctrl vs Old, p = 0.5932; and Fresh vs Old, p = 0.0074).

(E) The percentage of total freezing episodes after the first contact. One-way ANOVA (F (2,47) = 37.38, p < 0.0001) with Tukey's multiple comparisons test (Ctrl vs. Fresh, p < 0.0001; Ctrl vs Old, p = 0.0411; and Fresh vs Old, p < 0.0001).

(F) The percentage of total stretch sniff episodes. One-way ANOVA (F(2,49) = 0.007016, p = 0.9930).

(G) The percentage of total direct investigation episodes. One-way ANOVA (F (2,48) = 25.17, p < 0.0001) with Tukey's multiple comparisons test (Ctrl vs. Fresh, p < 0.0001; Ctrl vs Old, p < 0.0001; and Fresh vs Old, p = 0.5762).

(H) Plasma ACTH concentrations following exposure to clean (n = 10), fresh (n = 8), or old saliva (n = 10) swabs. One-way ANOVA (F (2,25) = 3.735, p = 0.0381) with Tukey's multiple comparisons test (Ctrl vs. Fresh, p = 0.0370; Ctrl vs Old, p = 0.1609; and Fresh vs Old, p = 0.6804).

In (D) - (H), the values are presented as means ± S.E.M., with individual dots representing individual mice. *, **, and **** denote significance levels of p < 0.05, 0.01, and 0.0001, respectively.

Figure 2 Cat saliva activates VNO neurons and modulates freezing behavior

(A) The percentage of total freezing episodes directed towards control (n = 12) or fresh saliva (n = 11) swabs in cassettes that prevent direct contact with the swabs. Two-tailed t test (t = 0.08745, df = 21, p = 0.9311).

(B) The percentage of total investigation episodes toward cassettes containing control or fresh saliva swabs. Two-tailed t test (t=1.662, df=21, p = 0.1114).

(C) The percentage of total freezing episodes directed toward control or fresh saliva swabs in Trpc2-WT (n = 10) or -KO (n = 9) mice. Two-way ANOVA ([genotype] F (1, 34) = 3.662, p = 0.0641, [swab type] F (1, 34) = 8.053, p = 0.0076, [genotype] and [swab type] F (1, 34) = 5.963, p = 0.0200) with Tukey's multiple comparisons test (WT-Ctrl vs KO-Ctrl, p = 0.9922; WT-Ctrl vs WT-Saliva, p = 0.0164; WT-Ctrl vs KO-Saliva, p = 0.9136; KO-Ctrl vs WT-Saliva, p = 0.0100; KO-Ctrl vs KO-Saliva, p = 0.9832; WT-Saliva vs KO-Saliva, p = 0.0037).

(D) The percentage of total direct investigation episodes toward control or fresh saliva swabs in Trpc2-WT or -KO mice. Two-way ANOVA ([genotype] F (1, 34) = 11.46, p = 0.0018, [swab type] F (1, 34) = 7.725, p = 0.0088, [genotype] and [swab type] F (1, 34) = 13.96, p = 0.0007) with Tukey's multiple comparisons test (WT-Ctrl vs KO-Ctrl, p = 0.9051; WT-Ctrl vs WT-Saliva, p < 0.0001; WT-Ctrl vs KO-Saliva, p = 0.9732; KO-Ctrl vs WT-Saliva, p = 0.0006; KO-Ctrl vs KO-Saliva, p = 0.9949; WT-Saliva vs KO-Saliva, p = 0.0003).

(E) Representative images illustrating the expression of pS6 (green) and Gαo (magenta) in the VNO of mice exposed to control, fresh, or old saliva swabs. DAPI was used as a nuclear counterstain. Scale bar: 50 μm.

(F, G) The number of pS6-IR positive cells in 1000 VNO cells in the apical (F) and basal (G) regions of the VNO neuroepithelium in mice stimulated with control (n = 8), fresh saliva (n = 8), and old saliva (n = 8) swabs. (F) One-way ANOVA (F (2, 21) = 2.562, p = 0.1010). (G) One-way ANOVA (F (2, 21) = 17.90, p < 0.0001) with Tukey's multiple comparisons test (Ctrl vs. Fresh, p < 0.0001; Ctrl vs Old, p = 0.2262; and Fresh vs Old, p = 0.0014).

(H) The percentage of total freezing episodes in relation to the number of pS6-positive cells in the VNO of individual mice (n = 8 per swab type). Correlation analysis was conducted using Spearman's rank correlation coefficient as total data points did not pass Shapiro-Wilk and Kolmogor-ov-Smirnov normality tests. Statistical results are reported in the graph.

(A - D, F, G) the values are presented as means ± S.E.M., with individual dots representing individual mice. *, **, ***, and **** denote significance levels of p < 0.05, 0.01, 0.001, and 0.0001, respectively.

Figure 3 Cat saliva activates the V2R-A4 subfamily of the vomeronasal receptors

(A) Phylogenetic tree of the V2R family and its subfamilies used for V2R cRNA probe design.

(B) Representative images displaying the expression of V2RA4 (magenta) and pS6 (green) in the VNO of a mouse exposed to a fresh saliva swab. DAPI was utilized as a nuclear counterstain. Scale bar: 50 μm.

(C) The percentage of cells co-labeled with V2R probes and anti-pS6 antibody among the total pS6-positive cells in the VNO of mice exposed to fresh saliva (top, from 3 – 6 animals) or old saliva (bottom, from 3 – 6 animals).

Figure 4 Cat saliva activates neurons in the AOB and defensive behavioral circuit

(A) Schematic diagrams illustrating the VNO sensory pathway that controls defensive behaviors.

(B) Representative images displaying the expression of cFos (magenta) in the mitral cell (MCL) and granule cell (GCL) layers of the AOB of mice exposed to control, fresh, or old saliva swabs. Gαi2 (green) visualizes the anterior region of the glomerular layer (GL). DAPI was utilized as a nuclear counterstain. Scale bar: 200 μm.

(C, D) The number of cFos-IR positive mitral cells per 1000 mitral cells in the anterior (C) and posterior (D) regions of the AOB (n = 10 per swab type). (C) One-way ANOVA (F (2, 27) = 0.6106, p = 0.5504). (D) One-way ANOVA (F (2, 27) = 6.665, p = 0.0044) with Tukey's multiple comparisons test (Ctrl vs. Fresh, p = 0.0210; Ctrl vs Old, p = 0.0059; and Fresh vs Old, p = 0.8593).

(E, F) The number of cFos-IR positive granule cells per 1000 granule cells in the anterior (E) and posterior (F) regions of the AOB (n = 10 per swab type). (E) One-way ANOVA (F (2, 27) = 0.5831, p = 0.5651). (F) One-way ANOVA F (2, 27) = 5.996, p = 0.0070) with Tukey's multiple comparisons test (Ctrl vs. Fresh, p = 0.0262; Ctrl vs Old, p = 0.0099; and Fresh vs Old, p = 0.9105).

(G) The number of cFos-IR positive neurons per 1000 cells in the BNST in mice stimulated with control (n = 10), fresh saliva (n = 5), and old saliva (n = 6) swabs. One-way ANOVA (F (2, 18) = 0.4375, p = 0.6524).

(H) The number of cFos-IR positive neurons per 1000 cells in the MeApv in mice stimulated with control (n = 9), fresh saliva (n = 6), and old saliva (n = 7) swabs. One-way ANOVA (F (2, 19) = 2.454, p = 0.1128).

(I) The number of cFos-IR positive neurons per 1000 cells in the VMH in mice stimulated with control (n = 6), fresh saliva (n = 6), and old saliva (n = 6) swabs. One-way ANOVA (F (2, 15) = 6.368, p = 0.0100) with Tukey's multiple comparisons test (Ctrl vs. Fresh, p = 0.0116; Ctrl vs Old, p = 0.0374; and Fresh vs Old, p = 0.8247).

(J) The number of cFos-IR positive neurons per 1000 cells in the dPAG in mice stimulated with control (n =5), fresh saliva (n = 6), and old saliva (n = 8) swabs. One-way ANOVA (F (2, 16) = 2.136, p = 0.1506).

In (C-J), the values are presented as means ± S.E.M., with individual dots representing individual mice. * and ** denote significance levels of p < 0.05 and 0.01, respectively.

Figure 5 Fresh and old cat saliva differentially activate VMH neurons to control defensive behavior

(A-C) Percentage of the total freezing episodes as a function of the number of cFos-positive posterior AOB mitral cells (A) and granule cells (B), as well as VMH neurons (C). Green, and yellow symbols represent values from mice exposed to fresh saliva and old saliva, respectively. Sample size is same as seen in figure 4. Correlation was analyzed using Pearson correlation coefficient as data points in each group pass Shapiro-Wilk and Kolmogorov-Smirnov normality tests. Statistical results are reported in the table below each graph. * denotes statistical significance with p < 0.05.

Figure 6 A population of VMH neurons is more sensitive to fresh saliva than old saliva

(A) Schematic diagram illustrating the cFos-IR distribution analysis. Each VMH region in five consecutive sections (140 μm apart) was prepared along the dorsomedial (dm)-ventrolateral (vl) axis and divided into 10 bins that are vertical to the dm-vl axis. cFos-IR signals in each bin were counted.

(B) The number of cFos-IR-positive cells in each bin in the VMH hemisphere sections. Sample images are the same as seen in figure 4 (n = 6 – 7 animals). The values are presented as means + S.E.M., with individual dots representing values from individual VMH hemisphere sections. Statistical analysis was performed using Two-Way ANOVA with Tukey's multiple comparisons. Statistical results are reported in Supplemental Table 1. Black * and ** denote significance levels of $p < 0.05$ and 0.01, respectively, between control and fresh or old saliva conditions. Red * and ** denote significance levels of $p < 0.05$ and 0.01, respectively, between fresh and old saliva conditions.

Figure 7 A model of the VMH-to-VMH pathway processing imminence of predator threat

The circuitry is selectively activated based on the varying imminence of predator cues. Fresh saliva activates more V2R-A4-expressing VNO neurons (blue), which results in stronger activation of the mitral cells innervating V2R-A4 glomeruli. This eventually triggers freezing-correlated neural activity in the VMH. In contrast, old saliva activates fewer V2R-A4-expressing VNO neurons (blue), which results in weaker activation of the mitral cells innervating V2R-A4 glomeruli, leading to freezing-non-correlated activity in the VMH.