Kinetochore and ionomic adaptation to whole genome duplication

Sian M. Bray[1], Tuomas Hämälä[1], Min Zhou[2], Silvia Busoms[3,4], Sina Fischer[1], Stuart D. Desjardins[5], Terezie Mandáková[6], Chris Moore[1], Thomas C. Mathers[7,8], Laura Cowan[1,9], Patrick Monnahan[3], Jordan Koch[3], Eva M. Wolf[10], Martin A. Lysak[6], Filip Kolar[11,12], James D. Higgins[5], Marcus A. Koch[10], and Levi Yant*[,1,11]

1. University of Nottingham, Nottingham NG7 2RD, United Kingdom
2. School of Life Sciences, Chongqing University, Chongqing 401331, China
3. Department of Cell and Developmental Biology, John Innes Centre, Norwich Research Park NR4 7UH, Norwich, UK
4. Department of Plant Physiology, Universitat Autònoma de Barcelona 08193 Barcelona, Spain
5. Department of Genetics and Genome Biology, University of Leicester, Leicester LE1 7RH, UK
6. CEITEC–Central European Institute of Technology, Masaryk University, Brno 625 00, Czech Republic
7. Department of Crop Genetics, John Innes Centre, Norwich Research Park NR4 7UH, Norwich, UK
8. Current address: Tree of Life, Welcome Sanger Institute, Hinxton, Cambridge, CB10 1SA, UK
9. Current address: Department of Plant Sciences, Cambridge University, Cambridge CB2 3EA, UK
10. Department of Biodiversity and Plant Systematics, Centre for Organismal Studies, Heidelberg University, 69120 Heidelberg, Germany
11. Department of Botany, Charles University, Benátská 2, CZ-12801 Prague, Czech Republic
12. Institute of Botany, The Czech Academy of Sciences, Zámek 1, CZ-252 43 Průhonice, Czech Republic

*Author for correspondence: levi.yant@nottingham.ac.uk

Keywords: polyploidy; evolution; kinetochore; meiosis; ion homeostasis; population genomics

Abstract

Whole genome duplication (WGD) brings challenges to key processes like meiosis, but
nevertheless is associated with diversification in all kingdoms. How is WGD tolerated, and what
processes commonly evolve to stabilize the new polyploid lineage? Here we study this in
Cochlearia spp., which have experienced multiple rounds of WGD in the last 300,000 years. We
first generate a chromosome-scale genome and sequence 113 individuals from 33 diploid,
tetraploid, hexaploid, and outgroup populations. We detect the clearest post-WGD selection
signatures in functionally interacting kinetochore components and ion transporters. We
structurally model these derived selected alleles, associating them with known WGD-relevant
functional variation and compare these results to independent recent post-WGD selection in
Arabidopsis arenosa and *Cardamine amara*. Some of the same biological processes evolve in all
three WGDs, but specific genes recruited are flexible. This points to a polygenic basis for
modifying systems that control the kinetochore, meiotic crossover number, DNA repair, ion
homeostasis, and cell cycle. Given that DNA management (especially repair) is the most salient
category with the strongest selection signal, we speculate that the generation rate of structural
genomic variants may be altered by WGD in young polyploids, contributing to their occasionally
spectacular adaptability observed across kingdoms.

Whole genome duplication (WGD, leading to polyploidy) is a dramatic mutation that disrupts

fundamental cellular processes. Yet, for those that can adapt to a transformed polyploid state,

WGD holds great promise[1-3]. Despite its importance to evolution, agriculture, and human health,

we do not yet know why some polyploids thrive, while others do not[1,4].

Immediately following WGD in autopolyploids (within-species WGD, not hybrid

allopolyploids), novel challenges arise. The most obvious concerns meiosis: the instant doubling

of homologs complicates chromosome pairing[5]. If a chromosome engages in crossing over with

more than one homolog, entanglements and breakage ensue at anaphase. WGD also disrupts

cellular equilibria, including ion homeostasis and cell size[6]. These challenges are insurmountable

for many nascent polyploids, although established autopolyploids can persist, indicating that they

can be overcome.

To date, work in two diploid-autotetraploid model systems has explicitly sought a basis of

adaptation to WGD in recent (< 300,000 year-old) autopolyploids. In *Arabidopsis arenosa,* a

handful of physically and functionally interacting meiosis proteins undergo adaptive evolution less

than 200,000 years post-WGD[7,8]. Derived alleles of these genes decrease chromosome

crossover rates, stabilizing meiosis[9-12]. Next, a pool-seq-based scan in *Cardamine*, a genus 17

million years diverged from *Arabidopsis,* showed only very modest convergence with *A. arenosa*

on the level of functional pathways for processes under selection shortly post-WGD[13]. Meiosis

showed little convergence and the signal of evolution of the meiosis genes obviously controlling

crossovers seen in *A. arenosa* was absent. These works gave insight but had important

limitations: there was very low sample number in *A. arenosa* (24 individuals[7])*,* and in *C. amara,*

pool-seq of only four populations using a highly fragmented reference[13]. More significantly, as

noted in the *C. amara* study, widespread vegetative reproduction in *C. amara* offers some escape

from selection for meiotic stability post-WGD, consistent with the results[13]. Thus, minimal

convergence between these systems leaves unresolved what salient processes stabilize recent

polyploids. This is important because the genomic changes that occur post-WGD may also help

explain why some polyploids are so successful and most are not.

Here we address this in a novel system that overcomes these limitations in a more

distantly related, independent, and successful set of WGD events within a single species flock (at

least 5 WGDs in *Cochlearia* the last 300,000 years)[14]. The *Cochlearia* species complex exhibits

diploid, autotetraploid, allohexaploid, octoploid and heptaploid cytotypes (Fig.1A)[14-17], with the

widespread autotetraploid cytotype[14], similar in age to *A. arenosa*[18,19]. *Cochlearia* is found across

Europe, from Spain to the Arctic, in a wide range of habitats including freshwater springs, coastal

cliffs, sand dunes, salt marshes, metal contaminated sites and roadside grit (Fig. 1A)[14,20-30]. A

broad habitat differentiation is evident by ploidy, with diploids typically found in upland freshwater
springs, autotetraploids overwhelmingly on coasts, often directly adjacent to seawater or
continuously submerged, and hexaploids in extreme salt marsh conditions. In fact, the hexaploid
Cochlearia danica is one of the most rapidly spreading invasive species in Europe, invading salted
roadways since the 1970's[23,31].

Here we first assess *Cochlearia* demography by individually resequencing 113 plants from
33 diploid, autotetraploid, hexaploid, and outgroup populations from across its ploidy-variable
range. We then focus our analysis on closely related diploids and autotetraploids in the UK and
scan for selective sweeps post-WGD. We dissect functional targets of adaptive evolution post-
WGD using protein modelling and identification of orthologous derived sites from functional
studies. Our results show convergence at the process level in very recent WGD adaptation events
in these three genera separated by greater spans of ~40 million years. This indicates that similar
processes adapt in response to WGD, but that specific genes recruited are far less constrained.
Surprisingly, also we also find strong signal of post-WGD evolution in several kinetochore
components, pointing to a novel mechanism of adaptation to polyploid meiosis and mitosis.

Results and Discussion

Chromosome-level assembly. To serve as a reference for our demographic analysis and
selection scans, we built a chromosome-level genome assembly of one diploid *Cochlearia*
excelsa individual (Styria, Gurktaler Alpen, Austria). We chose *C. excelsa* because it is an early
diverging diploid (2n = 12) and conveniently also a rare primarily selfing *Cochlearia* species. This
resulted in a highly contiguous primary assembly (contig N50=15 Mb; Fig.1B), generated from
Oxford Nanopore PromethION data (read N50=27 Kb). The primary assembly was performed
with Flye 2.9[32] and NECAT[33] with one round of polishing in Medaka[34] and Pilon[35] The assembled
contigs were then scaffolded to chromosome scale using Hi-C (Fig.1B) and a final cleanup
performed with Blobtools[36] (Fig. S1). Hi-C-guided chromosome arm orientations were confirmed
with concordant FISH and in silico mapping of telomeric and centromeric repeats (Fig.1B; Fig.
S2). This assembly consists of six primary scaffolds corresponding to the six *C. excelsa*
chromosomes with a scaffold N50 of 37 Mb and an overall genome size of 310 Mb (Fig.1B, C),
matching our estimated haploid genome size of 302 Mb (Fig. S3). Gene space representation
was very good, with 96.4% complete Brassicales BUSCOs[37] found in the assembly Fig.1C). We
performed an annotation incorporating RNA-seq data from the reference line (flower bud, leaf,
stem, and silique), protein homology information, and *ab initio* modelling with BRAKER2[38]. This
yielded 54,424 gene models across the six chromosomes.

113
Figure 1 | Ploidy variation, genome assembly, sampling, and genetic structure. **(A)** Three *Cochlearia*
species in this study (top: *C. pyrenaica* [2n=2x=12], middle: *C. officinalis* [2n=4x=24], bottom: *C. danica*
[2n=6x=42]. Scale bars, 10 μm; **(B)** Hi-C contact map and cytology used to orient chromosome arms.
Chromosomes are bounded in blue and centromeres indicated in green. Cytology shows DAPI-stained
chromosomes with heterochromatic pericentromeres (top) and 102 bp satellite (pink) and Arabidopsis-type
telomeric satellite (green) probes hybridizing to all (peri)centromeres and telomeres, respectively (bottom).
Scale bar, 10 μm; **(C)** Chromosome-scale assembly of the diploid *C. excelsa* genome; **(D)** The 30
Cochlearia populations included in the short read sequencing (locations in Dataset S2); **(E)** Nei's genetic
distances between the populations (hexaploids excluded) visualized in SplitsTree[39]; **(F)** fastSTRUCTURE[40]
analysis (*k*=3, min alleles=8) of all *Cochlearia* individuals in the study, with regions and ploidies indicated.
Blue=diploids; orange=tetraploids; green=hexaploids.

Geographic distribution, ploidy variation, cohort construction and population sequencing.
To determine optimal population contrasts for WGD-specific signatures of selection, we sampled populations across the reported range of the *Cochlearia* species throughout Europe[14,17,22-25,27,28,30] and conducted flow cytometry- and cytologically-based surveys of genome size and ploidy variation (Fig.1A; Dataset S1 Ploidy Survey; Fig. S4). Measurements were normalised against the diploid population with the most stable individual within-population genome size estimates, WOL (Dataset S1). We focus our demographic analysis on the three most abundant ploidies (Fig.1A): diploids = *Cochlearia pyrenaica* (inland UK and mainland Europe)*,* autotetraploids = *Cochlearia officinalis* (coastal UK and Norway)*,* hexaploids = *Cochlearia danica* (inland UK). Based on these ploidy surveys we chose 113 individuals from 33 populations for sequencing by Illumina PE (average per-individual depth = 17x; minimum = 4x; Fig.1D; Dataset S2 Sample Metrics). The final dataset consisted of 18,307,309 SNPs, on average one variant every 17 bp (quality and depth filtered; Methods).

Genetic structure. To assess structure in our dataset, we first performed fastSTRUCTURE[40] (Fig. 1F) on our 109 *Cochlearia* individuals, excluding outgroup sister genus Ionopsidium (23,733 LD-pruned, biallelic 4-fold-degenerate SNPs; max 20% missing data; min minor allele frequency = 0.02). *K*=3 maximized marginal likelihood and grouped samples by ploidy. Focusing on diploids vs tetraploids, PCA confirmed that ploidy dominates over geography (PC1 [ploidy] = 27% of variance explained; PC2 [UK vs Norwegian tetraploids] = 11% of variance explained; Fig. S5). This is reflected also in clean geographical groupings by SplitsTree[39] analyses, which visualize simple genetic distances (Fig. 1E). This is consistent with a rapid inter- or peri-glacial radiation and postglacial migration such as found in other *Cochlearia* species[14].

Meiotic and ion homeostasis-related phenotypic shifts upon WGD. In a young autopolyploid, initial maintenance of diploid-like crossover frequencies can lead to chromosome entanglements and breakage at meiosis. We therefore confirmed the establishment of meiotic stability of these autotetraploids (Fig. 2). Similar to *A. arenosa* autotetraploids[7], we found a significant per-chromosome reduction in class I mature crossovers, evidenced by HEI10 foci (p <0.00001; Mann-Whitney). This translated to a degree of meiotic stability in the *Cochlearia* tetraploids, although we observed considerably greater variation in meiotic stability in autotetraploids, which varied dramatically both within and between families for multivalent production, as seen in *A. arenosa*. This high variability suggests segregating variation for factors promoting stability (Fig. 2C, D).

Figure 2 | Reduction in crossover number, improved drought response, and high inter-lineage phenotypic variability upon WGD. (A and B) Quantification of mature meiotic crossover events by HEI10 staining (green) on pachytene chromosomes stained by ZIP1 (red), showing a significant per-chromosome downregulation of crossovers in *Cochlearia* autotetraploids (p <0.00001; Mann-Whitney); **(C)** Metaphase I chromosome spreads and **(D)** quantification of meiotic stability, showing decreased, but highly variable stability in tetraploids relative to diploids. Increased stomatal conductance **(E)** and photosynthetic **(F)** rates under dehydration stress conditions in tetraploids. Plots show median and variation of drought stressed plants in comparison to well-watered plants. (Sign differences between diploids and autotetraploids as seen in one-way ANOVA with posthoc Tukey; Table S4) Blue=diploids; orange=tetraploids. Scale bars, 10 μm.

Cochlearia in the UK exhibits a broadly disjunct geographic distribution by ploidy, with

diploids (n=6) deeply inland and autotetraploids inhabiting coastal regions of the highest salinity,

including full seawater submergence (Fig.1A; Fig. S4). A direct mechanistic link between WGD

and salinity tolerance was established in *Arabidopsis thaliana*, where first-generation

neoautopolyploids (otherwise isogenic with diploid siblings) show elevated salinity tolerance and

intracellular potassium[41]. We therefore tested for ploidy-related differences in salinity tolerance

and dehydration stress tolerance in wild *Cochlearia*. Interestingly, in terms of overall plant survival,

we found extreme salinity tolerance in all ecotypes tested, with even diploids tolerating up to

600mM NaCl (salinity level of seawater), along with all higher ploidies (Table S1). Tetraploids

showed signal of increased drought tolerance, with both elevated stomatal conductance and net

photosynthetic rates under drought, relative to diploids (Fig. 2E and F; Tables S2-S4). This

suggests an adaptive benefit specific to higher ploidies in response to drought. A benefit under

salinity stress may be tempered by preadaptation to a stringent ionic challenge even in the diploids

of this species flock, consistent with their halophyte and cold-loving nature[42,43].

Selective sweeps associated with WGD. To identify candidate genes and processes mediating

adaptation to WGD, we next focused on the 18 geographically proximal populations from the UK

(44 autotetraploid individuals and 29 diploid individuals) with good sequencing coverage.

Concentrating the selection scan on UK diploids and autotetraploids minimizes the effect of

genetic structure that would be introduced if we were to use the mainland European samples as

well. To guide our selection scan window size choice, we calculated pairwise linkage decay, which

was rapid in both diploids and tetraploids, with near complete lack of genotypic correlations within

2 kb to very low background levels (Fig. 3A). We thus calculated in 1kb windows a battery of

differentiation metrics (Dxy[44], Rho[45], Hudson's F_{ST}[46], Nei's F_{ST}[47], Weir-Cochran's F_{ST}[48], and

groupwise allele frequency difference [AFD]) genome-wide (minimum = 15; mean = 101 SNPs

per window). After filtering, these scans overlapped 40,245 of 54,424 total predicted genes, or

74% of gene coding loci with sufficient coverage for assessment.

To empirically determine which metric most reliably identified genomic regions that exhibit

localized peaks in AFD indicative of specific sweeps upon WGD, we performed an inspection of

all AFD plots in all outlier tails (see Methods). From this, we identified Hudson's F_{ST} as most

reliably identifying regions of localized AFD peaks (and not e.g., low diversity[44]). This formulation

of F_{ST} brings the added benefit of robustness for unequal population sizes and presence of rare

variants[49] and direct comparisons to the *C. amara* study, which used this same metric[13]. We

therefore extracted windows in the top 1% of this distribution as candidate outliers, consisting of

1,823 1kb windows, overlapping 753 gene-coding loci for which we could obtain functional

descriptions primarily from orthology (or, lacking this, close homology) to *Arabidopsis thaliana*

(Dataset S3 Selective Sweep Candidates). We focus on the most extreme 25 of these (to the right

of the dashed line in Fig 3B; Table 1), which we confirm exhibit elevated Dxy values (outlier F_{ST}

peak Dxy=0.17; mean outside peak Dxy=0.06; Mann Whitney U test: $P < 2 \times 10^{-16}$; Fig. 3C).

Figure 3 | Rapid linkage decay and empirical outlier analysis. **(A)** Immediate decay of genotypic
correlations (r^2) observed in both diploid (blue) and tetraploid (orange) *Cochlearia*; **(B)** Distribution of
genome wide F_{ST} values for 182,327 1kb windows. The dashed red line gives the extreme stringency F_{ST}
cutoff of the top 25 genome-wide outlier genes; the solid red line gives the 1% cutoff; windows inside the
CENP-E gene coding region are highlighted in orange; **(C)** Dxy values are significantly elevated inside F_{ST}
peaks (Mann Whitney U test: $P < 2 \times 10^{-16}$); **(D)** *CENP-E*, the #1 genome-wide F_{ST} outlier, also exhibits
greatly excess differentiation for its level of diversity in the tetraploids, a classical signal of selective sweep.
215

Complementing this approach, we also focused our 1% outlier list on gene coding regions

with a fineMAV approach[50]. Using Grantham scores to estimate functional impact of each non-

synonymous amino acid change encoded by a given SNP, this approach scales the severity of

predicted amino acid change by the AFD between groups. Of the 107,055 non-synonymous-

encoding SNPs assigned a MAV score, the top 1% outliers from the empirical distribution were

intersected with our F_{ST} outliers, yielding a protein-evolution-oriented list of 159 gene coding loci,

harboring 290 MAV SNPs (bold in Dataset S3 Selective Sweep Candidates; 1% F_{ST} outliers with

10 or more 1% extreme MAV outliers are given in Table 1). By these approaches, we could

resolve clear gene-specific peaks of F_{ST} (Fig. 4A) and candidate selective sweep alleles in our

top 25 genome-wide outliers (Fig. 4 B-G).

F_{ST} rank	*Cochlearia* ID	*A.thaliana* ID	Name	Description
1*	g7445	AT3G10180	CENP-E	**CENTROMERE PROTEIN E.** Kinetochore protein that moves mono-oriented chromosomes to the spindle equator[51]. Cooperates with chromokinesins and dynein to mediate chromosome congression. Activity is regulated by post-translational modifications, protein interactions and autoinhibition. Tetraploid cancers are far more susceptible to CENP-E inhibitors than diploids[52-54].
2	g31016	AT1G06670	NIH	NUCLEAR DEIH-BOX HELICASE. Binds DNA without clear specificity[55].
3	g7446	AT3G10070	TAF12	TATA-ASSOCIATED II 58. Controls stress responsive root growth[56]
4	g40185	AT2G26690	NPF6.2	NITRATE TRANSPORTER 6.2. Mediates drought stress response[57]
5	g25338	AT4G00060	MEE44	MATERNAL EFFECT EMBRO ARREST 44: Part of the RNA TRAMP complex
6	g49945	AT1G15660	CENP-C	**CENTROMERE PROTEIN C**. Kinetochore protein that is critical for centromere identity in both mitosis and meiosis. Loss of CENP-C results in aneuploidy and cell death[60]. Necessary in mitotic cells for kinetochore assembly centromere establishment. Mutants also fail to retain centromeric SC in late pachytene[58,59].
7	g25334	AT1G05940	CAT9	CATIONIC AMINO ACID TRANSPORTER 9
8	g25335	AT4G00026	SD3	**SEGREGATION DISTORTION 3**. Mutants have lower ploidy[60].
9	g6996	AT3G16730	CAP-H2 / HEB2	**CONDENSIN-II COMPLEX SUBUNIT H2**. Functions in DSB repair and arranges interphase chromatin[61,62]. Plays a role in alleviating DNA damage and genome integrity in *A.thaliana*[62]. Condensin II-depleted human cells have a defect in homologous recombination-mediated repair[63].
10	g24649	AT5G40595		Unknown protein
11	g24648	AT4G02000		Ta11-like non-LTR retrotransposon
12	g33330	AT1G54310		S-adenosyl-L-methionine-dependent methyltransferases superfamily protein
13	g40302	AT4G10310	HKT1	**HIGH-AFFINITY K+ TRANSPORTER 1**. Sodium transporter. Mediates salinity tolerance in wild *A. thaliana* populations[64]. Under selection post-WGD in *A. arenosa*[7].
14	g33311	AT5G61390	NEN2	NAC45/86-DEPENDENT EXONUCLEASE-DOMAIN PROTEIN 2
15	g10739	AT2G01980	SOS1	**SALT OVERLY SENSITIVE 1.** Plasma membrane-localized Na+/H+ antiporter that extrudes Na+ from cells. Is essential for plant salt and stress tolerance.
16	g46402	AT1G77990	SULTR2	SULPHATE TRANSPORTER 2;2. A low-affinity sulfate transporter
17	g4631	AT5G19270		Reverse transcriptase-like protein
18	g4632	AT2G07200		Cysteine proteinases superfamily protein
19	g25121	AT5G35970		P-loop containing nucleoside triphosphate hydrolases superfamily protein
20	g45503	AT5G08620	STRS2	STRESS RESPONSE SUPPRESSOR 2. A DEA(D/H)-box helicase involved in drought, salt and cold stress responses.
21	g25112	AT5G35980	YAK1	YAK1-related. Controls cell cycle and regulates drought tolerance[65,66].
22	g40349	AT4G11110	SPA2	SPA1-related 2. Convergent with WGD adaptation in *A. arenosa*[7].
23	g54387	AT3G01420	DOX1	An alpha-dioxygenase involved in protection against oxidative stress.
24	g25300	AT5G34940	GUS3	GLUCURONIDASE 3
25	g45090	AT1G65320	CBSX6	Cystathionine beta-synthase family protein
15 mav	g39361	AT5G55820	INCEP / WYRD	**INNER CENTROMERE PROTEIN**. The largest subunit of the Chromosome Passenger Complex (CPC), and directly binds to all other subunits in animals and yeast[67]. The CPC ensures that all kinetochores are attached to microtubules emanating from opposing poles. INCENP is necessary for normal mitotic divisions[68]. At mitosis and meiosis localises to kinetochores, and later, the phragmoplast[67].
12 mav	g25323	AT4G29090		Ribonuclease H-like superfamily protein
11 mav	g23778	AT4G25290		DNA photolyase
11 mav	g54385	AT1G56145	CORK1	A LRR receptor kinase required for cellooligomer-induced responses.
10 mav	g33400	AT1G50240	FU	**FUSED**. An ARM repeat domain-containing protein kinase involved in male meiosis. Tightly localized to nascent phragmoplast and with the expanding phragmoplast ring.

226 **Table 1 | Top selective sweep candidates following WGD in *Cochlearia*.** Top 25 of 40,245 genes
227 assessed, with 5 additional genes (bottom) with 10 or more MAV SNPs and in top 1% F_{ST} tail. * = contains
228 37/39 whole-genome fixed differences between diploids and tetraploids and is also the only gene with three
229 of the top 25 Dxy windows. Genome-wide F_{ST} rank is given in Column 1.
230

Figure 4 | Selective sweep signatures of DNA management and ion homeostasis alleles. (A) Ploidy-specific differentiation across the *Cochlearia* genome. The dashed orange line gives the extreme stringency F_{ST} cutoff of the top 25 outlier genes (of 40,245 assessed); the solid line gives a 1% F_{ST} cutoff; **(B-G)** Examples of selective sweep signal among 6 top outlier genes; **(B-E)** Represent kinetochore or DNA management; **(F-G)** ion homeostasis functional categories. The X axis gives genome position in megabases (Mb). The Y axis gives AFD values at single SNPs (dots) between diploid and tetraploid *Cochlearia*. Orange arrows indicate genes overlapping the top F_{ST} gene outliers, and grey lines indicate neighboring genes. Orange dots indicate MAV outliers.

Functional processes under selection post-WGD in *Cochlearia*. While we focus our
discussion below to the top 25 genome-wide outliers, a broader, 1% F_{ST} list of 753 selective sweep
candidates (Dataset S3) yield particularly informative gene ontology (GO) enrichments, with 181
significantly enriched categories (using a conservative 'elim' Exact Test; Dataset S4 Gene
Ontology Enrichment Cochlearia). Many of the enriched categories can be grouped into three
classes congruent with WGD-associated changes[1,2,4-7,69]: 1) DNA management: 29 categories
relate to DNA integration, cell division, meiotic chromosome segregation, mitosis, DNA repair,
and recombination; 2) Ion homeostasis: 26 categories relate to ion transport (principally
extrusion), cation homeostasis, salt stress, and stomata; and 3) Cell stoichiometry: 7 categories
relate to global gene expression, cell wall, and biosynthetics, pointing to both global gene
expression and 'nucleotypic'[6] changes upon WGD. These categories often overlap, for example
'cell stoichiometry' and 'DNA management' in terms of global RNA transcriptional changes post-
WGD, which should alter due to doubled DNA template.

Kinetochore evolution upon WGD. Genome-wide, the most dramatic selection signature is
directly over the coding region of the *CENTROMERE PROTEIN E* ortholog (*CENP-E*; Fig. 4A,
B)[70]. This gene overlaps the 5 top F_{ST} outlier windows (mean F_{ST} = 0.88) and includes 37 of the
39 genome-wide fixed differences between ploidies. An essential kinetochore protein, *CENP-E*
moves mono-oriented chromosomes to the spindle equator, mediating congression[51,71,72].
Strikingly, tetraploid cancers are far more susceptible to CENP-E inhibitors than diploids[52-54,73,74].

The *Cochlearia* CENP-E coding region contains 26 SNPs (synonymous or non-
synonymous) that are highly differentiated between diploids and tetraploids (>50% AFD). Six of
these are unique to *Cochlearia* tetraploids, at highly conserved sites across angiosperms (Fig.
5A). None are in characterized conserved functional regions of the kinase domain[75], meaning
motor activity is likely intact. Most of the tetraploid-specific changes are in the coiled coil regions,
which in animals are important for regulation of cell division via phosphorylation and protein-
protein interactions[51]. For example, point mutations in the coiled coils are associated with human
disease (e.g., microcephaly[76,77]). In humans and Xenopus, these regions are known to be
extensively phosphorylated during the cell cycle and may be involved in the autoinhibition of
CENP-E[78-81]. Indeed, we see four tetraploid-specific changes that may affect regulation via the
loss of phosphorylation (S717A, S821A, S1059L and S1169). Three additional changes toward
the C-terminus are in a cargo (chromosomes) binding region. Four of the tetraploid-specific
changes show remarkable conservation across plants, being otherwise absolutely conserved
across CENP-E-like kinesins (Fig. 5A; A607G, Q613D, R899Q and Q1024E). Taken together,

these data suggest changed regulation of CENP-E at mitosis and/or meiosis. This is consistent

with functional evidence from *A. thaliana* showing that *CENP-E* mutations extend the cell cycle[81].

Figure 5 | Tetraploid-derived protein structure changes in CENP-E and HKT1. (A) Kinetochore subunit
CENP-E structural prediction and pan-plant diversity alignment. Structure: diploid structure (blue),
consensus (>50% allele frequency) derived polymorphisms in tetraploids (orange), kinesin motor domain
(slate). Alignment: colour by percentage identity; consensus mutations at highly conserved sites (orange
boxes); **(B-D)** Ion channel HKT1: diploid structure (blue), mutation sites (orange; diploids, magenta:
tetraploids), pore domains (teal); **(C)** HKT1 extracellular surface. Salinity tolerance-mediating change
L344V in red and selectivity filter residues S/G1, G2, G3 and G4 (black); **(D)** superimposition of diploid
(blue) and tetraploid (grey) intracellular domains. Structures that are predicted to have rotated are
highlighted on the tetraploid structure (dark grey). Tetraploid consensus mutations (magenta).

Our top outlier list (Table 1) contains two additional orthologs of *A. thaliana* kinetochore

components: *CENP-C*, an essential kinetochore component in both mitosis and meiosis, needed

for centromere identity in plants, yeast, *Drosophila*, and humans[58,59,82] (Fig. 4C) and *INNER*

CENTROMERE PROTEIN (INCENP), which controls mitotic and meiotic chromosome

segregation and cytokinesis in plants, yeast and animals[68,83]. At mitosis and meiosis *INCENP*

localizes to kinetochores, and later, the phragmoplast as the main subunit of the Chromosome

Passenger Complex[67]. Both INCENP and CENP-C contain 1% outlier-MAV SNPs, with *INCENP*

harboring a remarkable 15 MAV outlier SNPs, the greatest number of any gene in the genome.

Evolution of DNA repair and transcription. Several of the top signals of selective sweeps are

in DNA repair-related genes, for example in *CONDENSIN-II COMPLEX SUBUNIT*

H2/HYPERSENSITIVITY TO EXCESS BORON 2 (Fig. 4D), which functions directly in double-

strand break (DSB) repair[62,84] and chromatin management in plants, mouse and *Drosophila[61,85]*.

Condensin II is required for proper DNA DSB repair by homologous recombination (HR) repair in

A. thaliana and humans[62,63], and has been implicated association and dissociation of

centromeres[85]. We find in our 1% F_{ST} outliers an outlier also at a homolog of *DAYSLEEPER*, an

essential domesticated transposase[86]. *DAYSLEEPER* binds the Kubox1 motif upstream of the

DNA repair gene *Ku70* to regulate non-homologous end joining double-strand break repair, the

only alternative to HR[87,88].

In a young polyploid, the total DNA content doubles but the protein content and cell size

does not scale accordingly[6], so we predicted that the control of global gene expression, like

meiosis, should undergo adaptive evolution post-WGD. Here we see signal of this, with a suite of

DNA or RNA polymerase-associated genes among our selective sweep outliers. In our 1% F_{ST}

outliers this includes *NRPB9*, an RNA polymerase subunit that is implicated in transcription

initiation, processivity, fidelity, proofreading and DNA repair[89-93] as well as the ortholog of *MED13*,

of the mediator complex, which is essential for the production of nearly all cellular transcripts[94]

(Dataset S3).

Evolution of ion homeostasis, transport and stress signaling. The ionomic equilibrium of the

cell is immediately disrupted at WGD[41]; in particular K^+ concentrations are increased instantly,

consistent with increases in salinity tolerance in synthetic *A. thaliana* autotetraploids[41]. In our

young tetraploids, among the top selective sweeps are ion channels that function explicitly to

remove K^+, Na^+ and other cations from the cell. At F_{ST} rank 13 genome-wide, we see the ortholog

of *HIGH-AFFINITY K^+ TRANSPORTER[64,95] (HKT1;* Fig 4G, 5B*),* and at rank 15, *SALT OVERLY*

SENSITIVE 1 (SOS1), a membrane Na^+/H^+ transporter that removes excessive Na^+ from the cell and is central to salt tolerance[96,97] (Fig 4F). Several studies have demonstrated adaptive natural variation in response to salinity in *HKT1*[64,95], but among natural systems there have been to date no works until now that implicate *SOS1,* although this gene is central to salt tolerance pathways (discussed in [98]).

Our modeling confirms that the *Cochlearia HKT* under selection is class 1 from its selectivity filter residue configuration (S-G-G-G), indicating it is likely Na^+ selective (and K^+ non-selective)[99]. There is a L344V mutation in the tetraploid relative to the diploid; remarkably, this is the identical site and amino acid change that is associated with salt tolerance in rice OsHKT1;5 (Fig 5B)[100,101]: functional confirmation in rice shows the orthologous site substitution to valine in our tetraploid to be associated with salt tolerance (including faster Na^+ transport), while the diploid leucine is associated with salt sensitivity (including slower Na^+ transport)[100]. Given that the tetraploids live overwhelmingly in coastal regions where they are exposed to extreme Na levels, while the diploid lives in low Na freshwater streams, this makes biological sense. While the close proximity of L344 alone is likely enough to disrupt pore rigidity via its larger side chain relative to V, in *Cochlearia* HKT1 we also see T220M (a highly conserved residue) and I346F mutations which are even closer to the P1-4 pore domains that hold S-G2-G3-G4 together to create the selectivity control. These mutations all introduce large side chains that likely affect structural dynamics at the P1-4 pore[100,101]. In addition, mutations F326I, R200P L180F, Q303H, M360I are all in (R200P and F326I) or in contact with the four alpha helices that stabilize the SGGG selectivity filter. While this may seem an excessive quantity of mutations, suggesting gene inactivation and relaxation of selection, all the sites except M360I, T220M and L344V are loosly conserved, suggesting flexibility in these regions.

There are also mutations on the cytosolic side of the protein, a few poorly conserved residues appear to induce small structural changes (Fig 5D). This includes a cluster of changed residues (I388N, S386T, N384K, R396G) which are predicted to break up an alpha helicase in the tetraploid, and P18S and S19R, which appear to induce a break in the first alpha helicase of the protein. To our knowledge this domain is not functionally characterized, though given this positioning, this could represent a change in signaling or regulation.

Congruent with its conserved central role in ion homeostasis, *SOS1* is highly conserved and tetraploid-specific changes are not near active transport (nucleotide or ion) binding sites, nor dimerization domains. Instead, a tight cluster of 3 mutations marks the boundary between the β-sheet-rich cytoplasmic domain (β-CTD) and the C-terminal autoinhibition (CTA) domain which contains a further 3 mutations (K1014E, T1075S and R1101Q). The CTA is unstructured;

however, it has been experimentally shown that truncation, or just two point mutations, can

radically change the behavior of the channel, presumably by releasing autoinhibition increasing

Na$^+$ transport and therefore increasing salt tolerance[102]. Notably, this includes one T1075S

substitution: S and T are two of the three amino acids that can be phosphorylated and it has been

shown that this exact residue behaves differently when phosphorylated, which suggests that the

choice of S or T may have evolutionary consequences[103].

Ion homeostasis shifts should be associated with changes in responses to salt, osmotic,

and cold stress, as all these stressors have a common osmotic basis. Such a link between

immediate ionomic changes in the polyploid cell may be a key functional basis for the observed

ecotypic differentiation of young polyploids, especially as observed in arctic and alpine conditions.

We accordingly see in our top outliers categories of relevant genes, for example among the top

candidates the ortholog of *DEAD-BOX RNA HELICASE 25 (STRS2)*, identified in *A. thaliana* as

a repressor of stress signaling for salt, osmotic, and cold stress[104,105]. This gene also controls

freezing tolerance[106], highly relevant to the cold-loving arctic and alpine history of *Cochlearia*[14].

To assess if perhaps these sweeps were better associated with ecotype differences

between ploidies, we performed a salt tolerance experiment on diploid and tetraploid plants.

Interestingly, given their divergent ecotype preferences (Fig. S4), with tetraploids found in more

saline conditions, we found that the diploid *Cochlearia* are in fact more salt tolerant than the

tetraploids (p = 2.178 x 10^{-05}; See Supplementary Text 1 and Table S5.). This finding also

contrasts with observations of increased salinity tolerance in neotetraploid *Arabidopsis thaliana*[41].

Again, however, this may be a signal of preadaptation to osmotic challenge (common to freezing,

salinity, and dehydration) across the halophyte *Cochlearia*[42].

Relevant also to these phenotypes, genes involved in stomatal function were outliers post-

WGD, such as the ortholog of *OPEN STOMATA2*, a target of ABA stress signaling to close the

stomata during drought response[107]. This gene is an ATPase in the plasma membrane that drives

hyperpolarization and initiates stomatal opening[105]. We also see *TOO MANY MOUTHS*, where a

mutation leads to disruption of asymmetric cell division during stomata development[108]. Finally,

we see selection signal in *STOMAGEN*, which acts on the epidermis to increase stomatal

formation[109]. Sweeps in these loci are consistent with the phenotypic shifts we observe of

increased stomatal conductance and net photosynthetic rate under drought conditions in

tetraploid *Cochlearia* populations relative to diploids (Fig. 2E; Supplementary Text 2).

Gene-level convergence. To test for convergence at the ortholog level, we first determined

orthogroups[110] between *Cochlearia, A. arenosa,* and *C. amara* (Methods). Top 1% F$_{ST}$ outliers for

Cochlearia (n=753; Dataset S3), *A. arenosa* (n=452; Dataset S5), and *C. amara* (n=229; Dataset

S6) were considered orthologues if they were part of the same orthogroup. By this criterion not a

single ortholog was under selection in all three species (Fig. S6A; Dataset S7 Gene Ortholog

Convergence). This approach depends on strict 3-way orthogroup assignment, so we then

searched for convergence by assigning all genes in the outlier lists to a nearest *A. thaliana*

homolog. By this 'nearest homolog' criterion, only one gene was selected in all three WGDs:

DAYSLEEPER, an essential domesticated transposase[86] with a role in regulating non-

homologous end joining double-strand break repair (Fig. S6B; Dataset S8).

Interestingly, by both homolog assignment methods, several of the best *Cochlearia* WGD

adaptation candidates are candidates also in *A. arenosa*: *ASY3,* functionally validated[9,12] to have

an primary role in stabilizing autotetraploid meiosis in *A. arenosa* is also in our 1% F_{ST} outlier list

in *Cochlearia*[9]. This gene, along with *ASY1*, is critical for formation of meiotic chromosome axes,

and tetraploid alleles of both genes result in fewer deleterious multichromosome associations and

more rod-shaped bivalents in metaphase I[12]. We also see *CYCD5;1*, which is a QTL for

endoreduplication[111]. Additionally, the salinity and osmotic genes *HKT1* and *OST2* in both top

candidate lists (Fig. S6A; Dataset S7 Gene Ortholog Convergence). All of these genes are

involved in processes that have been implicated in adaptation to WGD[1,2,4-7] and therefore stand

as good candidates in salient challenges to nascent polyploids. We note that overlap between

Cochlearia and both *A. arenosa* and *C. amara* candidates was greater than expected by chance

(SuperExactTest p=0.0024 and p=0.0047 respectively), but only marginally for *C. amara* and *A.*

arenosa (SuperExactTest p=0.014).

Process-level convergence. We reasoned that there may be similarities in processes under

selection between the three independent WGDs, despite modest gene-level convergence. To

estimate this, we first compared our GO results from those published in *A. arenosa* and *C. amara*.

The much greater signal of overlap (process-level convergence; Fig. S6C) was between

Cochlearia and *A. arenosa*: of the 113 GO biological process terms significantly enriched in

Cochlearia, 17 were among the 73 GO terms enriched in *A. arenosa* (Dataset S4). These were

high-level GO terms including representatives of ploidy-relevant categories, e.g. 'cell division',

'transmembrane transport, and 'regulation of RNA metabolic processes'.

Despite this evident convergence, in *Cochlearia* an array of DNA repair and kinetochore

genes were among top candidates, signaling a shift relative to *A. arenosa*, where a more focused

prophase I-oriented signal emerged primarily around Synaptonemal Complex (SC)-associated

proteins mediating lower crossover rates[7]. However, given that *CENP-C* mutants fail to retain an

SC at the centromere and *CENP-C* appears to have functions in synapsis, cohesion, and
centromere clustering[112], there may be a closer parallel between the obviously 'SC- focused'
adaptive response in *A. arenosa* and that in *Cochlearia*. Indeed, aside from *ASY3*, in the
Cochlearia 1% F_{ST} outlier list we also see *ATAXIA-TELANGIECTASIA MUTATED (ATM;* also with
6 MAV SNPs; Dataset S3). This gene controls meiotic DNA double-strand break formation and
recombination and affects synaptonemal complex organization[113]. We also see a *PDS5* cohesion
cofactor ortholog and *REC Q MEDIATED INSTABILITY 1* (*RMI1*), which suppresses somatic
crossovers and is essential for resolution of meiotic recombination intermediates[114]. Notable also
are *BRCA2-like B* (which is essential at meiosis and interacts with *Rad51, Dss1,* and *Dmc1*) and
SHUGOSHIN C (which protects meiotic centromere cohesion)*.*

Further evidence of functional association outliers found in *Cochlearia* with those in *A.*
arenosa can be observed in protein interaction information from the STRING database, which
provides an estimate of proteins' joint contributions to a shared function[115]. Using comparable 1%
F_{ST} outlier lists from the two species, we see many connections between these independent
WGDs. Here we see particularly large clusters (Fig. 6A,B) in center of the overall network (Fig.
6C). In particular, the endopolyploidy gene[111] *CYCD5;1,* and *DNA Pol V,* a shared outlier in both
species, interact with a broad array of other outliers in each selection scan (Fig. 6A). This analysis
reveals, for example, that *DNA Pol V* shares as a top candidate either *NRPB9A* (in *A. arenosa*)
or *NRPB9B* (in *Cochlearia*). These subunits are partly redundant interactors with *Pol II, IV*, and *V*
and the double mutant is fatal in *A. thaliana*[91].

Taken together, these results indicate that adaptive evolution in response to WGD is
focused on particular functions instead of specific genes. These functions involve DNA
management and ion homeostasis. However, it is unclear why particular solutions are favored in
one species relative to another. A degree of stochasticity depending on available standing
variation can be expected, but species histories likely play a role, offering preadaptations that may
'nudge' evolution. For example, our analysis of salinity tolerance in *Cochlearia* gave the surprising
result that the diploid was at least as tolerant to extreme salt concentrations as the tetraploid,
although the diploid is found predominantly inland, except for rare diploid coastal populations from
Spain. A postglacial and boreal spread of the diploid towards the UK may have brought salinity
and cold tolerance along the way[116], altering the genomic substrate upon which selection acted
in response to WGD-associated ionomic challenge.

Figure 6 | Evidence for functional convergence between *Cochlearia* and *Arabidopsis arenosa* following independent WGDs. STRING[115] plots show *Cochlearia* candidate genes (blue) and *A. arenosa* candidate genes (red). Convergent genes that are present in both species' outlier lists as selection candidates are in yellow. **(A)** A large shared cluster surrounding the endopolyploidy gene *CYCD5;1*, which has many connections to large cluster centering on *DNA pol V*, which is an outlier in both datasets; **(B)** ion transport-related genes with a highly interconnected cluster of top outliers in both genome scans, *HKT1, KUP9,* and *HA1;* **(C)** The entire set of candidates in both genome scans for which the STRING database has information.

Conclusion

Following whole genome duplication (WGD), the newly polyploid cell requires modifications to chromosome segregation, ion homeostasis, stomatal function, and diverse other processes[1,2,4,6]. Here we investigated the signals of adaptive evolution post-WGD in a successful novel polyploid system, *Cochlearia*. We discovered striking signals of selection in core kinetochore components, *CENP-E, CENP-C, INCEP, CAP-H2,* and others, as well as well-studied ion homeostasis loci, *HKT1* and *SOS1.* We detail specific changes in these proteins upon WGD which are known in model systems to directly modulate WGD-relevant function. We also compare our results to independent WGD adaptation events, finding convergence of these processes, but not genes, indicating a highly flexible array of adaptive mechanisms.

Our results also suggest a hypothesis for the occasionally spectacular adaptability of polyploids. We observed a broad array of DNA management and repair processes under selection in all species, and especially *Cochlearia*. This may signal a temporarily increased post-WGD susceptibility to DNA damage, due to suboptimal function of DNA repair genes during the process of adaptation to the WGD state. This may result in a relative 'mutator phenotype' in neopolyploids. Such a mutator phenotype has been plainly observed in polyploid metastatic human cancers, which not only exhibit SNP-level hypermutator phenotypes, but also dramatic structural variation in malignant aneuploid swarms that are associated with progression[3]. We speculate that a parallel to this may exists following other WGDs. Whether this hypothesis is further supported by future discoveries, the centrality of WGD to evolution, ecology and agriculture, underscores the importance of understanding the processes mediating adaptation to—and perhaps also by—WGD.

Methods

Plant material. We first located 89 *Cochlearia* populations throughout Europe and collected population samplings of plants from each, aiming for at least 10 plants per population, with each sampled plant a minimum of 2 meters from any other. Of these, we selected 33 geographically and ecotypically diverse representative populations for population resequencing, including the outgroup *Ionopsidium* (Dataset S2 Sample Metrics). An average of 4 individuals per population were sequenced. A total of 149 individuals were initially sequenced, which was narrowed down by a cutoff of minimum average depth of 4x, leaving 113 individuals from 33 populations in the final analysed dataset, including the outgroup *Ionopsidium*.

Ploidy Determination by flow cytometry. DNA content and ploidy were inferred for populations using flow cytometry (Dataset S1). Approximately 1 square cm of leaf material was diced alongside an internal reference using razor blades in 1 ml ice cold extraction buffer (either 45 mM $MgCl_2$, 30 mM sodium citrate, 20mM MOPS, 1% Triton-100, pH 7 with NaOH for relative staining or 0.1 M citric acid, 0.5% Tween 20 for absolute measurements). The resultant slurry was then filtered through a 40-μm nylon mesh before the nuclei were stained with the addition of 1 ml staining buffer (either CyStain UV precise P [Sysmex, Fluorescence emission: 435nm to 500nm] for relative ploidy, or Otto 2 buffer [0.4 M $Na_2HPO_4 \cdot 12H_2O$, Propidium iodide 50 μgmL−1, RNase 50 μgmL−1], for absolute DNA content). After 1 minute of incubation at room temperature the sample was run for 5,000 particles on either a Partec PA II flow cytometer or a BD FACS Melody. Histograms were evaluated using FlowJo software version 10.6.1.

HEI10 immunostaining. Pachytene chromosome spreads were prepared from fixed anthers, as described previously[117]. Immunostaining was conducted using two primary antibodies: anti-AtZYP1C rat, 1:500[118] and anti-HvHEI10 rabbit, 1:250[119], followed by two secondary antibodies: goat anti-rat Alexa Fluor® 594 (Invitrogen) and goat anti-rabbit Alexa Fluor® 488 (Invitrogen). A Nikon Eclipse Ci fluorescence microscope equipped with NIS elements software was used to capture and quantify images. HEI10 foci were counted at pachytene using NIS software and significance established using the Mann-Whitney U test (Minitab v 18.1.0.0).

Fluorescence *in situ* hybridization. Mitotic chromosome spreads from fixed root tips were prepared as described previously[120]. The *Arabidopsis*-type telomere repeat (TTTAGGG)n was prepared according to [121]. The *Cochlearia*-specific 102-bp (GTTAGATGTTTCATAAGTTCGTCAA ACTTGTACAAAGCTCATTGAGACACTTATAAGCACTCATGTTGCATGAAACTTGGTTTAGAG TCCTAGAAACGCGTT) tandem repeat was designed and prepared based on Mandáková et al. (2013) and used for identification of centromeres. The DNA probes were labeled by nick translation with biotin-dUTP and digoxigenin-dUTP according to [122], pooled and precipitated by adding 1/10 volume of 3 M sodium acetate, pH 5.2, and 2.5 volumes of ice-cold 96% ethanol and kept at -20°C for 30 min. The pellet was then centrifuged at 13,000 g at 4°C for 30 min. The pellet was resuspended in 20 μl of the hybridization mix (50% formamide and 10% dextran sulfate in 2×SSC) per slide. 20 μl of the probe was pipetted onto a chromosome-containing slide. The cover slips were framed with rubber cement. The probe and chromosomes were denatured together on a hot plate at 80°C for 2 min and incubated in a moist chamber at 37°C overnight. Post-hybridization washing was performed according to [122]. After immunodetection, chromosomes

528 were counterstained with 4', 6-diamidino-2-phenylindole (DAPI, 2 µg/ml) in Vectashield (Vector
529 Laboratories). The preparations were photographed using a Zeiss Axioimager Z2 epifluorescence
530 microscope with a CoolCube camera (MetaSystems). The three monochromatic images were
531 pseudocolored, merged and cropped using Photoshop CS (Adobe Systems) and Image J
532 (National Institutes of Health) softwares.
533
534 **Reference Genome Assembly and Alignment.** We generated a long read-based *de*
535 *novo* genome assembly using Oxford Nanopore and Hi-C approaches, below.
536
High Molecular Weight DNA isolation and Oxford Nanopore sequencing. A total of 0.4 g
Cochlearia excelsa leaf material from one individual plant was ground using liquid nitrogen before
the addition of 10 ml of CTAB DNA extraction buffer (100 mM Tris-HCl, 2% CTAB, 1.4 M NaCl,
20 mM EDTA, and 0.004 mg/ml Proteinase K). The mixture was incubated at 55°C for 1 hour then
cooled on ice before the addition of 5 ml Chloroform. This was then centrifuged at 3000 rpm for
30 minutes and the upper phase taken, this was added to 1X volume of
phenol:chloroform:isoamyl-alcohol and spun for 30 minutes at 3000 rpm. Again, the upper phase
was taken and mixed with a 10% volume of 3M NaOAc and 2.5X volume of 100% ethanol at 4
°C. This was incubated on ice for 30 minutes before being centrifuged for 30 minutes at 3000 rpm
and 4 °C. Three times the pellet was washed in 4ml 70% ethanol at 4 °C before being centrifuged
again for 10 minutes at 3000 rpm and 4°C. The pellet was then air dried and resuspend in 300 µl
nuclease-free water containing 0.0036 mg/ml RNase A. The quantity and quality of high molecular
weight DNA was checked on a Qubit Fluorometer 2.0 (Invitrogen) using the Qubit dsDNA HS
Assay kit. Fragment sizes were assessed using a Q-card (OpGen Argus) and the Genomic DNA
Tapestation assay (Agilent). Removal of short DNA fragments and final purification to HMW DNA
was performed with the Circulomics Short Read Eliminator XS kit.
553
Long read libraries were prepared using the Genomic DNA by Ligation kit (SQK-LSK109; Oxford
Nanopore Technologies) following the manufacturer's procedure. Libraries were then loaded onto
a R9.4.1 PromethION Flow Cell (Oxford Nanopore Technologies) and run on a PromethION Beta
sequencer. Due to the rapid accumulation of blocked flow cell pores or due to apparent read
length anomalies on some *Cochlearia* runs, flow cells used in runs were treated with a nuclease
flush to digest blocking DNA fragments before loading with fresh library according to the Oxford
Nanopore Technologies Nuclease Flush protocol, version NFL_9076_v109_revD_08Oct2018.
561
562 **Genome size estimation and computational ploidy inference.** We used KMC[123] to create a k-
563 mer frequency spectrum (Kmer length=21) of trimmed Illumina reads. We then used
564 GenomeScope 2.0 (parameters: -k 21 -m 61) and Smudgeplot[124] to estimate genome size and
565 heterozygosity from k-mer spectra.
566
Data processing and assembly. Fast5 sequences produced by PromethION sequencing were
base called using the Guppy 6 high accuracy base calling model (dna_r9.4.1_450bps_hac.cfg)
and the resulting fastq files were quality filtered by the base caller. A total of 17.2 GB base called
data were generated for the primary assembly, resulting in 60x expected coverage. Primary

assembly was performed in Flye[32] and Necat[33]. The contigs were polished to improve the single-
base accuracy in a single round of polishing in Medaka[34] and Pilon[35].

Pseudomolecule construction by Hi-C, assembly cleanup, and polishing. To scaffold the
assembled contigs into pseudomolecules, we performed chromosome conformation capture
using HiC. Leaves from a single plant were snap-frozen in liquid N and ground to a fine powder
using mortar and pestle. The sample was then homogenised, cross-linked and shipped to Phase
Genomics (Seattle, USA), who prepared and sequenced an in vivo Hi-C library. After filtering low-
quality reads with Trimmomatic[125], we aligned the Hi-C reads against the contig-level assembly
using bwa-mem[126] (settings -5 -S -P) and removed PCR duplicated using Picard Tools
(https://broadinstitute.github.io/picard/). We used 3D-DNA[127] to conduct the initial scaffolding,
followed by a manual curation in Juicebox[128]. After manually assigning chromosome boundaries,
we searched for centromeric and telomeric repeats to orient the chromosome arms and to assess
the completeness of the assembled pseudomolecules. To identify the centromeric repeat motif in
C. excelsa, we used the RepeatExplorer[129] pipeline to search for repetitive elements from short-
read sequence data originating from the reference individual. RepeatExplorer discovered a highly
abundant 102 nucleotide repeat element (comprising 21% of the short-read sequence), which we
confirmed as the centromeric repeat motif by fluorescence in situ hybridisation. Using BLAST, we
localised the centromeric and telomeric (TTTAGGG) repeats and used them to orient the
chromosome arms. We performed a final assembly cleanup in Blobtools[36] (Fig.S1). Gene space
completeness was assessed using BUSCO version 3.0.2)[37].

Assembly annotation and RNA-seq. Prior to gene annotation, we identified and masked
transposable element (TE) sequences from the genome assembly. To do so, we used the EDTA
pipeline[130], which combines multiple methods to comprehensively identify both retrotransposons
and DNA transposons. After running EDTA on our chromosome-level genome assembly, we
performed BLAST queries against a curated protein database from Swiss-Prot to remove putative
gene sequences from the TE library and masked the remaining sequences from the assembly
using RepeatMasker (https://www.repeatmasker.org).

We then used the BRAKER2[38] pipeline to conduct gene annotataion on the TE-masked genome
assembly. Evidence types included RNAseq data from the identical *C. excelsa* line and protein
data from related species. RNA-seq was generated from bud, stem and leaf tissue. Total RNA
was extracted from each tissue using the Qiagen RNeasy Extraction Kit. Stranded RNA libraries
with polyA were constructed Using NEB Next Ultra II Directional RNA Library Prep Kit for Illumina
and then evaluated by qPCR, TapeStation and Qubit at the DeepSeq facility (Nottingham, UK)
before being sequenced at PE 150 at Novogene, inc (Cambridge, UK). We mapped the RNA-seq
reads of each tissue to our reference genome using STAR[131] with default parameters (-
twopassMode Basic) before running BRAKER2. Running BRAKER2 without UTR prediction
generated more gene models and much better BUSCO metrics than with UTR prediction (97.8%
[raw, pre-Blobtools trimmed] complete BUSCOs without UTR prediction vs 91.7% with UTR
prediction), so for the final annotation we used the more complete set and ran BRAKER2 without
UTR prediction.

Population Resequencing and Analysis.

Library preparation and sequencing. DNA was prepared using the commercially available DNeasy Plant Mini Kit from Qiagen. DNA libraries were made using TruSeq DNA PCR-free Library kit from Illumina as per the manufacturer's instructions and were multiplexed based on concentrations measured with a Qubit Flourometer 2.0 (Invitrogen) using the Qubit dsDNA HS Assay kit. Sequencing was carried out on either NextSeq 550 (Illumina) in house (4 runs) or sent to Novogene for Illumina Hiseq X, PE150 sequencing (2 runs).

Data preparation, alignment, and genotyping. Reads were quality trimmed with Trimmomatic 0.39[125] (PE -phred33 LEADING:10 TRAILING:10 SLIDINGWINDOW:4:15 MINLEN:50) and then aligned to the *C. excelsa* reference using bwa-mem[132] and further processed with samtools[133]. Duplicate reads were removed and read group IDs added to the bam files using Picard (version 1.134). Indels were realigned with GATK (version 4.2.3.0)[134]. Samples were first genotyped individually with "HaplotypeCaller" (--emit-ref-confidence BP_RESOLUTION --min-base-quality-score 25 --minimum-mapping-quality 25) and were then genotyped jointly using "GenotypeGVCFs" in GATK (version 4.2.3.0). The resulting VCF files were then filtered for biallelic sites and mapping quality (QD < 2.0, FS > 60.0, MQ < 40.0, MQRankSum < -12.5, ReadPosRankSum < -8.0, HaplotypeScore < 13.0). The VCF was then filtered by depth. Based on this distribution a depth cutoff of 4,322 was applied to the VCF containing the dataset and this was then used as a mask for the final VCF containing all individuals.

Demographic analysis. We first inferred genetic relationships between individuals using principal component analysis (PCA). Following [135], we estimated a matrix of genetic covariances between each pair of individuals. For two individuals, i and j, covariance (C) was calculated as:

$$C_{ij} = \frac{1}{m} \sum_{s=1}^{m} \frac{(g_{is} - x_{ij}p_s)(g_{js} - x_{ij}p_s)}{x_{ij}p_s(1 - p_s)},$$

where m is the number of variable sites, g_{is} is the genotype of individual i in site s, x is the average ploidy level of the two individuals, and p is the alternate allele frequency. PCA was performed on the matrix using the R function *prcomp*, setting scaling to TRUE, and first two axes of the rotated data extracted for plotting. For fastSTRUCTURE we followed [136] by randomly subsampling two alleles from tetraploid and hexaploid populations using a custom script. We have previously demonstrated that results generated in this way are directly comparable to results generated with the full dataset in STRUCTURE[136]. We calculated Nei's distances among all individuals in StAMPP[137] and visualized these using SplitsTree[39]. Linkage disequilibrium was estimated using ldsep[138]. To avoid biasing the estimates with unequal sample sizes, we chose 39 diploids and tetraploids for the analysis. To reduce computation time, the analysis was performed on 4-fold sites from a single chromosome (chromosome 1). To visualize the decay of LD as a function of physical distance, we calculated average r^2 in 10 bp non-overlapping windows and fit a loess curve on the binned data.

Window-based scan for selective sweep signatures. We performed a window-based divergence scan for selection consisting of 1 kb windows that contained at least 15 SNPs. The

data were filtered as described above and in addition was filtered for no more than 20% missing
data and a depth of >= 8x. We calculated the following metrics: Rho[45], Nei's F_{ST}[47], Weir-Cochran's
F_{ST}[48], Hudson's F_{ST}[46], Dxy[44], number of fixed differences and average groupwise allele frequency
difference (AFD). To determine the best metric to detect localised peaks of divergence we
performed a quantitative analysis of AFD plot quality for all 1% outliers of each metric. Each
window was given a score of 0-4, with 0 being the lowest quality and 4 the highest. Scores were
based on two qualities: peak height and peak specificity. For peak height one point was awarded
if the window contained one SNP of AFD between 0.5 and 0.7, and two points were awarded for
any SNP of AFD > 0.7. Likewise, for peak specificity two points were awarded for an AFD peak
that was restricted to a single gene and one point was awarded for a peak that was restricted to
2-3 genes. Compared to all other single 1% outlier lists and all permutations of overlapped 1%
outlier lists, the top 1% outliers from Hudson's F_{ST} performed most favorably as it maximized the
number of '4' and '3' scores while minimizing the number of '1' and '0' scores. Finally, we masked
from downstream analysis a region of uniformly high differentiation marking a suspected inversion
at scaffold 6 (between 5,890,246 and 6,137,362 bp).
673
MAV analysis. Following [13], we performed a FineMAV[50]-like analysis on all biallelic, non-
synonymous SNPs passing the same filters as the window-based selection scan. SNPs were
assigned a Grantham score according to the amino acid change and this was scaled by the AFD
between ploidies. SNPs were first filtered for a minimum AFD of 0.25. The top 1% outliers of all
these MAV-SNPs were then overlapped with the genes in our 1% F_{ST} outlier windows to give a
refined list of candidate genes that contain potentially functionally significant non-synonymous
mutations at high AFD between cytotypes. The code outlining this can be found at
https://github.com/paajanen/meiosis_protein_evolution/tree/master/FAAD.
682
Orthogrouping and Reciprocal Best Blast Hits. We performed an orthogroup analysis using
Orthofinder version 2.5.5[110]. to infer orthologous groups (OGs) from four species (*C. amara, A.*
lyrata, A. thaliana, C. excelsa). A total of 25,199 OGs were found. Best reciprocal blast hits (RBHs)
for *Cochlearia* and *A. thaliana* genes were found using BLAST version 2.9.0. *Cochlearia* genes
were then assigned an *A. thaliana* gene ID for GO enrichment analysis in one of five ways. First
if the genes' OG contained only one *A. thaliana* gene ID, that gene ID was used. If the OG
contained more than one *A. thaliana* gene ID then the RBH was taken. If there was no RBH then
the OG gene with the lowest E-value in a BLAST versus the TAIR10 database was taken. If no
OG contained the *Cochlearia* gene then the RBH was taken. Finally, if there was no OG or RBH
then the gene with the lowest E-value in a BLAST versus the TAIR10 database was taken.
BLASTs were performed using the TAIR10.1 genome with data generated on 2023-01-02.
694
GO Enrichment Analysis. To infer functions significantly associated with directional selection
following WGD, we performed gene ontology enrichment of candidate genes in the R package
TopGO v.2.52[139], using our *Cochlearia* universe set. We tested for overrepresented Gene
Ontology (GO) terms within the three domains Biological Process (BP), Cellular Component (CC)
and Molecular Function (MF) using Fisher's exact test with conservative 'elim' method, which
tests for enrichment of terms from the bottom of the hierarchy to the top and discards any genes
that are significantly enriched in a descendant GO term. We used a significance cut-off of 0.05.

Generation of Consensus Sequences. Consensus sequences were generated for proteins of interest so that they could be closely inspected via MSAs and 3D protein structure prediction. Genomic regions were selected for either all diploids or all tetraploids present in the selection scan with GATK SelectVariants, while simultaneously being filtered for biallelic SNPs, "--max-nocall-fraction 0.2" and "-select 'AF > 0.5'". A consensus sequence was generated for exons by combining samtools faidx and bcftools consensus. Finally, a VCF containing only non-biallelic variation was manually inspected and any multiallelic variants at AF>0.5 and max-nocall-fraction<0.2 manually incorporated into the consensus.

Multiple Sequence Alignments (MSAs). We generated multiple sequence alignments using Clustal-Omega[140] in combination with amino acid sequences from the GenBank database. Sequences were selected either because the genes/proteins were well studied in other organisms or to give a phylogenetically broad coverage. Alignments were manually refined and visualised in JalView[141].

Protein modeling. Protein homology models were created using AlphaFold[142] version 2.1 on the Czech national HPC MetaCentrum. The full database was uses with a model preset of monomer and a maximum template data of 2020-05-14. Structures were visualised and images generated in the PyMOL Molecular Graphics System (Version 2.0 Schrödinger, LLC).

Data Availability
Sequence data for this study have been deposited in the European Nucleotide Archive (ENA) at EMBL-EBI (https://www.ebi.ac.uk/ena/browser/view/PRJEB66308) under accession number PRJEB66308.

Acknowledgements
We thank Kirsten Bomblies, Lara Hebberecht-Lopez, Mary Bray and Nigel Bray for assistance collecting plant material. We further acknowledge wet lab support from Anna Loreth (Heidelberg). This work was supported by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program [grant number ERC-StG 679056 HOTSPOT], via a grant to LY and under the Marie Skłodowska-Curie grant agreement No. 101022295 to TH. Nanopore sequencing was performed in the University of Nottingham DeepSeq sequencing facility. We are grateful for access to the University of Nottingham's Augusta HPC service and the CESNET LM2015042 and the CERIT Scientific Cloud LM2015085, provided under the programme 'Projects of Large Research, Development, and Innovations Infrastructures'. We thank local authorities for issuing permission to collect samples (Slovak Ministry for Environment, permission No 062-219/18).

Author Contributions: LY and SMB conceived the study. LY, SMB, MK, PM, and JK performed collections. SMB, SB, SDD, T.Mandáková, CM, LC, MZ, and SF performed laboratory experiments. SMB, TH, SDD, CM TM, T.Mathers, T.Mandáková, EW, MB, SB, SF, PP, MK and LY performed analyses. LY and SMB wrote the manuscript. All authors approved the final manuscript.

746
747 **Competing Interests statement:** The authors declare no competing interests.

References

References

1   Baduel, P., Bray, S., Vallejo-Marin, M., Kolář, F. & Yant, L. The "Polyploid Hop": Shifting
Challenges and Opportunities Over the Evolutionary Lifespan of Genome Duplications.
Frontiers in Ecology and Evolution **6** (2018). https://doi.org:10.3389/fevo.2018.00117

2   Yant, L. & Bomblies, K. Genome management and mismanagement--cell-level
opportunities and challenges of whole-genome duplication. *Genes & Development* **29**,
2405-2419 (2015). https://doi.org:10.1101/gad.271072.115

3   Van de Peer, Y., Mizrachi, E. & Marchal, K. The evolutionary significance of polyploidy.
Nat Rev Genet **18**, 411-424 (2017). https://doi.org:10.1038/nrg.2017.26

4   Bomblies, K. When everything changes at once: finding a new normal after genome
duplication. *Proc Biol Sci* **287**, 20202154 (2020). https://doi.org:10.1098/rspb.2020.2154

5   Bomblies, K. Learning to tango with four (or more): the molecular basis of adaptation to
polyploid meiosis. *Plant Reprod* **36**, 107-124 (2023). https://doi.org:10.1007/s00497-022-
00448-1

6   Doyle, J. J. & Coate, J. E. Polyploidy, the Nucleotype, and Novelty: The Impact of Genome
Doubling on the Biology of the Cell. *International Journal of Plant Sciences* **180**, 1-52
(2019). https://doi.org:10.1086/700636

7   Yant, L. *et al.* Meiotic adaptation to genome duplication in Arabidopsis arenosa. *Current
Biology* **23**, 2151-2156 (2013). https://doi.org:10.1016/j.cub.2013.08.059

8   Hollister, J. D. *et al.* Genetic adaptation associated with genome-doubling in autotetraploid
Arabidopsis arenosa. *PLoS Genetics* **8**, e1003093 (2012).

9   Seear, P. J. *et al.* A novel allele of ASY3 is associated with greater meiotic stability in
autotetraploid Arabidopsis lyrata. *PLoS Genetics* **16**, e1008900 (2020).
https://doi.org:10.1371/journal.pgen.1008900

10   Morgan, C., Knight, E. & Bomblies, K. The meiotic cohesin subunit REC8 contributes to
multigenic adaptive evolution of autopolyploid meiosis in Arabidopsis arenosa. *PLoS
Genetics* **18**, e1010304 (2022).

11   Morgan, C. *et al.* Evolution of crossover interference enables stable autopolyploidy by
ensuring pairwise partner connections in Arabidopsis arenosa. *Current Biology* **31**, 4713-
4726 e4714 (2021). https://doi.org:10.1016/j.cub.2021.08.028

12   Morgan, C., Zhang, H., Henry, C. E., Franklin, F. C. H. & Bomblies, K. Derived alleles of
two axis proteins affect meiotic traits in autotetraploid Arabidopsis arenosa. *Proceedings
of the National Academy of Sciences* **117**, 8980-8988 (2020).

13   Bohutinska, M. *et al.* Novelty and Convergence in Adaptation to Whole Genome
Duplication. *Molecular Biology and Evolution* **38**, 3910-3924 (2021).
https://doi.org:10.1093/molbev/msab096

14   Wolf, E., Gaquerel, E., Scharmann, M., Yant, L. & Koch, M. A. Evolutionary footprints of a
cold relic in a rapidly warming world. *Elife* **10**, e71572 (2021).

15   Gill JJB, M. H., Fearn, GM Cytotaxonomic studies on the Cochlearia officinalis L. group
from inland stations in Britain. *Watsonia* **12**, 15-21 (1978).

16   Gill, J. Cytogenetic studies in Cochlearia L.(Cruciferae). The origins of C. officinalis L. and
C. micacea Marshall. *Genetica* **44**, 217-234 (1973).

17   Gill, J. Cytogenetic studies in Cochlearia L. The chromosomal homogeneity within both
the 2 n= 12 diploids and the 2 n= 14 diploids and the cytogenetic relationship between the
two chromosome levels. *Annals of Botany* **35**, 947-956 (1971).

18   Marburger, S. *et al.* Interspecific introgression mediates adaptation to whole genome
duplication. *Nature Communications* **10**, 5218 (2019). https://doi.org:10.1038/s41467-
019-13159-5

19   Arnold, B., Kim, S.-T. & Bomblies, K. Single geographic origin of a widespread
autotetraploid Arabidopsis arenosa lineage followed by interploidy admixture. *Molecular
Biology and Evolution* **32**, 1382-1395 (2015).

20 Fearn, G. A morphological and cytological investigation of Cochlearia populations on the
Gower Peninsula, Glamorgan. *New Phytologist* **79**, 455-458 (1977).

21 Stace, C. A. Hybridization and the Flora of the British Isles. *Botanical Society of the British
Isles* (1975).

22 Koch, M. Genetic differentiation and speciation in prealpine Cochlearia: Allohexaploid
Cochlearia bavarica Vogt (Brassicaceae) compared to its diploid ancestor Cochlearia
pyrenaica DC. in Germany and Austria. *Plant Systematics and Evolution* **232**, 35-49
(2002).

23 Koch, M. Zur Ausbreitung des Dänischen Löffelkrautes (Cochlearia danica L.) als
Küstensippe in das niedersächsische Binnenland. *Florist. Rundbr* **30**, 20-23 (1996).

24 Koch, M. Kurznotiz zur südlichen Ausbreitung des Dänischen Löffelkrauts (Cochlearia
danica L.) in Nordrhein-Westfalen. *Flor. Rundbr* **30**, 136-138 (1997).

25 Koch, M., Dobeš, C., Bernhardt, K.-G. & Kochjarová, J. Cochlearia macrorrhiza
(Brassicaceae): A bridging species between Cochlearia taxa from the Eastern Alps and
the Carpathians? *Plant Systematics and Evolution* **242**, 137-147 (2003).

26 Koch, M., Hurka, H. & Mummenhoff, K. Chloroplast DNA restriction site variation and
RAPD-analyses in Cochlearia (Brassicaceae): Biosystematics and speciation. *Nordic
Journal of Botany* **16**, 585-603 (1996).

27 Koch, M., Huthmann, M. & Hurka, H. Isozymes, speciation and evolution in the polyploid
complex Cochlearia L.(Brassicaceae). *Botanica Acta* **111**, 411-425 (1998).

28 Koch, M., Mummenhoff, K. & Hurka, H. Molecular phylogenetics of Cochlearia
(Brassicaceae) and allied genera based on nuclear ribosomal ITS DNA sequence analysis
contradict traditional concepts of their evolutionary relationship. *Plant Systematics and
Evolution* **216**, 207-230 (1999).

29 Koch, M. A. Mid-Miocene divergence of Ionopsidium and Cochlearia and its impact on the
systematics and biogeography of the tribe Cochlearieae (Brassicaceae). *Taxon* **61**, 76-92
(2012).

30 Koch, M. A. *et al.* The Quaternary evolutionary history of Bristol rock cress (Arabis scabra,
Brassicaceae), a Mediterranean element with an outpost in the north-western Atlantic
region. *Annals of Botany* **126**, 103-118 (2020).

31 Fekete, R., Mesterhazy, A., Valko, O. & Molnár V, A. A hitchhiker from the beach: the
spread of the maritime halophyte Cochlearia danica along salted continental roads. *Preslia*
90, 23-37 (2018).

32 Kolmogorov, M., Yuan, J., Lin, Y. & Pevzner, P. A. Assembly of long, error-prone reads
using repeat graphs. *Nature Biotechnology* **37**, 540-546 (2019).

33 Chen, Y. *et al.* Fast and accurate assembly of Nanopore reads via progressive error
correction and adaptive read selection. *BioRxiv*, 2020.2002. 2001.930107 (2020).

34 Medaka (Available from: https://github.com/nanoporetech/medaka).

35 Walker, B. J. *et al.* Pilon: an integrated tool for comprehensive microbial variant detection
and genome assembly improvement. *PloS one* **9**, e112963 (2014).

36 Laetsch, D. R. & Blaxter, M. L. BlobTools: Interrogation of genome assemblies.
F1000Research **6**, 1287 (2017).

37 Manni, M., Berkeley, M. R., Seppey, M., Simao, F. A. & Zdobnov, E. M. BUSCO Update:
Novel and Streamlined Workflows along with Broader and Deeper Phylogenetic Coverage
for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes. *Molecular Biology and
Evolution* **38**, 4647-4654 (2021). https://doi.org:10.1093/molbev/msab199

38 Brůna, T., Hoff, K. J., Lomsadze, A., Stanke, M. & Borodovsky, M. BRAKER2: automatic
eukaryotic genome annotation with GeneMark-EP+ and AUGUSTUS supported by a
protein database. *NAR genomics and bioinformatics* **3**, lqaa108 (2021).

39 Huson, D. H. SplitsTree: analyzing and visualizing evolutionary data. *Bioinformatics* **14**,
68-73 (1998).

40   Raj, A., Stephens, M. & Pritchard, J. K. fastSTRUCTURE: variational inference of
851   population structure in large SNP data sets. *Genetics* **197**, 573-589 (2014).

852   41   Chao, D. Y. *et al.* Polyploids exhibit higher potassium uptake and salinity tolerance in
853   Arabidopsis. *Science* **341**, 658-659 (2013). https://doi.org:10.1126/science.1240561

854   42   Nawaz, I., Iqbal, M., Bliek, M. & Schat, H. Salt and heavy metal tolerance and expression
855   levels of candidate tolerance genes among four extremophile Cochlearia species with
856   contrasting habitat preferences. *Sci Total Environ* **584-585**, 731-741 (2017).
857   https://doi.org:10.1016/j.scitotenv.2017.01.111

858   43   Pegtel, D. M. Effect of ploidy level on fruit morphology, seed germination and juvenile
859   growth in scurvy grass (Cochlearia officinalis L. sl, Brassicaceae). *Plant Species Biology*
860   **14**, 201-215 (1999).

861   44   Cruickshank, T. E. & Hahn, M. W. Reanalysis suggests that genomic islands of speciation
862   are due to reduced diversity, not reduced gene flow. *Molecular ecology* **23**, 3133-3157
863   (2014).

864   45   Ronfort, J., Jenczewski, E., Bataillon, T. & Rousset, F. Analysis of population structure in
865   autotetraploid species. *Genetics* **150**, 921-930 (1998).

866   46   Hudson, R. R., Slatkin, M. & Maddison, W. P. Estimation of levels of gene flow from DNA
867   sequence data. *Genetics* **132**, 583-589 (1992).

868   47   Nei, M. Genetic distance between populations. *The American Naturalist* **106**, 283-292
869   (1972).

870   48   Weir, B. S. & Cockerham, C. C. Estimating F-statistics for the analysis of population
871   structure. *Evolution*, 1358-1370 (1984).

872   49   Bhatia, G., Patterson, N., Sankararaman, S. & Price, A. L. Estimating and interpreting
873   FST: the impact of rare variants. *Genome research* **23**, 1514-1521 (2013).

874   50   Szpak, M. *et al.* FineMAV: prioritizing candidate genetic variants driving local adaptations
875   in human populations. *Genome biology* **19**, 1-18 (2018).

876   51   Craske, B. & Welburn, J. P. Leaving no-one behind: how CENP-E facilitates chromosome
877   alignment. *Essays in biochemistry* **64**, 313-324 (2020).

878   52   Yoshizawa, K. *et al.* Tetraploidy-linked sensitization to CENP-E inhibition in human cells.
879   *Molecular oncology* (2023).

880   53   Pisa, R., Phua, D. Y. Z. & Kapoor, T. M. Distinct Mechanisms of Resistance to a CENP-E
881   Inhibitor Emerge in Near-Haploid and Diploid Cancer Cells. *Cell Chemical Biology* **27**, 850-
882   857 e856 (2020). https://doi.org:10.1016/j.chembiol.2020.05.003

883   54   Real, A. M., Marsiglia, W. M. & Dar, A. C. Ploidy Leads a Molecular Motor to Walk Different
884   Paths to Drug Resistance. *Cell Chemical Biology* **27**, 770-772 (2020).
885   https://doi.org:10.1016/j.chembiol.2020.06.019

886   55   Isono, K., Yamamoto, H., Satoh, K. & Kobayashi, H. An Arabidopsis cDNA encoding a
887   DNA-binding protein that is highly similar to the DEAH family of RNA/DNA helicase genes.
888   *Nucleic acids research* **27**, 3728-3735 (1999).

889   56   Kim, J. S. *et al.* Arabidopsis TBP-ASSOCIATED FACTOR 12 ortholog NOBIRO6 controls
890   root elongation with unfolded protein response cofactor activity. *Proceedings of the*
891   *National Academy of Sciences* **119** (2022). https://doi.org:10.1073/pnas.2120219119

892   57   Rolly, N. K. & Yun, B. W. Regulation of Nitrate (NO(3)) Transporters and Glutamate
893   Synthase-Encoding Genes under Drought Stress in Arabidopsis: The Regulatory Role of
894   AtbZIP62 Transcription Factor. *Plants* **10** (2021). https://doi.org:10.3390/plants10102149

895   58   Fellmeth, J. E. & McKim, K. S. Meiotic CENP-C is a shepherd: bridging the space between
896   the centromere and the kinetochore in time and space. *Essays Biochem* **64**, 251-261
897   (2020). https://doi.org:10.1042/EBC20190080

898   59   Ogura, Y., Shibata, F., Sato, H. & Murata, M. Characterization of a CENP-C homolog in
899   Arabidopsis thaliana. *Genes & genetic systems* **79**, 139-144 (2004).

60   Hamasaki, H. *et al.* SD3, an Arabidopsis thaliana homolog of TIM21, affects intracellular
ATP levels and seedling development. *Molecular Plant* **5**, 461-471 (2012).
https://doi.org/10.1093/mp/ssr088
61   Municio, C. *et al.* The Arabidopsis condensin CAP-D subunits arrange interphase
chromatin. *New Phytologist* **230**, 972-987 (2021). https://doi.org/10.1111/nph.17221
62   Sakamoto, T. *et al.* Condensin II alleviates DNA damage and is essential for tolerance of
boron overload stress in Arabidopsis. *Plant Cell* **23**, 3533-3546 (2011).
https://doi.org/10.1105/tpc.111.086314
63   Wood, J. L., Liang, Y., Li, K. & Chen, J. Microcephalin/MCPH1 associates with the
condensin II complex to function in homologous recombination repair∗. *Journal of*
Biological Chemistry **283**, 29586-29592 (2008).
64   Busoms, S. *et al.* Fluctuating selection on migrant adaptive sodium transporter alleles in
coastal Arabidopsis thaliana. *Proceedings of the National Academy of Sciences* **115**,
E12443-E12452 (2018). https://doi.org:10.1073/pnas.1816964115
65   Barrada, A. *et al.* A TOR-YAK1 signaling axis controls cell cycle, meristem activity and
plant growth in Arabidopsis. *Development* **146**, dev171298 (2019).
66   Kim, D., Ntui, V. O. & Xiong, L. Arabidopsis YAK 1 regulates abscisic acid response and
drought resistance. *FEBS letters* **590**, 2201-2209 (2016).
67   Komaki, S. *et al.* Functional Analysis of the Plant Chromosomal Passenger Complex.
Plant Physiol **183**, 1586-1599 (2020). https://doi.org/10.1104/pp.20.00344
68   Kirioukhova, O. *et al.* Female gametophytic cell specification and seed development
require the function of the putative Arabidopsis INCENP ortholog WYRD. *Development*
138, 3409-3420 (2011). https://doi.org/10.1242/dev.060384
69   Bomblies, K., Jones, G., Franklin, C., Zickler, D. & Kleckner, N. The challenge of evolving
stable polyploidy: could an increase in "crossover interference distance" play a central
role? *Chromosoma* **125**, 287-300 (2016).
70   Diaz, K. J. *A novel kinesin, Arabidopsis thaliana Centromere associated protein E2, is*
necessary for the progression of mitosis and meiosis. (University of Massachusetts
Boston, 2014).
71   Iemura, K. & Tanaka, K. Chromokinesin Kid and kinetochore kinesin CENP-E differentially
support chromosome congression without end-on attachment to microtubules. *Nature*
Communications **6**, 6447 (2015). https://doi.org/10.1038/ncomms7447
72   She, Z. Y. *et al.* Kinesin-7 CENP-E regulates chromosome alignment and genome stability
of spermatogenic cells. *Cell Death Discov* **6**, 25 (2020). https://doi.org/10.1038/s41420-
020-0261-8
73   Wood, K. W. *et al.* Antitumor activity of an allosteric inhibitor of centromere-associated
protein-E. *Proceedings of the National Academy of Sciences* **107**, 5839-5844 (2010).
https://doi.org/10.1073/pnas.0915068107
74   Matsuo, K. & Tamaoki, N. Rational design and development of a lit-active photoswitchable
inhibitor targeting CENP-E. *Org Biomol Chem* **19**, 6979-6984 (2021).
https://doi.org/10.1039/d1ob01332g
75   Marx, A., Hoenger, A. & Mandelkow, E. Structures of kinesin motor proteins. *Cell motility*
and the cytoskeleton **66**, 958-966 (2009).
76   Mirzaa, G. M. *et al.* Mutations in CENPE define a novel kinetochore-centromeric
mechanism for microcephalic primordial dwarfism. *Human genetics* **133**, 1023-1039
(2014).
77   Yen, T. J., Li, G., Schaar, B. T., Szilak, I. & Cleveland, D. W. CENP-E is a putative
kinetochore motor that accumulates just before mitosis. *Nature* **359**, 536-539 (1992).
78   Huang, Y. *et al.* BubR1 phosphorylates CENP-E as a switch enabling the transition from
lateral association to end-on capture of spindle microtubules. *Cell Research* **29**, 562-578
(2019).

Kim, Y., Holland, A. J., Lan, W. & Cleveland, D. W. Aurora kinases and protein phosphatase 1 mediate chromosome congression through regulation of CENP-E. *Cell* **142**, 444-455 (2010).

Nousiainen, M., Silljé, H. H., Sauer, G., Nigg, E. A. & Körner, R. Phosphoproteome analysis of the human mitotic spindle. *Proceedings of the National Academy of Sciences* **103**, 5391-5396 (2006).

Espeut, J. *et al.* Phosphorylation relieves autoinhibition of the kinetochore motor CENP-E. *Molecular cell* **29**, 637-643 (2008).

Killinger, K. *et al.* Auto-inhibition of Mif2/CENP-C ensures centromere-dependent kinetochore assembly in budding yeast. *The EMBO journal* **39**, e102938 (2020). https://doi.org:10.15252/embj.2019102938

Vagnarelli, P. & Earnshaw, W. C. Chromosomal passengers: the four-dimensional regulation of mitotic events. *Chromosoma* **113**, 211-222 (2004).

Smith, S. J., Osman, K. & Franklin, F. C. H. The condensin complexes play distinct roles to ensure normal chromosome morphogenesis during meiotic division in A rabidopsis. *The Plant Journal* **80**, 255-268 (2014).

Sakamoto, T., Sugiyama, T., Yamashita, T. & Matsunaga, S. Plant condensin II is required for the correct spatial relationship between centromeres and rDNA arrays. *Nucleus* **10**, 116-125 (2019).

Bundock, P. & Hooykaas, P. An Arabidopsis hAT-like transposase is essential for plant development. *Nature* **436**, 282-284 (2005).

Chang, W.-C. *et al.* Regulation of Ku gene promoters in Arabidopsis by hormones and stress. *Functional Plant Biology* **35**, 265-280 (2008).

Tamura, K., Adachi, Y., Chiba, K., Oguchi, K. & Takahashi, H. Identification of Ku70 and Ku80 homologues in Arabidopsis thaliana: evidence for a role in the repair of DNA double-strand breaks. *The Plant Journal* **29**, 771-781 (2002).

Li, S., Ding, B., Chen, R., Ruggiero, C. & Chen, X. Evidence that the transcription elongation function of Rpb9 is involved in transcription-coupled DNA repair in Saccharomyces cerevisiae. *Molecular and cellular biology* **26**, 9430-9441 (2006).

Hull, M. W., McKune, K. & Woychik, N. A. RNA polymerase II subunit RPB9 is required for accurate start site selection. *Genes & Development* **9**, 481-490 (1995).

Tan, E. H., Blevins, T., Ream, T. S. & Pikaard, C. S. Functional consequences of subunit diversity in RNA polymerases II and V. *Cell reports* **1**, 208-214 (2012).

Hemming, S. A. *et al.* RNA polymerase II subunit Rpb9 regulates transcription elongation in vivo. *Journal of Biological Chemistry* **275**, 35506-35511 (2000).

Nesser, N. K., Peterson, D. O. & Hawley, D. K. RNA polymerase II subunit Rpb9 is important for transcriptional fidelity in vivo. *Proceedings of the National Academy of Sciences* **103**, 3268-3273 (2006).

Allen, B. L. & Taatjes, D. J. The Mediator complex: a central integrator of transcription. *Nature reviews Molecular cell biology* **16**, 155-166 (2015).

Baxter, I. *et al.* A coastal cline in sodium accumulation in Arabidopsis thaliana is driven by natural variation of the sodium transporter AtHKT1; 1. *PLoS Genetics* **6**, e1001193 (2010).

Shi, H., Ishitani, M., Kim, C. & Zhu, J.-K. The Arabidopsis thaliana salt tolerance gene SOS1 encodes a putative Na+/H+ antiporter. *Proceedings of the National Academy of Sciences* **97**, 6896-6901 (2000).

Zhou, M. & Wang, W. SOS1 safeguards plant circadian rhythm against daily salt fluctuations. *Proceedings of the National Academy of Sciences* **119**, e2212950119 (2022). https://doi.org:10.1073/pnas.2212950119

Busoms, S., Fischer, S. & Yant, L. Chasing the mechanisms of ecologically adaptive salinity tolerance. *Plant Communications* (2023). https://doi.org:https://doi.org/10.1016/j.xplc.2023.100571

Furini, S. & Domene, C. K+ and Na+ conduction in selective and nonselective ion channels via molecular dynamics simulations. *Biophysical journal* **105**, 1737-1745 (2013).

Cotsaftis, O., Plett, D., Shirley, N., Tester, M. & Hrmova, M. A two-staged model of Na+ exclusion in rice explained by 3D modeling of HKT transporters and alternative splicing. *PloS one* **7**, e39865 (2012).

Almeida, P., Katschnig, D. & De Boer, A. H. HKT transporters—state of the art. *International Journal of Molecular Sciences* **14**, 20359-20385 (2013).

Wang, Y. *et al.* Architecture and autoinhibitory mechanism of the plasma membrane Na+/H+ antiporter SOS1 in Arabidopsis. *Nature Communications* **14**, 4487 (2023).

Pandey, A. K. *et al.* An Inherent Difference between Serine and Threonine Phosphorylation: Phosphothreonine Strongly Prefers a Highly Ordered, Compact, Cyclic Conformation. *ACS Chemical Biology* (2023).

Kant, P., Kant, S., Gordon, M., Shaked, R. & Barak, S. STRESS RESPONSE SUPPRESSOR1 and STRESS RESPONSE SUPPRESSOR2, two DEAD-box RNA helicases that attenuate Arabidopsis responses to multiple abiotic stresses. *Plant physiology* **145**, 814-830 (2007).

Khan, A. *et al.* The A rabidopsis STRESS RESPONSE SUPPRESSOR DEAD-box RNA helicases are nucleolar-and chromocenter-localized proteins that undergo stress-mediated relocalization and are involved in epigenetic gene silencing. *The Plant Journal* **79**, 28-43 (2014).

Kim, J. S., Kim, K. A., Oh, T. R., Park, C. M. & Kang, H. Functional characterization of DEAD-box RNA helicases in Arabidopsis thaliana under abiotic stress conditions. *Plant and cell physiology* **49**, 1563-1571 (2008).

Merlot, S. *et al.* Constitutive activation of a plasma membrane H+-ATPase prevents abscisic acid-mediated stomatal closure. *The EMBO journal* **26**, 3216-3226 (2007).

Nadeau, J. A. & Sack, F. D. Control of stomatal distribution on the Arabidopsis leaf surface. *Science* **296**, 1697-1700 (2002).

Sugano, S. S. *et al.* Stomagen positively regulates stomatal density in Arabidopsis. *Nature* **463**, 241-244 (2010).

Emms, D. M. & Kelly, S. OrthoFinder2: fast and accurate phylogenomic orthology analysis from gene sequences. *BioRxiv* **466201** (2018).

Sterken, R. *et al.* Combined linkage and association mapping reveals CYCD5; 1 as a quantitative trait gene for endoreduplication in Arabidopsis. *Proceedings of the National Academy of Sciences* **109**, 4678-4683 (2012).

Unhavaithaya, Y. & Orr-Weaver, T. L. Centromere proteins CENP-C and CAL1 functionally interact in meiosis for centromere clustering, pairing, and chromosome segregation. *Proceedings of the National Academy of Sciences* **110**, 19878-19883 (2013).

Kurzbauer, M.-T. *et al.* ATM controls meiotic DNA double-strand break formation and recombination and affects synaptonemal complex organization in plants. *The Plant Cell* **33**, 1633-1656 (2021).

Bonnet, S., Knoll, A., Hartung, F. & Puchta, H. Different functions for the domains of the Arabidopsis thaliana RMI1 protein in DNA cross-link repair, somatic and meiotic recombination. *Nucleic acids research* **41**, 9349-9360 (2013).

Szklarczyk, D. *et al.* STRING v11: protein–protein association networks with increased coverage, supporting functional discovery in genome-wide experimental datasets. *Nucleic acids research* **47**, D607-D613 (2019).

Eisenschmid, K., Jabbusch, S. & Koch, M. A. Evolutionary footprints of cold adaptation in arctic-alpine Cochlearia (Brassicaceae)–Evidence from freezing experiments and electrolyte leakage. *Perspectives in Plant Ecology, Evolution and Systematics* **59**, 125728 (2023).

117 Desjardins, S., Kanyuka, K. & Higgins, J. D. A cytological analysis of wheat meiosis
targeted by virus-induced gene silencing (VIGS). *Plant Meiosis: Methods and Protocols*,
319-330 (2020).
118 Higgins, J. D., Sanchez-Moran, E., Armstrong, S. J., Jones, G. H. & Franklin, F. C. H. The
Arabidopsis synaptonemal complex protein ZYP1 is required for chromosome synapsis
and normal fidelity of crossing over. *Genes & Development* **19**, 2488-2500 (2005).
119 Desjardins, S. D. *et al.* MutS homologue 4 and MutS homologue 5 maintain the obligate
crossover in wheat despite stepwise gene loss following polyploidization. *Plant Physiology*
183, 1545-1558 (2020).
120 Mandáková, T. & Lysak, M. A. Chromosome preparation for cytogenetic analyses in
Arabidopsis. *Current protocols in plant biology* **1**, 43-51 (2016).
121 Ijdo, J., Wells, R., Baldini, A. & Reeders, S. Improved telomere detection using a telomere
repeat probe (TTAGGG) n generated by PCR. *Nucleic acids research* **19**, 4780 (1991).
122 Mandáková, T. & Lysak, M. A. Painting of Arabidopsis chromosomes with chromosome-
specific BAC clones. *Current protocols in plant biology* **1**, 359-371 (2016).
123 Deorowicz, S., Debudaj-Grabysz, A. & Grabowski, S. Disk-based k-mer counting on a PC.
BMC bioinformatics **14**, 1-12 (2013).
124 Ranallo-Benavidez, T. R., Jaron, K. S. & Schatz, M. C. GenomeScope 2.0 and
Smudgeplot for reference-free profiling of polyploid genomes. *Nature Communications* **11**,
1432 (2020). https://doi.org:10.1038/s41467-020-14998-3
125 Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114-2120 (2014).
126 Li, H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
arXiv (2013).
127 Dudchenko, O. *et al.* De novo assembly of the Aedes aegypti genome using Hi-C yields
chromosome-length scaffolds. *Science* **356**, 92-95 (2017).
128 Durand, N. C. *et al.* Juicebox provides a visualization system for Hi-C contact maps with
unlimited zoom. *Cell systems* **3**, 99-101 (2016).
129 Novák, P., Neumann, P., Pech, J., Steinhaisl, J. & Macas, J. RepeatExplorer: a Galaxy-
based web server for genome-wide characterization of eukaryotic repetitive elements from
next-generation sequence reads. *Bioinformatics* **29**, 792-793 (2013).
130 Ou, S. *et al.* Benchmarking transposable element annotation methods for creation of a
streamlined. *BioRxiv* **657890** (2019).
131 Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21
(2013).
132 Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows–Wheeler
transform. *Bioinformatics* **25**, 1754-1760 (2009).
133 Li, H. *et al.* The sequence alignment/map format and SAMtools. *Bioinformatics* **25**, 2078-
2079 (2009).
134 McKenna, A. *et al.* The Genome Analysis Toolkit: a MapReduce framework for analyzing
next-generation DNA sequencing data. *Genome research* **20**, 1297-1303 (2010).
135 Patterson, N., Price, A. L. & Reich, D. Population structure and eigenanalysis. *PLoS*
genetics **2**, e190 (2006).
136 Monnahan, P. *et al.* Pervasive population genomic consequences of genome duplication
in Arabidopsis arenosa. *Nature Ecology and Evolution* **3**, 457-468 (2019).
https://doi.org:10.1038/s41559-019-0807-4
137 Pembleton, L. W., Cogan, N. O. & Forster, J. W. St AMPP: An R package for calculation
of genetic differentiation and structure of mixed-ploidy level populations. *Molecular*
Ecology Resources **13**, 946-952 (2013).
138 Gerard, D. Pairwise linkage disequilibrium estimation for polyploids. *Molecular Ecology*
Resources **21**, 1230-1242 (2021).

1103 139 Alexa, A. & Rahnenführer, J. Gene set enrichment analysis with topGO. *Bioconductor*
1104    *Improv* **27**, 1-26 (2009).
1105 140 Sievers, F. & Higgins, D. G. Clustal omega. *Current protocols in bioinformatics* **48**, 3.13.
1106    11-13.13. 16 (2014).
1107 141 Waterhouse, A. M., Procter, J. B., Martin, D. M., Clamp, M. & Barton, G. J. Jalview Version
1108    2—a multiple sequence alignment editor and analysis workbench. *Bioinformatics* **25**,
1109    1189-1191 (2009).
1110 142 Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**,
1111    583-589 (2021).
1112