The values of ecosystem services inside and outside of protected areas in Eastern and Southern Africa

Falko T. Buschke[1*], Claudia Capitani[1†],

1. European Commission, Joint Research Centre, Ispra, Italy.

* Corresponding author: falko.buschke@ec.europa.eu, ORCID: https://orcid.org/0000-0003-1167-7810

† claudia.capitani@ec.europa.eu, ORCID: https://orcid.org/0000-0002-1899-8679

Acknowledgements

We thank Mark Gerrard and Nuredin Juhar from the *Conservation Strategy Fund* for the collaboration on the 2nd edition of the *State of Protected and Conserved Areas in Eastern and Southern Africa* report.

Data availability

The data used in this study are available from the Ecosystem Services Valuation Database (https://www.esvd.net/) and the R-scripts to replicate our analysis are available on GitHub (https://github.com/falko-buschke/SOPACA/) and will be permanently deposited on Zenodo after peer-review.

Equity, Diversity, and Inclusion statement

The authors acknowledge that this study focuses on a region underrepresented in the scientific literature. While this specific manuscript does not include contributions by authors from the region, it only forms one portion of a chapter for the 2nd edition of the *State of Protected and Conserved Areas in Eastern and Southern Africa* report co-authored with researchers from South Africa and Ethiopia (named in the acknowledgments section of this manuscript). Other chapters of the report are authored primarily by researchers from within the region, so this manuscript forms part of a larger representative research process.

The values of ecosystem services inside and outside of protected areas in Eastern and Southern Africa

Abstract

Conservation policies often take for granted the importance of protected areas for supplying ecosystem services. The 1st edition of the *State of Protected and Conserved Areas in Eastern and Southern Africa* report contained limited information on ecosystem services, so for the upcoming 2nd edition we compared statistically 561 standardised economic values of various types of ecosystem services inside and outside of protected areas. We found that data from local and sub-national case-studies in the Ecosystem Service Valuation Database were biased geographically, highlighting major evidence gaps for most of the region. For well-studied countries (Botswana, Ethiopia, Kenya, South Africa, Tanzania, Uganda), the value of ecosystem services varied considerably across different types of services but were – on average – five times higher outside protected areas. This trend was not universal, however, given that opportunities for recreation and tourism tended to be 20 times higher within protected areas. Combined, these findings suggest that conservation authorities across Eastern and Southern Africa (1) prioritise ecosystem service valuation studies; (2) expand the focus of ecosystem service policies to include wider landscapes beyond protected area boundaries; and (3) avoid making generic assumptions about ecosystem services by identifying which types of services are most compatible with the broader goals of protection.

Key words: Africa, ecosystem services, generalised linear mixed-effect model, nature's contributions to people, protected areas.

1. Introduction

Protected areas are valuable for maintaining biodiversity (Watson et al., 2014; Wauchope et al., 2022), and it is often taken for granted that they also enhance ecosystem services. Consequently, global and regional conservation policies regularly link ecosystem services with protected areas. At the global level, Target 3 of the Kunming-Montréal Global Biodiversity Framework aims to conserve areas of particular importance to ecosystem services through protected areas and other effective area-based conservation measures (Convention on Biological Diversity, 2022). In Africa, the *Kigali Call to Action for People and Nature*, which emerged from the inaugural IUCN Africa Protected Areas Congress in 2022, appealed for unprotected areas of importance to ecosystem services to be built into conservation plans (IUCN, 2022).

Despite the implicit acceptance that protected and conserved areas (hereafter protected areas) are important for the continued supply of ecosystem services, empirical evidence from Africa is far from unequivocal. For instance, although the IPBES regional assessment report for Africa (IPBES, 2018) described protected areas

as "a driver of positive change" to ecosystem services (page 265), it did not present
empirical evidence comparing the ecosystem services inside and outside of protected
areas. Similarly, neither of the *State of Protected and Conserved Areas* reports for
Central Africa (Doumenge et al., 2021) or Eastern and Southern Africa (IUCN Eastern
and Southern Africa Regional Office, 2020) quantified the relative contributions of
protected areas to ecosystem services.

Regional protected area policies in Africa would benefit from a transparent scientific
foundation when it comes to ecosystem services. Existing empirical studies in Eastern
and Southern Africa tend to focus on single protected areas and their immediate
surroundings (e.g., Guerbois and Fritz, 2017; Ndayizeye et al., 2020), or on a
protected area network within a single country (e.g., Ament et al., 2017; Roux et al.,
2020). Other studies quantify the coverage of ecosystem services by a regional
protected area network (e.g., Neugarten et al., 2020; Wei et al., 2020), assuming
implicitly that ecosystem services are enhanced by protection. Despite this,
comparative studies of different types of ecosystems services inside and outside of
protected areas remain scarce.

In this study, we evaluate ecosystem services inside and outside of protected areas in
Eastern and Southern Africa. Our analysis is intended to feed into the 2nd edition of
the *State of protected and conserved areas in Eastern and Southern Africa*, to guide
how ecosystem services within protected areas and surrounding areas ought to be
acknowledged, managed, and communicated. To cover such a large geographic area,
we rely on existing ecosystem valuation studies from the Ecosystem Service Valuation
Database (ESVD) (Brander et al., 2023), one of the most comprehensive and
standardised collections of ecosystem service valuations studies (Schmidt and
Seppelt, 2018). Although the ESVD database is limited to studies that estimated the
monetary value of ecosystem services, it has the unique feature that its spatially
explicit terrestrial case-studies include information on protection status (Schmidt and
Seppelt, 2018). This allowed us to compare statistically the value of different kinds of
ecosystem services inside and outside of protected areas in Eastern and Southern
Africa.

Our analysis was primarily exploratory, without specific *a priori* hypotheses. However,
we started from the position that value probably varied across different types of
ecosystem services, without assuming which types were most valuable. In some
instances, protection likely shields ecosystems from negative human impacts,
enhancing the supply and value of ecosystem services. In other instances, protection
may restrict people's access to ecosystems services, thereby reducing their value.
Therefore, we did not have a specific hypothesis about whether the values of
ecosystem services were higher or lower in protected areas overall because this would
depend on an interaction between protection status and the relative value of different
types of ecosystem services. So rather than predicting the specific monetary value of
ecosystem services within protected areas, our analysis is intended to provide broad
guidance for protected area policies across the region.

2. Methods

2.1. Study area

This study focused on 26 countries (Figure 1) to be covered by the upcoming 2nd edition of the State of protected and conserved areas in Eastern and Southern Africa (IUCN Eastern and Southern Africa Regional Office, 2020). The 1st edition of the report covered 24 countries in Eastern and Southern Africa, but this edition will be expanded to also consider Burundi and the Democratic Republic of Congo (DRC) because both states are members of the East African Community (EAC) and the DRC is member of the Southern African Development Community (SADC).

2.2. Data sources and data inclusion rules

Data were from the version of the Ecosystem Service Valuation Database (ESVD) available on 31 May 2023 (Brander et al., 2023). This global database is comprised of 9,453 estimated values of ecosystem services from more than 1,100 case studies. We filtered the global dataset to the 26 focal countries of this study, resulting in 1,166 independent value estimates. We only included estimates from local (n = 866) and sub-national (n = 213) assessments. Each estimate included a standardised value of ecosystems services in International $.ha^{-1}.year^{-1} for the year 2020, and after excluding cases where standardised economic value was either unreported or reported as zero, 769 estimates remained (these are shown in Figure 1).

Each record in ESVD includes metadata on protection status. This information includes three levels of protection: "*No protection*" (n = 187); "*Partially Protected*" (n = 352); and "*Protected*" (n = 106), which we aggregated into two levels by combining "*Partially Protected*", and "*Protected*". We also assumed that records without protection information (n = 124) were unprotected, so we aggregated these into a single "*Unprotected*" category.

The ESVD identifies the specific type of ecosystem service for each estimate of value (Figure 1), but in some instances, estimates represent the aggregated value of multiple ecosystem services. The database guidelines advise against disaggregating values from multiple services, so we excluded all estimates based on multiple ecosystem services (n = 117).

To balance the data included in the statistical analysis, we only included data from countries that had estimates from both protected and unprotected sites (n = 592). This meant that we excluded data from Madagascar (n = 7 protected sites), Malawi (n = 3 protected sites), Mozambique (n = 5 unprotected sites), Namibia, (n = 10 protected sites), Rwanda (n = 1 protected site), Sudan (n = 21 protected sites), Zambia (n = 5 protected sites), and Zimbabwe (n = 8 unprotected sites). Similarly, we excluded types of ecosystem services that did not include estimates from both protected and unprotected sites. This resulted in the exclusion of "*unreported*" ecosystem service types (n = 12 unprotected sites); "*aesthetic information*" (n = 1 unprotected site); "*inspiration for culture, art, and design*" (n = 1 protected site); "*maintenance of genetic diversity*" (n = 2 protected sites); "*maintenance of life cycles*" (n = 1 protected site); and "*regulation of water flows*" (n = 2 protected sites). After these exclusions, Somalia no longer included estimates from unprotected sites, so we excluded the remaining

129 estimates from protected sites from this country (n =12). The final dataset used for
130 statistical modelling included 561 estimates of 12 ecosystem services types from both
131 protected and unprotected sites in six countries (Botswana, Ethiopia, Kenya, South
132 Africa, Tanzania, and Kenya).

Figure 1: The geographic location of local and sub-national case studies valuing ecosystem services
across Eastern and Southern Africa from the Ecosystem Services Valuation Database (Brander et al.,
2023). Points are coloured according to the type of ecosystem service and scaled according to the
standardised economic value (International $.ha^{-1}.year^{-1} for the year 2020). Shaded areas denote
protected areas from the World Database for Protected Areas (UNEP-WCMC and IUCN, 2022). The
two Indian Ocean island states of Mauritius and Seychelles are not displayed on this map because
there were no local or sub-national ecosystem service valuation case studies for either country.

2.3.   Model structure

Our aim was to examine whether protected areas are associated with higher values
of ecosystem services, while recognising that protection may be more effective at
conserving certain types of ecosystem services than others. As such, we included
protection level and ecosystem type, and their interactions, as categorical fixed effects
in our model. Moreover, we anticipated that the relationship between protected areas
and ecosystem services was possibly affected by several unmeasured spatially

autocorrelated variables – such as climate, biogeography, governance, or
management – which would have been too complex to model explicitly. Therefore, we
included countries as random effects in the mixed effects model.

The distribution of values of ecosystems services (International \$.ha^{-1}.year^{-1} for the
year 2020) were strongly right-skewed and dominated by many low values and few
high values. Therefore, we initially considered two modelling approaches: (1) a log-
linked Gamma generalised linear mixed effect model (*Gamma-glme*) using the `glmer`
function in the `lme4` package (Bates et al., 2015) in R version 4.1.3 (R Core team,
2022) and (2) a linear mixed effects model with a log-transformed response variable
(*log-lme:* `lmer` function in `lme4` package). Model diagnostics suggested that neither
modelling approach was clearly superior (Appendix 1, Figure S1). So we favoured the
Gamma-glme because the model residuals were distributed more normally than the
log-lme (Figure S1 a,b) and the model predictions are more intuitive (i.e., *Gamma-*
glme models the log of the expected mean value of ecosystem services, whereas the
log-lme models the expectation of the log-transformed mean value of ecosystem
services: Appendix 1). Our final model structure was:

$$Value_{ij} \sim Gamma\left(\mu_{ij}\right)$$

$$E\left(Value_{ij}\right) = \mu_{ij}$$

$$\log\left(\mu_{ij}\right) = Protected_{ij} + ES_{ij} + Protected_{ij} \times ES_{ij} + Country_i$$

$$Country_i \sim \mathcal{N}\left(0, \sigma^2\right)$$

Where $Value_{ij}$ is the estimated value (International \$.ha^{-1}.year^{-1} for the year 2020) of
ecosystem observation j in country i with an expected (E) mean value of μ_{ij}.
$Protected_{ij}$ (two levels of protection, with "*Unprotected"* as the baseline) and ES_{ij} (12
levels of ecosystem service type, with "*Food"* as the baseline) were included as
categorical predictor variables, along with their two-way interaction. $Country_i$ is the
random intercept in the model, which is assumed to be normally distributed with a
mean of zero and a variance of σ^2.

Alongside this full model, we also modelled three subset models by (i) only including
protection as a predictor, (ii) only including ecosystem service type as a predictor, (iii)
modelling protection level and ecosystem service type additively without an interaction
term. These subset models were compared using a likelihood-ratio χ^2 test (`anova`
command in `stats` package). Given the relatively large dataset – 561 data points –
we used a conservative threshold ($\alpha = 0.01$) to assess whether regression
coefficients differed significantly from zero.

3. Results

Estimates of ecosystem service values in the ESVD were clearly biased
geographically (Figure 1). Eight of the 26 countries (30%) in Eastern and Southern
Africa had no case study data, and a further seven countries had fewer than 10 value

estimates. Therefore, our statistical model should be interpreted cautiously; especially
in parts of the region that are poorly represented in the ESVD.

The log-linked Gamma generalised linear mixed effects model that included both
protection level, ecosystem service type and their interaction provided the best fit to
the empirical data from the ESVD (Table 1). This model formulation had the lowest
AIC (7734.9) and highest log-likelihood (-3841.5) compared to simpler model
formulations. This confirms that the variation in the standardised economic value of
ecosystem services is related statistically to protection status, the type of ecosystem
service and their interaction.

Table 1: An incremental comparison of candidate model formulations of increasing complexity using a
likelihood-ratio χ^2 test. Models are ordered based on decreasing Akaike's Information Criterion (AIC),
which corresponds with the addition of fixed effects – and their interaction – to the model.

Fixed effects	AIC	log-likelihood	χ^2 (degrees of freedom)*	P-value*
Protection	7822.2	-3907.1	-	
Ecosystem service type	7790.5	-3881.2	51.749 (10)	< 0.001
Protection + Ecosystem service type	7757.2	-3863.6	35.296 (1)	< 0.001
Protection + Ecosystem service type + (Protection x Ecosystem service type)	7734.9	-3841.5	44.255 (11)	< 0.001

* Values are based on a pair-wise comparisons with the preceding model

The variance explained by country identity as a random effect to the full model
accounted for 85.86 % of the total random variance (Variance$_{country}$ = 29.83 ± 5.5 sd;
Variance$_{residual}$ = 4.91 ± 2.2 sd). The average value of ecosystem services increased
sequentially across Botswana, Tanzania, Kenya, Ethiopia, Uganda, and South Africa
(Figure 2). It remains uncertain whether this is due to differences between the
countries *per se* (e.g., governance or management differences) or due to underlying
spatially autocorrelated environmental variables (e.g. climate, ecosystem type, or
biogeography). Nevertheless, it cautions against assuming that the standardised
economic value from one part of the region is transferable to another.

Model coefficients suggest that the standardised value of ecosystem services outside
of protected areas was three to six times higher than the value inside protected areas
($e^{-1.506(\pm 0.31)} = 0.22\ (0.16 - 0.30)$: Table 2). Such a large statistical effect was,
understandably, statistically significant (t = -4.89, P < 0.001). Standardised value also
varied across the type of ecosystem service (Table 2). "*Food*" had the highest value
on average and was set as the baseline category in the regression model (Table 2,
Figure 2). The values of "*existence, bequest values", "information for cognitive*
development", "opportunities for recreation and tourism", "waste treatment", and
"*water*" were statistically similar to the value of "*food*" (each of these categories had
negative coefficients suggesting that they were less valuable on average, but these
coefficients did not differ significantly from zero at the conservative threshold of 0.001:
Table 2). By contrast, the values of "*climate regulation*" (4.1%), "*erosion prevention*"
(5.6%), "*maintenance of soil fertility*" (1.2%), "*medicinal resources*" (0.1%),

"moderation of extreme events" (5.1%), and *"raw materials"* (10.4%) were all
considerably less than *"food"* (Table 2, Figure 2).

Table 2: Model coefficients of a log-linked Gamma generalised linear mixed effects model of the
standardised value of ecosystem services relative to protected status, the type of ecosystem service
and their interaction. Given the log-link in the model, exponents of the estimated coefficients reflect the
statistical effect on the value of ecosystem services. Statistically significant coefficients at the
conservative threshold of $\alpha = 0.01$ are shown as P-values in bold font.

Fixed effect	Additive terms			Interaction terms		
	Coefficient (sd)	t-value	P-value	Coefficient (sd)	t-value	P-value
Protection (Unprotected = 0)						
Protected	-1.506 (0.31)	-4.89	**< 0.001**			
Ecosystem service type (Food = 0)				**Interaction with protection**		
Climate regulation	-3.191 (0.67)	-4.75	**< 0.001**	2.003 (1.28)	1.57	0.116
Erosion prevention	-2.887 (0.61)	-4.71	**< 0.001**	-4.821 (1.25)	-3.86	**<0.001**
Existence, bequest values	-1.784 (0.71)	-2.51	0.012	0.933 (1.01)	0.92	0.358
Information for cognitive development	-2.185 (1.38)	-1.58	0.113	-3.382 (2.34)	-1.45	0.147
Maintenance of soil fertility	-4.458 (1.36)	-3.27	**0.001**	3.515 (1.75)	2.0	0.045
Medicinal resources	-6.324 (1.0)	-6.31	**< 0.001**	1.616 (1.47)	1.10	0.273
Moderation of extreme events	-2.969 (0.79)	-3.74	**< 0.001**	1.589 (2.04)	0.78	0.436
Opportunities for recreation and tourism	-1.182 (0.55)	-2.15	0.032	1.877 (0.61)	3.08	**0.002**
Raw materials	-2.260 (0.53)	-4.28	**<0.001**	-0.424 (0.56)	-0.76	0.448
Waste treatment	-0.714 (1.96)	-0.36	0.716	-4.042 (2.69)	-1.50	0.133
Water	-0.066 (1.96)	-0.09	0.932	-2.283 (0.93)	-2.46	0.014

The model's interaction coefficients suggested that protection status affected the value
of ecosystem services differently (Table 2). Specifically, the value of *"erosion*
prevention" in protected areas was approximately 15% of the value of the same
service outside of protected areas. In comparison, the average value of *"opportunities*
for recreation and tourism" inside protected areas was roughly 20 times higher than
unprotected areas (Table 2; Figure 2).

Figure 2: The standardised value (International $.ha⁻¹.year⁻¹) of different types of ecosystem services in protected and unprotected areas. Grey points show empirical values from 561 case studies. Coloured squares show predictions from a log-linked Gamma generalised linear mixed effects models with six countries included as random factors. The horizontal axis is log-scaled.

4. Discussion

This study set out to analyse the value of different types of ecosystem services inside and outside of protected areas in Eastern and South Africa. Our analysis was designed to inform the upcoming 2nd edition of the *State of protected and conserved areas in Eastern and Southern Africa.* However, the message most relevant to policy probably stems from what our analysis failed to show. The uneven geographic spread of records in the ESVD database highlights how little is known about the value of ecosystem services across most of the region and suggests that collecting representative data should be an urgent policy priority across the region.

A recent study by Duncanson et al., (2023) quantified the added value of protected areas for preserving carbon stocks globally. Five Eastern and Southern African countries (DRC, Mozambique, Tanzania, Zambia, and Madagascar) were amongst the top 20 countries globally in terms of the additionalilty of carbon storage by protected areas. Yet despite these countries being important for global climate mitigation, only Tanzania had sufficient ecosystem valuation case studies to inform our comparative analysis. Data gaps like these risk overlooking much of nature's

value, leading to its inadvertent devaluation in decision-making contexts (Costanza et
al., 2014). Such disregard is unfortunately already a common policy outcome even
when valuation data do exist (IPBES, 2022; Laurans et al., 2013). Although we did not
expressly set out to compare the value of ecosystem services across countries in
Eastern and Southern Africa, the random effect in our statistical model showed how
country identity accounted for 85% of the variation in ecosystem service value that
could not be attributed to the protection status or type of ecosystem service. This
suggests that filling data gaps in one country by extrapolating the value of ecosystem
services to another is not advisable.

The data we could access suggested that the value of ecosystem services were on
average five times higher outside of protected area than within them. This relationship
is unlikely causal, and does not imply that protected area expansion will reduce the
value of ecosystem services. The monetary value of ecosystem services is not just a
function of the aggregate supply of these services by nature, but also the aggregate
demand by people (Dasgupta, 2021). Aggregate demand is typically determined by
human population size, their *per capita* consumption, and the efficiency with which
nature's contributions are consumed (Dasgupta, 2021). All three of these variables
tend to be higher outside of protected areas. In our analysis, this is most clearly
illustrated by "*food*" ecosystem services, which had the highest value of the service
types considered in our model. Even though cropland is expanding into protected
areas globally, in Southern and Eastern Africa expansion rates are slower than
unprotected background rates (Meng et al., 2023). It stands to reason that the value
of food services will be higher outside of protected areas. That said, expanding
protected areas would not necessarily reduce cultivated area and the value of food
services. Instead, protected area expansion may redirect cultivation elsewhere in the
landscape (Ford et al., 2020; Waldron et al., 2020). So, our results should not be
misconstrued that protection reduces the value of ecosystem services.

Our results imply that policies specifically targeting ecosystem services should
consider expanding their focus beyond the boundaries of protected areas. This is
consistent with global calls for integrated management of "working" or "shared"
landscapes (Kremen and Merenlender, 2018; Obura et al., 2021) and regional
programmes, like the European Commission's *NaturAfrica* initiative (European
Commission, 2021). However, across Eastern and Southern Africa these kinds of
landscape-wide efforts often struggle for legitimacy and face challenges around
governance, power dynamics and environmental justice (Bourgeois et al., 2023;
Cassidy, 2021; Kicheleri et al., 2021). Although community-based conservation in
unprotected landscapes relies on many of the same tools as protected areas – notably,
clearly defined boundaries, the need for monitoring, restricted resource use, and
sanctions for rule-breaking (Mahajan et al., 2021) – it faces a unique tension between
the concentration of decision-making power in formal institutions and the local
communities who resist these trends (Kicheleri et al., 2021; Nelson et al., 2021).
Conservation authorities should not presuppose that existing protected area policies
will transfer smoothly in unprotected landscapes and should, therefore, plan
accordingly.

The significance of the unprotected landscapes for the supply of ecosystem services does not absolve protected area authorities from dealing with nature's contributions. On the contrary, our results support more nuanced inclusion of ecosystem service in protected area policies and strategies. This is demonstrated by the interaction term in our regression model, which showed, for example, how "*opportunities for recreation and tourism*" were 20 times higher in protected areas on average. Globally, the consumer surplus generated by protected area visitation dwarfs spending on protected areas (Balmford et al., 2015). The African Union's post-pandemic *Green Recovery Action Plan* recognises this opportunity*,* calling for the immediate support of ecotourism within the context of improving the management of protected areas (African Union, 2021). Protected area authorities could consider management interventions that enhance tourism opportunities because accessibility is a major determinant of tourist visitation to, as well as within, protected areas (Arbieu et al., 2018; Balmford et al., 2015). Such interventions should be prudent though, because unchecked tourism may exacerbate ecological degradation (Dasgupta, 2021; Hunter and Shaw, 2005).

In conclusion, protected areas remain a cornerstone of conservation policy. But our study, which is not without its caveats, advises against taking for granted the importance of protected areas for the supply of ecosystem services. In general, evidence for such claims is sparse, emphasising the urgent need for ecosystem service valuation studies that (i) address existing geographic biases in data; (ii) evaluate values beyond the narrow scope of monetary valuations studies (e.g., IPBES, 2022); (iii) include assessments of the marine realm, which was completely overlooked by our analyses. The limited data that do exist, however, suggest that ecosystem services provide considerable value beyond the boundaries of protected areas, probably because this is where nature's services are more accessible. Therefore, policies targeting ecosystem services need to look beyond protected areas to preserve nature's contributions in unprotected working landscapes. Within protected areas, authorities should avoid generic assumptions about ecosystem services. Tailored interventions require a more nuanced, evidence-based approach. If ecosystem services are indeed a policy priority for protected area authorities, then they should invest resources and attention into identifying the specific types of services most compatible with the broader goals of protection.

5. References

African Union, 2021. Green Recovery Action Plan 2021-2027. African Union Directorate of Sustainable Environment and Blue Economy, Addis Ababa, Ethiopia.

Ament, J.M., Moore, C.A., Herbst, M., Cumming, G.S., 2017. Cultural Ecosystem Services in Protected Areas: Understanding Bundles, Trade-Offs, and Synergies. Conserv Lett 10, 440–450. https://doi.org/10.1111/conl.12283

Arbieu, U., Grünewald, C., Martín-López, B., Schleuning, M., Böhning-Gaese, K., 2018. Large mammal diversity matters for wildlife tourism in Southern African Protected Areas: Insights for management. Ecosystem Services 31, 481–490. https://doi.org/10.1016/j.ecoser.2017.11.006

Balmford, A., Green, J.M.H., Anderson, M., Beresford, J., Huang, C., Naidoo, R., Walpole, M., Manica, A., 2015. Walk on the Wild Side: Estimating the Global Magnitude of Visits to Protected Areas. PLoS Biol 13, e1002074. https://doi.org/10.1371/journal.pbio.1002074

Bates, D., Mächler, M., Bolker, B., Walker, S., 2015. Fitting Linear Mixed-Effects Models Using **lme4**. J. Stat. Soft. 67. https://doi.org/10.18637/jss.v067.i01

Bourgeois, R., Guerbois, C., Giva, N., Mugabe, P., Mukamuri, B., Fynn, R., Daré, W., Motsholapheko, M., Nare, L., Delay, E., Ducrot, R., Bucuane, J., Mercandalli, S., Le Page, C., Caron, A., 2023. Using anticipation to unveil drivers of local livelihoods in Transfrontier Conservation Areas: A call for more environmental justice. People and Nature pan3.10446. https://doi.org/10.1002/pan3.10446

Brander, L.M., De Groot, R., Guisado Goñi, V., van 't Hoff, V., Schägner, P., Solomonides, S., McVittie, A., Eppink, F., Sposato, M., Do, L., Ghermandi, M., Sinclair, M., 2023. Ecosystem Services Valuation Database (ESVD). Foundation for Sustainable Development and Brander Environmental Economics.

Cassidy, L., 2021. Power dynamics and new directions in the recent evolution of CBNRM in Botswana. Conservat Sci and Prac 3. https://doi.org/10.1111/csp2.205

Convention on Biological Diversity, 2022. Kunming-Montreal Global biodiversity framework (No. CBD/COP/15/L.25). Conference of the Parties to the Convention on Biological Diversity, Montreal, Canada.

Costanza, R., de Groot, R., Sutton, P., van der Ploeg, S., Anderson, S.J., Kubiszewski, I., Farber, S., Turner, R.K., 2014. Changes in the global value of ecosystem services. Global Environmental Change 26, 152–158. https://doi.org/10.1016/j.gloenvcha.2014.04.002

Dasgupta, P., 2021. The economics of biodiversity: the Dasgupta review: full report, Updated: 18 February 2021. ed. HM Treasury, London.

Doumenge, C., Palla, F., Itsoua Madzous, G.-L., 2021. State of Protected Areas in Central Africa 2020. OFAC-COMIFAC, IUCN, Yaounde, Cameroon and Gland, Switzerland.

Duncanson, L., Liang, M., Leitold, V., Armston, J., Krishna Moorthy, S.M., Dubayah, R., Costedoat, S., Enquist, B.J., Fatoyinbo, L., Goetz, S.J., Gonzalez-Roglich, M., Merow, C., Roehrdanz, P.R., Tabor, K., Zvoleff, A., 2023. The effectiveness of global protected areas for climate change mitigation. Nat Commun 14, 2908. https://doi.org/10.1038/s41467-023-38073-9

European Commission, 2021. NaturAfrica: the Green Deal approach for EU support to biodiversity conservation in Africa. European Commission. Directorate General for International Partnerships, Brussels, Belgium.

Ford, S.A., Jepsen, M.R., Kingston, N., Lewis, E., Brooks, T.M., MacSharry, B., Mertz, O., 2020. Deforestation leakage undermines conservation value of tropical and subtropical forest protected areas. Global Ecol. Biogeogr. 29, 2014–2024. https://doi.org/10.1111/geb.13172

Guerbois, C., Fritz, H., 2017. Patterns and perceived sustainability of provisioning ecosystem services on the edge of a protected area in times of crisis. Ecosystem Services 28, 196–206. https://doi.org/10.1016/j.ecoser.2017.11.010

Hunter, C., Shaw, J., 2005. Applying the ecological footprint to ecotourism scenarios. Envir. Conserv. 32, 294–304. https://doi.org/10.1017/S0376892906002591

IPBES, 2022. Methodological assessment of the diverse values and valuation of nature of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. Zenodo. https://doi.org/10.5281/ZENODO.6522522

IPBES, 2018. The IPBES regional assessment report on biodiversity and ecosystem services for Africa. Secretariat of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services, Bonn, Germany. https://doi.org/10.5281/ZENODO.3236178

IUCN, 2022. Kigali Call to Action for People and Nature. IUCN World Commission on Protected Areas, Republic of Rwanda, African Wildlife Foundation, Kigali, Rwanda.

IUCN Eastern and Southern Africa Regional Office, 2020. The state of protected and conserved areas in Eastern and Southern Africa. IUCN, International Union for Conservation of Nature. https://doi.org/10.2305/IUCN.CH.2020.15.en

Kicheleri, R.P., Mangewa, L.J., Nielsen, M.R., Kajembe, G.C., Treue, T., 2021. Designed for accumulation by dispossession: An analysis of Tanzania's Wildlife Management Areas through the case of Burunge. Conservat Sci and Prac 3. https://doi.org/10.1111/csp2.360

Kremen, C., Merenlender, A.M., 2018. Landscapes that work for biodiversity and people. Science 362, eaau6020. https://doi.org/10.1126/science.aau6020

Laurans, Y., Rankovic, A., Billé, R., Pirard, R., Mermet, L., 2013. Use of ecosystem services economic valuation for decision making: Questioning a literature blindspot. Journal of Environmental Management 119, 208–219. https://doi.org/10.1016/j.jenvman.2013.01.008

Mahajan, S.L., Jagadish, A., Glew, L., Ahmadia, G., Becker, H., Fidler, R.Y., Jeha, L., Mills, M., Cox, C., DeMello, N., Harborne, A.R., Masuda, Y.J., McKinnon, M.C., Painter, M., Wilkie, D., Mascia, M.B., 2021. A theory-based framework for understanding the establishment, persistence, and diffusion of community-based conservation. Conservat Sci and Prac 3. https://doi.org/10.1111/csp2.299

Meng, Z., Dong, J., Ellis, E.C., Metternicht, G., Qin, Y., Song, X.-P., Löfqvist, S., Garrett, R.D., Jia, X., Xiao, X., 2023. Post-2020 biodiversity framework challenged by cropland expansion in protected areas. Nat Sustain. https://doi.org/10.1038/s41893-023-01093-w

Ndayizeye, G., Imani, G., Nkengurutse, J., Irampagarikiye, R., Ndihokubwayo, N., Niyongabo, F., Cuni-Sanchez, A., 2020. Ecosystem services from mountain forests: Local communities' views in Kibira National Park, Burundi. Ecosystem Services 45, 101171. https://doi.org/10.1016/j.ecoser.2020.101171

Nelson, F., Muyamwa-Mupeta, P., Muyengwa, S., Sulle, E., Kaelo, D., 2021. Progress or regression? Institutional evolutions of community-based conservation in eastern and southern Africa. Conservat Sci and Prac 3. https://doi.org/10.1111/csp2.302

Neugarten, R.A., Moull, K., Martinez, N.A., Andriamaro, L., Bernard, C., Bonham, C., Cano, C.A., Ceotto, P., Cutter, P., Farrell, T.A., Gibb, M., Goedschalk, J., Hole, D., Honzák, M., Kasecker, T., Koenig, K., Larsen, T.H., Ledezma, J.C., McKinnon, M., Mulligan, M., Nijbroek, R., Olsson, A., Rakotobe, Z.L., Rasolohery, A., Saenz, L., Steininger, M., Wright, T.M., Turner, W., 2020. Trends in protected area representation of biodiversity and ecosystem services in five tropical countries. Ecosystem Services 42, 101078. https://doi.org/10.1016/j.ecoser.2020.101078

Obura, D.O., Katerere, Y., Mayet, M., Kaelo, D., Msweli, S., Mather, K., Harris, J., Louis, M., Kramer, R., Teferi, T., Samoilys, M., Lewis, L., Bennie, A., Kumah, F., Isaacs, M., Nantongo, P., 2021. Integrate biodiversity targets from local to global levels. Science 373, 746–748. https://doi.org/10.1126/science.abh2234

R Core team, 2022. R: A Language and Environment for Statistical Computing.

Roux, D.J., Smith, M.K.S., Smit, I.P.J., Freitag, S., Slabbert, L., Mokhatla, M.M., Hayes, J., Mpapane, N.P., 2020. Cultural ecosystem services as complex outcomes of people–nature interactions in protected areas. Ecosystem Services 43, 101111. https://doi.org/10.1016/j.ecoser.2020.101111

Schmidt, S., Seppelt, R., 2018. Information content of global ecosystem service databases and their suitability for decision advice. Ecosystem Services 32, 22–40. https://doi.org/10.1016/j.ecoser.2018.05.007

UNEP-WCMC, IUCN, 2022. Protected Planet: The World Database on Protected Areas (WDPA). UNEP-WCMC and IUCN.

Waldron, A., Adams, V., Allan, J., Arnell, A., Asner, G., Atkinson, S., Baccini, A., Baillie, J., Balmford, A., Beau, J.A., others, 2020. Protecting 30% of the planet for nature: costs, benefits and economic implications.

Watson, J.E.M., Dudley, N., Segan, D.B., Hockings, M., 2014. The performance and potential of protected areas. Nature 515, 67–73. https://doi.org/10.1038/nature13947

Wauchope, H.S., Jones, J.P.G., Geldmann, J., Simmons, B.I., Amano, T., Blanco, D.E., Fuller, R.A., Johnston, A., Langendoen, T., Mundkur, T., Nagy, S., Sutherland, W.J., 2022. Protected areas have a mixed impact on waterbirds, but management helps. Nature 605, 103–107. https://doi.org/10.1038/s41586-022-04617-0

Wei, F., Wang, S., Fu, B., Liu, Y., 2020. Representation of biodiversity and ecosystem services in East Africa's protected area network. Ambio 49, 245–257. https://doi.org/10.1007/s13280-019-01155-4

Supplementary Information

Appendix 1: Model diagnostics

In the main manuscript, we present the results from a log-linked Gamma generalised linear mixed effect model (*Gamma-glme*), which we favoured over a linear mixed effects model with a log-transformed response variable (*log-lme*).

Model diagnostics suggested that neither approach was clearly superior. The residuals from the *Gamma-glme* model (Figure S1a) were distributed more normally than those from the *log-lme* model (Figure S1b), thus favouring the former model formulation. However, the residuals from the *Gamma-glme* model were biased downwards (mean residuals were negative: Figure S1c) compared to those from the *log-lme* model (Figure S1d). The modes from the random intercepts were approximately normal for both the *Gamma-glme* (Figure S1e) and the *log-lme* (Figure S1f), though in the former model intercepts for Botswana were lower than expected from normally distributed random effects.

A pertinent difference between the two modelling approaches, however, is how they model the response variable. The *Gamma-glme* model estimated the log of the expected mean value of ecosystem services, $log(E[Value])$, whereas the *log-lme* models the expected mean of the log-transformed value of ecosystem services, $E[log(Value)]$. These estimates are not the same. Given that the distribution of ecosystem service values is strongly right-skewed, modelling log-transformed response variables downplays ecosystems services with exceptionally high value. Therefore, we used the *Gamma-glme* model for the rest of our study.

Figure S1: Diagnostic plots comparing a log-linked generalised linear mixed effect model to linear mixed effects model with a log-transformed response variable. Q-Q normal plots for the model residuals (**a** and **b**), the residual plot relative to model predictions (**c** and **d**), Q-Q normal plots for the intercepts of the country identity as a random effect in the model (**e** and **f**).