From natural to sexual selection: Revealing a hidden preference for camouflage patterns

Yseult Hejja Brichard [1] *, Michel Raymond [2], Innes C. Cuthill [3], Tamra C. Mendelson [1] ¶, Julien P. Renoult [4] * ¶

[1] University of Maryland, Baltimore County, Baltimore, MD; [2] Institut des Sciences de l'Evolution (CNRS-UMR 5554), Montpellier, France; [3] School of Biological Sciences, University of Bristol, Bristol, United Kingdom; [4] CEFE, University of Montpellier, CNRS, EPHE, IRD, Montpellier, France

* Corresponding authors. E-mails: yseulthb@gmail.com ; julien.renoult@cefe.cnrs.fr
¶ These authors contributed equally to this work

Abstract

Natural and sexual selection can be in conflict in driving the evolution of sexual ornamentation. Sexual selection typically favors detectability to potential mates, whereas natural selection penalizes detectability to avoid predators. Considering signal efficiency in addition to detectability, however, suggests that natural and sexual selection need not be antagonistic. In the visual domain, considerable evidence supports a "processing bias" by which people prefer images that match the spatial statistics of natural scenes, likely because the brain has evolved to process such scenes efficiently. A direct but untested prediction of this bias is that background-matching camouflage can be favored by natural and sexual selection. We conducted an online experiment where we show for the first time human preference for camouflaged patterning and confirm a preference for the average image statistics of natural scenes. Because many of the underlying visual mechanisms are shared across vertebrates, our results suggest that camouflage patterns can serve as evolutionary precursors of sexual signals.

Introduction

Sexual ornamentation in animals is often pulled in opposite evolutionary directions. Sexual selection driven by mate preferences is generally thought to lead to detectability or conspicuousness, to draw the attention of potential mates (*1*), whereas natural selection should lead to the opposite, favoring camouflage over detectability to avoid attracting predators (*2*). Evidence for such a conflict has been shown in guppies, *Poecilia reticulata,* in which female preferences are biased towards detectable orange colouration (*3–5*) and males develop more intense orange colouration when predation is relaxed (*6*, *7*). While conflicts between natural and sexual selection are generally thought to play an important role in shaping sexual signal diversity (*8–10*), many of the best examples are based on ornament color (*11*). However, sexual signals also include patterns, which for visual images describe

the spatial organization of features, including colors and also shades of gray. Although visual patterns are receiving increased attention in research on sexual signals (*12*, *13*), the question of whether and to what extent they are shaped by conflicts between natural and sexual selection has so far remained little explored.

One problem is that conflicts between natural and sexual selection in the evolution of communication signals are primarily framed as a problem of signal *detection*, perhaps because of a historic focus on color. However, the attractiveness of a signal depends on more than its ability to be detected; it also depends on the ease with which the signal is processed by the brain (*14–17*). This "processing bias" (*18*) may stem from the well-supported adaptation of the visual system to efficiently process natural statistics of the environment (*19*, *20*). For visual signals, efficiency is achieved by matching signal design with the sensitivity (i.e., tuning) of visual neurons (*21*). As a result, stimuli that resemble the underlying spatial patterns of natural scenes will be processed more efficiently (i.e. more easily and at lower metabolic cost), generating a feeling of fluency that is experienced as pleasant and therefore rewarding (*22*). Indeed, a preference in humans for the statistical features of natural scenes is a robust finding in experimental psychology (*23*).

In nature, background-matching is a type of camouflage whereby the color and pattern of an animal's phenotype mimic those of its surroundings (see e.g. *25*). Given that background-matching relies on matching visual statistics of a stimulus with those of the habitat, the processing bias hypothesis predicts that, once detected, camouflage patterns should also be attractive to the observer. Small changes that increase the detectability of background-matching patterns to conspecifics could therefore increase the attractiveness of the signaler, thus aligning natural and sexual selection. This important but as yet untested prediction suggests that naturally selected camouflage patterns could serve as evolutionary precursors of some sexual signals.

Here, using humans as a model, we test the hypothesis that camouflage patterns are intrinsically visually attractive, and thus a pattern favored by natural selection can evoke preferences which then initiate sexual selection. In a first experiment, we used a detection task to assess the effectiveness of camouflage of different circular patterned stimuli that varied in their level of background matching. The backgrounds were artificial, but with spatial statistics either similar to those of many natural scenes or manipulated away from typical values. Participants were required to click on the circular pattern ('target') as quickly as possible to assess the detectability of the stimuli. In a second and third experiment, we evaluated the attractiveness of these targets, using the same patterned stimuli as for the detection task but removing the detectability constraint. Participants were asked to indicate which of the two patterns they preferred. In the second experiment, stimuli were presented in pairs against a uniform gray background to assess the baseline attractiveness of the target stimuli, independent of the background. In the third experiment, stimuli were presented in pairs against the same patterned backgrounds as for the detection task, so that the degree of background matching varied across stimuli, but stimuli were outlined to make them easily detectable. Our experiment therefore asked whether camouflage patterns are attractive when rendered detectable.

Results and Discussion

Generating variation in camouflage effectiveness

To evaluate the attractiveness of camouflage patterns, we first needed to generate patterned visual stimuli that differ in their camouflage effectiveness. To do so, we modified the Fourier slope of white noise images to manipulate the relative distribution of their spatial frequencies (from low to high spatial frequencies, corresponding to coarse-grained and fine-grained patterns, respectively). The Fourier slope is the relationship between power and spatial frequency when plotted on log-log axes, and that slope has been observed to lie close to -2 for many natural scenes. (e.g., *26*, *27*). With this approach, we could create abstract patterns and remove potential bias that would make interpretation more difficult (e.g. color or phase manipulation effects). Moreover, the Fourier slope is one of the most documented image statistics with effects on visual processing at different levels (i.e., on preferences, fluency, and neural encoding). We created circular images ("targets") and large square ("background") images and manipulated both the slope of the targets and the slope of their corresponding background in order to create different levels of background matching (Figure 1A). We designed the detection task with targets randomly positioned onto background images.

We postulated that targets with a slope closer to the slope of the background would be better camouflaged, that is, they would be less frequently detected and require more time to detect. Our results confirmed this prediction: these targets were the hardest to find (lower detection rates and higher detection times, Spearman correlation between detection rates and detection times, r = -0.98, N = 18).

Overall, results of the first experiment corroborate previous studies showing a link between Fourier slope and camouflage effectiveness. For example, octopuses and cuttlefish in camouflage mode modify their body pattern such that the Fourier slope of their body matches that of their visual environment (*27*, *28*). In damselfish, matching the spatial frequencies of the background reduces the likelihood of being attacked by predators (*29*). Our results complement these correlative studies, showing that the Fourier slope can be manipulated to generate variation in the camouflage effectiveness of background-matching stimuli.

***Figure 1: Detection task.** (A) Exemplars of presented stimuli with target values indicated below each image. (B) Detection rate (%) for the different conditions, computed as the number of trials for which the target was detected divided by the total number of presented trials. Each condition (i.e. one combination of background slope and target slope) is represented by five closed circles (made more visible using jittering) corresponding to five different stimuli versions (see Methods). Thin box plots indicate the amount of within-condition variation in detection rates between the stimuli versions. Background slope values are color-coded: blue for -1, orange for -2, and grey for -3. Dashed lines represent a quadratic fit to the data for each background separately (-1, -2, and -3). Arrows indicate the corresponding background value.*

Preference for natural statistics in abstract visual patterns

Our second experiment tested whether the patterns we generated for the targets in the first experiment also varied in their attractiveness. In a two-alternative forced-choice task (2-AFC), we assessed participants' preferences for the targets used in the detection task. The targets in this experiment were presented in pairs against a uniform grey background (Figure 2A). We calculated a preference score for each target using the Élő rating method, an iterative algorithm that performs pairwise comparisons and is commonly used to measure

preference (see e.g. *31*, *32*). All stimuli are then ranked based on their performance in paired comparisons, thereby obtaining a preference score.

Targets with differing Fourier slopes elicited markedly different preferences (Figure 2B). The resulting data can be fitted with a simple quadratic model (ranking~slope + slope²), which predicts a preference peak for a target slope of -2.36 (fit adjusted R^2: 0.9136). This preference peak matches the preferred peak value reported in previous studies (see e.g. (*23*)). The most widely accepted explanation for this value is that it matches the dominant Fourier slope of the natural visual environment (between -1.81 and -2.4 depending on the study, as reported by (*33*)).

To confirm that the slope with the peak preference score in our second experiment corresponds with the dominant slope of natural scenes, we estimated the slope of 4,319 images of the Landscape Pictures dataset (*33*), which contains images of different types of landscapes (e.g. mountain, beach, desert). Methods of Fourier slope calculation vary across the literature, thus any comparison of slope values across datasets must be based on the same calculation. We found an average slope of -2.39 (SD: 0.31) for those images. This is very similar to the preference peak inferred from our experiment on a gray background (-2.36), further corroborating the general finding that people prefer the Fourier slope of natural scenes. This result, combined with those from the detection task, validate the use of the Fourier slope to manipulate both the camouflage effectiveness and the attractiveness of stimuli. They also indicate a baseline pattern preference that must be accounted for when considering a preference for camouflaged stimuli in the third preference task.

Figure 2: Preference task on gray background. *(A) Exemplar of a presented stimulus. Target slopes are -1.25 for the left target and -0.25 for the right one. For display reasons, the background color here is lighter than that of the stimuli used in the experiment (see Methods). (B) Log-log plot of the averaged power spectrum from which the Fourier slope of an image is derived (red regression line), here representing the averaged Fourier slope of images from the Landscape Pictures dataset.*

(C) Élő rankings for the 2-AFC task with a grey background. Each condition (i.e. target slope values) is represented by five closed circles (made more visible using jittering) corresponding to five different stimuli versions. Dots indicate the mean ranking values for each of the five stimuli versions of each condition (indicated on the x-axis). Thin box plots give the variability of preference scores across those versions. The dotted line indicates a quadratic fit of the data.

A preference for the Fourier slope of natural scenes has been reported in numerous studies in experimental psychology. In a particularly compelling study, Spehar and collaborators (*23*) showed that stimuli to which people have the highest sensitivity (i.e. absolute detection, discrimination, and contrast sensitivities) are also the most attractive. Their result emphasizes the influence of efficient processing on visual preferences: patterns whose properties coincide with visual sensitivity are processed more efficiently by the visual system and are generally preferred over patterns that are further away from visual sensitivity peaks. These and our own results are therefore consistent with the processing bias hypothesis (*18*), which posits that the spatial statistics of natural scenes are attractive because they are the easiest to process by the brain. They also help explain a large body of results in empirical aesthetics, which show that visual artwork imitates natural spatial statistics (*34*, *35*). For example, faces painted by portrait artists across time and cultures tend to have a slope value matching that of natural landscapes even though real faces typically do not, suggesting that artists increase the attractiveness of their work by unconsciously mimicking the spatial statistics of natural scenes (*36*).

A preference for the Fourier slope of natural scenes is not only limited to humans. Hardenbicker and Tedore (*37*) recently discovered that peacock jumping spiders prefer images with the Fourier slope of natural scenes over two other options. In addition, in several fish species, Hulse and collaborators (*38*) found a correlation between the Fourier slope of species-typical habitats and the sexual signaling patterns of males (but not females). That study did not measure preferences, so whether fish are attracted to natural Fourier slopes remains to be tested, but a correlation between the spatial statistics of male sexual signals and their habitat is reminiscent of studies that demonstrate a correlation between the Fourier slope of human artwork and natural scenes.

Hidden preference for camouflage patterns
In our third experiment, we assessed the attractiveness of patterns that varied in their degree of background matching. We used the same stimuli as for the detection task (i.e., targets were displayed in front of a patterned background, rather than a gray background as in the second experiment), but we removed the detectability constraint by circling and centering the targets (Figure 3A). All targets therefore were assumed to be equally conspicuous. We assessed participants' preferences for different levels of background-matching patterns with a 2-AFC task and computed a preference score for each combination of background matching using the Élő-ranking method.

To explore the relationship between preference scores and background matching, we used linear regression to model preference scores as a response variable and the difference between target slope and background slope, as a measure of background matching, as an explanatory variable. We also included target slope and a quadratic term for background slope as additional potential explanatory variables. We found an effect of the difference between target slope and background slope on preference scores (estimate: -120.292 ±

12.015; t = -10.012, df = 89.413, p = 0.000, see Table S2). Stimuli with the lowest slope difference between target and background, i.e., with the highest background matching, tended to have higher preference scores. Similar significant effects were also obtained when data for each background were analyzed separately (Table S3). This, the main result of this study, indicates that targets that were more effectively camouflaged in the detection experiment were also more attractive when rendered detectable in the third preference experiment.

We also fitted a simple quadratic model (ranking~slope + slope^2) to find the target slope that predicts the highest attractiveness ranking for each background (data shown in Figure 3B). For a background slope of -1, a target slope of -1.92 predicts the highest ranking (77.67, fit adjusted R^2: 0.9993). For a background slope of -2, a target slope of -2.14 predicts the highest ranking (44.06, fit adjusted R^2: 0.9418). For a background slope of -3, a target slope of -2.51 predicts the highest ranking (80.32, fit adjusted R^2: 0.9943). These results indicate that preferences for visual patterns can be shifted away from a general baseline value (for Fourier slope -2.36, as determined in the second experiment) and toward the value of the background (Figure 3C), as predicted by a preference for background-matching stimuli and the processing bias hypothesis.

Notably, a similar modulation of visual attractiveness by background statistics was shown for human faces. Menzel and collaborators (*39*) varied the Fourier slope of the background against which faces were presented and asked participants to rate their attractiveness. They found that participants rated faces as more attractive when those faces were presented against a background that was closer to the Fourier slope of both natural scenes (-2) and faces (-3) compared to other background slopes (-1 and -4).

Figure 3. Preference task on a patterned background. *(A) Exemplars of presented stimuli with target values indicated below each image. (B) Élő rankings for the 2-AFC task with a patterned background. Each condition (i.e. one combination of background slope and target slope) is represented by five closed circles (made more visible using jittering) corresponding to five different stimuli versions. Thin box plots give the variability of preference scores across those 5 stimuli versions. Background slope values are colour-coded: blue for -1, orange for -2, and grey for -3. Dashed lines represent a quadratic fit to the data for each background separately (-1, -2, and -3) and*

arrows indicate the corresponding background value. C) Heuristic representation of the observed shift of preference of visual patterns away from the general baseline value of -2.36 (pink curve, a peak value obtained from Experiment 2) as a result of the modulation of the background slope.

Behavioral and evolutionary implications for animal communication

Given similarities in the early visual systems of humans and those of macaques (*40*), cats (*41*), mice (*42*) and birds (*43*), these results have important implications for animal sexual signaling. First, on a short time scale, they suggest that animals should track the pattern of the visual background against which they are displaying. If able, they should respond behaviourally or physiologically to match the statistical features of their patterns with those of the background. Similarly, if the sexual signal changes, as observed in some fish that change sex during development, individuals might choose another displaying site. Previous studies have shown that animals choose their display site in relation to their own sexual signals (*44*, *45*). Those studies, however, focused on coloration, showing that animals gain attractiveness by maximizing the contrast with the background and the ambient light. Our results extend these findings to signal patterns, but predict that, for this signal component, animals would benefit from mimicking the visual environment.

On an evolutionary scale, our results suggest that signal patterns will be shaped by the spatial statistics of the environment. Because the visual system adapts to efficiently process the spatial statistics of natural environments, patterns that mimic those statistics will also be efficiently processed and therefore attractive (*18*). This prediction of the processing bias hypothesis is an extension of "sensory exploitation" (*46*) or "sensory drive," which propose a similar mechanism for detectability, with animal signals evolving to be more easily detected (*47*). The African cichlid fish *Pundamilia nyererei,* for example, lives in an environment with red-shifted light. Females exhibit an increased expression of red-sensitive photoreceptors that render red stimuli more detectable, and males exhibit more reddish coloration (*48*). The sensory drive hypothesis is supported mainly by studies on coloration; our results suggest that expanding sensory drive beyond the framework of signal detection will allow us to predict the evolution of signal patterns, in addition to coloration.

Because traditional approaches to understanding the evolution of signal design are based on signal detection, natural and sexual selection are traditionally pitted in conflict along an axis of detectability. Our results suggest instead that these mechanisms can act synergistically. Natural selection generally favors camouflage, such as background-matching patterns. Due to their low detectability, the attractiveness of camouflage patterns is usually not considered, far less tested. However, if a mutation occurs that makes camouflaged patterns more easily detectable, for example by adding color or a sharp outline, their attractiveness should increase because of the efficiency with which they are processed. Hidden preferences for those efficiently processed patterns are predicted to lead to an increase in the frequency of attractive phenotypes if ecological conditions permit (Figure 4). In this way, camouflage patterns could be evolutionarily co-opted by sexual selection. This framework could therefore describe another pathway to the evolution of "magic traits" (*49*), i.e., traits that evolve due to divergent ecological selection and also affect mate choice.

One strategy to leverage the attractiveness of camouflage patterns might be seasonal variation in detectability. Individuals that are camouflaged most of the year could be rendered more attractive by the addition of a detectable feature during the mating season.

Male darter fish (genus *Etheostoma)*, for example, exhibit vivid and species-specific colors during the breeding season that appear to complement rather than conceal their underlying background-matching pattern (*53*, *54,* Figure 4C). Indeed, females in at least two species of darters are attracted similarly to conspecific color and pattern when these signal components are decoupled (*52*). An attractive and conspicuous seasonal addition could thus enhance, and be enhanced by, the underlying attractiveness of the camouflage pattern. Aposematic patterns can also be adaptively concealed, as in frogs, for which "hidden" aposematism on the belly represents an intermediate stage in the evolution of whole-body aposematic patterning (*53*). Interestingly, many aposematic patterns in frogs appear to be colorful embellishments of background-matching patterns (Figure 4B, *57*), and in *Oophaga pumilio,* for example, these patterns are attractive to the opposite sex (*55*). The attractiveness of aposematic patterns therefore also may be due to a combination of color and background-matching pattern, with sexual selection and natural selection similarly aligned in the evolution of warning signals.

In sum, we provide the first evidence that camouflage patterns matching the average image statistics of natural scenes are attractive, and although demonstrated here only in humans, this result is expected to generalize to other species. By providing evidence of a processing bias, the key mechanism by which camouflage patterns can be co-opted to function as sexual signals, this study highlights the importance of considering the psychological landscape (*56*) of signal receivers, and of interfacing research in evolutionary biology with the vast literature in human cognitive psychology.

Figure 4. Evolution from camouflage to sexual signalling through the exploitation of processing bias.
Panel A: 1. *Natural selection due to predation generally results in inconspicuous phenotypes. Phenotypes with a pattern that matches that of the environment ('inconspicuous pattern') are well-camouflaged and thus gradually increase in frequency in the population. Such phenotypes typically do not elicit strong preferences among conspecifics because of their low detectability.* **2.** *Any new mutation increasing conspicuousness should be negatively selected by predation (low survival). However, because they are highly detectable by potential mates and thus reduce their cost of mating, these phenotypes are potentially attractive. Among conspicuous phenotypes, camouflage patterns that match the visual structure of the habitat ('conspicuous patterned') should be preferred (ΔP) over those that do not ('conspicuous plain') because they are even more efficiently processed. In this study, we have provided the first empirical evidence supporting this prediction.* **3.** *These 'sleeping beauties' would remain negatively selected by natural selection until a change in ecological factors (e.g., colonization of habitats or regions with fewer predators, the appearance of new, alternative prey, predators' focus on new sensory cues) makes sexual selection strong enough to counteract the effects of predation on*

*phenotype evolution **4**. Thus, conspicuous phenotypes that have evolved from camouflage patterns can be evolutionarily co-opted by sexual selection. **Lower panels***: Examples of camouflage patterns underlying conspicuous color and patterns. B:* Etheostoma caeruleum *(Julien Renoult). C:* Ranitomeya reticulata *(DH Fischer). D: Painting of a peacock (*Pavo cristatus*) camouflaged in a tree by Abbott H. Thayer. Thayer, a naturalist painter of the 18th century, explored the idea that the primary function of animal patterns is camouflage. Although widely criticized at the time, Thayer's idea is consistent with the proposed framework, that even the most canonical nuptial patterns can evolve from, and continue to function at least in part as, camouflage. (Abbott Handerson Thayer, Richard S. Meryman, Peacock in the Woods, study for book Concealing Coloration in the Animal Kingdom, 1907, oil on canvas, Smithsonian American Art Museum).*

References and Notes

1. M. Andersson, *Sexual Selection* (Princeton University Press, 1994; https://press.princeton.edu/books/paperback/9780691000572/sexual-selection).
2. J. A. Endler, "Natural and sexual selection on color patterns in poeciliid fishes" in *Evolutionary Ecology of Neotropical Freshwater Fishes: Proceedings of the 1st International Symposium on Systematics and Evolutionary Ecology of Neotropical Freshwater Fishes, Held at DeKalb, Illinois, U.S.A., June 14–18, 1982*, T. M. Zaret, Ed. (Springer Netherlands, Dordrecht, 1984; https://doi.org/10.1007/978-94-015-7682-6_7)*Developments in environmental biology of fishes*, pp. 95–111.
3. F. Breden, G. Stoner, Male predation risk determines female preference in the Trinidad guppy. *Nature* **329**, 831–833 (1987).
4. J. A. Endler, A. E. Houde, Geographic Variation in Female Preferences for Male Traits in Poecilia Reticulata. *Evolution* **49**, 456–468 (1995).
5. A. K. Schwartz, A. P. Hendry, A test for the parallel co-evolution of male colour and female preference in Trinidadian guppies (Poecilia reticulata). *Evol. Ecol. Res.* **9**, 71–90 (2007).
6. J. A. Endler, Natural Selection on Color Patterns in Poecilia Reticulata. *Evolution* **34**, 76–91 (1980).
7. J. A. Endler, Natural and sexual selection on color patterns in poeciliid fishes. *Environ. Biol. Fishes* **9**, 173–190 (1983).
8. D. Gomez, C. Richardson, T. Lengagne, S. Plenet, P. Joly, J.-P. Léna, M. Théry, The role of nocturnal vision in mate choice: females prefer conspicuous males in the European tree frog (Hyla arborea). *Proc. R. Soc. B Biol. Sci.* **276**, 2351–2358 (2009).
9. M. J. Ryan, M. E. Cummings, Perceptual Biases and Mate Choice. *Annu. Rev. Ecol. Evol. Syst.* **44**, 437–459 (2013).
10. C. Waitt, A. C. Little, S. Wolfensohn, P. Honess, A. P. Brown, H. M. Buchanan-Smith, D. I. Perrett, Evidence from rhesus macaques suggests that male coloration plays a role in female primate mate choice. *Proc. R. Soc. Lond. B Biol. Sci.* **270**, S144–S146 (2003).
11. K. Delhey, M. Valcu, C. Muck, J. Dale, B. Kempenaers, Evolutionary predictors of the specific colors of birds. *Proc. Natl. Acad. Sci.* **120**, e2217692120 (2023).
12. J. A. Endler, D. A. Westcott, J. R. Madden, T. Robson, Animal Visual Systems and the Evolution of Color Patterns: Sensory Processing Illuminates Signal Evolution. *Evolution* **59**, 1795–1818 (2005).
13. M. C. Stoddard, D. Osorio, Animal Coloration Patterns: Linking Spatial Vision to Quantitative Analysis. *Am. Nat.* **193**, 164–186 (2019).
14. R. L. Fantz, Form preferences in newly hatched chicks. *J. Comp. Physiol. Psychol.* **50**, 422–430 (1957).
15. E. Munar, G. Gómez-Puerto, J. Call, M. Nadal, Common Visual Preference for Curved Contours in Humans and Great Apes. *PLOS ONE* **10**, e0141106 (2015).
16. R. Reber, N. Schwarz, P. Winkielman, Processing Fluency and Aesthetic Pleasure: Is Beauty in the Perceiver's Processing Experience?: *Personal. Soc. Psychol. Rev.*, doi: 10.1207/s15327957pspr0804_3 (2004).
17. J. P. Renoult, J. Bovet, M. Raymond, Beauty is in the efficient coding of the beholder. *R. Soc. Open Sci.* **3**, 160027 (2016).
18. J. P. Renoult, T. C. Mendelson, Processing bias: extending sensory drive to include efficacy and efficiency in information processing. *Proc. R. Soc. B Biol. Sci.* **286**, 20190165 (2019).
19. B. A. Olshausen, D. J. Field, Sparse coding with an overcomplete basis set: A strategy employed by V1? *Vision Res.* **37**, 3311–3325 (1997).
20. E. P. Simoncelli, B. A. Olshausen, Natural Image Statistics and Neural Representation. *Annu. Rev. Neurosci.* **24**, 1193–1216 (2001).
21. H. B. Barlow, "Possible Principles Underlying the Transformations of Sensory

Messages" in *Sensory Communication*, W. A. Rosenblith, Ed. (The MIT Press, 1961; http://mitpress.universitypressscholarship.com/view/10.7551/mitpress/9780262518420. 001.0001/upso-9780262518420-chapter-13), pp. 216–234.

22. H. A. Geller, R. Bartho, K. Thömmes, C. Redies, Statistical image properties predict aesthetic ratings in abstract paintings created by neural style transfer. *Front. Neurosci.* **16**, 999720 (2022).

23. B. Spehar, S. Wong, S. van de Klundert, J. Lui, C. Clifford, R. Taylor, Beauty and the beholder: the role of visual sensitivity in visual preference. *Front. Hum. Neurosci.* **9** (2015).

24. I. C. Cuthill, Camouflage. *J. Zool.* **308**, 75–92 (2019).

25. G. J. Burton, I. R. Moorhead, Color and spatial structure in natural scenes. *Appl. Opt.* **26**, 157–170 (1987).

26. B. A. Olshausen, D. J. Field, Natural image statistics and efficient coding. *Netw. Comput. Neural Syst.* **7**, 333–339 (1996).

27. N. Josef, P. Amodio, G. Fiorito, N. Shashar, Camouflaging in a Complex Environment—Octopuses Use Specific Features of Their Surroundings for Background Matching. *PLOS ONE* **7**, e37579 (2012).

28. S. Zylinski, M. J. How, D. Osorio, R. T. Hanlon, N. J. Marshall, To Be Seen or to Hide: Visual Characteristics of Body Patterns for Camouflage and Communication in the Australian Giant Cuttlefish Sepia apama. *Am. Nat.* **177**, 681–690 (2011).

29. G. A. C. Phillips, M. J. How, J. E. Lange, N. J. Marshall, K. L. Cheney, Disruptive colouration in reef fish: does matching the background reduce predation risk? *J. Exp. Biol.* **220**, 1962–1974 (2017).

30. A. P. Clark, K. L. Howard, A. T. Woods, I. S. Penton-Voak, C. Neumann, Why rate when you could compare? Using the "EloChoice" package to assess pairwise comparisons of perceived physical strength. *PLOS ONE* **13**, e0190393 (2018).

31. A.-S. Tribot, Q. Carabeux, J. Deter, T. Claverie, S. Villéger, N. Mouquet, Confronting species aesthetics with ecological functions in coral reef fish. *Sci. Rep.* **8**, 11733 (2018).

32. T. Pouli, E. Reinhard, D. W. Cunningham, *Image Statistics in Visual Computing* (CRC Press, 2013).

33. A. Rougetet, Landscape Pictures, version 1 (2020); https://www.kaggle.com/datasets/arnaud58/landscape-pictures.

34. D. Graham, D. Field, Statistical regularities of art images and natural scenes: Spectra, sparseness and nonlinearities. *Spat. Vis.* **21**, 149–164 (2008).

35. D. Graham, C. Redies, Statistical regularities in art: Relations with visual coding and perception. *Vision Res.* **50**, 1503–1509 (2010).

36. C. Redies, J. Hasenstein, J. Denzler, Fractal-like image statistics in visual art: similarity to natural scenes. *Spat. Vis.* **21**, 137–148 (2007).

37. M.-C. Hardenbicker, C. Tedore, Peacock spiders prefer image statistics of average natural scenes over those of male ornamentation. *Behav. Ecol.*, arad042 (2023).

38. S. V. Hulse, J. P. Renoult, T. C. Mendelson, Sexual signaling pattern correlates with habitat pattern in visually ornamented fishes. *Nat. Commun.* **11**, 2561 (2020).

39. C. Menzel, G. U. Hayn-Leichsenring, O. Langner, H. Wiese, C. Redies, Fourier Power Spectrum Characteristics of Face Photographs: Attractiveness Perception Depends on Low-Level Image Properties. *PLOS ONE* **10**, e0122801 (2015).

40. D. Fize, M. Cauchoix, M. Fabre-Thorpe, Humans and monkeys share visual representations. *Proc. Natl. Acad. Sci.* **108**, 7635–7640 (2011).

41. I. Kang, R. E. Reem, A. L. Kaczmarowski, J. G. Malpeli, Contrast Sensitivity of Cats and Humans in Scotopic and Mesopic Conditions. *J. Neurophysiol.* **102**, 831–840 (2009).

42. B. Williams, J. D. Rosario, T. Muzzu, K. Peelman, S. Coletta, E. K. Bichler, A. Speed, L. Meyer-Baese, A. B. Saleem, B. Haider, Spatial modulation of dark versus bright stimulus responses in the mouse visual system. *Curr. Biol.* **31**, 4172-4179.e6 (2021).

43. W. Clark, M. Chilcott, A. Azizi, R. Pusch, K. Perry, M. Colombo, Neurons in the pigeon visual network discriminate between faces, scrambled faces, and sine grating images.

Sci. Rep. **12**, 589 (2022).

44. J. A. Endler, M. Thery, Interacting Effects of Lek Placement, Display Behavior, Ambient Light, and Color Patterns in Three Neotropical Forest-Dwelling Birds. *Am. Nat.* **148**, 421–452 (1996).

45. L. J. Fleishman, C. A. Perez-Martinez, M. Leal, Can Sensory Drive Explain the Evolution of Visual Signal Diversity in Terrestrial Species? A Test with Anolis Lizards. *Am. Nat.*, 000–000 (2022).

46. M. J. Ryan, J. H. Fox, W. Wilczynski, A. S. Rand, Sexual selection for sensory exploitation in the frog Physalaemus pustulosus. *Nature* **343**, 66–67 (1990).

47. J. A. Endler, A. L. Basolo, Sensory ecology, receiver biases and sexual selection. *Trends Ecol. Evol.* **13**, 415–420 (1998).

48. O. Seehausen, Y. Terai, I. S. Magalhaes, K. L. Carleton, H. D. J. Mrosso, R. Miyagi, I. van der Sluijs, M. V. Schneider, M. E. Maan, H. Tachida, H. Imai, N. Okada, Speciation through sensory drive in cichlid fish. *Nature* **455**, 620–626 (2008).

49. M. R. Servedio, G. S. Van Doorn, M. Kopp, A. M. Frame, P. Nosil, Magic traits in speciation: "magic" but not rare? *Trends Ecol. Evol.* **26**, 389–397 (2011).

50. R. A. Kuehne, R. W. Barbour, *The American Darters* (University Press of Kentucky, 1983; https://www.jstor.org/stable/j.ctt130hpj2).

51. L. M. Page, *Handbook of Darters* (Tfh Pubns Inc, Neptune City, NJ, First Edition., 1983).

52. T. H. Williams, T. C. Mendelson, Female preference for male coloration may explain behavioural isolation in sympatric darters. *Anim. Behav.* **82**, 683–689 (2011).

53. K. Loeffler-Henry, C. Kang, T. N. Sherratt, Evolutionary transitions from camouflage to aposematism: Hidden signals play a pivotal role. *Science* **379**, 1136–1140 (2023).

54. D. Galindo Uribe, J. Sunyer, S. Hauswaldt, A. Amézquita, H. Proehl, M. Vences, Colour and pattern variation and Pleistocene phylogeographic origin of the strawberry poison frog, Oophaga pumilio, in Nicaragua. *Salamandra* **50**, 225–235 (2014).

55. M. E. Maan, M. E. Cummings, Female Preferences for Aposematic Signal Components in a Polymorphic Poison Frog. *Evolution* **62**, 2334–2345 (2008).

56. T. Guilford, M. S. Dawkins, Receiver psychology and the evolution of animal signals. *Anim. Behav.* **42**, 1–14 (1991).

57. J. R. de Leeuw, jsPsych: A JavaScript library for creating behavioral experiments in a Web browser. *Behav. Res. Methods* **47**, 1–12 (2015).

58. C. Neumann, EloChoice: Preference Rating for Visual Stimuli Based on Elo Ratings, version 0.29.4 (2019); https://rdrr.io/cran/EloChoice/.

59. R. Goodspeed, Research note: An evaluation of the Elo algorithm for pairwise visual assessment surveys. *Landsc. Urban Plan.* **157**, 131–137 (2017).

60. S. Thorpe, D. Fize, C. Marlot, Speed of processing in the human visual system. *Nature* **381**, 520–522 (1996).

61. D. Bates, M. Mächler, B. Bolker, S. Walker, Fitting Linear Mixed-Effects Models Using lme4. *J. Stat. Softw.* **67**, 1–48 (2015).

62. J. Henrich, S. J. Heine, A. Norenzayan, The weirdest people in the world? *Behav. Brain Sci.* **33**, 61–83; discussion 83-135 (2010).

Acknowledgements:

We would like to thank Claudia Ximena Restrepo-Ortiz for her help with the Spanish translation of the online experiment.

Funding: This work was supported by the National Science Foundation grant NSF IOS 2026334.

Author contributions:
Conceptualization: YHB, MR, ICC, TCM, JPR
Methodology: YHB, MR, TCM, JPR
Investigation: YHB, TCM, JPR
Visualization: YHB, TCM, JPR
Funding acquisition: TCM, JPR
Project administration: TCM, JPR
Supervision: TCM, JPR
Writing – original draft: YHB, TCM, JPR
Writing – review & editing: YHB, MR, ICC, TCM, JPR

Competing interests: Authors declare that they have no competing interests.

Data and materials availability: All data and code, and materials used in the analysis are available on the following GitLab repository:
https://gitlab.com/yseulthb/camo_exp.