TITLE: Gas5 regulates early life stress-induced anxiety and spatial memory

Dipanjana Banerjee, Sania Sultana and Sourav Banerjee

National Brain Research Centre, NH – 8, Nainwal Mode, Manesar 122052, Haryana,

India

Correspondence: souravnbrc@gmail.com or sourav@nbrc.ac.in

Keywords: Maternal separation, stress, long non-coding RNA, synapse, anxiety,

spatial memory.

Abbreviations

- ***Ago****: Argonaute*
- ***AAV****: Adeno-Associated Virus*
- ***AMPA****: α-Amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid*
- ***ANOVA****: Analysis of Variance*
- ***BCA****: Bicinchoninic Acid*
- ***BSA****: Bovine Serum Albumin*
- ***CA1****: Cornu Ammonis 1*
- ***CA2****: Cornu Ammonis 2*
- ***CA3****: Cornu Ammonis 3*
- ***CIRTs****: CRISPR-Cas-inspired RNA targeting systems*
- ***CD1****: Outbred Swiss mice*
- ***CLIP-seq****: Cross-Linking Immunoprecipitation followed by high-throughput sequencing*
- ***CMV****: Cytomegalovirus*
- ***CPCSEA****: Committee for the Purpose of Control and Supervision of Experiments on Animals*
- ***CRISPR****: Clustered Regularly Interspaced Short Palindromic Repeats*
- ***cDNA****: Complementary DNA*
- ***DH5α****: Escherichia coli strain*
- ***DI****: Discrimination Index*
- ***DMEM****: Dulbecco's Modified Eagle Medium*
- ***DIV****: Days In Vitro*
- ***DNA****: Deoxyribonucleic Acid*
- ***dF****: Degrees of Freedom*
- ***ECL****: Enhanced Chemiluminescence*
- ***EDTA****: Ethylenediaminetetraacetic Acid*
- ***EF1-α****: Elongation Factor 1-alpha*
- ***EGFP****: Enhanced Green Fluorescent Protein*
- ***ES****: Early life stress*
- ***FBS****: Fetal Bovine Serum*
- ***FDR****: False Discovery Rate*
- ***Gas5****: Growth Arrest Specific-5*
- ***GFP****: Green Fluorescent Protein*
- ***GABAergic****: Gamma-Aminobutyric Acidergic*

•   *GO*: Gene Ontology

•   *HEK293T*: Human Embryonic Kidney 293 Cells, expressing large T antigen of SV40

•   *HC*: Homecaged

•   *IAEC*: Institutional Animal Ethics Committee

•   *LSD*: Least Significant Difference

•   *MOI*: Multiplicity of Infection

•   *MRI*: Magnetic Resonance Imaging

•   *MS*: Maternal separation

•   *NcA*: Nucleus accumbens

•   *NEB*: New England Biolabs

•   *NMDA*: N-Methyl-D-Aspartate

•   *PBS*: Phosphate Buffered Saline

•   *PCR*: Polymerase Chain Reaction

•   *PFA*: Paraformaldehyde

•   *P*: Postnatal Day

•   *POSTAR3*: Post-transcriptional Regulation Atlas of lncRNAs and mRNAs

•   *Pvt1*: Plasmacytoma Variant Translocation 1

•   *qPCR*: Quantitative Polymerase Chain Reaction

•   *RBPs*: Ribosome Binding Proteins

•   *qRT-PCR*: Quantitative Reverse Transcription Polymerase Chain Reaction

•   *RISC*: RNA-Induced Silencing Complex

•   *Rmst1*: rhabdomyosarcoma 2-associated transcript 1

•   *RNAs*: Ribonucleic acids

•   *RNP*: Ribonucleoprotein

•   *SDS*: Sodium Dodecyl Sulfate

•   *SEM*: Standard Error of the Mean

•   *shRNAs*: Short hairpin RNAs

•   *Snhg8*: Small Nucleolar RNA Host Gene 8

•   *Snhg12*: Small Nucleolar RNA Host Gene 12

•   *SV40*: Simian Virus 40

•   *SynLAMP*: Synaptic Lysosome-associated Membrane Protein

•   *Tubulinβ-1*: Beta-1 tubulin

•   *UC*: Universal Control

•   *Zfas1*: Zinc finger antisense 1

•   *ZsGreen*: Zoanthus sp. Green Fluorescent Protein

ABSTRACT:

Early life stress (ES) induced by maternal separation (MS) remains a proven causality of anxiety and memory deficits at later stages of life. Emerging studies have shown that MS-induced gene expression in the hippocampus is operated at the level of transcription. However, the extent of involvement of non-coding RNAs in MS-induced behavioural deficits remains unexplored. Here, we have investigated the role of synapse-enriched long non-coding RNAs (lncRNAs) in anxiety and memory upon MS. We observed that MS led to an enhancement of expression of the lncRNA Gas5 (Growth Arrest Specific-5) in the hippocampus; accompanied by increased levels of anxiety and deficits in spatial memory. Gas5 knockdown in early life was able to reduce anxiety and partially rescue the spatial memory deficits of maternally separated adult mice. However, the reversal of MS-induced anxiety and memory deficits is not attributed to Gas5 activity during neuronal development as Gas5 RNAi did not influence spine development. Gene Ontology analysis revealed that Gas5 exerts its function by regulating RNA metabolism and translation. Our study highlights the importance of MS-regulated lncRNA in anxiety and spatial memory.

INTRODUCTION

Maternal separation, a well-established model of early-life stress, has been extensively studied to investigate the long-term consequences of adverse early life events during this critical period of brain function and behaviour (Taylor et al., 2010; Korosi et al., 2012). Stress due to maternal separation (MS) in rodents leads to anxiety-like behaviours in adulthood as characterized by heightened fear, avoidance, and stress sensitivity (Banqueri et al., 2017; Lundberg et al., 2020; Balouek *et al.*, 2023). The ventral hippocampus, a key brain area affected by MS (Dimatelis et al., 2014; Marais et al., 2009), has been shown to regulate emotions (Jimenez et al., 2018; H. Li et al., 2022; McNaughton, 1997). MS-induced anxiety can lead to changes in the total volume of the hippocampus (Kalisch et al. 2006) and alterations in the expression of glutamatergic receptors (Pickering et al. 2006), both NMDA (Barkus et al. 2010) and AMPA (Katsouli et al. 2014); GABAergic receptors (Sterley, Howells, and Russell 2013), and serotonergic as well as cholinergic receptors (File et al., 2000). Moreover, early-life stress can impact spatial memory that involves the dorsal hippocampus

(Morris *et al.*, 2006; Blum *et al.*, 1999). Such behavioural deficits are linked with

structural and functional modifications in the hippocampus (Izquierdo and Medina,

1997; Revest *et al.*, 2009; Kremmyda *et al.*, 2016). These observations indicate that

the enduring effects of maternal separation on anxiety and memory involve alterations

in synaptic plasticity (Monroy et al., 2010; Bannerman *et al.*, 2014) and

neurotransmitters, for example, noradrenaline and serotonin (Daniels et al. 2004).

More recently, transcriptomics analysis following MS identified a distinct set of miRNAs

in specific brain areas including the hippocampus (Mckibben and Dwivedi 2021).

Some of these miRNAs are localized at the synapse and have been shown to regulate

synaptic plasticity and memory (Mckibben and Dwivedi 2021). A recent study from our

laboratory has identified a set of synapse-enriched lncRNAs from the hippocampus.

Among these transcripts, a specific lncRNA has been shown to regulate hippocampus-

dependent memory (Samaddar et al., 2023). LncRNA functions as scaffolds, decoys,

guides, or enhancers to modulate chromatin remodelling, transcription, mRNA

stability, and RNA-protein interactions (Mattick et al., 2023; Mercer et al., 2009).

Recent attention has focused on lncRNA activity at synapses that govern synaptic

plasticity, neuronal development, and neurological disorders (Zhang, Hamblin, and Yin

2017) (Shi, Zhang, and Qin 2017) (A. Wang et al. 2017). The regulatory function of

lncRNA has also been attributed to anxiety (Spadaro et al. 2015) and memory

(Soutschek et al. and Schratt et al., 2023; Cui et al., 2022).

Prompted by these observations, we have investigated the impact of MS on the

expression of lncRNAs and their involvement in MS-induced anxiety and memory. Our

study identifies that MS enhances the expression of the lncRNA Gas5. The knockdown

of Gas5 alone shows no significant changes in spine development in adult mice.

However, when accompanied by early-life stress, Gas5 RNAi partially alleviates the

enduring consequences of the stress as detected by the reduction in anxiety and the

occlusion of memory deficits following MS.

MATERIAL AND METHODS

Animal ethics, acquisition, and maintenance

All animals were issued from the National Brain Research Centre (NBRC) animal house facility from the protocol numbers NBRC/IAEC/2015/171, NBRC/IAEC/2015/172 and NBRC/IAEC/2015/178 approved by IAEC (Institutional Animal Ethics Committee) of NBRC registered with the CPCSEA (Committee for the Purpose of Control and Supervision of Experiments on Animals) (Registration number: 464/GO/ReBi-S/Re-L/01/CPCSEA), Government of India.

Animals were housed in standard housing facilities in cages with food water *ad libitum*. A temperature of around 24± 0.5 °C and humidity of 40 ± 5% was maintained along with a 12-hour light/dark cycle for all animals.

Maternal Separation

In our study, we have focused on the aspects of induction of stress only due to maternal separation and not early social isolation, where pups are separated from the mother as well as from each other (Alves et al., 2022). The maternal separation paradigm was used as a chronic early-life stress paradigm for C57BL/6 mice pups (Sood et al. 2018). A litter size of 5-7 pups was selected for the experiment. 1 litter of pups or 1 dam was considered 'N' and was arbitrarily allocated into homecaged (HC) and maternally separated (MS) groups. A total of 32 dams (120 pups (n)) were used for maternal separation throughout the study. Postnatal day 2 (P2) pups were separated from their mother for a period of 3 hours (0900 to 1200) till P14. The mother was first moved into a different fresh cage and kept in a separate room. The pups were then placed in a beaker with soft bedding and protective paper lining and the temperature was maintained at 37˚C throughout separation. After 3 hours, first, the pups were kept back in the original cage followed by the mother. No separation was done from P15 till weaning on P21. Handling of the pups was done very gently. Cage changes were done every three days starting from P4. For the control group, designated as HC, all litters and mothers were kept undisturbed from P0 to P21 except for regular maintenance and cage changes every three days. Male pups were

190 sacrificed at P21 (euthanization with CO_2 gas), and hippocampi were isolated for
191 further biochemical processes.

Screening of Lnc RNAs

P21 male and female pups (N=14) were sacrificed in a CO_2 euthanasia chamber at a
30-70% fill rate of CO_2 gas and death was confirmed by tail/feet pinch and lack of
respiration. Euthanasia was followed by a quick decapitation of the animals and the
whole hippocampi was harvested in Trizol reagent (Invitrogen) for total RNA isolation.
SuperScript III First Strand Synthesis System (Life Technologies, Invitrogen) kit was
used for synthesizing cDNA following the manufacturer's protocol. Further, qPCR of
the cDNA prepared was done using SyberGreen (Applied Biosystems). See Table 1
for qPCR primer details.

TABLE 1. Primers for qPCR

No	LncRNA	Forward Primer (5' – 3')	Reverse Primer (5' – 3')
1	Tubulinβ-1	GCGCCTTTGGACACCTATTCA	CCGCGCCCTCCGTATAGTGC
2	GAPDH	CGTATTGGGCGCCTGGTCAC	CGGCCTCACCCCATTTGATG
4	Gas5	AGGAGGATGAAGGCTTACGA	AACTGTCTATCCAGCCTCAA
5	Pvt1	TTGCCTGGAGATGTTACAAG	AACATTCTCATTGAGTCAAG
6	SynLAMP (2410006H16Rik)	AAGAGTGGAAGATTGCCAAC	GATTACCAAATGTGCCCACC
7	2900052NO1 Rik	ACAATCCAAATCTACCCACGAG	TGGGATTGTAGATTGTGCTGTC
	1700020IRik	TGACTGCATCTCCGAGGAAT	GGCCTGCTACTATTGCTTACA
9	Zfas1	CGCCGCCAAGGTCTATATAA	CAGGTCAACCATCTGCAAGT
10	Pantr1	ATTTCCTGAGTCAACATGGC	AGAGTATGCCACATGGATGATG
11	Snhg8	CAGTGGAGGAAGTGTTAAGGT	CCACTACAGTACCTGCAAGAA
12	Snhg12	TCTCCGGATGATTCGTGAGA	TCTGCTGTATCGTTGTCCAT
13	Rmst1	CATGAATGTTGTCGACACTGT	GAGCTTGGCAGATCCTTCAA

Cloning and Virus preparation

EF1-α promoter from pLVTHM (Addgene # 12247) was used to replace the CMV
promoter in pAAV.U6.shRLuc.CMV.ZsGreen.SV40 (Penn Core Vector) plasmid by
excising out the CMV promoter using the EcoRI (NEB #R3101) and NHE-I (NEB
#R3131) cut sites. Infusion cloning (Takara-Bio In-Fusion HD cloning Kit; 639650) was
used to clone in EF1-alpha promoter generating a pAAV.U6.shRLuc.EF1-
α.ZsGreen.SV40 plasmid and express ZsGreen under EF1-α promoter (Table 2).
Universal Control (UC) and Gas5 shRNAs (shRNA # 1= G5C1 and shRNA # 2= G5C2)
were cloned by replacing the existing Luciferase shRNA (shRLuc) under a U6
promoter using BamHI (NEB #R0136L) and EcoRI restriction sites (Table 3). The final
vector was coined as pAAV.U6.shRNA(UC).EF1-α.ZsGreen.SV40,
pAAV.U6.shRNA(G5C1).EF1-α.ZsGreen.SV40 and pAAV.U6. shRNA(G5C2).EF1-
α.ZsGreen.SV40 respectively.

ZsGreen was replaced with EGFP in all vectors to achieve a diffused fluorescence for
imaging. All vectors were digested with NcoI-HF (NEB #R3193) and NotI-HF (NEB
#R3189) to excise out ZsGreen. EGFP from the pLVTHM vector was inserted by
infusion cloning (Table 4).

All clones were verified by Sanger sequencing.

TABLE 2. Infusion primers for replacing CMV with EF1-α in AAV backbone

No.	Primers 5'-3'	Oligo sequence
1.	Forward	TTTTACGCGTGAATTCGGCTCCGGTGCCCGTCAG
2.	Reverse	GACCGGTAGCGCTAGCTCACGACACCTGAAATGGAAG

224 TABLE 3. shRNA Antisense Sequence

No.	Lnc-RNA	Oligo Sequence (5' – 3')
1.	UC shRNA	ATCTCGCTTGGGCGAGAGTAAG
2.	G5C1 shRNA	GAGGCTGGATAGACAGTTTGA
3.	G5C2 shRNA	GGTCCTTCATTCTGAATTTCA

TABLE 4. Infusion primers for replacing ZsGreen with EGFP in AAV backbone

No.	Primers 5'-3'	Oligo sequence
1.	Forward primer 5'-3'	CCGGTCGCCACCATGATGGTGAGCAAGGGCGAGG
2.	Reverse primer 5'-3'	CTGCTCGAAGCGGCCCTTGTACAGCTCGTCCATGCC

The expression vector was propagated using the *E.coli* Stbl3 strain and the AAV packaging plasmids pAdDeltaF6 and pAAV2.9 (Penn vector Core) were propagated using the DH5α strain. Endotoxin-free Maxiprep kit (Qiagen) was used to prepare purified plasmid for HEK293T cell transfection. Calcium phosphate transfection was done using 10µg of the above-mentioned transfer vector, 10µg pAAV2/9n and 20µg of pAdDeltaF6 into HEK293T cells of density 2×10^6 cells. The culture was maintained at 37℃, 5% CO_2 in low-glucose DMEM (Gibco) with 10% foetal bovine serum with one-time media change after 8 hours. Post 96 hours of infection, NaCl (final concentration: 500 mM) was added to the plates for 3 hours to lyse the cells. The supernatant was collected and stored at 4℃. The remaining cells on the plates were scraped and collected separately. The cells went through three freeze-thaw cycles where they were frozen at -80℃ and thawed at 37℃ for complete dissociation of virus particles from the cell membrane. The filtrate was added to the supernatant collected before; together they were filtered through a 100µ and a 0.45µ filter successively. The final filtrate, containing the virus, was then centrifuged using Amicron-Ultra-15 (Merk Millipore #UFC910024) filters to concentrate the virus and determine the titre by fluorescence-based cell counter. The multiplicity of infection (MOI) was calculated by infecting 80K HEK293T cells with serially diluted AAV solutions. An MOI of 1-2 was used for further experiments.

Primary Neuronal Culture

Hippocampal neurons were cultured following the previous protocol (Kaech & Banker, 2006). CD1 dams were euthanized in a CO_2 gas chamber (fill rate 30-70%) followed by cervical dislocation and death was confirmed by tail/feet pinch. The uterus containing the embryonic day 16 (E16) pups was dissected and taken out on ice-cold dissection media (Gibco) to render the pups unresponsive to painful stimuli. An incision

was made in the uterus to extract the pups for decapitation and the whole brain was
collected from the sacrificed pups. Hippocampi from all pups from a single CD1 (N=4)
were pooled for further experiments. Briefly, hippocampi from embryonic pups were
dissected and collected in dissection media (Gibco). It was then digested with the
addition of 2.5% trypsin to the dissection media at 37℃ for 10 mins. Trypsin-treated
tissue was washed with dissection media and collected in Glial Minimum Essential
Media (GMEM; Gibco). Further, trituration was done with a fire-polished Pasteur
pipette to obtain a single-cell suspension which was plated on poly-L-lysine coated
60mm dishes (1 mg/mL, Sigma) at a concentration of 200-250 cells/mm^2. The culture
was maintained at 37℃, 5% CO_2 in Neurobasal Media (Gibco) with B27 supplement
(Gibco) throughout the experiment. Cells were infected with AAV at DIV 7 and
harvested at DIV 21 for further experiments.

Stereotaxic injection

Surgery was done in P0 pups (N=56, n=168) 6-8 hours post-birth only after the
appearance of the milk spot (Kim et al. 2014). Half of the litter was taken at a time and
kept on a heating pad at 37°C. For anaesthesia, the pup was kept on a cold metal
plate kept on the ice for 2-3 mins till there was no movement and the body colour
turned whitish pink from reddish pink. It was then lightly taped to the platform with its
head erect. The platform was kept at 7-8℃ to maintain the anaesthesia. In case the
pup's body colour started to show a bluish tinge, the temperature was raised to 12-
15℃. Stereotaxic arms were moved to (X, Y) = (0.8, 1.5) mm coordinates and the
needle was lowered till at Z= -1.8 mm (Christensen et al. 2010). Each hippocampus
was injected with 0.5 µl of AAV with MOI=1. A volume of 0.1 µl was injected per minute
with a gap of 10 mins. On completion of 0.5 µl virus injection, an additional gap of 10
mins was given before the needle was pulled out very slowly to prevent any back
suction. We proceeded to inject into the second hippocampus after 15 minutes during
which the first wound site was allowed to close. The pup was then kept on a 37℃
heating pad and warmed till movement and colour were regained. They were returned
to the mother together to minimise disturbances. The cage was left completely
undisturbed for the next day. (n=163; surgeries performed for western blot, spine

analysis, light dark assay, spatial object recognition assay. 'n' determines the number

of individual animals; males).

Light/dark box assay

The apparatus consists of a smaller dark box with a light intensity of 2 Lux (one-third

of the total area) and a bigger light box with a light intensity of 390 Lux (two-thirds of

the total area) separated by a door (Bourin and Hascoët 2003) (Takao and Miyakawa

2006). P42 mice (N=26, n=79) were habituated for 2 minutes to handling and to a

transport vessel in dim light (2-5 Lux) for 1 week in the experimental room. Animals

were housed in standard cages and a 12-hour light/dark cycle was maintained as

usual. On test day, the mouse was kept in the dark box with a closed door. The door

was opened 10 seconds later and the mouse was left free to explore the light box for

10 minutes. The behaviour assays and habituations were performed between 0900

hours to 1800 hours. Arenas were cleaned using 70% alcohol between tests to

prevent any odour-related bias (Takao and Miyakawa 2006). The video was recorded

using a Logitech HD webcam C270 overhead-mounted camera and analysed using

online free software (Toxtrac) (Rodriguez et al. 2018), which was also blind tested

manually. The total distance covered, time spent in the light box, and the number of

transitions made between the two boxes were considered. The litter effect for animals

used in light/dark box assay was also analysed (Figure S 3A-C).

Spatial object recognition assay

All animals were housed in single cages a week before the experiment. P50 male

(N=27, n=84) C57Bl/6 mice were habituated for 2 minutes to handling and a transport

vessel for 1 week followed by spatial object recognition assay which comprised a total

of 5 sessions, each 10 minutes long. The process is similar to the classical object

recognition assay with a few variations (Wimmer et al. 2012). A light intensity of around

15 Lux was maintained at the centre of the arena (Leger et al. 2013). Session 1

involved introducing the animal to the empty arena (12 cm X 15 cm with spatial cues)

for short habituation to the open field. In sessions 2 to 4, the animal was trained to

explore two objects of different shapes kept in their fixed coordinates, with a 5 cm gap

between wall and object, through all the sessions. The animal was returned to its original cage for 5 minutes to relieve stress and the arena and objects were wiped with 70% alcohol between each session. The test session (5th session) was carried out after 24 hours of the training sessions when only one object was moved to a new coordinate with leu to the spatial cues and the animal was allowed to inspect for 10 minutes. All recordings were done using a Logitech HD webcam C270 overhead-mounted camera and a MATLAB-based autotyping software was used to calculate the number of bouts on each object (T. P. Patel et al. 2014), which was also blind tested manually. A bout was considered when the animal moved its nose within a space 2 cm radially from the object. Any attempt to climb or jump on the object was not considered a bout. The Discrimination Index (DI), which is the ratio of the difference in time spent with either object to the total time spent with both objects during the test session, was calculated to verify the ability of the animal to successfully recognize the displaced object (Denninger, Smith, and Kirby 2018).

DI= [time spent with object displaced- time spent with not displaced]/ [time spent with displaced+ time spent with not displaced]

The litter effect for animals used in spatial object recognition assay was also analysed (Figure S 3D).

Cryo-sectioning

Following behaviour assessments, animals underwent sacrifice at P60 (N=53, n=163). Mice were sedated with a heavy dose of Ketamine (60mg/kg) and Xylazine (10mg/kg) cocktail injected intraperitoneally. They were perfused intracardially with a solution of 1XPBS (phosphate buffer saline) followed by 4% paraformaldehyde, for maintaining tissue integrity. To ensure thorough fixation, brains were extracted and immersed in 4% paraformaldehyde for an additional 5 days post-perfusion. Subsequently, the fixed brains were incubated in 40% sucrose solution for a week till they settled to the bottom, enabling optimal cryo-sectioning. Brains were flash-frozen in dry ice and coronal sections of 50µ thickness were obtained and mounted on slides pre-coated with gelatin. The tissue sections were washed 3 times with autoclaved Mili-Q water and mounted with Vectasheild DAPI to stain the nucleus and visualized under the microscope.

Spine analysis

To measure neuronal development, recombinant AAV expressing two distinct shRNAs were injected into the hippocampus of male mice (P0) (N=3, n=5) and these neurons were imaged at P38-P40 (16-bit) using a Nikon A1 HD25 point scanning confocal microscope with Nikon Plan Apochromat NA=1.4 oil immersion objective at 1024X1024 pixel resolution at 0.75X. 2X optical zoom was used to get high-magnification images of neurons with 10 -15 optical sections with 0.25μm step size. The GFP signal was captured by a 488 nm solid-state laser. Images were obtained under uniform parameters of laser power; pinhole diameter and detector gain across all conditions (Figure 6A-B).

Neurolucida 360 (MBF Bioscience) was used for spine morphology analysis. A 3-D skeletal image was generated using the Directional Kernel tracing algorithm and spines were detected and classified. Spine density was obtained by using Neurolucida Explorer (Figure 6C-E).

Coronal sections of the mouse brain were visualized using a Leica LAS V4.2 fluorescence microscope and Zeiss Apotome3 (magnified image) (Figure S 1G-H). All imaging parameters were kept uniform across experiments. Images were stitched post-acquisition using Fiji image analysis software (MosaicJ plugin).

Western blot

The hippocampal tissue was obtained from C57BL/6 male P21 pups (N=8, n=25), previously subjected to injections of UC, G5C1, or G5C2 shRNAs at P0. The pups were euthanized in a CO_2 chamber at a 30-70% fill rate of CO_2 gas and death was confirmed by tail/feet pinch. The tissue was lysed in RIPA buffer (composed of 150mM NaCl, 0.1% NP-40, 0.5% sodium deoxycholate, 0.1% SDS, 50 mM Tris, pH 7.4) and subsequently centrifuged at 4°C for 10 minutes at 4000 g to remove debris. Protein quantification was performed using the Pierce BCA Protein Assay Kit (Thermofisher) with the resulting clear supernatant.

For experimental purposes, specific sample lysates were mixed with Laemmli Buffer

and heated to 95°C. An amount of 5 µg of each sample was utilized for housekeeping

gene tubulin-β1 (Tuj-1) blot, while 30 µg of each sample was employed for pan-Ago

blot. The samples were resolved by electrophoresis on a 10% polyacrylamide gel and

then transferred onto a nitrocellulose membrane. Subsequently, the membrane was

blocked with 5% BSA and incubated with primary antibodies against Tuj-1 (dilution

1:10,000, Sigma, Catalog # T8578) and pan-Ago (dilution 1:1200, MerkMillipore,

Catalog # MABE56) overnight at 4°C. Following the primary antibody incubation, the

membrane was washed 6 times, 10 minutes/wash, with 1X TBST (Tris-buffered saline

with 1% Tween 20) at RT. The blot was then exposed to secondary antibodies (dilution

1:10,000 for Tuj-1 and 1:5000 for pan-Ago, Goat Anti-Mouse, Thermofisher Scientific,

Catalog# A11030) for 3 hours. Further washing was done with 1X TBST (6 washes,

10 minutes/wash) to obtain cleaner background. Luminescence on the blot was

detected using the ECL chemiluminescence kit (Millipore). Band intensities and the

expression of pan-Ago were quantified using ImageJ software, with normalization

against tubulin-β1.

395   Statistics

Statistical analyses were performed for all experiments. Data shown are represented

as mean±SEM. The number of pregnant dams/ litters used is defined by 'N' and the

total number of pups (obtained from the dams) used is designated as 'n'. The number

of neurons considered for spine analysis is represented as '*n"*. qRT-PCR data for

lncRNA screening, Argonaute (Ago) immunoblots and spine density were analyzed

using unpaired Student's t-test with Welch's correction for unequal variance. qRT-PCR

data for Gas5 knockdown efficiency were analyzed using paired Student's t-test. Two-

way ANOVA with Fisher's LSD test was used for analyzing light/dark and SOR data

analysis. GPower 3.1 was used for post-hoc power analysis for all experiments. The

outlier test was done using the ROUT method (0.1% aggressive). Immunoblot band

intensity analysis and image stitching were done using ImageJ software. Gene

Ontology plots were computed using RStudio (Figure 5 A-C) and SRplot (Figure 7 B-

E) (Tang et al. 2023). Data with a p-value less than 0.05% were considered significant.

All data were analysed using GraphPad Prism 9. No randomisation technique was

used animals were grouped arbitrarily. No prior sample size was calculated. Post-hoc

power analysis was done for all experiments. Detailed statistical analysis and criteria

for the exclusion of animals is provided in Supplementary data and Figure S5.

RESULTS

Maternal separation alters the differential expression of synapse-enriched

lncRNAs

To investigate the impact of early life stress, this study used the maternal separation

(MS) paradigm (Figure 1A) and the differential expression of a subset of synapse-

enriched lncRNAs was analyzed by qPCR in males and females separately. In males,

we observed a significant upregulation in the expression of specific lncRNAs, such as

Pantr1 (146.3 ± 56.78% increase, p=0.041) and Gas5 (79.6 ± 19.14% increase,

p=0.006) and without any change in expression of 1700020I14Rik (8.30 ± 13.14%

increase, p=0.5522) upon MS. Although, we observed an increase in expression of

2410006H16Rik (105.5 ± 75.28% increase, p=0.210), 2900052NO1Rik (100.3 ±

70.88% increase, p=0.206), Pvt1 (82.9 ± 46.95% increase, p=0.127), RMST_1 (49.9

± 51.05% increase, p=0.366), Snhg6 (95.4 ± 81.12% increase, p=0.284), Snhg8 (80.5

± 73.63% increase, p=0.316), Snhg12 (86.3 ± 48.60% increase, p=0.126) and Zfas1

(37.4 ± 23.57% increase, p=0.163);   these enhancement in expression are not

statistically significant (Figure 1B). Gas5 was selected as the target lncRNA and two

shRNAs (G5C1 and G5C2) against Gas5 were cloned into an AAV vector (Figure

S1A). Knockdown efficacy was assessed in primary hippocampal neuronal culture and

successful knockdown was evaluated for both the constructs (82.82 ± 9.33 %

decrease, p= 0.003 for G5C1 and 84.93 ± 13.68 % decrease, p= 0.25 for G5C2 w.r.t

GAPDH; 81.3 ± 8.46 % decrease, p=0.0024 for G5C1 and 83.71 ± 6.9 % decrease,

p=0.0067 for G5C2 w.r.t Tubulin-β1) (Figure S1C-D).

Figure 1. Maternal separation affects the expression of synapse-enriched lncRNAs. (**A**). Maternal separation paradigm followed: P2 pups separated from their mother for 3 hours during P2 – P14 and following MS pups were kept till P21. (**B**). Differential expression of synapse-enriched lnc-RNAs in male hippocampus (P21) (a) 1700020I14Rik, (b) 2410006H16Rik, (c) 2900052NO1Rik, (d) Pantr1, (e) Pvt1, (f) RMST_1, (g) Snhg6, (h) Snhg8, (i) Gas5, (j) Snhg12, (k) Zfas1 as obtained through qPCR, *p < 0.05, **p < 0.01, ns = not significant. N=7, Data shown as mean ± SEM. Unpaired Student's t-test with Welch's correction.

The underlying data is also shared in https://figshare.com/articles/dataset/_b_Gas5_regulates_early_life_stress-induced_anxiety_and_spatial_memory_b_/26047969 .

____

In females, no significant alteration of Gas5 expression was seen after MS (24.77 ± 21.91, p= 0.3095). A significant downregulation of Zfas1 (47.50 ± 5.535, p= 0.0004) was observed. 2900052NO1Rik (164.9 ± 54.88, p= 0.0300), Snhg6 (101.9 ± 39.48, p= 0.0494) and Snhg8 (19.09 ± 6.565, p= 0.0335) were significantly upregulated upon MS. Other lncRNAs like 1700020I14Rik (5.670 ± 10.08, p= 0.5982), 2410006H16Rik (-28.74 ± 16.25, p= 0.1373), Pantr1 (-13.13 ± 18.85, p= 0.5172), Pvt1 (-20.40 ± 18.66, p= 0.3241), RMST_1 (219.2 ± 210.8, p= 0.3460) and Snhg12 (-5.823 ± 37.76, p= 0.8835) were not differentially regulated in the hippocampus after early life stress. The data showed the sexual dimorphic expression of lncRNAs between males and females.

Figure 2. Maternal separation affects the expression of synapse-enriched lncRNAs in hippocampus of female P21 mice. Differential expression of synapse-enriched lnc-RNAs (A). 1700020I14Rik (B)2410006H16Rik, (C) 2900052NO1Rik, (D) Pantr1, (E) Pvt1, (F) RMST_1, (G) Snhg6, (H) Snhg8, (I) Snhg12, (J) Gas5, (K) Zfas1 as obtained through qPCR, *p < 0.05, **p < 0.01, ns = not significant. N=6, Data shown as Mean ± SEM. Unpaired Student's t-test with Welch's correction.

The underlying data is also shared in https://figshare.com/articles/dataset/_b_Gas5_regulates_early_life_stress-induced_anxiety_and_spatial_memory_b_/26047969 .

Gas5 knockdown partially alleviates MS-induced anxiety

We have analysed open-space anxiety of mice that are injected with AAV expressing shRNAs against Gas5 or non-targeting control. Mice were bilaterally injected with AAV expressing either G5C1 or G5C2 in the hippocampus at P0. These mouse pups were maternally separated or homecaged (P2 – P14); following which they were kept with their respective mother till P21 (day of weaning). Young-adult male mice (P48) were subjected to the light/dark box assay (Figure 3A) (X. Wang et al. 2022). Maternally separated mice exhibited avoidance of brightly-lit open areas as detected by subsequent reduction in time spent (102.2 ± 24.35 second (s) decrease, p<0.0001); distance covered (18.26 ± 2.814 meters (m) decrease, p<0.0001), and fewer transitions (13.55 ± 3.55 decrease, p= 0.0003) into the brightly lit arena as compared to the homecaged control. However, the maternally separated Gas5 RNAi mice showed a partial rescue in anxiety level as measured by time spent (65.47 ± 27.03 s increase, p= 0.0179 for G5C1 and 66.62 ± 25.21 s increase, p<0.0101 for G5C2), distance covered (17.45 ± 3.123 m increase, p <0.0001 for G5C1 and 13.38 ± 2.912 m increase, p<0.0001 for G5C2) and number of transitions (11.42 ± 3.943 increase, p= 0.005 for G5C1 and 8.615 ± 3.677 increase, p=0.0219 for G5C2) into the brightly lit arena with respect to maternally separated mouse expressing non-targeting control shRNA (Figure 3C-E). We have also observed that homecaged mice expressing Gas5 shRNA did not show any change in time spent, distance covered and the number of transitions into the brightly lit arena as compared to homecaged mice expressing non-targeting shRNA (Figure 3C-E). These observations indicate that Gas5 RNAi does not

495   alter any physiological phenotype of the animal. Litter effect was also statistically

496   analysed and no significant difference in behaviour among the litters was found (Figure

497   S 3 A-C).

Figure 3. Gas5 knockdown reduces anxiety following maternal separation. A. Paradigm depicting light/dark box assay to measure anxiety. **B**. Representative trace showing the movement of the mouse in the bright-lit arena in the light/dark box. **C.** HC = Homecaged, MS = Maternally Separated, G5C1= shRNA # 1, G5C2 = shRNA # 2. Distance covered in the light box (in meters). **D.** Time spent in the light box (in seconds). **E.** Total number of transitions made between light and dark box. Data shown as mean±SEM, N=3-7, n=10 - 18, *p<0.05 **p< 0.01, ***p<0.005, ****p<0.0001, ns not significant. Two-way ANOVA with Fisher's LSD test. The underlying data is also shared in https://figshare.com/articles/dataset/_b_Gas5_regulates_early_life_stress-induced_anxiety_and_spatial_memory_b_/26047969 .

____

Gas5 RNAi rescues MS-induced spatial memory deficit

We have analysed the impact of early-life stress on memory using spatial object recognition (SOR) test and evaluated the role of Gas5 following its knockdown (Figure S1A). Mice were injected (P0) with AAV expressing shRNA against Gas5 or non-targeting control. These mice were maternally separated or homecaged (P2 – P14) and subjected to SOR (P56). Mice were allowed to explore two distinct objects placed in two specific locations within an arena during the training session (Figure 4A, S2). On the test day, 24 hours after training, one of the objects was displaced within the same arena and the exploration time of non-displaced and displaced objects was analysed (Figure 4A-3B). We observed that the maternally separated mice expressing non-targeting shRNA displayed reduced recognition of the displaced object during the test session as determined by the discrimination index (DI) (0.4152 ± 0.1204 decrease, p=0.0009) when compared to the homecaged control (Figure 4D). Next, we examined the effects of maternal separation on memory following the knockdown of Gas5. We observed that the maternally separated mice having Gas5 RNAi could partially enhance DI (0.5568 ± 0.116 increase, p<0.0001 for G5C1 and 0.5186 ± 0.1181 increase, p<0.0001 for G5C2) as compared to the maternally separated mice injected with non-targeting control shRNA. Also, we have detected comparable DI of homecaged mice injected with AAV expressing either Gas5 shRNAs or non-targeting control shRNA (Figure 4D). Furthermore, the exploration of objects that were not displaced within a separate arena remains unaffected following Gas5 RNAi and maternal separation as we did not observe any change in DI across experimental

534 conditions (Figure 4C). These observations indicate that the reduction of Gas5

535 expression could partially rescue deficits in spatial memory following maternal

536 separation. Litter effect was also statistically analysed and no significant difference in

537 behaviour among the litters was found (Figure S 3D).

A.

Habituation to Context 10 mins

Training (3X, 10 mins/session)

Testing after 24 hours

Not Displaced

Displaced

1 week habituation (2 mins/day)

24 hours

B.

C.   Heat map of arena with displaced object during test session

Homecaged

Maternally Separated

Control

Gas5 C1 RNAi

Gas5 C2 RNAi

C.

ns

ns

ns

ns

ns

Discrimination Index (DI)

HC   MS   HC   MS   HC   MS

Control RNAi   G5C1 RNAi   G5C2 RNAi

Not displaced in test session

D.

****

ns

ns

***

****

Discrimination Index (DI)

HC   MS   HC   MS   HC   MS

Control RNAi   G5C1 RNAi   G5C2 RNAi

Displaced in test session

Figure 4. Gas5 reverses the effect of MS-induced deficits in memory. **A.** Schematic showing paradigm for SOR. **B.** Representative heat maps in the arena with the displaced object during the test session, circular (glass) object represents the displaced object. **C.** HC= Homecaged, MS= Maternally Separated, G5C1= shRNA # 1, G5C2= shRNA # 2. Discrimination Index (DI) in the arena with no displaced object during the test session. **D.** DI in the arena with a displaced object during the test session. Data shows mean±SEM. N=4-6, n=12-15, *p< 0.05, **p< 0.01, ***p<0.005, ns = not significant. Two-way ANOVA with Fisher's LSD test.

The underlying data is also shared in https://figshare.com/articles/dataset/_b_Gas5_regulates_early_life_stress-induced_anxiety_and_spatial_memory_b_/26047969 .

____

Gas5 is potentially involved in regulating RNA metabolism

To gain a plausible mechanistic insight into the Gas5-mediated regulation of anxiety and memory upon maternal separation, we took an *in silico* approach to identify the expression of transcripts influenced by Gas5. First, we have selected Gas5 interacting coding transcripts from the available CLIP-seq data (POSTAR3 database – http://postar.ncrnalab.org./) (Hu et al. 2017). We then performed a Gene Ontology (GO) analysis of these interacting coding transcripts and focused on significant enrichment (FDR < 0.05) of transcripts in specific GO terms (Figure 5A-C). The cellular processes included enrichment of GO-terms for ribonucleoprotein complex (FRD= 1.05E-31), cytoplasmic stress granule (FDR= 1.41E-17), polysome (FDR= 3.50E-05), somatodendritic compartment (FDR= 0.010067571) and dendrite (FDR= 0.004478929) (Figure 5B) etc. The molecular functions are enriched with GO-terms representing ribonucleoprotein complex binding (FDR= 1.22E-05), mi-RNA binding (FDR= 6.05E-17) and translation regulator activity (FDR= 7.53E-05) (Figure 5C). The biological process displays significant enrichment for GO-terms, such as RNA processing (FDR= 1.58E-43), RNA splicing (FDR= 8.08E-45), regulation of translation (FDR= 2.52E-18) etc. alongside some not significant but enriched terms like dendritic transport of ribonucleoprotein complex (FDR= 0.094670904) and miRNA transport (FDR= 0.063158655) (Figure 5A) (For details see Table S1 and Supplementary data). Overall, the GO analysis revealed that Gas5 is primarily involved in RNA metabolism

including stability and regulation of translation. Prompted by this observation, we
focused on the transcript encoding Argonaute (Ago) family of proteins that are involved
in translation and RNA stability. As a proof-of-concept, we have analyzed the
expression of Ago following Gas5 knockdown. Mice were injected with AAV expressing
shRNA against Gas5 or UC (P0), and Ago expression in the hippocampus was
analyzed at P21. We observed a significant increase in Ago expression in G5C1 (0.88
± 0.25 increase, p=0.0084) and G5C2 (0.78 ± 0.29 increase, p=0.0335) injected
animals as compared to Control RNAi individuals (Figure 5 D-E).

Figure 5. **Involvement of Gas5 in RNA metabolism as revealed by GO-analysis.** Available

CLIP-seq data of Gas5 interacting coding transcripts were used to perform gene ontology

(GO) analysis showing clusters of transcripts involved in **A**. Biological processes, **B**. Cellular components, and **C**. Molecular functions. Red dots show significant enrichment (FDR< 0.05), blue dots show highly enriched but not significant **D.** Immunoblot of Pan-Ago and Tuj-1 expression at the hippocampus of P21 male animals injected with Control (UC), Gas5 C1 (G5C1) or Gas5 C2 (G5C2) shRNA. **E.** Quantification of Pan-Ago with Tuj-1 as housekeeping gene; N=2-3, n=7-10, *p<0.05; Data shows mean±SEM. Unpaired t-test with Welch's correction.

The underlying data is also shared in https://figshare.com/articles/dataset/_b_Gas5_regulates_early_life_stress-induced_anxiety_and_spatial_memory_b_/26047969 .

____

Gas5 does not regulate spine development

To analyse the importance of Gas5 in spine development, we effectively knocked down Gas5 using two shRNAs at P0 and studied the changes in spine density at P30. We studied the spine density in neurons of the CA3 region of the hippocampus. No significant difference was observed in total spine density between the Control RNAi and either of the Gas5 RNAi neurons (0.01281±0.05731, p= 0.825 for G5C1 and 0.02916±0.06329, p=0.649 for G5C2) (Figure 6C). Additionally, we observed no significant difference in the density of stubby and mushroom spines between the Control RNAi group and the Gas5 RNAi individuals (Stubby: 0.0029± 0.034, p= 0.931 for G5C1 and 0.0533±0.034, p= 0.138 for G5C2; Mushroom: 0.0232±0.0171, p= 0.1869 for G5C1 and 0.0265±0.0177, p= 0.1504 for G5C2) (Figure 6D-E).

Figure 6. **Effect of Gas5 knockdown on spine development. A, B.** High magnification images of AAV injected mouse brain sections showing EGFP expressing hippocampal neurons; Control RNAi, G5C1 RNAi and G5C2 RNAi (left to right) respectively. **B**. Synapse density in hippocampal neurons of P30 mice injected with AAV at P0; Control RNAi, G5C1 RNAi and G5C2 RNAi (left to right) respectively. Quantification of **D.** total spine density, **E.** stubby spine density and **F.** mushroom spine density per μm in Control RNAi (UC), Gas5 C1 (G5C1) or Gas5 C2 (G5C2) shRNA; N=1, n=1-2, *n'*=13-19, *p<0.05; Data shows mean±SEM. Unpaired t-test with Welch's correction.

The underlying data is also shared in https://figshare.com/articles/dataset/_b_Gas5_regulates_early_life_stress-induced_anxiety_and_spatial_memory_b_/26047969 .

DISCUSSION

LncRNAs emerged as key regulatory RNAs involved in diverse cognitive functions (Mercer, Dinger, and Mattick 2009) (Samaddar and Banerjee 2021) (Liau et al. 2021). A recent study from our laboratory identified a set of synapse-enriched lncRNAs from the hippocampus (Samaddar et al., 2023). This study highlighted the importance of one such synapse-enriched lncRNA, Gas5, which in early life stress, evoked behavioural deficits. We observed that early life stress of maternal separation induces the expression of Gas5 in the hippocampus of male mice (Figure 1B). The reduction of Gas5 expression by RNAi alleviates MS-induced anxiety and rescues memory deficits (Figure 3–4). Apart from Gas5, maternal separation also induces the expression of another synapse-enriched lncRNA, Pantr1 (Figure 1B). The hippocampi of female pups (P21) were collected similarly to the male pups. The screen revealed a very different regulation of lncRNAs in female pups upon MS revealing the sexually dimorphic outcomes due to stress. Further behavioural experiments were done in males only as female behaviour is evidenced to be influenced by the oestrous cycle (Figure 2).

The *in situ* hybridization (Allen Brain Atlas) data showed that Gas5 is expressed in the CA1, CA2, CA3 and DG of both the dorsal and the ventral hippocampus (R. S. Patel et al. 2023), whereas, Pantr1 expression is mainly restricted to the CA1 region of the dorsal hippocampus (Goff et al. 2015). Differential expression of other lncRNAs is also observed in different neuronal populations of the hippocampus (CA1, CA2, CA3 and DG). For example, GM9968 is a lncRNA that is enriched in the hippocampus. Its expression spans through CA1, CA3 and DG in both the ventral and dorsal hippocampus, with a higher expression, especially in the ventral hippocampus. Meanwhile, another hippocampus-enriched lncRNA GM13644 is equally expressed in the ventral and dorsal hippocampus but higher at the DG compared to the CA1-CA3 regions (Kadakkuzha et al., 2015). These subregions of the hippocampus have been shown to have different functions (Tsien et al., 1996, Hitti and Siegelbaum, 2014, Nakashiba et al., 2008) and can also play a unique role in regulating the behaviour of an individual (Cui et al., 2022, Grinman et al., 2019, Butler et al., 2019). Existing literature shows that the dorsal hippocampus is involved in spatial memory formation (Moser and Moser 1998) and the ventral hippocampus has a prominent role in anxiety

(Fanselow and Dong 2010). Therefore, to probe the lncRNA function in MS-induced
anxiety and memory, we have focused on Gas5.

The Gas5 lncRNA is involved in aggression (Feldker et al. 2003) and novelty-induced
behaviour (Xu et al. 2020). Consistent with these observations, our study
demonstrates the involvement of Gas5 in MS-induced anxiety (Figure 3). The
maternally separated mouse displays a significant reduction in exploring the displaced
object in our spatial object recognition test, indicating deficits in memory upon early
life stress (Figure 4). These memory deficits are partially rescued by the Gas5
knockdown (Figure 4). Our observation of impaired spatial learning following maternal
separation is in agreement with previous findings (Huot et al. 2002)(Grace et al.
2009)(Hulshof et al. 2011). However, the impaired cognitive performance with
accompanied enhancement in anxiety differs from other studies that demonstrated
improved spatial learning along with elevated levels of anxiety following maternal
separation (Suri et al. 2013) (McGrath, Records, and Rice 2008). This discrepancy
could arise due to the use of different rodent strains and paradigms to test
hippocampus-dependent memory. We speculate that heightened anxiety upon
maternal separation could contribute to hyperactivity that may lead to the exploration
of both the displaced as well as the non-displaced object in our memory test (Figure
4). Our data shows that there is no bias between home-caged and maternally
separated mice in their preference towards any object. Also, no preferential bias for
any object is observed between mice expressing Gas5 shRNA or non-targeting control
(Figure 4).

The role of direct Gas5 in hippocampus development is yet to be evaluated *in vivo,*
although previous studies have shown that Gas5 has different expression patterns in
young and old mice (Verbitsky et al., 2004). Aged mice showed a higher expression of
Gas5. This finding was later supported by the fact that Alzheimer's disease AD patients
had an elevated level of Gas5 in their hippocampus accompanied by a decrease in
hippocampal volume (Chen et al., 2022). The molecular mechanism of regulation of
Gas5 in the development of the hippocampus is still unexplored. A study done in
Mouse Embryonic Hippocampal Progenitor cells showed an elevated Gas5 level
accompanies neuronal commitment, while Gas5 downregulation occurs during
terminal differentiation, suggesting a regulatory role in conjunction with connexin
expression during neurogenesis (Rozental et al., 2000).

It is important to note that the Gas5 RNAi alone does not influence anxiety or memory

without any exposure to early life stress (Figure 3-4). This observation indicates that

the MS-induced Gas5 level above the physiological set point is a prerequisite for

anxiety and memory deficits due to early life stress. To evaluate whether the reversal

of ES-induced anxiety and memory deficits occurs due to reduced level of Gas5 in the

adult or developing brain, we have measured spine density and type from the adult

hippocampus (P38 – P40) following shRNA-mediated knockdown of the transcript

during development (P0). The spine density and type upon Gas5 knockdown were

assessed as these parameters are key determinants of anxiety and memory (Melo et

al., 2020; Leuner and Shors, 2013; Restivo et al., 2009; Kasai et al., 2010). No

significant changes in the density of the total spine, stubby spine and mushroom spine

were detected in the dendrites of hippocampal neurons (P38 - 40) following Gas5

knocked down (P0) (Figure 6). This observation confirms that the Gas5 activity during

neuronal development does not influence the reversal of MS-induced anxiety and

spatial memory deficit. The use of two distinct shRNAs against Gas5 (Figure S1) for

all experiments eliminates the possibility of an off-target effect and consequent

artefactual behavioural rescue or developmental changes. Taken together, our study

highlights an RNA-centric approach that could potentially ameliorate the debilitating

effect of early life stress.

For future investigations, employing alteration of Gas5 at distinct developmental

stages (P7, P14, P21, P35, P50) would be an ideal choice to show the temporal

resolution of Gas5 activity and its implications in reversing ES-induced elevated level

of anxiety and memory deficit during adolescence. This region-specific knockdown of

Gas5 through stereotaxic surgery, targeting specific regions such as CA1, CA2, CA3,

and DG, could provide deeper insights into its function within the dorsal and ventral

hippocampus. This can unravel the intricate role of Gas5 in modulating behaviour

through tailored assays, shedding light on its potential therapeutic implications of

Gas5. Further investigation is required to dissect a detailed mechanistic view of

lncRNA-dependent reversal of elevated anxiety and memory deficit upon ES. To

further understand the role of Gas5 in development, we can use the *in utero*-

electroporation technique to knockdown Gas5 at the embryonic stage and thereafter

study the development of the different types of the neural circuit of the hippocampus

(i.e. Perforant pathway, Mossy fibre pathway, Schaffer collateral pathway etc.). To

further understand the effect of Gas5 on functional changes at the synapse, we can study the changes in electrophysiological properties of neurons using the patch-clamp technique. This can help in elucidating the role of Gas5 in shaping the synaptic landscape of the hippocampus during crucial developmental periods.

To gain a plausible mechanistic insight into the Gas5-mediated regulation of anxiety and memory deficit upon maternal separation, we analyzed the functions of the Gas5 interacting transcripts identified from the POSTAR3 database. Our GO analysis revealed that Gas5 interacting transcripts are involved in RNA metabolism and synaptic functions. We have noted the regulation of transcripts associated with translation, RISC loading complex, miRNA transport and RNA splicing. The GO analysis also detected regulation of transcripts encoding proteins necessary for dendritic transport of ribonucleoprotein complex, somatodendritic compartment and ionotropic glutamate receptor binding. These observations impinge on the fact that Gas5 could, potentially, regulate synaptic function *via* post-transcriptional regulation of gene expression within subcellular spaces upon exposure to early life stress. Our data shows that the expression of Argonaute, a component necessary for miRISC loading and miRNA function, is reduced following the knockdown of Gas5. Consistent with previous observations showing the involvement of miRNAs in early life stress-induced gene expression (Allen and Dwivedi 2020), we speculate that Gas5-mediated regulation of anxiety and memory upon maternal separation involves components of the miRNA pathway. The enhanced expression of Argonaute may shift the miRNA loading between coding transcript and Gas5 *via* promoting molecular decoy activity to alter the translation status (Eulalio, Huntzinger, and Izaurralde 2008) (J. Li, Lv, and Che 2020). In our investigation, we identified six transcripts modulated by Gas5 (Fus, Hnrnph1, Hnrnpa2b1, Hnrnph2, Cpeb4 and Pcbp1), all pivotal in synaptic activity, using existing CLIP-seq data from the POSTAR3 database. Concurrently, other studies have focused on Gas5-mediated molecular framework regulating RNA processing operating in the infralimbic cortex for fear-memory extinction (Liau et al., 2023). The study proposes an interaction of Gas5 with Capirin1 (Cell Cycle Regulated Protein 1) and G3BP2 (Stress Granule Assembly Factor 2) at the synapse during fear extinction learning. The interaction of Gas5 with these RNA-granule proteins leads to the sequestering of Capirini and G3BP2 at the synapse. Caprin1 is evidenced to be important for the transport and translation of proteins essential for synaptic functions

(Pavinato et al., 2023) and G3BP2 helps in RNA localization and stabilization (Jin et al., 2022). Therefore, sequestering of Caprin1 and G3BP2 alters synaptic plasticity through local translation of proteins. A separate study exploring Gas5's role in addiction enlisted its targets in the nucleus accumbens (NcA), uncovering 16,053 potential interactors (Xu et al., 2020). Our comparative analysis revealed an overlap of 354 Gas5 targets between the NcA and hippocampus datasets, hinting at shared regulatory mechanisms across distinct brain regions.

We have performed a KEGG pathway and Gene Ontology analysis to determine the functional activities of these interacting partners of Gas5. The GO analysis revealed that Gas5 is predominantly involved in the regulation of RNA processing and translation of transcripts encoding proteins influencing synaptic function through the regulation of biological processes (pre and postsynaptic endocytosis, stress granule assembly, dendritic spine morphogenesis, regulation of axon extension, neuronal differentiation etc.), cellular components (polysome, growth cone, synapse, neuron projection etc.), and molecular functions (dopamine receptor binding, glutamate receptor binding, mRNA binding, GTPase activity etc.). KEGG pathway analysis indicated the involvement of Gas5 in key signalling pathways, c e.g. cAMP, mTOR and Wnt, that regulate synaptic functions in developing and adult brains (Ma et al., 2022, Grinman et al., 2021, Chen *et al.*, 2020). These pathways have been shown to directly influence anxiety and memory (Gafford et al., 2013, Zhang et al., 2011, Moore and Loprinzi, 2021, Wang et al., 2020, Argyrousi et al., 2020, Bertagna et al., 2021) (Figure 7) For details see Table S2 and Supplementary data).

Figure 7. **Gas5 targets are involved in the modulation of synapse and translation in the nucleus accumbens and the hippocampus. A**. An overlap of the number of Gas5 targeted coding transcripts regulating addiction in the nucleus accumbens (NcA) (Xu et al. 2020) and fear memory extinction in the hippocampus (Liau et al. 2023). Gas5 interacting coding transcripts were used to analyse their involvement in the **B**. KEGG pathway and Enrichment pathways in Gene Ontology analysis of **C**. Biological processes, **D**. Cellular components, and **E.** Molecular functions. FDR<0.05.

The underlying data is also shared in https://figshare.com/articles/dataset/_b_Gas5_regulates_early_life_stress-induced_anxiety_and_spatial_memory_b_/26047969 .

In our study, the knockdown of synapse-enriched transcripts of Gas5 could lead to the sequestering of RNA-binding proteins (RBPs), disrupting translation dynamics at these neuronal junctions. The Gas5 transcript's interaction with RBPs might influence how these proteins are compartmentalized within distinct synaptic subcellular spaces. This reorganization could restrict RBPs' access to other mRNAs, causing shifts in local protein synthesis at the synapse and ultimately impacting synaptic plasticity.

Gas5 immunoprecipitation, coupled with Mass spectrometry analysis from hippocampal tissue of stressed animals, holds promise for uncovering novel protein candidates and pathways. By examining the varied expression of Gas5 targets, manipulating their levels via overexpression or loss of function can offer critical insights into the potential reversibility of early-life stress impacts. Leveraging CRISPR-Cas-inspired RNA targeting systems (CIRTs), Gas5 targets can be precisely targeted at the synapse, shedding light on their role in synapse development and plasticity. Additionally, it can also be used to rescue stress-induced anxiety and loss of memory offering a scope for therapeutic intervention.

The use of CO_2 gas for euthanasia might be considered a limitation of our study as the high level of CO2 may trigger the expression of early-effect gene cFos and can also

affect other gene expressions due to a panic attack at the moment of euthanasia
(Johnson et al., 2011, Khokhlova et al., 2022). However, maternal separation data is
normalised as the ratio between homecaged control and maternally separated
animals. Hence, the probable artefact due to the euthanasia method is significantly
minimized.

Overall, our study highlights the importance of evaluating diverse lncRNA activities
during early life stress. To gain mechanistic insight, it would be important to
understand how lncRNA activity is modulated by early life stress and its consequence
on cognitive performance. We believe that this study will promote further research to
probe how early life stress across temporal windows could lead to behavioural
diversity *via* non-coding RNA-mediated regulation of gene expression.

AUTHOR CONTRIBUTION

D.B. and S.B. conceptualized the project. D.B. performed all experiments. S.S.
performed spatial memory and anxiety analysis. D.B. and S.B. wrote the manuscript.

ACKNOWLEDGEMENT

We thank Penn Vector Core for AAV vectors. We acknowledge Balakumar Srinivasan
for GO analysis, Gourav Sharma for spatial memory set-up, Smruti Mhalgi for blind
scoring of SOR data, Kamakshi Garg for assisting in confocal microscopy and Sarbani
Samaddar for conceptual discussions and proofreading of this manuscript. This work
is supported by extramural research grant BT/PR14071/GET/119/36/2015 (to S.B.)
from the Department of Biotechnology, Government of India and the core fund of the
National Brain Research Centre.

All art has been created with BioRender.com (Figure 1-4, S 1A-B, S 5 and Graphical
Abstract).

CONFLICTS OF INTEREST

The authors declare no conflict of interest.

DATA AVAILABILITY

The data supporting this study's findings are available at https://figshare.com/articles/dataset/_b_Gas5_regulates_early_life_stress-induced_anxiety_and_spatial_memory_b_/26047969. Additionally, a preprint of the manuscript is available on Biorxiv https://www.biorxiv.org/content/biorxiv/early/2023/09/27/2023.09.27.559778.full.pdf posted on 27th of September 2023.

REFERENCES

Allen, Lauren, and Yogesh Dwivedi. 2020. "MicroRNA Mediators of Early Life Stress Vulnerability to Depression and Suicidal Behavior." *Molecular Psychiatry* 25(2): 308–20. http://dx.doi.org/10.1038/s41380-019-0597-8.

Balouek, Julie-anne et al. 2023. "Reactivation of Early-Life Stress-Sensitive Neuronal Ensembles Contributes to Lifelong Stress Hypersensitivity." 43(34): 5996–6009.

Bannerman, David M. et al. 2014. "Hippocampal Synaptic Plasticity, Spatial Memory and Anxiety." *Nature Reviews Neuroscience* 15(3): 181–92.

Banqueri, María, Marta Méndez, and Jorge L. Arias. 2017. "Behavioral Effects in Adolescence and Early Adulthood in Two Length Models of Maternal Separation in Male Rats." *Behavioural Brain Research* 324: 77–86. http://dx.doi.org/10.1016/j.bbr.2017.02.006.

Barkus, Christopher et al. 2010. "Hippocampal NMDA Receptors and Anxiety: At the Interface between Cognition and Emotion." *European Journal of Pharmacology* 626(1): 49–56. http://dx.doi.org/10.1016/j.ejphar.2009.10.014.

Blum, Sonja, Anthony N. Moore, Frank Adams, and Pramod K. Dash. 1999. "A Mitogen-Activated Protein Kinase Cascade in the CA1/CA2 Subfield of the Dorsal Hippocampus Is Essential for Long-Term Spatial Memory." *Journal of Neuroscience* 19(9): 3535–44.

Bourin, Michel, and Martine Hascoët. 2003. "The Mouse Light/Dark Box Test." *European Journal of Pharmacology* 463(1–3): 55–65.

Christensen, Mette, Lars A. Larsen, Sakari Kauppinen, and Gerhard Schratt. 2010.
"Recombinant Adeno-Associated Virus-Mediated MicroRNA Delivery into the Postnatal
Mouse Brain Reveals a Role for MiR-134 in Dendritogenesis in Vivo." *Frontiers in*
Neural Circuits 3(JAN): 1–10.

Cui, Xiaoli et al. 2022. "Identification and Characterization of Long Non-Coding RNA Carip in
Modulating Spatial Learning and Memory." *Cell Reports* 38(8): 110398.
https://doi.org/10.1016/j.celrep.2022.110398.

Daniels, W. M.U., C. Y. Pietersen, M. E. Carstens, and D. J. Stein. 2004. "Maternal
Separation in Rats Leads to Anxiety-like Behavior and a Blunted ACTH Response and
Altered Neurotransmitter Levels in Response to a Subsequent Stressor." *Metabolic*
Brain Disease 19(1–2): 3–14.

Denninger, Jiyeon K., Bryon M. Smith, and Elizabeth D. Kirby. 2018. "Novel Object
Recognition and Object Location Behavioral Testing in Mice on a Budget." *Journal of*
Visualized Experiments 2018(141): 1–10.

Dimatelis, J. J., V. A. Russell, D. J. Stein, and W. M. Daniels. 2014. "Methamphetamine
Reversed Maternal Separation-Induced Decrease in Nerve Growth Factor in the Ventral
Hippocampus." *Metabolic Brain Disease* 29(2): 433–39.

Eulalio, Ana, Eric Huntzinger, and Elisa Izaurralde. 2008. "GW182 Interaction with
Argonaute Is Essential for MiRNA-Mediated Translational Repression and MRNA
Decay." *Nature Structural and Molecular Biology* 15(4): 346–53.

Fanselow, Michael S., and Hong Wei Dong. 2010. "Are the Dorsal and Ventral Hippocampus
Functionally Distinct Structures?" *Neuron* 65(1): 7–19.
http://dx.doi.org/10.1016/j.neuron.2009.11.031.

Feldker, Dorine E.M. et al. 2003. "GeneChip Analysis of Hippocampal Gene Expression
Profiles of Short- and Long-Attack-Latency Mice: Technical and Biological Implications."
Journal of Neuroscience Research 74(5): 701–16.

File, Sandra E., Paul J. Kenny, and Survjit Cheeta. 2000. "The Role of the Dorsal
Hippocampal Serotonergic and Cholinergic Systems in the Modulation of Anxiety."
Pharmacology Biochemistry and Behavior 66(1): 65–72.

Goff, Loyal A. et al. 2015. "Spatiotemporal Expression and Transcriptional Perturbations by

Long Noncoding RNAs in the Mouse Brain." *Proceedings of the National Academy of*
Sciences of the United States of America 112(22): 6855–62.

Grace, Laurian et al. 2009. "Effect of Exercise on Learning and Memory in a Rat Model of
Developmental Stress." *Metabolic Brain Disease* 24(4): 643–57.

Hu, Boqin et al. 2017. "POSTAR: A Platform for Exploring Post-Transcriptional Regulation
Coordinated by RNA-Binding Proteins." *Nucleic Acids Research* 45(D1): D104–14.

Hulshof, Henriëtte J. et al. 2011. "Maternal Separation Decreases Adult Hippocampal Cell
Proliferation and Impairs Cognitive Performance but Has Little Effect on Stress
Sensitivity and Anxiety in Adult Wistar Rats." *Behavioural Brain Research* 216(2): 552–
60.

Huot, Rebecca L., Paul M. Plotsky, Robert H. Lenox, and Robert K. McNamara. 2002.
"Neonatal Maternal Separation Reduces Hippocampal Mossy Fiber Density in Adult
Long Evans Rats." *Brain Research* 950(1–2): 52–63.

Izquierdo, Ivan, and Jorge H. Medina. 1997. "Memory Formation: The Sequence of
Biochemical Events in the Hippocampus and Its Connection to Activity in Other Brain
Structures." *Neurobiology of Learning and Memory* 68(3): 285–316.

Jimenez, Jessica C. et al. 2018. "Anxiety Cells in a Hippocampal-Hypothalamic Circuit."
Neuron 97(3): 670-683.e6.

Kaech, Stefanie, and Gary Banker. 2006. "Culturing Hippocampal Neurons." *Nature*
Protocols 1(5): 2406–15.

Kalisch, Raffael et al. 2006. "Anxiety and Hippocampus Volume in the Rat."
Neuropsychopharmacology 31(5): 925–32.

Kasai, Haruo et al. 2010. "Structural Dynamics of Dendritic Spines in Memory and
Cognition." *Trends in Neurosciences* 33(3): 121–29.
http://dx.doi.org/10.1016/j.tins.2010.01.001.

Katsouli, S. et al. 2014. "Sexually Dimorphic Long-Term Effects of an Early Life Experience
on AMPA Receptor Subunit Expression in Rat Brain." *Neuroscience* 257: 49–64.
http://dx.doi.org/10.1016/j.neuroscience.2013.10.073.

Kim, Ji Yoen et al. 2014. "Intracerebroventricular Viral Injection of the Neonatal Mouse Brain
for Persistent and Widespread Neuronal Transduction." *Journal of Visualized*
Experiments (91): 1–7.

Korosi, A. et al. 2012. "Early-Life Stress Mediated Modulation of Adult Neurogenesis and
Behavior." *Behavioural Brain Research* 227(2): 400–409.
http://dx.doi.org/10.1016/j.bbr.2011.07.037.

Kremmyda, Olympia et al. 2016. "Beyond Dizziness: Virtual Navigation, Spatial Anxiety and
Hippocampal Volumein Bilateral Vestibulopathy." *Frontiers in Human Neuroscience*
10(MAR2016): 1–12.

Leger, Marianne et al. 2013. "Object Recognition Test in Mice." *Nature Protocols* 8(12):
2531–37.

Leuner, B., and T. J. Shors. 2013. "Stress, Anxiety, and Dendritic Spines: What Are the
Connections?" *Neuroscience* 251: 108–19.
http://dx.doi.org/10.1016/j.neuroscience.2012.04.021.

Li, Hongzhu et al. 2022. "Loss of SST and PV Positive Interneurons in the Ventral
Hippocampus Results in Anxiety-like Behavior in 5xFAD Mice." *Neurobiology of Aging*
117: 165–78. https://doi.org/10.1016/j.neurobiolaging.2022.05.013.

Li, Jie, Hui Lv, and Yu Qin Che. 2020. "Long Non-Coding RNA Gas5 Potentiates the Effects
of MicroRNA-21 Downregulation in Response to Ischaemic Brain Injury." *Neuroscience*
437: 87–97. https://doi.org/10.1016/j.neuroscience.2020.01.014.

Liau, Wei Siang et al. 2023. "Fear Extinction Is Regulated by the Activity of Long Noncoding
RNAs at the Synapse." *Nature Communications* 14(1).

Liau, Wei Siang, Sarbani Samaddar, Sourav Banerjee, and Timothy W. Bredy. 2021. "On the
Functional Relevance of Spatiotemporally-Specific Patterns of Experience-Dependent
Long Noncoding RNA Expression in the Brain." *RNA Biology* 18(7): 1025–36.
https://doi.org/10.1080/15476286.2020.1868165.

Lundberg, Stina, Ingrid Nylander, and Erika Roman. 2020. "Behavioral Profiling in Early
Adolescence and Early Adulthood of Male Wistar Rats After Short and Prolonged
Maternal Separation." *Frontiers in Behavioral Neuroscience* 14(March): 1–10.

Marais, Lelanie, Suzél M. Hattingh, Dan J. Stein, and Willie M.U. Daniels. 2009. "A
Proteomic Analysis of the Ventral Hippocampus of Rats Subjected to Maternal
Separation and Escitalopram Treatment." *Metabolic Brain Disease* 24(4): 569–86.

Mattick, John S. et al. 2023. "Long Non-Coding RNAs: Definitions, Functions, Challenges
and Recommendations." *Nature Reviews Molecular Cell Biology* 24(June): 430–47.

McGrath, J. M., K. Records, and M. Rice. 2008. "基因的改变NIH Public Access." *Infant*
behavior and development 31(1): 71–80.

Mckibben, Lauren A., and Yogesh Dwivedi. 2021. "Early-Life Stress Induces Genome-Wide
Sex-Dependent MiRNA Expression and Correlation across Limbic Brain Areas in Rats."
Epigenomics 13(13): 1031–56.

McNaughton, Neil. 1997. "Cognitive Dysfunction Resulting from Hippocampal Hyperactivity -
A Possible Cause of Anxiety Disorder?" *Pharmacology Biochemistry and Behavior*
56(4): 603–11.

Melo, Silvana Regina de, Caren Tatiane de David Antoniazzi, Shakhawat Hossain, and
Bryan Kolb. 2020. "Short Predictable Stress Promotes Resistance to Anxiety Behavior
and Increases Dendritic Spines in Prefrontal Cortex and Hippocampus." *Brain*
Research 1746(July): 147020. https://doi.org/10.1016/j.brainres.2020.147020.

Mercer, Tim R., Marcel E. Dinger, and John S. Mattick. 2009. "Long Non-Coding RNAs:
Insights into Functions." *Nature Reviews Genetics*.

Monroy, Elibeth, Elizabeth Hernández-Torres, and Gonzalo Flores. 2010. "Maternal
Separation Disrupts Dendritic Morphology of Neurons in Prefrontal Cortex,
Hippocampus, and Nucleus Accumbens in Male Rat Offspring." *Journal of Chemical*
Neuroanatomy 40(2): 93–101.

Morris, Richard G.M. et al. 2006. "Memory Reconsolidation: Sensitivity of Spatial Memory to
Inhibition of Protein Synthesis in Dorsal Hippocampus during Encoding and Retrieval."
Neuron 50(3): 479–89.

Moser, May Britt, and Edvard I. Moser. 1998. "Functional Differentiation in the
Hippocampus." *Hippocampus* 8(6): 608–19.

Patel, Rekha S. et al. 2023. "Small Molecule Targeting Long Noncoding RNA GAS5
Administered Intranasally Improves Neuronal Insulin Signaling and Decreases
Neuroinflammation in an Aged Mouse Model." *Scientific Reports* 13(1): 1–18.
https://doi.org/10.1038/s41598-022-27126-6.

Patel, Tapan P. et al. 2014. "An Open-Source Toolbox for Automated Phenotyping of Mice in
Behavioral Tasks." *Frontiers in Behavioral Neuroscience* 8(OCT): 1–16.

Pickering, Chris et al. 2006. "Repeated Maternal Separation of Male Wistar Rats Alters
Glutamate Receptor Expression in the Hippocampus but Not the Prefrontal Cortex."
Brain Research 1099(1): 101–8.

Restivo, Leonardo, Gisella Vetere, Bruno Bontempi, and Martine Ammassari-Teule. 2009.
"The Formation of Recent and Remote Memory Is Associated with Time-Dependent
Formation of Dendritic Spines in the Hippocampus and Anterior Cingulate Cortex."
Journal of Neuroscience 29(25): 8206–14.

Revest, J. M. et al. 2009. "Adult Hippocampal Neurogenesis Is Involved in Anxiety-Related
Behaviors." *Molecular Psychiatry* 14(10): 959–67.

Rodriguez, Alvaro et al. 2018. "ToxTrac: A Fast and Robust Software for Tracking
Organisms." *Methods in Ecology and Evolution* 9(3): 460–64.

Samaddar Sarbani. 2023. "Transcriptomic Analysis Identifies Synapse-Enriched LncRNAs
Required for Excitatory Synapse Development and Fear Memory." *bioRxiv*: 1–50.
https://www.biorxiv.org/content/10.1101/2023.07.14.549055v2.full.pdf.

Samaddar, Sarbani, and Sourav Banerjee. 2021. "Far from the Nuclear Crowd: Cytoplasmic
LncRNA and Their Implications in Synaptic Plasticity and Memory." *Neurobiology of
Learning and Memory* 185(April): 107522. https://doi.org/10.1016/j.nlm.2021.107522.

Shi, Changhua, Ling Zhang, and Chuan Qin. 2017. "Long Non-Coding RNAs in Brain
Development, Synaptic Biology, and Alzheimer's Disease." *Brain Research Bulletin*
132(5): 160–69. http://dx.doi.org/10.1016/j.brainresbull.2017.03.010.

Sood, Ankit et al. 2018. "Early Emergence of Altered 5-HT2A Receptor-Evoked Behavior,
Neural Activation and Gene Expression Following Maternal Separation." *International
Journal of Developmental Neuroscience* 65(May 2017): 21–28.
https://doi.org/10.1016/j.ijdevneu.2017.10.005.

Soutschek, Michael, and Gerhard Schratt. 2023. "Non-Coding RNA in the Wiring and
Remodeling of Neural Circuits." *Neuron*: 1–15.
http://www.ncbi.nlm.nih.gov/pubmed/37230080.

Spadaro, Paola A. et al. 2015. "Long Noncoding RNA-Directed Epigenetic Regulation of
Gene Expression Is Associated with Anxiety-like Behavior in Mice." *Biological*
Psychiatry 78(12): 848–59. http://dx.doi.org/10.1016/j.biopsych.2015.02.004.

Sterley, Toni Lee, Fleur M. Howells, and Vivienne A. Russell. 2013. "Maternal Separation
Increases GABAA Receptor-Mediated Modulation of Norepinephrine Release in the
Hippocampus of a Rat Model of ADHD, the Spontaneously Hypertensive Rat." *Brain*
Research 1497: 23–31. http://dx.doi.org/10.1016/j.brainres.2012.12.029.

Suri, Deepika et al. 2013. "Early Stress Evokes Age-Dependent Biphasic Changes in
Hippocampal Neurogenesis, Bdnf Expression, and Cognition." *Biological Psychiatry*
73(7): 658–66. http://dx.doi.org/10.1016/j.biopsych.2012.10.023.

Takao, Keizo, and Tsuyoshi Miyakawa. 2006. "Light/Dark Transition Test for Mice." *Journal*
of Visualized Experiments (1): 1–3.

Tang, Doudou et al. 2023. "SRplot: A Free Online Platform for Data Visualization and
Graphing." *PLoS ONE* 18(11 October): 1–8.
http://dx.doi.org/10.1371/journal.pone.0294236.

Taylor, Shelley E. 2010. "Mechanisms Linking Early Life Stress to Adult Health Outcomes."
Proceedings of the National Academy of Sciences of the United States of America
107(19): 8507–12.

Wang, Andi, Junbao Wang, Ying Liu, and Yan Zhou. 2017. "Mechanisms of Long Non-
Coding RNAs in the Assembly and Plasticity of Neural Circuitry." *Frontiers in Neural*
Circuits 11(October).

Wang, Xiaobing et al. 2022. "Maternal Separation Affects Anxiety-like Behavior Beginning in
Adolescence and Continuing through Adulthood and Related to Dnmt3a Expression."
Journal of Neurophysiology 128(3): 611–18.

Wimmer, Mathieu E., Pepe J. Hernandez, Jennifer Blackwell, and Ted Abel. 2012. "Aging
Impairs Hippocampus-Dependent Long-Term Memory for Object Location in Mice."
Neurobiology of Aging 33(9): 2220–24.

http://dx.doi.org/10.1016/j.neurobiolaging.2011.07.007.

Xu, Haiyang et al. 2020. "Role of Long Noncoding RNA Gas5 in Cocaine Action." *Biological*

Psychiatry 88(10): 758–66. https://doi.org/10.1016/j.biopsych.2020.05.004.

Zhang, Xuejing, Milton H. Hamblin, and Ke Jie Yin. 2017. "The Long Noncoding RNA

Malat1: Its Physiological and Pathophysiological Functions." *RNA Biology* 14(12):

1705–14. https://doi.org/10.1080/15476286.2017.1358347.

1046