Anticipatory brain activity depends on the sensory modality of the expected stimulus: insights from single-trial analyses

Abstract

Perceptual anticipation is known to impact the reaction time of decisions. While anticipatory patterns have been identified in human brain activity, the single-trial neural signature of anticipation remains unexplored. Similarly, past studies have underlined an effect of pre-stimulus alpha-band activity on reaction times. Still, it remains unknown whether this activity is stimulus-specific or rather acts as a general indicator of readiness. This study aimed to determine whether human participants expected a visual or an auditory stimulus at the single-trial level in both cued and uncued trials. We show that pre-stimulus brain activity entails information about the expected upcoming stimulus, and that the information content can be extracted from single-trial brain activity. Behavioral analyses helped uncover the link between correct anticipation and shifts in decision strategy, and additionally validated the classification of uncued trials. This understanding of single-trial stimulus-specific neural signatures of anticipation can significantly impact cognitive neuroscience, human-computer interaction, and neuroergonomics research, enabling the development of real-time systems capable of predicting and adapting to an individual's anticipated stimuli, thereby enhancing task performance and user experience in diverse applications ranging from adaptive interfaces to clinical interventions.

Keywords: decision-making, anticipation, EEG, Riemannian Geometry, single-trial, behavior, diffusion-decision model

Novelty and Significance

With a focus on uncovering single-trial neural signatures of anticipation, this study aims to explore the specificity of pre-stimulus brain activity to expected visual or auditory stimuli in both cued and

uncued trials, i.e. whether participants are informed about the upcoming stimulus or not. Our

research seeks to contribute to the fundamental understanding of human cognition by elucidating

the informational content embedded in anticipatory brain patterns. The insights gained hold promise

in advancing fields like cognitive neuroscience and human-computer interaction, paving the way for

innovative applications in adaptive interface design, real-time systems, and personalized

interventions.

29   Introduction

Perceptual anticipation in decision-making shapes behavioral responses by allowing individuals to

prepare for forthcoming stimuli (Poulton, 1950). Correctly anticipating a stimulus typically results in

faster and more accurate responses (Petro et al., 2019; Poulton, 1950). Moreover, pre-stimulus brain

states impact judgement confidence (Samaha et al., 2017) and stimuli perceptibility (Railo et al.,

2021). Despite its critical role, exploring pre-stimulus brain states, particularly anticipation, remains

to be developed due to methodological challenges.

Traditional analyses align brain activity to specific events, posing significant hurdles when capturing

anticipatory processes that manifest at undefined time points before stimulus onset. While aligning

all trials to a consistent point in time is complex with well-identified brain activity (Barthélemy et al.,

2013), it becomes highly challenging to interpret complex single-trial patterns. Moreover, the

unconscious nature of anticipation (Koch and Preuschoff, 2007) complicates the accessibility to the

ground-truth knowledge of what was expected, as direct participant questioning might alter neural

activity (Trevena and Miller, 2002). Electroencephalography (EEG) emerges as a favored neuro-

recording technique due to its fine temporal resolution and ability to capture conscious and

unconscious cortical activities. To circumvent the challenges above, researchers have additionally

employed paradigms utilizing cues to induce specific expectations and pinpoint their onset

(Chavarriaga et al., 2012; Petro et al., 2019). However, using cues may only reflect idealized

scenarios, as real-world decisions are often uncued, presenting a gap in ecological validity.

A neuronal component of expectation, called the Contingent Negative Variation (CNV), has been extensively studied. The CNV is an event-related potential (ERP) that emerges after a cue announcing a target stimulus is displayed. Upon several presentations of this association of stimuli, the CNV appears, with temporal jitter weakening it (Walter et al., 1964). Initially identified as a neural marker of temporal expectation, the CNV was later associated with stimulus expectation and motor preparation (Chavarriaga et al., 2012; Garipelli et al., 2011, 2009; Khaliliardali et al., 2015, 2012). In Go/NoGo tasks, anticipation was successfully classified with above-chance accuracy (Garipelli et al., 2009), revealing anticipatory brain activity patterns among half of the participants (Khaliliardali et al., 2015). While these studies focused on discerning whether participants expected to take action, decoding the semantic content of anticipation at the single-trial level, particularly its dependence on the anticipated stimulus, remains an open avenue for exploration.

Besides the CNV, pre-stimulus alpha band activity $(8 - 13\,Hz)$ has attracted attention for its presumed role in sensory information gating (Jensen and Mazaheri, 2010). Linked to top-down processing (Min and Herrmann, 2007), its precise influence on decision-making remains unclear. Studies correlating alpha activity with confidence ratings, albeit not perception, in visual (Iemi et al., 2017; Samaha et al., 2017) and auditory (Wöstmann et al., 2019) tasks suggest its potential modulatory effect on higher cognitive functions. Conversely, other studies have implied plausible connections between alpha activity and sensory encoding (Lou et al., 2014) and decision formation (Barik et al., 2019). Despite these insights, whether alpha-band activity serves as a specific neural correlate of anticipation remains unknown.

This work shows the possibility of distinguishing between anticipation of visual and auditory stimuli at the single-trial level. We implemented an EEG experiment wherein participants engaged in a discrimination task between visual and auditory stimuli, cued in one experimental session, revealing the existence of anticipatory processes. The core analysis of this study involved differentiating EEG activity depending on participants' expectations (sound or image) using the CNV or the pre-stimulus

alpha activity of cued trials only. Classifiers trained on cued data were subsequently applied to

uncued trials, successfully deciphering the semantic content of anticipation even without explicit

cues. To understand which stage of decision formation anticipation modulates and assess the

classification performance on uncued trials, we related the classification performance and

predictions to behavioral modelling parameters (Ratcliff, 1978). This work marks a substantial stride

toward understanding the dynamics of anticipation and its influence on decision-making processes.

Materials and methods

Participants

81 42 participants (24 males, 18 females; 5 left-handed; aged $20 - 64$, mean: 30.43 ± 10.78) with

82 normal or corrected-to-normal hearing and vision took part in this study. The experiment was

83 approved by the Comité d'Ethique de la Recherche Paris-Saclay, under the application number 321.

84 Each participant was informed about the purpose of the study and signed informed consent forms

85 upon participation. All participants fully completed the experiment.

Experimental design

The experiment consisted of two parts, denominated "test" and "calibration" phases thereafter,

which the participants completed within the same recording session. During the test phase,

participants were presented randomly at each trial with either the sketch of a face ("face" trial) or a

sound ("sound" trial). They were instructed to respond as fast as possible to each stimulus by

pressing with their dominant hand on the right arrow of a keyboard for "face" stimuli and on the left

arrow for "sound" stimuli. Each trial started with the apparition of a red cross in the middle of the

screen, indicating a $1.5\,s$ rest period to the participant. The cross then became white to instruct

participants to start focusing on the task and avoid parasitic movements that can blur EEG signals

(blinking, jaw and head movements in particular). This baseline period lasted for $1.5 - 3\,s$, with its

duration varying randomly across trials. After that, a square visual noise clip appeared for $0.9\ s$ in the

middle of the screen, frames being updated at a rate of $10\ Hz$. While the last frame remained on the

screen until the end of the trial, the stimulus was displayed for $200\ ms$ at the end of the noise clip.

The trial terminated upon participant response or after a timeout of $2\ s$ after stimulus onset.

Trials in the calibration phase were identical to those in the test phase, with the addition that a cue

appeared for $200\ ms$ at the beginning of the noise clip, indicating which stimulus would be

displayed. A pictogram of an eye indicated a "face" trial, and an ear icon indicated a "sound" trial. To

eliminate the possibility of having created a delayed response instead of sensory anticipation, we set

the cues to be inaccurate in 20% of the cases, excluding the ten first trials. We chose this catch

probability according to the threshold at which oddball paradigms function, hence ensuring that the

cue was still reliable while introducing uncertainty and surprise among subjects (Chavarriaga et al.,

2014; Ehrlich and Cheng, 2018; Noah et al., 2020). A summary of the trial sequence for these two

phases is given in Figure 1.

Participants performed 3 test and 4 calibration blocks of 60 trials each, resulting in 180 test trials

and 240 calibration trials. 20 training trials preceded the test phase, and 60 training trials were

performed before the calibration phase. The training phase preceding the calibration phase was

longer so that participants had time to learn the impact of the cue, the correct response to each

target stimulus, and to avoid making mistakes in catch trials. Half of the participants performed the

test blocks first, while the other group of participants started with the calibration blocks to prevent

any influence of the order of the phases. Participants could take a break between each block and

could decide when to continue with the next one. The experiment was performed in a dark room to

enhance the sight of the visual stimulus.

Figure 1

Summary schematic of the sequence of a trial. Each trial starts with a $1.5\ s$ rest period, followed by a $1.5 - 3\ s$

baseline period, during which participants are instructed to limit movement and focus on the screen. The pre-

stimulus period then lasts for $0.9\ s$, a period during which visual noise (refresh rate of $10\ Hz$) is presented. In

the "cued" condition, the cue is presented during the first $200\ ms$ of this period. Then the stimulus, either an

image of a face or a sound, is presented for $200\ ms$ in addition to a single frame of visual noise which lasts until

the end of the trial. Participants are instructed to report the perceived stimulus as fast and accurately as

possible by pressing on the left arrow for a sound and the right arrow for a face stimulus. The trial is terminated

upon participant's response or after a $2\ s$ timeout.

Stimuli

129 Target stimuli consisted of a sound pulse at $1000\ Hz$ for the sound stimulus, and a sketch of a face

130 for the visual stimulus. The sketches were generated by Yang et al. (2020) from the Radboud Faces

131 Database (Langner et al., 2010). Cue stimuli consisted of an open-source drawing of an eye and one

132 of an ear, displayed at the center of a screen over a random dot square cloud. All stimuli lasted

133 for $200\ ms$.

EEG data acquisition and pre-processing

EEG signals were recorded at $1000\ Hz$ using 32 active AgCl electrodes, and the actiCHamp Plus

amplifier (Brain Products GmbH, Gilching, Germany). The electrodes were placed according to the

10/20 international system. Electrode Fz served as the reference electrode upon acquisition.

To avoid contamination by stimulus-related activity, we applied offline a non-causal finite impulse

response band-pass filter with cut-off frequencies $0.1 - 35\ Hz$ to remove high-frequency artifacts

(muscle activity in particular) and low-frequency artifacts, such as sweat. Data was subsequently

epoched on the relevant periods of interest (see below).

EEG data comparisons: periods of interest

We wanted to see if we could distinguish the ERP patterns between anticipation versus baseline,

anticipation and no anticipation (i.e., pre-stimulus period with and without a cue), and visual versus

auditory anticipation. The pre-stimulus period was defined as the window $[-400{:}\ 0]\ ms$ preceding

the target stimulus for both stimuli and in both cued and non-cued conditions. It can equivalently be

defined as the $[500{:}\ 900]\ ms$ time window following the onset of the noise clip and cue. 2

comparisons were performed: anticipation vs. no anticipation (defined as cued vs. uncued trials), and

visual vs. auditory anticipation.

Computing ERP components

151   For the initial investigation of significant differences between the different periods of interest

152   described above, ERP components were computed by averaging trials within each category (cued

153   trials, uncued trials, face anticipation, and sound anticipation). A comparison of the grand average

154   was performed, averaging the signal both across trials and channels within participants. Note that

155   only the $0.1 - 35\ Hz$ bandpass filter was applied for this analysis.

Classification pipelines

Classifying ERPs using covariance matrices

The CNV is a candidate for discriminating the type of anticipation at the single-trial level. To use it as a feature for classification, we first applied a $1 - 4\,Hz$ band-pass filter on raw calibration data and epoched it from 500 to $900\,ms$ after noise clip onset, corresponding to the last $400\,ms$ before stimulus onset. we used it as a feature to classify single trials according to the type of anticipation (face vs. sound anticipation). Trials were then split into 10 stratified folds, using cue labels for stratification to ensure that the label distribution is preserved in each fold. For each cross-validation round, 9 of these folds were used as the train set and the remaining one as the test set, so that each fold was used once as the test set. Using the PyRiemann Python toolbox (Barachant et al., 2023; Congedo et al., 2017), extended covariance matrices (Barachant and Congedo, 2014) with XDawn components (Rivet et al., 2009) were computed for each class using the train set, selecting 8 components, and the Minimum Distance to Mean was used to assess the class of each trial. Using covariance matrices for EEG activity classification was proven efficient in improving classification accuracy on ERP data (Barachant and Congedo, 2014) and motor imagery (Barachant et al., 2013, 2012). The classification accuracy was then computed for each test set and then averaged across test sets within participant. As pre-stimulus alpha-band activity has also been shown to be predictive of behavior in other works (Barik et al., 2019; Bode et al., 2012; Iemi et al., 2017; Lou et al., 2014; Min et al., 2008; Min and Herrmann, 2007; Petro et al., 2019; Railo et al., 2021; Samaha et al., 2017; van Dijk et al., 2008; Wöstmann et al., 2019), we repeated this analysis with a $[8 - 13]\,Hz$ band-pass filter.

Classifying pre-stimulus activity of uncued trials

The afore-described classification pipelines are based on supervised learning, i.e. they require the prior knowledge of a ground-truth. In this study, it requires a prior knowledge of the type (visual or auditory) of anticipation. However, the ground-truth label of anticipation is unknown on trials of the test phase of the experiment, i.e. on uncued trials. To circumvent this issue, two solutions are

proposed here. First, if we assume the stationarity of the anticipation phenomenon across tasks, the

brain activity patterns of anticipation over uncued trials should be similar to those found on the cued

trials. Therefore, classifiers trained on cued data can be applied on uncued data of the same

participant. We hence re-trained the classifiers obtained on all the trials of the calibration phase of

the experiment for each participant, and predicted the corresponding anticipation class using the

thus trained classifier on uncued trials.

The second solution consists of analyzing the post-stimulus effects of anticipation. Indeed,

challenged expectations should result in potential variations in the post-stimulus period (Kappenman

and Luck, 2011; Lou et al., 2014). Consequently, detecting mismatch signals in the post-stimulus ERP

should indicate the type of anticipation produced by the participant in the pre-stimulus period,

hence providing "ground-truth" labels on the type of anticipation on uncued trials. We therefore

trained a classifier to detect incorrect anticipations in the post-stimulus period of the cued trials,

using the same pipeline as described in "Classifying ERPs using covariance matrices". The ROC curves

were plotted for each participant to verify the correctness of the classification, then the classifiers

were trained again over all cued trials and were subsequently used to detect incorrect anticipations

on uncued trials. The anticipation labels were then obtained by combining the stimulus labels and

the labels obtained from this pipeline. These labels were then used as the cue labels in the pipelines

described earlier.

The different pipelines used to generate the classes of anticipation on both cued and uncued trials

are summarized Figure 2. In the following, we will refer to the anticipation on cued trials as "ground"

anticipation when the cue label is used as the anticipation class, or as "classifier" when the

anticipation class is the one predicted by the pre-stimulus classifier. On uncued trials, we will refer to

"cued-pre-stimulus" anticipation when the anticipation is the one predicted by the aforementioned

classifier trained on cued trials, or to "post-stimulus" when this anticipation is the one predicted by

the pre-stimulus classifier trained using the labels predicted by the post-stimulus classifier trained on

cued data and subsequently applied to uncued trials.

Figure 2: Processing pipeline

Different anticipation classification methods, on cued and uncued trials. We use two types of classifiers,

symbolized by grey ellipsoids: a classifier trained to detect whether the anticipation is visual or auditory based

on pre-stimulus EEG activity, and a classifier trained to detect whether the anticipation was correct or incorrect

(i.e., whether anticipation matched the following stimulus or not) based on post-stimulus EEG activity. The final

labels that were used as the anticipation class are frames in light grey, with their denomination in the article

underlined in the vicinity.

Linking brain and behavior

One of the main assumptions of the effects of anticipation on behavior is the reduction of response

times and error rates subsequent to correct anticipations. If the classification algorithm was indeed

trained to distinguish the type of anticipation, we should observe that correct anticipations are more

represented among shorter reaction times relative to long ones. Conversely, incorrect anticipations

should result in longer reaction times.

Response time quantiles

We first split the predicted anticipations depending on their correctness and behavioral outcome:

the anticipation can either match the stimulus or not, corresponding to correct and incorrect

anticipations respectively, and the response can match the stimulus or not, corresponding to correct

and incorrect responses respectively. In addition, the trials were split into three groups depending on

their response time: for each participant, we computed the 33^{th} and 66^{th} percentiles of response

times and hence split the trials into short, medium, and long response times.

We then studied the distribution of each of the four pairs of anticipation+response outcomes across

response time quantiles. This was done separately for cued and uncued trials in order to compare

the results on uncued trials to the more reliable ones over cued trials.

DDM fitting

To analyze the effects of anticipation on behavior further, we fitted diffusion decision models (DDMs)

(Ratcliff, 1978; Ratcliff and McKoon, 2008; Ratcliff and Tuerlinckx, 2002) to behavioral data.

According to DDMs, sensory evidence is accumulated linearly from a starting point z_r until reaching a

fixed decision boundary a, at which time a decision is made.

The DDM is defined by the equation:

$$dx = v\, dt \, + N(t), \tag{2}$$

where the decision variable x varies by dx in infinitesimal time dt. The decision state can be viewed

as a particle subject to Brownian motion with drift v, Gaussian white noise term $N(t)$. Additionally, a

non-decision time T_0 is fitted to account for the biological delay of sensory encoding and motor

preparation explaining the difference between the decision time and the observed response time.

Our DDM analysis aimed at identifying whether changes in anticipation resulted in changes in the

accumulation rate v, or rather in a bias in the starting point z_r. More specifically, correct anticipation

resulting in faster and more accurate decisions than incorrect anticipation, one could expect that

correct anticipation either increases the rate of evidence accumulation, or that the decision is initially

biased towards the correct decision.

On cued data, two hypotheses were tested:

-   Correct anticipations should result in larger drifts or larger starting points than incorrect

anticipations

-   Participants who displayed above-chance classification performance should also display a

larger parameter difference between correct and incorrect anticipations

To test these hypotheses, we fitted a DDM for each participant, fixing the boundary $a = 1$ and non-

decision time $T_0 = 0.3\,s$ for all participants. This was necessary in order to compare model

parameters across participants, since these parameters are interdependent with the drift and the

starting point. For each participant, both the drift term and the starting point were fitted separately

depending on whether ground-truth anticipation was correct or incorrect. Starting point, drift, and

non-decision time variability were additionally fitted for each subject, but not further analysed. The

software fast-dm (Voss and Voss, 2007) was used, using the Kolmogorov-Smirnov fitting method.

To test hypothesis 1, we performed paired-sample t-tests to assess whether the drifts and starting

points significantly depended on the correctness of anticipation.

To test hypothesis 2, we followed two complementary approaches. First, we performed independent

sample t-tests to assess whether the parameter difference was significantly greater on participants

displaying above-chance classification accuracies. Second, we performed a correlation analysis to

assess a possible linear dependency of classification accuracy on either of the parameter differences.

The same DDM fitting was performed on uncued data, using this time the anticipation class predicted

by either classifier (i.e., using "cued-pre-stimulus"- and "post-stimulus"-based labels) to assess the

correctness of anticipation. This time, we assessed whether one method provided a greater

parameter difference than the other. Based on the DDM analysis of cued trials (see Results), this

measure would indicate a better classification performance from one classifier or the other.

Statistical analyses

Behavioral effects

We first tested the effects of the presence of the cue and the stimulus on the response time and

accuracy at the group level using a repeated measures ANOVA procedure, with "Stimulus type"

(auditory, visual) and "Condition" (cued, uncued) as within-subject factors. We further tested the

effect of congruent and incongruent cues and stimuli on the response time and accuracy on the cued

condition only, using repeated measures ANOVA again, with "Stimulus type" and "Cue congruency"

as within-subject factors. Note that the tests were done separately in each condition due to the

unbalanced number of trials.

Significance of ERP differences

To test the group-level effects of anticipation on time series, we performed cluster-based

permutation testing. The idea of this type of testing is explained in (Maris and Oostenveld, 2007), and

the methodology is summarized here. This analysis aims to test the statistical significance of effects

appearing in ERPs, all while correcting for multiple comparisons. Since the number of tests to

perform is very big (901 time points, or pixels), classical corrections such as the Bonferroni correction

are not adapted as they would result in meaningless reference thresholds.

F-statistics of each point are computed to identify clusters of (uncorrected) significant activity. The

pre-cluster threshold is set to $p < 0.05$. The signals are then shuffled (i.e., in the case of the

comparison of cued and uncued signals, the attribution of each signal to the "cued" or "uncued"

category is randomized. Similarly, the assignment to the "auditory" and "visual" categories is

randomized at this stage when comparing auditory and visual anticipation) and the statistics are

computed again on the permutation. The statistics of the bigger cluster thus obtained are stored, and

the whole process is repeated over several iterations to form the null hypothesis distribution, against

which all the clusters identified in the original signals are tested.

We performed 1000 permutations. This number allowed us to obtain stable null hypothesis

distributions while remaining time-efficient.

298   Significance of classification performance

We compute a mean classification score from the k-fold cross-validation for each participant and

pipeline. The classification scores are compared to chance level that we obtained through

permutation testing over 100 iterations. At each iteration, the labels are shuffled, and the classifier is

re-run to compute the classification accuracy over the randomly labeled data. The null-hypothesis

distribution of the chance level classification performance is thus obtained for each participant, and

the classification performance is compared against this distribution. The p-value of the classification

accuracy against the null-hypothesis distribution is computed for each participant and further

compared across all participants by performing one-sided paired samples $t-$ tests of the accuracy

against the individual chance level 95% confidence interval. The accuracy indeed needs to lie above

the individual chance level to consider that the classification succeeded.

Results

We have analyzed EEG recordings from 42 participants who performed a sensory categorization task. At each trial, participants had to decide whether the stimulus they were randomly presented with was a sound or a visual stimulus consisting of a drawing of a face. The trials were preceded by a pre-stimulus period of 0.9 seconds, and on some trials a cue was presented at the beginning of this pre-stimulus period, indicating with 80% confidence the class of the upcoming stimulus. In that case, the pre-stimulus period was called the *anticipation* period.

Our analyses aimed to characterize the brain activation patterns related to specific stimulus anticipation at the group level and at the single-trial level.

Behavioral results

The descriptive statistics of response times and accuracies for each condition and stimulus type are summarized on Figure 3 (A and B). We first tested the effects of the stimulus and the condition (i.e. presence or not of an informative cue) on the response times and accuracy at the group level using ANOVA. We observe a significant effect of the stimulus ($RT_{auditory} = 526ms$, $RT_{visual} = 462ms$, $F_{\{1,41\}} = 119.783$, $p < 0.001$) and the condition ($RT_{cued} = 470ms$, $RT_{uncued} = 526ms$, $F_{\{1,41\}} = 70.304$, $p < 0.001$) on the response time. We also note an interaction effect between stimulus and condition ($F_{\{1,41\}} = 30.024$, $p < 0.001$), with responses of cued visual trials faster than any other trial type ($p < 0.001$), and uncued visual trials significantly faster than uncued auditory trials ($t = 12.236, p < 0.001$), but not than cued auditory trials ($t = 1.058, p = 0.293$). However, the accuracy was impacted by neither the stimulus type nor the presence of the cue (see Figure 3B). Note that there is no interaction effect between the two factors on the accuracy. These results show that the presence of the cue effectively reduced the response times (see also Figure 3A).

Then, we tested the effect of a congruent or incongruent cue and the stimulus on the response times

and mean accuracy (the descriptive statistics can be found in Figure 3C and 3D) within the cued

condition using ANOVA. We observed that incongruent trials had a significant effect on both the

response time ($RT_{congruent} = 452ms$, $RT_{incongruent} = 558ms$, $F_{\{1,41\}} = 131.890$, $p < 0.001$) and

the accuracy ($Acc_{congruent} = 99.1\%$, $Acc_{incongruent} = 91.1\%$, $F_{\{1,41\}} = 31.140$, $p < 0.001$), and

observe as previously that the stimulus type had an effect on the response times ($RT_{auditory} =$

$496ms$, $RT_{visual} = 445ms$, $F_{\{1,41\}} = 67.009$, $p < 0.001$, see also Figure 3C) but not on the

accuracy (Figure 3D). We observed no interaction effect on either the response times or the

accuracy. These results show that the cue was considered informative by the participants.

Figure 3

Mean response times in ms (A, C) and accuracy (B, D) for all participants for each stimulus type, depending on

(A, B) the cued/uncued condition and (C, D) whether the cue was congruent or not in cued trials.

Together, these results support the emergence of anticipatory effects after the presentation of an

informative but partially unreliable cue, which validates the experimental paradigm implemented

here.

Anticipatory activity is distinct from temporal expectation

We first wanted to show that the anticipation activity is distinct from temporal expectation of the

coming event. Therefore, we compared at the group level the brain activity in the anticipation period

to the activity in the pre-stimulus period of uncued trials.

Grand average ERPs (Figure 4A) show a significant difference in two time periods: $[-718: -546]ms$

pre-stimulus ($p = 0.017$) and $[-268: 0]ms$ pre-stimulus ($p = 0.023$).

The first one corresponds to the visual-evoked potential due to the presentation of the cue and is not

relevant for our next analyses because it is not related to anticipation but rather to the sensory

processing of information. The second one, however, is quite similar to the Contingent Negative

Variation (CNV) that appears during temporal expectations. Since this signal is significantly different

from the uncued one, where little perceptual anticipation can be made, and temporal expectation

should prevail, it indicates that the content of anticipation shapes the related brain activity.

Figure 4A also shows the areas involved in anticipation by representing the difference of evoked

activity in the pre-stimulus period in the calibration versus test phase, i.e. with and without an

informative cue. We observe a general ramping of EEG activity, and a more positive difference in the

central areas. These observations are consistent with the definition of the CNV. Later activity is

dominated by occipital activation, again due to the presence of SSVEPs, which, in addition to the

global ramping of activity, bestride activity in other channels.

Figure 4

Grand average ERP traces (average across all trials, participants, and channels) with 95% bootstrapped

confidence interval, and topoplots of the difference of activity at 400, 267, 133 and 0 ms pre-stimulus. (A) cued

vs. uncued trials. The topoplots represent cued activity minus uncued activity. (B) cued trials are split on

whether the cue was an eye ("visual" anticipation) or an ear ("auditory" anticipation). The topoplots represent

auditory minus visual anticipatory activity.

Discriminating anticipation-specific neural signatures

We first tried to test whether the pre-stimulus ERP could be distinguished at the group level between

visual and auditory anticipation, that is, whether the pre-stimulus ERP differed at the group level

when an eye and an ear cue were presented. As shown in Figure 4B, we observed no significant

difference between the two ERPs.

Qualitatively, however, we note a difference between the two grand averages, which indicates that

some participants may display stronger differences between the two types of anticipation. The

topographies also present some differences (Figures 4B). We note a difference over the central

areas, suggesting a difference between the CNVs and indicating that the CNV holds information

about the type of expectation, but also a difference in the parietal-occipital electrodes. This

difference might be due to a pre-activation of sensory areas in preparation of the upcoming stimulus.

Single-trial decoding of anticipation

Decoding anticipation from the CNV is feasible on some participants on cued trials

The core test of this study was to determine whether single-trial anticipation could be classified. In

the context of our experiment, we hypothesized that participants would display differences in their

CNV component depending on whether they expected a sound or a visual stimulus. To this aim, we

filtered the signal to keep frequencies between 1 and 4 Hz, hence keeping only the lower frequency

signals that characterize the CNV. As a first step, we only considered cued trials, as they allow for

direct computation of the classification accuracy.

The table of classification performances is shown in Table 1. Out of the 42 participants, 25 displayed

classification accuracies, as computed from a 10-fold cross-validation procedure, above their

individual empirical chance level. At the group level, the classification accuracy is significantly above

chance level (mean $\pm$ standard error of the mean: $57.6 \pm 0.7\%$ accuracy, $56 \pm 0.1\%$ chance 95%

threshold; one-sided paired-sample $t-$test between classification accuracy and individual empirical

chance levels: $t(41) = 2.188, p = 0.017$, Cohen's $d = 0.338$).

As the CNV appears after several presentations of the cue-stimulus association, we assess the effect

of the condition order on the classification performance. We observed no significant difference

between the two groups in the classification accuracy (Welch t-test, $p = 0.341, df = 40$). This result

was expected since all the participants received 60 training trials prior to the cued condition.

Table 1

Classification accuracy obtained from the covariance matrices of the $1-4Hz$-filtered pre-stimulus activity of

cued trials ("accuracy"), compared to the empirical chance level ("chance (95%)"). The p-value corresponds to

the position of the classification accuracy relative to the empirical distribution of chance level, obtained by

permutation.

Subject	accuracy	chance (95%)	p-value
pilot3	0.612500	0.558542	0.009901
136	0.595833	0.558542	0.009901
358	0.554167	0.554375	0.099010
916	0.616667	0.554167	0.009901
347	0.629167	0.541875	0.009901
766	0.591667	0.554167	0.009901
661	0.520833	0.562500	0.257426
959	0.562500	0.562708	0.079208
205	0.604167	0.558542	0.009901
400	0.666667	0.546250	0.009901
756	0.525000	0.554167	0.207921
897	0.516667	0.554583	0.376238
420	0.479167	0.562708	0.663366
804	0.608333	0.566875	0.009901
207	0.491667	0.558542	0.574257
196	0.608333	0.566667	0.019802
295	0.604167	0.575000	0.019802
402	0.554167	0.571042	0.118812
722	0.558333	0.566875	0.148515
720	0.529167	0.567083	0.217822
966	0.504167	0.554375	0.475248
568	0.562500	0.558542	0.059406
482	0.470833	0.558542	0.742574
307	0.566667	0.558542	0.049505
730	0.612500	0.558333	0.019802
228	0.595833	0.566667	0.019802
785	0.583333	0.550000	0.009901
640	0.650000	0.558542	0.009901
457	0.558333	0.579375	0.118812
183	0.575000	0.566875	0.049505
759	0.625000	0.558542	0.009901
278	0.562500	0.546458	0.049505
732	0.608333	0.571250	0.009901
441	0.600000	0.571042	0.019802
930	0.562500	0.562500	0.069307
755	0.625000	0.558542	0.009901
689	0.654167	0.562917	0.009901
877	0.529167	0.550000	0.188119
312	0.608333	0.575208	0.009901
751	0.587500	0.558750	0.029703
443	0.529167	0.554375	0.158416
223	0.612500	0.558958	0.019802

Alpha-band activity for estimating the single-trial anticipation class

Following the results of past studies, we also assessed whether the alpha-band power spectral density could characterize anticipation at the single-trial level. Using the same classification pipeline as for the CNV classification, this time filtering the raw signal between 8 and 13 Hz, we found that 10 participants showed above-chance classification accuracies, representing again a smaller proportion of all participants compared to the classification of ERPs. Moreover, the group-level accuracy lies within chance-level (mean $\pm$ standard error of the mean: $54 \pm 0.8\%$ accuracy, $56.3 \pm 0.1\%$ chance 95% threshold, one-sided paired-samples $t-$ test: $t(41) = -2.741, p = 0.995$, Cohen's $d = 0.423$).

Given the better results obtained using ERP classification, the analyses performed on uncued trials are using the pipeline described in the ERP section.

Towards decoding single-trial anticipation on uncued trials

On uncued trials, we do not have the ground truth about what the participant expected. Therefore, a behavioral proxy for anticipation is necessary to check the validity of our classification.

From the behavioral analyses, we found that incorrect anticipation led to longer response times, as seen in the effect of catch trials on the response times. Therefore, we assume in the following that incorrect anticipations lead to longer correct responses, while correct anticipations lead to shorter reaction times on correct responses.

Uncued trials were classified for each participant using two pipelines. We first describe how post-stimulus activity was used to infer a ground truth for the state of anticipation. We additionally applied the classifier trained on the covariance matrices of ERPs of cued trials.

Using post-stimulus brain activity to decipher anticipation

Post-stimulus activity holds information about the prior expectations about the stimulus. We

therefore trained a classifier to detect when the anticipation did not match the stimulus, using the

same pipeline as previously.

The classification performance on cued trials was significantly above chance level (mean AUC across

participants: 0.79 ± 0.12), with trials of one participant classified under chance level.

Given the satisfactory results of the classification on cued trials, the classifiers were trained again on

all the cued trials for each participant and applied to uncued data. The labels thus obtained were

converted to anticipation labels, themselves used as the ground-truth labels of anticipation for

classification. Applying the same pipeline as described earlier on CNV classification, we computed a

classification score for each participant and then combined the p-values across participants using the

Fisher method. The classifiers yielded above-chance classification accuracy for 5 participants, which

was not significant at the group level (one-sided paired-sample $t-$test, $t(41) = 7.064, p = 1$).

Combining the $p-$values indicated that an effect may have nevertheless been captured by the

classifiers (Fisher method, $\chi^2 = 121.1, p = 0.005$).

We expected a predicted correct anticipation rate close to 50% for all the participants. However, we

observed that participants anticipate the trials correctly $31 \pm 5\%$ (mean±standard deviation) of the

time, which is significantly different from 50% (one-sample t-test, $t(41) = 27.6, p < 0.001$).

In the following, we will refer to this pre-stimulus classification pipeline as "post-stimulus"-based.

Using the CNV classifier trained on cued trials

While further qualitative assessment of the classification performance is given later in the "Brain and

behavior" results section, the rate of predicted-correct anticipations can quantitatively indicate the

quality of the fit, as we expect again that around 50% of the trials are correctly anticipated. We

observed that $53 \pm 4\%$ of the trials are correctly anticipated by each participant on average,

according to the classifiers. This value also significantly differed from 50% (one-sample t-test,

$t(41) = 4.746, p < 0.001$). Note that in that case, individual chance levels could not be computed

because we used pre-trained models, and have no ground for comparison to an individual chance

level.

In the following, we will refer to these classification results as "cued-pre-stimulus"-based.

We observe in that the classification fails at the group level with both algorithms. This could indicate

either a failure of the post-stimulus classification or a failure of this pre-stimulus classification,

although it is not possible to conclude on that based on the results presented above. The next

analysis, whose results are presented thereafter, compared the classification predictions to

behavioral observations to substantiate these results.

Relation to behavior

Behavioral predictions

Behavior in decision-making tasks is characterized by the mean response time of correct and

incorrect responses and the response accuracy. The previous analyses yielded four different

anticipation determination method: on cued data, we either used the cue as the class of anticipation

(the "ground" method) or the label predicted by the classifier (the "classifier" method). On uncued

data, we either trained the classifiers on pre-stimulus uncued data using labels predicted on post-

stimulus activity (the "post-stimulus" method) or used the classifiers created during the "classifier"

method trained on cued data to generate the anticipation labels of the uncued trials (the "cued-pre-

stimulus" method). For each anticipation determination method, we therefore computed these

quantities for each participant separately. The mean and standard deviation of these values across

participants are presented in Table 2.

	Cued		Uncued	
	Ground	Classifier	Post-stimulus	Cued-pre-stimulus
Correct anticipation				
RT correct	450 ± 76	463 ± 76	480 ± 76	526 ± 77
RT incorrect	750 ± 490	596 ± 397	569 ± 422	542 ± 219
Response accuracy	99.1 ± 1.0	97.8 ± 1.7	98.1 ± 2.1	97.8 ± 1.8
Incorrect anticipation				
RT correct	564 ± 98	474 ± 82	544 ± 81	523 ± 85
RT incorrect	522 ± 310	576 ± 398	556 ± 161	563 ± 232
Response accuracy	91.1 ± 9.1	97.6 ± 2.6	97.5 ± 1.8	97.5 ± 2.2

We additionally performed repeated measures ANOVAs on the response times and response

accuracy, taking Response (correct or incorrect), Anticipation (correct or incorrect) and Anticipation

determination method (Ground, Classifier, Post-stimulus or Cued-pre-stimulus) as within-subject

factors. On response times, we observed a significant impact of anticipation ($F(1,41) = 10.510$,

$p = 0.01$), as well as an interaction effect of anticipation and response ($F(1,41) = 8.012, p = 0.02$).

Post-hoc analyses revealed that on correct responses, incorrect anticipations yielded significantly

longer response times than correct anticipations ($t(41) = 4.302, p_{Holm} = 0.003$). On response

accuracy, we observed an effect of both the anticipation ($F(1,41) = 30.329, p < 0.001$) and the

anticipation determination method ($F(3,123) = 17.692, \ p < 0.001$), as well as an interaction effect

of those two factors ($F(3,123) = 26.045, \ p < 0.001$). Post-hoc analyses revealed that only

incorrectly-anticipated cued trials, as determined by the Ground method, were responded to less

accurately than any other trial type ($p_S < 0.001$).

Distribution of anticipation and behavioral outcomes in response time quantiles

For each participant, the trials were split into three groups depending on their tercile of response

times. These terciles were computed individually, so that each participant had the same number of

short, medium, and long response times. We then also assessed whether trials were anticipated

correctly (i.e., if the anticipation and stimulus labels matched) and if they were responded to

correctly (i.e., if the stimulus and response labels matched), yielding four pairs of possible

anticipation-behavior outcomes.

Figure 5 represents the distribution of anticipation-behavior outcomes in each response time tercile

when the anticipation label is drawn directly from the cue label ("ground" method) (Figure 5A), on

cued trials when the anticipation is predicted by the classifier (Figure 5B), on uncued trials using the

post-stimulus method (Figure 5C), and on uncued trials using the cued-pre-stimulus method (Figure

5D).

On cued data (Figure 5A), we observed that, among correct responses, correct anticipations were

prominently represented among shorter reaction times relative to longer ones, while on the contrary

incorrect anticipations were denser on long reaction times. On the other hand, error responses

tended to be more represented among longer reaction times relative to shorter reaction times

regardless of whether the anticipation was correct or not. Only the proportion of incorrectly-

anticipated correctly-responded trials was significantly higher in long response times than in short

($t = 12.816, p < 0.001$) and medium ($t = 10.125, p < 0.001$) ones, and correctly-anticipated

incorrectly-responded trials were significantly more represented in long relative to short response

time trials ($t = 5.227, p < 0.001$). These global trends were also found using the classifier method

instead of the ground method on cued trials (Figure 5B). However, these differences were not

significant at the group level.

On uncued data, the two sources of anticipation labels yielded qualitatively different results. While

anticipation predicted by the post-stimulus method raised qualitatively similar results to those

obtained on cued data, the trends are reversed when using anticipation predicted by the cued-pre-

stimulus method. In particular, correct anticipations with correct responses qualitatively display

longer reaction times while incorrect anticipations with correct responses tend to be shorter. Note

that the post-stimulus-based classifier yielded that correctly-anticipated correctly-responded trials

were significantly more represented in short than in long response times ($t = 6.310, p < 0.001$). The

same qualitative difference can be observed on the ground-truth cued trials (Figure 5A), although not

significant.

These observations lead us to the interpretation that the post-stimulus method may have performed

better than the cued-pre-stimulus method.

An interesting difference that emerges between cued and uncued trials lies in the distribution of

correctly-anticipated error trials. These tend to have longer reaction times in the cued condition

while they are more represented among the shorter reaction times in the uncued condition (Post-

stimulus-based classifier (Figure 5C): short-medium: $t = 5.608, p < 0.001$, short-long: $t =$

$5.720, p < 0.001$; cued-pre-stimulus-based (Figure 5D): short-medium: $t = 4.356, p = 0.014$). While

it is possible that the classification algorithms failed on uncued trials, we propose the alternative

view that error trials on correct anticipations result from different mechanisms depending on the

condition: in the cued condition, these errors emerge when there is doubt either on the cue that has

been seen or the motor command corresponding to the matching response. In contrast, as no

indication about the upcoming stimulus is provided to the participant in uncued trials, shorter

reaction times and errors could emerge in a repetitive task when motivation drops.

Figure 5

Distribution of trials within each anticipation-behavior outcome type across response time terciles, for cued

(A,B) and uncued (C, D) trials. In each plot, the anticipation label was extracted differently. (A) anticipation

labels correspond to cue labels ("ground" method). (B) anticipation labels are the ones predicted by the

classifier ("classifier" method). (C) anticipation labels are the ones predicted from pre-stimulus activity by the

classifier trained on uncued data: the training was done using post-stimulus-predicted labels to build the

"ground-truth" labels of anticipation ("post-stimulus" method). (D) anticipation labels are the ones predicted

from pre-stimulus activity by the classifier trained on cued data ("cued-pre-stimulus" method). $*: p < 0.05$, $***$

$: p < 0.005$.

Anticipation and diffusion-decision models

We fitted the DDM to individual behavioral data in the cued condition to assess the mechanism by

which response times and response accuracy are improved upon correctly anticipating the upcoming

stimulus. For this, the boundary $a = 1$ and non-decision time $T_0 = 0.3s$ were fixed for all

participants, while the drift v and starting point z_r was fitted for each participant depending on

whether the cue matched the stimulus or not. This resulted in 2 drifts and starting points per

participant.

We first tested whether the drift or the starting point varied significantly depending on the

correctness of anticipation. The drift was significantly greater on correctly anticipated trials

compared to incorrectly anticipated trials (paired-sample t-test: $t(41) = 4.322, p < 0.001$, Cohen's

$d = 0.667$). We also observed a significant effect of anticipation correctness on the starting point

value, whereby the starting point was closer to the correct decision boundary when the stimulus was

correctly anticipated (paired-sample t-test: $t(41) = 2.465, p = 0.018$, Cohen's $d = 0.380$).

We then tested whether these differences also related to the classification accuracy on cued trials.

We observed that participants that displayed above-chance classification performances had a

significantly more positive difference in starting point between correctly and incorrectly-anticipated

trials, compared to participants whose classification performance remained at chance-level

(independent sample t-test: $t(40) = 2.102, p = 0.042$, Cohen's $d = 0.668$). Moreover, we noted a

significant correlation between classification accuracy and difference in starting point (Pearson's

$r: 0.346, p = 0.025$). Note that neither analysis revealed a significant relation between drift

difference and classification accuracy.

Applying these results to assess classification performance on uncued data of the post-stimulus-

based and the cued-pre-stimulus-based pipelines, we repeated the subjective fitting, this time using

uncued trials. We then computed the difference between parameters of correctly and incorrectly

anticipated trials, and compared the differences within and across classification pipelines. Within

pipeline, none of the pipelines returned a significant difference between parameters, which we could

allocate to a poor fitting compared to cued trials. However, the comparison of these differences

across pipelines signals that the cued-pre-stimulus-based pipeline returns significantly more positive

starting point differences than the post-stimulus-based pipeline (Shapiro-Wilk test: $W = 0.615, p <$

0.001, Wilcoxon signed-rank test: $W = 346, p = 0.02$, Cohen's $d = 0.488$). Given our analysis on

cued trials, a better pipeline should display a more positive starting point difference, indicating that

the cued-pre-stimulus-based pipeline could have been more accurate than the post-stimulus-based

one.

Discussion

In this work, our objective was to delve into the neural underpinnings of anticipation within the

brain, focusing on scrutinizing its manifestations in both cued and uncued perceptual decision

contexts. Our work's core idea involved assessing the selectivity of the CNV and pre-stimulus alpha-

band power concerning specific perceptual anticipation. We established that the CNV contained

information about the expected stimulus at the single-trial level on cued trials. We subsequently

attempted single-trial decoding on uncued trials using two techniques. DDM modeling also allowed

us to link anticipatory activity to shifts in decision strategy, additionally providing a method to assess

the classification performance of anticipation on uncued trials.

The CNV emerged as a promising indicator, with 25 out of 42 participants displaying above-chance

single-trial classification performance on cued trials. This discernment significantly contrasts with

historical views associating CNV with non-specific temporal expectation (Walter et al., 1964) or

estimation (Kononowicz and Penney, 2016). More recently, works have shown that it was possible to

distinguish anticipation in Go/NoGo tasks at the single-trial level, that is, to distinguish anticipation

between trials where a movement is required relative to when movement has to be withheld

(Chavarriaga et al., 2012; Garipelli et al., 2011, 2009; Khaliliardali et al., 2015, 2012). While

Chavarriaga et al. (2012) emancipated their analyses from motor preparation by requiring no motor

output in one of their tasks, the authors argued that the differences are ascribable to attentional

changes following an erroneous proposition. The present work goes further by showing that the CNV

is specific to the anticipated type of stimulus, opening new avenues for understanding the richness of

anticipatory processes.

Second, pre-stimulus alpha power has historically been considered a cornerstone in examining the

pre-stimulus period. Despite significant differences in alpha-band activation between types of

anticipation, the single-trial classification performance using alpha-band activity yielded

comparatively less robust results than CNV-based classification. Several reasons could explain this.

First, alpha band activity was previously thought to be related to idling states (Pfurtscheller et al.,

1996), thus resulting in general rather than modality-specific brain signals. This, however, contradicts

more recent evidence that sensory gating (Jensen et al., 2012; Jensen and Mazaheri, 2010) and

selective attention (Foxe and Snyder, 2011) are stimulus-specific. The more likely alternative is that

the paradigms implemented in previous studies used stimuli at different perception levels, including

stimuli close to the individual perception threshold of each observer (Barik et al., 2019; Bode et al.,

2012; Samaha et al., 2017). In these instances, stimulus perception would highly depend on the

network excitability, represented by the alpha-band activity level. Here, we used distinguishable

stimuli. However, Samaha et al. (2017) showed that the alpha-band activity biased the confidence

ratings of participants and not the behavioral performance, which would mean that regardless of

their alpha activity, participants could still sense the visual information they were presented with. In

the same line, Benwell et al. (2021) showed that discrimination accuracy was unaltered by

fluctuations of pre-stimulus alpha power, while subjective ratings of awareness correlated negatively

with alpha power. This is still compatible with our explanation, as it would mark a difference

between sensed and perceived information, the latter requiring a conscious grasp of the stimulus.

A question that can arise from these results is the performance of the classifiers. Indeed, the

literature often reports classification accuracies of the order of 90% in supervised learning. Here, we

reported a classification accuracy of the order of 60%, which can appear relatively anecdotical.

However, it is important to note that the studies reporting such high classification accuracies observe

well-aligned, stimulus-evoked post-stimulus brain activity, whose typical amplitudes is of the order of

10 µV in the case of P300. In comparison, while we tempered the difficulty of timing anticipation

onset by introducing an anticipation window, it remains an endogenous process, i.e. not induced by

external stimulation. Moreover, the typical amplitude of the CNV is of the order of 2µV, which is an

order of magnitude smaller than well-studied stimulus-evoked activity. Remarkably, no study

currently reports above-chance classification performances between CNV types. Despite the

seemingly low classification performances, robust statistical analyses show that they lie well beyond

chance level.

The DDM analyses revealed an association between correct anticipation and shifts of the starting

point toward the correct decision boundary (similar to Bode et al. (2012)) and an augmentation in

the drift rate (similar to Urai et al. (2019)). This is consistent with the theory of premature sampling

(Grosjean et al., 2001; Laming, 1979a, 1979b), whereby information is integrated before the stimulus

appears, and the idea of sensory facilitation (Walz et al., 2015), whereby information is more easily

integrated when expected. Premature sampling results in a starting point shift, while sensory

facilitation could explain the increased drift rate. Interestingly, only the difference in starting point

between correct and incorrect anticipation significantly varied between participants whose CNVs

were classified with above-chance accuracy and participants with chance-level classification

accuracy. In that sense, the classifier could have captured premature sampling more efficiently in

those participants with good classification performance compared to those for which the classifier

yielded chance-level classification accuracy. We further used this result to infer the classification

performance on pre-stimulus activity of uncued trials.

The exploration extended into single-trial decoding of anticipation on uncued trials, where two

distinct techniques were employed. The first method relied on classifiers trained on cued pre-

stimulus data, while the second method involved a separate classifier trained on cued post-stimulus

activity trials to discern incorrect anticipations and an additional classifier trained on uncued pre-

stimulus data. This multifaceted approach allowed for a nuanced examination of the dynamics at

play during uncued trials. Remarkably, the classification performance on uncued trials exhibited

variability among participants, with 5 out of 42 showcasing above-chance accuracy with the post-

stimulus-based pipeline. This variation emphasizes the complexity of anticipatory processes in the

absence of cues, suggesting individual differences in the ability to decode anticipation.

Classifying uncued pre-stimulus activity is particularly challenging due to the lack of ground-truth

information regarding the anticipation class. The DDM analysis on these trials revealed that the

difference in starting points between correct and incorrect anticipations (as predicted by either

classifier) was greater when the anticipation class was predicted using a classifier trained on cued

data and applied directly on uncued data ("cued-pre-stimulus" pipeline) than when it was predicted

from a classifier trained on uncued data, using labels predicted from post-stimulus EEG activity

("post-stimulus" pipeline). However, upon comparing the distribution of anticipation and behavioral

outcomes across response time quantiles, we observed that the "post-stimulus" pipeline yielded

patterns more similar to cued trials than the "cued-pre-stimulus" pipeline, suggesting conversely a

better performance of the "post-stimulus" pipeline. Note, however, that this analysis is qualitative

and that some uncertainty remains, as a quantitative assessment of the quality of the classification of

anticipation based on post-stimulus activity on uncued data is equally challenging and was not

performed here.

Alternative explanations for the observed differences could be put forward. First, one could argue

that the differences reflect different motor preparations as the CNV is indeed traditionally known for

its motor component (Rockstroh et al., 1990; Rohrbaugh and Gaillard, 1983; van Boxtel and Böcker,

2004). To eliminate the motor factor from our analyses, participants had to respond to both stimuli

using the same motor output: a click from their dominant hand. Second, one could believe that the

differences are ascribable to the sensory processing of the different visual cues. While our analyses

do not completely discard this possibility, applying the classification pipelines only to the late part of

the anticipation period allows to remove the initial part of sensory processing, which was shown to

evoke distinguishable brain activity patterns at the single trial level (see Barachant and Congedo,

2014; Kalafatovich et al., 2020 for example). The later part of this activity, we argue, reflects

anticipation and while this anticipation is induced by the visual cue, the difference in brain activity

between the two types of anticipation is independent from the initial processing stages. In addition,

the uncued trials were introduced to completely discard any influence from the sensory processing

of cues, and gives a qualitative assessment that the features studied do in fact reflect anticipation

without any sensory processing influence.

Anticipation could emerge from a variety of factors. Among them, sequential effects have been

shown to impact subsequent decisions (Abrahamyan et al., 2016; Bode et al., 2012; Palminteri et al.,

2017; Urai et al., 2019; Yu and Cohen, 2008). Our experiment was designed to present the stimuli

randomly to each participant at each trial. Moreover, the stimuli were well distinguishable, reducing

the uncertainty. However, it has been shown that past stimuli and decisions impacted subsequent

decisions even without explicit sequences in the experiment (Yu and Cohen, 2008). We did not

consider such effects here, in particular not in the cued experiment, where we implicitly assumed

that the effect of the cue was greater than possible sequential effects. Indeed, the presence of the

cue reduces the perceived uncertainty, as underlined by the reduction of response times compared

to uncued trials. We argue, however, that these effects should be considered in more uncertain

contexts, for example, when attempting to classify uncued pre-stimulus periods if the stimuli are

closer to the perceptual threshold. The effect of global levels of attention on subsequent behavior

could also be evaluated, as drowsiness is reflected in post-stimulus EEG patterns, possibly reflecting

an alteration of the evidence integration stage of decision-making (Jagannathan et al., 2022). Future

studies could weigh the contribution of the specific anticipation, sequential effects, and attention on

behavior, and in particular, response times, using regression analyses, for instance.

The uncovered anticipatory processes are likely implicit for the main part. Indeed, while our

experiment did not include a feedback questionnaire, some participants reported deliberately

ignoring the cue in the calibration phase to avoid making mistakes. However, our behavioral analysis

showed that participants were systematically faster in the calibration phase compared to the test

phase, regardless of the order in which they performed the blocks. The accuracy, however, did not

decrease with increasing speed, although the cue was faulty 20% of the time. It indicates that

participants finalized their decisions upon stimulus presentation but still considered the cue

informative, as they tended to make more mistakes in catch trials, i.e., when the cue and the

subsequent stimulus were mismatched. A deeper analysis, which could include a scaling of the

reliability of the cue combined with behavioral measurements and subjective ratings of the

usefulness of the cue, could help investigate this effect.

In conclusion, our study has successfully unveiled the neural correlates of perceptual anticipation,

emphasizing the CNV's specificity to the anticipated stimulus type. Applying the DDM and exploring

pre-stimulus alpha power provided a nuanced understanding of the underlying mechanisms, further

enriching our comprehension of anticipatory processes. Future research directions could involve

deeper modeling techniques to refine the description of anticipation's influence on behavior,

emphasizing individual differences. Additionally, investigations into the interplay between specific

anticipation, sequential effects, and attention on behavior could offer a more comprehensive

understanding of the multifaceted nature of anticipatory processes. Our work sets the stage for

further exploration in ecologically valid scenarios and opens avenues for refining the classification of

anticipation in uncued trials. The multidimensional nature of anticipation, embedded in neural

processes, calls for continued exploration to unravel its complexities fully.

747 Data availability

748 The codes and data produced for this study will be made available upon publication.

749 Conflicts of interest

750 The authors declare no competing financial interests.

751 Bibliography

Abrahamyan, A., Silva, L.L., Dakin, S.C., Carandini, M., Gardner, J.L., 2016. Adaptable history biases in
human perceptual decisions. Proc. Natl. Acad. Sci. 113, E3548–E3557.
https://doi.org/10.1073/pnas.1518786113
Barachant, A., Barthélemy, Q., Gramfort, A., Jean-Rémi KING, Rodrigues, P.L.C., Dave, Olivetti, E.,
Goncharenko, V., Maxdolle, Berg, G.W.V., G.Reguig, Yamamoto, M.S., Artim436, Beasley, B.,
Bjäreholt, E., Clisson, P., Jliersch, Sassenhagen, J., Mccorsi, Stevenmortier, Stonebig, 2023.
pyRiemann/pyRiemann: v0.4. https://doi.org/10.5281/ZENODO.7642689
Barachant, A., Bonnet, S., Congedo, M., Jutten, C., 2013. Classification of covariance matrices using a
Riemannian-based kernel for BCI applications. Neurocomputing 112, 172–178.
https://doi.org/10.1016/j.neucom.2012.12.039

Barachant, A., Bonnet, S., Congedo, M., Jutten, C., 2012. Multiclass Brain–Computer Interface
Classification by Riemannian Geometry. IEEE Trans. Biomed. Eng. 59, 920–928.
https://doi.org/10.1109/TBME.2011.2172210
Barachant, A., Congedo, M., 2014. A Plug&Play P300 BCI Using Information Geometry.
Barik, K., Daimi, S.N., Jones, R., Bhattacharya, J., Saha, G., 2019. A machine learning approach to
predict perceptual decisions: an insight into face pareidolia. Brain Inform. 6, 2.
https://doi.org/10.1186/s40708-019-0094-5
Barthélemy, Q., Gouy-Pailler, C., Isaac, Y., Souloumiac, A., Larue, A., Mars, J.I., 2013. Multivariate
temporal dictionary learning for EEG. J. Neurosci. Methods 215, 19–28.
https://doi.org/10.1016/j.jneumeth.2013.02.001
Benwell, C.S.Y., Coldea, A., Harvey, M., Thut, G., 2021. Low pre-stimulus EEG alpha power amplifies
visual awareness but not visual sensitivity. Eur. J. Neurosci. ejn.15166.
https://doi.org/10.1111/ejn.15166
Bode, S., Sewell, D.K., Lilburn, S., Forte, J.D., Smith, P.L., Stahl, J., 2012. Predicting Perceptual Decision
Biases from Early Brain Activity. J. Neurosci. 32, 12488–12498.
https://doi.org/10.1523/JNEUROSCI.1708-12.2012
Chavarriaga, R., Perrin, X., Siegwart, R., del R Millan, J., 2012. Anticipation- and error-related EEG
signals during realistic human-machine interaction: A study on visual and tactile feedback, in:
2012 Annual International Conference of the IEEE Engineering in Medicine and Biology
Society. Presented at the 2012 34th Annual International Conference of the IEEE Engineering
in Medicine and Biology Society (EMBC), IEEE, San Diego, CA, pp. 6723–6726.
https://doi.org/10.1109/EMBC.2012.6347537
Chavarriaga, R., Sobolewski, A., Millán, J. del R., 2014. Errare machinale est: the use of error-related
potentials in brain-machine interfaces. Front. Neurosci. 8.
https://doi.org/10.3389/fnins.2014.00208
Congedo, M., Barachant, A., Bhatia, R., 2017. Riemannian geometry for EEG-based brain-computer
interfaces; a primer and a review. Brain-Comput. Interfaces 4, 155–174.
https://doi.org/10.1080/2326263X.2017.1297192
Ehrlich, S.K., Cheng, G., 2018. Human-agent co-adaptation using error-related potentials. J. Neural
Eng. 15, 066014. https://doi.org/10.1088/1741-2552/aae069
Foxe, J.J., Snyder, A.C., 2011. The Role of Alpha-Band Brain Oscillations as a Sensory Suppression
Mechanism during Selective Attention. Front. Psychol. 2.
https://doi.org/10.3389/fpsyg.2011.00154
Garipelli, G., Chavarriaga, R., del R. Millan, J., 2009. Fast Recognition of Anticipation-Related
Potentials. IEEE Trans. Biomed. Eng. 56, 1257–1260.
https://doi.org/10.1109/TBME.2008.2005486
Garipelli, G., Chavarriaga, R., Millán, J. del R., 2011. Single Trial Recognition of Anticipatory Slow
Corticla Potentials: the role of spatio-spectral filtering. Presented at the Proceedings of the
5th International IEEE EMBS Conference on Neural Engineering, Cancun, Mexico.
https://doi.org/10.1109/ner.2011.5910573
Grosjean, M., Rosenbaum, D.A., Elsinger, C., 2001. Timing and reaction time. J. Exp. Psychol. Gen.
130, 256–272. https://doi.org/10.1037/0096-3445.130.2.256
Iemi, L., Chaumon, M., Crouzet, S.M., Busch, N.A., 2017. Spontaneous Neural Oscillations Bias
Perception by Modulating Baseline Excitability. J. Neurosci. 37, 807–819.
https://doi.org/10.1523/JNEUROSCI.1432-16.2016
Jagannathan, S.R., Bareham, C.A., Bekinschtein, T.A., 2022. Decreasing Alertness Modulates
Perceptual Decision-Making. J. Neurosci. 42, 454–473.
https://doi.org/10.1523/JNEUROSCI.0182-21.2021
Jensen, O., Bonnefond, M., VanRullen, R., 2012. An oscillatory mechanism for prioritizing salient
unattended stimuli. Trends Cogn. Sci. 16, 200–206.
https://doi.org/10.1016/j.tics.2012.03.002

Jensen, O., Mazaheri, A., 2010. Shaping Functional Architecture by Oscillatory Alpha Activity: Gating
by Inhibition. Front. Hum. Neurosci. 4. https://doi.org/10.3389/fnhum.2010.00186
Kalafatovich, J., Lee, M., Lee, S.-W., 2020. Decoding Visual Recognition of Objects from EEG Signals
based on Attention-Driven Convolutional Neural Network.
Kappenman, E.S., Luck, S.J. (Eds.), 2011. The Oxford Handbook of Event-Related Potential
Components, 1st ed. Oxford University Press.
https://doi.org/10.1093/oxfordhb/9780195374148.001.0001
Khaliliardali, Z., Chavarriaga, R., Gheorghe, L.A., del Millan, J.R., 2012. Detection of anticipatory brain
potentials during car driving, in: 2012 Annual International Conference of the IEEE
Engineering in Medicine and Biology Society. Presented at the 2012 34th Annual
International Conference of the IEEE Engineering in Medicine and Biology Society (EMBC),
IEEE, San Diego, CA, pp. 3829–3832. https://doi.org/10.1109/EMBC.2012.6346802
Khaliliardali, Z., Chavarriaga, R., Gheorghe, L.A., Millán, J. del R., 2015. Action prediction based on
anticipatory brain potentials during simulated driving. J. Neural Eng. 12, 066006.
https://doi.org/10.1088/1741-2560/12/6/066006
Koch, C., Preuschoff, K., 2007. Betting the house on consciousness. Nat. Neurosci. 10, 140–141.
https://doi.org/10.1038/nn0207-140
Kononowicz, T.W., Penney, T.B., 2016. The contingent negative variation (CNV): timing isn't
everything. Curr. Opin. Behav. Sci. 8, 231–237. https://doi.org/10.1016/j.cobeha.2016.02.022
Laming, D., 1979a. Autocorrelation of choice-reaction times. Acta Psychol. (Amst.).
https://doi.org/10.1016/0001-6918(79)90032-5
Laming, D., 1979b. Choice reaction performance following an error. Acta Psychol. (Amst.) 43, 199–
224. https://doi.org/10.1016/0001-6918(79)90026-X
Langner, O., Dotsch, R., Bijlstra, G., Wigboldus, D.H.J., Hawk, S.T., van Knippenberg, A., 2010.
Presentation and validation of the Radboud Faces Database. Cogn. Emot. 24, 1377–1388.
https://doi.org/10.1080/02699930903485076
Lou, B., Li, Y., Philiastides, M.G., Sajda, P., 2014. Prestimulus alpha power predicts fidelity of sensory
encoding in perceptual decision making. NeuroImage 87, 242–251.
https://doi.org/10.1016/j.neuroimage.2013.10.041
Maris, E., Oostenveld, R., 2007. Nonparametric statistical testing of EEG- and MEG-data. J. Neurosci.
Methods 164, 177–190. https://doi.org/10.1016/j.jneumeth.2007.03.024
Min, B.-K., Herrmann, C.S., 2007. Prestimulus EEG alpha activity reflects prestimulus top-down
processing. Neurosci. Lett. 422, 131–135. https://doi.org/10.1016/j.neulet.2007.06.013
Min, B.-K., Park, J.Y., Kim, E.J., Kim, J.I., Kim, J.-J., Park, H.-J., 2008. Prestimulus EEG alpha activity
reflects temporal expectancy. Neurosci. Lett. 438, 270–274.
https://doi.org/10.1016/j.neulet.2008.04.067
Noah, S., Powell, T., Khodayari, N., Olivan, D., Ding, M., Mangun, G.R., 2020. Neural Mechanisms of
Attentional Control for Objects: Decoding EEG Alpha When Anticipating Faces, Scenes,and
Tools. J. Neurosci. 40, 4913–4924. https://doi.org/10.1523/JNEUROSCI.2685-19.2020
Palminteri, S., Lefebvre, G., Kilford, E.J., Blakemore, S.-J., 2017. Confirmation bias in human
reinforcement learning: Evidence from counterfactual feedback processing. PLOS Comput.
Biol. 13, e1005684. https://doi.org/10.1371/journal.pcbi.1005684
Petro, N.M., Thigpen, N.N., Garcia, S., Boylan, M.R., Keil, A., 2019. Pre-target alpha power predicts
the speed of cued target discrimination. NeuroImage 189, 878–885.
https://doi.org/10.1016/j.neuroimage.2019.01.066
Pfurtscheller, G., Stancák, A., Neuper, Ch., 1996. Event-related synchronization (ERS) in the alpha
band — an electrophysiological correlate of cortical idling: A review. Int. J. Psychophysiol. 24,
39–46. https://doi.org/10.1016/S0167-8760(96)00066-9
Poulton, E.C., 1950. Perceptual Anticipation and Reaction Time. Q. J. Exp. Psychol. 2, 99–112.
https://doi.org/10.1080/17470215008416582

Railo, H., Piccin, R., Lukasik, K.M., 2021. Subliminal perception is continuous with conscious vision
and can be predicted from prestimulus electroencephalographic activity. Eur. J. Neurosci. 54,
4985–4999. https://doi.org/10.1111/ejn.15354

Ratcliff, R., 1978. A Theory of Memory Retrieval. Psychol. Rev. 85, 59–108.
https://doi.org/10.1037/0033-295X.85.2.59

Ratcliff, R., McKoon, G., 2008. The Diffusion Decision Model: Theory and Data for Two-Choice
Decision Tasks. Neural Comput. 20, 873–922. https://doi.org/10.1162/neco.2008.12-06-420

Ratcliff, R., Tuerlinckx, F., 2002. Estimating parameters of the diffusion model: Approaches to dealing
with contaminant reaction times and parameter variability. Psychon. Bull. Rev. 9, 438–481.
https://doi.org/10.3758/BF03196302

Rivet, B., Souloumiac, A., Attina, V., Gibert, G., 2009. xDAWN Algorithm to Enhance Evoked
Potentials: Application to Brain–Computer Interface. IEEE Trans. Biomed. Eng. 56, 2035–
2043. https://doi.org/10.1109/TBME.2009.2012869

Rockstroh, B., Elbert, T., Birbaumer, N., Lutzenberger, W., 1990. Biofeedback-produced hemispheric
asymmetry of slow cortical potentials and its behavioural effects. Int. J. Psychophysiol. 9,
151–165. https://doi.org/10.1016/0167-8760(90)90069-P

Rohrbaugh, J.W., Gaillard, A.W.K., 1983. 13 Sensory and Motor Aspects of the Contingent Negative
Variation, in: Advances in Psychology. Elsevier, pp. 269–310. https://doi.org/10.1016/S0166-
4115(08)62044-0

Samaha, J., Iemi, L., Postle, B.R., 2017. Prestimulus alpha-band power biases visual discrimination
confidence, but not accuracy. Conscious. Cogn. 54, 47–55.
https://doi.org/10.1016/j.concog.2017.02.005

Trevena, J.A., Miller, J., 2002. Cortical Movement Preparation before and after a Conscious Decision
to Move. Conscious. Cogn. 11, 162–190. https://doi.org/10.1006/ccog.2002.0548

Urai, A.E., de Gee, J.W., Tsetsos, K., Donner, T.H., 2019. Choice history biases subsequent evidence
accumulation. eLife 8, e46331. https://doi.org/10.7554/eLife.46331

van Boxtel, G.J.M., Böcker, K.B.E., 2004. Cortical Measures of Anticipation. J. Psychophysiol. 18, 61–
76. https://doi.org/10.1027/0269-8803.18.23.61

van Dijk, H., Schoffelen, J.-M., Oostenveld, R., Jensen, O., 2008. Prestimulus Oscillatory Activity in the
Alpha Band Predicts Visual Discrimination Ability. J. Neurosci. 28, 1816–1823.
https://doi.org/10.1523/JNEUROSCI.1853-07.2008

Voss, A., Voss, J., 2007. Fast-dm: A free program for efficient diffusion model analysis. Behav. Res.
Methods 39, 767–775. https://doi.org/10.3758/BF03192967

Walter, W.G., Cooper, R., Aldridge, V.J., McCALLUM, W.C., Winter, A.L., 1964. Contingent Negative
Variation : An Electric Sign of Sensori-Motor Association and Expectancy in the Human Brain.
Nature 203, 380–384. https://doi.org/10.1038/203380a0

Walz, J.M., Goldman, R.I., Carapezza, M., Muraskin, J., Brown, T.R., Sajda, P., 2015. Prestimulus EEG
alpha oscillations modulate task-related fMRI BOLD responses to auditory stimuli.
NeuroImage 113, 153–163. https://doi.org/10.1016/j.neuroimage.2015.03.028

Wöstmann, M., Waschke, L., Obleser, J., 2019. Prestimulus neural alpha power predicts confidence in
discriminating identical auditory stimuli. Eur. J. Neurosci. 49, 94–105.
https://doi.org/10.1111/ejn.14226

Yang, Y.-F., Brunet-Gouet, E., Burca, M., Kalunga, E.K., Amorim, M.-A., 2020. Brain Processes While
Struggling With Evidence Accumulation During Facial Emotion Recognition: An ERP Study.
Front. Hum. Neurosci. 14, 340. https://doi.org/10.3389/fnhum.2020.00340

Yu, A.J., Cohen, J.D., 2008. Sequential effects: Superstition or rational behavior? Adv. Neural Inf.
Process. Syst. 21, 1873–1880.

910