Individual variation in male pheromone production in *Xylocopa sonorina* correlates with size and gland color.

ANDREW J. GOFFINET[1+], KATHY DARRAGH[1+], NICHOLAS SALEH[1a],

MADELEINE M. OSTWALD[2], STEPHEN L. BUCHMANN[3], SANTIAGO R. RAMIREZ[1*]

[1]*Department of Evolution and Ecology, University of California, Davis, CA, 95616*

[2]*Cheadle Center for Biodiversity & Ecological Restoration, University of California,*

Santa Barbara, CA, 93106

[3]*Department of Ecology & Evolutionary Biology, University of Arizona, Tucson, AZ,*

85721

ORCID Kathy Darragh 0000-0002-9215-014X

ORCID Nicholas W. Saleh 0000-0002-5520-3396

ORCID Madeleine M. Ostwald 0000-0002-9869-8835

ORCID Santiago R. Ramírez 0000-0003-1306-1315

[+] These authors contributed equally

[a] Current address: School of Natural Sciences, Fresno Pacific University, Fresno, CA, 93702

*Corresponding author (Email: sanram@ucdavis.edu)

Acknowledgements

We acknowledge the support and feedback of the Ramirez lab with this project.

Abstract- Sex pheromones are species-specific chemical signals that facilitate the location,

identification, and selection of mating partners. These pheromones can vary between individuals,

and act as signals of mate quality. Here, we investigate the variation of male pheromones in the

mesosomal glands of the large carpenter bee *Xylocopa sonorina*, within a Northern California

population. We tested the hypothesis that morphological traits are correlated with the observed

variation in chemical blend composition of these bees. We also conducted behavioral assays to

test whether these male pheromones act as long-range attractants to conspecifics. We found that

larger males with darker mesosomal glands have a higher pheromone amount in their glands. Our

analysis also shows initial evidence that this pheromone blend serves as a long-range attractant to

both males and females. We show that both male body size and sexual maturation are important

factors influencing pheromone abundance, and that this pheromone blend acts as a long-range

attractant. We hypothesize that this recorded variation in male pheromone could be important for

female choice.

Key words- Pheromones, sexual selection, mate choice, chemical communication.

INTRODUCTION

Detection of chemical signals is considered to be amongst the most ancient and widespread of sensory systems, playing a major role in both inter- and intraspecific communication (Ache and Young 2005; Amo and Bonadonna 2018). In insects, pheromones are interspecific chemical signals, which are important for a diverse range of behaviors including aggregation (Wertheim et al. 2005), trail recruitment, and social organization (Czaczkes et al. 2015), and mating (Wyatt 2014). The chemical signals themselves can be comprised of a single compound (Daimon et al. 2012) or a mix of multiple compounds (Andersen et al. 1988)**.** A particularly well-studied group of chemical signals are those involved in mating, known as sex pheromones (Wyatt 2014).

Sex pheromones can serve a multitude of purposes during mating and mate choice. In some cases they are important for the initial step of mate attraction, acting over long ranges to attract individuals of the opposite sex (Itagaki and Conner 1988; Baker 1989). They are often species-specific signals, maintaining reproductive isolation by reducing hybridization between sympatric species (Daimon et al. 2012). Sex pheromones can also play key roles in intraspecific mate choice and/or signaling mate quality to the receiver, two roles that are not mutually exclusive (Juárez et al. 2015).Sex pheromones are thought to act as honest signals of mate quality, as diet can impact pheromone production (Darragh et al. 2019), with males fed a higher quality diet showing increased reproductive success (Rantala et al. 2003; Liedo et al. 2013). Therefore, while pheromones are often species-specific, they also show variation both between and within populations due to genetic drift, selection, or variation in diet quality or environmental conditions (Allison and Cardé 2016).

The large carpenter bees (Apidae: *Xylocopa*) are a cosmopolitan genus of wood nesting bees, which display a range of mating behaviors and male secondary sexual traits. Ancestrally, *Xylocopa* males defend resources to gain access to females, which is correlated with small or no mesosomal glands and a lack of sexual dimorphism (Leys and Hogendoorn 2008). However, in some subgenera, males defend non-resource based territories (Minckley 1994). These species tend to have sexually dimorphic males, with males exhibiting enlarged mesosomal glands, potentially some of the largest exocrine glands in insects (Minckley 1994). These glands release long-range sex pheromones to attract females and are thought to signal indirect benefits. An

especially well-studied species in this group is *Xylocopa sonorina*, the Valley Carpenter Bee,

found from California, Arizona, New Mexico, to Texas and into Sonora, Mexico, and introduced

to Hawaii and various Pacific Islands. This species displays extreme sexual dimorphism, with

all-black females and golden males with green eyes (Figure 1) (Marshall and Alcock 1981). In

early spring, males leave their nests daily in the late afternoon, where they defend a non-resource

based territory (Marshall and Alcock 1981; Alcock and Smith 1987). These are often referred to

as a lek mating system (as in some birds) where one or more male carpenter bees simultaneously

display to arriving females as potential mates (Marshall and Alcock 1981). Despite female

interactions being relatively infrequent, the majority of these interactions do not result in

copulation (Marshall and Alcock 1981; Alcock and Johnson 1990). Additionally, given that

females must land to commence copulation, and that the males do not pursue unmated females,

mate choice appears to be driven primarily by the female (Alcock and Smith 1987; Alcock and

Johnson 1990).

Sex pheromones play an important role in male *X. sonorina* mating displays. Males

choose a focal point, either a branch or a sprig of a non-flowering shrub or tree, which they

scent-mark and hover above, either chasing away or flying in loops with conspecific males

(Marshall and Alcock 1981; Alcock and Smith 1987). The pheromone released by males is easily

detectable to human observers (Marshall and Alcock 1981), who perceive it as a distinctive rose-

scent, with the primary constituent being a common compound found in rose volatiles (Dani et

al. 2021). One hypothesis is that the similarity of this pheromone to floral scents could have

evolved to exploit pre-existing sensory systems used to find flowers for nectar sources (Schiestl

2010). The pheromone itself is a blend of three primary compounds: all trans-geranylgeraniol, all

trans-farnesal and 3,7,11-methyl-2,7,10-dodecatrienal in a 9:6:1 ratio (Andersen et al. 1988). In

a previous field bioassay, the two most abundant components seem to attract females as

successfully as wild males (Minckley et al. 1991). Producing the pheromone is thought to be

costly as the mesosomal glands, which are lacking in females, are very large, occupying almost

20% of thoracic volume, otherwise occupied by the indirect flight muscles in other bee species

(Ostwald et al. 2022). The quantity of male sex pheromone produced potentially indicates male

quality, with females seeming to assess several males before making a decision (Marshall and

Alcock 1981). Across *Xylocopa* species, enlarged glands are associated with long-distance mate

attraction(Minckley et al. 1991); within a species, larger glands may similarly store greater

quantities of pheromones and facilitate attraction of females at greater distances.

The mesosomal gland undergoes a distinct ontogenetic development process throughout

the adult lifespan of *X. sonorina* males. In pre-sexual males, who overwinter as adults with

siblings in their natal nests, the gland exists in a static, non-secretory phase. The following

spring, the gland enters a secretory phase and pheromone production begins approximately a

month before they begin to leave their nests to seek mating opportunities at lek sites (Buchmann

pers. obs.). At some point during the mating season, the gland darkens in color as cuticular

sclerotization replaces the pale-colored former secretory tissue cell layers with hard, setal-lined

hollow tubules (Ostwald et al. 2022). This transition marks the ends of the gland's secretory

phase, with the gland now functioning as a storage organ for the accumulated pheromone blend

(Buchmann pers. obs). In this way, gland color can indicate its developmental stage.

In this study, we investigate how the chemical profile of *Xylocopa sonorina* males varies

within its population from Northern California. We analyze the mesosomal gland contents

released by displaying males during their active season. We describe the pheromone blend found

within the glands and identify factors correlated with pheromone variation, including body size

and gland color, both of which may covary with mating success.. We also carry out field

behavioral assays to test whether male pheromones serve as long range attractants for both males

and females.

METHODS AND MATERIALS

Sampling

Male bees were collected between April 2020 and May 2022 while actively displaying in

the late afternoon. Samples were collected throughout the city of Davis, California, USA. For a

detailed list of sites, see supplementary materials (Table S1). In addition, two males were

collected by hand from their nest in a log prior to the commencement of their mating season.

These latter samples were excluded from the overall analysis as they were collected using a

different method and later found to lack the main pheromone compounds. Upon collection, the

samples were frozen at at -20ºC. First, we measured the intertegular distance, which we use as a

proxy for body size (Cane 1987). We also recorded the wing wear (nicked wing edges), which

we used as a proxy for male age (Mueller and Wolf-Mueller 1993). To dissect the thoracic

pheromone gland from the mesothorax, the recently frozen bees were pinned onto an agar

dissection dish, with one pin placed through the prothorax and an additional pin placed through

the abdomen. Using forceps, the dorsal mesosomal cuticle was cut away, exposing the

mesosomal gland below. The glands were removed from the thorax, placed onto a white

microscope stage, and photographed, to measure their color. The camera, a Cannon EOS 60D,

was mounted on a tripod, approximately 40 centimeters above the microscope stage, and

manually focused during each sample. Lighting was provided using a portable microscope lamp

with two bulbs, placed on opposing sides of the stage, and diffused through wax paper.

Coloration Analysis

The Fiji distribution of ImageJ was used to collect color data (Schindelin et al. 2012). In

each photo, the gland was hand selected, five random points were chosen across the gland, and

across these five points, the program provided both the RGB (red-green-blue color model) values

and the HSB (hue-saturation-brightness color model) values for the image. These values were

then averaged. To assess color, a ratio of R (red) and G (green) was calculated, as the different

color values are only meaningful relative to the other values, and this ratio was used in our

analysis (Bergman and Beehner 2008).

Chemical Analysis

Each thoracic gland was placed in a glass vial containing 500μl hexane with 5μg of 2-

undecanone as an internal standard and solvent extracted for one hour. Following this, the

solvent was transferred to a new vial, and stored at -20°C. Samples were analyzed using an

Agilent model 5977A mass-selective detector connected to Agilent GC model 7890B, with a HP-

5 Ultra Inert column (Agilent, 117 30 m × 0.25 mm, 0.25 μm). 1μl of each sample was injected

using Agilent ALS 7694 autosampler in split mode with a 5:1 ratio with helium as the carrier gas

(250°C injector temperature, split flow of 3.5 ml/min). The temperature program started at 80°C

for 3 minutes, and then rose at 10°C/min to 300°C. The temperature was held at 300°C for 3

minutes and 315°C for 3 minutes. We quantified components using an internal standard of 2-

undecanone, and compounds not found in at least 10% of individuals were disregarded.

Compounds were identified by comparing to prior work done on this system (Andersen et al.

1988) and to NIST reference libraries. Furthermore, we identified compounds all trans-

geranylgeraniol and all trans-farnesal using authentic standards purchased from Sigma-
Aldrich chemicals.

Behavioral Assay

We conducted field behavioral bioassays in the spring of 2023 at a pair of test sites in the
UC Davis Arboretum (38.5328, -121.7547). The sites were 10 meters apart, on the same species
of shrubby plant. On each study day, a coin was flipped to randomly assign one of two
treatments to each site. During the study, each site had a qualitative cellulose filter paper disc
(Sterlitech, Catalog # CFP3-090), with a diameter of 9 cm, suspended from a branch of the shrub
approximately 1.5 meters off the ground using an odorless, green lab tape. To create the
experimental mixture, we used a mixture of the actual extracts from the top 75% of males in
terms of total pheromone abundance (n=30) from the samples included in this study. We applied
1ml of this mixture to the filter paper in the field. Since each male's gland was extracted into 0.5
ml of solvent, each 1ml aliquot represents two males' worth of pheromone. The second treatment
was a control, which consisted of 1 ml of pure hexane, also applied to the filter paper discs in the
field.

Following the application of the two treatments, the site was observed for the following
60 minutes and all interactions from both male and female bees were recorded. Interactions were
counted if an individual altered their flight path and approached the treated papers within 1
meter. We also noted if a "strong" interaction occurred, in which case the individual stopped and
hovered within 0.5 meter of the filter paper for any length of time. We chose to observe changes
in flight path and hovering as these are likely to be the initial stages of mate finding. Mating
attempts would not be expected with solely chemical cues because visual cues are also required
in this system for mating (Marshall and Alcock 1981). Initially, observations took place from
16:00-17:00 in the afternoon, coinciding with male *X. sonorina* activity. However, throughout the
season, this shifted back to 16:30-17:30 and eventually 17:00-18:00, following the observed
male display behavior, which progressed later into the afternoon as the season advanced (Table
S7). We carried out 14 assays between mid-May and the end of June 2023. Previous observations
found that male activity only happens at higher temperatures (>26.5ºC). We, therefore, only
carried out the assay on days with temperatures high enough to observe males displaying on
nearby plants.

Statistical Analysis: Variation in Compound Amount

Firstly, we were interested in determining the factors correlated with variation in total compound amount produced by the thoracic gland. We tested the effects of intertegular distance, wing wear, and various characteristics of the glands appearance, including color (R:G ratio), hue, saturation, and brightness, on total compound abundance, using ANOVA. We then evaluated our model fit with Akaike's information criteria (AIC) using the *AICcmodavg* package (Mazerolle 2020). The model with the lowest AIC score was chosen as the final model, containing the most likely explanatory variables.

Statistical Analysis: Variation in Chemical Profile Composition

The univariate analysis only includes total compound amount and does not consider which compounds are present in the chemical profile. To include this information in the analysis, we carried out a multivariate analysis to analyze the chemical mixture of the males, taking all the compounds into account. Due to the zeros present in our data matrix, we visualized divergence across the samples using nonmetric multidimensional scaling (NMDS) in three dimensions, based on a Bray–Curtis similarity matrix created using absolute peak areas. We created the NMDS with the "metaMDS" function in *vegan,* and this was plotted using the *ade4* package (Thioulouse et al. 2018; Oksanen et al. 2022).

To investigate factors that explain pheromone composition, we used a PERMANOVA (permutational multivariate analysis of variance) on a Bray-Curtis similarity matrix, using the "adonis2" function in *vegan* with 1000 permutations (Oksanen et al. 2022). We assessed each term individually using the option by="margin", this ensures that the order of terms in the model does not affect the results (in contrast to the by="sequential" option where each term is considered in order). Model fit was evaluated using AIC scores, with each model being tested sequentially. The model with the lowest AIC score was selected as the final model and identified as the model containing the explanatory factors. However, we chose the simplest model as the best fit if we found that two models were within two AIC values of each other. We also repeated this analysis with the relative compound abundances.

Statistical Analysis: Variation in the Most Abundant Compounds

The chemical profile of the bees is dominated by three main compounds, as described

previously (Andersen et al. 1988). These compounds appear to be biologically important to the

reproduction of *X. sonorina*, since an artificial mixture of two of these compounds, all trans-

farnesal and all trans-geranylgeraniol, seems to be sufficient to attract females (Minckley et al.

1991). Therefore, we re-analyzed our data including just these three primary compounds to

determine which factors are correlated with their abundance and composition as described for the

analysis of the entire dataset. We also carried out Pearson's correlation tests for these compounds

with both size and brightness.

Plotting and Data Manipulation

To make plots we used the following packages: *ggpubr* (Kassambara 2020), *cowplot*

(Wilke 2020) and *ggplot2* (Wickham 2016). In addition we used *tidyr* (Wickham and Girlich

2022), *plyr* (Wickham 2011), and *readr* (Wickham et al. 2023) for data transformation. Analyses

were carried out in R version 4.2.1 (R Core Team 2022).

Behavioral Assay Analysis

To determine if the likelihood of interaction was related to either sex of the individuals or

treatment (male gland extract vs. hexane control), we used a binomial GLM (generalized linear

model), with the package *MASS* (Venables and Ripley 2002). The response variable was whether

at least one interaction was observed in a given trial. We included sex and treatment as

explanatory variables as well as an interaction term between the two. To test significance of

factors, we used likelihood ratio tests on models with and without the variable of interest.

All raw data and analyses are available on the OSF

(https://osf.io/uf87d/?view_only=ab07af2f3c4e4c05a5f9f5fad680dd64).

236   RESULTS

We sampled a total of 40 males and identified 43 different compounds in their chemical

profile. On average, 284.5±168.7 *μ*g (SD) of total compound were present in each sample.

Across all the samples, the three main compounds described by prior studies in an

Arizona population were also found in the Davis population, with all trans-geranylgeraniol, all

trans-farnesal, and 3,7,11-trimethyl-2,7,10-dodecatrienal being the three most abundant

compounds (Figure 1, Panel B, labels 1, 2, 3) (Andersen et al. 1988).

In the sampled displaying males, we observed a wide range of gland color, with some

much darker than others (Figure S1). In addition to the displaying males, we also sampled two

adult males which were collected from their nest in a log prior to the start of the mating season.

Upon dissection, we observed that these males had extremely pale, almost white mesosomal

glands, and when chemically analyzed, these males had an average total compound amount of

2.08 ± 1.1 μg (SD). Additionally, these males lacked the three main compounds seen in the

actively displaying males.

Variation in Compound Amount

To investigate the impact of different factors on total compound amount, we carried out a

univariate ANOVA analysis. We found that both brightness and size were significantly related to

the total amount of compound (Brightness: ANOVA, $F_{1,36}= 10.99$, p=0.002094; Size: ANOVA,

$F_{1,36}=14.02$, p=0.000632) (Figure 2, Table S2). We found that males with brighter glands have an

overall lower total compound amountThese results agree with similar findings about the glands

progressing through a pre-secretory (lightest = brightest color) phase, then into an active secretor

phase, and finally a dark reddish brown storage phase, at which point males will have

accumulated their maximum store of pheromones (Buchmann, unpubl.). We also found that

smaller males, measured by their intertegular distance, produce a lower total compound amount.

We did not find that wing wear significantly explained variation in compound amount.

Variation in Chemical Profile Composition

To identify which factors drive overall compositional differences in the chemical profile

of male glands, we carried out a multivariate PERMANOVA analysis. As in the univariate

analysis, the best model included both brightness (PERMANOVA, Brightness, $F1,36 = 5.1835$, p

$=0.001998$) and size (PERMANOVA, Size, $F1,36 = 7.5555$, $p =0.000999$) (Figure 3, Table S3).

We found that size accounts for 16% of observed variation in chemical profile, with a further

11% explained by gland brightness. We did not find that wing wear significantly explained

variation in chemical profile.

When we reanalyzed the data using relative compounds abundances, we found that none

of our variables explained variation in chemical profile (Table S4).

Variation in the Most Abundant Compounds

The prior two analyses included every compound which was present in at least 10%

sampled males. However, prior studies have identified three main compounds, two of which can

attract females, and so we re-ran the previous analyses on a reduced dataset of these three main

compounds (Andersen et al. 1988; Minckley et al. 1991). Combined, these three main

compounds made up an average of 250.9 ± 157.7 μg (SD) in each male gland, which is $86.4\% \pm$

6.02% (SD) of total gland pheromone amount.

We found that both brightness and size was correlated with the total amount of compound

(Brightness: ANOVA, $F_{(1,36)}= 11.32$, p=0.001832; Size: ANOVA, $F_{(1,36)}=13.66$,

p=0.000724)(Figure S2, Table S5). To identify which factors drive overall compositional

differences in the three main compounds, we carried out a PERMANOVA analysis. The best

model fit again included both brightness (PERMANOVA, Brightness, $F1,36 = 5.1614$, p

$=0.003996$) and male bee size (PERMANOVA, Size, $F1,36 = 7.9076$, $p =0.000999$), with size

explaining 17% of observed variation in the three main compounds, and brightness explaining a

further 11% (Figure S3, Table S6). We did not find that wing wear significantly explained

variation in the three main compounds. These results were coincident with the entire chemical

blend, suggesting that these compounds are responsible for the patterns seen when analyzing the

entire dataset.

To determine if all three of these compounds are correlated with brightness and size, we

analyzed each compound individually. We found that variation in both all trans-farnesal and all

trans-geranylgeraniol is related to size and brightness (all trans-farnesal: Size, ANOVA, $F_{(1,36)}=$

9.1, p=0.0047; Brightness, ANOVA, $F_{(1,36)}= 10.9$, p=0.0022)(all trans-geranylgeraniol: Size,

ANOVA, $F_{(1,36)}= 11.88$, p=0.0015; Brightness, ANOVA, $F_{(1,36)}= 7.72$, p=0.0086). We did not

find that variation in 3,7,11-trimethyl-2,7,10-dodecatrienal was explained by size or brightness.

We also calculated the strength of correlation between all trans-farnesal and all trans-

geranylgeraniol with size and brightness. We found that all trans-farnesal is significantly

correlated with both size (Pearson's correlation, r=0.4, p=0.011) and brightness (Pearson's

correlation, r=-0.37, p=0.019). In contrast, all trans-geranylgeraniol is only significantly

correlated with size (Pearson's correlation, r=0.45, p=0.0032), but not brightness (Pearson's

correlation, r=-0.3, p=0.068). Taken together, these two compounds are responsible for the main

patterns detected in the overall dataset (Figure 4).

Behavioral Assay Results

In total, we conducted 14 behavioral field trials (bioassays), and interactions by either

males or females occurred in 13 of the 14 trials, resulting in 37 total interactions (Table 1, Table

S7). Females interacted with the control treatment on 4/14 days, while males interacted with the

control on 1/14 days. These numbers were much higher for the filter paper treated with extract,

with females interacting 11/14 days, and males 8/14 days. Treatment was found to significantly

explain the likelihood of interaction (2ΔlnL= 16.016, d.f.=1, p=0.000063).

We also observed what we have termed "strong responses". These consist of individuals

stopping and hovering within 0.5 meters of the filter paper. This occurred on 3 separate days for

the males, and 5 separate days for the females, and was observed solely on the filter paper with

the male's extract applied, while this hovering behavior did not occur towards the control (Table

S7).

DISCUSSION

Xylocopa sonorina males release large amounts of their sex pheromone during mating

displays which are detectable to humans as a rose-like scent (Alcock and Smith 1987). Here, we

investigated the chemical ecology of *X. sonorina*, describing pheromone variation within a

population in California and identified morphological factors which could explain the observed

variation. As previously identified in Arizona individuals, we found that the three main

constituents of this gland were all trans-geranylgeraniol, all trans-farnesal, and an isomer of

3,7,11-trimethyl-2,7,10-dodecatrienal (Andersen et al. 1988). This shows a high level of

consistency and stability in this chemical signal across the range of this species, and in two areas

with different climates. We show that larger males with darker gland coloration (*i.e.* mature

glands) have more total pheromone in their glands. We found that this is mainly by variation in

the abundance of all trans-farnesal and all-trans geranylgeraniol, two of the major compounds

found in the gland. These two compounds have also previously been shown to be sufficient to

attract females (Minckley et al. 1991), a result which we further support here, and provide new

evidence that males are also attracted to male pheromones.

We found that male size explained the most observed variation in male pheromone

amount in *X. sonorina*. In bees, as with other holometabolous insects, the male's adult body size

is determined during the larval stage, and is heavily impacted by both abiotic (temperature,

latitude, altitude, season) and biotic (primarily diet) factors (Chole et al. 2019). Larger male body

size is thought to signal a higher quality diet during development, which corresponds to

increased mating success (Agosta 2010). We found that larger *X. sonorina* produce more sex

pheromone, which we hypothesize could be related to a higher mating success, as seen across a

range of insects. For example, in the tobacco moth, *Ephestia elutella*, larger males produce more

pheromone when compared to smaller males, and these pairings result in larger offspring, with

size being positively correlated with mating success in these moths (Phelan and Baker 1986). We

hypothesize that larger male *X. sonorina* which produce a greater abundance of their sex

pheromones could allow these larger males to attract females from a greater distance than

smaller males, or potentially to release pheromones for a longer period of time. However, further

work is required to discern if females show a preference for these larger males.

We also found that male gland color was correlated with pheromone amount. In some

cases, age could be signaled by sex pheromones, ensuring that mating occurs in the window

between sexual maturation and senescence (Gomez-Diaz, Carolina and Benton, Richard 2013).

In the butterfly *Bicyclus anynana*, the male's sex pheromone composition changes as they age,

and females exhibit preferences for the pheromones of middle aged males (Nieberding et al.

2012), with similar patterns documented in other species (Lassance and Löfstedt 2009; Chemnitz

et al. 2015). We did not find a relationship between male *X. sonorina* sex pheromone and age

when using wing wear as a proxy; however this may be a poor indicator of age due to variability

in wing degradation (Eltz et al. 1999). We decided to use the color of the gland as a proxy for

male maturation as previous work on *X. sonorina* has found the gland is initially light in color

and subsequently darkens into a sclerotized storage organ with many hair lined tubules

(Buchmann pers. obs.; Minckley et al. 1991; Ostwald et al. 2022). During our dissections, we

observed a range of colors of the mesosomal glands, from near white in some that were collected

prior to the commencement of the males display season to very dark red-brown in some

displaying males. The two males collected from their nests before their display season began had
nearly white mesosomal gland tissues containing only trace amounts of pheromone. We found
that the brightness of the gland correlates with pheromone amount, with darker glands containing
higher amounts of pheromone, in agreement with the prediction that older males have higher
pheromone amounts.

The effect of higher sex pheromone production by larger, more sexually mature male *X.*
sonorina on female choice is unclear. Two ways in which sex pheromone production could
impact female choice are through the effectiveness of the pheromone as a long-range attractant
or as an indicator of mate quality. Based on our behavioral assays, we present preliminary
evidence that these compounds serve as a long-range attractant for both males and females. In
hymenopterans, it has long been accepted that sex pheromones serve as important facilitators of
sexual reproduction by acting as long-range attractants (Ayasse et al. 2001). Within *X. sonorina,*
an early study showed that the primary constituents from the male gland seemed to be sufficient
to attract females (Minckley et al. 1991). The results from our behavioral assay provide further
evidence to support those observations.

Nonetheless, this does not discount the possibility that females are assessing male quality
through their sex pheromones. *Xylocopa sonorina* exhibits a non-resource-based mating system.
This is in contrast to the mating system of two other *Xylocopa* species found in California, *X.*
tabaniformis and *X. californica.* In these species, males defend resource-based territories, have
enlarged eyes and lack the large mesosomal glands found in *X. sonorina.* This suggests that as
Xylocopa bees shifted from a resource based, territorial mating strategy to a non-resource-based
mating strategy, they shifted from a visual based mating system to one that is facilitated by
pheromones. In the *X. sonorina* system, females must land for mating to commence (Marshall
and Alcock 1981), and mating occurs relatively infrequently, with only 24% of female
interactions with displaying males resulting in copulation (Alcock and Johnson 1990). It appears
that females are assessing multiple males, often in a lekking display, with status communicated
through chemical signals. However, further studies are required to test this hypothesis and
determine the effect that the male's mesosomal glands play in the female assessment.

As well as being used by females to locate and assess mate quality, sex pheromones may
also be used to facilitate male-male interactions. When conspecific, displaying *X. sonorina* males

encounter each other at the same display location, they may exhibit one of two different

behavioral responses. Sometimes, one male will emerge dominant and chase off their competitor,

while in other cases, up to several males may share a display site, flying in loose circles around

each other and the focal point for some length of time (Alcock and Smith 1987). Further studies

are needed to determine if this variation in behavior is reflected in the chemistry of the

pheromones released by displaying male *X. sonorina*. Based on results from our behavioral

assay, we provide evidence that male *X. sonorina* are attracted to the mesosomal chemical blend

released by males during their mating display. Additionally, similar to what has been described in

other hymenopterans, such as sphecid (Coelho and Holliday 2001) and pompilid wasps (Alcock

and Kemp 2006), larger males may be better fliers or a have a better ability to defend their

territory. *Xylocopa sonorina* may provide a viable system to investigate how sex pheromones

facilitate male-male interactions during their complex mating displays.

Our study reveals variation in sex pheromone amounts are correlated with both body size

and gland color in male *X. sonorina*. Larger, more sexually mature males produce higher

amounts of sex pheromone. In addition, we show preliminary evidence that this blend of male

pheromones serves as a long-range attractant to both the opposite and same sex. Future studies

could clarify the fitness consequences of variation in pheromone amount, whether by impacting

attraction range, female choice, or male-male competitive interactions.

REFERENCES

Ache BW, Young JM (2005) Olfaction: Diverse Species, Conserved Principles. Neuron 48:417–430. https://doi.org/10.1016/j.neuron.2005.10.022

Agosta SJ (2010) Male Body Size and Mating Success and Their Relation to Larval Host Plant History in the Moth *Rothschildia lebeau* in Costa Rican Dry Forest: Body Size and Mating Success. Biotropica 42:201–207. https://doi.org/10.1111/j.1744-7429.2009.00556.x

Alcock J, Johnson MD (1990) Female mate choice in the carpenter bee *Xylocopa varipuncta* (Hymenoptera: Anthophoridae). J Zool 221:195–204. https://doi.org/10.1111/j.1469-7998.1990.tb03991.x

Alcock J, Kemp DJ (2006) The Behavioral Significance of Male Body Size in the Tarantula Hawk Wasp *Hemipepsis ustulata* (Hymenoptera: Pompilidae). Ethology 112:691–698. https://doi.org/10.1111/j.1439-0310.2006.01204.x

Alcock J, Smith AP (1987) Hilltopping, leks and female choice in the Carpenter bee *Xylocopa (Neoxylocopa) varipuncta*. J Zool 211:1–10. https://doi.org/10.1111/j.1469-7998.1987.tb07448.x

Allison JD, Cardé RT (eds) (2016) Pheromone communication in moths: evolution, behavior, and application. University of California Press, Oakland, California

Amo L, Bonadonna F (2018) Editorial: The Importance of Olfaction in Intra- and Interspecific Communication. Front Ecol Evol 6:71. https://doi.org/10.3389/fevo.2018.00071

Andersen JF, Buchmann SL, Weisleder D, et al (1988) Identification of thoracic gland constituents from male *Xylocopa spp. latreille* (Hymenoptera: Anthophoridae) from Arizona. J Chem Ecol 14:1153–1162. https://doi.org/10.1007/BF01019343

Ayasse M, Paxton RJ, Tengö J (2001) Mating Behavior and Chemical Comminication in the order Hymenoptera. Annu Rev Entomol 46:31–78. https://doi.org/10.1146/annurev.ento.46.1.31

Baker TC (1989) Sex pheromone communication in the lepidoptera: New research progress. Experientia 45:248–262. https://doi.org/10.1007/BF01951811

Bergman TJ, Beehner JC (2008) A simple method for measuring colour in wild animals: validation and use on chest patch colour in geladas (*Theropithecus gelada*): MEASURING COLOUR IN WILD ANIMALS. Biol J Linn Soc 94:231–240. https://doi.org/10.1111/j.1095-8312.2008.00981.x

Cane (1987) Estimation of Bee Size Using Intertegular Span (Apoidea). J Kans Entomol Soc 60:145–147

Chemnitz J, Jentschke PC, Ayasse M, Steiger S (2015) Beyond species recognition: somatic state affects long-distance sex pheromone communication. Proc R Soc B Biol Sci 282:20150832. https://doi.org/10.1098/rspb.2015.0832

Chole H, Woodard SH, Bloch G (2019) Body size variation in bees: regulation, mechanisms, and relationship to social organization. Curr Opin Insect Sci 35:77–87. https://doi.org/10.1016/j.cois.2019.07.006

Coelho JR, Holliday CW (2001) Effects of Size and Flight Performance on Intermale Mate Competition in the Cicada Killer, *Sphecius speciosus* Drury (Hymenoptera: Sphecidae). J Insect Behav

Czaczkes TJ, Grüter C, Ratnieks FLW (2015) Trail Pheromones: An Integrative View of Their Role in Social Insect Colony Organization. Annu Rev Entomol 60:581–599. https://doi.org/10.1146/annurev-ento-010814-020627

Daimon T, Fujii T, Fujii T, et al (2012) Reinvestigation of the Sex Pheromone of the Wild Silk Moth *Bombyx mandarina* : The Effects of Bombykal and Bombykyl Acetate. J Chem Ecol 38:1031–1035. https://doi.org/10.1007/s10886-012-0164-0

Dani KGS, Fineschi S, Michelozzi M, et al (2021) Diversification of petal monoterpene profiles during floral development and senescence in wild roses: relationships among geraniol content, petal colour, and floral lifespan. Oecologia 197:957–969. https://doi.org/10.1007/s00442-020-04710-z

Darragh K, Byers KJRP, Merrill RM, et al (2019) Male pheromone composition depends on larval but not adult diet in *Heliconius melpomene*. Ecol Entomol 44:397–405. https://doi.org/10.1111/een.12716

Eltz T, Whitten WM, Roubik DW, Linsenmair KE (1999) Fragrance Collection, Storage, and Accumulation by Individual Male Orchid Bees. J Chem Ecol 25:157–176

Gomez-Diaz, Carolina, Benton, Richard (2013) The joy of sex pheromones. Eur Mol Biol Organ 14:874–883. https://doi.org/10.1038/embor.2013.140

Itagaki H, Conner WE (1988) Calling Behavior of *Manduca sexta* (L.) (Lepidoptera: Sphingidae) with Notes on the Morphology of the Female Sex Pheromone Gland. Ann Entomol Soc Am 81:798–807. https://doi.org/10.1093/aesa/81.5.798

Juárez ML, Devescovi F, Břízová R, et al (2015) Evaluating mating compatibility within fruit fly cryptic species complexes and the potential role of sex pheromones in pre-mating isolation. ZooKeys 540:125–155. https://doi.org/10.3897/zookeys.540.6133

Kassambara A (2020) ggpubr: "ggplot2" Based Publication Ready Plots

Lassance J-M, Löfstedt C (2009) Concerted evolution of male and female display traits in the European corn borer, *Ostrinia nubilalis* . BMC Biol 7:10. https://doi.org/10.1186/1741-7007-7-10

478   Leys R, Hogendoorn K (2008) Correlated evolution of mating behaviour and morphology in
479   large carpenter bees (*Xylocopa*). Apidologie 39:119–132.
480   https://doi.org/10.1051/apido:2007044

481   Liedo P, Orozco D, Cruz-López L, et al (2013) Effect of post-teneral diets on the performance of
482   sterile *Anastrepha ludens* and *Anastrepha obliqua* fruit flies: Pre-release feeding on
483   sterile *Anastrepha* male performance. J Appl Entomol 137:49–60.
484   https://doi.org/10.1111/j.1439-0418.2010.01568.x

485   Marshall LD, Alcock J (1981) The evolution of the mating system of the Carpenter bee *Xylocopa*
486   *varipuncta* (Hymenoptera: Anthophoridae). J Zool 193:315–324.
487   https://doi.org/10.1111/j.1469-7998.1981.tb03447.x

488   Mazerolle MJ (2020) AICcmodavg: Model selection and multimodel inference based on
489   (Q)AIC(c)

490   Minckley (1994) Comparative morphology of the mesosomal 'gland' in male large carpenter
491   bees (Apidae: Xylocopini). Biol J Linn Soc 53:291–308. https://doi.org/10.1111/j.1095-
492   8312.1994.tb01014.x

493   Minckley RL, Buchmann SL, Wcislo WT (1991) Bioassay evidence for a sex attractant
494   pheromone in the large carpenter bee, *Xylocopa varipuncta* (Anthophoridae:
495   Hymenoptera). J Zool 224:285–291. https://doi.org/10.1111/j.1469-7998.1991.tb04805.x

496   Mueller UG, Wolf-Mueller B (1993) A method for estimating the age of bees: Age-dependent
497   wing wear and coloration in the Wool-Carder bee *Anthidium manicatum* (hymenoptera:
498   Megachilidae). J Insect Behav 6:529–537. https://doi.org/10.1007/BF01049530

499   Nieberding CM, Fischer K, Saastamoinen M, et al (2012) Cracking the olfactory code of a
500   butterfly: the scent of ageing: The scent of ageing in a butterfly. Ecol Lett 15:415–424.
501   https://doi.org/10.1111/j.1461-0248.2012.01748.x

502   Oksanen J, Simpson GL, Blanchet FG, et al (2022) vegan: Community Ecology Package

503   Ostwald MM, Alba-Tercedor J, Minckley RL, Buchmann SL (2022) Three-dimensional
504   morphology of the hypertrophied sex pheromone gland in a lek-mating carpenter bee
505   (*Xylocopa sonorina*) revealed by micro computed tomography and scanning electron
506   microscopy. Apidologie 53:60. https://doi.org/10.1007/s13592-022-00967-w

507   Phelan PL, Baker TC (1986) Male-size-related courtship success and intersexual selection in the
508   tobacco moth, *Ephestia elutella* . Experientia 42:1291–1293.
509   https://doi.org/10.1007/BF01946428

510   R Core Team (2022) R: A Language and Environment for Statistical Computing. R Foundation
511   for Statistical Computing, Vienna, Austria

Rantala MJ, Kortet R, Kotiaho JS, et al (2003) Condition dependence of pheromones and
immune function in the grain beetle *Tenebrio molitor*. Funct Ecol 17:534–540.
https://doi.org/10.1046/j.1365-2435.2003.00764.x

Schiestl FP (2010) The evolution of floral scent and insect chemical communication: Evolution
of floral scent. Ecol Lett 13:643–656. https://doi.org/10.1111/j.1461-0248.2010.01451.x

Schindelin J, Arganda-Carreras I, Frise E, et al (2012) Fiji: an open-source platform for
biological-image analysis. Nat Methods 9:676–682. https://doi.org/10.1038/nmeth.2019

Thioulouse J, Dray S, Dufour A-B, et al (2018) Multivariate Analysis of Ecological Data with
ade4. Springer

Venables WN, Ripley BD (2002) Modern Applied Statistics with S, Fourth. Springer, New York

Wertheim B, van Baalen E-JA, Dicke M, Vet LEM (2005) PHEROMONE-MEDIATED
AGGREGATION IN NONSOCIAL ARTHROPODS: An Evolutionary Ecological
Perspective. Annu Rev Entomol 50:321–346.
https://doi.org/10.1146/annurev.ento.49.061802.123329

Wickham H (2016) ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York

Wickham H (2011) The Split-Apply-Combine Strategy for Data Analysis. J Stat Softw 40:1–29

Wickham H, Girlich M (2022) tidyr: Tidy Messy Data

Wickham H, Hester J, Bryan J (2023) readr: Read Rectangular Text Data

Wilke CO (2020) cowplot: Streamlined Plot Theme and Plot Annotations for "ggplot2"

Wyatt TD (2014) Pheromones and animal behavior: chemical signals and signatures, 2nd edition.
Cambridge University Press, Cambridge

STATEMENTS AND DECLARATIONS

Funding

S.R.R. was funded by the David and Lucile Packard Foundation (2014-40378).

Competing Interests

The authors have no relevant financial or non-financial interests to disclose.

Author Contributions

All authors contributed to the study conception and design. Material preparation, data collection were performed by Andrew J. Goffinet, Kathy Darragh, and Nicholas Saleh. The analysis was carried out by Andrew J. Goffinet and Kathy Darragh. The first draft of the manuscript was written by Andrew J. Goffinet, and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

                                        FIGURE LEGENDS

Figure 1: (a) Male *X. sonorina*. Photo credit: Sebastian Scofield. (b) Representative

chromatogram of a male's mesosomal gland extract, showcasing the three primary compounds

and the range of minor constituents. The peak labeled as 1 is 3,7,11-methyl-2,7,10-dodecatrienal,

the peak labeled as 2 is all trans-farnesal, and the peak labeled as 3 is all trans-geranylgeraniol.

Figure 2: (a) Amount of pheromone is positively related to male *X. sonorina* body size (ANOVA,

$F_{1,36}$=14.02, p=0.000632), using intertegular distance as a proxy. (b) Amount of pheromone is

negatively related with gland brightness (ANOVA, $F_{1,36}$= 11.32, p=0.001832), according to the

HSB scale. A higher value indicates a lighter color of the gland.

Figure 3: NMDS (nonmetric multidimensional scaling) plot illustrating in two dimensions the

variation in mesosomal gland contents of male *X. sonorina.* Size of each circle corresponds to

the absolute amount of pheromone. The impact of body size (with intertegular distance as a

proxy) and brightness are shown.

Figure 4: (a) Scatterplot of the relationship between male *X. sonorina* body size, with the

intertegular distance as a proxy, and the absolute abundance of all trans-farnesal. (b) Scatterplot

of the relationship between the absolute abundance of all trans-farnesal and brightness of the

mesothoracic gland, measured on the HSB scale, with a higher value corresponding to a lighter

colored gland. (c) Scatterplot of the relationship between male *X. sonorina* body size, with the

intertegular distance as a proxy, and the absolute abundance of all trans-geranylgeraniol. (d)

Scatterplot of the relationship between the absolute abundance of all trans-geranylgeraniol and

brightness of the mesothoracic gland, measured on the HSB scale, with a higher value

corresponding to a lighter colored gland.